Main Manuscript for

Human deep sleep facilitates faster cerebrospinal fluid dynamics
linked to brain oscillations for sleep homeostasis and memory

Makoto Uji[1], Xuemei Li[1], An Saotome[1,2], Ryosuke Katsumata[1,3], R. Allen Waggoner[1], Chisato
Suzuki[1], Kenichi Ueno[1], Sayaka Aritake[1,2], Masako Tamaki[1,4]*

1 RIKEN Center for Brain Science, Saitama, Japan

2 Saitama Prefectural University, Saitama, Japan

3 Chiba University, Chiba, Japan

4 RIKEN Cluster for Pioneering Research, Saitama, Japan

*Masako Tamaki

Email: Masako.tamaki@riken.jp

Author Contributions: MU and MT designed the research. MU, XL, AS, RK, RAW, CS, KU, SA
and MT performed the experiments. MU, SA and MT analyzed the data. MU and MT wrote the
paper. All confirmed the content.

Competing Interest Statement: The authors declare no competing interests.

Classification:

Major: Biological Sciences

Minor: Psychological and Cognitive Sciences

Keywords: Sleep, Cerebrospinal fluid dynamics, Brain oscillations, Slow-wave sleep, REM sleep

This PDF file includes:

Main Text
Figures 1 to 5

Abstract

How sleep maintains our healthy brain function has remained one of the biggest mysteries in neuroscience, medical settings, and daily lives. While cerebrospinal fluid (CSF) during sleep have been implicated in metabolic waste reduction in animals, how CSF dynamics are driven in the healthy human brain during deep sleep remains elusive. A myriad of research has shown that crucial cognitive processing manifests in slow wave and rapid-eye movement (REM) sleep, suggesting that a key to maintaining brain functions lies in deep sleep. By leveraging a simultaneous sparse-fMRI and polysomnography method, we demonstrate that deep sleep-specific faster CSF dynamics are associated with spontaneous brain oscillations in healthy young human participants. Slow waves and sleep spindles during slow-wave sleep and rapid eye movements and sawtooth waves during rapid eye movement (REM) sleep are tightly linked to low-amplitude faster CSF fluctuations. In contrast, slow waves during light sleep and arousals produced large but slower CSF signal changes. Furthermore, CSF signals are significantly faster in frequency during deep than light sleep. These brain oscillations during light and deep sleep recruited essentially different brain networks, with deep sleep involving memory and homeostatic circuits. Thus, human deep sleep has a unique way of enabling faster CSF dynamics that are distinctive from arousal mechanisms.

Significance Statement

Sleep is indispensable to our life, but its functions remain a significant mystery in the field of neuroscience. One of the most enigmatic issues in sleep is whether and how sleep regulates the CSF. The present study demonstrates deep sleep-specific faster CSF dynamics time locked to sleep brain oscillations in healthy young human participants. Slow waves and sleep spindles during slow-wave sleep and rapid eye movements and sawtooth waves during REM sleep are tightly linked to CSF fluctuations, contributing to faster CSF signals. Our results consistently demonstrate that human deep sleep has a unique way of enabling faster CSF dynamics that are distinctive from arousal mechanisms.

Main Text

Introduction

Sleep is crucial for the maintenance of healthy brain functions (1–3). Poor sleep results in a wide variety of health problems, including neurodegenerative diseases such as Alzheimer's disease (3, 4). However, how sleep maintains human brain functions has remained one of the biggest mysteries in neuroscience, medicine, and daily life. Cerebrospinal fluid (CSF) during sleep has recently been proposed as one of the essential mechanisms for supporting brain functions (5–9). Research on animal models has suggested that brain metabolite waste is reduced specifically during sleep (5). This process is proposed to rely on CSF flows to reduce metabolic waste from the brain (6). In rodents, the amount of CSF tracer influx increases during deep nonrapid-eye-movement (NREM) sleep or slow-wave sleep compared to during wakefulness (5). Clearance and CSF movement were facilitated by vasomotion associated with norepinephrine level during sleep (9). Additionally, accumulating body of evidence suggests that CSF during sleep plays an important role in metabolic clearance also in humans. Tracer substances administered to CSF were reduced from the brain after sleep, whereas sleep deprivation impaired the clearance (8). Another study has reported that impaired sleep quality was associated with increased CSF tracer enrichment (7).

While recent findings suggest the contribution of sleep to CSF clearance, how CSF dynamics are maintained in healthy human sleep has yet to be clarified. Due to safety concerns when MRI methods are simultaneously measured with polysomnography (PSG), which is necessary for sleep measurement (10), functional magnetic resonance imaging (fMRI) is a commonly used noninvasive

method that allows measuring CSF dynamics or strengths concurrently with PSG (11–13), while
several other MRI techniques have been proposed for measuring CSF dynamics independent of
PSG (14, 15). Hereafter we refer to the fMRI signals measured from ventricles as CSF signals. By
measuring fMRI simultaneously with PSG during sleep, the slow frequency component (~0.05 Hz)
in the CSF signals was found to be stronger during light sleep than during wakefulness in correlation
with EEG delta activity (11). What during light sleep has caused such slow fluctuations has been
unclear. Since K-complexes, a large abrupt biphasic EEG activity, and EEG arousals occur
frequently during light nonrapid-eye-movement (NREM) sleep (16, 17), neural or physiological
activities induced by these events may contribute to the slow CSF signal changes (9, 18). However,
crucially, spontaneous brain oscillations and events, including slow waves and sleep spindles,
continuously occur during deep sleep (19). If neural activity caused by spontaneous brain activities
contributes to modulation of CSF signals, the CSF signals should oscillate in link with these
oscillations. Additionally, the CSF signals should oscillate faster during slow-wave sleep than those
during wakefulness or light sleep due to the abundancy of these sleep events, especially slow
waves, when arousal is largely suppressed (20). These rationales led to our hypothesis that human
deep sleep plays a specific role in accelerating CSF signal changes, which should be tightly linked
to sleep brain oscillations and events.
72
We demonstrate that sleep brain oscillations during slow-wave sleep and REM sleep are indeed
associated with significant changes in the CSF signals in healthy young human participants. We
found that CSF signals fluctuate significantly faster during slow-wave sleep than during light NREM
sleep, time-locked to slow waves and sleep spindles. During REM sleep, rapid eye movements and
sawtooth waves were associated with the significant CSF signal increase. Since we did not find
correlation between the CSF signal changes and any of respiration, heat rates, and head motions,
none of respirations, heart rates, or head motions were likely to cause the differences in the CSF
signal changes that were found among sleep stages and brain oscillations. We further confirmed
that characteristics of brain networks activated during slow-wave and REM sleep are distinct from
those during light NREM sleep or arousals. The present study suggests a fundamental property of
deep sleep for the maintenance of brain functions and offers a clue as to why deep sleep matters
for the human brain.
85
86 **Results**
87
One of the challenges in using fMRI in sleep research is the acoustic noise in MRI scans that cause
frequent arousals and prolonged wakefulness, inhibiting participants from reaching deeper sleep
(10, 21). To acquire slow-wave sleep and REM sleep data efficiently, we utilized a sparse fMRI
sequence which had a 2.1 s quiet period between 0.9 s of MRI image acquisitions measured
simultaneously with PSG. This protocol was inspired by previous studies that successfully acquired
slow-wave sleep and REM sleep data using magnetic resonance spectroscopy (MRS), where the
duration of MRI signal acquisition was approximately 1 s, separated by quiet periods (22–24).
Leveraging the sparse method (**Fig. S1**; see *Experimental design* and *MRI acquisition* in
Materials and Methods), we acquired a total of 50 datasets from twenty-five young healthy
participants (mean 23.6 ± 4.8 years old, 15 female) during mid-afternoon nap. The average sleep
duration was 69.6 ± 2.9 min (*mean ± SE*), including sessions with slow-wave sleep and/or REM
sleep (**Table S1**). The fMRI data were denoised using cardiac and respiration signals (25, 26). The
CSF signals were extracted from each participant's lateral ventricles to measure the signal changes
caused by CSF, because these are the closest ventricles from the choroid plexi that have been
generally regarded as one of the main sources of CSF production (27, 28) (**Fig. 1a**). Any epochs
containing arousals (29) or motion artifacts, defined as an epoch containing artifacts that lasted
longer than 1s or a participant showing motion in the video recording (see *Sleep stage scoring* in

Materials and Methods), were removed from the analyses (except for the analyses targeting
arousals; **Fig. 3a**).
107
Slow waves and sleep spindles accompany CSF signal fluctuations during slow-wave sleep
If sleep brain oscillations are linked to the CSF signal changes, CSF signal peaks should be present
when the signals are time-locked to brain oscillations. Since slow-wave sleep is characterized by
ample slow waves in addition to sleep spindles (19, 29), the peak latency should be shorter during
slow-wave sleep than during light NREM sleep due to the differences in abundancy of these events.
To test whether CSF signal changes follow sleep brain oscillations and whether the latency differs
between different sleep stages, EEG signals recorded simultaneously with fMRI data had artifacts
removed, sleep-stage scored in accordance with American Academy of Sleep Medicine criteria
(29). Then slow waves and sleep spindles were detected. Specifically, high-amplitude slow waves
with peak-to-peak amplitude of larger than 140 μV (30, 31) and sleep spindles of frequency
between 12-15Hz with the duration of 0.5–3s (32) were detected from the EEG signals during light
NREM (N2 sleep stage) and slow-wave sleep (N3 sleep stage; for more details see **Sleep event**
detection and **fMRI time course analysis** in **Materials and Methods**, **Fig. 1b**). The CSF signal
changes were measured for each event. If sleep brain oscillations are associated with CSF signals,
significant changes in the CSF signals would be observed after the onset of oscillations.
123
The number of slow waves were significantly larger during slow-wave sleep than during light NREM
sleep (**Fig. 1c**; mean intervals between events, light NREM sleep, 73.9 ± 13.05 s; slow-wave sleep,
14.8 ± 3.52 s, *mean ± SE*; Wilcoxson signed-rank test, two-sided with Bonferroni correction: n =
27, z = 3.9401, P = 8.1448e-05), whereas spindles were detected more frequently during light
NREM sleep than during slow-wave sleep (**Fig. 1c**; mean intervals between events, light NREM
sleep, 5.3 ± 0.29 s; slow-wave sleep, 6.8 ± 0.41 s, Wilcoxson signed-rank test, two-sided with
Bonferroni correction: n = 30, z = 3.9289, P = 8.5348e-05).
131
CSF signal changes that occurred after brain oscillations exhibit different characteristics depending
on the depth of sleep. During light NREM sleep, in addition to sleep spindles that were correlated
with significant CSF signal changes at 6 s after onset (**Fig. 1d**, dark green), slow waves preceded
a large CSF signal peak at 8 s after onset (**Fig. 1d**, light green). During slow-wave sleep, slow
waves triggered CSF signal changes peaking at 5.5 s **(Fig. 1e**, light blue), which was shorter in
latency (**Fig. 1f**, Wilcoxson signed-rank test, two-sided with Bonferroni correction: n = 28, z =
3.0060, P = 0.0026) and smaller in amplitude (**Fig. 1g**, Wilcoxson signed-rank test, two-sided with
Bonferroni correction: n = 28, z = 3.8028, P = 1.4305e-04) than light NREM sleep. During slow-
wave sleep, sleep spindles again preceded CSF signal changes at 4 s after their onset, which
lasted for approximately 10 s (**Fig. 1e**, dark blue). The CSF signal changes were significantly
correlated with GM signal changes with GM preceding CSF signal changes (**Fig. S2**). These
findings demonstrate that CSF signal changes are indeed correlated with brain oscillations during
sleep and that deep sleep was characterized by shorter latency and smaller amplitude CSF signal
changes that follow GM signal changes.
146
To further confirm the relationship between the CSF signal changes and sleep brain oscillations,
we conducted an inverse analysis; we investigated the EEG topographical changes based on CSF
signal positive peak events (see **CSF peak locked EEG topography analysis** in **Materials and**
Methods). The segmented EEG data were averaged for each sleep stage (**Fig. S3**). The EEG
topographic map indicated a negative peak centered around the frontal region, resembling a
topography of K-complexes (33, 34), which peaked 8 s *before* the CSF positive peak events during
light NREM sleep (note that slow waves during light NREM sleep were followed by a CSF peak 8s
after the onset of slow wave events, **Fig. 1d**). During slow-wave sleep, a negative peak spread
through the midline of frontal to parietal regions resembling those found typically in slow waves (35)
which peaked around 5 s *before* the CSF positive peak events during slow-wave sleep (note that
slow waves during slow-wave sleep were followed by a CSF peak 5.5s *after* the onset of slow wave

events, **Fig. 1e**). These inverse analyses further confirmed that sleep brain oscillations, mainly slow
waves, and CSF signal fluctuations are linked.
160
CSF signal changes follow rapid eye movements and sawtooth waves during REM sleep
Rapid eye movements and sawtooth waves during REM sleep have been associated with memory
processing and conscious experiences (36, 37). Thus, if brain activities during REM sleep, not
limited to NREM sleep, contribute to CSF signal changes, CSF signals should show significant
changes that are time-locked to these events. To this end, we detected typical REM sleep events,
such as rapid eye movements and sawtooth waves observed during REM sleep (***Sleep event***
detection in **Materials and Methods**; **Fig. 2a**). We combined all of the REM sleep events together
since the number of events was limited. We found that REM sleep events accompanied the CSF
signal fluctuations 30 s after the onset of the events (**Fig. 2b**; see **Fig. S4a & b** for the results for
each event). In contrast to NREM sleep events, REM sleep events took a longer time to induce a
smaller CSF signal fluctuation (**Fig. 2c**). We next tested whether global brain activation occurs prior
to the CSF peaks, similarly to NREM sleep. We found that REM sleep events were also followed
by a GM signal increase at 20 s after onset (**Fig. 2d, Fig. S4c & d**), approximately 10 s before the
CSF peak. However, REM sleep events showed a significant positive, not negative, correlation
between the CSF and GM signals (**Fig. 2e**, n = 14, Pearson's r = 0.31 at lag -6.1 s, P < 1e-10, *CI*
= [0.16, 0.46]), suggesting different mechanisms for CSF and GM coupling between NREM and
REM sleep (see also **Fig. S2c & d** for CSF and GM coupling during NREM sleep).
178
Different contributions of sleep brain oscillations and arousal mechanisms to CSF dynamics
Thus far, our results indicate that brain oscillations and neural events during deep sleep are tightly
linked to CSF signal changes. Could these results be explained solely by cortical arousals, as
arousals can occur throughout during sleep? If so, similar changes in the CSF signals could follow
sleep-related brain oscillations and arousals. To investigate whether the contributions of sleep-
related brain oscillations to CSF fluctuations are based on arousals, we examined arousal-
associated CSF signal changes and directly compared them with slow wave-associated CSF signal
changes. Arousal events in the EEG signals were detected by sleep experts in accordance with
the standard criteria (29). The arousal events were defined as an abrupt shift in EEG frequency,
which was clearly visible in the EEG signals as in the standard AASM criteria (**Fig. 3a**; ***Sleep event***
detection in **Materials and Methods**). The CSF and GM signals were subsequently segmented
on the basis of each arousal onset.
191
The arousals occurred significantly more often during light NREM sleep (21.2 ± 1.9) than during
slow-wave sleep (0.3 ± 0.1; **Fig. 3b**; Wilcoxson signed-rank test, two-sided with Bonferroni
correction: n = 30, z = 4.7832, P = 1.7257e-06) or REM sleep (0.9 ± 0.2; **Fig. 3b**; Wilcoxson signed-
rank test, two-sided with Bonferroni correction: n = 19, z = 3.8242, P = 1.3121e-04). These findings
confirm that deep sleep is the least prone to arousals. Next, CSF signal changes time-locked to
arousal events were measured. The average CSF signals increased 8 s after the onset of the
arousal events and then returned to baseline 20 s after onset (**Fig. 3c**, black). Arousal-related CSF
and GM signal changes were similar across sleep stages (**Fig. 3c, d**), with the exception that the
CSF signal changes to arousals were barely visible during slow-wave sleep (**Fig. 3c**). Notably, the
CSF and GM signal changes that followed arousals and slow waves during light NREM sleep had
similar signal properties (**Fig. 3e, f**). The latency to reach the peak CSF signal was approximately
8 s (**Fig. 3e, h**), and CSF signal changes were followed by a large GM undershoot (**Fig. 3f**). In
contrast, slow waves during slow-wave sleep had a shorter latency (**Fig. 3e, h**; Wilcoxson signed-
rank test, two-sided with Bonferroni correction: n = 28, z = 3.3704, P = 7.5062e-04) and smaller
amplitude changes (**Fig. 3e, g**; Wilcoxson signed-rank test, two-sided with Bonferroni correction: n
= 28, z = 4.2810, P = 1.8603e-05) in the CSF signals as compared to the arousals (see **Fig. S5** for
each comparison). One may wonder whether the differences in the signal properties between
arousals and slow waves are simply due to the frequencies of the events, as a larger number of
events may induce larger accumulative changes. However, since the total number of slow waves
during slow-wave sleep (77.8 ± 21.8) was significantly greater than the total number of arousals

(22.7 ± 1.9; Wilcoxson signed-rank test, two-sided with Bonferroni correction: n = 28, z = 3.3933,
P = 6.9063e-04), this is unlikely.
214
215 These findings consistently indicate that arousals and slow waves during light sleep are followed
216 by slow and large CSF fluctuations, whereas slow waves during slow-wave sleep accompany fast
217 yet small CSF signal changes, suggesting that sleep depth-dependent brain oscillations and
218 arousals contribute differently to the CSF fluctuations.
219
Faster frequency CSF signal components during slow-wave sleep and REM sleep
Our results thus far indicate that CSF signal changes are time-locked to brain oscillations and
neural events independent of arousals. Furthermore, our results suggest CSF signals fluctuate
faster during slow-wave sleep than during light sleep. If so, this should result in stronger faster
frequency component in CSF-signal based power spectrum (38). To test this, we examined the
periodic oscillation components of CSF signals across different sleep depths (see ***Power spectrum***
analysis of fMRI data in **Materials and Methods**; **Fig. S6**). Our results revealed that the faster
component power (approximately 0.06-0.12 Hz) in the CSF signals was significantly greater during
slow-wave sleep than during wakefulness or light NREM sleep (**Fig. S6a**, blue). In contrast, the
slow component power (<0.06 Hz) in the CSF signals was significantly greater during light NREM
sleep than during wakefulness or slow-wave sleep (**Fig. S6a**, green). CSF dynamics during REM
sleep were characterized by a mixture of slow and very fast (>0.12Hz) components (**Fig. S6a**, red).
Similarly to light NREM sleep, the slow component power was significantly greater during REM
sleep than during slow-wave sleep or wakefulness. However, the power of the very fast component
was significantly greater during REM sleep than during light NREM sleep (**Fig. S6a**). These findings
demonstrate that both slow-wave sleep and REM sleep are indeed characterized by higher
frequency CSF dynamics than light sleep.
237
CSF signal changes associated with respiration, cardiac rhythms, and head motions
Since CSF signals are known to couple with respiration and cardiac pulses (39–41), we performed
control analyses and confirmed that none of the respiratory or cardiac rhythms were likely to have
caused the differences in the CSF signal changes among sleep stages and brain oscillations in this
study. First, the intervals of the events were longer during the deep sleep stages than during the
light sleep stages. Cardiac cycles were significantly longer during slow-wave sleep or REM sleep
than during wakefulness (slow-wave sleep vs. wakefulness: Wilcoxson signed-rank test, two-sided
with Bonferroni correction: n = 25, z = 4.3747, P = 1.2157e-05; REM sleep vs. wakefulness:
Wilcoxson signed-rank test, two-sided with Bonferroni correction: n = 15, z = 3.3004, P = 9.7656e-
04; **Fig. S7a**). There was no significant difference in respiratory cycles across different sleep depths
(**Fig. S7b**). Second, the periodic frequency changes across different sleep depths were not
produced by denoising processes during the fMRI preprocessing. To this end, we investigated the
periodic component of CSF signals across different sleep depths without denoising processes of
respiration and cardiac pulses following previous studies (25, 26). The key findings in the periodic
frequency changes remained regardless of the existence of respiration and cardiac cycles in the
fMRI signals (**Figs. S8**). Third, although a gradual increase occurred in the CSF signals after the
respiration and cardiac pulses, these changes were significantly smaller than the CSF signal
changes that occurred after brain oscillations (**Fig. S9**). Fourth, the cardiac and respirations had
higher frequency signals (cardiac: around 1Hz; respiration: around 0.3Hz; **Fig. S10**) that did not
overlap with the frequency range induced by sleep brain oscillations (<0.15 Hz). Additional control
experiments and analyses were conducted to confirm that the aliasing noises from the rhythm of
respirations and cardiac pulses (42) were not the main factor inducing the frequency peak observed
at ~0.06Hz in the CSF signals (**Fig. S11; *The control experiment and analyses to test aliasing***
noises in **Materials and Methods**). Thus, these physiological signals are not the main causes of
the faster oscillating CSF signals found during slow-wave sleep and REM sleep.
263
Additional control analyses on head motions indicated that head motions were unlikely to have
caused the association between sleep brain oscillations and CSF signal changes. First, we

examined differences in the estimated distance of head motion during fMRI scans for each of sleep
stages (see *MRI data analysis* in **Materials and Methods** for head motion measurement). We
found that head motion during slow-wave sleep (0.021 ± 0.002 mm/volume) was significantly
smaller than that during light NREM sleep (0.027 ± 0.002 mm/volume, Wilcoxson signed-rank test,
two-sided with Bonferroni correction: n = 29, z = 4.7030, P = 2.5631e-06) or wakefulness (0.025 ±
0.002 mm/volume, Wilcoxson signed-rank test, two-sided with Bonferroni correction: n = 20, z =
3.9199, P = 8.8575e-05). Thus, the effect of head motion, if any, would have been minimal during
deep sleep compared with light sleep. Next, correlation coefficients between head motion values
and CSF and GM signals were measured by Pearson's correlation. The group mean correlation
coefficients were 0.06±0.02 (mean ± SE) between CSF signals and head motions and -0.09±0.03
(mean ± SE) between GM signals and head motions, indicating very small correlation with head
motions.
278
Relationship between CSF signal in the lateral ventricles and other regions
How might the CSF signals in the lateral ventricles relate to signals in the other regions? The
present study extracted CSF signals from the lateral ventricles instead of the fourth ventricles.
Thus, one might wonder whether these signals represent the ventricle signal itself or residuals
produced by some artifacts (head motion artifacts). To check the relationship between the lateral
ventricle signals and fMRI signals in other regions, we first examined zero-lag correlations between
the CSF signals from the lateral ventricles and signals in each of the whole-brain voxels (see *Voxel-*
wise correlation analyses: zero-lag correlation and cross-correlation time-lag map in
Materials and Methods). In the group mean voxel-wise zero-lag correlation map (**Fig. S12a**), we
found positive correlations within the ventricular regions, including the lateral and the fourth
ventricles. We also observed wide-spread negative correlations between the lateral ventricle
signals and grey matter regions. Importantly, the strongest positive correlations (r = 0.3)
concentrate in the lateral ventricles.
292
Next, we examined the time-lag of the strongest positive correlation between the lateral ventricle
signals and signals in each of the whole-brain voxels (see *Voxel-wise correlation analyses: zero-*
lag correlation and cross-correlation time-lag map in **Materials and Methods**). We found that
signals in the grey matter regions generally precede lateral ventricle signals, whereas small lags
can be found within in the ventricular regions (**Fig. S12b**). Additionally, positive lags are found in
the subarachnoid space, meaning the lateral ventricle signals precede the subarchnoid signal
changes. These findings are consistent with a previous study investigated the fourth ventricle
signals (43). Thus, the CSF signals in the present study represent ventricle signals, having tight
temporal relationship with grey matter and subarchnoid signal changes.
302
Sleep-depth-dependent and arousal-associated changes in the brain network
Because brain oscillations during slow-wave sleep are involved in plasticity and homeostatic
regulation (44–46), while K-complexes during light sleep can accompany arousals (16, 17, 33, 34),
we next asked whether the different modulations of CSF and GM signals found during light sleep,
deep sleep, and arousals could be associated with different brain circuits involved during different
depths of sleep. To this end, we applied a generalized linear model (GLM) analysis on the fMRI
signals to each event. We found that different cortical networks were recruited depending on the
depth of sleep (**Fig. 4**, **Tables S3–9**). The prefrontal and hippocampal regions, striatum, and
amygdala were more strongly recruited to slow waves during slow-wave sleep than during light
NREM sleep, consistent with previous studies investigating high-amplitude slow waves during
slow-wave sleep (31) (**Fig. 4**, **Table S4**), whereas sensorimotor areas, including the parietal and
visual cortices and cerebellum, were more strongly activated with slow waves during light NREM
sleep than during slow-wave sleep, again replicating previous studies (47) (**Fig. 4**, **Table S3**).
These findings indicate that different cortical processing underlies slow waves during light NREM
vs. slow-wave sleep. Additionally, we found that more localized cortical brain regions were recruited
to sleep spindles during slow-wave sleep than during light NREM sleep (48, 49) (**Tables S7 & S8**).
REM sleep events activated brain regions, including visual areas, the thalamus, the hippocampus,

the striatum, and the amygdala, also consistent with previous studies using fMRI and PET (50–52)
(**Fig. 4, Tables S5 & S9**). In contrast to slow-wave sleep or REM sleep, arousals involved whole-
brain activations not localized to any regions (**Fig. 4**, **Table S6**). Taken together, these results
demonstrate that brain oscillations involved in regulating CSF dynamics recruit essentially different
networks depending on the depth of sleep.
325
326 **Discussion**
327
The present results robustly indicate that deep human sleep has a unique way of interacting with
CSF signals, as indicated in **Table S2**. Slow waves and sleep spindles during slow-wave sleep are
followed by faster changes in CSF fluctuations. Neural events during REM sleep, such as rapid eye
movements and sawtooth waves, also accompany CSF signal changes. We further revealed
distinct frequency profiles in CSF signals among different sleep stages. The frequency of CSF
signals become faster during deep sleep than during wakefulness or light sleep. In contrast,
arousals and brain oscillations during light sleep were followed by infrequent, sharp CSF changes.
We confirmed that different brain networks are activated after brain oscillations or events during
different depths of sleep. Deep sleep recruits brain regions involved in memory and homeostatic
regulation, whereas light sleep and arousal recruit whole-brain activation, including the sensory-
motor network.
339
The present study suggests that at least two different processes are involved in regulating CSF
dynamics in human sleep. The first process is prominent during light sleep when arousals and
arousal-related brain activities, such as K-complexes, take the lead in modulating CSF oscillations.
The second process is facilitated specifically during deep stable sleep when arousals are largely
suppressed. These results further suggest that during deep sleep, CSF oscillations are facilitated
by spontaneous brain oscillations and neural events, such as slow waves, sleep spindles, and REM
sleep events. Since sleep events occur more often than arousals during deep sleep, it is highly
likely that sleep events, not arousals, are the major contributors to CSF oscillations during deep
sleep. We propose that these two processes are supported by distinctive mechanisms, as shown
in **Fig. 5**. The process during light NREM sleep is based on arousal mechanisms ("arousal-based
mechanism"), in which transient increases in brain activity facilitate CSF oscillations by arousing
stimuli to subcortical regions. Another process that we propose here functions specifically during
deep sleep ("deep sleep-based mechanism"), which is tightly linked to sleep brain oscillations to
drive the CSF oscillations. This deep-sleep mechanism may work continuously, producing faster
CSF oscillations by tightly interacting with sleep-related brain oscillations. In contrast to the arousal
mechanism that induces sharp and transient CSF signal increase, the deep-sleep mechanism may
induce low-amplitude oscillations, potentially causing minimal load to the vascular system while
enabling continuous faster oscillations.
358
Although the direct relationship between CSF signal changes during deep sleep and waste
clearance is unclear from the present study, the present results provide some important
implications. First, we found that slow waves are among the main oscillations that produce faster
CSF circulation during deep sleep. Given the tight relationship between slow waves and
homeostatic sleep needs (7, 8, 44), faster CSF circulation associated with slow waves might work
well for clearing waste products that might have accumulated during prolonged wakefulness.
Second, the present results suggest that sleep spindles also accelerate CSF oscillations. Sleep
spindles are known to facilitate episodic and motor learning through plastic changes in cortical and
subcortical regions (53–55). Our analysis has indeed shown that crucial brain network regions
involved in episodic and motor learning (e.g., hippocampus and striatum) were recruited during
sleep spindles during slow-wave sleep, consistent with previous studies (48). Thus, it is possible
that CSF oscillations are coupled not only to sleep homeostasis but also to plasticity processes,
where sleep spindles might play a major role.
372

While we found faster CSF dynamics during REM sleep as well as during slow-wave sleep, the
roles of CSF oscillations might differ. First, the latency to the CSF peak was much longer during
REM sleep than during slow-wave sleep. Second, we found that not only the very fast component
(> 0.12 Hz) but also the slow component (< 0.06 Hz) was present in the CSF frequency profiles
during REM sleep. Third, the CSF peak did not accompany the GM undershoot during REM sleep.
In addition to the rapid eye movements and sawtooth waves we analyzed in this study, EEG theta
and alpha activities can be observed, and muscle twitches occur intermittently during REM sleep,
which were not examined here. Since sympathetic nervous system activity, which regulates heart
rates and blood pressure, is increased during REM sleep, it is possible that autonomic variability
(e.g. respiratory and cardiac variabilities) in REM sleep also contributes to the CSF signal changes
during REM sleep, as was proposed previously (39, 56).
384
One may wonder whether cardiac and respiratory rhythms cause differences in the CSF peaks
during different sleep stages and brain oscillations since CSF signals are known to be coupled to
cardiac and respiratory rhythms (40, 57, 58). However, these signals are unlikely to be the sources
of the differences in the CSF signal changes among sleep stages and brain oscillations found in
the present study for the following three reasons. First, the intervals of the cardiac signals were
significantly longer during slow-wave sleep or REM sleep than during wakefulness. There was no
significant difference in the respiratory signals across different sleep depths. If these factors
contribute to changes in the CSF signal, the CSF signals should be slower during deep sleep than
during light sleep. Thus, these rhythms are not in accordance with the significantly faster CSF
signals found during slow-wave sleep and REM sleep. Second, we investigated the effects of
respiration and cardiac pulses in fMRI signals on the periodic frequency changes in CSF signals.
The key findings of the periodic frequency changes in the CSF signals remained regardless of the
existence of respiration and cardiac pulses in the fMRI signals. Third, previous studies have
indicated that cardiac- and respiration-driven CSF signal changes occur 0.45–1.58 s after the
events (40). However, CSF signal changes caused by slow waves were in the range of 5.5–8 s
after the events, much later than the cardiac and respiration-driven changes that might have
occurred. Additionally, we investigated the CSF signal changes that followed respiration and
cardiac pulses. We found that respiration- and cardiac-driven signal changes were significantly
smaller than sleep-related brain oscillations and arousals. Thus, the CSF peaks caused by cardiac
and respiratory rhythms, if any, should not overlap with the CSF peaks that appeared after brain
oscillations. Finally, the results of control analyses showed that the aliasing noises from respirations
and cardiac pulses were unlikely to be the main factor inducing the frequency peak observed at
~0.06Hz in the CSF signals found in the present study.
408
The present study at least partially suggests why and how deep sleep matters for healthy human
cognitive functions. Sleep disturbances are a major issue worldwide and are associated with
various health-related issues, including cognitive impairments, neurodegenerative diseases and
aging (59–62). Importantly, the most striking change found in these cases is the reduction in deep
stable sleep. In other words, light NREM sleep most likely remains regardless of the severity of the
disorder. Along with evidence showing a strong relationship between the stages of sleep, CSF
dynamics, and metabolite clearance (5, 7, 8, 63), dysfunctions in deep sleep may have severe
impacts on cognitive functions at least partially via altered CSF flows.
417
Although the present study did not directly measure the underlying mechanisms of the tight link
between sleep brain oscillations and the CSF signals, one of the likely mechanisms may involve
vascular dynamics. The vascular dynamics may change the size of the perivascular spaces of the
penetrating arterioles where the CSF circulate throughout the brain (8, 64). Since neural activity,
vascular dynamics, and CSF dynamics tightly interact with each other (11, 65), synchronous neural
activities, such as slow waves, sleep spindles, and various REM-sleep events may trigger
vasomotion, which may promote CSF circulation. Based on the present findings, which showed
that deep sleep involves accelerated CSF signal changes, these underlying processes may be
more pronounced during deep stable sleep than during light sleep or wakefulness. These further

suggest that deep sleep has a specific role in maintaining the tight physiological and neural balance
and that light sleep may not be able to compensate for the loss of deep sleep caused by sleep
restrictions, aging, or diseases.
430
The following four points need to be addressed in future research. First, our data were obtained
from midafternoon naps. Thus, what happens during nightly sleep when sleep needs are greater is
unclear. As the duration of slow-wave sleep is the longest and the number of slow waves is the
maximum during the first half of nightly sleep (29), it is possible that CSF oscillations are much
greater. REM sleep, in contrast, is longest during the early morning, and REM density is greater
(29). Thus, REM-related CSF dynamics could benefit from early morning studies. Additionally, our
data are restricted to approximately 1.5 hours, corresponding to the first sleep cycle. Thus, how
CSF signal fluctuations change over multiple sleep cycles is unclear. Second, the present study
used a sparse fMRI sequence with longer TR (3s) than previously used (~1 s) to prioritize the
acquisition of deep sleep data, rather than using a fast fMRI method to measure the inflow effects
in the CSF signals from the fourth ventricle (11, 66, 67). Nevertheless, consistent with the previous
studies investigated the lateral ventricles (68–70), our data clearly indicate that CSF signals are
differently modulated during wakefulness vs. sleep. Third, it was not optimized to measure the
signal changes driven by autonomic activity, including the cardiac and respiratory rhythms that
were associated with CSF pulsations (12, 13, 39, 70). Fourth, since the present study did not
directly measure the amount of metabolite waste or investigate the causal relationship between
CSF and metabolites, how CSF during deep sleep contributes to waste clearance in humans needs
to be clarified in future research.
449
Taken together, the results of the present study reveal that CSF signal changes are tightly
associated with sleep-brain oscillations and events during slow-wave sleep and REM sleep. These
CSF dynamics during deep sleep could be distinct from the arousal-based CSF dynamics for the
following reasons. First, the latency and amplitude characteristics are different. Second, the
frequency components are different among different sleep stages. Third, the relationships with
whole GM signals are different. Fourth, the brain network changes after sleep brain oscillations
during light vs. deep sleep and after arousals are different. While the causal relationship between
CSF signals during sleep and brain functions is unclear, the present study collectively suggests
that deep sleep-based CSF dynamics are the key to understanding how healthy brain functions are
maintained through sleep in humans.
460
461   **Materials and Methods**
462
Participants
Twenty-five young healthy participants (mean age of 23.6 ± 4.8 years, 15 female) participated in
this study. All of the participants provided written informed consent for their participation prior to
the start of the study. This study was approved by the institutional review board at RIKEN. There
were no studies that tested CSF dynamics during deep sleep prior to the present study. Previous
studies using fMRI during sleep in human participants (11, 47, 71) have reported n=11–20
subjects. However, to ensure that the results of our study were reliable and replicable, we used
25 subjects.
471
Screening process for eligibility
Before consent was obtained, the screening process for eligibility took place to confirm the
following aspects. First, the participants were between 18 and 35 years old. Second, on the basis
of a self-report questionnaire (23), anyone who had neurological or psychiatric conditions (e.g.,
epilepsy, migraine, stroke, chronic pain, major depression, anxiety disorder, psychotic disorder),
was currently using medication, or was suspected of having a sleep disorder (including insomnia,
sleep apnea syndrome, hypersomnolence, restless legs syndrome, parasomnias) was not eligible
to participate. Third, participants who had an irregular sleep schedule, i.e., those whose
sleep/wake time differed by more than 2 hours between weekdays and weekends, were not

eligible. We also ensured that the subjects had not taken a trip to a different time zone during the
last six months prior to the experiment. Participants who worked night shifts or who had a habit of
taking a nap were not eligible for this study. Fourth, a Pittsburgh Sleep Quality Index (PSQI) was
used to assess sleep quality and disturbances over a one-month time interval before the
experiments, and a PSQI score within 5 was used as an inclusion criterion (72). Fifth, those who
failed to complete an MRI safety questionnaire (i.e., claustrophobic, pregnant, having metallic
objects in the body, such as vascular clips, prosthetic valves, metal prostheses, metal fragments,
pacemakers, or dental braces) were not eligible for this study.
489
Experimental design
All eligible subjects participated in two sleep sessions approximately one week apart. The
participants arrived at the experimental room at approximately noon. The participants first
answered a questionnaire to report their bedtime, wake-up time, and duration of sleep the night
before. They also answered the Stanford Sleepiness Scale (SSS) rating (73), which ranges from
1 (feeling active, vital, alert, or wide awake) to 7 (no longer fighting sleep, sleep onset soon;
having dream-like thoughts). The subjects were asked to choose the scale rating that described
their state of sleepiness. Then, the electrodes for PSG were attached, which took approximately
one hour (see the PSG measurement section below). After the electrodes were attached, each
sleep session started in the early afternoon for simultaneous measurement of MRI and PSG (see
the MRI acquisition section below). The sleep session was conducted for less than two hours in
the MRI scanner including structural and functional image acquisition. After the sleep session,
they answered a questionnaire about the sleep content (data not shown).
503
The participants were instructed to maintain their regular sleep-wake habits, i.e., their daily
wake/sleep time and sleep duration, until the study was over. The sleep-wake habits of the
participants were monitored by a sleep log and an actigraphy device (MTN-221, ACOS, CO.,
LTD, Japan; SleepSign Act ver.2, KISSEI COMTEC, CO., LTD, Japan) during the study.
Throughout their participation, the subjects were instructed to refrain from taking naps. One day
prior to the experiments, alcohol consumption was not allowed. Unusual excessive physical
exercise and caffeine consumption were not allowed on the day of the sleep sessions.
511
MRI acquisition
MRI data were acquired with a 3T Siemens Prisma MRI scanner (Siemens Healthineers,
Germany) using a 64-channel head and neck coil at RIKEN. An anatomical T1-weighted image
was acquired using a 3D magnetization-prepared rapid gradient echo (MPRAGE) sequence (TR
= 2180 ms, TI = 1100 ms, TE = 2.95 ms, FA = 8°, voxel size = 0.7×0.7×0.7 mm^3, FOV = 290 ×
320 × 256 mm). Functional MRI data were acquired using a sparse gradient-echo echo-planar
imaging sequence (TR/TE=3000/30 ms, FA=85°, Multiband factor=3, 1800 volumes, GRAPPA
factor=3, bandwidth=752 Hz/pixel, B1+RMS=0.5 μT, MR acquisition time=0.9 s, and quiet
period=2.1 s) (74). 42 transverse slices (voxel size = 3x3x3 mm^3) orientated parallel to the AC-PC
plane were acquired if successfully covering the entire cerebrum. If not, the slice alignments were
adjusted accordingly to cover the entire cerebrum. Cardiac and respiratory cycles were
simultaneously recorded using a fingertip PPG sensor and respiratory belt (BIOPAC, US) with a
100 Hz sampling rate. Anatomical and functional MRI data were acquired simultaneously with
polysomnography (see the ***PSG measurement*** section below). Cushions and gauze were used
to stabilize the subjects' heads to reduce discomfort and head motion. Back and knee cushions
were used upon the participants' request to further reduce discomfort. Several blankets were
used to keep the participants warm and to initiate sleep during the scan (22, 23).
529
PSG measurement
EEG data were acquired using a BrainAmp MRplus EEG amplifier (Brain Products, Gilching,
Germany) with a 5 kHz sampling rate and an MR-compatible 32-channel EEG cap (EasyCap,
Wörthsee, Germany). The hardware bandpass filters were set to a 0.016–250 Hz range, with a
roll-off of 30 dB/octave at high frequency. The electrode layout followed the extended

international 10-20 system with additional drop-down channels or bipolar electrodes for recording
the electrocardiogram (ECG), electrooculogram (EOG), and chin electromyogram (EMG). EEG
was measured from either 27 electrodes (16 participants) or 31 electrodes (9 participants). For all
of the participants, EOG was measured from electrodes placed at the outer canthi. EMG was
measured from the mentum. ECG signals were measured from the lower shoulder blade. FCz
was used as the reference electrode, whereas AFz was used as the ground electrode. EEG data
were transferred outside the scanner room through fiber optic cables to a computer running the
BrainVision Recorder software (Brain Products, Gilching, Germany). MR-EEG scanner clocks
were synchronized (Brain Products Synchbox) for all EEG data acquisitions, and the onset of
every scanner repetition time (TR) period was recorded in the EEG data for MR gradient artifact
correction. The impedance of all electrodes was initially maintained below 10 kΩ and was
maintained below a maximum of 20 kΩ for the duration of the study.
547
PSG data preprocessing
PSG data were contaminated by artifacts due to the strong magnetic field. To remove the artifacts
from the PSG data, gradient artifacts were first corrected using BrainVision Analyzer2 (Version
2.2.0, Brain Products GmbH, Gilching, Germany) with the standard average artifact subtraction
(AAS) method. The AAS method uses sliding window templates formed from the averages of 21
TRs, which are subtracted from each occurrence of the respective artifacts for each electrode
(75, 76). The gradient-artifact corrected data were subsequently bandpass filtered (EEG: 0.1–50
Hz with a notch filter of 50 Hz) and downsampled to 250 Hz. Following the GA correction, cardiac
R-peaks were automatically detected from the ECG recording in BrainVision Analyzer2, checked
visually and corrected manually if necessary. These R-peak events were used to inform
ballistocardiogram correction. The BCG artifacts were further corrected via the AAS method with
sliding window templates formed from the averages of 21 R-peaks in BrainVision Analyzer2 (10,
77). These procedures were conducted for the whole duration of the fMRI scan to obtain artifact-
removed PSG data for further processing (see ***Sleep stage scoring*** and ***Sleep event detection***
below).
563
Sleep stage scoring
Sleep staging was conducted on the artifact-removed PSG data following the standard criteria
(29). Prior to sleep staging, PSG data were further bandpass filtered between 0.3 and 40 Hz for
EEG, EOG and ECG and between 10 and 50 Hz for EMG to remove low-frequency drift and high-
frequency noise and were re-referenced to the linked mastoids (TP9 & TP10). Scoring of the
sleep stages was performed every 30 s in conjunction by two trained sleep scorers with over
twenty years of experiences (MT & AS). We obtained sleep parameters, including the total sleep
time and duration of stage wakefulness (W), NREM 1 (N1), NREM 2 (N2), NREM 3 (N3), and
stage REM (REM) (**Table S1**). For further analysis, N1 and N2 were combined as light NREM
sleep, and N3 was categorized as slow-wave sleep. Epochs with motion artifacts and arousal
events (29) were marked and removed from further analyses, whereas the arousal events were
used only for arousal analysis. Motion artifacts were defined as an epoch containing artifacts that
lasted longer than 1s or a participant showing motion in the video recording.
577
578 ***Sleep event detection***
579 Using the artifact-removed PSG data, sleep events, including slow waves, sleep spindles, rapid
580 eye movements, sawtooth waves, and arousal events, were detected as follows.
581
Slow waves
Slow waves were detected semiautomatically as follows. First, EEG signals were bandpass
filtered between 0.1 and 4 Hz with a zero-phase 4th-order Butterworth filter. Then, slow waves
were detected automatically in accordance with a previous study (30) using Wonambi software
(https://wonambi-python.github.io/) from the F3, Fz, F4, C3, Cz, C4, P3, Pz, and P4 electrodes
during the N2 and N3 stages. Only the waves that met the following three criteria were selected
for detection of well-defined large amplitude slow waves: 1) a positive-to-negative zero crossing

and a subsequent negative-to-positive zero crossing separated between 0.3 s and 1 s, 2) a
negative peak between the two zero crossings with voltages smaller than -80 µV, and 3) a peak-
to-peak amplitude greater than 140 µV. The automatically detected slow waves were then
reviewed and removed by an experimenter if necessary. The number of slow waves was
measured for each electrode, and the electrode that had the maximum number of slow waves
was used as the most sensitive electrode for slow wave detection. The onset and duration of slow
waves were measured from the specific electrode for further analysis of CSF and GM signals
(see *fMRI time course analysis* and *fMRI GLM analysis* below).
597
<u>Sleep spindles</u>
Spindles were detected semiautomatically. As the first step, spindles were detected automatically
(32) using Wonambi software from the F3, Fz, F4, C3, Cz, C4, P3, Pz, and P4 electrodes during
the N2 and N3 stages in the following steps: EEG signals were filtered between 12–15 Hz, which
corresponded to the sleep spindle frequency band. Then, the root mean square (RMS) of the
filtered signals was subsequently calculated at every sample point using a moving window of 0.2
s. The threshold for spindle detection in the RMS signal was set to 1.5 standard deviations of the
filtered signal for each electrode. A spindle was detected when the RMS signal remained above
the threshold for 0.5–3 s, and then the beginning and end of the spindle were marked at the
threshold crossing points. When sleep spindles from multiple electrodes overlapped within a 5 s
time window, those sleep spindles were treated as trains of continuous sleep spindles and
merged. In the second step, the automatically detected spindles were reviewed and removed by
an experimenter if necessary. The onset and duration of sleep spindles were used for further
analysis of CSF and GM signals (see *fMRI time course analysis* and *fMRI GLM analysis*
below).
613
<u>Rapid eye movements</u>
Rapid eye movement events were detected during REM sleep. Rapid eye movement events were
automatically detected on two EOG signals (78). Specifically, to separate rapid eye movements
from slow eye movements (SEMs), two channels of EOG data are filtered using a fourth-order
Butterworth bandpass filter with cutoffs at 1 and 9 Hz to yield the filtered EOG data. The
candidate REM sleep events were generated as the negative instantaneous product of the two
filtered EOG sequences, and the REM sleep events were detected on the basis of their amplitude
threshold. The amplitude threshold was arbitrarily determined on the basis of each individual
dataset, since, owing to differences in the setup and/or the quality of the artifact removal, there
were individual differences in the amplitude of the EOG data. The onset and duration of rapid eye
movements were used for further analysis of the CSF and GM signals (see *fMRI time course*
analysis below).
626
<u>Sawtooth waves</u>
Sawtooth waves were manually marked on the scalp EEG independently by two trained sleep
scores (MT & SA) for the REM sleep episodes. These waves were defined as bursts of
consecutive surface-positive 2–5 Hz frontocentral bilateral synchronous symmetric waves with
amplitudes of 20–100 µV and with a slow incline to a negative peak with a subsequent steep
linear decline ending at a positive peak (79–81). The minimum number of consecutive waves was
set as 3. The consensus of the markings obtained from the two raters was selected, and the
onset and duration of sawtooth waves were used for further analysis of the CSF and GM signals
(see *fMRI timecourse analysis* below).
636
<u>Arousals</u>
An arousal was defined as an abrupt shift in EEG frequency, which may include theta, alpha
and/or frequencies greater than 16 Hz (but not spindles) that were at least 3 seconds in duration,
with at least 10 seconds of stable sleep preceding the change, in accordance with the standard
AASM criteria (29). Arousal events were manually scored by two trained sleep experts (MT & SA)
by visually checking the EEG signals. The consensus of the arousal scoring was used, and the

onset and duration of arousal were used for further analysis of the CSF and GM signals (see
fMRI time course analysis and *fMRI GLM analysis* below).
645
MRI data analysis
High-resolution T1-weighted images were processed using FreeSurfer 7.4.0 (82)
(https://surfer.nmr.mgh.harvard.edu/fswiki/FreeSurferWiki) to segment the head into five tissues
(i.e., the scalp, skull, cerebrospinal fluid (CSF), gray matter and white matter), and an anatomical
atlas. Anatomical surfaces such as the pial surface, gray/white matter interface and mid surface
(i.e., a middle layer of the gray matter) were estimated using FreeSurfer (83). The lateral
ventricles and whole gray matter region (GM) were extracted as the two main regions of interest
(ROIs) from each individual participant's anatomical image. The fMRI data were motion-corrected
and slice-timing corrected using the AFNI (https://afni.nimh.nih.gov) and RIKEN fMRI processing
programs (http://www.fmri.brain.riken.jp/~mri/Software.html). Furthermore, the fMRI data were
denoised using cardiac and respiration signals (25, 26). The two ROI images (GM and lateral
ventricles) were subsequently registered to each fMRI image using FSL FLIRT
(https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FLIRT), and the fMRI signals were z score transformed and
averaged across the voxels inside each ROI for each sleep session and each participant. The
distances of head motion during fMRI scans were estimated by MCFLIRT in FSL (84, 85).
661
fMRI time course analysis
fMRI data in each ROI were segmented based on sleep events (slow waves, spindles, rapid eye
movements, sawtooth waves, and arousals). First, -9–0 s (3TRs) before the onset of the events
was defined as a baseline. This baseline signal was averaged and subtracted from 0–30 s after
the onset of the events. These data were averaged across the number of each event for each
sleep session and then averaged across the sleep sessions and participants.
668
Cardiac and respiration peak-related fMRI time course analysis
Cardiac and respiratory cycles during each sleep session were recorded using a fingertip PPG
sensor and respiratory belt (BIOPAC, US) with a 100 Hz sampling rate. Peaks of the cardiac and
respiration cycles were detected using RIKEN fMRI processing programs
(http://www.fmri.brain.riken.jp/~mri/Software.html). fMRI data in each ROI were segmented based
on the peaks of the cardiac and respiration signals. First, -9–0 s (3TRs) before the onset of the
peak events was defined as a baseline. This baseline signal was averaged and subtracted from
0–30 s after the onset of the peak events. These data were averaged across the number of each
event for each sleep session and then averaged across the sleep sessions and participants.
678
Power spectrum analysis of fMRI data
Time-frequency analysis was applied to the whole fMRI signals during a sleep session of the two
ROIs using a multitaper approach (74, 86) in the Fieldtrip toolbox (87)
(http://www.ru.nl/neuroimaging/fieldtrip). Time windows of 200 s duration were moved across the
data in steps of 3 s corresponding to 1 TR, resulting in a frequency resolution of 0.005 Hz. The
calculated time-frequency representations (TFRs) were then transformed to the scale of dB by
10*log10, then demeaned by subtracting the mean power at each frequency during the whole
recording to calculate power fluctuations. Then, they were segmented based on sleep stages
every 30 s and averaged across each 30 s to extract each epoch power spectrum after removing
arousal- and motion-contaminated epochs. On the basis of the resulting power spectrum, periodic
components of the power spectrum were extracted after the estimated 1/f aperiodic components
were removed by fitting oscillations and one-over-f (FOOOF) algorithms (88). The resulting
periodic power spectra were averaged for each sleep stage first and then across sleep sessions
for each ROI.
693
694   *Correlation analysis between CSF and GM signals*
695   Sleep-specific event-locked (i.e., slow waves, spindles, and REM sleep events) time courses of
696   the CSF and GM signals based on their onset times (-9 s to 30 s) were used to investigate the

697 correlations between the CSF and GM signals. The correlation coefficients and time lags were
698 extracted when the absolute coefficient values were at their maximum between the two signals.
699
Voxel-wise correlation analyses: zero-lag correlation and cross-correlation time-lag map
To calculate voxel-wise zero-lag correlation, voxel-wise maps of correlation between the CSF
signal from the lateral ventricles and all voxels were computed using Pearsons' correlation
separately for each nap session. Next, we applied a Fisher Z transform prior to averaging then
group mean group mean voxel-wise zero-lag correlation map was generated by averaging the
individual maps. We also performed two-tailed one-sample t-tests (uncorrected $P < 0.001$) to
identify voxel regions with significant differences in correlation with the CSF signal compared with
zero.
708
709 To generate cross-correlation time-lag map, time-lag of the maximum positive correlation
710 between the CSF signal from the lateral ventricles and all voxels were computed using cross-
711 correlation separately for each nap session. Next, group mean group mean voxel-wise time-lag
712 map was generated by averaging the individual maps.
713
CSF peak locked EEG topography analysis
Positive peaks of the CSF signal from the lateral ventricles during each sleep session were
detected using a MATLAB function (findpeaks.m). The MRI-related artifact-removed EEG data
were segmented based on the positive peak events of the CSF signals (-10 s to 0 s). The
segmented EEG data were averaged across the sleep sessions and participants for each sleep
stage (N2 and N3 sleep stages). EEG topographies across time were generated in the Fieldtrip
toolbox (87) (http://www.ru.nl/neuroimaging/fieldtrip).
721
722 ***Power spectrum analysis of cardiac and respiration signals***
723 Cardiac and respiratory cycles during each sleep session were recorded using a fingertip PPG
724 sensor and respiratory belt (BIOPAC, US) with a 100 Hz sampling rate. The same power
725 spectrum analysis steps were applied to the cardiac and respiration signals (see ***Power***
726 ***spectrum analysis of fMRI data*** above).
727
fMRI GLM analysis
MRI data were processed using FSL v6.0.7.7 (https://fsl.fmrib.ox.ac.uk/fsl/). Data from each
participant were motion corrected (MCFLIRT), spatially smoothed (5 mm FWHM Gaussian
kernel), high-pass temporally filtered (100 s cutoff), registered to their T1 anatomical brain image
(FLIRT), and normalized to the MNI 2 mm standard brain. GLM analyses were performed using
FEAT v6.00. On the basis of each sleep-specific event (onset and duration for slow waves, sleep
spindles, and arousals) detected from the simultaneous EEG data, event-specific first-level
analysis was performed employing one regressor: 1) the event onset and duration. The regressor
of interest was convolved with the double-gamma HRF. Motion parameters estimated during
image realignment (3 translations and 3 rotations around the x, y, and z axes) were included in
the matrix as confounding regressors of no interest. Additionally, since MRI BOLD signal
amplitudes vary across different vigilance states (11, 89–92), the mean MRI signal inside the
white matter region and the mean MRI signal in the cerebrospinal fluid region were included in
the GLM as nuisance regressors (93). For each event of each participant, the event-specific first-
level results were combined across sleep sessions using a second-level, fixed effect analysis to
calculate an average response per subject. For each event, these second-level results were
subsequently combined and analyzed at the group level using a FLAME mixed-effect analysis
(94). In a cluster-based threshold statistical analysis, the resulting group-level BOLD Z statistic
images were thresholded using $Z > 3.1$ and a cluster-corrected significance threshold of $P < 0.05$.
For visualization, the group-level thresholded significant BOLD Z statistic image for each event
(slow waves, sleep spindles, and arousals) was transformed onto a standard template surface
map (fsaverage) using feat2surf and a standard template subcortical brain region, and the group-
level thresholded significant BOLD Z statistic image was shown as an isosurface in 3D view in

FreeSurfer freeview software. The automated anatomical atlas 3 (AAL3 (95)) was used for
anatomical labeling.
753
The control experiment and analyses to test aliasing noises
We conducted the additional experiment using TR=2.5s to check the effect of the aliasing noise
from the respirations and cardiac pulses on the CSF signals (42). If the spectrum peaks of the
CSF signal (around 0.06Hz) were due to the aliasing noise, the 0.06Hz peak should disappear
when the data were acquired with TR=2.5s since the aliasing noise between MRI acquisition rate
(0.4Hz) and respiration peak (around 0.27Hz) would be 0.13Hz (0.4-0.27=0.13Hz) as compared
to 0.06Hz (0.33-0.27=0.06Hz) when using TR=3s (0.33Hz).
761
Three young healthy participants (mean age of 25.3 ± 4.0 years, 3 female) participated in the
control experiment. All of the participants provided written informed consent for their participation
prior to the start of the study. The experimental design and data analyses were the same as the
main experiment except that the TR for MRI acquisition was 2.5s instead of 3s. Time-frequency
analysis was applied to the whole fMRI signals during a sleep session of the two ROIs using a
multitaper approach (see ***Power spectrum analysis of fMRI data*** in **Materials and Methods**).
We confirmed that the major spectrum peak of the CSF signals was still visible at ~0.06Hz
(**Supplementary Fig. S9**). These indicate that this spectrum peak of the CSF signals is not due
to the aliasing noises between the MRI acquisition rate and respiration peak.
771
Statistical analysis
The alpha level was set as 0.05. All tests conducted in this study were two-tailed. For the fMRI
power spectrum analysis, nonparametric cluster-based permutation tests were used to find
significant differences ($P < 0.05$) in the periodic power spectrum across different sleep stages in
each ROI using the Fieldtrip toolbox (96–98). For the fMRI time course analysis, two-tailed one-
sample t-tests ($P <0.05$) were applied to each time point of the fMRI signals compared with zero.
Two-tailed paired t-tests were used to compare each time point of event-locked fMRI signals
between sleep stages. For the statistical inference of the correlation analysis and the numbers
and time intervals of each event across different sleep depths, the Shapiro-Wilk test was
conducted to test whether the data were normally distributed. If data were not normally
distributed, the Wilcoxon signed-rank tests were used. When normality was not rejected by the
Shapiro-Wilk test, two-tailed paired t-tests were used. To control for multiple comparison
problems, Bonferroni corrections were applied to minimize the false-positive rates in the statistical
inference of the correlation analysis and the numbers and time intervals of each event across
different sleep depths. False discovery rate (FDR) corrections (99) were used for other statistical
comparisons, such as comparing each time point of the fMRI signals and the fMRI power
spectrum. We also included a 95% CI for a significant t-test and Pearson's correlation analysis.
Statistical tests were run by MATLAB (2020b, MathWorks, Inc.).
790
791 ***Data availability***
792 The datasets generated and/or analyzed in the current study are available from the
793 corresponding author upon request.
794
795 ***Code availability***
796 No custom algorithm or software was used to that is central to the present findings. The computer
797 codes that were used to generate the results of this study are available from the corresponding
798 author upon request.
799
800 ***Inclusion & Ethics***
801 This study was approved by the institutional review board at RIKEN. All of the participants
802 provided written informed consent for obtaining and sharing individual-level data.
803
804

Acknowledgments

The authors thank M. A. Carskadon and H. Lau for helpful comments on an earlier draft. The authors also thank H. Nakatomi, M. Gomyo, T. Yoshioka, K. Fukushima, M. Koyanagi for fruitful discussions on CSF measurements. This work was supported by JSPS KAKENHI Grant Numbers JP22H01107 (MT) and JP22K18664 (MT), the Naito Foundation (MT), Takeda Science Foundation (MT), Life Science Foundation of JAPAN (MU).

References

1. R. Stickgold, Sleep-dependent memory consolidation. *Nature* 437, 1272–1278 (2005).

2. L. Marshall, H. Helgadóttir, M. Mölle, J. Born, Boosting slow oscillations during sleep potentiates memory. *Nature* 444, 610–613 (2006).

3. A. J. Krause, *et al.*, The sleep-deprived human brain. *Nat Rev Neurosci* 18, 404–418 (2017).

4. M. R. Irwin, M. V Vitiello, Implications of sleep disturbance and inflammation for Alzheimer's disease dementia. *Lancet Neurol* 18, 296–306 (2019).

5. L. Xie, *et al.*, Sleep Drives Metabolite Clearance from the Adult Brain. *Science (1979)* 342, 373–377 (2013).

6. J. J. Iliff, *et al.*, A Paravascular Pathway Facilitates CSF Flow Through the Brain Parenchyma and the Clearance of Interstitial Solutes, Including Amyloid β. *Sci Transl Med* 4 (2012).

7. P. K. Eide, *et al.*, Altered glymphatic enhancement of cerebrospinal fluid tracer in individuals with chronic poor sleep quality. *Journal of Cerebral Blood Flow & Metabolism* 42, 1676–1692 (2022).

8. P. K. Eide, V. Vinje, A. H. Pripp, K.-A. Mardal, G. Ringstad, Sleep deprivation impairs molecular clearance from the human brain. *Brain* 144, 863–874 (2021).

9. N. L. Hauglund, *et al.*, Norepinephrine-mediated slow vasomotion drives glymphatic clearance during sleep. *Cell* 188, 606-622.e17 (2025).

10. T. Warbrick, Simultaneous EEG-fMRI: What Have We Learned and What Does the Future Hold? *Sensors* 22, 2262 (2022).

11. N. E. Fultz, *et al.*, Coupled electrophysiological, hemodynamic, and cerebrospinal fluid oscillations in human sleep. *Science (1979)* 366, 628–631 (2019).

12. V. Kiviniemi, *et al.*, Ultra-fast magnetic resonance encephalography of physiological brain activity – Glymphatic pulsation mechanisms? *Journal of Cerebral Blood Flow & Metabolism* 36, 1033–1045 (2016).

13. H. Helakari, *et al.*, Human NREM Sleep Promotes Brain-Wide Vasomotor and Respiratory Pulsations. *The Journal of Neuroscience* 42, 2503–2515 (2022).

14. B. Battal, *et al.*, Cerebrospinal fluid flow imaging by using phase-contrast MR technique. *Br J Radiol* 84, 758–765 (2011).

15. A. M. Wright, Y. Wu, L. Feng, Q. Wen, Diffusion magnetic resonance imaging of cerebrospinal fluid dynamics: Current techniques and future advancements. *NMR Biomed* 37 (2024).

16. M. de Zambotti, *et al.*, K-Complexes: Interaction between the Central and Autonomic Nervous Systems during Sleep. *Sleep* 39, 1129–1137 (2016).

17. J. Tank, *et al.*, Relationship between blood pressure, sleep K-complexes, and muscle sympathetic nerve activity in humans. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* 285, R208–R214 (2003).

18. A. Lüthi, M. Nedergaard, Anything but small: Microarousals stand at the crossroad between noradrenaline signaling and key sleep functions. *Neuron* 113, 509–523 (2025).

19. A. Rechtschaffen, A. Kales, *A manual of standardized terminology, techniques and scoring system of sleep stages in human subjects* (Public Health Service, US Government Printing Office, 1968).

20. M. Boselli, L. Parrino, A. Smerieri, M. G. Terzano, Effect of Age on EEG Arousals in
Normal Sleep. *Sleep* 21 (1998).
21. M. Uji, M. Tamaki, Sleep, learning, and memory in human research using noninvasive
neuroimaging techniques. *Neurosci Res* 189, 66–74 (2023).
22. M. Tamaki, T. Watanabe, Y. Sasaki, Coregistration of magnetic resonance spectroscopy
and polysomnography for sleep analysis in human subjects. *STAR Protoc* 2, 100974
(2021).
23. M. Tamaki, *et al.*, Complementary contributions of non-REM and REM sleep to visual
learning. *Nat Neurosci* 23, 1150–1156 (2020).
24. M. Tamaki, *et al.*, First-night effect reduces the beneficial effects of sleep on visual
plasticity and modifies the underlying neurochemical processes. *Sci Rep* 14, 14388
(2024).
25. X. Hu, T. H. Le, T. Parrish, P. Erhard, Retrospective estimation and correction of
physiological fluctuation in functional MRI. *Magn Reson Med* 34, 201–212 (1995).
26. G. H. Glover, T.-Q. Li, D. Ress, Image-based method for retrospective correction of
physiological motion effects in fMRI: RETROICOR. *Magn Reson Med* 44, 162–167 (2000).
27. O. C. Reddy, Y. D. van der Werf, The Sleeping Brain: Harnessing the Power of the
Glymphatic System through Lifestyle Choices. *Brain Sci* 10, 868 (2020).
28. T. O. Wichmann, H. H. Damkier, M. Pedersen, A Brief Overview of the Cerebrospinal
Fluid System and Its Implications for Brain and Spinal Cord Diseases. *Front Hum*
Neurosci 15 (2022).
29. R. Berry, S. Quan, A. Abreu, Manual for the Scoring of Sleep and Associated Events:
Rules, Terminology and Technical Specifications. *American Academy of Sleep Medicine*
Version 2.6 (2020).
30. M. Massimini, R. Huber, F. Ferrarelli, S. Hill, G. Tononi, The Sleep Slow Oscillation as a
Traveling Wave. *The Journal of Neuroscience* 24, 6862–6870 (2004).
31. T. T. Dang-Vu, *et al.*, Spontaneous neural activity during human slow wave sleep.
Proceedings of the National Academy of Sciences 105, 15160–15165 (2008).
32. M. Mölle, T. O. Bergmann, L. Marshall, J. Born, Fast and Slow Spindles during the Sleep
Slow Oscillation: Disparate Coalescence and Engagement in Memory Processing. *Sleep*
34, 1411–1421 (2011).
33. V. Latreille, *et al.*, The human K-complex: Insights from combined scalp-intracranial EEG
recordings. *Neuroimage* 213, 116748 (2020).
34. Y. L. Wang, *et al.*, Intracerebral dynamics of sleep arousals: a combined scalp-intracranial
EEG study. *The Journal of Neuroscience* e0617232024 (2024).
https://doi.org/10.1523/JNEUROSCI.0617-23.2024.
35. A. R. Adamantidis, C. Gutierrez Herrera, T. C. Gent, Oscillating circuitries in the sleeping
brain. *Nat Rev Neurosci* 20, 746–762 (2019).
36. R. Boyce, S. D. Glasgow, S. Williams, A. Adamantidis, Causal evidence for the role of
REM sleep theta rhythm in contextual memory consolidation. *Science (1979)* 352, 812–
816 (2016).
37. M. Dresler, *et al.*, Neural Correlates of Dream Lucidity Obtained from Contrasting Lucid
versus Non-Lucid REM Sleep: A Combined EEG/fMRI Case Study. *Sleep* 35, 1017–1020
(2012).
38. A. Elabasy, *et al.*, Sleep increases propagation speed of physiological brain pulsations.
[Preprint] (2025).
39. D. Picchioni, *et al.*, Autonomic arousals contribute to brain fluid pulsations during sleep.
Neuroimage 249, 118888 (2022).
40. S. D. Williams, *et al.*, Neural activity induced by sensory stimulation can drive large-scale
cerebrospinal fluid flow during wakefulness in humans. *PLoS Biol* 21, e3002035 (2023).
41. C. Strik, U. Klose, M. Erb, H. Strik, W. Grodd, Intracranial oscillations of cerebrospinal fluid
and blood flows: Analysis with magnetic resonance imaging. *Journal of Magnetic*
Resonance Imaging 15, 251–258 (2002).

42. L. Raitamaa, *et al.*, Spectral analysis of physiological brain pulsations affecting the BOLD
signal. *Hum Brain Mapp* 42, 4298–4313 (2021).
43. J. Gonzalez-Castillo, I. S. Fernandez, D. A. Handwerker, P. A. Bandettini, Ultra-slow fMRI
fluctuations in the fourth ventricle as a marker of drowsiness. *Neuroimage* 259, 119424
(2022).
44. A. A. Borbely, A two process model of sleep regulation. *Hum Neurobiol* 1 (1982).
45. S. Brodt, M. Inostroza, N. Niethard, J. Born, Sleep—A brain-state serving systems
memory consolidation. *Neuron* 111, 1050–1075 (2023).
46. G. Tononi, C. Cirelli, Sleep and the Price of Plasticity: From Synaptic and Cellular
Homeostasis to Memory Consolidation and Integration. *Neuron* 81, 12–34 (2014).
47. M. Betta, *et al.*, Cortical and subcortical hemodynamic changes during sleep slow waves
in human light sleep. *Neuroimage* 236, 118117 (2021).
48. M. Schabus, *et al.*, Hemodynamic cerebral correlates of sleep spindles during human non-
rapid eye movement sleep. *Proceedings of the National Academy of Sciences* 104,
13164–13169 (2007).
49. D. Baena, *et al.*, Functional differences in cerebral activation between slow wave-coupled
and uncoupled sleep spindles. *Front Neurosci* 16 (2023).
50. R. Wehrle, *et al.*, Functional microstates within human REM sleep: first evidence from
fMRI of a thalamocortical network specific for phasic REM periods. *European Journal of
Neuroscience* 25, 863–871 (2007).
51. T. T. Dang-Vu, *et al.*, Functional Neuroimaging Insights into the Physiology of Human
Sleep. *Sleep* 33, 1589–1603 (2010).
52. S. Miyauchi, M. Misaki, S. Kan, T. Fukunaga, T. Koike, Human brain activity time-locked
to rapid eye movements during REM sleep. *Exp Brain Res* 192, 657–667 (2009).
53. S. Fogel, *et al.*, Reactivation or transformation? Motor memory consolidation associated
with cerebral activation time-locked to sleep spindles. *PLoS One* 12, e0174755 (2017).
54. M. Tamaki, T. Matsuoka, H. Nittono, T. Hori, Fast Sleep Spindle (13–15 Hz) Activity
Correlates with Sleep-Dependent Improvement in Visuomotor Performance. *Sleep* 31,
204–211 (2008).
55. M. Schabus, *et al.*, Interindividual sleep spindle differences and their relation to learning-
related enhancements. *Brain Res* 1191, 127–135 (2008).
56. P. S. Özbay, *et al.*, Sympathetic activity contributes to the fMRI signal. *Commun Biol* 2,
421 (2019).
57. L. Chen, A. Beckett, A. Verma, D. A. Feinberg, Dynamics of respiratory and cardiac CSF
motion revealed with real-time simultaneous multi-slice EPI velocity phase contrast
imaging. *Neuroimage* 122, 281–287 (2015).
58. V. Vijayakrishnan Nair, *et al.*, Human CSF movement influenced by vascular low
frequency oscillations and respiration. *Front Physiol* 13 (2022).
59. I. Slutsky, Linking activity dyshomeostasis and sleep disturbances in Alzheimer disease.
Nat Rev Neurosci 25, 272–284 (2024).
60. Y.-E. S. Ju, *et al.*, Sleep Quality and Preclinical Alzheimer Disease. *JAMA Neurol* 70, 587
(2013).
61. M. V. Vitiello, P. N. Prinz, D. E. Williams, M. S. Frommlet, R. K. Ries, Sleep Disturbances
in Patients With Mild-Stage Alzheimer's Disease. *J Gerontol* 45, M131–M138 (1990).
62. B. A. Mander, J. R. Winer, M. P. Walker, Sleep and Human Aging. *Neuron* 94, 19–36
(2017).
63. G. Liu, A. Ladrón-de-Guevara, Y. Izhiman, M. Nedergaard, T. Du, Measurements of
cerebrospinal fluid production: a review of the limitations and advantages of current
methodologies. *Fluids Barriers CNS* 19, 101 (2022).
64. L. Bojarskaite, *et al.*, Sleep cycle-dependent vascular dynamics in male mice and the
predicted effects on perivascular cerebrospinal fluid flow and solute transport. *Nat
Commun* 14, 953 (2023).
65. L.-F. Jiang-Xie, *et al.*, Neuronal dynamics direct cerebrospinal fluid perfusion and brain
clearance. *Nature* 627, 157–164 (2024).

66.   T. C. Diorio, *et al.*, Real-time quantification of in vivo cerebrospinal fluid velocity using the
966   functional magnetic resonance imaging inflow effect. *NMR Biomed* 37 (2024).
967   67.   V. V. Nair, T. C. Diorio, Q. Wen, V. L. Rayz, Y. Tong, Using respiratory challenges to
968   modulate CSF movement across different physiological pathways: An fMRI study. *Imaging*
969   *Neuroscience* 2, 1–14 (2024).
970   68.   T. Martins, *et al.*, Characterization of pulsations in the brain and cerebrospinal fluid using
971   ultra-high field magnetic resonance imaging. *Front Neurosci* 18 (2024).
972   69.   J. Tuunanen, *et al.*, Cardiovascular and vasomotor pulsations in the brain and periphery
973   during awake and NREM sleep in a multimodal fMRI study. *Front Neurosci* 18 (2024).
974   70.   H. Helakari, *et al.*, Effect of sleep deprivation and NREM sleep stage on physiological
975   brain pulsations. *Front Neurosci* 17 (2023).
976   71.   C. Song, M. Boly, E. Tagliazucchi, H. Laufs, G. Tononi, fMRI spectral signatures of sleep.
977   *Proceedings of the National Academy of Sciences* 119 (2022).
978   72.   D. J. Buysse, C. F. Reynolds, T. H. Monk, S. R. Berman, D. J. Kupfer, The Pittsburgh
979   sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry*
980   *Res* 28, 193–213 (1989).
981   73.   E. Hoddes, V. Zarcone, H. Smythe, R. Phillips, W. C. Dement, Quantification of
982   Sleepiness: A New Approach. *Psychophysiology* 10, 431–436 (1973).
983   74.   M. Uji, R. Wilson, S. T. Francis, K. J. Mullinger, S. D. Mayhew, Exploring the advantages
984   of multiband fMRI with simultaneous EEG to investigate coupling between gamma
985   frequency neural activity and the BOLD response in humans. *Hum Brain Mapp* 39, 1673–
986   1687 (2018).
987   75.   P. J. Allen, O. Josephs, R. Turner, A Method for Removing Imaging Artifact from
988   Continuous EEG Recorded during Functional MRI. *Neuroimage* 12, 230–239 (2000).
989   76.   R. Wilson, K. J. Mullinger, S. T. Francis, S. D. Mayhew, The relationship between negative
990   BOLD responses and ERS and ERD of alpha/beta oscillations in visual and motor cortex.
991   *Neuroimage* 199, 635–650 (2019).
992   77.   M. Bullock, G. D. Jackson, D. F. Abbott, Artifact Reduction in Simultaneous EEG-fMRI: A
993   Systematic Review of Methods and Contemporary Usage. *Front Neurol* 12 (2021).
994   78.   R. Agarwal, T. Takeuchi, S. Laroche, J. Gotman, Detection of Rapid-Eye Movements in
995   Sleep Studies. *IEEE Trans Biomed Eng* 52, 1390–1396 (2005).
996   79.   B. Frauscher, *et al.*, Rapid Eye Movement Sleep Sawtooth Waves Are Associated with
997   Widespread Cortical Activations. *The Journal of Neuroscience* 40, 8900–8912 (2020).
998   80.   L. Peter-Derex, *et al.*, Enhanced thalamocortical functional connectivity during rapid-eye-
999   movement sleep sawtooth waves. *Sleep* 46 (2023).
1000   81.   S. Sato, *et al.*, Relationship between muscle tone changes, sawtooth waves and rapid eye
1001   movements during sleep. *Electroencephalogr Clin Neurophysiol* 103, 627–632 (1997).
1002   82.   B. Fischl, FreeSurfer. *Neuroimage* 62, 774–781 (2012).
1003   83.   B. Fischl, A. M. Dale, Measuring the thickness of the human cerebral cortex from
1004   magnetic resonance images. *Proceedings of the National Academy of Sciences* 97,
1005   11050–11055 (2000).
1006   84.   M. Jenkinson, P. Bannister, M. Brady, S. Smith, Improved Optimization for the Robust and
1007   Accurate Linear Registration and Motion Correction of Brain Images. *Neuroimage* 17,
1008   825–841 (2002).
1009   85.   S. M. Smith, Fast robust automated brain extraction. *Hum Brain Mapp* 17, 143–155
1010   (2002).
1011   86.   R. Scheeringa, *et al.*, Neuronal Dynamics Underlying High- and Low-Frequency EEG
1012   Oscillations Contribute Independently to the Human BOLD Signal. *Neuron* 69, 572–583
1013   (2011).
1014   87.   R. Oostenveld, P. Fries, E. Maris, J.-M. Schoffelen, FieldTrip: Open Source Software for
1015   Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Comput Intell*
1016   *Neurosci* 2011, 1–9 (2011).
1017   88.   T. Donoghue, *et al.*, Parameterizing neural power spectra into periodic and aperiodic
1018   components. *Nat Neurosci* 23, 1655–1665 (2020).

89.   M. Fukunaga, *et al.*, Large-amplitude, spatially correlated fluctuations in BOLD fMRI
signals during extended rest and early sleep stages. *Magn Reson Imaging* 24, 979–992
(2006).
90.   S. G. Horovitz, *et al.*, Low frequency BOLD fluctuations during resting wakefulness and
light sleep: A simultaneous EEG-fMRI study. *Hum Brain Mapp* 29, 671–682 (2008).
91.   S. Olbrich, *et al.*, EEG-vigilance and BOLD effect during simultaneous EEG/fMRI
measurement. *Neuroimage* 45, 319–332 (2009).
92.   L. J. Larson-Prior, *et al.*, Cortical network functional connectivity in the descent to sleep.
Proceedings of the National Academy of Sciences 106, 4489–4494 (2009).
93.   C. Caballero-Gaudes, R. C. Reynolds, Methods for cleaning the BOLD fMRI signal.
Neuroimage 154, 128–149 (2017).
94.   M. W. Woolrich, T. E. J. Behrens, C. F. Beckmann, M. Jenkinson, S. M. Smith, Multilevel
linear modelling for FMRI group analysis using Bayesian inference. *Neuroimage* 21,
1732–1747 (2004).
95.   E. T. Rolls, C.-C. Huang, C.-P. Lin, J. Feng, M. Joliot, Automated anatomical labelling
atlas 3. *Neuroimage* 206, 116189 (2020).
96.   E. Maris, Statistical testing in electrophysiological studies. *Psychophysiology* 49, 549–565
(2012).
97.   E. Maris, R. Oostenveld, Nonparametric statistical testing of EEG- and MEG-data. *J
Neurosci Methods* 164, 177–190 (2007).
98.   T. E. Nichols, A. P. Holmes, Nonparametric permutation tests for functional neuroimaging:
A primer with examples. *Hum Brain Mapp* 15, 1–25 (2002).
99.   C. R. Genovese, N. A. Lazar, T. Nichols, Thresholding of Statistical Maps in Functional
Neuroimaging Using the False Discovery Rate. *Neuroimage* 15, 870–878 (2002).

1043
1044 **Figures and Tables**
1045

1047
Figure 1. Sleep-specific brain oscillations are followed by faster and milder CSF signal changes during slow-wave sleep. a, A schematic of the regions of interest. Blue, lateral ventricles. Red, whole gray matter (GM). **b,** Example MRI signals (top; blue, CSF signals. red, GM signals), corresponding sleep stages (middle) during a 90-min sleep session, and example slow waves and sleep spindles (bottom). Wake, wakefulness. SWS, slow-wave sleep. Light, light NREM sleep. REM, REM sleep. **c,** Intervals of slow waves (left) and sleep spindles (right) during light NREM (green) and slow-wave sleep (blue). Two-tailed Wilcoxson signed-rank tests, *P < 0.05, ****P < 0.001. LN, light NREM sleep. SWS, slow-wave sleep. **d,** CSF signal changes during light NREM sleep time-locked to the onset of slow waves (light green) and sleep spindles (dark green). The horizontal bars (light green, slow waves; dark green, spindles) indicate statistical significance compared with zero baselines via two-tailed one-sample t-tests (FDR corrected, Ps <0.05). **e,** CSF signal changes during slow-wave sleep time-locked to the onset of slow waves (light blue) and sleep spindles (dark blue). The horizontal bars (light blue: slow waves; dark blue: spindles) indicate statistical significance compared with zero baselines via two-tailed one-sample t-tests (FDR corrected Ps < 0.05). **f,** Peak latencies of CSF signals during light NREM (green) and slow-wave sleep (blue) induced by slow waves. Two-tailed Wilcoxson signed-rank tests, ****P < 0.001. **g,** Peak amplitudes of CSF signal changes during light NREM (green) and slow-wave sleep (blue) induced by slow waves. Two-tailed Wilcoxson signed-rank tests, ****P < 0.001. SWS, slow-wave sleep. See **Supplementary Fig. S2** for the relationship between the CSF and GM signals. SWS, slow-wave sleep.
1068
1069

1070
1071
Figure 2. REM sleep events induce changes in CSF dynamics. a, Examples of horizontal
(Hor) and vertical (Ver) rapid eye movements (top) and sawtooth waves detected from the frontal
electrodes (bottom). **b,** Intervals of rapid eye movements (REMs), sawtooth waves (STWs) and
all REM sleep events (All). **c,** CSF signal changes time-locked to the onset of REM sleep events.
The horizontal bars represent statistical significance against zero baselines according to two-
tailed one-sample *t*-tests (FDR-corrected *P* < 0.05). **d,** GM signal changes time-locked to the
onset of REM sleep events. The horizontal bars represent statistical significance against zero
baselines according to two-tailed one-sample *t*-tests (FDR-corrected *P* <0.05). **e**, Cross-
correlation between CSF and GM signal changes time-locked to all REM events including rapid
eye movements and sawtooth waves (max |r| = 0.31 at lag = -6s).
1082
1083

Figure 3. Arousals induce large CSF signal changes during sleep. a, Example of an epoch
containing an arousal event (red). The embedded diagram shows an EEG spectrogram (O1 and
O2 electrodes averaged) during arousal events (number of samples = 50 sleep sessions),
indicating EEG activations across a wide range of frequency bands. **b,** Number of arousals during
light NREM sleep, slow-wave sleep and REM sleep stages. Two-tailed Wilcoxson signed-rank
tests with Bonferroni correction, ****$Ps < 0.001$. **c,** CSF signal changes time-locked to arousal
events for each sleep stage. The horizontal bars indicate significance against zero baselines via
two-tailed one-sample *t*-tests (FDR-corrected $Ps < 0.05$). Because the majority of arousals
occurred from light NREM sleep (as indicated in **b**), the plot for light NREM sleep (light green)
and that for all arousal events (black) overlap. **d,** GM signal changes time-locked to arousal
events. The horizontal bars indicate significance against zero baselines via two-tailed one-sample
t-tests (FDR-corrected $Ps < 0.05$). As in **c**, because the majority of arousals occurred from light
NREM sleep, the plot for light NREM sleep (light green) and that for all arousal events (black)
overlap. **e,** CSF signal changes to arousal (black), slow waves during light NREM sleep (green,
same as **Fig. 1d**), and slow waves during slow-wave sleep (blue, same as **Fig. 1e**). The
horizontal bars indicate significance against zero baselines via two-tailed one-sample *t*-tests
(FDR-corrected $Ps < 0.05$). **f,** GM signal changes to arousal (black), slow waves during light
NREM sleep (green), and slow waves during slow-wave sleep (blue). The horizontal bars indicate
significance against zero baselines via two-tailed one-sample *t*-tests (FDR-corrected $Ps < 0.05$).
g, Peak amplitudes of CSF signals induced by slow waves during light NREM sleep (green),
slow-wave sleep (blue), and arousal (black). Two-tailed Wilcoxson signed-rank tests with

 Bonferroni correction, ****Ps < 0.001). The plots for light NREM and slow-wave sleep are the
1108 same as those in **Fig. 1g**. **h,** Mean latencies to reach the peak CSF signals induced by slow
1109 waves during light NREM sleep (green), slow-wave sleep (blue), and arousal (black). Two-tailed
1110 Wilcoxson signed-rank tests with Bonferroni correction, ****Ps < 0.001. SWS, slow-wave sleep.
1111 The plots for light NREM and slow-wave sleep are the same as those in **Fig. 1f**.
1112
1113

1114
1115
Figure 4. Sensory and motor regions are recruited during light sleep, whereas plasticity
and homeostatic circuits dominate deep sleep. A generalized linear model (GLM) analysis
was performed on the fMRI data corresponding to each event. A cluster-based statistical analysis
was conducted at the group level, with the cluster-defining threshold ($Z > 3.1$) corrected at P
< 0.05. Left, lateral view of the left hemisphere of a cortical surface in an inflated format. Middle,
medial view of the left hemisphere of a cortical surface in an inflated format. Right, subcortical
areas. The red highlighted areas in the first row indicate the areas that significantly increased
during slow waves in light NREM sleep compared with those in slow-wave sleep. The red
highlighted areas in the second row indicate the areas that significantly increased during slow
waves in slow-wave sleep compared with those in light NREM sleep. The red highlighted areas in
the third row indicate the areas that increased significantly with rapid eye movements (REMs)
during REM sleep. The red highlighted areas in the fourth row indicate the areas that increased
significantly in response to arousals. Light NREM sleep induces sensory and motor regions,
whereas slow-wave sleep induces regions involved in plasticity and homeostatic circuits. REM
sleep recruits visual and plasticity circuits. Arousals accompany widespread brain activation.
Details of brain regions recruited for each sleep event can be found in **Supplementary Tables**
S3-9.
1133
1134

1135
1136
1137 **Figure 5. An illustration of the hypothetical relationships between CSF and brain circuits**
1138 **during light and deep human sleep.** During light sleep (left), CSF oscillations are characterized
1139 by intermittent sharp peaks induced by cortical arousals or arousal-related waves (K-complexes).
1140 During deep sleep (right), CSF oscillations become faster, but their amplitudes are low (mild).
1141 These faster CSF oscillations are induced by sleep-related brain oscillations.
1142
1143