1

SARS-CoV-2 nsp16 is regulated by host E3 ubiquitin ligases, UBR5, and MARCHF7

Li Tian[1#], Zongzheng Zhao[2#], Wenying Gao[1], Zirui Liu [2], Xiao Li[2*], Wenyan Zhang[1*] and Zhaolong Li[1*]

[1]Department of Infectious Diseases, Infectious Diseases and Pathogen Biology Center, Institute of Virology and AIDS Research, Key Laboratory of Organ Regeneration and Transplantation of The Ministry of Education, The First Hospital of Jilin University, Changchun 130000, Jilin, China

[2]Research Unit of Key Technologies for Prevention and Control of Virus Zoonoses, Chinese Academy of Medical Sciences, Changchun Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Changchun 130000, Jilin, China

[#]These authors contributed equally to this article.

*Correspondence:

Wenyan Zhang (zhangwenyan@jlu.edu.cn), Xiao Li (skylee6226@163.com), and Zhaolong Li (lizhaolong@jlu.edu.cn)

Competing Interests: The authors declare no conflicts of interest.

Abstract

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the virus responsible for coronavirus disease 2019 (COVID-19), remains a global public health threat with major economic implications. The non-structural protein 16 (nsp16) of SARS-CoV-2, in complex with nsp10, catalyses the final step of viral mRNA capping via its 2'-O-methylase activity, enabling the virus to evade host immunity and protect viral mRNA from degradation. However, host factors regulating nsp16 have not been thoroughly explored. Although various E3 ubiquitin ligases are known to interact with SARS-CoV-2 proteins, their specific roles in targeting nsp16 for degradation remain unclear. In this study, we demonstrate that nsp16 is ubiquitinated and degraded by host E3 ubiquitin ligases UBR5 and MARCHF7, acting through the ubiquitin-proteasome system (UPS). UBR5 and MARCHF7 induce nsp16 degradation via K48- and K27-linked ubiquitination, respectively, in independent processes that inhibit SARS-CoV-2 replication both in vitro and in vivo. Furthermore, UBR5 and MARCHF7 exhibited broad-spectrum antiviral activity by degrading nsp16 variants from different SARS-CoV-2 strains. Our findings uncover novel mechanisms by which the UPS antagonises SARS-CoV-2 replication and provide promising targets for therapeutic intervention against COVID-19.

Key words: SARS-CoV-2, nsp16, ubiquitination, UBR5, MARCHF7, proteasomal degradation

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has emerged as a global public health threat, causing over 700 million coronavirus disease 2019 (COVID-19) cases and 7 million deaths (Carabelli et al., 2023; Markov et al., 2023). SARS-CoV-2 encodes four structural proteins, 16 non-structural proteins (Nsps) that assist in viral replication and transcription, and a series of accessory proteins (ORFs) associated with immune evasion (V'kovski et al., 2021; Wang et al., 2020). Some viruses such as Ebola virus (Valle et al., 2021), dengue virus (Jung et al., 2018), and reovirus (Furuichi et al., 1975) encode 2'-O methyltransferases (2'-O-MTases) homologous to those found in eukaryotes. This allows them to mimic the host 5' cap structure on their RNA, hindering recognition by the innate immune system (Daffis et al., 2010; Züst et al., 2011). Similarly, SARS-CoV-2 nsp16, in complex with nsp10, functions as a 2'-O-MTase, methylating the first nucleotide of a capped viral RNA strand at the 2′-O position, converting the 5' cap of RNA from 'cap-0' to 'cap-1' (Benoni et al., 2021; Park et al., 2022). This modification shields SARS-CoV-2 from host antiviral responses such as MDA5 recognition and IFIT1 restriction (Bergant et al., 2022; Russ et al., 2022). Importantly, nsp16 inhibits global host mRNA splicing, which reduces host protein and mRNA levels (Banerjee et al., 2020). Additionally, nsp16 enhances SARS-CoV-2 cell entry by promoting TMPRSS2 expression (Han et al., 2023), highlighting its multifaceted role as a virulence factor. Moreover, nsp16-deficient strains exhibit reduced pathogenicity and induce a strong immune response, suggesting their potential as a live-attenuated vaccine candidate (Ye et al., 2022). Given nsp16's critical functions, drugs targeting nsp16 or the nsp16-nsp10 complex have been developed (Klima et al., 2022; Nguyen et al., 2022). Therefore, understanding host factors that interact with nsp16 is crucial for uncovering novel therapeutic strategies.

The ubiquitin-proteasome system (UPS) is an important pathway for the targeted degradation of intracellular proteins and is referred to as "the molecular kiss of death" (Dongdem & Wezena, 2021; Park et al., 2020). Substrate ubiquitination requires a cascade of three ligases: E1, E2, and E3 (Zheng & Shabek, 2017). Some E3 ubiquitin

ligases function as monomers, while others form multi-subunit E3 complexes. Regardless of their form, all E3 ligases share the same functional specificity in recognizing substrates and determining the specificity of the UPS. In some complexes, an additional protein may be required as a substrate receptor to recognize the substrate (Iconomou & Saunders, 2016; Yang et al., 2021). E3 ligases are indispensable for direct binding to substrates and determining the specificity of the UPS (Li et al., 2021; Wang et al., 2022). Researchers have identified over 600 human genome-encoded E3 ligases that operate in diverse cellular environments, respond to a wide range of cellular signals, and regulate a vast array of protein substrates (Dongdem & Wezena, 2021; Garcia-Barcena et al., 2020). Dysregulation of E3 ligase activity has been implicated in various human diseases, making them promising drug targets (Humphreys et al., 2021). E3 ligases are classified into three classes based on conserved domains and ubiquitin (Ub) transfer mechanisms: Really Interesting New Gene (RING) E3s, homologous to the E6AP carboxyl terminus (HECT) E3s, and RING-between-RING (RBR) E3s (Garcia-Barcena et al., 2020; Morreale & Walden, 2016). RING E3 ligases directly transfer Ub from the E2-Ub complex to the substrate (Deshaies & Joazeiro, 2009), whereas HECT-type E3s transfer Ub to their own catalytic cystine residue before attaching it to the substrate (Huibregtse et al., 1995). RBR E3s combine the characteristics of both RING and HECT types (Walden & Rittinger, 2018). RBR and RING E3s share the RING-binding domains, but RBR family members can generate thioester intermediates with Ub, as is the case with HECT-type E3s (Garcia-Barcena et al., 2020).

To investigate the relationship between SARS-CoV-2 and host factors, we focused on the interaction between nsp16 and the UPS. In this study, we present a novel regulatory mechanism targeting nsp16 via the host UPS. We demonstrate, for the first time, that nsp16 is ubiquitinated and degraded by the host proteasome. Interestingly, two E3 ligases from distinct families, the RING-type MARCHF7 and the HECT-type UBR5, independently recognise and target nsp16 for degradation in the cytoplasm and nucleus. Since nsp16 possesses 2'-O-MTase activity, its enzymatic function converts the viral RNA 5' cap structure from cap-0 to cap-1. This dual targeting of nsp16 by MARCHF7 and UBR5 leads

to its degradation, disrupting its function and ultimately demonstrating effective antiviral

activity against SARS-CoV-2. Our findings identify novel therapeutic targets for developing

effective strategies to prevent SARS-CoV-2 and treat COVID-19.

Results

Ubiquitination and proteasomal degradation of SARS-CoV-2 nsp16 by the UPS

Ubiquitination plays an important role in SARS-CoV-2 infection and pathogenesis

(Gao et al., 2022; Guo et al., 2021; Li et al., 2023; Zhang et al., 2021; Zhang et al., 2024;

Zhang et al., 2023). To explore whether the non-structural proteins of SARS-CoV-2 are

regulated by the UPS, we examined the effect of the proteasome inhibitor MG132 on the

protein expression of all 16 non-structural proteins. MG132 treatment markedly increased

the abundance of nsp8, nsp11, and nsp16 proteins in HEK293T cells (Fig. 1A), suggesting

their potential degradation by the UPS. It has been reported that nsp8 undergoes

UPS-mediated degradation via TRIM22-mediated ubiquitination (Fan et al., 2024). In this

study, we focused on nsp16 for further investigation. To confirm the role of the UPS in

nsp16 degradation, we employed additional proteasome inhibitors, Bortezomib and

Carfilzomib. Unlike the lysosomal inhibitors Bafilomycin A1 and NH_4Cl and the

autophagy-lysosomal inhibitor vinblastine, these proteasome inhibitors enhanced nsp16

stability (Fig. 1B). The impact of the USP on the half-life of nsp16 was investigated. To this

end, nsp16-expressing cells were treated with a protein synthesis inhibitor, cycloheximide,

and the kinetics of nsp16 decay were examined. We observed that the half-life of nsp16 in

MG132-treated samples was significantly longer than that in MG132-untreated samples

(15 h versus 2 h) (Fig. 1C and 1D).

To identify proteins interacting with nsp16, we performed co-immunoprecipitation

(Co-IP) followed by mass spectrometry (MS) analysis, comparing nsp16-expressing cells

treated with and without MG132 (Fig. 1E). KEGG pathway and Gene Ontology analyses

of the MS data (Figure 1—figure supplement 1A). Biological process enrichment analysis

showed that blocking nsp16 degradation resulted in an increase in the number of proteins

that interact with it, including 43 proteins involved in virus-associated processes and 22

proteins involved in regulating mRNA stability. Five proteins were related to the regulation

of the cellular antiviral defence. Additionally, interactions were associated with mRNA

splicing and RNA methylation (Banerjee et al., 2020), consistent with nsp16's role as a

2'-O-MTase. Furthermore, SARS-CoV-2 uses nsp16 to disrupt host mRNA splicing to

facilitate its infection (Park et al., 2022; Russ et al., 2022; Züst et al., 2011). These findings

corroborate the reliability of our MS data.

As expected, we found that several nsp16-binding proteins were related to

ubiquitination and degradation pathways (Figure 1 — figure supplement 1B). Four

deubiquitinases (DUBs), the E2 ligase UBE2D3, and 14 proteasomal enzymes were

identified. Of these, six E3 ligases, UBR5, MARCHF7, HECTD1, TRIM32, MYCBP2, and

TRIM21, were chosen for further investigation.

E3 ligases UBR5 and MARCHF7 independently mediate nsp16 degradation

To identify the E3 ligases responsible for nsp16 degradation, we designed small

interfering RNA (siRNA) targeting the six candidate E3 ligases and evaluated their effects

on nsp16 abundance. Knockdown of UBR5 and MARCHF7 stabilised the protein levels of

nsp16 (Fig. 2A). Furthermore, we generated stable cell lines with UBR5 or MARCHF7

knockdowns to further investigate their effects on nsp16 stability and confirm silencing

efficiency (Fig. 2B). To determine whether UBR5 and MARCHF7 cooperate in nsp16

degradation, we knocked down UBR5 or MARCHF7 using siRNA in MARCHF7- or

UBR5-knockdown cells, respectively. Knockdown of UBR5 in MARCHF7-knockdown cells,

and vice versa, further enhanced nsp16 stability. This indicates that UBR5 and MARCHF7

act independently to degrade nsp16 (Fig. 2C). We further validated the independent

function of these E3 ligases by examining nsp16 degradation. MARCHF7 overexpression

induced nsp16 degradation in both MARCHF7-knockdown and

MARCHF7/UBR5-double-knockdown cells. Similarly, UBR5 overexpression degraded

nsp16 even in the presence of MARCHF7 knockdown. Knockdown efficiencies were

confirmed for all experiments (Figure 2 — figure supplement 1A and 1B). Notably,

overexpression of wild-type MARCHF7 or UBR5, but not mutants lacking the functional

RING domain (1-542 aa) or the HECT domain inactivated mutant, disrupts the stability of

nsp16 in cells with MARCHF7 or UBR5 knockdown (Fig. 2D and 2E). These findings

demonstrate that UBR5 and MARCHF7 independently ubiquitinate nsp16 via their HECT

or RING domains, targeting it for proteasomal degradation.

165 UBR5 and MARCHF7 mediate distinct Ub linkages on nsp16

We first confirmed the ubiquitination of nsp16, both in the presence and absence of

exogenous Ub (Fig. 3A-B). To further investigate the regulation by E3 ligases, we

examined the impact of UBR5 or MARCH7 knockdown on nsp16 ubiquitination.

Knockdown of either UBR5 or MARCHF7 decreased the ubiquitination level of nsp16

compared to that in the negative control (Fig. 3C). Ub contains seven lysine residues (K6,

K11, K27, K29, K33, K49, and K63) that form polyubiquitin chains with distinct functions,

ultimately dictating the fate of the attached target protein (Grice & Nathan, 2016). To

determine the specific type of polyubiquitin chain modification of nsp16 mediated by UBR5

and MARCH7, we constructed a series of Ub mutants retaining only a single lysine

residue. Except for K33, all single-lysine Ub mutants promoted nsp16 ubiquitylation to

varying extents, suggesting a complex polyubiquitin chain structure on nsp16 potentially

regulated by multiple E3 ligases or E2-E3 ligase pairs (Fig. 3D). Notably, analysis of

specific lysine linkages revealed that MARCHF7 knockdown primarily reduced the

K27-linked ubiquitination of nsp16, while UBR5 knockdown primarily reduced the

K48-linked ubiquitination of nsp16 (Fig. 3E and 3F). These findings indicate that

MARCHF7 and UBR5 induce K27- and K48-linked ubiquitination on nsp16, respectively.

182 UBR5 and MARCHF7 directly interact and colocalise with nsp16 in the endoplasmic

183 reticulum (ER)

MS analysis suggested interactions between nsp16 and both UBR5 and MARCH7.

Co-IP experiments confirmed that both Myc-tagged MARCHF7 and endogenous UBR5

interact with nsp16 (Figure 4 — figure supplement 1A and 1B). To determine whether

MARCHF7 or UBR5 directly interact with nsp16, fluorescence resonance energy transfer

(FRET) assays were performed. When nsp16-YFP was bleached, the fluorescence

signals from CFP-UBR5 or CFP-MARCHF7 fusion proteins increased, indicating a direct

interaction between these E3 ligases and nsp16 (Figure 4—figure supplement 1C). Image

J software was used to quantify the relative fluorescence intensities (Figure 4—figure

supplement 1D). We next investigated whether UBR5 and MARCHF7 require each other

for nps16 binding. We compared the binding ability of UBR5 to nsp16 in the presence or

absence of MARCHF7 using Co-IP. MARCHF7 knockdown had no effect on the

interaction between UBR5 and nsp16, and vice versa (Fig. 4A and 4B), indicating

independent binding of UBR5 and MARCHF7 to nsp16. Previous studies have shown that

nsp16 is localised to both the nucleus and cytoplasm (Zhang et al., 2020). UBR5

(containing two nuclear localisation signals) and MARCHF7 are present in both

compartments (Muñoz-Escobar et al., 2015; Shearer et al., 2018) (Nathan et al., 2008).

Immunofluorescence staining revealed that UBR5 and MARCHF7 colocalised primarily

with nsp16 in the cytoplasm, with some colocalisation in the nucleus of Hela cells (Figure

4—figure supplement 1E). Similar results were observed in nsp16-transfected HEK293T

cells using antibodies against endogenous UBR5 or MARCHF7 (Figure 4 — figure

supplement 1F).

To identify the specific cellular compartment where UBR5 or MARCH7 interacts with

nsp16, we co-transfected cells with UBR5-CFP or MARCH7-CFP along with nsp16-YFP.

Immunostaining was performed with organelle-specific antibodies to label the

mitochondrial marker COX5A, the ER marker PDI, and the Golgi apparatus marker

GM130. Notably, both UBR5 and MARCHF7 interacted with nsp16 and colocalised with

PDI, but not with COX5A or GM130 (Fig. 4C and 4D). These findings indicate that UBR5

and MARCHF7 directly interact with nsp16 in the ER.

Functional domains of UBR5 and MARCHF7 are required for nsp16 interaction and

ubiquitination

UBR5 is a four-domain E3 ligase containing two nuclear localisation signals. The four

domains are UBA, UBR, PABC, and HECT (Muñoz-Escobar et al., 2015) (Figure 4—figure

supplement 2A). Notably, the HECT domain is essential for its E3 ligase activity, where

UBR5 must be conjugated to Ub before transferring it to the substrate (Kim et al., 2021).

To identify the UBR5 domain responsible for nsp16 ubiquitination, we utilised UBR5

mutants with inactivated individual domains. Only the HECT domain mutant failed to degrade nsp16 (Figure 4—figure supplement 2B), consistent with previous findings (Zhou et al., 2022), and was unable to induce nsp16 ubiquitination, further supporting its essential function (Figure 4—figure supplement 2C).

To identify the MARCHF7 region responsible for nsp16 degradation, we constructed a series of truncation mutants, as previously described (Figure 4—figure supplement 2D) (Zhao et al., 2018) (Nathan et al., 2008). All mutants lost the ability to degrade nsp16 (Figure 4—figure supplement 2E). Furthermore, we explored the ability of these mutants to interact with nsp16 and their impact on nsp16 ubiquitination. Only the mutant with an intact N-terminal region (aa 1–542) retained strong binding to nsp16, while mutants with the active RING domain region (aa 543–616) did not. Consistently, only the wild-type MARCHF7 induced the formation of K27-linked Ub chains on nsp16 (Figure 4—figure supplement 2F). These findings suggest that the N-terminal of MARCH7 mediates binding to nsp16, while the RING domain is required for its K27-linked ubiquitination activity.

UBR5 and MARCHF7 mediate antiviral activity against SARS-CoV-2 by targeting nsp16 for degradation

To examine the functional impact of nsp16 degradation by UBR5 and MARCHF7 on viral replication, we used a biosafety level 2 cell culture system to generate SARS-CoV-2 transmissible virus-like particles capable of infecting and replicating in Caco2-N^{int} cells (Ju et al., 2021). Knockdown of either UBR5 or MARCHF7 in these cells significantly increased SARS-CoV-2 replication compared with that in the control group. Notably, simultaneous knockdown of both UBR5 and MARCHF7 further increased SARS-CoV-2 replication (Figure 5—figure supplement 1A and 1B).

Given the critical antiviral roles of UBR5 and MARCH7, we investigated the modification of nsp16 following viral infection. HEK293T cells stably expressing the SARS-CoV-2 entry receptor ACE2 (HEK293T-ACE2) were infected with the Wuhan strain for 48 h. Co-IP using an anti-nsp16 antibody revealed that nsp16 interacts with

endogenous MARCHF7 and UBR5 and undergoes ubiquitination (Fig. 5A). We further

validated the antiviral effects of UBR5 and MARCH7 in a biosafety level 3 facility using the

Wuhan strain at a multiplicity of infection (MOI) of 0.01 and the Omicron strain at an MOI

of 0.001. Compared to the control group, UBR5 or MARCH7 knockdown in Caco2 cells

significantly increased intracellular and secreted viral mRNA levels (M and E genes) of

both strains. Additionally, with the Wuhan strain, MARCH7 knockdown resulted in a

0.6-log increase in viral titre in the supernatants, while UBR5 knockdown caused an over

1-log increase. With the Omicron strain, knockdown of the two E3 ligases resulted in an

even greater increase in viral titres (Fig. 5C–E for the Wuhan strain and 5F–H for the

Omicron strain). Immunoblotting (IB) confirmed these findings, demonstrating elevated

intracellular and secreted N protein levels upon UBR5 or MARCH7 knockdown (Fig. 5I).

The knockdown efficiency of MARCHF7 and UBR5 was also confirmed (Fig. 5B). Due to

low transfection efficiency in Caco2 cells, we overexpressed UBR5-Myc or MARCH7-Myc

in HEK293T-ACE2 cells. Overexpression of UBR5 or MARCH7 significantly decreased

viral mRNA levels of the M and E genes in both the Wuhan and Omicron strains, as

measured by real-time quantitative polymerase chain reaction (RT-qPCR). IB analysis

also showed decreased N protein levels in cells and supernatants, accompanied by a

decrease in viral titres of more than 0.5-log. However, co-transfection with increasing

amounts of nsp16 significantly abrogated the inhibitory effects of UBR5 and MARCH7 on

SARS-CoV-2 replication (Fig. 6A–H and Figure 6—figure supplement 1A–H). To eliminate

the potential influence of nsp16's intrinsic function on experimental outcomes, we

overexpressed UBR5 with an inactive HECT domain (UBR5-ΔHECT) or MARCHF7 with a

RING domain deletion (MARCHF7-1-542), along with nsp16. These E3 ligase mutants

failed to inhibit viral replication. Moreover, a gradual enhancement of nsp16 protein

expression did not enhance viral replication. While there was a slight increase in the

mRNA levels of M, protein levels of E and N remained unchanged (Figure 6—figure

supplement 2A–H). These results suggest that UBR5 and MARCH7 exhibit antiviral

activity against SAR-CoV-2 of UBR5 and MARCHF7 through their enzymatic function,

significantly suppressing viral replication by targeting nsp16 for degradation.

UBR5 and MARCHF7 mediate broad-spectrum degradation of nsp16 variants

Given the ongoing emergence of SARS-CoV-2 variants, broad-spectrum antiviral

activity remains critical. To investigate whether UBR5 and MARCH7 can mediate

broad-spectrum degradation of various nsp16 variants, we aligned the amino acid

sequences of nsp16 proteins obtained from the National Center for Biotechnology

Information (Figure 6 — figure supplement 3A). Among these, only three variants,

XBB.1.9.1, XBB.1.5, and XBB.1.16, exhibited a higher number of mutation sites compared

to other variants, which showed only one or two mutations, suggesting a more conserved

sequence. Consequently, using mutagenesis techniques, we synthesised the nsp16

sequences for these three variants along with several single-site mutants from other

variants. Treatment with the proteasome inhibitor MG132 restored nsp16 protein levels for

all variants (Figure 6—figure supplement 3B), confirming that degradation occurs through

the UPS. Furthermore, knockdown of either UBR5 or MARCHF7 increased the stability of

nsp16 proteins from these variants, albeit to varying degrees, indicating that these nsp16

proteins are sensitive to UBR5- or MARCHF7-mediated degradation (Figure 6—figure

supplement 3C). Taken together, these findings suggest that UBR5 and MARCHF7 may

confer broad-spectrum antiviral activity by targeting nsp16 from diverse SARS-CoV-2 for

degradation.

SARS-CoV-2 infection reduces UBR5 and MARCHF7 expression

To explore the relationship between SARS-CoV-2 infection and the expression levels

of UBR5 and MARCHF7, we analysed changes in their mRNA and protein levels after

infection with either the Wuhan or Omicron strains at varying MOIs, using RT-qPCR and

IB, respectively. Both mRNA and protein levels of MARCFH7 decreased with increasing

viral titres. However, UBR5 expression initially increased at low titres but continuously

declined with increasing titres (Figure 6 — figure supplement 4A-4C). Previous studies

have reported a role for UBR5 in regulating interferon-γ-mediated pathways, which might

explain the observed increase in UBR5 expression at low titres (Wu et al., 2022). To confirm these findings *in vivo*, we examined UBR5 and MARCHF7 mRNA levels in peripheral blood mononuclear cells from SARS-CoV-2-infected patients with varying disease severity. While UBR5 mRNA levels negatively correlated with disease progression, MARCHF7 levels showed no significant correlation (Figure 6 — figure supplement 4D). These results suggest that SARS-CoV-2 may suppress host antiviral defences mediated by UBR5 and MARCHF7.

UBR5 and MARCHF7 protect mice from SARS-CoV-2 challenge

Given the lack of known activators or inhibitors for E3 ligases, we developed a transient overexpression method for UBR5 and MARCH7 in mice using high-pressure tail vein injection of plasmids (Bonamassa et al., 2011). To assess their *in vivo* antiviral effects, mice were injected with plasmids encoding UBR5 or MARCH7, followed by SARS-CoV-2 challenge (Fig. 7A). Both MARCHF7 and UBR5 overexpression significantly decreased viral *E* gene copy numbers in mouse lungs compared to those in the control group, as well as markedly decreased viral titres in lung tissue homogenates (Fig. 7C-D). Additionally, treated mice experienced less weight loss (Fig. 7E). Histopathological analysis of major organs 5 d post-infection revealed that lungs from mice treated with UBR5 or MARCH7 displayed attenuated lesions, including alveolar contraction and pulmonary oedema, compared to those typically observed in control mice (Fig. 7F). These treatments also reduced the abundance of N proteins (Fig. 7G). Collectively, these results suggest that UBR5 or MARCH7 overexpression inhibits SARS-CoV-2 virulence *in vivo* by promoting nsp16 protein degradation.

SARS-CoV-2 infection can trigger a severe cytokine storm, thought to be responsible for its high mortality rates (Song et al., 2020) (Hojyo et al., 2020). To examine the effect of UBR5 and MARCH7 on inflammatory responses, we measured levels of key cytokines, including interleukin-6, interleukin-1 receptor antagonist, and interleukin-1β, in the spleens of SARS-CoV-2-infected mice (Makaremi et al., 2022) (Hu et al., 2021). Consistent with their antiviral activity, treatment with UBR5 or MARCH7 significantly reduced the

329   production of these cytokines (Fig. 7H).

Discussion

SARS-CoV-2 nsp16 functions as a 2'O-MTase, catalysing the methylation of the

penultimate nucleotide of the viral RNA cap. This modification produces a 5'-RNA cap

structure that mimics host RNA, thereby evading host immune detection and responses

(Lin et al., 2020) (Balieiro et al., 2022; Russ et al., 2022). High-resolution structures of

nsp16 and nsp16-nsp10 heterodimers have been characterised (Rosas-Lemus et al.,

2020) (Klima et al., 2022; Lugari et al., 2010), paving the way for antiviral drug

development targeting these heterodimers (Balieiro et al., 2022; Melo-Filho et al., 2022).

However, the host factors that interact with and regulate nsp16 remain largely unknown.

Previously, our group and others have demonstrated the importance of the UPS in

modulating SARS-CoV-2 infection (Xu et al., 2022). For example, host E3 ligases, such as

RNF5, Cullin4-DDB1-PRPF19, ZNF598, and TRIM7, ubiquitinate and degrade key

SARS-CoV-2 proteins, thereby suppressing viral replication (Li et al., 2023; Liang et al.,

2022; Maimaitiyiming et al., 2022; Zhang et al., 2024). Conversely, SARS-CoV-2 employs

DUBs to counteract these host defences and enhance its replication (Chen et al., 2024;

Gao et al., 2024; Gao et al., 2022; Guo et al., 2021). In this study, we identify nsp16 as a

short-lived protein with a half-life of only 2–3 h that is targeted for ubiquitination and

degradation by the UPS. Using MS and bioinformatics screening, we identified two E3

ligases, UBR5 and MARCHF7, that directly interact with and target nsp16 for

ubiquitination and degradation.

UBR5 is a member of the UBR box protein family and contains a HECT domain

essential for its E3 ligase activity (Kim et al., 2021). It is known to regulate tumour growth,

metastasis (Xiang et al., 2022) (Qiao et al., 2020), and viral infections. For example, URB5

targets Middle East respiratory syndrome coronavirus ORF4b protein for degradation,

dampening the host immune response (Zhou et al., 2022). Conversely, UBR5 acts as a

cofactor for Zika virus replication by helping TER94/VCP degrade the capsid protein to expose the viral genome to the cytoplasm (Gestuveo et al.). MARCHF7 is a RING E3 Ub ligase that regulates T cell proliferation, neuronal development, inflammasomes, and tumour progression (Zheng, 2021) (Zhang et al., 2016) (Cai et al., 2022) (Zhao et al., 2018). However, its role in viral infections has been largely unexplored, with only one study identifying it as a host protein interacting with the 2C of foot-and-mouth disease virus (Mahajan et al., 2021).

Our findings reveal that UBR5 and MARCHF7 independently interact with nsp16 and facilitate its degradation, primarily within the cytoplasmic side of the ER. This reflects an evolutionary arms race between viruses, which develop mechanisms to resist host antiviral responses, and hosts, which deploy multiple strategies to counteract the virus. One such strategy is seen with multiple E3 ligases that target the same viral protein for degradation. Functional analyses demonstrate that UBR5 or MARCHF7 knockdown enhanced SARS-CoV-2 replication, whereas their overexpression impaired replication. Furthermore, we found that when the enzymatic activity of these E3 ligases is inactivated, their antiviral effect is abolished, indicating that their antiviral effects are dependent on their E3 ligase activity. Additionally, we observed that in the absence of functional UBR5 or MARCH7, overexpression of nsp16 does not increase viral levels. This suggests that although nsp16 plays an important role, it cannot independently promote viral replication without the involvement of E3 ligases. This underscores the critical function of UBR5 and MARCHF7 as antiviral factors that suppress SARS-CoV-2 replication by promoting nsp16 degradation.

Notably, while nsp16 possesses 18 lysine residues, identifying specific ubiquitination sites proved challenging. Based on structural data of nsp16 (Decroly et al., 2011), we constructed truncated nsp16 mutants to assess their degradation profiles. We found that nsp16 could still be degraded to varying degrees (Appendix-figure S2A-B), suggesting that multiple lysine residues serve as recognition sites for E3 ligases. Nonetheless, the results from these truncated mutants regarding the MG132 recovery provided us with valuable insights. For instance, the Δ2-17 mutant, which lacks this segment, showed more

than a two-fold increase in nsp16 expression levels. Conversely, the Δ204-212 mutant exhibited significantly reduced recovery after MG132 treatment compared to the wild-type. These observations offer important clues for further investigation into the interaction regions between nsp16 and MARCHF7 or UBR5, as well as potential ubiquitination sites. MS analysis identified K76 of nsp16 as a ubiquitination site (Appendix-figure S2C). However, an nsp16 K76R mutant was still degraded by E3 ligases, although the degree of ubiquitination was reduced compared to that of the wild-type, indicating that K76 might be one of several potential ubiquitination sites (Appendix-figure S2D-E). Therefore, we hypothesise that nsp16 undergoes ubiquitination at multiple sites and may involve non-canonical ubiquitination pathways at non-lysine residues (McClellan et al., 2019). A comprehensive understanding of the ubiquitination landscape of nsp16 requires further investigation. In addition to UBR5 and MARCHF7, other E3 ligases could contribute to nsp16 regulation, which warrants further studies.

Currently, no specific activators or inhibitors exist for UBR5 or MARCH7. To validate their antiviral effects *in vivo*, we employed high-pressure tail vein injection, a method commonly used for gene delivery to the liver (Kamimura et al., 2014). This method has also been shown to facilitate gene delivery to other tissues, including the lungs (Bonamassa et al., 2011). Gene expression was successfully detected in lung tissues. While this method induced mild immune activation, its effect is relatively minimal compared with that of other techniques (Suda et al., 2023). Moreover, studies suggest that mice and non-human primates exhibit higher resistance to innate immune activation than humans, suggesting that our findings were only minimally influenced by immune activation (Raper et al., 2003). However, this highlights that translating new drug targets into therapeutic treatments will require considerable time and effort. For example, human trials involving the injection of naked DNA (e.g., OTC cDNA) have shown variable outcomes. While no severe immune responses were observed in most participants, one case of fatal immune activation was reported (Raper et al., 2003), highlighting the variability of human immune responses and a key limitation in extrapolating findings from animal models to human trials. This variability might also explain why we did not observe a clear trend for

MARCHF7 mRNA levels in peripheral blood samples from patients with varying disease

severities. The limited number of patients in our study could further contribute to this

observation.

In this study, we identified the host E3 ligases, UBR5 and MARCHF7, as critical regulators

of SARS-CoV-2 nsp16 through ubiquitination and subsequent degradation. We

demonstrated a negative correlation between the expression levels of these ligases and

the degree of infection, underscoring their antiviral roles. Furthermore, both ligases

significantly inhibit SARS-CoV-2 replication. Our findings pave the way for the

development of novel therapeutic strategies targeting the UPS for the treatment of

COVID-19.

Materials and Methods

Reagents and antibodies

The drugs used in this study were as follows: MG132 (catalog no. S2619),

Bafilomycin A1 (catalog no. S1413), Bortezomib (catalog no. S1013), Carfilzomib (catalog

no. S2853) and Vinblastine (catalog no. S4504) were purchased from Selleck (Houston,

TX, USA). Cycloheximide (catalog no. 66-81-9) was purchased from Sigma (Saint Louis,

MO, USA). The antibodies used for immunoblotting analysis (IB) were as follows:

anti-UBR5 mAb (Proteintech, Rosemont, IL, USA, catalog no. 66937-1-Ig),

anti-MARCHF7 mAb (Santa Cruz Biotechnology, Dallas, TX , USA, catalog no.

sc-166945), SARS-CoV-2 2'-O-ribose Methytransferase Antibody (nsp16) (Cell signaling,

Danvers, Massachusetts, USA, catalog no. #70811), anti-Flag mAb (Sigma, catalog no.

F1804), anti-tubulin mAb (Abcam, Cambridge, Cambridgeshire, UK, catalog no. ab11323),

anti-hemagglutinin (anti-HA) pAb (Invitrogen, catalog no. 71-5500), anti-Myc pAb

(Proteintech, Rosemont, IL, USA, catalog no. 16286-1-AP), and anti-GFP mAb (Abcam,

catalog no. ab1218). Primary antibody: anti-COX5A pAb (Sangon Biotec, Shanghai, CHN,

catalog no. D261450), anti-PDI-mAb (Proteintech, catalog no. 2E6A11) and Human

GM130/GOLGA2 Antibody (R&D Systems, Minneapolis, MN, USA, catalog no. AF8199)

were used for immunofluorescence, and SARS-CoV-2 nucleocapsid monoclonal antibody

(mAb) (GeneTex, Irvine, CA, USA, catalog no. GTX635679) were used for

Immunohistochemistry. Fluorescent secondary antibodies: goat anti-Rabbit IgG (H+L)

Highly Cross Adsorbed Secondary Antibody, Alexa Fluor Plus 488 (Invitrogen, catalog no.

A11001), and goat anti-Rabbit IgG (H+L) Highly Cross Adsorbed Secondary Antibody,

Alexa Fluor Plus 568 (catalog no. A-11011) and immunohistochemical secondary

antibodies: Streptavidin-Peroxidase Anti-Rabbit IgG kit (catalog no. KIT-9706) were

purchased from Invitrogen and maixin (Fuzhou, CHN) respectively.

Cell lines and viruses

The cell lines used in this study include HEK293T cells (catalog no. CRL-11268), Hela

cells (catalog no.CRM-CCL-2), Caco2 cells (catalog no. HTB-37). All cell lines were

purchased from American Type Culture Collection (ATCC; Manassas, VA, USA) and

tested for mycoplasma. In order to conduct virus related experiments, stable cell lines

including Caco2-N^{int} (stably expressing SARS-CoV-2 N gene) and 293T-ACE2 (stably

expressing ACE2) were generated in our laboratory. The cells were cultured in a cell

incubator containing 5% CO_2 at 37°C using Dulbecco's modified Eagle's medium (DMEM;

HyClone, Logan, UT, USA) containing 10% heat-inactivated fetal calf serum (FCS, GIBCO

BRL, Grand Island, NY, USA), 100 mg/ml penicillin, and 100 ug/ml streptomycin to provide

nutrition. We expanded SARS-CoV-2 virus-like-particles (Wuhan-Hu-1, MN908947,

Clade:19A) with high replication capacity in Caco2-N^{int} using a biosafety level-2 (BSL-2)

cell culture system. The virus infecting the cells include Omicron BA. 1 strain

(human/CHN_CVRI-01/2022) and SARS-CoV-2 Wuhan strain

(BetaCoV/Beijing/IME-BJ05-2020, GenBank access no. MT291831.1), and mouse

adapted SARS-CoV-2/C57MA14 variant (GenBank: OL913104.1) used for experiments in

mice were provided by Key Laboratory of Jilin Province for Zoonosis Prevention and

Control, and all experiments for virus infection were performed in Biosafety level 3 (BSL-3)

cell culture system.

Plasmids

Eukaryotic expression plasmids encoding SARS-CoV-2 nsp protein were provided by

Professor Wang Peihui (Shandong University). The nsp16-HA expression vector was

constructed by adding HA tag at the C-terminus. UBR5 (Gnen ID: 51366) and its mutants

(UBR5-ΔHECT, UBR5-ΔPABC, UBR5-ΔUBR) expression plasmids were constructed

using purchased plasmids from Addgene (Watertown, MA, USA) as templates with no tag

or a MYC tag at the N-terminus. The cDNA of 293T cells was used as a template to

construct MARCHF7(Gene ID: 64844) and its truncated mutant with a Myc tag at the

C-terminus. All the above expression plasmids were inserted into the VR1012 vector. For

immunofluorescence (IF) and Fluorescence Resonance Energy Transfer (FRET)

experiments, pCDNA3.1-YFP vector (catalog no. 13033) and pECFP-C1 vector (catalog

no. 6076-1) were purchased from Addgene and BD (Biosciences Clontech), and

MARCHF7 or UBR5 was constructed on pECFP-C1, and nsp16 was constructed on

pCDNA3.1-YFP. Single point mutants of nsp16 of different virus subtypes were obtained

by point mutagenesis. Multi-site combined mutants were synthesized by Sangon Biotec

Company. Human ubiquitin protein and its mutants have been previously described.

Primers required for PCR were listed in Appendix Table S1.

RNA extraction and real-time quantitative PCR

We used the way of Trizol (Invitrogen, catalog no. 15596018CN) to extract RNA. The

RNA was subsequently reverse transcribed using a High-Capacity cDNA Reverse

Transcription kit (Applied Biosystems, Carlsbad, CA, USA, catalog no. 4368814). Finally,

PrimeScript RT Master Mix (Takara, Shiga, JPN, catalog no. RR036A) and relative

real-time primer were used for qPCR on an Mx3005P instrument (Agilent Technologies,

Stratagene, La Jolla, CA, USA). Amplification procedure of the target fragment is as

follows: predenaturation at 95°C for 2min, denaturation at 95 °C for 30s, annealing at

55 °C for 30s, and extension at 72 °C for 30s, total of 40 cycles. Real-time primer

sequences used in this study were shown in appendix table S1.

Immunoblotting analysis (IB)

Cells or supernatant cultured for a certain time were collected, cell precipitates were

resuspended with lysis buffer (50 mM Tris–HCl [pH 7.8], 150 mM NaCl, 1.0% NP-40, 5%

glycerol and 4 mM EDTA), 4xloading buffer was added (0.08M Tris [pH 6.8], 2.0%SDS,

10% glycerol, 0.1 M dithiothreitol, and 0.2% bromophenol blue), and samples were lysed

by heating at 100℃ for 30min. After removal of cell debris by centrifugation, the

supernatant was taken to separate proteins of different sizes by SDS-PAGE. The proteins

were transferred onto polyvinylidene fluoride (PVDF) membranes, incubated overnight

with the indicated primary antibody, and after 1 hour at room temperature with

HRP-conjugated secondary antibodies (Jackson Immunoresearch, West Grove, NJ, USA,

catalog no. 115-035-062 for anti-mouse and 111-035-045 for anti-rabbit), the proteins

were visualized through Ultrasensitive ECL Chemiluminescence Detection Kit

(Proteintech, catalog no. B500024).

Stable cell line generation

We purchased the lentiviral vector pLKO.1-puro (catalog no. 8453) from Adggen.

ShRNA targeting target genes were designed and synthesized by Sangon Biotech. After

annealing, it was inserted into the vector. Lentivirus was generated by transfection with

shRNA (cloned in pLKO.1) or control vector, RRE, VSVG and REV, and the cells were

screened with 1/1000 puromycin (3 μg/ml, Sigma, catalog no. P8833) after infection 48

hours. The knockdown efficiency was detected by RT-qPCR or IB. The shRNA target

sequences used in this study are shown in appendix table S1.

RNAi

Short interfering RNA (siRNA) used in this study were purchased from RiboBio Co.

Ltd. (Guangzhou, CHN). The siRNA sequences for knockdown of MARCHF7 or UBR5 are

provided in appendix table S1. The siRNA was transfected into the cells by transfection

reagent Lipofectamine 2000 (Invitrogen, catalog no. 11668-019), and the corresponding

plasmids were transfected 24 hours later using Lipofectamine 3000 Reagent (Invitrogen,

Carlsbad, CA, USA, catalog no. L3000-008). The expression of target genes was detected

by RT-qPCR or immunoblotting (IB) analysis 48 hours later.

Immunoprecipitation

After 10 hours of MG132 (10 µM) treatment, the cells were harvested, resuspended

in 1ml lysis buffer (50 mM Tris-HCl [pH 7.5], 150 mM NaCl, 0.5% NP-40) containing

protease inhibitor (Roche, catalog no. 11697498001), placed on the shaker oscillator for 4

hours to lyse, then centrifuged to remove cell debris, and the supernatant was incubated

with the antibody and protein G agarose beads (Roche, Basel, Basel-City, CH, catalog no. 11243233001) at 4°C overnight. Protein G agarose beads was washed 6-8 times with wash buffer (20 mM Tris-HCL [pH 7.5], 100 mM NaCl, 0.1 mM EDTA, and 0.05% Tween 20) at 4°C, centrifuged at 800 g for 1 min. The proteins were eluted by adding loading buffer (0.08 M Tris [pH 6.8], 2.0% SDS, 10% glycerol, 0.1 M dithiothreitol, and 0.2% bromophenol blue), the samples were boiled at 100°C for 10min to elate the proteins. The lysates and immunoprecipitations were detected by IB.

Mass spectrometry (MS)

HEK293T were transfected with SARS-CoV-2-nsp16-Flag for 36 hours, and then harvested after 10 hours of treatment with MG132 or DMSO. Proteins were enriched by co-immunoprecipitation assay. And the elutions were analyzed by MS. MS analysis was performed by the national center for protein science (Beijing, CHN).

Immunofluorescence

At 48 hours after transfection, the solution was discarded and cells were washed twice with preheated PBS for 5min each, followed by fixation with 4% paraformaldehyde at 37°C for 10min. After three washes with PBS, permeabilized with 0.2% Triton X-100 for 5 min at 37°C, and then immediately followed by blocking with 10% fetal bovine serum at room temperature for 1 hour. The blocked cells were incubated overnight with the indicated primary antibodies. The next day, after three washes with PBS, the cells were incubated with corresponding secondary antibodies for 1 hour at room temperature in the dark. The nuclei were stained with DAPI (49,6-diamidino-2-phenylindole, Sigma, catalog no. 9542) and then stored in 90% glycerol. The fluorescence was detected by a laser scanning confocal microscope (FV 3000, Olympus, Tokyo, Japan).

Viral infectivity assay

The Caco2 cells were transfected with siRNA by Lipofectamine RNAiMAX Reagent (Invitrogen, catalog no. 13778150) to knockdown UBR5 or MARCHF7 and infected with Wuhan strain (MOI=0.01) or Omicron-BA.1 strain (MOI=0.001) at 24 hours after transfection. After two hours of infection, the Caco2 cells were cultured in fresh medium containing 2% fetal bovine serum for 48 hours. The cells and supernatants were collected.

Viral levels were characterized by RT-qPCR and IB analysis of viral structural proteins. The overexpression plasmid UBR5-Myc, MARCHF7-Myc and nsp16-Flag was transfected in 293T-ACE2 by Lipofectamine 3000 Reagent and infected with virus 24 hours later.

Mouse lines and infections

BALB/C mice, 6 weeks, were purchased from Charles River Laboratories, Beijing. Mice were cultured in groups according to experimental groups. Our study examined male and female animals, and similar findings are reported for both sexes. To reduce animal suffering, all welfare and experimental procedures were carried out in strict accordance with the Guide for the Care and Use of Laboratory Animals and relevant ethical regulations during the experiment. The mice were randomly divided into 6 groups and injected with 40 μg/500 μl MARCHF7 or UBR5 plasmids via the tail vein at high pressure. Three groups of mice were infected 50ul with SARS-CoV-2 isolates by nasal challenge at a dose of $10^{5.5}$ $TCID_{50}$/mL, three groups were not infected with virus as control, and the empty vector injection group was also used as control.

Immunohistochemical analysis

Three mice in each group, a total of 18 mice, were sacrificed after anesthesia. The lungs of mice were collected and fixed in 4% paraformaldehyde fixative. After 2 days, the lungs were progressively dehydrated through different concentrations of ethanol solution, transparently treated, soaked in xylene, and finally embedded in paraffin. Some thin sections obtained through a microtome for hematoxylin and eosin (H&E) staining. The other fraction was incubated with 3% hydrogen peroxide for 5-10 min at room temperature to eliminate endogenous peroxidase activity. SARS-CoV-2 nucleocapsid monoclonal antibody (mAb) (GeneTex, catalog no. GTX635679) and a Streptavidin-Peroxidase Anti-Rabbit IgG kit (Maixin, Fuzhou, CHN, catalog no. KIT-9706) was used to quantify the viral level in lung tissue.

Statistical analysis

The statistical analyses used in the figures have been described in detail in the figure legends. All data were expressed as mean $\pm$ standard deviations (SDs). Statistical comparisons were performed using student's t-test, one-way ANOVA, or repeated

measurement ANOVA. The differences were statistically significant as follows: $*P < 0.05$,

$**P < 0.01$, $***P < 0.001$; ns is for no meaning

Ethics statement

Collection of inpatient blood was approved by the Ethics Committee of the First

Hospital of Jilin University (21K105-001) in accordance with the guidelines and principles

of WMA Declaration of Helsinki and the Department of Health and Human Services

Belmont Report. All study participants signed an informed consent form. PBMCS were

obtained from 9 COVID-19 patients with mild disease, 6 patients with severe disease, and

5 patients with critical disease (Appendix table S2). All animal experiments were approved

by the ethics committee of Research Unit of Key Technologies for Prevention and Control

of Virus Zoonoses, Chinese Academy of Medical Sciences, Changchun Veterinary

Research Institute, Chineses Academy of Agricultural Sciences (IACUC of

AMMS-11-2020-006).

Acknowledgments

We thank C.Y. Dai for providing critical reagents.

Funding information

This work was supported by funding from the National Natural Science Foundation of

China (82341072, 81930062 and 82272316 to ZWY, 82341062 to LZL), the National Key

R&D Program of China (2021YFC2301900 and 2301904, 2023YFC2306603), the

Science and Technology Department of Jilin Province (YDZJ202301ZYTS521 and

YDZJ202201ZYTS587), the Key Laboratory of Molecular Virology, Jilin Province

(20102209), and Bethune Project, Jilin University (2023B03). The funding sources were

involved in study design, data collection and interpretation, and   the decision to submit

the work for publication.

Authors' contributions

L.T.: Conceptualization; Data curation; Formal Analysis; Investigation; Methodology;

Writing – original draft. Z.Z.: Methodology; Data curation; Writing – review & editing. Z.L.:

Conceptualization; Data curation; Formal Analysis; Investigation; Methodology; Writing –

review & editing; Funding acquisition. X.L., Z.L. and W.G.: Methodology; Resources. W.Z.:

Conceptualization; Formal Analysis; Funding acquisition; Methodology; Project

administration; Resources; Writing – review & editing.

Data availability

The authors declare that [the/all other] data supporting the findings of this study are

available within the paper [and its supplementary information files]. The mass

spectrometry proteomics data have been deposited to the ProteomeXchange Consortium

(http://proteomecentral.proteomexchange.org) via the iProX partner repository with the

dataset identified PXD053961.

References

Balieiro, A. M., Anunciação, E. L., Costa, C. H., Qayed, W. S., & Silva, J. R. A. (2022). Computational
Analysis of SAM Analogs as Methyltransferase Inhibitors of nsp16/nsp10 Complex from
SARS-CoV-2. *International Journal of Molecular Sciences*, *23*(22), 13972.

Banerjee, A. K., Blanco, M. R., Bruce, E. A., Honson, D. D., Chen, L. M., Chow, A., Bhat, P., Ollikainen,
N., Quinodoz, S. A., & Loney, C. (2020). SARS-CoV-2 disrupts splicing, translation, and
protein trafficking to suppress host defenses. *Cell*, *183*(5), 1325-1339. e1321.

Bonamassa, B., Hai, L., & Liu, D. (2011). Hydrodynamic gene delivery and its applications in
pharmaceutical research. *Pharm Res*, *28*(4), 694-701.
https://doi.org/10.1007/s11095-010-0338-9

Cai, B., Zhao, J., Zhang, Y., Liu, Y., Ma, C., Yi, F., Zheng, Y., Zhang, L., Chen, T., & Liu, H. (2022).
USP5 attenuates NLRP3 inflammasome activation by promoting autophagic degradation
of NLRP3. *Autophagy*, *18*(5), 990-1004.

Chen, X., Tian, L., Zhang, L., Gao, W., Yu, M., Li, Z., & Zhang, W. (2024). Deubiquitinase USP39
promotes SARS-CoV-2 replication by deubiquitinating and stabilizing the envelope
protein. *Antiviral Res*, *221*, 105790. https://doi.org/10.1016/j.antiviral.2023.105790

Decroly, E., Debarnot, C., Ferron, F., Bouvet, M., Coutard, B., Imbert, I., Gluais, L., Papageorgiou, N.,
Sharff, A., & Bricogne, G. (2011). Crystal structure and functional analysis of the
SARS-coronavirus RNA cap 2′-O-methyltransferase nsp10/nsp16 complex. *PLoS*
pathogens, *7*(5), e1002059.

Fan, L., Zhou, Y., Wei, X., Feng, W., Guo, H., Li, Y., Gao, X., Zhou, J., Wen, Y., Wu, Y., Shen, X., Liu, L.,
Xu, G., & Zhang, Z. (2024). The E3 ligase TRIM22 restricts SARS-CoV-2 replication by
promoting proteasomal degradation of NSP8. *mBio*, *15*(2), e0232023.
https://doi.org/10.1128/mbio.02320-23

Gao, W., Wang, L., Cui, W., Wang, H., Huang, G., Li, Z., Li, G., & Zhang, W. (2024). Deubiquitinase
USP1 regulates sarbecovirus ORF6 protein function. *J Virol*, *98*(1), e0143723.
https://doi.org/10.1128/jvi.01437-23

Gao, W., Wang, L., Ju, X., Zhao, S., Li, Z., Su, M., Xu, J., Wang, P., Ding, Q., & Lv, G. (2022). The
deubiquitinase USP29 promotes SARS-CoV-2 virulence by preventing proteasome
degradation of ORF9b. *Mbio*, *13*(3), e01300-01322.

[Record #31 is using a reference type undefined in this output style.]

Grice, G. L., & Nathan, J. A. (2016). The recognition of ubiquitinated proteins by the proteasome. *Cellular and molecular life sciences*, *73*, 3497-3506.

Guo, G., Gao, M., Gao, X., Zhu, B., Huang, J., Luo, K., Zhang, Y., Sun, J., Deng, M., & Lou, Z. (2021). SARS-CoV-2 non-structural protein 13 (nsp13) hijacks host deubiquitinase USP13 and counteracts host antiviral immune response. *Signal transduction and targeted therapy*, *6*(1), 119.

Hojyo, S., Uchida, M., Tanaka, K., Hasebe, R., Tanaka, Y., Murakami, M., & Hirano, T. (2020). How COVID-19 induces cytokine storm with high mortality. *Inflamm Regen*, *40*, 37. https://doi.org/10.1186/s41232-020-00146-3

Hu, B., Huang, S., & Yin, L. (2021). The cytokine storm and COVID-19. *J Med Virol*, *93*(1), 250-256. https://doi.org/10.1002/jmv.26232

Ju, X., Zhu, Y., Wang, Y., Li, J., Zhang, J., Gong, M., Ren, W., Li, S., Zhong, J., Zhang, L., Zhang, Q. C., Zhang, R., & Ding, Q. (2021). A novel cell culture system modeling the SARS-CoV-2 life cycle. *PLoS Pathog*, *17*(3), e1009439. https://doi.org/10.1371/journal.ppat.1009439

Kamimura, K., Kanefuji, T., Yokoo, T., Abe, H., Suda, T., Kobayashi, Y., Zhang, G., Aoyagi, Y., & Liu, D. (2014). Safety assessment of liver-targeted hydrodynamic gene delivery in dogs. *PLoS One*, *9*(9), e107203. https://doi.org/10.1371/journal.pone.0107203

Kim, J. G., Shin, H.-C., Seo, T., Nawale, L., Han, G., Kim, B. Y., Kim, S. J., & Cha-Molstad, H. (2021). Signaling pathways regulated by UBR box-containing E3 ligases. *International journal of molecular sciences*, *22*(15), 8323.

Klima, M., Khalili Yazdi, A., Li, F., Chau, I., Hajian, T., Bolotokova, A., Kaniskan, H. Ü., Han, Y., Wang, K., & Li, D. (2022). Crystal structure of SARS-CoV-2 nsp10–nsp16 in complex with small molecule inhibitors, SS148 and WZ16. *Protein Science*, *31*(9), e4395.

Li, Z., Hao, P., Zhao, Z., Gao, W., Huan, C., Li, L., Chen, X., Wang, H., Jin, N., & Luo, Z.-Q. (2023). The E3 ligase RNF5 restricts SARS-CoV-2 replication by targeting its envelope protein for degradation. *Signal Transduction and Targeted Therapy*, *8*(1), 53.

Liang, X., Xiao, J., Li, X., Liu, Y., Lu, Y., Wen, Y., Li, Z., Che, X., Ma, Y., & Zhang, X. (2022). A C-terminal glutamine recognition mechanism revealed by E3 ligase TRIM7 structures. *Nature Chemical Biology*, *18*(11), 1214-1223.

Lin, S., Chen, H., Ye, F., Chen, Z., Yang, F., Zheng, Y., Cao, Y., Qiao, J., Yang, S., & Lu, G. (2020). Crystal structure of SARS-CoV-2 nsp10/nsp16 2′-O-methylase and its implication on antiviral drug design. *Signal transduction and targeted therapy*, *5*(1), 131.

Lugari, A., Betzi, S., Decroly, E., Bonnaud, E., Hermant, A., Guillemot, J.-C., Debarnot, C., Borg, J.-P., Bouvet, M., & Canard, B. (2010). Molecular mapping of the RNA Cap 2′-O-methyltransferase activation interface between severe acute respiratory syndrome coronavirus nsp10 and nsp16. *Journal of Biological Chemistry*, *285*(43), 33230-33241.

Mahajan, S., Sharma, G. K., Bora, K., & Pattnaik, B. (2021). Identification of novel interactions between host and non-structural protein 2C of foot-and-mouth disease virus. *J Gen Virol*, *102*(3). https://doi.org/10.1099/jgv.0.001577

Maimaitiyiming, Y., Yang, T., Wang, Q. Q., Feng, Y., Chen, Z., Björklund, M., Wang, F., Hu, C., Hsu, C.-H., & Naranmandura, H. (2022). Heat treatment promotes ubiquitin-mediated proteolysis of SARS-CoV-2 RNA polymerase and decreases viral load. *Research*.

Makaremi, S., Asgarzadeh, A., Kianfar, H., Mohammadnia, A., Asghariazar, V., & Safarzadeh, E. (2022). The role of IL-1 family of cytokines and receptors in pathogenesis of COVID-19.

Inflamm Res, *71*(7-8), 923-947. https://doi.org/10.1007/s00011-022-01596-w

McClellan, A. J., Laugesen, S. H., & Ellgaard, L. (2019). Cellular functions and molecular mechanisms of non-lysine ubiquitination. *Open Biology*, *9*(9), 190147.

Melo-Filho, C. C., Bobrowski, T., Martin, H.-J., Sessions, Z., Popov, K. I., Moorman, N. J., Baric, R. S., Muratov, E. N., & Tropsha, A. (2022). Conserved coronavirus proteins as targets of broad-spectrum antivirals. *Antiviral Research*, *204*, 105360.

Muñoz-Escobar, J., Matta-Camacho, E., Kozlov, G., & Gehring, K. (2015). The MLLE domain of the ubiquitin ligase UBR5 binds to its catalytic domain to regulate substrate binding. *Journal of Biological Chemistry*, *290*(37), 22841-22850.

Nathan, J. A., Sengupta, S., Wood, S. A., Admon, A., Markson, G., Sanderson, C., & Lehner, P. J. (2008). The ubiquitin E3 ligase MARCH7 is differentially regulated by the deubiquitylating enzymes USP7 and USP9X. *Traffic*, *9*(7), 1130-1145.

Park, G. J., Osinski, A., Hernandez, G., Eitson, J. L., Majumdar, A., Tonelli, M., Henzler-Wildman, K., Pawłowski, K., Chen, Z., & Li, Y. (2022). The mechanism of RNA capping by SARS-CoV-2. *Nature*, *609*(7928), 793-800.

Qiao, X., Liu, Y., Prada, M. L., Mohan, A. K., Gupta, A., Jaiswal, A., Sharma, M., Merisaari, J., Haikala, H. M., & Talvinen, K. (2020). UBR5 is coamplified with MYC in breast tumors and encodes an ubiquitin ligase that limits MYC-dependent apoptosis. *Cancer research*, *80*(7), 1414-1427.

Raper, S. E., Chirmule, N., Lee, F. S., Wivel, N. A., Bagg, A., Gao, G. P., Wilson, J. M., & Batshaw, M. L. (2003). Fatal systemic inflammatory response syndrome in a ornithine transcarbamylase deficient patient following adenoviral gene transfer. *Mol Genet Metab*, *80*(1-2), 148-158. https://doi.org/10.1016/j.ymgme.2003.08.016

Rosas-Lemus, M., Minasov, G., Shuvalova, L., Inniss, N. L., Kiryukhina, O., Brunzelle, J., & Satchell, K. J. (2020). High-resolution structures of the SARS-CoV-2 2′-O-methyltransferase reveal strategies for structure-based inhibitor design. *Science signaling*, *13*(651), eabe1202.

Russ, A., Wittmann, S., Tsukamoto, Y., Herrmann, A., Deutschmann, J., Lagisquet, J., Ensser, A., Kato, H., & Gramberg, T. (2022). Nsp16 shields SARS–CoV-2 from efficient MDA5 sensing and IFIT1-mediated restriction. *EMBO reports*, *23*(12), e55648.

Shearer, R. F., Frikstad, K.-A. M., McKenna, J., McCloy, R. A., Deng, N., Burgess, A., Stokke, T., Patzke, S., & Saunders, D. N. (2018). The E3 ubiquitin ligase UBR5 regulates centriolar satellite stability and primary cilia. *Molecular biology of the cell*, *29*(13), 1542-1554.

Song, P., Li, W., Xie, J., Hou, Y., & You, C. (2020). Cytokine storm induced by SARS-CoV-2. *Clin Chim Acta*, *509*, 280-287. https://doi.org/10.1016/j.cca.2020.06.017

Suda, T., Yokoo, T., Kanefuji, T., Kamimura, K., Zhang, G., & Liu, D. (2023). Hydrodynamic Delivery: Characteristics, Applications, and Technological Advances. *Pharmaceutics*, *15*(4). https://doi.org/10.3390/pharmaceutics15041111

Wu, B., Song, M., Dong, Q., Xiang, G., Li, J., Ma, X., & Wei, F. (2022). UBR5 promotes tumor immune evasion through enhancing IFN-γ-induced PDL1 transcription in triple negative breast cancer. *Theranostics*, *12*(11), 5086.

Xiang, G., Wang, S., Chen, L., Song, M., Song, X., Wang, H., Zhou, P., Ma, X., & Yu, J. (2022). UBR5 targets tumor suppressor CDC73 proteolytically to promote aggressive breast cancer. *Cell Death & Disease*, *13*(5), 451.

Xu, G., Wu, Y., Xiao, T., Qi, F., Fan, L., Zhang, S., Zhou, J., He, Y., Gao, X., & Zeng, H. (2022).

Multiomics approach reveals the ubiquitination-specific processes hijacked by
SARS-CoV-2. *Signal Transduction and Targeted Therapy*, *7*(1), 312.

Zhang, H., Zheng, H., Zhu, J., Dong, Q., Wang, J., Fan, H., Chen, Y., Zhang, X., Han, X., & Li, Q.
(2021). Ubiquitin-modified proteome of SARS-CoV-2-infected host cells reveals insights
into virus–host interaction and pathogenesis. *Journal of Proteome Research*, *20*(5),
2224-2239.

Zhang, J., Cruz-Cosme, R., Zhuang, M.-W., Liu, D., Liu, Y., Teng, S., Wang, P.-H., & Tang, Q. (2020).
A systemic and molecular study of subcellular localization of SARS-CoV-2 proteins.
Signal transduction and targeted therapy, *5*(1), 269.

Zhang, L., Hao, P., Chen, X., Lv, S., Gao, W., Li, C., Li, Z., & Zhang, W. (2024). CRL4B E3 ligase
recruited by PRPF19 inhibits SARS-CoV-2 infection by targeting ORF6 for
ubiquitin-dependent degradation. *mBio*, *15*(2), e0307123.
https://doi.org/10.1128/mbio.03071-23

Zhang, L., Wang, H., Tian, L., & Li, H. (2016). Expression of USP7 and MARCH7 is correlated with
poor prognosis in epithelial ovarian cancer. *The Tohoku journal of experimental medicine*,
239(3), 165-175.

Zhang, X., Yang, Z., Pan, T., Sun, Q., Chen, Q., Wang, P. H., Li, X., & Kuang, E. (2023). SARS-CoV-2
Nsp8 suppresses MDA5 antiviral immune responses by impairing TRIM4-mediated
K63-linked polyubiquitination. *PLoS Pathog*, *19*(11), e1011792.
https://doi.org/10.1371/journal.ppat.1011792

Zhao, K., Yang, Y., Zhang, G., Wang, C., Wang, D., Wu, M., & Mei, Y. (2018). Regulation of the
Mdm2–p53 pathway by the ubiquitin E3 ligase MARCH 7. *EMBO reports*, *19*(2), 305-319.

Zheng, C. (2021). The emerging roles of the MARCH ligases in antiviral innate immunity.
International Journal of Biological Macromolecules, *171*, 423-427.

Zhou, Y., Zheng, R., Liu, D., Liu, S., Disoma, C., Li, S., Liao, Y., Chen, Z., Du, A., & Dong, Z. (2022).
UBR5 Acts as an Antiviral Host Factor against MERS-CoV via Promoting Ubiquitination
and Degradation of ORF4b. *Journal of Virology*, *96*(17), e00741-00722.

Züst, R., Cervantes-Barragan, L., Habjan, M., Maier, R., Neuman, B. W., Ziebuhr, J., Szretter, K. J.,
Baker, S. C., Barchet, W., & Diamond, M. S. (2011). Ribose 2′-O-methylation provides a
molecular signature for the distinction of self and non-self mRNA dependent on the RNA
sensor Mda5. *Nature immunology*, *12*(2), 137-143.

Figure legends

Figure 1. The non-structural protein nsp16 of SARS-CoV-2 was identified that can be degraded through the proteasome pathway.

A. The non-structural proteins nsp8, nsp11 and nsp16 could be restored by the proteasome inhibitor MG132. HEK293T cells in 12-well plates were transfected with the plasmids of 16 nonstructural proteins (nsp1-16) encoded by SARS-CoV-2. Thirty-six hours later, the cells were treated with MG132 (10 µM) or DMSO for 12 h before collection. The protein level was detected by Immunoblotting (IB). Quantification of nsp protein levels relative to the control protein is shown. Data are representative of three independent experiments and shown as average ±SD (n = 3). Significance was determined by a two-tailed t-test: *P < 0.05; **P < 0.01; ***P < 0.001.

B. Proteasomal inhibitors but no other inhibitors stabilized nsp16 protein. HEK293T cells transfected with the nsp16-Flag expression vector were treated with dimethyl sulfoxide (DMSO), MG132 (10 µM), Bortezomib (10 µM), Carfilzomib (10 µM), Bafilomycin A1 (5 µM), Vinblastine (2.5 µM), or NH_4CL (2.5 µM) for 12 h prior to harvest. The cell lysates were analyzed by anti-Flag antibody.

(C-D). The half-life of nsp16 was prolonged by the proteasome inhibitor MG132. C. HEK293T cells were transfected with the nsp16-Flag-expressing plasmids. 12 hours later, the cells were treated with DMSO or MG132 (10 µM) for 12 h, then 50 µg/mL cycloheximide (CHX) was added. Cells were harvested at the indicated times to detect the level of viral protein by anti-Flag antibody. D. Quantification of nsp16 protein levels relative to tubulin at different time points is shown. The half-life of the nsp16 protein was determined based on protein quantification using Image J, combined with the protein half-life formula for calculation. Results are shown as mean ± SD (n = 3 independent experiments). ***, P < 0.001 by by a two-tailed t-test.

E. Samples were prepared for mass spectrometry, and nsp16 interacting proteins were obtained by immunoprecipitation (IP) (created with BioRender.com and the agreement no. is LC27PCBEOF). The plasmids were transfected into HEK293T cells for 48 h. Treat cells

with or without MG132 (10 µM) for 12 h prior to harvest. The whole-cell lysates were incubated with protein G agarose beads conjugated with anti-Flag antibodies and used for IB with anti-Flag antibodies to detect the nsp16 protein. Samples enriched for proteins were analyzed by mass spectrometry.

Figure 1—figure supplement 1. Analysis of mass spectrometry results.

A. The DAVID bioinformatics website (https://david.ncifcrf.gov/tools.jsp) was used for KEGG pathway analysis and GO analysis of the differential proteins bound by nsp16 protein after treating with or without MG132. The bubble plots or circle diagram depict the results of the analysis separately. In bubble plots, a change from yellow to purple indicates a decreased p-value, while the size of the circles indicates the number of enriched genes. In the circle plot, depicting the BP (Biological process) analysis in the GO term, the length of the purple rectangle indicates the number of genes included in the term. The length of the green rectangle indicates the number of overlapping genes between the genes included in the term and the genes entered in the gene enrichment analysis, a change from deep to shallow of purple indicates a decreased p-value (created with chiplot.com).

B. Schematic representation of proteins degraded by the proteasome pathway, as well as proteins associated with the proteasome in the MS enrichment analysis are shown (created with BioRender.com and the agreement no. is VO27O5XHHC).

Figure 2. MARCHF7 and UBR5 were identified as E3 ubiquitin ligases involves in nsp16 protein degradation.

A. Knockdown of MARCHF7 or UBR5 resulted in nsp16 restoration. HEK293T cells were transfected with siRNA of E3 ligase candidates for 24 h, followed by co-incubation with the nsp16-Flag-expressing plasmids for 48 h, treated with MG132 (10 µM) for 16h before harvesting, lysed, and subjected to IB assay using anti-Flag antibody. RT-qPCR was conducted to determine the mRNA expression levels of E3 ligase candidates. The si-RNA targeting regions for the candidate E3 ubiquitin ligase proteins and the targeted regions for RT-qPCR are shown in Appendix-figure S1A. Data are representative of three independent experiments and shown as average ± SD (n = 3). Significance was

determined by a two-tailed t-test: ***P < 0.001.

B. RNA levels of UBR5 or MARCHF7 from HEK293T cells infected with lentivirus

containing control or shRNA targeting UBR5 or MARCHF7 for 48 h and screened with

antibiotics for 48 h. Knockdown cell lines were transfected with plasmids expressing

nsp16-Flag, collected at the indicated times, and the protein levels of nsp16, MARCHF7,

and UBR5 were detected by IB.C. MARCHF7 and UBR5 acted separately and did not

depend on each other. HEK293T cells stably expressing UBR5 shRNA or MARCHF7

shRNA were transfected with siRNA of MARCHF7 or UBR5 for 24 h, respectively, followed

by co-incubation with the nsp16-Flag-expressing plasmids for 48 h. The protein levels and

the RNA levels of nsp16, UBR5 and MARCHF7 were measured by IB and RT-qPCR,

respectively.

(D-E). In HEK293T cells stably expressing UBR5 shRNA or MARCHF7 shRNA, nsp16

was degraded by overexpressed UBR5 or MARCHF7, respectively, whereas the mutant

failed to degrade nsp16. The cell lysates were analyzed by anti-Flag antibody.

Figure 2—figure supplement 1. MARCHF7 and UBR5 degrade nsp16 independently.

(A-B). UBR5 siRNA was transfected into shMARCHF7 cells to knock down UBR5. After 24

hours, MARCHF7 and nsp16 expression vectors were co-transfected, and the cells were

harvested 72 hours later. The levels of nsp16 were characterized by IB with anti-Flag

antibody. Whether MARCHF7 was dependent on UBR5 to degrade of nsp16 was

determined by further transfection of MARCHF7 siRNA into shUBR5 cells, followed by

co-transfection of UBR5 and nsp16 expression vectors 24 h later. The other operations

are the same as above. Data are representative of three independent experiments and

shown as average $\pm$SD (n = 3). Significance was determined by a two-tailed t-test. *P >

0.05; **P < 0.01; ***P < 0.001.

Figure 3. MARCHF7 or UBR5 catalyze the formation of K-27 type or K-48 type

ubiquitin chains of nsp16 respectively.

A. Nsp16 can be ubiquitinated. HEK293T cells co-transfected with ubiquitin-Myc and

nsp16-Flag or transfected with nsp16-Flag alone. The cells were treated with MG132 for

12 h before collection. The whole-cell lysates were incubated with anti-Flag beads and used for IB with anti-Myc or anti-Flag antibodies to detect the polyubiquitination chain of nsp16.

B. Assess the endogenous ubiquitination level of nsp16 protein. Cells were transfected with nsp16-Flag or an empty vector, and collected 48 hours later. Prior to harvesting, cells were treated with MG132 for 16 hours. Co-IP experiments were then performed to analyze the endogenous ubiquitination level of nsp16.

C. The level of ubiquitination of nsp16 decreased with decreasing the protein levels of MARCHF7 or UBR5. E3 was knocked down by transfection with siRNA targeting UBR5 or MARCHF7, and 24 h later ubiquitin-Myc and nsp16-HA were co-transfected or nsp16-HA alone. Cells were treated with MG132 for 16 h before collection. Whole cell lysates were incubated with anti-HA beads, and polyubiquitinated chains of nsp16 were detected by IB with anti-Myc or anti-HA antibodies.

D. Nsp16 can be modified by a variety of ubiquitin chains. HEK293T cells were transfected with either nsp16-HA alone or together with plasmids encoding various mutants of ubiquitin (K6 only, K11 only, K27 only, K29 only, K33 only, K48 only, K63 only). Thirty-six hours later, cells were treated with MG132 for 12 h. Cell lysates were then subjected to immunoprecipitation, followed by IB to analysis.

(E-F) MARCHF7 or UBR5 causes nsp16 to be modified by the K27 type or K48 type ubiquitin chain. 293T cell lines with or without MARCHF7 or UBR5 knockdown were co-transfected with plasmids encoding ubiquitin-WT or various mutants of ubiquitin (K6 only, K11 only, K27 only, K29 only, K33 only, K48 only, K63 only). The other experimental methods were the same as C.

Figure 4. MARCHF7 and UBR5 directly interact with nsp16 respectively.

(A-B). The binding of MARCHF7 or UBR5 to nsp16 was not mutually dependent. The binding of nsp16 to UBR5 or MARCHF7 was identified by co-immunoprecipitation in HEK293T cells transfected siMARCHF7 or siUBR5, respectively. The immunoprecipitates and input were analyzed by IB. The knockdown efficiency was detected by RT-qPCR and IB.

(C-D). MARCHF7 or UBR5 co-localized with nsp16 in the endoplasmic reticulum. Hela

cells were co-transfected with YFP-nsp16(yellow) and CFP-UBR5(cyan) or

CFP-MARCHF7(cyan). The organelles were labeled with antibodies against marker

proteins of endoplasmic reticulum, Golgi apparatus and mitochondria respectively(red).

The cells were analyzed by confocal microscopy (C). Scale bars, 20 um. The ratio of

colocalization was quantified by measuring the fluorescence intensities using Image J (D).

Figure 4—figure supplement 1. Interaction of MARCHF7 or UBR5 with nsp16.

A. HEK293T cells were transfected with either nsp16-Flag alone or together with

MARCHF7-Myc. Thirty-six hours after transfection, the cells were treated with MG132 (10

μM) for 12 h. Cell lysates were subjected to immunoprecipitation with anti-Flag antibody.

Using IB to analyze the precipitates and input.

B. HEK293T cells were transfected with nsp16-Flag. Cell lysates were subjected to

immunoprecipitation with anti-UBR5 or IgG antibody.

(C-D). Hela cells were co-transfected with YFP-nsp16 and CFP-UBR5 or CFP-MARCHF7.

A representative image of YFP-nsp16 (yellow) and ECFP-MARCHF7 (cyan) or

ECFP-UBR5 (cyan) expressing cells before and after photobleaching the acceptor

fluorophore, YFP. The region chosen for photobleaching is marked (white open box), Bars,

10um. The quantization of fluorescence brightness was analyzed by Image J. Data are

representative of three independent experiments and shown as average $\pm$ SD (n = 3).

Significance was determined by a two-tailed t-test. *P > 0.05; **P < 0.01; ***P < 0.001.

E. HeLa cells transfected with nsp16-Flag were analyzed by confocal microscopy. The

Flag-tagged nsp16 labeled with anti-Flag antibody (red). MARCHF7 or UBR5 were

labeled with endogenous antibodies (green). Cell nuclei were stained using DAPI (blue).

Representative images were shown. Scale bars, 20 um. The ratio of colocalization was

quantified by measuring the fluorescence intensities using Image J.

F. nsp16 was stably transfected into HEK293T cells. The cells were analyzed by confocal

microscopy. The other operations are the same as above.

Figure 4—figure supplement 2. Domains in which MARCHF7 or UBR5 functions.

A. The schematic represents UBR5 WT or mutants used in the study.

B. The HECT domain of UBR5 is required for nsp16 degradation. After co-transfection

with UBR5 WT or mutants and nsp16-HA, cells were harvested 48 hours later, and cell

lysates were examined by IB.

C. The HECT domain of UBR5 affects K48-type ubiquitin chain of nsp16.HEK293T cells

were transfected with the assigned plasmids. After 36 hours, cells were treated with 10 μM

MG132 for 12 hours, harvested, and cell lysates were incubated with protein G agarose

beads conjugated with anti-HA antibodies. Cell lysates and precipitated samples were

analyzed by IB.

D. The schematic represents wild-type and truncated forms of MARCHF7 used in the

study. E. Only MARCHF7 wild-type degraded nsp16.

F. The N-terminal region of MARCHF7 interacted with nsp16, and only the wild type could

catalyze the K27-type ubiquitin chain of nsp16.

Figure 5. Knockdown of MARCHF7 or UBR5 promotes viral replication.

A. The virus-encoded nsp16 protein interacts with endogenous MARCHF7 and UBR5 and

undergoes ubiquitination modification. In 293T-ACE2 cells, with or without Wuhan strain

infection (MOI: 0.01), the medium was changed 2 hours post-infection, and cells were

harvested 48 hours later, with MG132 treatment added 16 hours before harvesting. nsp16

protein was enriched using Protein-G beads coupled with the nsp16 antibody, and

interactions and ubiquitination were analyzed by immunoblotting (IB) with endogenous

antibodies against MARCHF7, UBR5, and ubiquitination.

(B-I). MARCHF7 and UBR5 were knocked down by siRNA in Caco2 cells. 24 h after

transfection, the cells were infected with Wuhan strain (MOI:0.01) (C-E) or Omicron BA.1

strain (MOI: 0.001) (F-H), respectively. 2 h post infection, the supernatant was discarded,

and the cells were cultured in DMEM containing 3% fetal bovine serum for 48 h. The

mRNA levels of SARS-CoV-2 *M* and *E* genes in the cells (C, F) and *E* genes in

supernatant (D, G) were detected by RT-qPCR and the viral titers in supernatant (E, H)

were measured. The N protein levels of Wuhan or Omicron viruses were detected by IB (I).

Knock-down efficiencies of MARCHF7 and UBR5 were detected by RT-qPCR or IB (B, I).

Data are representative of three independent experiments and shown as average ±SD (n

= 3). Significance was determined by one-way ANOVA, followed by a Tukey multiple

comparisons posttest: *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$.

Figure 5—figure supplement 1. Effect of MARCHF7 or UBR5 on SARS-CoV-2 trVLP.

(A-B). Knocking down MARCHF7 or UBR5 enhances SARS-CoV-2 trVLP infectivity.

MARCHF7 or UBR5 was knocked down by siRNA in Caco2 cells with stable expression of

SARS-CoV-2 N protein. Twenty-four hours later, cells were infected with SARS-CoV-2

virus-like-particles (MOI:0.1), the medium was changed two hours after infection, and the

eGFP-positive cells were detected by flow cytometry 48 hours later (A). Protein content

was determined by RT-qPCR (B).

Figure 6. Increased levels of nsp16 rescued viral inhibition by UBR5 or MARCHF7'

(A-H) UBR5 or MARCHF7 was transfected in 293T cells stably overexpressed with ACE2,

and the increased doses of nsp16-Flag was transfected simultaneously. After 24 h, the

cells were infected with Wuhan strains. The mRNA levels of *M* and *E* genes of Wuhan

strain in the cells (A, D) and *E* gene in supernatant (B, E) were detected by RT-qPCR, as

well as the detection of viral titers in supernatant (C, F). The N protein of the virus and the

overexpression efficiency was detected by IB (G, H). Data are representative of three

independent experiments and shown as average ±SD (n = 3). Significance was

determined by one-way ANOVA, followed by a Tukey multiple comparisons posttest. $P >$

0.05; **$P < 0.01$; ***$P < 0.001$. Figure 6 — figure supplement 1 shows data related to

infection with Omicron BA.1.

Figure 6 — figure supplement 1. Effect of MARCHF7 or UBR5 on Omicron BA.1

infectivity.

(A-H). Showing data related to infection with Omicron BA.1. The experimental procedure

was the same as Figure 6. Data are representative of three independent experiments and

shown as average $\pm$ SD (n = 3). Significance was determined by one-way ANOVA,

followed by a Tukey multiple comparisons posttest. $P > 0.05$; **$P < 0.01$; ***$P < 0.001$.

Figure 6—figure supplement 2. The enzyme activity-deficient mutants do not exhibit

antiviral activity, and overexpression of nsp16 does not promote viral replication.

(A-H) In 293T-ACE2 cells, the RING domain deletion mutant of MARCHF7

(MARCHF7-1-542) or the HECT domain inactivated mutant of UBR5 (UBR5-△HECT)

were transfected, along with a gradient of nsp16-Flag overexpression. The cells were

infected with the Wuhan strain (MOI: 0.01), medium was changed 2 hours post-infection,

and cells and supernatants were collected 48 hours after infection.

Figure 6—figure supplement 3. MARCHF7 or UBR5 have effects on mutant of nsp16

in different subtypes of SARS-CoV-2.

A. This diagram shows the mutation of nsp16 in different virus subtypes. The amino acids

sequences of different SARS-CoV-2 strains were obtained from NCBI, and the amino

acids sequences of nsp16 of different strains were compared by DNAMAN software.

B. Nsp16 mutants can still be regulated by MG132. The mutated nsp16 plasmids were

transfected into HEK293T cells. After 36 hours of culture, cells were treated with 10 μm

MG132 or DMSO, harvested 12 hours later, and cell lysates were examined by IB.

C. MARCHF7 or UBR5 can degrade nsp16 mutants. After transfecting MARCHF7 or

UBR5 siRNA and the mutated nsp16 plasmids, the cells were harvested 48 h later. The

cell lysates were detected by IB.

Figure 6—figure supplement 4. The expression level of MARCHF7 was negatively

correlated with the viral titer, while the expression level of UBR5 was increased at

low titer and decreased at high titer.

(A-C). The protein and mRNA levels of MARCHF7 or UBR5 upon infection with different

titers. Endogenous MARCHF7 and UBR5 RNA levels were detected by RT-qPCR 48

hours after infection with different titers of Wuhan strain (MOI:0, 0.0001, 0.001, 0.01) or

omicron BA.1 strain (MOI:0, 0.0001, 0.001). Protein levels were examined by IB.

D. The expression level of UBR5 was negatively correlated with the severity of the

disease but MARCHF7 expression levels were not. PBMC cells were extracted from

common, severe and critical COVID-19 patients. RT-qPCR was used to detect the mRNA

level of UBR5 or MARCHF7 in patients. Significance was determined by one-way ANOVA,

followed by a Tukey multiple comparisons posttest. ns, P > 0.05; **P < 0.01; ***P < 0.001.

Figure 7. In a mouse infection model, overexpression of MARCHF7 or UBR5 exerted

inhibitory effects on virus.

(A-G) BLAB/C mice were injected with the corresponding plasmids at 40ug/500ul via the

high-pressure tail vein, followed by nasal inoculation with 50µl SARS-CoV-2 virus at a

dosage of $10^{5.5}$ TCID50/mL (created with BioRender.com and the agreement no. is

FO27O5X39A). IB was used to detect the expression of MARCHF7 or UBR5 in the lung

tissues (B). Viral RNA loads in mouse lung tissues were detected by measuring the mRNA

levels of the E genes by RT-qPCR (C). Lung tissue was collected, homogenized, and the

residue was removed by centrifugation to collect the supernatant. The viral titer was then

measured using the TCID50 method (D). Mouse body weight was monitored during the

experimental period (E). Representative images of H&E staining of lungs of mice with

different treatments. Magnification, ×40. Bars, 20 µm (F). The staining of viral N proteins.

Magnification, ×63. Bars, 20 µm. n =3 in each group (G). RT-qPCR was used to measure

the expression of cytokines and chemokines in the spleens of mice in each group (H).

Statistical significance was analyzed using a one-way analysis of variance with Tukey's

multiple comparisons test. (NS, no significance, *p < 0.05, **p < 0.01, ***p < 0.001).

Figure 8. Schematic diagram of MARCHF7 and UBR5 ubiquitinate the SARS-CoV-2

non-structural protein nsp16, leading to its degradation via the proteasomal

pathway, thereby affecting viral replication (created with BioRender.com and the

agreement no. is FM27O5ZS3U).

1023

Figure 1

Figure 1 supplement 1

Figure 2

Figure 2 supplement 1

A

B

Figure 3

Figure 4

Figure 4 supplement 1

Figure 4 supplement 2

Figure 5

Figure 5 supplement 1

Figure 6

Figure 6 supplement 1

Figure 6 supplement 2

Figure 6 supplement 3

A

B

C

Figure 6 supplement 4

Figure 7

Figure 8