Spatial Heterogeneity of Climate Fluctuations Shapes Ecological Processes and Community Dynamics in Riverine Invertebrates Across a Landscape

Xiaowei Lin[1,2], Jialu Wan[1,2], Qingyi Luo[1,2], Qinghua Cai[1*], Ming-Chih Chiu[1] and Vincent H. Resh[3]

[1]Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China

[2]College of Advanced Agricultural Sciences, University of Chinese Academy of Sciences, Beijing, China

[3]Department of Environmental Science, Policy & Management, University of California Berkeley, Berkeley 94720, USA

[*]Corresponding Author:

QC: qhcai@ihb.ac.cn; MCC: mingchih.chiu@gmail.com

Other authors' emails

XL: linxiaowei@ihb.ac.cn; VHR: resh@berkeley.edu

QL: luoqingyi@ihb.ac.cn; JW: wanjial@ ihb.ac.cn

Conflict of Interest

The authors declare no conflict of interest.

Author Contributions

XL, QL, QC, MCC, and VHR drafted the article and conceptualized the project. XL

and MCC devised the research methodology. XL, JW and MCC conducted data

processing and analysis. All authors reviewed and revised the article.

Data availability statement

The data that supports the findings of this study are available in the supporting

information Data S4 at https://onlinelibrary.wiley.com/doi/10.1111/jbi.13913.

Funding

Qinghua Cai was funded by the Second Tibetan Plateau Scientific Expedition and

Research Program (2019QZKK0402). Ming-Chih Chiu was funded by the Chinese

Academy of Sciences Taiwan Young Talent Programme (Grant No. 2017TW2SA0004-

Y-Y).

Acknowledgments

We gratefully acknowledge the authors (Canedo-Arguelles, M., Gutierrez-Canovas, C.,

Acosta, R., Castro-Lopez, D., Cid, N., Fortuno, P., et al.) of the study titled "As time

goes by: 20 years of changes in the aquatic macroinvertebrate metacommunity of

Mediterranean river networks" for generously sharing their data.

Abstract

Climate fluctuations exhibit spatial heterogeneity in influencing ecological processes and community dynamics. However, the mechanisms underlying these impacts on biodiversity remain largely unexplored. This study investigates how spatial variations in climate fluctuations affect the relative significance of ecological processes shaping invertebrate community dynamics across 15 rivers in the European Iberian Peninsula over 21 years. Spatial shifts in community dynamics were driven by a combination of deterministic (climatic filtering) and stochastic (temporal variability) processes. Notably, the dominance of deterministic versus stochastic processes showed an elevation-dependent pattern characterized by a single peak. Analyzing community dynamics across different elevations provides a foundational basis for predicting and mitigating the biodiversity impacts of climate change, guiding effective conservation strategies.

Keywords: climatic fluctuation, community dynamics, temporal beta diversity, elevation patterns, stochastic processes, deterministic processes

1 Introduction

Overwhelming evidence suggests an unprecedented decline in global biodiversity and the vital ecosystem services it delivers [1]. This trend is particularly pronounced for species facing varied fates due to adaptations in their life histories, morphology, and physiology amidst the complexities of a changing global environment. For instance, species inhabiting high-altitude or latitude regions often struggle to adapt to shifting climatic conditions [2]. Spatial heterogeneity within landscapes is crucial for understanding how diverse biological communities respond to both anthropogenic and natural environmental changes through various ecological processes.

Spatial variations in climatic fluctuations are pivotal in shaping the geographic distribution of community dynamics. For instance, the influence of climate variability on community dynamics has been extensively documented [3-5]. Moreover, with climate fluctuations intensifying at higher latitudes and elevations [6], disparities in community dynamics may arise due to geographical distinctions. Existing research has predominantly focused on two main areas: the spatial patterns of climate fluctuations [7,8] and the impacts of these fluctuations on biogeographic patterns and community dynamics [5,9,10]. However, the interaction between spatial patterns of climate variability and the geographic distribution of community dynamics remains largely unexplored from a mechanistic perspective involving ecological processes.

The importance of ecological processes is integral to understanding the underlying mechanisms that result in the resulting spatial patterns in community dynamics. Deterministic processes are typically associated with environmental and biological

filtration, whereas stochastic processes are more commonly linked to probability distributions and ecological drift [11-13]. For instance, the relative importance of ecological processes undergoes a transformation with community succession. Over time, there is a shift from stochastic to deterministic processes, which become the dominant forces shaping community composition [12,14]. Additionally, different time scales influence the relative importance of ecological processes. Some studies have indicated that deterministic processes tend to prevail over longer time scales, influencing community dynamics [15,16].

This study represents a pioneering effort to investigate the intricate relationship between spatial climate fluctuations and community dynamics from a mechanistic standpoint rooted in ecological processes. We conducted an analysis of ecological processes influencing community dynamics using extensive long-term data on benthic macroinvertebrates from a geographically expansive and continuously monitored area (Fig. 1). Our primary hypothesis examined whether heightened climate fluctuations, particularly in higher elevations, significantly influence the geographical patterns observed in community dynamics. Furthermore, we assessed how changes in elevation might modify the relative contributions of stochastic and deterministic processes driving biological community dynamics.

2 Materials and Methods

2.1 Study area and Macroinvertebrate data

The benthic macroinvertebrate data used in this study were originally collected and analyzed by Cañedo-Argüelles, et al. [17] from 15 rivers located in the northeastern

part of the Iberian Peninsula in Spain. This region is characterized by a Mediterranean climate, with annual precipitation ranging from 500 to 1400 mm and mean annual temperatures between 7 and 15°C. According to the Assessment Criteria of Mediterranean Rivers [18], the human pressures in the studied river reaches are assessed as relatively low.

Figure 1. Sampling sites in the Iberian Peninsula studied by Cañedo-Argüelles, et al. [17]

2.2 Climate variables and Time variables

In addition to the biological data analyzed by Cañedo-Argüelles, et al. [17], which focused on the spatial variation of community structure, we incorporated additional variables from diverse sources to explore various aspects. Elevation data and monthly

climatic variables such as mean minimum temperature, mean maximum temperature,

and total precipitation were sourced from the WorldClim 2 database [19] for each

sampling site over the period spanning 1997 to 2017. These data were further processed

using the "biovars" function within the R package "dismo" [20] to derive 19 physical

and climatic variables, including annual precipitation, mean annual temperature,

minimum temperature of the coldest month, and maximum temperature of the hottest

month (see Appendix S1 for details).

Temporal variables were generated to represent stochastic processes in time (e.g.,

ecological drift) using the asymmetric eigenvector map (AEM) method, which is a

filtering approach to model the effects produced by directed processes (e.g., time) [21].

Studies of time-series dynamics often necessitate logical consistency, which posits that

a previous point in time will have an effect on a later point in time. Consequently, the

AEM approach represents the optimal methodology for such studies [22]. The sampling

times of each sample point were sorted, placed in the same frame, and the AEMs are

computed in such a way that each sampling time point corresponds to the corresponding

AEM. In calculating the AEM, the sampling time interval was employed to determine

the weights. The formula was as follows：

$$Weight = 1 - (d / d\max)^2$$

[1]

In this context, the notation "d" represents the sampling time interval, while

"dmax" denotes the maximum value of the sampling time interval. Given a sample size

of n, the AEM produces a total of n-1 positive eigenroots, with no negative eigenroots.

The eigenfunctions can be divided into two sets of variables with positive and negative temporal correlation [23]. Based on the Moran I coefficients, we selected positively correlated AEMs [21], as detailed in Appendix S1. The AEMs were computed with the "aem.time" function from the "adespatial" package [24].

2.3 Data analysis

The objective of this study was to examine how climate fluctuations vary with altitude. To this end, the temporal dynamics of the altitudinal patterns of total monthly precipitation, mean monthly minimum temperature, and mean monthly maximum temperature were examined. Additionally, the coefficients of variation (CVs) of total monthly precipitation, mean monthly minimum temperature, and mean monthly maximum temperature were calculated and their variation with altitude was explored. The test is performed using the generalized additive model (GAM).

The GAM model was employed to investigate elevation patterns in species richness across sample sites over a 21-year period. Subsequently, the presence and absence data were utilized to calculate three distance matrices for each sample point based on the Jaccard coefficient of variation. These matrices represent temporal beta diversity (Btotal), its decomposed species substitution (Brepl), and richness difference (Brich). The distance matrices were obtained using the "BAT" package in R, as outlined by Cardoso [25]. Subsequently, the GAM model was employed to investigate the elevation patterns of temporal beta diversity and its components, with the relative importance of the different components of temporal beta diversity then calculated. The Wilcoxon test was employed to ascertain which of the components of temporal beta

diversity (Brep and Brich) was the primary driver.

To validate how elevational change in the relative significance of stochastic and deterministic processes can drive the shift of community dynamics, we employed generalized dissimilarity models (GDMs) to estimate temporal beta diversity and its components, while accounting for climatic and temporal variables [26,27]. GDMs transform environmental variables through an iterative maximum likelihood fitting process with the objective of identifying the optimal fit between environmental discrepancies and community differences [28].

The GDM assumes that biome variance increases monotonically with environmental variance and employs a link function to transform linear predictors in order to adjust for the nonlinear relationship between the two [28]. The GDM model fit is constrained by the number of i-spline basis functions, with a default value of three i-splines [29]. We conducted variance decomposition at each sample point to assess the rate of explanation of temporal beta diversity and its components by climatic and temporal variables. This was accomplished using the the "gdm.partition.deviance" function in the R package "gdm" [28,30]. Then, the GAMs were employed to investigate the relationship between the ratio of pure explanatory rates of climatic and temporal variables and elevation, with the objective of identifying patterns of elevation on the various ecological processes that shape community dynamics.

3 Results

3.1. Climatic fluctuations

We computed the total monthly precipitation and the average monthly minimum- and

maximum- temperatures in relation to elevation, and found that precipitation gradually

increases with elevation through the GAM model, and that both minimum and

maximum temperatures diminish with rising elevation (Fig. 2a, 2d, 2g). When

calculating the coefficient of variation (CV) for precipitation, minimum temperature,

and maximum temperature over the study period, we found a decrease in precipitation

fluctuation with increasing altitude, but that the fluctuation of minimum and maximum

temperature increased gradually (Fig. 2b, 2e, 2h). The results indicated that the

frequency of extreme weather events (precipitation, minimum and maximum

temperatures), although different at different altitudes, did not show a clear altitudinal

pattern (Fig. 2c, 2f, 2i). For the definition and frequency of extreme events, see

Appendix S3 for details: Table S2, and further statistical information on the GAM

model, please refer to Appendix S3.

Figure 2. Elevation patterns of (a) monthly total precipitation, (b) the variability of monthly

total precipitation, (c) the frequency of monthly total precipitation, (d) monthly mean

minimum temperature, (e) the variability of monthly mean minimum temperature, (f) the

frequency of monthly mean minimum temperature, (g) monthly mean maximum

temperature, (h) the variability of monthly mean maximum temperature, (i) the frequency of

monthly mean maximum temperature.

3.2 Biotic dynamics

Using generalized additive modeling (GAM), species richness followed a bimodal

pattern with elevation over the study period (Fig. 3a). The temporal beta diversity (Btoal)

gradually decreased with increasing elevation, but its components Brepl and Brich did

not show a clear or regular change with elevation (Fig. 3b, 3c, 3d). Species turnover

(Brepl) was significantly higher than richness difference (Brich) when a wilcoxon test

for Brepl and Brich was performed. This result suggested that species turnover

dominated the temporal beta diversity in the region (Refer to Appendix S3 for further

details).

Figure 3. Elevation patterns of (a) Richness, (b) temporal beta diversity (Btotal), (c) species

replacement (Brepl), and (d) richness difference in temporal beta diversity (Brich).

3.3 Ecological processes

The GDM model's explanatory power for temporal beta diversity and its

components (Brepl and Brich) had notable variation across sample points. We

discovered that for temporal beta diversity and its components (Brepl and Brich), AEM

was usually more important than the climate variable during the study period. The ratio

of climate to AEM for Btotal exhibits a unimodal pattern of increasing and then

decreasing. In contrast, the ratios for Brepl and Brich do not demonstrate a clear

elevation pattern (Fig. 4a, 4b, 4c). See Appendix S2 for further details.

We conducted a separate analysis of the changes in the relative importance of

climate and AEM with elevation. Our findings revealed that for Btotal, the importance

of climate exhibited a single-peaked pattern of increasing and then decreasing with

elevation. In contrast, for Brepl and Brich, there was no clear pattern of elevation in the

importance of climate (Fig. 4d, 4e, 4f). In contrast, for Brepl, the relative importance of

AEM exhibited an elevation pattern of decreasing and then increasing. In contrast, for

Btotal and Brich, the relative importance of AEM underwent a change, yet there was

no discernible elevation pattern (Fig. 4g, 4h, 4i).

Figure 4. Elevation patterns of the relative importance of different ecological processes

driving (a) temporal beta diversity (Btotal), (b) species replacement in temporal beta diversity

(Brepl), and (c) richness difference in temporal beta diversity (Brich). Elevation patterns of the

relative importance of climate driving (d) Btotal, (e) Brepl, and (f) Brich. Elevation patterns of

the relative importance of AEM driving (g) Btotal, (h) Brepl, and (i) Brich.

4 Discussion

We examined the association between the spatial pattern of climate fluctuations and the geographic pattern of community dynamics to attempt to explore the influence of elevation and the relative importance of deterministic and stochastic processes in community dynamics. Our hypothesis that climate fluctuations increase with altitude is partially confirmed. The results show that temperature fluctuation increases with elevation, but that the elevation pattern of precipitation fluctuation is in the opposite direction. We also found that the main influence on community dynamics was determined by stochastic processes, and the importance of deterministic versus stochastic processes exhibited a single-peak pattern with elevation.

4.1 Spatial variation of community dynamics

Temporal beta diversity gradually declined with increasing elevation, suggesting that macroinvertebrate community fluctuations gradually decreased and community stability increased. A growing body of evidence suggested that as species richness increased, the community became more stable, which is an example of the "portfolio effect", where the stability of the community increases with diversity. This finding has been extensively supported by studies of the diversity-stability ratio (DSR) of ecosystems [31]. For instance, with increasing species richness, the temporal synchronization of populations in a community diminishes, indicating that the community became more resistant to fluctuations and became more stable [32]. However, our findings revealed a bimodal pattern of species richness, with no

discernible trend of increase with elevation. Elevational patterns of species richness

demonstrated a diverse pattern as the study taxa and study area changed. However,

differences in elevational gradients in this study precluded a complete demonstration of

the bimodal pattern. The discrepancy in the results may be attributed to the influence of

species abundance. Communities with greater abundance generally have fewer

fluctuations in the community, and the distribution of population-abundance

heterogeneity also impacts DSR [33,34].

The results of this study indicated that temporal beta diversity was primarily driven

by species replacement, which is consistent with previous findings [35-37]. The

frequent turnover of macroinvertebrate species may be attributed to the high temporal

heterogeneity of the climate at the study site. Climate clearly played a crucial role in

the worldwide distribution of biodiversity [38]. The region's pronounced environmental

fluctuations have introduced novel temporal ecological niches for macroinvertebrates,

thereby promoting local species survival and facilitating species turnover [36,37,39].

The rapid turnover observed among macroinvertebrates often coincided with changes

in rare taxa [40-42], which are particularly sensitive and vulnerable to environmental

fluctuations [43,44]. The frequency of species occurrence during the study period is

detailed in Appendix S1. Additionally, macroinvertebrates possess unique life history

responses and evolutionary strategies that allow them to exploit these temporal

ecological niches, potentially leading to significant species turnover [37].

4.2 Inconsistent ecological processes across the landscape

The results revealed a single peak pattern regarding the relative importance of

deterministic and stochastic processes in the community dynamics that we observed in

our study area. With regard to total temporal beta diversity (Btotal), the significance of

deterministic processes undergoes a process of increasing and then decreasing

importance. In relation to the turnover component of the main driver of temporal beta

diversity (Brepl), the significance of stochastic processes exhibited a decreasing and

then increasing trend. This results may be closely related to climate fluctuations, which,

when within the tolerance range of organisms, filter species more and more strongly as

they are enhanced, leading to stronger and stronger effects on biotic communities [45].

Once a certain threshold is surpassed, heightened climatic fluctuations result in species

mortality, thereby resetting the community and introducing stochastic processes such

as stochastic colonization, which further underscores the importance of stochastic

processes [46]. Our study suggested that stochastic and deterministic processes jointly

influenced macroinvertebrate community dynamics. We found that stochastic processes

were the main drivers of community dynamics. Whereas ecological niche theory

highlights the influence of environmental filtering and species interactions on

communities [47], this study supports the neutral theory. In our study, the time variable

was the predominant factor driving the elevational patterns of macroinvertebrate

community dynamics, this may be related to the scale of our study. Deterministic

processes played an increasingly essential role as the scale of the study increased, as

documented by the spatial and temporal distribution of biodiversity [48,49]. Our

findings indicated that for Btotal, the ratio of deterministic to stochastic processes

followed a single-peaked model, similar to the behavior of the deterministic process.

The stochastic variables (AEMs in Appendix 1) were linked to stochastic ecological processes that include fluctuations in species abundance and richness among other characteristics. Temporary variation in species abundance can result in the generation of temporal stochastic processes by failure to detect rare species. Alternatively, sample sites with high total species numbers (temporal γ-diversity) throughout the study period were less likely to collect rare species and those rare species may not always be detected. However, our results suggested that the relative importance of deterministic and stochastic processes in macroinvertebrate community dynamics exhibited a single-peak pattern. This result was not surprising because biodiversity elevational patterns vary greatly worldwide.

Frequent extreme climate events, such as droughts affecting our rivers, have resulted in stochastic patterns of species extinction and colonization, albeit tempered by ecosystem resilience. The frequency, intensity, and periodicity of these events, including floods and droughts, significantly influence community dynamics [50]. These disturbances disrupt macroinvertebrate communities from returning to their optimal ecological niches, thereby reducing interspecific competition. Consequently, these disruptions may manifest as an amplification of stochastic processes, obscuring their spatial and temporal patterns. Community dynamics often exhibit temporal lags, where gains or losses projected in a given year are influenced by the community's preceding status. For example, population losses during unfavorable years may be compensated by gains in more favorable periods, thereby altering expected community dynamics

327   [47].

5 Conclusions

Amid the current era of heightened global climate variability, analyzing the spatial

trends of these fluctuations can offer valuable insights into the worldwide distribution

of biodiversity, such as migrations to higher elevations in search of refuges. We believe

that continued investigations will establish a solid theoretical basis for predicting the

geographic spread of community dynamics influenced by diverse ecological processes

amidst ongoing global transformations. Consequently, long-term biomonitoring

remains essential, particularly in regions vulnerable to climatic variability, such as

mountainous areas. Our findings emphasize the importance of distinguishing between

temporal community dynamics and observed spatial variations to enhance local

biodiversity management and conservation efforts across diverse landscapes.

For instance, shifts in community dynamics are intricately linked to changes in the

frequency, intensity, and periodicity of extreme events, which exhibit spatial

differentiation, including variations with elevation. Thus, further exploration of

seasonal and cyclical fluctuations in climatic patterns could significantly enhance the

predictability and effectiveness of regional conservation strategies through the

implementation of locally tailored practices for biodiversity protection.

References

348   1.   Liu, X.J.; Ferreira-Rodríguez, N.; Wu, R.W.; Ouyang, S.; Wu, X.P. Sixty years of species
349   diversity and population density decline of freshwater mussels in a global biodiversity

350   hotspot. *Global Ecology and Conservation* **2023**, *46*, doi:10.1016/j.gecco.2023.e02573.

2. Doxa, A.; Kamarianakis, Y.; Mazaris, A.D. Spatial heterogeneity and temporal stability characterize future climatic refugia in Mediterranean Europe. *Global Change Biology* **2022**, *28*, 2413-2424, doi:10.1111/gcb.16072.

3. Zhang, Z.; Hautier, Y.; Bao, T.J.; Yang, J.; Qing, H.; Liu, Z.L.; Wang, M.; Li, T.K.; Yan, M.; Zhang, G.L. Species richness and asynchrony maintain the stability of primary productivity against seasonal climatic variability. *Frontiers in Plant Science* **2022**, *13*, doi:10.3389/fpls.2022.1014049.

4. Jyvasjarvi, J.; Hamalainen, H. Profundal benthic invertebrate communities in boreal lakes vary with climate fluctuation. *Aquatic Sciences* **2015**, *77*, 261-269, doi:10.1007/s00027-014-0384-1.

5. Chan, W.P.; Chen, I.C.; Colwell, R.K.; Liu, W.C.; Huang, C.; Shen, S.F. Seasonal and daily climate variation have opposite effects on species elevational range size (vol 351, pg 1437, 2016). *Science* **2018**, *360*, doi:10.1126/science.aat9919.

6. Antunez, P. Evidence of the variation in the rate of change of temperature and precipitation. *Ecological Informatics* **2023**, *73*, doi:10.1016/j.ecoinf.2022.101928.

7. Kozlovsky, D.Y.; Branch, C.L.; Pitera, A.M.; Pravosudov, V.V. Fluctuations in annual climatic extremes are associated with reproductive variation in resident mountain chickadees. *Royal Society Open Science* **2018**, *5*, doi:10.1098/rsos.171604.

8. Whitenack, L.E.; Welklin, J.F.; Branch, C.L.; Sonnenberg, B.R.; Pitera, A.M.; Kozlovsky, D.Y.; Benedict, L.M.; Heinen, V.K.; Pravosudov, V.V. Complex relationships between climate and reproduction in a resident montane bird. *Royal Society Open Science* **2023**, *10*, doi:10.1098/rsos.230554.

9. Santillán, V.; Quitián, M.; Tinoco, B.A.; Zárate, E.; Schleuning, M.; Böhning-Gaese, K.; Neuschulz, E.L. Spatio-temporal variation in bird assemblages is associated with fluctuations in temperature and precipitation along a tropical elevational gradient. *Plos One* **2018**, *13*, doi:10.1371/journal.pone.0196179.

10. Wayman, J.P.; Sadler, J.P.; Pugh, T.A.M.; Martin, T.E.; Tobias, J.A.; Matthews, T.J. Assessing taxonomic and functional change in British breeding bird assemblages over time. *Global Ecology and Biogeography* **2022**, *31*, 925-939, doi:10.1111/geb.13468.

11. He, X.L.; Brown, C.; Lin, L.X. Relative importance of deterministic and stochastic processes for beta diversity of bird assemblages in Yunnan, China. *Ecosphere* **2021**, *12*, doi:10.1002/ecs2.3545.

12. Dini-Andreote, F.; Stegen, J.C.; van Elsas, J.D.; Salles, J.F. Disentangling mechanisms that mediate the balance between stochastic and deterministic processes in microbial succession. *Proceedings of the National Academy of Sciences of the United States of America* **2015**, *112*, E1326-E1332, doi:10.1073/pnas.1414261112.

13. Gao, J.; Liu, Y.H. Climate stability is more important than water-energy variables in shaping the elevational variation in species richness. *Ecology and Evolution* **2018**, *8*, 6872-6879, doi:10.1002/ece3.4202.

14. Bontemps, Z.; Moenne-Loccoz, Y.; Hugoni, M. Stochastic and deterministic assembly processes of microbial communities in relation to natural attenuation of black stains in Lascaux Cave. *Msystems* **2024**, *9*, doi:10.1128/msystems.01233-23.

15. Vanwonterghem, I.; Jensen, P.D.; Dennis, P.G.; Hugenholtz, P.; Rabaey, K.; Tyson, G.W.

Deterministic processes guide long-term synchronised population dynamics in replicate
anaerobic digesters. *Isme Journal* **2014**, *8*, 2015-2028, doi:10.1038/ismej.2014.50.

16.   Aguilar, P.; Sommaruga, R. The balance between deterministic and stochastic processes
in structuring lake bacterioplankton community over time. *Molecular Ecology* **2020**, *29*,
3117-3130, doi:10.1111/mec.15538.

17.   Cañedo-Argüelles, M.; Gutiérrez-Cánovas, C.; Acosta, R.; Castro-López, D.; Cid, N.;
Fortuño, P.; Munné, A.; Múrria, C.; Pimentao, A.R.; Sarremejane, R.; et al. As time goes by:
20 years of changes in the aquatic macroinvertebrate metacommunity of Mediterranean
river networks. *Journal of Biogeography* **2020**, *47*, 1861-1874, doi:10.1111/jbi.13913.

18.   Sanchez-Montoya, M.M.; Vidal-Abarca, M.R.; Punti, T.; Poquet, J.M.; Prat, N.; Rieradevall,
M.; Alba-Tercedor, J.; Zamora-Munoz, C.; Toro, M.; Robles, S.; et al. Defining criteria to
select reference sites in Mediterranean streams. *Hydrobiologia* **2009**, *619*, 39-54,
doi:10.1007/s10750-008-9580-0.

19.   Fick, S.E.; Hijmans, R.J. WorldClim 2: new 1-km spatial resolution climate surfaces for
global land areas. *International Journal of Climatology* **2017**, *37*, 4302-4315,
doi:10.1002/joc.5086.

20.   Hijmans, R.J.P., S.; Leathwick, J.; Elith, J. *dismo: Species Distribution Modeling*, R package
version 1.3-14; 2023.

21.   Blanchet, F.G.; Legendre, P.; Maranger, R.; Monti, D.; Pepin, P. Modelling the effect of
directional spatial ecological processes at different scales. *Oecologia* **2011**, *166*, 357-368,
doi:10.1007/s00442-010-1867-y.

22.   Baho, D.L.; Futter, M.N.; Johnson, R.K.; Angeler, D.G. Assessing temporal scales and
patterns in time series: Comparing methods based on redundancy analysis. *Ecological*
Complexity **2015**, *22*, 162-168, doi:10.1016/j.ecocom.2015.04.001.

23.   Blanchet, F.G.; Legendre, P.; Borcard, D. Modelling directional spatial processes in
ecological data. *Ecol. Model.* **2008**, *215*, 325-336, doi:10.1016/j.ecolmodel.2008.04.001.

24.   Dray, S.B., D.; Blanchet, G.; Borcard, D.; Clappe, S.; Guénard, G.; Jombart, T.; Larocque, G.;
Legendre, P.; Madi, N.; Wagner, H.H. *adespatial: Multivariate Multiscale Spatial Analysis*, R
package version 0.3-19.; 2022.

25.   Cardoso, P.M., S.; Rigal, F.; Carvalho, J. *BAT: Biodiversity Assessment Tools*, R package
version 2.9.2.; 2022.

26.   Ferrier, S.; Drielsma, M.; Manion, G.; Watson, G. Extended statistical approaches to
modelling spatial pattern in biodiversity in northeast New South Wales. II. Community-
level modelling. *Biodiversity and Conservation* **2002**, *11*, 2309-2338,
doi:10.1023/a:1021374009951.

27.   Ferrier, S.; Manion, G.; Elith, J.; Richardson, K. Using generalized dissimilarity modelling to
analyse and predict patterns of beta diversity in regional biodiversity assessment. *Diversity*
and Distributions **2007**, *13*, 252-264, doi:10.1111/j.1472-4642.2007.00341.x.

28.   Jones, M.M.; Ferrier, S.; Condit, R.; Manion, G.; Aguilar, S.; Perez, R. Strong congruence in
tree and fern community turnover in response to soils and climate in central Panama.
Journal of Ecology **2013**, *101*, 506-516, doi:10.1111/1365-2745.12053.

29.   Fitzpatrick, M.M., K.; Manion, G.; Nieto-Lugilde, D.; Ferrier, S. *gdm: Generalized*
Dissimilarity Modeling, R package version 1.5.0-9.1.; 2022.

30.   Borcard, D.; Legendre, P.; Drapeau, P. Partialling out the spatial component of ecological

438   variation. *Ecology* **1992**, *73*, 1045-1055, doi:10.2307/1940179.

439 31. Campbell, V.; Murphy, G.; Romanuk, T.N. Experimental design and the outcome and
440   interpretation of diversity-stability relations. *Oikos* **2011**, *120*, 399-408,
441   doi:10.1111/j.1600-0706.2010.18768.x.

442 32. Roscher, C.; Weigelt, A.; Proulx, R.; Marquard, E.; Schumacher, J.; Weisser, W.W.; Schmid,
443   B. Identifying population- and community-level mechanisms of diversity-stability
444   relationships in experimental grasslands. *Journal of Ecology* **2011**, *99*, 1460-1469,
445   doi:10.1111/j.1365-2745.2011.01875.x.

446 33. Thibaut, L.M.; Connolly, S.R. Understanding diversity-stability relationships: towards a
447   unified model of portfolio effects. *Ecology Letters* **2013**, *16*, 140-150,
448   doi:10.1111/ele.12019.

449 34. Kilpatrick, A.M.; Ives, A.R. Species interactions can explain Taylor's power law for ecological
450   time series. *Nature* **2003**, *422*, 65-68, doi:10.1038/nature01471.

451 35. Angeler, D.G. Revealing a conservation challenge through partitioned long-term beta
452   diversity: increasing turnover and decreasing nestedness of boreal lake metacommunities.
453   *Diversity and Distributions* **2013**, *19*, 772-781, doi:10.1111/ddi.12029.

454 36. Tonkin, J.D.; Bogan, M.T.; Bonada, N.; Rios-Touma, B.; Lytle, D.A. Seasonality and
455   predictability shape temporal species diversity. *Ecology* **2017**, *98*, 1201-1216,
456   doi:10.1002/ecy.1761.

457 37. Wu, N.C.; Wang, Y.C.; Wang, Y.X.; Sun, X.M.; Faber, C.; Fohrer, N. Environment regimes
458   play an important role in structuring trait- and taxonomy-based temporal beta diversity
459   of riverine diatoms. *Journal of Ecology* **2022**, *110*, 1442-1454, doi:10.1111/1365-
460   2745.13859.

461 38. Jocque, M.; Field, R.; Brendonck, L.; De Meester, L. Climatic control of dispersal-ecological
462   specialization trade-offs: a metacommunity process at the heart of the latitudinal diversity
463   gradient? *Global Ecology and Biogeography* **2010**, *19*, 244-252, doi:10.1111/j.1466-
464   8238.2009.00510.x.

465 39. Heino, J.; Tolonen, K.T. Ecological drivers of multiple facets of beta diversity in a lentic
466   macroinvertebrate metacommunity. *Limnol. Oceanogr.* **2017**, *62*, 2431-2444,
467   doi:10.1002/lno.10577.

468 40. Mori, T.; Murakami, M.; Saitoh, T. Latitudinal gradients in stream invertebrate assemblages
469   at a regional scale on Hokkaido Island, Japan. *Freshwater Biology* **2010**, *55*, 1520-1532,
470   doi:10.1111/j.1365-2427.2009.02363.x.

471 41. Castro, D.M.P.; da Silva, P.G.; Solar, R.; Callisto, M. Unveiling patterns of taxonomic and
472   functional diversities of stream insects across four spatial scales in the neotropical savanna.
473   *Ecol. Indic.* **2020**, *118*, 8, doi:10.1016/j.ecolind.2020.106769.

474 42. Rocha, M.P.; Bini, L.M.; Gronroos, M.; Hjort, J.; Lindholm, M.; Karjalainen, S.-M.; Tolonen,
475   K.E.; Heino, J. Correlates of different facets and components of beta diversity in stream
476   organisms. *Oecologia* **2019**, *191*, 919-929, doi:10.1007/s00442-019-04535-5.

477 43. Nyirabuhoro, P.; Liu, M.; Xiao, P.; Liu, L.M.; Yu, Z.; Wang, L.N.; Yang, J. Seasonal Variability
478   of Conditionally Rare Taxa in the Water Column Bacterioplankton Community of
479   Subtropical Reservoirs in China. *Microbial Ecology* **2020**, *80*, 14-26, doi:10.1007/s00248-
480   019-01458-9.

481 44. Monaghan, M.T.; Robinson, C.T.; Spaak, P.; Ward, J.V. Macroinvertebrate diversity in

fragmented Alpine streams: implications for freshwater conservation. *Aquatic Sciences*
2005, *67*, 454-464, doi:10.1007/s00027-005-0787-0.

45.   Vorste, R.V.; Corti, R.; Sagouis, A.; Datry, T. Invertebrate communities in gravel-bed,
braided rivers are highly resilient to flow intermittence. *Freshwater Science* **2016**, *35*, 164-
177, doi:10.1086/683274.

46.   Chase, J.M. Community assembly: when should history matter? *Oecologia* **2003**, *136*, 489-
498, doi:10.1007/s00442-003-1311-7.

47.   Chesson, P. Mechanisms of maintenance of species diversity. *Annual Review of Ecology*
and Systematics **2000**, *31*, 343-366, doi:10.1146/annurev.ecolsys.31.1.343.

48.   Pajunen, V.; Luoto, M.; Soininen, J. Climate is an important driver for stream diatom
distributions. *Global Ecology and Biogeography* **2016**, *25*, 198-206,
doi:10.1111/geb.12399.

49.   Whittaker, R.J.; Willis, K.J.; Field, R. Scale and species richness: towards a general,
hierarchical theory of species diversity. *Journal of Biogeography* **2001**, *28*, 453-470,
doi:10.1046/j.1365-2699.2001.00563.x.

50.   Chiu, M.C.; Ao, S.C.; Resh, V.H.; He, F.Z.; Cai, Q.H. Species dispersal along rivers and
streams may have variable importance to metapopulation structure. *Science of the Total*
Environment **2021**, *760*, doi:10.1016/j.scitotenv.2020.144045.