1 **Microtubule depolymerization at kinetochores restricts anaphase spindle**
2 **elongation**
3

4 Geng-Yuan Chen[1], Changfeng Deng[2], David M. Chenoweth[2], Michael A. Lampson[1]*
5
6 [1]Department of Biology, University of Pennsylvania, Philadelphia, PA, 19104
7 [2]Department of Chemistry, University of Pennsylvania, Philadelphia, PA, 19104
8
9 *Corresponding author. Email: lampson@sas.upenn.edu

10
11

Abstract

Anaphase chromosome segregation depends on forces exerted by spindle
microtubules. Current models propose two mechanisms of force generation:
kinetochore microtubules (kMTs) depolymerize to pull chromosomes toward the spindle
poles (anaphase A), while antiparallel microtubule sliding in the central spindle further
separates the chromosomes by elongating the spindle (anaphase B). Experimental
evidence in cells supports the sliding mechanism, but contributions of the
depolymerization mechanism remain unclear. Here we show that kMT depolymerization
limits spindle elongation rather than moving chromosomes apart. We developed a
chemical optogenetic approach to recruit a microtubule depolymerase to kinetochores
at anaphase onset, thereby increasing the rate of kMT depolymerization without
perturbing earlier stages of mitosis. We find that increased depolymerization slows the
velocity at which spindle poles move apart without changing kinetochore separation
velocities. Our findings support a model in which kinetochores selectively couple to
central spindle microtubules parallel to their kMTs, so that antiparallel sliding drives
chromosome segregation while kMT depolymerization pulls poles inward.

Introduction

During cell division, sister chromosomes segregate in anaphase to partition the two
replicated genomes into daughter cells. Three mechanisms of force generation in
anaphase have been described. First, kinetochore-microtubule (kMT) bundles
depolymerize to pull chromosomes and spindle poles toward each other. Second,
antiparallel microtubule sliding in the central spindle pushes the two half-spindles apart.
Third, astral microtubules pull spindle poles apart. Connections between these different
microtubule populations make it difficult to dissect the contributions of potential force-
generating mechanisms to chromosome segregation.[1]
Current "chromosome-pulled" models propose that depolymerizing kMTs pull each
chromosome toward its attached pole (anaphase A) (**Fig. 1a**).[2-4] As chromosomes are
pulled outward, antiparallel microtubule sliding pushes poles apart or pushes kMTs
apart to elongate the spindle (anaphase B).[5,6] Astral microtubules can also pull poles
apart, but antiparallel sliding is thought to be the primary mechanism driving anaphase
B in human cells.[6] Molecular motors slide antiparallel microtubules apart *in vitro*,
mimicking spindle elongation,[7-9] and laser ablation of the central spindle between sister

kinetochores in early anaphase slows kinetochore separation.[10,11] Furthermore,
manipulations that increase or decrease antiparallel sliding velocities affect both
anaphase spindle elongation and chromosome segregation velocities.[12-18] These
findings demonstrate the importance of antiparallel sliding for chromosome segregation.
In contrast, although depolymerizing microtubules can move tip-coupled objects
(chromosomes or beads) in vitro,[3,4] the contribution of kMT depolymerization to
anaphase chromosome segregation remains unclear. Manipulations that slow kMT
depolymerization in cells, by inhibiting microtubule depolymerase activity, have the
expected effect of decreasing the velocity of kinetochores and poles moving toward
each other (anaphase A). However, chromosome segregation velocities have not been
reported in these conditions.[19-24] In addition, depolymerases localize to microtubules
beyond kMTs, so inhibition can affect multiple populations of spindle microtubules.[4]
In an alternative model, the central spindle is coupled to chromosomes so that
antiparallel sliding directly drives anaphase chromosome segregation.[25] Neither kMT
depolymerization (anaphase A) nor spindle elongation (anaphase B) is necessary for
chromosome movements in this model. The spindle elongates because poles are
attached to kMTs, so sliding moves poles outward, but chromosome movements are
independent of spindle poles. We refer to this model as the "pole-pulled" model because
kMT depolymerization at either kinetochores (plus-ends) or poles (minus-ends) pulls
poles inward toward the chromosomes (**Fig. 1a**).[25] Spindle elongation is thus a
combination of outward forces on spindle poles from antiparallel sliding and astral
microtubules and inward forces from kMT depolymerization and potentially central
spindle depolymerization. For example, if kMT depolymerization is fast enough,
chromosomes can segregate without spindle elongation as in *C. elegans* meiosis.[26]
To distinguish between the chromosome-pulled and pole-pulled models, we
designed a chemical optogenetic assay to acutely increase microtubule
depolymerization at kinetochores in anaphase in human cells. We show that increased
kMT depolymerization restricts pole separation without affecting chromosome
separation, consistent with the pole-pulled model. We propose a molecular model in
which each kinetochore selectively binds to the sides of central spindle microtubules in
parallel orientation to its kMTs. Antiparallel sliding moves central spindle microtubules of
opposite orientations apart, along with their bound kinetochores. As kMTs
depolymerize, poles are pulled inward toward kinetochores anchored to the central
spindle. To test this model, we developed additional chemical optogenetic assays to
manipulate spindle checkpoint signaling and antiparallel sliding in the central spindle.
We inhibited the checkpoint to induce anaphase on monopolar spindles.[27] Some sister
kinetochore pairs localize near antiparallel microtubule bundles and move away from
other in anaphase, consistent with antiparallel sliding driving their movements. We also
recruited a microtubule-binding protein to the central spindle in anaphase to suppress
antiparallel sliding. We find reduced kMT depolymerization, consistent with kinetochore
linkage to the central spindle to constrain plus-end depolymerization. Overall, these
results support our model in which kinetochores are mechanically coupled to central
spindle microtubules oriented parallel to their kMTs, so that antiparallel sliding moves
sister chromosomes apart while kMT depolymerization pulls poles inward.

89
Results
kMT depolymerization regulates spindle elongation
The chromosome-pulled model predicts that increasing kMT depolymerization in
anaphase should increase kinetochore separation velocity, with no effect on pole
separation velocity. Conversely, the pole-pulled model predicts decreased pole
separation velocity as poles are pulled inward, with no effect on kinetochore separation
velocity. To distinguish between these models, we designed an assay to increase the
rate of microtubule depolymerization at kinetochores at anaphase onset. We recruited
the motor domain of the microtubule depolymerase MCAK (Mitotic Centromere-
Associated Kinesin; kinesin-13) to kinetochores using a photocaged small molecule that
heterodimerizes HaloTag and *Escherichia coli* dihydrofolate reductase (eDHFR) fusion
proteins.[28-30] We fused HaloTag to the kinetochore protein SPC25 (the anchor) and
fused eDHFR to the MCAK motor domain (the effector), so that uncaging leads to
dimerization and MCAK recruitment to kinetochores (**Fig. 1b-d**). This chemical
optogenetic approach provides temporal precision by light activation and molecular
specificity by using genetically tagged constructs.[31] By activating immediately after
anaphase onset, our manipulations do not affect earlier stages of spindle formation and
kinetochore attachment. As MCAK is active only at microtubule ends,[32] we expect that
recruitment of the motor domain to kinetochores increases depolymerization of kMTs
but not other proximal microtubules in the central spindle.
We first tested the kMT depolymerization assay in metaphase spindles, in which
cohesion prevents sister kinetochores from moving farther apart. We find that MCAK
recruitment leads to rapid spindle shortening, as poles move inward toward the
kinetochores within 2 min of uncaging (**Fig. 1e-f, Supplementary Video 1**). This result
indicates that MCAK recruitment increases kMT depolymerization as expected, and that
kinetochores remain coupled to their attached kMTs.
To determine the effects of increased kMT depolymerization in anaphase, we
recruited MCAK to kinetochores within 1 min after sister kinetochores start to move
apart, indicating anaphase onset (**Fig. 2a-b, Supplementary Video 2**). We measured
three distances at each time point: between separating kinetochores, between poles,
and between kinetochores and their attached pole (**Fig. 2c**). For each cell, we fit a slope
to the earliest time window exhibiting a linear range for all distances to measure
velocities of pole separation (V_{PP}), kinetochore separation (V_{KK}), and kMT
depolymerization (V_{PK}) (**Fig. 2d**). MCAK overexpression alone does not affect any of
these velocities (**Supplementary Fig. 1**), and MCAK recruitment to kinetochores
increases the rate of kMT depolymerization, as expected (V_{PK}, from 0.52 ± 0.03 μm/min
to 0.77 ± 0.04 μm/min, mean $\pm$ SEM, **Fig. 2e**). Pole separation velocity decreases (V_{PP},
from 1.07 ± 0.08 μm/min to 0.40 ± 0.10 μm/min, mean $\pm$ SEM) with increased kMT
depolymerization, but kinetochore separation velocity is not affected (**Fig. 2f-g**). The
decrease in V_{PP} is double the increase in V_{PK}, which represents only half of the spindle,
indicating that poles are pulled inward toward the kinetochores by depolymerizing kMTs
as in metaphase (**Fig. 1e-f**). We conclude that kMT depolymerization restricts spindle
elongation rather than moving chromosomes, consistent with the pole-pulled model.

To explain both outward forces on chromosomes and inward forces on poles, we
propose a "sort and grip" model: each kinetochore sorts central spindle microtubules
and selectively grips (binds tightly to the sides of) those in parallel orientation to the
kMTs (**Fig. 2h**). As a result, antiparallel sliding moves kinetochores and their attached
kMTs outward, while kMT depolymerization pulls poles inward relative to the central
spindle where kinetochores are anchored. Forces exerted by kMT plus-end
depolymerization rely on the well-established ability of kinetochores to maintain
attachments to depolymerizing kMT plus-ends.[3] Sorting can be mediated by molecular
motors, crosslinkers, or branching microtubule arrays that sort microtubules for the
same polarity,[33,34] so that kinetochores selectively bind parallel central spindle
microtubules based on proximity.
144
Anti-poleward motions in monopolar anaphase
To test the sort-and-grip model, we designed two additional chemical optogenetic
assays to manipulate spindle checkpoint signaling and antiparallel sliding in the central
spindle. First, we initiated anaphase on monopolar spindles by inhibiting the spindle
assembly checkpoint using conditional pharmacology. The sort-and-grip model predicts
that kinetochore motions depend on the arrangement of kMTs and adjacent antiparallel
bundles. For some sister kinetochore pairs, kMTs orient in opposite directions on a
monopolar spindle (**Fig. 3a**).[35] In this configuration, antiparallel sliding would move
sister kinetochores away from each other rather than moving both sisters toward the
pole. To induce anaphase, we used a photocaged inhibitor of the checkpoint kinase
MPS1 (Monopolar Spindle 1) linked to the Halo ligand, which binds covalently to the
Halo-tagged kinetochore anchor SPC25 (**Fig. 3b**).[36] MPS1 is inhibited only after
uncaging with a pulse of light. As the inhibitors are bound to SPC25 and do not diffuse
between cells, local uncaging triggers anaphase in one cell at a time. This approach
allows sequential live imaging of individual cells with high temporal resolution.
After MPS1 inhibition, we examined localization of the antiparallel microtubule
crosslinker PRC1 (Protein Regulator of Cytokinesis 1), which marks antiparallel
bundles.[14,37,38] PRC1 signals gradually decrease in length and increase in intensity,
indicating antiparallel sliding that decreases the overlap length in each bundle
(**Supplementary Fig. 2**), as observed in the central spindle in normal bipolar
anaphase[14,37,38] and in reconstituted systems.[39-41] We analyzed kinetochore motions
separately for sister pairs either colocalized to (on-bundle) or away from (off-bundle)
PRC1 signals (**Fig. 3c-f, Supplementary Videos 3-4**). Off-bundle pairs exhibit minimal
movement of sisters away from each other, consistent with a lack of separation forces
by antiparallel sliding. Conversely, some on-bundle sisters move apart, with the distal
kinetochore of these pairs moving away from the pole (**Fig. 3g**). This result is consistent
with the sort-and-grip model, in which antiparallel sliding drives chromosome
movements depending on local spindle organization near kinetochores, independent of
spindle poles. As monopolar spindles are generated by inhibiting kinesin-5 motors,
antiparallel sliding is likely due to kinesin-4 (KIF4A) motors recruited to PRC1 bundles.[15]
Previous findings, using laser ablation to detach one kinetochore from its attached pole

in a bipolar spindle, also show sister kinetochore separation consistent with the sort-
and-grip model.[11]
178
Suppressed central spindle sliding inhibits kMT depolymerization
In the sort-and-grip model, kMT plus ends are coupled to the central spindle via
side-attached kinetochores. Without antiparallel sliding to move kinetochores outward,
kMT depolymerization would require shortening the spindle by pulling poles inward,
against outward forces exerted on poles by astral MTs. Thus, we predict that
suppressed antiparallel sliding would reduce kMT depolymerization because both kMT
plus and minus ends are constrained.
To suppress antiparallel sliding, we generated "brake" complexes in the central
spindle. We fused HaloTag to full-length dimeric PRC1 (the anchor) and fused eDHFR
to a monomeric PRC1 truncation mutant (PRC1M, the effector) that binds to
microtubules without crosslinking activity on its own. Uncaging generates brake
complexes by recruiting PRC1M to dimeric PRC1 bound to antiparallel bundles, which
should strengthen PRC1 binding to microtubules and thus increase friction to suppress
sliding within the bundle (**Fig. 4a-b**). Before uncaging, PRC1M binds to microtubules
proportional to local microtubule density, with a subtle enrichment around the equator
(**Fig. 4c, Supplementary Video 5**). This enrichment increases after activation,
consistent with recruitment to PRC1-labeled antiparallel bundles in the central
spindle.[14,37,38]
PRC1 itself acts as a molecular brake, as it restricts microtubule sliding *in vitro,*[39-41]
and its depletion in cells increases the separation velocity of the two half-spindles.[14,42]
Thus, we predict increased friction and suppressed antiparallel sliding due to PRC1
overexpression alone, followed by a further increase in friction after uncaging and
PRC1M recruitment at anaphase onset. We measured kinetochore and pole velocities
and kMT depolymerization as in Fig. 2c-d. We find that PRC1 overexpression
decreases pole and kinetochore separation velocities, proportional to PRC1 levels on
spindles, and PRC1M recruitment further decreases these velocities (**Fig. 4d-e**,
Supplementary Fig. 3a-c, Supplementary Video 6). These results indicate
progressively greater suppression of antiparallel sliding as expected. The rate of kMT
depolymerization decreases as antiparallel sliding is suppressed, with the slowest rate
observed in the presence of both PRC1 overexpression and PRC1M recruitment (**Fig.**
4f, Supplementary Fig. 3d). This result is consistent with the sort-and-grip model. The
plus-ends of kMTs are coupled to the central spindle, while the minus-ends are
anchored at the poles and constrained by astral MTs, so kMTs are not free to
depolymerize.
213
Discussion
Using chemical tools to manipulate spindle organization and specific microtubule
populations within the spindle, our experiments support a model in which antiparallel
sliding directly drives anaphase chromosome segregation while kMT depolymerization
limits spindle elongation. We first show that enhanced kMT depolymerization slows pole
separation without increasing kinetochore separation (**Fig. 2**). Based on these findings,

we propose a sort-and-grip model (**Fig. 5**): parallel microtubule sorters position
kinetochores close to central spindle microtubules oriented parallel to the kMTs, so that
kinetochores selectively grip these central spindle microtubules upon anaphase onset. If
sorting is established before anaphase onset, then sorters can be absent in anaphase
or allow sliding of kMTs relative to the central spindle. Kinetochores grip tightly in
anaphase, however, without sliding on the central spindle, which has two key outcomes.
First, antiparallel sliding in the central spindle moves sister kinetochores apart together
with their attached kMTs. Second, kMT plus-end depolymerization pulls kMTs and poles
inward toward kinetochores that are tightly coupled to the central spindle, against
outward movement of kMTs driven by anti-parallel sliding. Depolymerization of kMT
minus-ends also pulls poles inward in this model. Our results from two additional assays
support the sort-and-grip model. First, sister kinetochores can move apart on monopolar
spindles, consistent with antiparallel sliding moving sorted microtubules together with
their gripped kinetochores (**Fig. 3**). Second, suppressed antiparallel sliding slows kMT
depolymerization, indicating that kinetochores couple to the central spindle (**Fig. 4**).
Velocity measurements in multiple model systems are consistent with the sort-and-
grip model. In human cells, *C. elegans* oocytes and embryos, and *Xenopus* egg
extracts, antiparallel sliding velocity within the central spindle matches kinetochore
velocity in anaphase (**Supplementary Fig. 4a**)[10,43], suggesting that antiparallel sliding
moves chromosomes without requiring a contribution from kMT depolymerization.
Sliding velocity also correlates with poleward movements of kMTs.[11,15] In addition,
slowing kMT depolymerization, by depleting microtubule depolymerases or severing
enzymes in human cells or adding non-degradable cyclin B to *Xenopus* egg extracts,
increases spindle elongation velocity.[20,24,44] These results are consistent with kMT
depolymerization pulling poles inward and with our findings that increasing kMT plus-
end depolymerization slows spindle elongation (**Fig. 2**).
Microtubule severing experiments in multiple systems are also consistent with the
sort-and-grip model. Cutting the central spindle early in anaphase transiently halts
kinetochore separation in human cells and *C. elegans* oocytes and embyros,[10] whereas
kinetochores continue to separate after ablating centrosomes or microtubules midway
between kinetochores and their attached poles in human, *C. elegans*, rat-kangaroo,
newt, grasshopper, and crane-fly cells.[10,11,45-50] These findings indicate that antiparallel
sliding directly moves chromosomes. Structurally, central spindle microtubules are close
to kinetochores and their attached kMTs in high-resolution optical and EM images,[10,51,52]
and kMT perturbations by microneedle manipulation or microtubule severing suggest
crosslinking to the central spindle near chromosomes.[11,53-55] Kinetochores can thus
directly bind to the central spindle or interact with microtubule-associated proteins on
the central spindle to make the coupling robust.
To move chromosomes toward their attached poles correctly, the sort-and-grip
model relies on kinetochores attaching selectively to the sides of central spindle
microtubules oriented parallel to their kMTs (**Fig. 5**). This selectivity can be mediated by
microtubule sorting mechanisms acting earlier in mitosis that position kMTs adjacent to
parallel microtubules in the central spindle. In prometaphase, short microtubules bind
kinetochores at their plus-ends, while the minus-ends interact with antiparallel bundles

in the central spindle.[56] Minus-end directed motors sort these nascent kMTs and
establish parallel coupling to central spindle microtubules to biorient sister kinetochores
(**Supplementary Fig. 4b**).[57] Alternatively, minus-end directed motors also sort newly-
grown microtubules in the central spindle poleward through kMTs (**Supplementary Fig.**
4c).[58] Furthermore, parallel microtubule arrays can assemble by growing microtubules
from existing microtubule templates.[33,59-61] Microtubules nucleated from sister kMTs can
thus grow toward the equator to generate a "bridging fiber" by antiparallel crosslinking
(**Supplementary Fig. 4d**).[11] We propose that parallel microtubules sorted before
anaphase persist so that kinetochores couple to these microtubules and move by
antiparallel sliding in anaphase. Inhibition or depletion of potential candidates for these
sorting mechanisms (e.g. augmin,[62] NuMA,[63] ZW10,[64] and kinesin-14[65]) commonly
causes anaphase chromosome segregation errors, which can arise from incorrect
kinetochore coupling to microtubules antiparallel to their kMTs.
Restricting spindle elongation may be important for later stages of division, when
excessive spindle elongation is associated with midzone buckling and defocused
cytokinetic furrows.[14,37,38,66-70] In addition, spindle poles act as hubs for numerous
biochemical and mechanical cues,[71-73] so restricted spindle elongation may regulate
pole communication with spindle midbodies or the cytokinetic furrow. Hence, telophase
biology likely requires coordination between pole separation and chromosome
separation velocities via depolymerizing kMTs.
284
Methods
Synthesis, characterization, and storage of chemical optogenetic probes. The
photoactivatable dimerizer (CTH)[29,30] and photoactivatable localized MPS1 inhibitor
(CRH)[36] were prepared as previously described. Both compounds were dissolved in
DMSO at 20 mM and stored as 1 μL aliquots in amber plastic microcentrifuge tubes at -
80 °C.
291
292 **Small molecule inhibitors.** All stocks were dissolved in DMSO and stored at -20 °C.
293 Stock concentrations include 20 mM S-Trityl-L-Cysteine (STLC, Eg5 inhibitor; Sigma
294 Aldrich) and 10 mM Verapamil (Tocris). Fresh aliquots were used for all experiments.
295
Plasmids. The pEM705 backbone with a CAG promoter was used to clone all plasmids
for anchors and effectors.[74] The kinetochore anchor SPC25, fused to three tandem
copies of HaloTag and an eGFP on its N-terminus (3×Halo-GFP-SPC25), was used as
previously described.[29,30] To manipulate central spindle mechanics, PRC1 was used as
an anchor, fused to 3× HaloTag with the far-red fluorescent reporter, miRFP670nano3
(hereafter denoted 670nano3), on its N-terminus (3×Halo-670nano3-PRC1).
All effectors were designed based on the construct with the red fluorescent protein
mScarlet fused to the N terminus of *Escherichia coli* dihydrofolate reductase DHFR
(mScarlet-eDHFR). For targeted microtubule depolymerization, the phospho-resistant
motor domain of human MCAK (MCAK$_{179-583;S192A}$) was fused to the N terminus of
mScarlet-eDHFR (MCAK-mScarlet-eDHFR).[30,75] To suppress microtubule sliding, a

monomeric PRC1 mutant (PRC1M, PRC1$_{341-620}$)[76] was fused to the C terminus of
mScarlet-eDHFR to make mScarlet-eDHFR-PRC1M.
To label spindle poles, a single copy or two tandem copies of the PACT (Pericentrin
and AKAP Centrosome Targeting)[77] domain was fused to eGFP (PACT-GFP or
2×PACT-GFP) in the pcDNA3.1 backbone containing cytomegalovirus and T7
promoters. To label the central spindle, full-length PRC1 was fused to two tandem
copies of 670nano3 (2×670nano3-PRC1), using pEM705 as the backbone.
314
Cell cultures and transfection. HeLa RMCE (recombination-mediated cassette
exchange) acceptor cells were passaged at 37°C in the growth medium containing
DMEM (Gibco) supplemented with 10% Fetal Bovine Serum (Clontech) and 1%
penicillin/streptomycin (Life Technologies), within 5% CO_2 atmosphere and humidity.
For assays, cells were cultured on coverslips (Bellco) coated by poly-D-lysine (Sigma-
Aldrich) for at least 16 h, and then transfected with the desired constructs using
Lipofectamine 3000 (Invitrogen) in Opti-MEM (Gibco) for at least 24 h.
Manipulations of mechanics and biochemistry at kinetochores were conducted with
cells stably expressing 3×Halo-GFP-SPC25, as previously described.[29,30] For the
monopolar anaphase assay, cells were transfected with 0.5-1 μg 2×670nano3-PRC1
and 250 ng PACT-GFP to mark the central spindle and spindle poles, respectively. For
kMT depolymerization, cells were transfected with 1-1.5 μg MCAK-mScarlet-eDHFR
and 250 ng 2×PACT-GFP.
Manipulations of central spindle mechanics were conducted with cells transiently
transfected with 0.6 μg Scarlet-eDHFR-PRC1M and 0.6 μg 3×Halo-670nano3-PRC1,
together with 100 ng CenpB-GFP and 250 ng PACT-GFP for labeling centromeres and
poles, respectively.
332
Photoactivation and dimerization. For recruitment experiments, cells were placed in a
magnetic chamber (Chamlide CM-S22-1; LCI) and incubated with growth medium and
CTH for 1 h. Unbound CTH was washed out by incubating for 20 min in imaging
medium containing L15 without phenol red (Gibco) (imaging medium) plus 10% FBS
and 1% penicillin/streptomycin. The medium was then replaced with fresh imaging
medium, and the chamber was moved to a stage in a 37°C environmental chamber
(PeCon). For dimerization experiments, cells were imaged live to determine the time of
anaphase onset. CTH was activated within 1 min after sister kinetochores started to
move apart, using a 405-nm laser (CrystaLaser LC) controlled by iLas2 software (Roper
Scientific) installed on MetaMorph (Molecular Devices), as previously described.[30]
Activation was performed over the entire cell at 25% laser power with 5 repetitions. To
induce anaphase in monopolar cells, cells were incubated with 10 μM Verapamil (an
efflux pump inhibitor) and 20 μM STLC in growth medium for 2 h, then 20 μM CRH was
added for 1 h. Unbound CRH was washed out by incubating in imaging medium with
STLC for 30 min. CRH was uncaged by a 5-second light pulse using a mercury arc
lamp filtered at 385/11 nm, and imaging started 15-20 min later. Procedures with light-
sensitive molecules were performed under red light in the room.
350

Image acquisition. Live cells at 37 °C were imaged with a confocal microscope (DM4000; Leica), equipped with a 100× 1.4 NA oil immersion objective (Leica), an XY Piezo-Z stage (Applied Scientific Instrumentation), a CSU10 spinning disk (Yokogawa), an electron multiplier charge-coupled device camera (ImageEM; Hamamatsu Photonics), and a 4-line laser module (405 nm, 488 nm, 561 nm, and 639 nm; Vortran Stradus VersaLase 4), as previously described.[30] Images were acquired with a 10-60s time interval, with 0.5-2 μm spacing for z-stacks covering 6-15 μm total. For dimerization experiments, cells were activated after image acquisition at t=0. For monopolar anaphase assay, images were acquired 15 minutes after uncaging.

Data analyses and statistical tests.

For monopolar anaphase cells, both kinetochores and poles were tracked by the Fiji::MTrackJ plug-in. In each frame, the distances of two sisters relative to the pole were calculated (D_{PK}). The distance difference between the two sisters (ΔD_{PK}) represents the inter-kinetochore distance along the axis between the pole and kinetochores (D_{KK}). For each trajectory, the maximum kinetochore separation velocity (max V_{KK}) represents the maximal displacement of inter-kinetochore distance (D_{KK}) across all possible one-minute time windows. A cutoff was defined as the 95% percentile of max V_{KK} of off-bundle sister kinetochore pairs.

For bipolar anaphase cells, the coordinates of kinetochores and spindle poles were measured by the Fiji::TrackMate and Manual Tracking plug-ins, respectively. Kinetochore coordinates were projected to the mitotic axis defined by poles, and then grouped based on their closest pole. In each frame, the median position of each group represents the centroid of projected kinetochores. Initial velocities were defined by the time window exhibiting linear displacement between the two poles (V_{PP}), between the two kinetochore centroids (V_{KK}), and between a kinetochore centroid and its corresponding pole ($-V_{PK}$). Cells were included in the analysis only if they exhibit constant kinetochore separation velocity at early time points, representing complete loss of cohesion and separation of sister chromosomes. To measure PRC1 intensity on a bipolar spindle, image stacks were mean-projected along z. Spindle-bound PRC1 signal was defined by the intensity within the minimal circle encompassing both spindle poles, with cytosolic intensity subtracted as background, summed for each cell. For each set of experiments, PRC1 intensities across different perturbations were normalized relative to the mean intensity of cells expressing 3×Halo-670nano3-PRC1.

Data from measurements were analyzed by MATLAB. Two-tailed Student's t-test was used to compare measurements between groups.

Acknowledgments: We thank the Philly ChromoClub and the Penn Center for Genome Integrity for discussions. This work was supported by the NIH (GM122475 and P01-CA265794) and the Basser Center for BRCA.

Author contributions: GYC designed and conducted cell biology experiments and wrote the manuscript. CD and DMC synthesized and characterized chemical optogenetic probes. MAL designed experiments and edited the manuscript.

Competing interests: The authors claim no competing financial interests.

Materials & Correspondence. Cell lines, chemicals, and protocols are available upon request. The corresponding authors adhere to the NIH Grants Policy and Sharing of Unique Research Resources. Correspondence and requests for materials should be addressed to MAL (lampson@sas.upenn.edu).

Supplementary Information is available for this paper.

Data availability. The datasets generated and analyzed during the current study are available from the corresponding author on reasonable request without restriction.

Figure 1. Induced kMT depolymerization shortens metaphase spindle. (a)
Mechanical models for anaphase chromosome segregation. Polarity of antiparallel
microtubules in the central spindle (blue), kMTs (red), and astral microtubules (gray) is
indicated, with minus-ends (–) and plus-ends (+) facing toward and away from poles,
respectively. In the chromosome-pulled model, antiparallel sliding in the central spindle
pushes poles or kMTs apart (blue arrows), while kMT depolymerization pulls each
chromosome outward toward its attached pole (red arrows). In the pole-pulled model,
the central spindle couples to chromosomes, so antiparallel sliding directly moves sister
chromosomes apart (blue arrows), and kMT depolymerization pulls poles inward toward
the chromosomes (red arrows). **(b)** Schematic of light-inducible dimerization. The
photocaged dimerizer (CTH) includes a Halo ligand (green) linked to the eDHFR ligand
trimethoprim (TMP, red) and a photocage (purple). The dimerizer is activated by
uncaging to make TMP accessible to eDHFR, recruiting the eDHFR-tagged effector to
the Halo-tagged anchor. **(c)** Schematic of kMT shortening assay in metaphase. The
microtubule depolymerase MCAK (orange) is recruited to kinetochores (green) after
dimerizer uncaging at t=0 to enhance kMT depolymerization. **(d)** Constructs for kMT
shortening assay. The minimal microtubule depolymerization domain (MCAK$_{179-583}$) was
used, with a mutation to prevent phosphorylation: S192A. SPC25 is the kinetochore
anchor, and PACT labels spindle poles. **(e-f)** Metaphase kMT shortening assay for cells
expressing Halo-GFP-SPC25, PACT-GFP, and MCAK-mScarlet-eDHFR.
Representative images (e) show kinetochores, poles, and MCAK recruitment after
dimerization at t=0. Pole-pole distances (f) were measured for each cell and averaged
at every time point (n=29 cells pooled from four independent experiments). Time
stamps, min:s. Scale bars, 10 μm. Error bars: mean ± SEM.

Figure 2. Enhanced kMT depolymerization slows pole separation in anaphase. (a)
Schematic of kMT shortening assay in anaphase. The microtubule depolymerase
MCAK (orange) is recruited to kinetochores (green) after dimerizer uncaging at t=0 at
anaphase onset. **(b)** Representative images show cells expressing Halo-GFP-SPC25,
MCAK-mScarlet-eDHFR, and PACT-GFP, with or without dimerization at t=0,
immediately after sister kinetochores start to move apart. Contrast in the mScarlet
channel was adjusted in each image to visualize MCAK recruitment. Insets for
+dimerizer cells show kinetochores and MCAK recruitment at higher magnification with
enhanced contrast. Time stamps, min:s. Scale bars, 5 μm or 500 nm in insets. **(c-d)**
Analyses of kinetochore and pole motions. Example snapshot at t=0 (c) shows
coordinates of poles (black) and kinetochores (gray circles). A pair of kinetochore
centroids (green) represents the median positions of the two separating kinetochore
masses projected to the mitotic axis (gray dashed lines, defined by the two poles).
Three distances were calculated based on these coordinates: between kinetochore
centroids (D_{KK}, green), between poles (D_{PP}, black), and between each kinetochore
centroid and its adjacent pole (D_{PK}, red). Example traces (d) show distances over time,
with initial velocities (dashed lines) measured in the earliest time window exhibiting
linear increases of all three distances. **(e-g)** Initial velocities were calculated for kMT
depolymerization (e), kinetochore separation (f), and pole separation (g) (n=58 cells –
dimerizer pooled from six independent experiments, 30 cells +dimerizer pooled from
seven independent experiments). Each data point represents a single cell. Lines: mean
± SD. **(h)** The sort-and-grip model proposes that parallel sorters (gray dumbbells)
position kMTs (red) adjacent to parallel microtubules in the central spindle (blue).
Kinetochores (green) grip adjacent microtubules, which are in the correct orientation for
antiparallel sliding (mediated by antiparallel couplers/motors, magenta dumbbells) to

460 move sister kinetochores apart (blue arrows). Depolymerization at kMT plus-ends pulls
461 poles inward toward kinetochores (red arrows) that are coupled to the central spindle.
462

Figure 3. Anti-poleward kinetochore motions in monopolar anaphase. (a)
Schematic of monopolar anaphase. Sister kinetochore pairs can localize away from or
close to antiparallel bundles (off-bundle or on-bundle, respectively). Both kinetochores
can attach to the pole (top panels), or the kMTs of the distal kinetochore can orient
away from the pole (bottom panels). In the latter case, the sort-and-grip model predicts
that antiparallel sliding can move the proximal and distal kinetochores toward and away
from the pole, respectively, as kinetochores grip microtubules in parallel orientation to
their kMTs (bottom right panel). Green arrows represent directions of kinetochore
movement. **(b)** Schematic illustrates the conditional pharmacology approach for spindle
checkpoint silencing in single cells. The MPS1 inhibitor (reversine, brown) is linked to a
photocage (purple) and a Halo ligand (light green). The caged inhibitor localizes to
kinetochores by binding to the Halo-tagged kinetochore protein SPC25 (green) and
inhibits MPS1 (brown) only after uncaging. **(c-d)** Representative images (c) show HeLa
cells expressing Halo-GFP-SPC25 (kinetochores), PACT-GFP (spindle poles), and
670nano3-PRC1 (antiparallel bundles), incubated with the kinesin-5 inhibitor STLC to
create monopolar spindles. Time-lapse images (d) show representative off-bundle and
on-bundle pairs, boxed in c. Yellow line: distance between the off-bundle sisters; time
stamps (min:s): time after uncaging at t=0. **(e)** Individual kinetochore movements
relative to the pole for the off-bundle (gray) and on-bundle (orange) sister pairs shown in
d. Kinetochore velocity relative to the pole (V_{PK}) was calculated as the displacement
within a one-minute time window, with poleward motion defined as the positive direction.
Dashed lines represent the maximal velocity of sister kinetochore separation (Max V_{KK}).
(f) Max V_{KK} plotted for off-bundle (n=168) and on-bundle (n=204) sister kinetochore
pairs, from 23 cells pooled from seven independent experiments. Dashed line: upper
bound for pairs that behave as off-bundle (95th percentile of the Max V_{KK} distribution for
off-bundle pairs). Some on-bundle pairs separate with high V_{KK} as shown in e (orange
traces). **(g)** For each on-bundle pair with Max V_{KK} above the dashed line (n=93 pairs),

491   indicating functional interactions with the bundle, V_{PK} is plotted for the proximal and

492   distal sisters. Black lines: mean ± SD. Scale bars, 5 μm in c or 2 μm in d.

Figure 4. Suppressed bundle sliding inhibits kMT depolymerization. (a) Schematic
of braking assay: monomeric PRC1 (PRC1M, magenta half-dumbbells) is recruited to
dimeric PRC1 (magenta dumbbells) after uncaging at t=0, forming "brake" complexes to
suppress antiparallel sliding (shown by reduced size of blue arrows). **(b)** Constructs for
braking assay. PRC1M ($PRC1_{341-620}$) is truncated to remove the N-terminus. PACT and
CenpB label spindle poles and kinetochores, respectively. **(c)** Representative images of
cells expressing CenpB-GFP, PACT-GFP, Halo-670nano3-PRC1, and mScarlet-
eDHFR-PRC1M with or without dimerization at t=0, immediately after anaphase onset.
Time stamps min:s; scale bars, 5 μm. **(d-f)** Initial velocity analyses for pole separation
(d), kinetochore separation (e), and kMT depolymerization (f). From top to bottom in
each panel: n=39 control cells (without Halo-670nano3-PRC1 or mScarlet-eDHFR-
PRC1M), n=28 with Halo-670nano3-PRC1, n=42 with Halo-670nano3-PRC1 and
mScarlet-eDHFR-PRC1M but no dimerization, n=27 with Halo-670nano3-PRC1 and
mScarlet-eDHFR-PRC1M and dimerization. All cells express CenpB-GFP and PACT-
GFP. Cells were pooled from three independent experiments; each data point
represents a single cell. Lines: mean ± SD.

Figure 5. Proposed sort-and-grip processes. Before anaphase onset, microtubule

sorting mechanisms position kMTs (red) close to parallel microtubules in the central

spindle (blue) by parallel sorters (gray dumbbells). Upon anaphase onset, kinetochores

(green) grip adjacent microtubules, which are correctly oriented for antiparallel sliding to

move sister kinetochores apart. Parallel sorters are not needed after anaphase onset

and are not shown for simplicity. Two concurrent processes drive pole and kinetochore

movements in anaphase: antiparallel sliding directly separates gripped kinetochores,

while microtubule depolymerization at kinetochores pulls poles inward toward

kinetochores. The processes are shown sequentially for clarity but can occur

simultaneously.

References

1. Maiato, H. & Lince-Faria, M. The perpetual movements of anaphase. *Cell Mol Life Sci* **67**,
2251-69 (2010).

2. Alberts, B. et al. in *Molecular Biology of the Cell*, Vol. seventh edition 1027-88 (W. W.
Norton & Company, 2022).

3. Asbury, C.L. Anaphase A: Disassembling Microtubules Move Chromosomes toward
Spindle Poles. *Biology (Basel)* **6**, 15 (2017).

4. McIntosh, J.R. Anaphase A. *Semin Cell Dev Biol* **117**, 118-126 (2021).

5. Scholey, J.M., Civelekoglu-Scholey, G. & Brust-Mascher, I. Anaphase B. *Biology (Basel)* **5**,
51 (2016).

6. Vukusic, K. & Tolic, I.M. Anaphase B: Long-standing models meet new concepts. *Semin*
Cell Dev Biol **117**, 127-139 (2021).

7. Kapitein, L.C. et al. The bipolar mitotic kinesin Eg5 moves on both microtubules that it
crosslinks. *Nature* **435**, 114-8 (2005).

8. Wijeratne, S. & Subramanian, R. Geometry of antiparallel microtubule bundles regulates
relative sliding and stalling by PRC1 and Kif4A. *Elife* **7**, e32595 (2018).

9. Lim, W.M. et al. Regulation of minimal spindle midzone organization by mitotic kinases.
Nat Commun **15**, 9213 (2024).

10. Yu, C.H. et al. Central-spindle microtubules are strongly coupled to chromosomes during
both anaphase A and anaphase B. *Mol Biol Cell* **30**, 2503-2514 (2019).

11. Vukusic, K. et al. Microtubule Sliding within the Bridging Fiber Pushes Kinetochore Fibers
Apart to Segregate Chromosomes. *Dev Cell* **43**, 11-23 e6 (2017).

12. Ris, H. The anaphase movement of chromosomes in the spermatocytes of the
grasshopper. *Biol Bull* **96**, 90-106 (1949).

13. Nicklas, R.B. Chromosome movement and spindle birefringence in locally heated cells:
interaction versus local control. *Chromosoma* **74**, 1-37 (1979).

14. Pamula, M.C. et al. High-resolution imaging reveals how the spindle midzone impacts
chromosome movement. *J Cell Biol* **218**, 2529-2544 (2019).

15. Vukusic, K., Ponjavic, I., Buda, R., Risteski, P. & Tolic, I.M. Microtubule-sliding modules
based on kinesins EG5 and PRC1-dependent KIF4A drive human spindle elongation. *Dev*
Cell **56**, 1253-1267 e10 (2021).

16. Orr, B. et al. An anaphase surveillance mechanism prevents micronuclei formation from
frequent chromosome segregation errors. *Cell Rep* **37**, 109783 (2021).

17. Pereira, A.L. et al. Mammalian CLASP1 and CLASP2 cooperate to ensure mitotic fidelity
by regulating spindle and kinetochore function. *Mol Biol Cell* **17**, 4526-42 (2006).

18. Brust-Mascher, I., Sommi, P., Cheerambathur, D.K. & Scholey, J.M. Kinesin-5-dependent
poleward flux and spindle length control in Drosophila embryo mitosis. *Mol Biol Cell* **20**,
1749-62 (2009).

19. Rogers, G.C. et al. Two mitotic kinesins cooperate to drive sister chromatid separation
during anaphase. *Nature* **427**, 364-70 (2004).

20. Ganem, N.J., Upton, K. & Compton, D.A. Efficient mitosis in human cells lacking
poleward microtubule flux. *Curr Biol* **15**, 1827-32 (2005).

21.   Zhang, D., Rogers, G.C., Buster, D.W. & Sharp, D.J. Three microtubule severing enzymes
contribute to the "Pacman-flux" machinery that moves chromosomes. *J Cell Biol* **177**,
231-42 (2007).

22.   Rath, U. et al. The Drosophila kinesin-13, KLP59D, impacts Pacman- and Flux-based
chromosome movement. *Mol Biol Cell* **20**, 4696-705 (2009).

23.   Mukherjee, S. et al. Human Fidgetin is a microtubule severing the enzyme and minus-
end depolymerase that regulates mitosis. *Cell Cycle* **11**, 2359-66 (2012).

24.   Guerreiro, A. et al. WDR62 localizes katanin at spindle poles to ensure synchronous
chromosome segregation. *J Cell Biol* **220**, e202007171 (2021).

25.   Anjur-Dietrich, M.I., Kelleher, C.P. & Needleman, D.J. Mechanical Mechanisms of
Chromosome Segregation. *Cells* **10**, 465 (2021).

26.   Zimyanin, V. & Redemann, S. Microtubule length correlates with spindle length in C.
elegans meiosis. *Cytoskeleton (Hoboken)* **81**, 356-368 (2024).

27.   Canman, J.C. et al. Determining the position of the cell division plane. *Nature* **424**, 1074-
8 (2003).

28.   Ballister, E.R., Aonbangkhen, C., Mayo, A.M., Lampson, M.A. & Chenoweth, D.M.
Localized light-induced protein dimerization in living cells using a photocaged dimerizer.
Nat Commun **5**, 5475 (2014).

29.   Zhang, H. et al. Optogenetic control of kinetochore function. *Nat Chem Biol* **13**, 1096-
1101 (2017).

30.   Chen, G.Y. et al. Tension promotes kinetochore-microtubule release by Aurora B kinase.
J Cell Biol **220**, e202007030 (2021).

31.   Chen, G.Y. & Lampson, M.A. Chemical tools for dissecting cell division. *Nat Chem Biol* **17**,
632-640 (2021).

32.   Hunter, A.W. et al. The kinesin-related protein MCAK is a microtubule depolymerase
that forms an ATP-hydrolyzing complex at microtubule ends. *Mol Cell* **11**, 445-57 (2003).

33.   Valdez, V.A., Neahring, L., Petry, S. & Dumont, S. Mechanisms underlying spindle
assembly and robustness. *Nat Rev Mol Cell Biol* **24**, 523-542 (2023).

34.   Oriola, D., Needleman, D.J. & Brugues, J. The Physics of the Metaphase Spindle. *Annu*
Rev Biophys **47**, 655-673 (2018).

35.   Khodjakov, A., Copenagle, L., Gordon, M.B., Compton, D.A. & Kapoor, T.M. Minus-end
capture of preformed kinetochore fibers contributes to spindle morphogenesis. *J Cell*
Biol **160**, 671-83 (2003).

36.   Deng, C. et al. Conditional Localization Pharmacology Manipulates the Cell Cycle with
Spatiotemporal Precision. *BioRxiv*, 612697 (2024).

37.   Neef, R. et al. Choice of Plk1 docking partners during mitosis and cytokinesis is
controlled by the activation state of Cdk1. *Nat Cell Biol* **9**, 436-44 (2007).

38.   Mollinari, C. et al. PRC1 is a microtubule binding and bundling protein essential to
maintain the mitotic spindle midzone. *J Cell Biol* **157**, 1175-86 (2002).

39.   Lansky, Z. et al. Diffusible crosslinkers generate directed forces in microtubule networks.
Cell **160**, 1159-68 (2015).

40.   Gaska, I., Armstrong, M.E., Alfieri, A. & Forth, S. The Mitotic Crosslinking Protein PRC1
Acts Like a Mechanical Dashpot to Resist Microtubule Sliding. *Dev Cell* **54**, 367-378 e5
(2020).

41. Braun, M. et al. Adaptive braking by Ase1 prevents overlapping microtubules from
sliding completely apart. *Nat Cell Biol* **13**, 1259-64 (2011).
42. Janson, M.E. et al. Crosslinkers and motors organize dynamic microtubules to form
stable bipolar arrays in fission yeast. *Cell* **128**, 357-68 (2007).
43. Desai, A., Maddox, P.S., Mitchison, T.J. & Salmon, E.D. Anaphase A chromosome
movement and poleward spindle microtubule flux occur At similar rates in Xenopus
extract spindles. *J Cell Biol* **141**, 703-13 (1998).
44. Murray, A.W., Desai, A.B. & Salmon, E.D. Real time observation of anaphase in vitro.
Proc Natl Acad Sci U S A **93**, 12327-32 (1996).
45. Laband, K. et al. Chromosome segregation occurs by microtubule pushing in oocytes.
Nat Commun **8**, 1499 (2017).
46. Dumont, J., Oegema, K. & Desai, A. A kinetochore-independent mechanism drives
anaphase chromosome separation during acentrosomal meiosis. *Nat Cell Biol* **12**, 894-
901 (2010).
47. Nahaboo, W., Zouak, M., Askjaer, P. & Delattre, M. Chromatids segregate without
centrosomes during Caenorhabditis elegans mitosis in a Ran- and CLASP-dependent
manner. *Mol Biol Cell* **26**, 2020-9 (2015).
48. Spurck, T.P. et al. UV microbeam irradiations of the mitotic spindle. II. Spindle fiber
dynamics and force production. *J Cell Biol* **111**, 1505-18 (1990).
49. Spurck, T., Forer, A. & Pickett-Heaps, J. Ultraviolet microbeam irradiations of epithelial
and spermatocyte spindles suggest that forces act on the kinetochore fibre and are not
generated by its disassembly. *Cell Motil Cytoskeleton* **36**, 136-48 (1997).
50. Nicklas, R.B. The motor for poleward chromosome movement in anaphase is in or near
the kinetochore. *J Cell Biol* **109**, 2245-55 (1989).
51. Redemann, S. et al. C. elegans chromosomes connect to centrosomes by anchoring into
the spindle network. *Nat Commun* **8**, 15288 (2017).
52. Carlini, L., Renda, F., Pamula, M.C., Khodjakov, A. & Kapoor, T.M. Coupling of
microtubule bundles isolates them from local disruptions to set the structural stability of
the anaphase spindle. *Proc Natl Acad Sci U S A* **119**, e2204068119 (2022).
53. Suresh, P., Long, A.F. & Dumont, S. Microneedle manipulation of the mammalian spindle
reveals specialized, short-lived reinforcement near chromosomes. *Elife* **9**, e53807
(2020).
54. Elting, M.W., Prakash, M., Udy, D.B. & Dumont, S. Mapping Load-Bearing in the
Mammalian Spindle Reveals Local Kinetochore Fiber Anchorage that Provides
Mechanical Isolation and Redundancy. *Curr Biol* **27**, 2112-2122 e5 (2017).
55. Kajtez, J. et al. Overlap microtubules link sister k-fibres and balance the forces on bi-
oriented kinetochores. *Nat Commun* **7**, 10298 (2016).
56. Sikirzhytski, V. et al. Microtubules assemble near most kinetochores during early
prometaphase in human cells. *J Cell Biol* **217**, 2647-2659 (2018).
57. Renda, F. et al. Non-centrosomal microtubules at kinetochores promote rapid
chromosome biorientation during mitosis in human cells. *Curr Biol* **32**, 1049-1063 e4
(2022).
58. Lecland, N. & Luders, J. The dynamics of microtubule minus ends in the human mitotic
spindle. *Nat Cell Biol* **16**, 770-8 (2014).

59.   Petry, S., Groen, A.C., Ishihara, K., Mitchison, T.J. & Vale, R.D. Branching microtubule
nucleation in Xenopus egg extracts mediated by augmin and TPX2. *Cell* **152**, 768-77
(2013).
60.   Alfaro-Aco, R., Thawani, A. & Petry, S. Biochemical reconstitution of branching
microtubule nucleation. *Elife* **9**, e49797 (2020).
61.   Almeida, A.C. et al. Augmin-dependent microtubule self-organization drives kinetochore
fiber maturation in mammals. *Cell Rep* **39**, 110610 (2022).
62.   Stimac, V., Koprivec, I., Manenica, M., Simunic, J. & Tolic, I.M. Augmin prevents
merotelic attachments by promoting proper arrangement of bridging and kinetochore
fibers. *Elife* **11**, e83287 (2022).
63.   Haren, L., Gnadt, N., Wright, M. & Merdes, A. NuMA is required for proper spindle
assembly and chromosome alignment in prometaphase. *BMC Res Notes* **2**, 64 (2009).
64.   Yang, Z., Tulu, U.S., Wadsworth, P. & Rieder, C.L. Kinetochore dynein is required for
chromosome motion and congression independent of the spindle checkpoint. *Curr Biol*
17, 973-80 (2007).
65.   Yoshida, M.W., Yamada, M. & Goshima, G. Moss Kinesin-14 KCBP Accelerates Chromatid
Motility in Anaphase. *Cell Struct Funct* **44**, 95-104 (2019).
66.   Hu, C.K., Coughlin, M., Field, C.M. & Mitchison, T.J. KIF4 regulates midzone length during
cytokinesis. *Curr Biol* **21**, 815-24 (2011).
67.   Kunchala, P. et al. Plasticity of the mitotic spindle in response to karyotype variation.
Curr Biol **34**, 3416-3428 e4 (2024).
68.   Uehara, R. et al. Aurora B and Kif2A control microtubule length for assembly of a
functional central spindle during anaphase. *J Cell Biol* **202**, 623-36 (2013).
69.   Rizk, R.S., Discipio, K.A., Proudfoot, K.G. & Gupta, M.L., Jr. The kinesin-8 Kip3 scales
anaphase spindle length by suppression of midzone microtubule polymerization. *J Cell*
Biol **204**, 965-75 (2014).
70.   Straight, A.F., Sedat, J.W. & Murray, A.W. Time-lapse microscopy reveals unique roles
for kinesins during anaphase in budding yeast. *J Cell Biol* **143**, 687-94 (1998).
71.   Arquint, C., Gabryjonczyk, A.M. & Nigg, E.A. Centrosomes as signalling centres. *Philos*
Trans R Soc Lond B Biol Sci **369**, 20130464 (2014).
72.   Pollard, T.D. Nine unanswered questions about cytokinesis. *J Cell Biol* **216**, 3007-3016
(2017).
73.   Dumont, S. & Mitchison, T.J. Force and length in the mitotic spindle. *Curr Biol* **19**, R749-
61 (2009).
74.   Khandelia, P., Yap, K. & Makeyev, E.V. Streamlined platform for short hairpin RNA
interference and transgenesis in cultured mammalian cells. *Proc Natl Acad Sci U S A* **108**,
12799-804 (2011).
75.   Andrews, P.D. et al. Aurora B regulates MCAK at the mitotic centromere. *Dev Cell* **6**, 253-
68 (2004).
76.   Subramanian, R., Ti, S.C., Tan, L., Darst, S.A. & Kapoor, T.M. Marking and measuring
single microtubules by PRC1 and kinesin-4. *Cell* **154**, 377-90 (2013).
77.   Gillingham, A.K. & Munro, S. The PACT domain, a conserved centrosomal targeting motif
in the coiled-coil proteins AKAP450 and pericentrin. *EMBO Rep* **1**, 524-9 (2000).
697
698

Supplementary Figure 1. MCAK overexpression does not affect motions of chromosomes and poles. Velocities of kMT depolymerization (a), kinetochore separation (b), and pole separation (c) were measured for cells expressing mScarlet-eDHFR (–MCAK, n=40 cells), or MCAK-mScarlet-eDHFR (+MCAK, n=58 cells) as in Fig. 2c-d. For +MCAK cells, data were replotted from –dimerizer cells in Fig. 2e-g. Each data point represents a single cell. Lines: mean ± SD.

Supplementary Figure 2. PRC1 morphology in monopolar spindles. (a)
Representative images of PRC1 bundles before (left) or after (right) anaphase onset.
Time stamps indicate the time after uncaging. Scale bars, 5 μm. **(b)** An example of
PRC1 intensity analyzed by line scan (box in a). Bundle length and intensity are defined
by the full width at half-maximum and the peak height, respectively. **(c-d)** Analyses for
bundle length (c) and intensity (d). Each dot represents a PRC1 bundle (n = 162 for pre-
anaphase bundles and 148 for anaphase bundles from 23 cells). Lines: mean ± SD.

716
Supplementary Figure 3. PRC1 overexpression and formation of brake complexes
inhibit kMT depolymerization. (a) Representative images of cells expressing CenpB-
GFP and PACT-GFP, with or without expression of Halo-670nano3-PRC1. Time stamps
(min:s) indicate time after first observing sister kinetochore separation at t=0. Scale
bars, 5 μm. **(b-d)** Dependence of pole separation (b), kinetochore separation (c), and
kMT depolymerization (d) velocities on spindle-bound Halo-670nano3-PRC1. Colored
data points show control cells without Halo-670nano3-PRC1 or mScarlet-eDHFR-
PRC1M (gray), cells expressing Halo-670nano3-PRC1 (orange), cells expressing Halo-
670nano3-PRC1 and mScarlet-eDHFR-PRC1M without dimerization (blue), and cells
expressing Halo-670nano3-PRC1 and mScarlet-eDHFR-PRC1M with dimerization
(pink). All cells express CenpB-GFP and PACT-GFP. Each data point represents a
single cell.
729

730
Supplementary Figure 4. Velocity relationships predicted by the sort-and-grip
model and proposed mechanisms of parallel sorting. (**a**) Schematic shows
kinetochore velocity (V_K), pole velocity (V_P), and antiparallel sliding in the central spindle
($V_{sliding}$). Kinetochore and pole velocities are half of the velocities of kinetochore
separation (V_{KK}) and pole separation (V_{PP}), respectively, as measured in Figs. 2 and 4.
Kinetochores, poles, and fiducial marks that label antiparallel sliding are shown in
green, black, and brown, respectively. In the sort-and-grip model, kinetochore velocity
matches sliding velocity, and pole velocity is sliding velocity (moving poles outward)
subtracting kMT depolymerization from either end (V_{PK}, moving poles inward).
Manipulations of kMT depolymerization affect pole velocity but not kinetochore velocity
in this model. (**b**) Kinetochore-nucleated kMT fragments (red) have their plus-ends at
kinetochores. Minus-end directed motors (gray) at the minus-ends of these short kMTs
preferentially guide kMT growth toward the minus ends of parallel microtubules in the
central spindle (blue) to biorient sister kinetochores. (**c**) Minus-end directed motors
(gray) guide the minus ends of newly-grown microtubules in the central spindle toward
spindle poles through kMTs. (**d**) A fraction of the central spindle is grown by
microtubules branching from kMTs (gray). These branching microtubules form
antiparallel bundles (bridging fibers) between the two sister kinetochores, bound by
antiparallel couplers.
750
751

752 **Supplementary Video 1.** Max-intensity projection over 1 μm for the cell shown in Fig.
753 1e. Time stamps (min:s) indicate time after uncaging at t=0. White arrows: spindle pole;
754 scale bar: 10 μm.
755
756 **Supplementary Video 2.** Max-intensity projection over 4 μm for the cells shown in Fig.
757 2b, with (right) or without (left) dimerization. Time stamps (min:s) indicate time after
758 uncaging at t=0. White arrows: spindle pole; scale bar: 5 μm.
759
760 **Supplementary Video 3.** Max-intensity projection over 2 μm showing the off-bundle
761 sister kinetochore pair (yellow arrow) indicated by the gray box in Fig. 3c. Time stamps
762 (min:s) indicate time after uncaging at t=0. Scale bar: 5 μm.
763
764 **Supplementary Video 4.** Max-intensity projection over 2 μm showing the on-bundle
765 sister kinetochore pair (yellow arrow) indicated by the orange box in Fig. 3c. Time
766 stamps (min:s) indicate time after uncaging at t=0. Scale bar: 5 μm.
767
768 **Supplementary Video 5.** Max-intensity projection over the whole cell for the cells
769 shown in Fig. 4c. Left: –dimerizer; Right: +dimerizer. Time stamps (min:s) indicate time
770 after uncaging at t=0. White arrows: spindle pole; scale bar: 5 μm.
771
772 **Supplementary Video 6.** Max-intensity projection over the whole cell for the cells
773 shown in Supplementary Fig. 3a, with (right) or without (left) expression of Halo-
774 670nano3-PRC1. Time stamps (min:s) indicate time after first observing sister
775 kinetochore separation at t=0. White arrows: spindle pole; scale bar: 5 μm.