A Comprehensive Meta-Analysis of Breast Cancer Gene Expression

Ifeanyichukwu O. Nwosu

Department of Biology

Brigham Young University

Provo, UT, USA

inwosu@byu.edu

Stephen R. Piccolo*

Department of Biology

Brigham Young University

Provo, UT, USA

stephen_piccolo@byu.edu

14
15   *Please address correspondence to S.R.P.

Abstract

Background

Triple-negative breast cancers (TNBC) occur more frequently in African Americans and are

associated with worse outcomes when compared to other subtypes of breast cancer. These

cancers lack expression of estrogen receptor (ER), progesterone receptor (PR) and human

epidermal growth factor receptor 2 (HER2) and have limited treatment options. To shed light

on mechanisms behind these differences and suggest novel treatments, we used a meta-

analytic approach to identify gene expression differences in breast tumors for people with

self-reported African or European ancestry; additionally, we compared gene expression

levels based on ER, PR, HER2 and TNBC status.

Methods

After gathering and standardizing gene expression data and metadata from 106 datasets

(representing 27,000 samples), we identified genes that were expressed differently between

these groups via random-effects meta-analyses. To evaluate the robustness of these gene

lists, we devised a novel computational methodology that uses cross validation and

classification. We also computed overlaps between the most significant genes and known

signaling pathways.

Results

Using a false discovery rate threshold of 0.05, we identified genes that are known to play a

significant role in their respective breast cancer subtypes (e.g., *ESR1* for ER status and

ERBB2 for HER2 status), thus confirming the validity of our findings. We also discovered

genes that have not been reported previously and may be new targets for breast cancer

therapy. *GATA3*, *CA12*, *TBC1D9*, *XBP1* and *FOXA1* were among the most significant genes

for ER, PR, and TNBC. However, none of these genes overlapped with HER2 status,

supporting prior research that HER2 tumors are mechanistically different from endocrine

breast cancers. The genes identified from the race meta-analysis—including *DNAJC15*,

HLA-DPA1, *STAP2*, *CEP68*, *MOGS*—have not been associated previously with race-

specific breast-cancer outcomes, highlighting a potential area of further research.

Conclusions

We have carried out a large meta-analysis of breast cancer gene expression data,

identifying novel genes that may serve as potential biomarkers for breast cancer in diverse

populations. We have also developed a computational method that identifies gene sets small

enough to be analyzed and explored in future studies. This method has the potential to be

applied to other cancers.

Keywords

Breast cancer, Racial disparity, Triple negative Breast Cancer, Meta-analysis, Machine

Learning.

Background

Cancers arise as a result of alterations in groups of genes known as proto-oncogenes and tumor suppressor genes[1,2]. These definitions cover hundreds of genes[3,4] that have been implicated in tumorigenic processes, with an average of 33 to 66 mutations found in solid tumors[5]. This represents an enormous amount of genetic variation, even within the same tissue types[6], making it difficult to develop a "one size fits all" approach to treatments, or to understanding cancer biology in general. Among cancers, breast cancer is the most common type of cancer in women worldwide, and accounts for 12.5% of new cancer cases in the world each year[7,8]. It is also the second most common cause of cancer deaths among women in the U.S. In 2023, the American Cancer Society estimated 297,790 new cases of invasive breast cancer diagnosed in women in the US alone[9]. This is an enormous public health burden. Advances in treatment, coupled with early diagnoses, have improved survival rates and lowered mortality[10,11] for breast cancer patients. However, researchers continue to search for methods to assign the most appropriate types of therapies[12,13] and thus to tailor treatments to patients.

Despite these advancements, the heterogeneous nature of the disease[14] and the complex genetic landscape[15] still make it difficult to unravel the underlying molecular mechanisms governing its pathogenesis and progression. Furthermore, there exist racial disparities among population groups[16]. For example, the American Cancer Society reports that survival rates for all cancers are lower for individuals of African descent when compared to those of Caucasian descent[17]. These differences are largely due to cancers of the breast in women, but this is also true for prostate[18] and lung[19] cancers. Various studies[20–22] have highlighted the multifactorial causes of these disparities. These include socioeconomic factors such as access to healthcare, income levels, as well as treatment delays and trust in physicians. However, these factors do not fully explain these disparities[23,24]. The question thus arises as to factors potentially driven by biological causes.

Gene expression is the process by which the information encoded in genes is used to direct

the synthesis of proteins via RNA transcripts. It is assessed by measuring the number of

RNA transcripts for each gene in a tissue sample; the sum of a sample's RNA transcripts is

referred to as its transcriptome. High-throughput genomic technologies have enabled

researchers to produce huge amounts of gene-expression data[25–27]. The analysis of gene

expression data is a powerful tool for deciphering molecular signatures associated with

different breast cancer subtypes[28], predicting patient prognoses[29] and guiding therapeutic

decision-making[30]. Extensive research has shown that studying the transcriptome can help

to identify functional mechanisms behind biological processes. This is validated by a number

of studies which have explored gene expression patterns in cancer; these have found

patterns associated with race[31–34] and identified connections between race and hormone

receptor status[35,36]. These studies, however, have used a limited number of samples

(between 52 and 627 per study). Some of these datasets are publicly available, while others

are not.

Traditionally, clinical subtypes of breast cancer have been defined based on a patient's

immunohistochemistry (IHC) profile. These profiles are determined by the combined

expression status of three receptors: estrogen receptor (ER), progesterone receptor (PR),

and human epidermal growth factor receptor 2 (HER2)[37]. It has also been observed that

women of African descent more commonly have tumors negative for ER, PR and HER2[38].

These cancers are typically referred to as triple negative breast cancers (TNBC). TNBCs are

associated with worse outcomes when compared to other subtypes[39] and occur more

frequently in African Americans[40,41] than other groups. Some studies have identified

biological processes that may contribute to enhanced disease aggressiveness in African-

American patients, including angiogenesis, chemotaxis[42], and the insulin-like growth factor 1

(IGF1) pathway[43].

Building on a large curation effort that we performed with publicly available data[44], we

identified gene-expression datasets that include race metadata and/or metadata for ER, PR

and HER2 status. In total, we have data for over 27,000 samples across 47 datasets with ER

status, 35 datasets with PR status, 28 datasets with ER status and 7 datasets with race

information. With these datasets, we applied a meta-analytic method that combines data

from multiple independent scientific studies, with the aim of identifying consistent patterns[45].

Our research questions can be broadly divided into: A) which genes are most consistently

expressed differently between individuals who self-identify as having European ancestry or

African ancestry, B) which genes are consistently differentially expressed between

individuals who are positive or negative for ER, PR, HER2, or TNBC status, and C) which

signaling pathways are associated with these genes. By addressing these questions, we aim

to uncover molecular mechanisms underlying breast cancer disparities and hormone

receptor statuses, potentially informing personalized treatment strategies and improving

outcomes across diverse populations.

166
167
168   **Methods**

169
Data Acquisition

In a prior study[44], we gathered 114 gene expression datasets from public repositories

including Gene Expression Omnibus[46,47], ArrayExpress[48], the Molecular Taxonomy of Breast

Cancer International Consortium (METABRIC) project[49], TCGA[50] and the SCAN-B

Initiative[51,52] in a bid to enable researchers to address a variety of questions related to

breast-cancer transcriptomics. For this study, we used a subset of those datasets (Additional

File 1). We identified datasets for which one or more of the following metadata variables

were available: self-reported race/ancestry, ER status, PR status, and/or HER2 status.

Data Preparation

For the comparison based on self-reported ancestry, we filtered the data to focus on people

described as "White", "Caucasian" or "European American" and people described as "Black"

or "African American". We focused on these groups because they were most common in the

data and because an understanding of disparities between these groups is critically needed.

We removed samples that did not fall into either of these categories. Next, we standardized

the metadata variable names because different studies used different terms to represent the

same variables. Samples that were labeled "Caucasian", "W", "w", "European American", or

"white" were all categorized as "White", while samples labeled "African American", "B", "b",

"Black or African American", or "black" were categorized as "Black". For the cell-surface

receptors (ER, PR and HER2), we standardized each value as "positive" or "negative," as

determined by the original authors. We excluded samples with missing values for these

variables. For ambiguous data where we could not readily identify what the values referred

to, we referred to the journal article(s) associated with each dataset. If we found more

information to clarify what those values meant, we standardized and included them in our

study; otherwise, we filtered them out. For example, in one study, we found that the values

"0/1", "0-1", "1+", and "1+ (only 1 core pos.)" referred to HER2 negative samples. When

determining TNBC status, we implemented logic to handle missing values. We marked a

tumor as TNBC-negative only if it was known that the tumor was negative for all three

markers (no missing data). If a tumor was positive for any marker, we inferred that the tumor

was *not* TNBC-negative, even if other markers were missing. Additionally, we identified

tumors for which race and TNBC status were available and performed a comparison

between individuals of African or European ancestry.

For each comparison (Black and White, ER+ vs ER-, PR+ vs PR-, HER2+ vs HER2-, TNBC

vs non-TNBC, and Black TNBC vs White TNBC), we selected datasets that had at least 10

samples in each group. For example, when comparing ER status, we required a minimum of

10 "ER+" samples and 10 "ER-" samples. We also determined the relative proportion of the

groups and ensured the ratio between groups was no larger than 20:1 (in either direction).

We did this to prevent our results from being skewed due to class imbalance.

Meta-analysis

Different gene-expression profiling platforms and annotations produce data for different

genes. Therefore, for each comparison, we first selected genes that were common across all

datasets being used for that comparison. There were 11,049 common genes for race,

10,809 common genes for ER status, 10,960 common genes for PR status, 10,926 common

genes for HER2 status, 10,960 for triple negative status, and 11,049 for race + TNBC.

We carried out the meta-analyses using the DExMA[53] package (version 1.10.7) in

R[54](version 4.3.2). We used the following parameters:

i) typeMethod: random effects model;

ii) missAllow: The maximum proportion of missing values allowed in a sample = 10%;

iii) proportionData: The minimum proportion of datasets in which a gene must be present =

90%.

After the meta-analysis was completed, we filtered the results based on False Discovery

Rate (FDR) values less than 0.05 and arranged them in descending order based on the

absolute value of the effect size. This ensured that the top-ranked genes were both

statistically significant and had a relatively large effect size.

Meta-analyses commonly result in thousands of statistically significant genes. It is difficult to

interpret such large lists. We wished to estimate the number of significant genes that would

effectively discriminate between the groups while being as parsimonious as possible.

Furthermore, Venet et al[55] discuss that empirically identified gene-expression signatures are

sometimes no more predictive of breast-cancer outcomes than random gene expression

signatures, even when the specified genes are statistically significant. Thus, we wished to verify that our identified gene lists were more biologically relevant than what we could obtain using randomly selected genes. To address these goals, we developed a process that used leave-one-dataset-out cross validation. Cross validation is a statistical method to evaluate the performance of a model on unseen data in an unbiased manner[56,57]. Rather than holding out individual samples, we held out one dataset at a time. First, we carried out a meta-analysis using *n - 1* datasets at a time (Figure 1). We assumed that if genes identified through an n - 1 meta-analysis are biologically relevant, expression values for those genes would be effective at classifying the two groups being compared on the held-out dataset. The second step was to perform classification analyses and quantify the predictive performance relative to randomly selected genes. For this step, we performed within-dataset cross validation for the held-out dataset. We used modules from the scikit-learn library (version 1.3.1)[58] for this step. We split the datasets using the StratifiedKFold method because it ensures that each fold of the dataset has a similar proportion of observations with a given label. It is particularly useful when dealing with classification tasks that have imbalanced class distributions, as observed in our datasets. We also used the Random Forest[59] classifier as it has been shown to perform well on gene expression data[60]. For *m* top-ranked genes (5, 10, 20, 50, 100, 500, or 1000), we performed classification and calculated a score for each cross-validation fold. We used the "balanced accuracy" scoring metric to account for the imbalanced nature of the datasets. We then repeated this process 100 times for *m* genes selected at random from the held-out dataset. We ensured that each random gene had not been identified as relevant in the meta-analysis. Consider the race datasets as an example. For each round of meta-analysis, we selected six of the seven data sets, leaving one out, and we performed the meta-analysis on those datasets. The meta-analysis produced a list of genes. From this list, we selected the top 5, 10, 20, 50, 100, 500 and 1000 genes and evaluated how accurately the expression data for those genes predicted race for the seventh data set.

263
264
Figure 1. Diagram illustrating the leave-one-dataset-out cross-validation method. For a
given fold, one dataset was held out for testing. Via meta-analysis, significant genes were
identified and ranked according to fold change. Within-dataset cross validation was
performed for the held-out test dataset, and the labels were predicted using a classification
algorithm for m top-ranked genes. The same process was performed for randomly selected
genes and repeated 100 times. We compared the balanced accuracy for the genes identified
via meta-analysis vs. the genes identified via random selection.

272
273

Pathway Analysis

Based on the cross-validation results, we investigated the gene lists using the Molecular

Signatures Database (MSigDB)[61,62] which has thousands of annotated gene sets. Examining

the overlaps of these gene sets can help identify dysregulated pathways and underlying

biological themes. To compute these overlaps, we pasted the top 20 genes from our gene

lists into the *Investigate* tab on the MSigDB website (https://www.gsea-

msigdb.org/gsea/msigdb/human/annotate.jsp). In particular, we computed overlaps using the

"H: hallmark" gene sets"[63] and the "C2: curated gene sets," using an FDR q-value threshold

of 0.05.

Results

Results

We performed meta-analyses to compare gene-expression levels between people of African

or European descent and between people with positive or negative tumor status (ER, PR,

HER2, or TNBC). There was some overlap across the analyses; some datasets had more

than one of these variables (Table 1). Additionally, we compared TNBC tumors from people

of African descent versus people of European descent.

291
292

Variable	Positive	Negative	Equivocal	Unknown	Total Number of Samples	Number of Datasets
	White	**Black**				
Race	1,175	295	NA	NA	1,470	7
Estrogen Receptor Status	16,174	4,426	26	1,334	21,960	47
Progesterone Receptor Status	11,472	5,359	96	1,214	18,141	35
HER2/Neu Status	2,471	13,429	179	1,482	17,561	28
	Triple Negative	**Non Triple Negative**				
Triple Negative Status	1,982	13,614	NA	NA	15,596	22

Table 1. Dataset and sample counts for the meta-analyses. For some datasets, the
original authors reported hormone receptor status but were unable to determine whether
certain samples were positive or negative due to ambiguous immunohistochemistry results
or when these values were marked as intermediate. These samples are listed as
"Equivocal." In other cases, the status was reported for some samples but not reported for
others. These samples are listed as "Unknown."

299
300
For the comparison between individuals of African or European ancestry, 850 genes were

significantly differentially expressed (Additional File 2). For the comparisons between ER-

positive and ER-negative breast tumors, 9,729 genes were differentially expressed

(Additional File 3). We observed similarly high numbers for PR and HER2: 9,047 and 7,236,

respectively (Additional Files 4-5). For the TNBC versus non-TNBC comparison, 8,939

genes were differentially expressed (Additional File 6). For the comparison between people

of African or European ancestry specific to TNBC tumors, six genes were differentially

expressed (Additional File 7).

To identify a smaller number of genes that should be considered relevant for each meta-

analysis, we developed a new computational method. Leaving out one dataset at a time, we

used the remaining datasets to identify significant genes and then performed classification

via cross validation on the held-out dataset. We compared the predictive performance of *m*

genes (as ranked by each meta-analysis) against the predictive performance of randomly

selected genes, to estimate the number of significant genes that would be most predictive.

This analysis revealed that 20 genes often performed well compared to other thresholds

(Figures 2-6). Although this threshold did not always perform better than other thresholds,

we chose 20 genes as the threshold for all six meta-analyses for consistency. Twenty genes

is few enough that researchers can interpret such a list and potentially create a panel to use

in a laboratory setting. Table 2 shows the top 20 genes for each comparison. *GATA3, CA12,*

XBP1 and *FOXA1* were among the top 20 genes for ER, PR, and TNBC. However, none of

these overlapped with the genes for HER2 status (or race).

Rank	Race	ER Status	PR Status	Her2 Status	TNBC	Race + TNBC
1	DNAJC15	**GATA3**	PGR	ERBB2	**FOXA1**	RETN
2	HLA-DPA1	ESR1	**CA12**	PGAP3	MLPH	ADAMTS13
3	STAP2	**CA12**	MAPT	STARD3	FOXC1	ACOXL
4	CEP68	TBC1D9	**GATA3**	GRB7	**GATA3**	FAM3A
5	MOGS	**FOXA1**	ESR1	CDK12	EN1	GPR35
6	TNS1	MLPH	DNALI1	PSMD3	GABRP	PGLYRP1
7	TXNL4A	DNALI1	GREB1	MED1	**CA12**	
8	PSPH	NAT1	NAT1	GSDMB	BCL11A	

9	EIF5A	PSAT1	SCUBE2	MED24	SFT2D2
10	RPP25	SLC39A6	IL6ST	PNMT	**XBP1**
11	PKP3	**XBP1**	TBC1D9	TCAP	AGR2
12	POP7	ABAT	ABAT	MSL1	FAM171A1
13	CENPA	GREB1	GFRA1	CASC3	DSC2
14	TTBK2	MAPT	**XBP1**	CWC25	TBC1D9
15	CXCL12	THSD4	KDM4B	PSMB3	UGT8
16	PYCR1	BCL11A	THSD4	PIP4K2B	SPDEF
17	HSD17B10	GFRA1	CELSR1	LASP1	PSAT1
18	TIMM23	AGR2	SLC39A6	RPL23	SOX10
19	PCGF1	PHGDH	**FOXA1**	CLCA2	ART3
20	ABCA6	DSC2	ERBB4	KMO	MICALL1

Table 2. Top-ranked genes identified in the meta-analyses. These genes represent our
findings when performing meta-analyses across all datasets for which each variable was
available (i.e., not as part of our cross-validation experiments). Fewer than 20 genes are
shown for "Race + TNBC" because this meta-analysis identified only 6 genes with statistical
significance. Genes bolded and highlighted in red were common across more than one of
the meta-analyses.

For the comparison between tumors from individuals of European or African descent, the

median balanced accuracy for the top 20 genes was 0.57 (the maximum possible accuracy

is 1.0). The median balanced accuracy for the randomly selected genes was 0.50. Using a

one-sided, Mann-Whitney U test, we compared the balanced accuracy values when using

the top 20 meta-analysis genes for classification, versus the balanced accuracies for the

randomly selected genes; the meta-analysis accuracies were *not* significantly higher (p-

value = 0.23; Figure 2). These findings suggest that gene-expression differences between

these populations were not significantly different from randomly selected genes; however,

sample sizes may have been a limiting factor.

For the ER-positive versus ER-negative comparison, the balanced accuracies were

significantly higher for the meta-analysis genes than for the randomly selected genes (p <

0.001; Figure 3). The same was true for the PR, HER2, and TNBC comparisons (p < 0.001).

However, the overall patterns were markedly distinct for HER2 relative to the other

comparisons (Figures 3-6). For nearly all thresholds, the balanced accuracy was above 0.8

when using the meta-analysis genes; however, for ER, PR, and TNBC, the random-gene

accuracies started low (around 0.6) and increased considerably as the number of genes

increased. This pattern suggests that ER and/or PR positivity cause a cascade of

downstream effects, possibly altering the expression of thousands of genes. Thus, when we

selected genes at random (not including the top-ranked genes), the classification algorithm

was frequently able to classify the tumors as positive or negative. In contrast, for HER2, the

median balanced accuracy was below 0.55 for all thresholds when randomly selected genes

were used. This pattern suggests that overexpression of HER2 protein, which typically

results from DNA amplification or somatic mutations[66], thus activating oncogenic

pathways[67,68], causes relatively few downstream effects compared to endocrine breast

cancers, which rely on estrogen or progesterone signaling for tumor growth[69,70].

366
367

Figure 2. Cross-validation results for race status. The left panel represents the balanced accuracy, identified via cross validation on each held-out test set, for different numbers of top-ranked genes from the race meta-analysis. Each data point represents the balanced accuracy for a particular dataset. The right panel represents the balanced accuracy for the same numbers of genes selected at random in 100 iterations. Each data point represents the mean balanced accuracy for a particular dataset, across the iterations. The balanced accuracy for the meta-analysis genes was relatively low (between 0.5 and 0.6) when using 5 to 200 genes; it was equivalent to baseline levels (0.5) when 500 or 1000 genes were used. The balanced accuracy for the randomly selected genes was consistently close to baseline levels.

Figure 3. Cross-validation results for ER Status. The left panel represents the balanced accuracy, identified via cross validation on each held-out test set, for different numbers of top-rank genes from the ER meta-analysis. Each data point represents the balanced accuracy for a particular dataset. The right panel represents the average balanced accuracy for the same numbers of genes selected at random in 100 iterations. Each data point represents the mean balanced accuracy for a particular dataset, across the iterations. The balanced accuracy for the meta-analysis genes was consistently high (around 0.85 to 0.9) regardless of the number of genes used, while the balanced accuracy for the random genes started low and increased as the number of genes increased.

395
396
Figure 4. Cross-validation results for PR Status. The left panel represents the balanced
accuracy, identified via cross validation on each held-out test set, for different numbers of
top-rank genes from the PR meta-analysis. Each data point represents the balanced
accuracy for a particular dataset. The right panel represents the average balanced accuracy
for the same numbers of genes selected at random in 100 iterations. Each data point
represents the mean balanced accuracy for a particular dataset, across the iterations. The
balanced accuracy for the meta-analysis genes was consistently high (around 0.8)
regardless of the number of genes used, while the balanced accuracy for the random genes
started low and increased as the number of genes increased.
406
407
408

410
Figure 5. Cross-validation results for HER2 status. The left panel represents the
balanced accuracy, identified via cross validation on each held-out test set, for different
numbers of top-rank genes from the HER2 meta-analysis. Each data point represents the
balanced accuracy for a particular dataset. The right panel represents the average balanced
accuracy for the same numbers of genes selected at random in 100 iterations. Each data
point represents the mean balanced accuracy for a particular dataset, across the iterations.
The balanced accuracy for the meta-analysis genes was consistently higher than the
balanced accuracy for the random genes.
419
420
421
422

423 Number of genes used for cross-validation predictions
424
Figure 6. Cross-validation results for TNBC status. The left panel represents the
balanced accuracy, identified via cross validation on each held-out test set, for different
numbers of top-rank genes from the TNBC meta-analysis. Each data point represents the
balanced accuracy for a particular dataset. The right panel represents the average balanced
accuracy for the same numbers of genes selected at random in 100 iterations. Each data
point represents the mean balanced accuracy for a particular dataset, across the iterations.
The balanced accuracy for the meta-analysis genes was consistently higher than the
balanced accuracy for the random genes.
433
434
For the comparison between individuals of African or European descent, some of the top

genes were *DNAJC15*, *RPP25*, *CXCL12*, *EIF5A*, and *TIMM23*. These genes play diverse

roles in cellular functioning. *DNAJC15* has been found to be expressed in breast cancer

cells[71] and may play a role in apoptosis[72]. *RPP25* enables ribonuclease P RNA binding

activity and is involved in tRNA 5'-leader removal[73,74]. *CXCL12* is the ligand for the G-protein

coupled receptor and chemokine and can affect tumor progression, angiogenesis,

metastasis, and survival in cancer cells[75,76]. *EIF5A* is a translation elongation factor[77] and

plays a role in regulating tumor cell growth[78]. *TIMM23* mediates the transport of transit

peptide-containing proteins across the membrane[79]. Elvidge et al. showed that some of

these genes may play a role in modulating changes in gene expression induced by hypoxia

in MCF-7 breast cancer cell lines[80]. However, hypoxia and these other cellular processes do

not have a clear, known function that would explain gene-expression differences in breast

tumors between people of African or European descent specifically.

The top genes identified in the ER-positive versus ER-negative comparison were *GATA3*,

ESR1, and *CA12*. These genes exhibited approximately two-fold higher expression in ER-

positive patients compared to ER-negative patients. *ESR1* encodes the estrogen receptor

alpha (ERα) protein, which mediates the effects of estrogen in various tissues throughout the

body[81]. Finding this gene near the top of our list when comparing ER positive versus ER

negative breast cancers is evidence of the validity of our data and methods. *GATA3* is

crucial for the differentiation of the mammary epithelium and is a key regulator in breast

cancer[82]. It also plays a role in mediating the binding of estrogen receptor elements to

ESR1[83]. Furthermore, co-expression of *GATA3*, *ESR1,* and *FOXA1* in breast tumors has

been reported[84,85]. *CA12* is involved in cellular proliferation and pH regulation and has been

associated with a good prognosis in breast cancer[86]. Other significant genes include *FOXA1*

and *TBC1D9. FOXA1* is a transcriptional regulator that modulates the effects of estrogen

signaling by influencing *ESR1* activity[87,88]. *TBC1D9* is involved in cellular trafficking

processes that can influence cancer cell behavior[89]. These and other top-ranked genes merit

further study to provide mechanistic insight into the ER protein's role in breast

tumorigenesis, and suggest their potential roles both as biomarkers and therapeutic targets.

For the comparison between PR-positive and PR-negative tumors, a common thread among

the top genes identified, is their involvement in various aspects of known breast cancer

biology. Other studies[90] have shown that many of these genes (*PGR, ESR1, FOXA1,*

GATA3, GREB1) are differentially expressed and that *PGR*[91] and *CA12*[86] may be good

prognostic markers for breast cancer.

Among the top genes for the comparison between HER2-positive and HER2-negative

tumors were *ERBB2* (which was expected because this gene encodes the HER2 protein) as

well as *PGAP3, STARD3, GRB7, CDK12,* and *PSMD3. PGAP3* is overexpressed in breast

cancer patients, is associated with lymph node metastasis, and correlates with HER2

overexpression[92]. *STARD3* is necessary for the transfer of cholesterol and metabolism in

tumor cells[93], and is often co-amplified with *HER2*[94]. *GRB7* is responsible for signal

transduction from cell surface receptors via signaling pathways in the cell[95]. It is sometimes

co-amplified and overexpressed with HER2 and is associated with a more aggressive

phenotype[96]. *CDK12* is a transcription-associated cyclin-dependent kinase, which plays

various roles in the regulation of translation[97], cellular proliferation[98], and cell cycle

progression[99]; it acts as a tumor promoter in HER2-positive breast cancer[100]. *PSMD3* is an

enzyme regulator by stabilizing *HER2* and preventing it from degradation[101]. These genes

may serve as alternative markers for HER2 positive breast cancer[92,102].

For the comparison between TNBC and non-TNBC patients, the most significant genes

sometimes overlapped with what we identified in the earlier analyses. *FOXA1*, *MLPH*, and

GATA3 were expressed at approximately two-fold higher levels in non-TNBCs compared to

TNBCs, while *FOXC1* had approximately 2-fold lower expression in non-TNBC than TNBC

tumors.

Our last comparison focused on individuals of African or European ancestry who had TNBC.

The statistically significant genes were *RETN, ADAMTS13, ACOXL, FAM3A, GPR35*, and

PGLYRP1. RETN encodes the hormone resistin that plays a role in inflammation[103] and

insulin resistance[104], which are critical factors in obesity-related cancers[105]. Increased

resistin levels have been associated with breast cancer[106,107]. *ADAMTS13* encodes an

enzyme that processes von Willebrand factor, a protein involved in blood clotting.

Dysregulation of *ADAMTS13* has been implicated in cancer metastasis and tumor

angiogenesis due to its role in extracellular matrix remodeling[108,109]. *ACOXL* is involved in

fatty acid β-oxidation and metabolic processes[110]. Although there is no research linking it to

breast cancer, its expression has been linked to prostate cancer, where it is a potential

biomarker due to differential expression in benign versus malignant prostate tissues[111].

FAM3A is involved in regulating of both glucose and lipid metabolism[112]. Given its metabolic

functions and the link between metabolism and cancer progression, it has been suggested to

play a role in metabolic diseases[113] and possibly cancers[114]. *GPR35* is involved in

inflammatory responses and is highly expressed in various cancers, including

gastrointestinal and prostate. It is considered a potential therapeutic target due to its role in

tumorigenesis and cancer progression[115,116]. *PGLYRP1* encodes a protein involved in the

innate immune response and plays a role in regulating acquired immunity[117]. *PGLYRP1* has

been implicated in inflammation and immune responses in the tumor

microenvironment[118,119], which can influence cancer development and progression. These

genes share involvement in inflammation, metabolic regulation, and the tumor

microenvironment, which are critical processes in the development and progression of

cancer even though they are not all specific to breast cancer. We did not find evidence to

provide mechanistic insight about these expression differences between people of African or

European descent. Yet they may merit further exploration as potential therapeutic targets.

To gain further insight into the biology behind the selected genes for each comparison, we

performed pathway enrichment analysis using MSigDB[61,62] (Additional Files 8 - 12). Some of

these results further confirm the validity of our methods. For example, in the ER-positive

versus ER-negative comparison, curated pathways that reflect ER signaling were statistically

significant. Similarly, for the HER2-positive versus HER2-negative comparison, pathways

associated with HER2 amplification were significant. For the PR comparison, ER pathways

were statistically significant, but we did not observe pathways specific to PR signaling; this

suggests that ER and PR positivity have similar effects but may also reflect that MSigDB

lacks pathways specific to PR. When a gene is statistically significant but not associated with

any of the statistically significant pathways for a given comparison, it serves as a hypothesis

for further investigation. Additionally, for significant pathways, genes in those pathways

might be prioritized for investigation, even when they were not among the most statistically

significant genes.

Discussion

Researchers have used diverse methodologies to analyze and combine gene expression

datasets[120]. Some have focused on identifying prognostic markers[43,121] or genes[122,123],

predicting treatment outcomes[124–126], or deriving more specific breast cancer subtypes[49,127].

Others have attempted to differentiate between primary and metastatic tumors[128], as well as

differential gene-expression analysis[43,129]. In this study, we carried out meta-analyses for

tens of thousands of breast tumors. To the best of our knowledge, no meta-analyses of this

magnitude has been performed previously; nor has a meta-analysis been performed to

compare breast cancer patients of African and European ancestry. Schneider et al. focused

on ER patients and identified a subset of genes that also show up in our analysis[130]. Al-Ejeh

et al. focused on TNBCs and identified a set of 206 "aggressive genes"[30], some of which

also were significant in our study. D'Arcy et al. compared Luminal A breast tumors between

African American and Caucasian women and arrived at a list of 6 genes; but only one of

those genes (*PSPH*) overlaps with our list. These differences may be due to their modeling

approach and the fact that they had a smaller sample size (n = 167 samples) and used an

imputation method to account for missing data.

Race and Gene Expression

One goal of this research was to understand the biology behind disparities in breast cancer

between women of African and Caucasian descent. Our analysis revealed 850 genes that

were significantly differentially expressed between these populations. Among the top 20

genes, we observed significant roles for inflammation (*RETN*)[103], cell cycle regulation

(*CENPA* and *TTBK2*)[131,132], immune responses (*HLA-DPA1, CXCL12,* and *STAP2*)[76,133,134],

and cellular metabolism (*PSPH, PYCR1,* and *HSD17B10*)[135−137]. These can influence tumor

behavior and responses to treatments. For instance, *RETN* (resistin) is associated with

inflammation and insulin resistance, which are critical in the pathogenesis of obesity-related

cancers, including breast cancer. Elevated resistin levels have been linked to poor prognosis

in breast cancer patients, underscoring the potential of RETN as a biomarker for breast

cancer in diverse populations.

Black women are more likely to be diagnosed with TNBC, which is more aggressive and has

fewer targeted treatment options, while Caucasian women are more frequently diagnosed

with hormone receptor-positive breast cancers, which have better prognoses and more

effective hormone-targeted therapies[138,139]. The genes we identified when we merged TNBC

and race are involved in inflammation and immune responses, which are critical pathways in

cancer development, including breast cancer. ADAMTS13, an enzyme involved in blood

clotting, plays a role in cancer metastasis and angiogenesis through extracellular matrix

remodeling. Dysregulation of ADAMTS13 has been implicated in various cancers,

suggesting its potential as a therapeutic target in breast cancer[140]. Similarly, genes such as

GPR35, involved in inflammatory responses, and *PGLYRP1*, implicated in the innate

immune response, highlight the importance of inflammation and immune regulation in cancer

development[141−145]. Variations in these genes and their expression can vary by race,

potentially influencing disease susceptibility and progression[146,147].

A key limitation of our analysis is that categories based on race and ethnicity are self-

described and may not adequately represent a person's ancestry. By performing secondary

577 analyses of publicly available data, we were limited to the descriptions provided by the

578 original authors. For simplicity, we used broad groupings (African descent, European

579 descent) to categorize the patients. However, by aggregating data from many studies, we

580 anticipated that meaningful patterns would be identifiable, despite this noise. Although our

581 analyses identified genes that have plausible connections to breast tumorigenesis and may

582 lead to insights about population disparities, gene-expression differences between these

583 categories were relatively minor, providing support for the argument that race is a social

584 construct more than a biological concept[148].

586 *ER, PR, and HER2 Status*

587 Compared to ER-negative breast tumors, genes like *ESR1, GATA3,* and *CA12* were more

588 highly expressed in ER-positive tumors. *ESR1*, encoding the estrogen receptor alpha (ERα),

589 is crucial for mediating estrogen's effects and is a well-established marker in breast cancer

590 diagnostics and treatment[81,149]. *GATA3* is essential for mammary epithelial differentiation

591 and is a key regulator in breast cancer, while *CA12* is involved in cellular proliferation and pH

592 regulation and is associated with a good prognosis[150,151].

594 For PR status, genes such as *PGR, ESR1,* and *FOXA1* were differentially expressed,

595 consistent with their roles in breast cancer biology. Progesterone receptor (PGR) and

596 estrogen receptor (ESR1) are critical markers for hormone receptor-positive breast cancers

597 and are important for determining therapeutic strategies[152,153].

599 Our HER2-positive vs. HER2-negative comparisons identified genes like *ERBB2*, *PGAP3*,

600 and *GRB7*, which were prominently expressed in HER2-positive tumors. *ERBB2*

601 overexpression is a hallmark of HER2-positive breast cancers and a target for therapies

602 such as trastuzumab[154]. The co-amplification of genes like *GRB7* with HER2 suggests their

603 roles in aggressive breast cancer phenotypes and potential as additional therapeutic

604 targets[155,156].

TNBC vs. Non-TNBC

TNBC vs. non-TNBC analysis identified *FOXA1, MLPH,* and *GATA3,* which showed higher expression in non-TNBC, while *FOXC1* was expressed lower in non-TNBC. These findings highlight the distinct molecular profiles of TNBC, characterized by the lack of hormone receptor and HER2 expression[157].

Common Genes Across Comparisons

GATA3, CA12, TBC1D9, XBP1, and *FOXA1* were consistently significant across ER, PR, and TNBC comparisons but not in the HER2 comparison. This supports the notion that the mechanisms behind ER-positive and PR-positive tumors are overlapping and that HER2-positive tumors are mechanistically distinct from endocrine breast cancers, necessitating different therapeutic strategies[158–160].

Computational Validation

It is common for differential-expression studies, such as this meta-analysis, to identify thousands of statistically significant genes. Such findings may be reassuring because they imply that the compared phenotypes are truly distinct at the molecular level. However, a larger goal of a differential-expression study is to reduce the search space for genes that are most important in driving the phenotype. Lab-based, experimental validation can provide assurances that genes are truly differentially expressed between conditions; however, such procedures do not scale beyond a few genes. Furthermore, the complexity of cancers likely cannot be captured adequately using data from a few genes. Therefore, it is desirable to find a balance between characterizing the complexity of cancers and identifying a gene set that is small enough to be analyzed and interpreted by humans and that can be explored feasibly in subsequent, mechanistic studies. Our novel computational method addresses these goals, while also showing that most of our identified gene sets are more relevant than randomly selected genes. Moreover, it provides insight into the potential to use these genes

to predict phenotypes. A limitation of differential-expression studies is that commonly used

statistical methods consider evidence from one gene at a time, whereas our classification

approach constructs models from many genes simultaneously. We believe this method has

potential to be applied in many other contexts. Further validation will be necessary.

Conclusion

Findings from our meta-analyses elucidate gene expression differences between racial

groups and clinically relevant breast cancer subtypes, highlighting the roles of diverse

cellular processes. They provide additional evidence about mechanisms that may drive

breast cancer biology. Such studies are necessary to pave the way for developing targeted

diagnostics and therapies in a personalized-medicine context. Further research is warranted

to explore the mechanistic pathways of these genes and their potential as biomarkers and

therapeutic targets in diverse patient populations.

List of Abbreviations

ER : estrogen receptor

ESR1 : estrogen receptor

HER2 : human epidermal growth factor receptor 2

IGF1 : insulin-like growth factor 1

IHC : immunohistochemistry

MSigDB : Molecular Signatures Database

OSF : Open Science Framework

PGR : Progesterone receptor

PR : progesterone receptor

SCAN-B : The Sweden Cancerome Analysis Network - Breast

TCGA : The Cancer Genome Atlas

TNBC : Triple-negative breast cancers

Additional Data Files

Additional File 1

Format: xlsx

Title: Identifiers for datasets used in the meta-analyses.

Description: This spreadsheet indicates the accession number or other unique identifiers for each dataset analyzed in this study.

Additional File 2

Format: xlsx

Title: Genes that were significantly differentially expressed between individuals of African or European ancestry.

Additional File 3

Format: xlsx

Title: Genes that were significantly differentially expressed between ER-positive and ER-negative breast tumors.

Additional File 4

Format: xlsx

Title: Genes that were significantly differentially expressed between PR-positive and PR-negative breast tumors.

Additional File 5

Format: xlsx

Title: Genes that were significantly differentially expressed between HER2-positive and HER2-negative breast tumors.

Additional File 6

Format: xlsx

Title: Genes that were significantly differentially expressed between TNBC versus

non-TNBC breast tumors.

Additional File 7

Format: xlsx

Title: Genes that were significantly differentially expressed in TNBC tumors between

individuals of African or European ancestry.

Additional File 8

Format: xlsx

Title: Gene-set overlap results for the top-ranked genes identified in the race meta-

analysis.

Additional File 9

Format: xlsx

Title: Gene-set overlap results for the top-ranked genes identified in the ER status

meta-analysis.

Additional File 10

Format: xlsx

Title: Gene-set overlap results for the top-ranked genes identified in the PR status

meta-analysis.

Additional File 11

Format: xlsx

Title: Gene-set overlap results for the top-ranked genes identified in the HER2 status

meta-analysis.

Additional File 12.

Format: xlsx

Title: Gene-set overlap results for the top-ranked genes identified in the TNBC status

meta-analysis.

722
723 **Declarations**
724
725 *Ethics approval and consent to participate*
726 Not applicable
727
728
729 *Consent for publication*
730 Not applicable
731
732
733 *Availability of data and materials*
734 All source code, gene expression data, and metadata are available in a publicly accessible

735 Open Science Framework (OSF)[64] repository (https://osf.io/mr8sb/). The code is packaged

736 and executable using Docker[65] software containers for easier reproducibility.

737
738
739 *Competing interests*
740 The authors declare that they have no competing interests
741
742
743 *Funding*
744 The authors received no external funding for this research.
745
746
747 *Authors' contributions*
748 ION prepared and analyzed data. Both authors designed the study and interpreted the

749 results. SRP supervised the work. Both authors wrote the manuscript and approved the final

750 manuscript.

751
752 *Acknowledgements*
753 Not applicable
754
755

References

1. Vogelstein, B. & Kinzler, K. W. Cancer genes and the pathways they control. *Nat. Med.*

10, 789–799 (2004).

2. Ali, M. A. & Sjöblom, T. Molecular pathways in tumor progression: from discovery to

functional understanding. *Mol. Biosyst.* **5**, 902–908 (2009).

3. Futreal, P. A. *et al.* A census of human cancer genes. *Nat. Rev. Cancer* **4**, 177–183

(2004).

4. Cosmic. Cancer Gene Census. https://cancer.sanger.ac.uk/census (2018).

5. Vogelstein, B. *et al.* Cancer genome landscapes. *Science* **339**, 1546–1558 (2013).

6. Pennisi, E. Human Genetic Variation. *Science* **318**, 1842–1843 (2007).

7. Breast cancer facts and statistics 2023. https://www.breastcancer.org/facts-statistics.

8. CDCBreastCancer. Breast cancer statistics. *Centers for Disease Control and Prevention*

https://www.cdc.gov/cancer/breast/statistics/index.htm (2023).

9. Breast Cancer Statistics. https://www.cancer.org/cancer/types/breast-cancer/about/how-

common-is-breast-cancer.html.

10. Arranz, E. E., Vara, J. Á. F., Gámez-Pozo, A. & Zamora, P. Gene signatures in breast

cancer: current and future uses. *Transl. Oncol.* **5**, 398–403 (2012).

11. Berry, D. A. *et al.* Effect of screening and adjuvant therapy on mortality from breast

cancer. *N. Engl. J. Med.* **353**, 1784–1792 (2005).

12. Lal, S., McCart Reed, A. E., de Luca, X. M. & Simpson, P. T. Molecular signatures in

breast cancer. *Methods* vol. 131 135–146 Preprint at

https://doi.org/10.1016/j.ymeth.2017.06.032 (2017).

13. Ringnér, M., Fredlund, E., Häkkinen, J., Borg, Å. & Staaf, J. GOBO: gene expression-

based outcome for breast cancer online. *PLoS One* **6**, e17911 (2011).

14. Guo, L. *et al.* Breast cancer heterogeneity and its implication in personalized precision

therapy. *Exp. Hematol. Oncol.* **12**, 3 (2023).

15. Melchor, L. & Benítez, J. The complex genetic landscape of familial breast cancer. *Hum.*

Genet. **132**, 845–863 (2013).

16. Keenan, T. *et al.* Comparison of the Genomic Landscape Between Primary Breast

Cancer in African American Versus White Women and the Association of Racial

Differences With Tumor Recurrence. *Journal of Clinical Oncology* vol. 33 3621–3627

Preprint at https://doi.org/10.1200/jco.2015.62.2126 (2015).

17. American Cancer Society. Cancer Facts & Figures for African Americans 2016-2018.

American Cancer Society (2016).

18. Lowder, D. *et al.* Racial disparities in prostate cancer: A complex interplay between

socioeconomic inequities and genomics. *Cancer Lett.* **531**, 71–82 (2022).

19. Ryan, B. M. Lung cancer health disparities. *Carcinogenesis* **39**, 741–751 (2018).

20. Tannenbaum, S. L., Koru-Sengul, T., Miao, F. & Byrne, M. M. Disparities in survival

after female breast cancer diagnosis: a population-based study. *Cancer Causes Control*

24, 1705–1715 (2013).

21. Nurgalieva, Z. Z. *et al.* Impact of timing of adjuvant chemotherapy initiation and

completion after surgery on racial disparities in survival among women with breast

cancer. *Med. Oncol.* **30**, 419 (2013).

22. Polite, B. N., Dignam, J. J. & Olopade, O. I. Colorectal cancer model of health

disparities: understanding mortality differences in minority populations. *J. Clin. Oncol.*

24, 2179–2187 (2006).

23. Smith, E. C., Ziogas, A. & Anton-Culver, H. Delay in surgical treatment and survival after

breast cancer diagnosis in young women by race/ethnicity. *JAMA Surg.* **148**, 516–523

(2013).

24. Parise, C. A. & Caggiano, V. Disparities in race/ethnicity and socioeconomic status: risk

of mortality of breast cancer patients in the California Cancer Registry, 2000--2010.

BMC Cancer **13**, 449 (2013).

25. Goodwin, S., McPherson, J. D. & McCombie, W. R. Coming of age: ten years of next-

generation sequencing technologies. *Nat. Rev. Genet.* **17**, 333–351 (2016).

26. Hansen, A. R. & Bedard, P. L. Clinical application of high-throughput genomic

technologies for treatment selection in breast cancer. *Breast Cancer Res.* **15**, R97

(2013).

27. Mardis, E. R. The impact of next-generation sequencing technology on genetics. *Trends*

Genet. **24**, 133–141 (2008).

28. Parker, J. S. *et al.* Supervised risk predictor of breast cancer based on intrinsic

subtypes. *J. Clin. Oncol.* **27**, 1160–1167 (2009).

29. Sørlie, T. *et al.* Distinct molecular mechanisms underlying clinically relevant subtypes of

breast cancer: gene expression analyses across three different platforms. *BMC*

Genomics **7**, 127 (2006).

30. Al-Ejeh, F. *et al.* Meta-analysis of the global gene expression profile of triple-negative

breast cancer identifies genes for the prognostication and treatment of aggressive

breast cancer. *Oncogenesis* **3**, e100 (2014).

31. Sturtz, L. A., Melley, J., Mamula, K., Shriver, C. D. & Ellsworth, R. E. Outcome

disparities in African American women with triple negative breast cancer: a comparison

of epidemiological and molecular factors between African American and Caucasian

women with triple negative breast cancer. *BMC Cancer* **14**, 62 (2014).

32. D'Arcy, M. *et al.* Race-associated biological differences among Luminal A breast

tumors. *Breast Cancer Research and Treatment* vol. 152 437–448 Preprint at

https://doi.org/10.1007/s10549-015-3474-4 (2015).

33. Field, L. A. *et al.* Identification of differentially expressed genes in breast tumors from

African American compared with Caucasian women. *Cancer* **118**, 1334–1344 (2012).

34. Stewart, P. A., Luks, J., Roycik, M. D., Sang, Q.-X. A. & Zhang, J. Differentially

expressed transcripts and dysregulated signaling pathways and networks in African

American breast cancer. *PLoS One* **8**, e82460 (2013).

35. Byun, J. S. *et al.* Racial Differences in the Association Between Luminal Master

Regulator Gene Expression Levels and Breast Cancer Survival. *Clin. Cancer Res.* **26**,

1905–1914 (2020).

36. Chavez-MacGregor, M. *et al.* Differences in gene and protein expression and the effects

of race/ethnicity on breast cancer subtypes. *Cancer Epidemiol. Biomarkers Prev.* **23**,

316–323 (2014).

37. Howlader, N. *et al.* US incidence of breast cancer subtypes defined by joint hormone

receptor and HER2 status. *J. Natl. Cancer Inst.* **106**, (2014).

38. Wright, J. L. *et al.* Racial disparity in estrogen receptor positive breast cancer patients

receiving trimodality therapy. *Breast* **21**, 276–283 (2012).

39. Dent, R. *et al.* Triple-negative breast cancer: Clinical features and patterns of

recurrence. *Clin. Cancer Res.* **13**, 4429–4434 (2007).

40. Perez, C. A. *et al.* Black race as a prognostic factor in triple-negative breast cancer

patients treated with breast-conserving therapy: a large, single-institution retrospective

analysis. *Breast Cancer Res. Treat.* **139**, 497–506 (2013).

41. Gleason, M. X., Mdzinarishvili, T. & Sherman, S. Breast cancer incidence in black and

white women stratified by estrogen and progesterone receptor statuses. *PLoS One* **7**,

e49359 (2012).

42. Martin, D. N. *et al.* Differences in the tumor microenvironment between African-

American and European-American breast cancer patients. *PLoS One* **4**, e4531 (2009).

43. Lindner, R. *et al.* Molecular phenotypes in triple negative breast cancer from African

American patients suggest targets for therapy. *PLoS One* **8**, e71915 (2013).

44. Nwosu, I. O., Tabler, D. D., Chipman, G. & Piccolo, S. R. Annotated Compendium of

102 Breast Cancer Gene-Expression Datasets. *bioRxiv* 2023.09.22.559045 (2023)

doi:10.1101/2023.09.22.559045.

45. Mikolajewicz, N. & Komarova, S. V. Meta-Analytic Methodology for Basic Research: A

875   Practical Guide. *Front. Physiol.* **10**, 203 (2019).

876   46. Edgar, R., Domrachev, M. & Lash, A. E. Gene Expression Omnibus: NCBI gene

877   expression and hybridization array data repository. *Nucleic Acids Res.* **30**, 207–210

878   (2002).

879   47. Barrett, T. *et al.* NCBI GEO: archive for functional genomics data sets--update. *Nucleic*

880   *Acids Res.* **41**, D991–5 (2013).

881   48. Parkinson, H. *et al.* ArrayExpress--a public database of microarray experiments and

882   gene expression profiles. *Nucleic Acids Res.* **35**, D747–50 (2007).

883   49. Curtis, C. *et al.* The genomic and transcriptomic architecture of 2,000 breast tumours

884   reveals novel subgroups. *Nature* **486**, 346–352 (2012).

885   50. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast

886   tumours. *Nature* **490**, 61–70 (2012).

887   51. Saal, L. H. *et al.* The Sweden Cancerome Analysis Network - Breast (SCAN-B)

888   Initiative: a large-scale multicenter infrastructure towards implementation of breast

889   cancer genomic analyses in the clinical routine. *Genome Med.* **7**, 20 (2015).

890   52. Staaf, J. *et al.* RNA sequencing-based single sample predictors of molecular subtype

891   and risk of recurrence for clinical assessment of early-stage breast cancer. *NPJ Breast*

892   *Cancer* **8**, 94 (2022).

893   53. Villatoro-García, J. A. *et al.* DExMA: An R Package for Performing Gene Expression

894   Meta-Analysis with Missing Genes. *Sci. China Ser. A Math.* **10**, 3376 (2022).

895   54. Core Team, R. R. No title. https://apps.dtic.mil/sti/citations/AD1039033.

896   55. Venet, D., Dumont, J. E. & Detours, V. Most random gene expression signatures are

897   significantly associated with breast cancer outcome. *PLoS Comput. Biol.* **7**, (2011).

898   56. Refaeilzadeh, P., Tang, L. & Liu, H. Cross-Validation. in *Encyclopedia of Database*

899   *Systems* (eds. Liu, L. & Özsu, M. T.) 532–538 (Springer US, Boston, MA, 2009).

57. Lei, J. Cross-Validation With Confidence. *J. Am. Stat. Assoc.* **115**, 1978–1997 (2020).

58. Pedregosa, F. *et al.* Scikit-learn: Machine learning in Python. *the Journal of machine*

Learning research **12**, 2825–2830 (2011).

59. Breiman, L. Random Forests. *Mach. Learn.* **45**, 5–32 (2001).

60. Piccolo, S. R., Mecham, A., Golightly, N. P., Johnson, J. L. & Miller, D. B. The ability to

classify patients based on gene-expression data varies by algorithm and performance

metric. *PLoS Comput. Biol.* **18**, e1009926 (2022).

61. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for

interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**,

15545–15550 (2005).

62. Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**,

1739–1740 (2011).

63. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set

collection. *Cell Syst* **1**, 417–425 (2015).

64. Foster, E. D. & Deardorff, A. Open Science Framework (OSF). *J. Med. Libr. Assoc.* **105**,

203 (2017).

65. Merkel, D. Docker: lightweight Linux containers for consistent development and

deployment. *Linux J.* **2014**, 2 (2014).

66. Bose, R. *et al.* Activating HER2 mutations in HER2 gene amplification negative breast

cancer. *Cancer Discov.* **3**, 224–237 (2013).

67. Baselga, J. Why the epidermal growth factor receptor? The rationale for cancer therapy.

Oncologist **7 Suppl 4**, 2–8 (2002).

68. Godoy-Ortiz, A. *et al.* Deciphering HER2 Breast Cancer Disease: Biological and Clinical

Implications. *Front. Oncol.* **9**, 1124 (2019).

69. Lanari, C. & Molinolo, A. A. Progesterone receptors - animal models and cell signaling

in breast cancer Diverse activation pathways for the progesterone receptor - possible

implications for breast biology and cancer. *Breast Cancer Res.* **4**, 1–4 (2002).

70. Hua, H., Zhang, H., Kong, Q. & Jiang, Y. Mechanisms for estrogen receptor expression

in human cancer. *Exp. Hematol. Oncol.* **7**, 24 (2018).

71. Navasa, N. *et al.* Ikaros mediates the DNA methylation-independent silencing of

MCJ/DNAJC15 gene expression in macrophages. *Sci. Rep.* **5**, 14692 (2015).

72. Sinha, D. & D'Silva, P. Chaperoning mitochondrial permeability transition: regulation of

transition pore complex by a J-protein, DnaJC15. *Cell Death Dis.* **5**, e1101 (2014).

73. Jarrous, N. & Gopalan, V. Archaeal/eukaryal RNase P: subunits, functions and RNA

diversification. *Nucleic Acids Res.* **38**, 7885–7894 (2010).

74. Abu-Zhayia, E. R. *et al.* A role of human RNase P subunits, Rpp29 and Rpp21, in

homology directed-repair of double-strand breaks. *Sci. Rep.* **7**, 1002 (2017).

75. Guyon, A. CXCL12 chemokine and its receptors as major players in the interactions

between immune and nervous systems. *Front. Cell. Neurosci.* **8**, 65 (2014).

76. Meng, W., Xue, S. & Chen, Y. The role of CXCL12 in tumor microenvironment. *Gene*

641, 105–110 (2018).

77. Ning, L., Wang, L., Zhang, H., Jiao, X. & Chen, D. Eukaryotic translation initiation factor

5A in the pathogenesis of cancers. *Oncol. Lett.* **20**, 81 (2020).

78. Nakanishi, S. & Cleveland, J. L. Targeting the polyamine-hypusine circuit for the

prevention and treatment of cancer. *Amino Acids* **48**, 2353–2362 (2016).

79. Bauer, M. F., Sirrenberg, C., Neupert, W. & Brunner, M. Role of Tim23 as voltage

sensor and presequence receptor in protein import into mitochondria. *Cell* **87**, 33–41

(1996).

80. Elvidge, G. P. *et al.* Concordant regulation of gene expression by hypoxia and 2-

oxoglutarate-dependent dioxygenase inhibition: the role of HIF-1alpha, HIF-2alpha, and

other pathways. *J. Biol. Chem.* **281**, 15215–15226 (2006).

81. Miziak, P. *et al.* Estrogen Receptor Signaling in Breast Cancer. *Cancers* **15**, (2023).

82. Takaku, M. *et al.* GATA3 zinc finger 2 mutations reprogram the breast cancer

transcriptional network. *Nat. Commun.* **9**, 1059 (2018).

83. Theodorou, V., Stark, R., Menon, S. & Carroll, J. S. GATA3 acts upstream of FOXA1 in

mediating ESR1 binding by shaping enhancer accessibility. *Genome Res.* **23**, 12–22

(2013).

84. Badve, S. *et al.* FOXA1 expression in breast cancer--correlation with luminal subtype A

and survival. *Clin. Cancer Res.* **13**, 4415–4421 (2007).

85. Wilson, B. J. & Giguère, V. Meta-analysis of human cancer microarrays reveals GATA3

is integral to the estrogen receptor alpha pathway. *Mol. Cancer* **7**, 49 (2008).

86. Li, Y. *et al.* High expression of carbonic anhydrase 12 (CA12) is associated with good

prognosis in breast cancer. *Neoplasma* **66**, 420–426 (2019).

87. Hurtado, A., Holmes, K. A., Ross-Innes, C. S., Schmidt, D. & Carroll, J. S. FOXA1 is a

key determinant of estrogen receptor function and endocrine response. *Nat. Genet.* **43**,

27–33 (2011).

88. Seachrist, D. D., Anstine, L. J. & Keri, R. A. FOXA1: A Pioneer of Nuclear Receptor

Action in Breast Cancer. *Cancers* **13**, (2021).

89. Kothari, C. *et al.* TBC1D9: An Important Modulator of Tumorigenesis in Breast Cancer.

Cancers **13**, (2021).

90. Wu, J.-R., Zhao, Y., Zhou, X.-P. & Qin, X. Estrogen receptor 1 and progesterone

receptor are distinct biomarkers and prognostic factors in estrogen receptor-positive

breast cancer: Evidence from a bioinformatic analysis. *Biomed. Pharmacother.* **121**,

109647 (2020).

91. Kurozumi, S. *et al.* Power of PgR expression as a prognostic factor for ER-

positive/HER2-negative breast cancer patients at intermediate risk classified by the Ki67

labeling index. *BMC Cancer* **17**, 354 (2017).

92. Hao, N. *et al.* High PGAP3 expression is associated with lymph node metastasis and

low CD8+T cell in patients with HER2+ breast cancer. *Pathol. Res. Pract.* **251**, 154890

(2023).

93. Fararjeh, A. F. S., Al Khader, A., Kaddumi, E., Obeidat, M. & Al-Fawares, O. 'la.

Differential Expression and Prognostic Significance of STARD3 Gene in Breast

Carcinoma. *Int J Mol Cell Med* **10**, 34–41 (2021).

94. Kauraniemi, P. & Kallioniemi, A. Activation of multiple cancer-associated genes at the

ERBB2 amplicon in breast cancer. *Endocr. Relat. Cancer* **13**, 39–49 (2006).

95. Vermehren-Schmaedick, A., Mhawech-Fauceglia, P., Park, B. S., Pejovic, T. & Luoh,

S.-W. The prognostic significance of GRB7 protein expression and localization in human

breast and ovarian cancers. *Oncotarget* **11**, 2273–2289 (2020).

96. Bivin, W. W., Yergiyev, O., Bunker, M. L., Silverman, J. F. & Krishnamurti, U. GRB7

Expression and Correlation With HER2 Amplification in Invasive Breast Carcinoma.

Appl. Immunohistochem. Mol. Morphol. **25**, 553–558 (2017).

97. Choi, S. H. *et al.* CDK12 phosphorylates 4E-BP1 to enable mTORC1-dependent

translation and mitotic genome stability. *Genes Dev.* **33**, 418–435 (2019).

98. Lei, T. *et al.* Cyclin K regulates prereplicative complex assembly to promote mammalian

cell proliferation. *Nat. Commun.* **9**, 1876 (2018).

99. Malumbres, M. & Barbacid, M. Cell cycle, CDKs and cancer: a changing paradigm. *Nat.*

Rev. Cancer **9**, 153–166 (2009).

100.Liang, S. *et al.* CDK12: A Potent Target and Biomarker for Human Cancer Therapy.

Cells **9**, (2020).

101.Fararjeh, A. S. *et al.* Proteasome 26S Subunit, non-ATPase 3 (PSMD3) Regulates

Breast Cancer by Stabilizing HER2 from Degradation. *Cancers* **11**, (2019).

102.Lodi, M. *et al.* STARD3: A New Biomarker in HER2-Positive Breast Cancer. *Cancers*

15, (2023).

103.Park, H. K., Kwak, M. K., Kim, H. J. & Ahima, R. S. Linking resistin, inflammation, and

cardiometabolic diseases. *Korean J. Intern. Med.* **32**, 239–247 (2017).

104.Steppan, C. M. *et al.* The hormone resistin links obesity to diabetes. *Nature* **409**, 307–

312 (2001).

105.Dalamaga, M. Resistin as a biomarker linking obesity and inflammation to cancer:

potential clinical perspectives. *Biomark. Med.* **8**, 107–118 (2014).

106.Lee, J. O. *et al.* Resistin, a fat-derived secretory factor, promotes metastasis of MDA-

MB-231 human breast cancer cells through ERM activation. *Sci. Rep.* **6**, 18923 (2016).

107.Deb, A., Deshmukh, B., Ramteke, P., Bhati, F. K. & Bhat, M. K. Resistin: A journey from

metabolism to cancer. *Transl. Oncol.* **14**, 101178 (2021).

108.Kumar, S., Rao, N. & Ge, R. Emerging Roles of ADAMTSs in Angiogenesis and Cancer.

Cancers **4**, 1252–1299 (2012).

109. Redondo-García, S., Peris-Torres, C., Caracuel-Peramos, R. & Rodríguez-

Manzaneque, J. C. ADAMTS proteases and the tumor immune microenvironment:

Lessons from substrates and pathologies. *Matrix Biol Plus* **9**, 100054 (2021).

110. Van Veldhoven, P. P. Biochemistry and genetics of inherited disorders of peroxisomal

fatty acid metabolism. *J. Lipid Res.* **51**, 2863–2895 (2010).

111. O'Hurley, G. *et al.* Analysis of the Human Prostate-Specific Proteome Defined by

Transcriptomics and Antibody-Based Profiling Identifies TMEM79 and ACOXL as Two

Putative, Diagnostic Markers in Prostate Cancer. *PLoS One* **10**, e0133449 (2015).

112. Wang, C. *et al.* FAM3A activates PI3K p110α/Akt signaling to ameliorate hepatic

gluconeogenesis and lipogenesis. *Hepatology* **59**, 1779–1790 (2014).

113. Yan, H. *et al.* FAM3A maintains metabolic homeostasis by interacting with F1-ATP

synthase to regulate the activity and assembly of ATP synthase. *Metabolism* **139**,

155372 (2023).

114. Dong, Q.-T. *et al.* FAM3 family genes are associated with prognostic value of human

cancer: a pan-cancer analysis. *Sci. Rep.* **13**, 15144 (2023).

115. Takkar, S. *et al.* From orphan to oncogene: The role of GPR35 in cancer and immune

modulation. *Cytokine Growth Factor Rev.* (2024) doi:10.1016/j.cytogfr.2024.03.004.

116. Milligan, G. GPR35: from enigma to therapeutic target. *Trends Pharmacol. Sci.* **44**, 263–

273 (2023).

117. Yashin, D. V., Sashchenko, L. P. & Georgiev, G. P. Mechanisms of Action of the

PGLYRP1/Tag7 Protein in Innate and Acquired Immunity. *Acta Naturae* **13**, 91–101

(2021).

118. Bhusal, A. *et al.* The microglial innate immune protein PGLYRP1 mediates

neuroinflammation and consequent behavioral changes. *Cell Rep.* **43**, 113813 (2024).

119. Schnell, A. *et al.* Targeting PGLYRP1 promotes antitumor immunity while inhibiting

autoimmune neuroinflammation. *Nat. Immunol.* **24**, 1908–1920 (2023).

120. Nwosu, I. O. & Piccolo, S. R. A systematic review of datasets that can help elucidate

relationships among gene expression, race, and immunohistochemistry-defined

subtypes in breast cancer. *Cancer Biol. Ther.* **22**, 417–429 (2021).

121.Mehra, R. *et al.* Identification of GATA3 as a breast cancer prognostic marker by global

gene expression meta-analysis. *Cancer Res.* **65**, 11259–11264 (2005).

122.Goldhirsch, A. *et al.* Personalizing the treatment of women with early breast cancer:

highlights of the St Gallen International Expert Consensus on the Primary Therapy of

Early Breast Cancer 2013. *Ann. Oncol.* **24**, 2206–2223 (2013).

123.Dowsett, M. *et al.* Prognostic value of Ki67 expression after short-term presurgical

endocrine therapy for primary breast cancer. *J. Natl. Cancer Inst.* **99**, 167–170 (2007).

124.Stover, D. G. *et al.* The Role of Proliferation in Determining Response to Neoadjuvant

Chemotherapy in Breast Cancer: A Gene Expression-Based Meta-Analysis. *Clin.*

Cancer Res. **22**, 6039–6050 (2016).

125.Julka, P. K. *et al.* A phase II study of sequential neoadjuvant gemcitabine plus

doxorubicin followed by gemcitabine plus cisplatin in patients with operable breast

cancer: prediction of response using molecular profiling. *Br. J. Cancer* **98**, 1327–1335

(2008).

126.Chang, J. C. *et al.* Gene expression profiling for the prediction of therapeutic response

to docetaxel in patients with breast cancer. *Lancet* **362**, 362–369 (2003).

127.Burstein, M. D. *et al.* Comprehensive genomic analysis identifies novel subtypes and

targets of triple-negative breast cancer. *Clin. Cancer Res.* **21**, 1688–1698 (2015).

128.Ellsworth, R. E. *et al.* A gene expression signature that defines breast cancer

metastases. *Clin. Exp. Metastasis* **26**, 205–213 (2009).

129.Loi, S. *et al.* Gene expression profiling identifies activated growth factor signaling in poor

prognosis (Luminal-B) estrogen receptor positive breast cancer. *BMC Med. Genomics*

2, 37 (2009).

130.Schneider, J. *et al.* Identification and meta-analysis of a small gene expression

signature for the diagnosis of estrogen receptor status in invasive ductal breast cancer.

Int. J. Cancer **119**, 2974–2979 (2006).

131. McKinley, K. L. & Cheeseman, I. M. The molecular basis for centromere identity and function. *Nat. Rev. Mol. Cell Biol.* **17**, 16–29 (2016).

132. Bauer, N. G., Richter-Landsberg, C. & Ffrench-Constant, C. Role of the oligodendroglial cytoskeleton in differentiation and myelination. *Glia* **57**, 1691–1705 (2009).

133. Neefjes, J., Jongsma, M. L. M., Paul, P. & Bakke, O. Towards a systems understanding of MHC class I and MHC class II antigen presentation. *Nat. Rev. Immunol.* **11**, 823–836 (2011).

134. Matsuda, T. *et al.* Signal-transducing adaptor protein-2 modulates T-cell functions. *Explor. Immunol.* 771–782 (2022).

135. He, X.-Y., Isaacs, C. & Yang, S.-Y. Roles of Mitochondrial 17β-Hydroxysteroid Dehydrogenase Type 10 in Alzheimer's Disease. *J. Alzheimers. Dis.* **62**, 665–673 (2018).

136. Amelio, I., Cutruzzolá, F., Antonov, A., Agostini, M. & Melino, G. Serine and glycine metabolism in cancer. *Trends Biochem. Sci.* **39**, 191–198 (2014).

137. Phang, J. M., Liu, W., Hancock, C. & Christian, K. J. The proline regulatory axis and cancer. *Front. Oncol.* **2**, 60 (2012).

138. Siddharth, S. & Sharma, D. Racial disparity and triple-negative breast cancer in African-American women: A multifaceted affair between obesity, biology, and socioeconomic determinants. *Cancers* **10**, (2018).

139. Dietze, E. C., Sistrunk, C., Miranda-Carboni, G., O'Regan, R. & Seewaldt, V. L. Triple-negative breast cancer in African-American women: disparities versus biology. *Nat. Rev. Cancer* **15**, 248–254 (2015).

140. Zheng, L., Cao, L. & Zheng, X. L. ADAMTS13 protease or lack of von Willebrand factor protects irradiation and melanoma-induced thrombotic microangiopathy in zebrafish. *J. Thromb. Haemost.* **20**, 2270–2283 (2022).

141. López-Gil, J. C. *et al.* The Peptidoglycan Recognition Protein 1 confers immune evasive properties on pancreatic cancer stem cells. *Gut* (2024) doi:10.1136/gutjnl-2023-330995.

142. Cui, X. *et al.* TNF upregulates peptidoglycan recognition protein 1 in esophageal cancer

cells to clear the path to its signaling: Making the 'enemy' a friend. *Arch. Biochem.*

Biophys. **722**, 109192 (2022).

143. Quon, T., Lin, L.-C., Ganguly, A., Tobin, A. B. & Milligan, G. Therapeutic Opportunities

and Challenges in Targeting the Orphan G Protein-Coupled Receptor GPR35. *ACS*

Pharmacol Transl Sci **3**, 801–812 (2020).

144. Guo, Y. J., Zhou, Y. J., Yang, X. L., Shao, Z. M. & Ou, Z. L. The role and clinical

significance of the CXCL17-CXCR8 (GPR35) axis in breast cancer. *Biochem. Biophys.*

Res. Commun. **493**, 1159–1167 (2017).

145. Pitteri, S. J. *et al.* Tumor microenvironment-derived proteins dominate the plasma

proteome response during breast cancer induction and progression. *Cancer Res.* **71**,

5090–5100 (2011).

146. Wang, C.-Q. *et al.* Impacts of RETN genetic polymorphism on breast cancer

development. *J. Cancer* **11**, 2769–2777 (2020).

147. Aziz, M. A., Akter, T., Sarwar, M. S. & Islam, M. S. The first combined meta‑analytic

approach for elucidating the relationship of circulating resistin levels and RETN gene

polymorphisms with colorectal and breast cancer. *Egyptian Journal of Medical Human*

Genetics **23**, 1–19 (2022).

148. Smedley, A. 'race' and the construction of human identity. *Am. Anthropol.* **100**, 690–702

(1998).

149. Rusidzé, M. *et al.* Estrogen receptor-α signaling in post-natal mammary development

and breast cancers. *Cell. Mol. Life Sci.* **78**, 5681–5705 (2021).

150. Asselin-Labat, M.-L. *et al.* Gata-3 is an essential regulator of mammary-gland

morphogenesis and luminal-cell differentiation. *Nat. Cell Biol.* **9**, 201–209 (2007).

151. Watson, P. H. *et al.* Carbonic anhydrase XII is a marker of good prognosis in invasive

breast carcinoma. *Br. J. Cancer* **88**, 1065–1070 (2003).

152. Allison, K. H. *et al.* Estrogen and Progesterone Receptor Testing in Breast Cancer:

American Society of Clinical Oncology/College of American Pathologists Guideline

Update. *Arch. Pathol. Lab. Med.* **144**, 545–563 (2020).

153. Scabia, V. *et al.* Estrogen receptor positive breast cancers have patient specific

hormone sensitivities and rely on progesterone receptor. *Nat. Commun.* **13**, 3127

(2022).

154. Swain, S. M., Shastry, M. & Hamilton, E. Targeting HER2-positive breast cancer:

advances and future directions. *Nat. Rev. Drug Discov.* **22**, 101–126 (2023).

155. Madrid-Paredes, A., Cañadas-Garre, M., Sánchez-Pozo, A. & Calleja-Hernández, M. Á.

Non-HER2 signaling pathways activated in resistance to anti-HER2 therapy in breast

cancer. *Breast Cancer Res. Treat.* **153**, 493–505 (2015).

156. Sahlberg, K. K. *et al.* The HER2 amplicon includes several genes required for the

growth and survival of HER2 positive breast cancer cells. *Mol. Oncol.* **7**, 392–401

(2013).

157. Lehmann, B. D. *et al.* Identification of human triple-negative breast cancer subtypes and

preclinical models for selection of targeted therapies. *J. Clin. Invest.* **121**, 2750–2767

(2011).

158. Pegram, M., Jackisch, C. & Johnston, S. R. D. Estrogen/HER2 receptor crosstalk in

breast cancer: combination therapies to improve outcomes for patients with hormone

receptor-positive/HER2-positive breast cancer. *NPJ Breast Cancer* **9**, 45 (2023).

159. Burguin, A., Diorio, C. & Durocher, F. Breast Cancer Treatments: Updates and New

Challenges. *J Pers Med* **11**, (2021).

160. Mazumder, A., Shiao, S. & Haricharan, S. HER2 Activation and Endocrine Treatment

Resistance in HER2-negative Breast Cancer. *Endocrinology* **162**, (2021).