A Comprehensive Meta-Analysis of Breast Cancer Gene Expression

Ifeanyichukwu O. Nwosu

Department of Biology

Brigham Young University

Provo, UT, USA

inwosu@byu.edu

Stephen R. Piccolo*

Department of Biology

Brigham Young University

Provo, UT, USA

stephen_piccolo@byu.edu

*Please address correspondence to S.R.P.

Abstract

Background

Triple-negative breast cancers (TNBC) occur more frequently in African Americans and are associated with worse outcomes when compared to other subtypes of breast cancer. These cancers lack expression of estrogen receptor (ER), progesterone receptor (PR) and human epidermal growth factor receptor 2 (HER2) and have limited treatment options. To shed light on mechanisms behind these differences and suggest novel treatments, we used a meta-analytic approach to identify gene expression differences in breast tumors for people with self-reported African or European ancestry; additionally, we compared gene expression levels based on ER, PR, HER2 and TNBC status.

Methods

After gathering and standardizing gene expression data and metadata from 106 datasets (representing 27,000 samples), we identified genes that were expressed differently between these groups via random-effects meta-analyses. To evaluate the robustness of these gene lists, we devised a novel computational methodology that uses cross validation and classification. We also computed overlaps between the most significant genes and known signaling pathways.

Results

Using a false discovery rate threshold of 0.05, we identified genes that are known to play a significant role in their respective breast cancer subtypes (e.g., *ESR1* for ER status and *ERBB2* for HER2 status), thus confirming the validity of our findings. We also discovered genes that have not been reported previously and may be new targets for breast cancer therapy. *GATA3*, *CA12*, *TBC1D9*, *XBP1* and *FOXA1* were among the most significant genes for ER, PR, and TNBC. However, none of these genes overlapped with HER2 status, supporting prior research that HER2 tumors are mechanistically different from endocrine

breast cancers. The genes identified from the race meta-analysis—including *DNAJC15*, *HLA-DPA1*, *STAP2*, *CEP68*, *MOGS*—have not been associated previously with race-specific breast-cancer outcomes, highlighting a potential area of further research.

Conclusions

We have carried out a large meta-analysis of breast cancer gene expression data, identifying novel genes that may serve as potential biomarkers for breast cancer in diverse populations. We have also developed a computational method that identifies gene sets small enough to be analyzed and explored in future studies. This method has the potential to be applied to other cancers.

Keywords

Breast cancer, Racial disparity, Triple negative Breast Cancer, Meta-analysis, Machine Learning.

Background

Cancers arise as a result of alterations in groups of genes known as proto-oncogenes and tumor suppressor genes[1,2]. These definitions cover hundreds of genes[3,4] that have been implicated in tumorigenic processes, with an average of 33 to 66 mutations found in solid tumors[5]. This represents an enormous amount of genetic variation, even within the same tissue types[6], making it difficult to develop a "one size fits all" approach to treatments, or to understanding cancer biology in general. Among cancers, breast cancer is the most common type of cancer in women worldwide, and accounts for 12.5% of new cancer cases in the world each year[7,8]. It is also the second most common cause of cancer deaths among women in the U.S. In 2023, the American Cancer Society estimated 297,790 new cases of invasive breast cancer diagnosed in women in the US alone[9]. This is an enormous public health burden. Advances in treatment, coupled with early diagnoses, have improved survival rates and lowered mortality[10,11] for breast cancer patients. However, researchers continue to search for methods to assign the most appropriate types of therapies[12,13] and thus to tailor treatments to patients.

Despite these advancements, the heterogeneous nature of the disease[14] and the complex genetic landscape[15] still make it difficult to unravel the underlying molecular mechanisms governing its pathogenesis and progression. Furthermore, there exist racial disparities among population groups[16]. For example, the American Cancer Society reports that survival rates for all cancers are lower for individuals of African descent when compared to those of Caucasian descent[17]. These differences are largely due to cancers of the breast in women, but this is also true for prostate[18] and lung[19] cancers. Various studies[20–22] have highlighted the multifactorial causes of these disparities. These include socioeconomic factors such as access to healthcare, income levels, as well as treatment delays and trust in physicians. However, these factors do not fully explain these disparities[23,24]. The question thus arises as to factors potentially driven by biological causes.

Gene expression is the process by which the information encoded in genes is used to direct the synthesis of proteins via RNA transcripts. It is assessed by measuring the number of RNA transcripts for each gene in a tissue sample; the sum of a sample's RNA transcripts is referred to as its transcriptome. High-throughput genomic technologies have enabled researchers to produce huge amounts of gene-expression data[25–27]. The analysis of gene expression data is a powerful tool for deciphering molecular signatures associated with different breast cancer subtypes[28], predicting patient prognoses[29] and guiding therapeutic decision-making[30]. Extensive research has shown that studying the transcriptome can help to identify functional mechanisms behind biological processes. This is validated by a number of studies which have explored gene expression patterns in cancer; these have found patterns associated with race[31–34] and identified connections between race and hormone receptor status[35,36]. These studies, however, have used a limited number of samples (between 52 and 627 per study). Some of these datasets are publicly available, while others are not.

Traditionally, clinical subtypes of breast cancer have been defined based on a patient's immunohistochemistry (IHC) profile. These profiles are determined by the combined expression status of three receptors: estrogen receptor (ER), progesterone receptor (PR), and human epidermal growth factor receptor 2 (HER2)[37]. It has also been observed that women of African descent more commonly have tumors negative for ER, PR and HER2[38]. These cancers are typically referred to as triple negative breast cancers (TNBC). TNBCs are associated with worse outcomes when compared to other subtypes[39] and occur more frequently in African Americans[40,41] than other groups. Some studies have identified biological processes that may contribute to enhanced disease aggressiveness in African-American patients, including angiogenesis, chemotaxis[42], and the insulin-like growth factor 1 (IGF1) pathway[43].

Building on a large curation effort that we performed with publicly available data[44], we identified gene-expression datasets that include race metadata and/or metadata for ER, PR and HER2 status. In total, we have data for over 27,000 samples across 47 datasets with ER status, 35 datasets with PR status, 28 datasets with ER status and 7 datasets with race information. With these datasets, we applied a meta-analytic method that combines data from multiple independent scientific studies, with the aim of identifying consistent patterns[45]. Our research questions can be broadly divided into: A) which genes are most consistently expressed differently between individuals who self-identify as having European ancestry or African ancestry, B) which genes are consistently differentially expressed between individuals who are positive or negative for ER, PR, HER2, or TNBC status, and C) which signaling pathways are associated with these genes. By addressing these questions, we aim to uncover molecular mechanisms underlying breast cancer disparities and hormone receptor statuses, potentially informing personalized treatment strategies and improving outcomes across diverse populations.

Methods

Data Acquisition

In a prior study[44], we gathered 114 gene expression datasets from public repositories including Gene Expression Omnibus[46,47], ArrayExpress[48], the Molecular Taxonomy of Breast Cancer International Consortium (METABRIC) project[49], TCGA[50] and the SCAN-B Initiative[51,52] in a bid to enable researchers to address a variety of questions related to breast-cancer transcriptomics. For this study, we used a subset of those datasets (Additional File 1). We identified datasets for which one or more of the following metadata variables were available: self-reported race/ancestry, ER status, PR status, and/or HER2 status.

Data Preparation

For the comparison based on self-reported ancestry, we filtered the data to focus on people described as "White", "Caucasian" or "European American" and people described as "Black"

or "African American". We focused on these groups because they were most common in the data and because an understanding of disparities between these groups is critically needed. We removed samples that did not fall into either of these categories. Next, we standardized the metadata variable names because different studies used different terms to represent the same variables. Samples that were labeled "Caucasian", "W", "w", "European American", or "white" were all categorized as "White", while samples labeled "African American", "B", "b", "Black or African American", or "black" were categorized as "Black". For the cell-surface receptors (ER, PR and HER2), we standardized each value as "positive" or "negative," as determined by the original authors. We excluded samples with missing values for these variables. For ambiguous data where we could not readily identify what the values referred to, we referred to the journal article(s) associated with each dataset. If we found more information to clarify what those values meant, we standardized and included them in our study; otherwise, we filtered them out. For example, in one study, we found that the values "0/1", "0-1", "1+", and "1+ (only 1 core pos.)" referred to HER2 negative samples. When determining TNBC status, we implemented logic to handle missing values. We marked a tumor as TNBC-negative only if it was known that the tumor was negative for all three markers (no missing data). If a tumor was positive for any marker, we inferred that the tumor was *not* TNBC-negative, even if other markers were missing. Additionally, we identified tumors for which race and TNBC status were available and performed a comparison between individuals of African or European ancestry.

For each comparison (Black and White, ER+ vs ER-, PR+ vs PR-, HER2+ vs HER2-, TNBC vs non-TNBC, and Black TNBC vs White TNBC), we selected datasets that had at least 10 samples in each group. For example, when comparing ER status, we required a minimum of 10 "ER+" samples and 10 "ER-" samples. We also determined the relative proportion of the groups and ensured the ratio between groups was no larger than 20:1 (in either direction). We did this to prevent our results from being skewed due to class imbalance.

Meta-analysis

Different gene-expression profiling platforms and annotations produce data for different genes. Therefore, for each comparison, we first selected genes that were common across all datasets being used for that comparison. There were 11,049 common genes for race, 10,809 common genes for ER status, 10,960 common genes for PR status, 10,926 common genes for HER2 status, 10,960 for triple negative status, and 11,049 for race + TNBC.

We carried out the meta-analyses using the DExMA[53] package (version 1.10.7) in R[54](version 4.3.2). We used the following parameters:

i) typeMethod: random effects model;

ii) missAllow: The maximum proportion of missing values allowed in a sample = 10%;

iii) proportionData: The minimum proportion of datasets in which a gene must be present = 90%.

After the meta-analysis was completed, we filtered the results based on False Discovery Rate (FDR) values less than 0.05 and arranged them in descending order based on the absolute value of the effect size. This ensured that the top-ranked genes were both statistically significant and had a relatively large effect size.

Meta-analyses commonly result in thousands of statistically significant genes. It is difficult to interpret such large lists. We wished to estimate the number of significant genes that would effectively discriminate between the groups while being as parsimonious as possible. Furthermore, Venet et al[55] discuss that empirically identified gene-expression signatures are sometimes no more predictive of breast-cancer outcomes than random gene expression signatures, even when the specified genes are statistically significant. Thus, we wished to verify that our identified gene lists were more biologically relevant than what we could obtain using randomly selected genes. To address these goals, we developed a process that used leave-one-dataset-out cross validation. Cross validation is a statistical method to evaluate

the performance of a model on unseen data in an unbiased manner[56,57]. Rather than holding out individual samples, we held out one dataset at a time. First, we carried out a meta-analysis using $n - 1$ datasets at a time (Figure 1). We assumed that if genes identified through an n - 1 meta-analysis are biologically relevant, expression values for those genes would be effective at classifying the two groups being compared on the held-out dataset. The second step was to perform classification analyses and quantify the predictive performance relative to randomly selected genes. For this step, we performed within-dataset cross validation for the held-out dataset. We used modules from the scikit-learn library (version 1.3.1)[58] for this step. We split the datasets using the StratifiedKFold method because it ensures that each fold of the dataset has a similar proportion of observations with a given label. It is particularly useful when dealing with classification tasks that have imbalanced class distributions, as observed in our datasets. We also used the Random Forest[59] classifier as it has been shown to perform well on gene expression data[60]. For m top-ranked genes (5, 10, 20, 50, 100, 500, or 1000), we performed classification and calculated a score for each cross-validation fold. We used the "balanced accuracy" scoring metric to account for the imbalanced nature of the datasets. We then repeated this process 100 times for m genes selected at random from the held-out dataset. We ensured that each random gene had not been identified as relevant in the meta-analysis. Consider the race datasets as an example. For each round of meta-analysis, we selected six of the seven data sets, leaving one out, and we performed the meta-analysis on those datasets. The meta-analysis produced a list of genes. From this list, we selected the top 5, 10, 20, 50, 100, 500 and 1000 genes and evaluated how accurately the expression data for those genes predicted race for the seventh data set.

Figure 1. Diagram illustrating the leave-one-dataset-out cross-validation method. For a given fold, one dataset was held out for testing. Via meta-analysis, significant genes were identified and ranked according to fold change. Within-dataset cross validation was performed for the held-out test dataset, and the labels were predicted using a classification algorithm for *m* top-ranked genes. The same process was performed for randomly selected genes and repeated 100 times. We compared the balanced accuracy for the genes identified via meta-analysis vs. the genes identified via random selection.

Pathway Analysis

Based on the cross-validation results, we investigated the gene lists using the Molecular Signatures Database (MSigDB)[61,62] which has thousands of annotated gene sets. Examining the overlaps of these gene sets can help identify dysregulated pathways and underlying biological themes. To compute these overlaps, we pasted the top 20 genes from our gene lists into the *Investigate* tab on the MSigDB website (https://www.gsea-msigdb.org/gsea/msigdb/human/annotate.jsp). In particular, we computed overlaps using the "H: hallmark" gene sets"[63] and the "C2: curated gene sets," using an FDR q-value threshold of 0.05.

Results

We performed meta-analyses to compare gene-expression levels between people of African or European descent and between people with positive or negative tumor status (ER, PR, HER2, or TNBC). There was some overlap across the analyses; some datasets had more than one of these variables (Table 1). Additionally, we compared TNBC tumors from people of African descent versus people of European descent.

Variable	Positive	Negative	Equivocal	Unknown	Total Number of Samples	Number of Datasets
	White	**Black**				
Race	1,175	295	NA	NA	1,470	7
Estrogen Receptor Status	16,174	4,426	26	1,334	21,960	47
Progesterone Receptor Status	11,472	5,359	96	1,214	18,141	35
HER2/Neu Status	2,471	13,429	179	1,482	17,561	28
	Triple Negative	**Non Triple Negative**				
Triple Negative Status	1,982	13,614	NA	NA	15,596	22

Table 1. Dataset and sample counts for the meta-analyses. For some datasets, the original authors reported hormone receptor status but were unable to determine whether certain samples were positive or negative due to ambiguous immunohistochemistry results or when these values were marked as intermediate. These samples are listed as "Equivocal." In other cases, the status was reported for some samples but not reported for others. These samples are listed as "Unknown."

For the comparison between individuals of African or European ancestry, 850 genes were significantly differentially expressed (Additional File 2). For the comparisons between ER-positive and ER-negative breast tumors, 9,729 genes were differentially expressed (Additional File 3). We observed similarly high numbers for PR and HER2: 9,047 and 7,236,

respectively (Additional Files 4-5). For the TNBC versus non-TNBC comparison, 8,939 genes were differentially expressed (Additional File 6). For the comparison between people of African or European ancestry specific to TNBC tumors, six genes were differentially expressed (Additional File 7).

To identify a smaller number of genes that should be considered relevant for each meta-analysis, we developed a new computational method. Leaving out one dataset at a time, we used the remaining datasets to identify significant genes and then performed classification via cross validation on the held-out dataset. We compared the predictive performance of *m* genes (as ranked by each meta-analysis) against the predictive performance of randomly selected genes, to estimate the number of significant genes that would be most predictive. This analysis revealed that 20 genes often performed well compared to other thresholds (Figures 2-6). Although this threshold did not always perform better than other thresholds, we chose 20 genes as the threshold for all six meta-analyses for consistency. Twenty genes is few enough that researchers can interpret such a list and potentially create a panel to use in a laboratory setting. Table 2 shows the top 20 genes for each comparison. *GATA3, CA12, XBP1* and *FOXA1* were among the top 20 genes for ER, PR, and TNBC. However, none of these overlapped with the genes for HER2 status (or race).

Rank	Race	ER Status	PR Status	Her2 Status	TNBC	Race + TNBC
1	DNAJC15	**GATA3**	PGR	ERBB2	**FOXA1**	RETN
2	HLA-DPA1	ESR1	**CA12**	PGAP3	MLPH	ADAMTS13
3	STAP2	**CA12**	MAPT	STARD3	FOXC1	ACOXL
4	CEP68	TBC1D9	**GATA3**	GRB7	**GATA3**	FAM3A
5	MOGS	**FOXA1**	ESR1	CDK12	EN1	GPR35
6	TNS1	MLPH	DNALI1	PSMD3	GABRP	PGLYRP1
7	TXNL4A	DNALI1	GREB1	MED1	**CA12**	
8	PSPH	NAT1	NAT1	GSDMB	BCL11A	

9	EIF5A	PSAT1	SCUBE2	MED24	SFT2D2	
10	RPP25	SLC39A6	IL6ST	PNMT	**XBP1**	
11	PKP3	**XBP1**	TBC1D9	TCAP	AGR2	
12	POP7	ABAT	ABAT	MSL1	FAM171A1	
13	CENPA	GREB1	GFRA1	CASC3	DSC2	
14	TTBK2	MAPT	**XBP1**	CWC25	TBC1D9	
15	CXCL12	THSD4	KDM4B	PSMB3	UGT8	
16	PYCR1	BCL11A	THSD4	PIP4K2B	SPDEF	
17	HSD17B10	GFRA1	CELSR1	LASP1	PSAT1	
18	TIMM23	AGR2	SLC39A6	RPL23	SOX10	
19	PCGF1	PHGDH	**FOXA1**	CLCA2	ART3	
20	ABCA6	DSC2	ERBB4	KMO	MICALL1	

Table 2. Top-ranked genes identified in the meta-analyses. These genes represent our findings when performing meta-analyses across all datasets for which each variable was available (i.e., not as part of our cross-validation experiments). Fewer than 20 genes are shown for "Race + TNBC" because this meta-analysis identified only 6 genes with statistical significance. Genes bolded and highlighted in red were common across more than one of the meta-analyses.

For the comparison between tumors from individuals of European or African descent, the median balanced accuracy for the top 20 genes was 0.57 (the maximum possible accuracy is 1.0). The median balanced accuracy for the randomly selected genes was 0.50. Using a one-sided, Mann-Whitney U test, we compared the balanced accuracy values when using the top 20 meta-analysis genes for classification, versus the balanced accuracies for the randomly selected genes; the meta-analysis accuracies were *not* significantly higher (p-value = 0.23; Figure 2). These findings suggest that gene-expression differences between these populations were not significantly different from randomly selected genes; however, sample sizes may have been a limiting factor.

For the ER-positive versus ER-negative comparison, the balanced accuracies were significantly higher for the meta-analysis genes than for the randomly selected genes (p < 0.001; Figure 3). The same was true for the PR, HER2, and TNBC comparisons (p < 0.001).

However, the overall patterns were markedly distinct for HER2 relative to the other comparisons (Figures 3-6). For nearly all thresholds, the balanced accuracy was above 0.8 when using the meta-analysis genes; however, for ER, PR, and TNBC, the random-gene accuracies started low (around 0.6) and increased considerably as the number of genes increased. This pattern suggests that ER and/or PR positivity cause a cascade of downstream effects, possibly altering the expression of thousands of genes. Thus, when we selected genes at random (not including the top-ranked genes), the classification algorithm was frequently able to classify the tumors as positive or negative. In contrast, for HER2, the median balanced accuracy was below 0.55 for all thresholds when randomly selected genes were used. This pattern suggests that overexpression of HER2 protein, which typically results from DNA amplification or somatic mutations[64], thus activating oncogenic pathways[65,66], causes relatively few downstream effects compared to endocrine breast cancers, which rely on estrogen or progesterone signaling for tumor growth[67,68].

Figure 2. Cross-validation results for race status. The left panel represents the balanced accuracy, identified via cross validation on each held-out test set, for different numbers of top-ranked genes from the race meta-analysis. Each data point represents the balanced accuracy for a particular dataset. The right panel represents the balanced accuracy for the same numbers of genes selected at random in 100 iterations. Each data point represents the mean balanced accuracy for a particular dataset, across the iterations. The balanced accuracy for the meta-analysis genes was relatively low (between 0.5 and 0.6) when using 5 to 200 genes; it was equivalent to baseline levels (0.5) when 500 or 1000 genes were used. The balanced accuracy for the randomly selected genes was consistently close to baseline levels.

Figure 3. Cross-validation results for ER Status. The left panel represents the balanced accuracy, identified via cross validation on each held-out test set, for different numbers of top-rank genes from the ER meta-analysis. Each data point represents the balanced accuracy for a particular dataset. The right panel represents the average balanced accuracy for the same numbers of genes selected at random in 100 iterations. Each data point represents the mean balanced accuracy for a particular dataset, across the iterations. The balanced accuracy for the meta-analysis genes was consistently high (around 0.85 to 0.9) regardless of the number of genes used, while the balanced accuracy for the random genes started low and increased as the number of genes increased.

Figure 4. Cross-validation results for PR Status. The left panel represents the balanced accuracy, identified via cross validation on each held-out test set, for different numbers of top-rank genes from the PR meta-analysis. Each data point represents the balanced accuracy for a particular dataset. The right panel represents the average balanced accuracy for the same numbers of genes selected at random in 100 iterations. Each data point represents the mean balanced accuracy for a particular dataset, across the iterations. The balanced accuracy for the meta-analysis genes was consistently high (around 0.8) regardless of the number of genes used, while the balanced accuracy for the random genes started low and increased as the number of genes increased.

Figure 5. Cross-validation results for HER2 status. The left panel represents the balanced accuracy, identified via cross validation on each held-out test set, for different numbers of top-rank genes from the HER2 meta-analysis. Each data point represents the balanced accuracy for a particular dataset. The right panel represents the average balanced accuracy for the same numbers of genes selected at random in 100 iterations. Each data point represents the mean balanced accuracy for a particular dataset, across the iterations. The balanced accuracy for the meta-analysis genes was consistently higher than the balanced accuracy for the random genes.

Figure 6. Cross-validation results for TNBC status. The left panel represents the balanced accuracy, identified via cross validation on each held-out test set, for different numbers of top-rank genes from the TNBC meta-analysis. Each data point represents the balanced accuracy for a particular dataset. The right panel represents the average balanced accuracy for the same numbers of genes selected at random in 100 iterations. Each data point represents the mean balanced accuracy for a particular dataset, across the iterations. The balanced accuracy for the meta-analysis genes was consistently higher than the balanced accuracy for the random genes.

For the comparison between individuals of African or European descent, some of the top genes were *DNAJC15*, *RPP25*, *CXCL12*, *EIF5A*, and *TIMM23*. These genes play diverse roles in cellular functioning. *DNAJC15* has been found to be expressed in breast cancer cells[69] and may play a role in apoptosis[70]. *RPP25* enables ribonuclease P RNA binding activity and is involved in tRNA 5'-leader removal[71,72]. *CXCL12* is the ligand for the G-protein coupled receptor and chemokine and can affect tumor progression, angiogenesis, metastasis, and survival in cancer cells[73,74]. *EIF5A* is a translation elongation factor[75] and plays a role in regulating tumor cell growth[76]. *TIMM23* mediates the transport of transit peptide-containing proteins across the membrane[77]. Elvidge et al. showed that some of these genes may play a role in modulating changes in gene expression induced by hypoxia in MCF-7 breast cancer cell lines[78]. However, hypoxia and these other cellular processes do

not have a clear, known function that would explain gene-expression differences in breast tumors between people of African or European descent specifically.

The top genes identified in the ER-positive versus ER-negative comparison were *GATA3*, *ESR1,* and *CA12*. These genes exhibited approximately two-fold higher expression in ER-positive patients compared to ER-negative patients. *ESR1* encodes the estrogen receptor alpha (ERα) protein, which mediates the effects of estrogen in various tissues throughout the body[79]. Finding this gene near the top of our list when comparing ER positive versus ER negative breast cancers is evidence of the validity of our data and methods. *GATA3* is crucial for the differentiation of the mammary epithelium and is a key regulator in breast cancer[80]. It also plays a role in mediating the binding of estrogen receptor elements to *ESR1*[81]. Furthermore, co-expression of *GATA3*, *ESR1,* and *FOXA1* in breast tumors has been reported[82,83]. *CA12* is involved in cellular proliferation and pH regulation and has been associated with a good prognosis in breast cancer[84]. Other significant genes include *FOXA1* and *TBC1D9*. *FOXA1* is a transcriptional regulator that modulates the effects of estrogen signaling by influencing *ESR1* activity[85,86]. *TBC1D9* is involved in cellular trafficking processes that can influence cancer cell behavior[87]. These and other top-ranked genes merit further study to provide mechanistic insight into the ER protein's role in breast tumorigenesis, and suggest their potential roles both as biomarkers and therapeutic targets.

For the comparison between PR-positive and PR-negative tumors, a common thread among the top genes identified, is their involvement in various aspects of known breast cancer biology. Other studies[88] have shown that many of these genes (*PGR, ESR1, FOXA1, GATA3, GREB1*) are differentially expressed and that *PGR*[89] and *CA12*[84] may be good prognostic markers for breast cancer.

Among the top genes for the comparison between HER2-positive and HER2-negative tumors were *ERBB2* (which was expected because this gene encodes the HER2 protein) as

well as *PGAP3, STARD3, GRB7, CDK12,* and *PSMD3. PGAP3* is overexpressed in breast cancer patients, is associated with lymph node metastasis, and correlates with HER2 overexpression[90]. *STARD3* is necessary for the transfer of cholesterol and metabolism in tumor cells[91], and is often co-amplified with *HER2*[92]. *GRB7* is responsible for signal transduction from cell surface receptors via signaling pathways in the cell[93]. It is sometimes co-amplified and overexpressed with HER2 and is associated with a more aggressive phenotype[94]. *CDK12* is a transcription-associated cyclin-dependent kinase, which plays various roles in the regulation of translation[95], cellular proliferation[96], and cell cycle progression[97]; it acts as a tumor promoter in HER2-positive breast cancer[98]. *PSMD3* is an enzyme regulator by stabilizing *HER2* and preventing it from degradation[99]. These genes may serve as alternative markers for HER2 positive breast cancer[90,100].

For the comparison between TNBC and non-TNBC patients, the most significant genes sometimes overlapped with what we identified in the earlier analyses. *FOXA1*, *MLPH*, and *GATA3* were expressed at approximately two-fold higher levels in non-TNBCs compared to TNBCs, while *FOXC1* had approximately 2-fold lower expression in non-TNBC than TNBC tumors.

Our last comparison focused on individuals of African or European ancestry who had TNBC. The statistically significant genes were *RETN, ADAMTS13, ACOXL, FAM3A, GPR35*, and *PGLYRP1. RETN* encodes the hormone resistin that plays a role in inflammation[101] and insulin resistance[102], which are critical factors in obesity-related cancers[103]. Increased resistin levels have been associated with breast cancer[104,105]. *ADAMTS13* encodes an enzyme that processes von Willebrand factor, a protein involved in blood clotting. Dysregulation of *ADAMTS13* has been implicated in cancer metastasis and tumor angiogenesis due to its role in extracellular matrix remodeling[106,107]. *ACOXL* is involved in fatty acid β-oxidation and metabolic processes[108]. Although there is no research linking it to breast cancer, its expression has been linked to prostate cancer, where it is a potential biomarker due to

differential expression in benign versus malignant prostate tissues[109]. *FAM3A* is involved in regulating of both glucose and lipid metabolism[110]. Given its metabolic functions and the link between metabolism and cancer progression, it has been suggested to play a role in metabolic diseases[111] and possibly cancers[112]. *GPR35* is involved in inflammatory responses and is highly expressed in various cancers, including gastrointestinal and prostate. It is considered a potential therapeutic target due to its role in tumorigenesis and cancer progression[113,114]. *PGLYRP1* encodes a protein involved in the innate immune response and plays a role in regulating acquired immunity[115]. *PGLYRP1* has been implicated in inflammation and immune responses in the tumor microenvironment[116,117], which can influence cancer development and progression. These genes share involvement in inflammation, metabolic regulation, and the tumor microenvironment, which are critical processes in the development and progression of cancer even though they are not all specific to breast cancer. We did not find evidence to provide mechanistic insight about these expression differences between people of African or European descent. Yet they may merit further exploration as potential therapeutic targets.

To gain further insight into the biology behind the selected genes for each comparison, we performed pathway enrichment analysis using MSigDB[61,62] (Additional Files 8 - 12). Some of these results further confirm the validity of our methods. For example, in the ER-positive versus ER-negative comparison, curated pathways that reflect ER signaling were statistically significant. Similarly, for the HER2-positive versus HER2-negative comparison, pathways associated with HER2 amplification were significant. For the PR comparison, ER pathways were statistically significant, but we did not observe pathways specific to PR signaling; this suggests that ER and PR positivity have similar effects but may also reflect that MSigDB lacks pathways specific to PR. When a gene is statistically significant but not associated with any of the statistically significant pathways for a given comparison, it serves as a hypothesis for further investigation. Additionally, for significant pathways, genes in those pathways might

be prioritized for investigation, even when they were not among the most statistically significant genes.

Discussion

Researchers have used diverse methodologies to analyze and combine gene expression datasets[118]. Some have focused on identifying prognostic markers[43,119] or genes[120,121], predicting treatment outcomes[122–124], or deriving more specific breast cancer subtypes[49,125]. Others have attempted to differentiate between primary and metastatic tumors[126], as well as differential gene-expression analysis[43,127]. In this study, we carried out meta-analyses for tens of thousands of breast tumors. To the best of our knowledge, no meta-analyses of this magnitude has been performed previously; nor has a meta-analysis been performed to compare breast cancer patients of African and European ancestry. Schneider et al. focused on ER patients and identified a subset of genes that also show up in our analysis[128]. Al-Ejeh et al. focused on TNBCs and identified a set of 206 "aggressive genes"[30], some of which also were significant in our study. D'Arcy et al. compared Luminal A breast tumors between African American and Caucasian women and arrived at a list of 6 genes; but only one of those genes (*PSPH*) overlaps with our list. These differences may be due to their modeling approach and the fact that they had a smaller sample size (n = 167 samples) and used an imputation method to account for missing data.

Race and Gene Expression

One goal of this research was to understand the biology behind disparities in breast cancer between women of African and Caucasian descent. Our analysis revealed 850 genes that were significantly differentially expressed between these populations. Among the top 20 genes, we observed significant roles for inflammation (*RETN*)[101], cell cycle regulation (*CENPA* and *TTBK2*)[129,130], immune responses (*HLA-DPA1, CXCL12,* and *STAP2*)[74,131,132], and cellular metabolism (*PSPH, PYCR1,* and *HSD17B10*)[133–135]. These can influence tumor behavior and responses to treatments. For instance, *RETN* (resistin) is associated with

inflammation and insulin resistance, which are critical in the pathogenesis of obesity-related cancers, including breast cancer. Elevated resistin levels have been linked to poor prognosis in breast cancer patients, underscoring the potential of RETN as a biomarker for breast cancer in diverse populations.

Black women are more likely to be diagnosed with TNBC, which is more aggressive and has fewer targeted treatment options, while Caucasian women are more frequently diagnosed with hormone receptor-positive breast cancers, which have better prognoses and more effective hormone-targeted therapies[136,137]. The genes we identified when we merged TNBC and race are involved in inflammation and immune responses, which are critical pathways in cancer development, including breast cancer. ADAMTS13, an enzyme involved in blood clotting, plays a role in cancer metastasis and angiogenesis through extracellular matrix remodeling. Dysregulation of ADAMTS13 has been implicated in various cancers, suggesting its potential as a therapeutic target in breast cancer[138]. Similarly, genes such as *GPR35*, involved in inflammatory responses, and *PGLYRP1*, implicated in the innate immune response, highlight the importance of inflammation and immune regulation in cancer development[139–143]. Variations in these genes and their expression can vary by race, potentially influencing disease susceptibility and progression[144,145].

A key limitation of our analysis is that categories based on race and ethnicity are self-described and may not adequately represent a person's ancestry. By performing secondary analyses of publicly available data, we were limited to the descriptions provided by the original authors. For simplicity, we used broad groupings (African descent, European descent) to categorize the patients. However, by aggregating data from many studies, we anticipated that meaningful patterns would be identifiable, despite this noise. Although our analyses identified genes that have plausible connections to breast tumorigenesis and may lead to insights about population disparities, gene-expression differences between these

categories were relatively minor, providing support for the argument that race is a social construct more than a biological concept[146].

ER, PR, and HER2 Status

Compared to ER-negative breast tumors, genes like *ESR1, GATA3,* and *CA12* were more highly expressed in ER-positive tumors. *ESR1*, encoding the estrogen receptor alpha (ERα), is crucial for mediating estrogen's effects and is a well-established marker in breast cancer diagnostics and treatment[79,147]. *GATA3* is essential for mammary epithelial differentiation and is a key regulator in breast cancer, while *CA12* is involved in cellular proliferation and pH regulation and is associated with a good prognosis[148,149].

For PR status, genes such as *PGR, ESR1,* and *FOXA1* were differentially expressed, consistent with their roles in breast cancer biology. Progesterone receptor (PGR) and estrogen receptor (ESR1) are critical markers for hormone receptor-positive breast cancers and are important for determining therapeutic strategies[150,151].

Our HER2-positive vs. HER2-negative comparisons identified genes like *ERBB2*, *PGAP3*, and *GRB7*, which were prominently expressed in HER2-positive tumors. *ERBB2* overexpression is a hallmark of HER2-positive breast cancers and a target for therapies such as trastuzumab[152]. The co-amplification of genes like *GRB7* with HER2 suggests their roles in aggressive breast cancer phenotypes and potential as additional therapeutic targets[153,154].

TNBC vs. Non-TNBC

TNBC vs. non-TNBC analysis identified *FOXA1, MLPH,* and *GATA3,* which showed higher expression in non-TNBC, while *FOXC1* was expressed lower in non-TNBC. These findings highlight the distinct molecular profiles of TNBC, characterized by the lack of hormone receptor and HER2 expression[155].

Common Genes Across Comparisons

GATA3, CA12, TBC1D9, XBP1, and *FOXA1* were consistently significant across ER, PR, and TNBC comparisons but not in the HER2 comparison. This supports the notion that the mechanisms behind ER-positive and PR-positive tumors are overlapping and that HER2-positive tumors are mechanistically distinct from endocrine breast cancers, necessitating different therapeutic strategies[156–158].

Computational Validation

It is common for differential-expression studies, such as this meta-analysis, to identify thousands of statistically significant genes. Such findings may be reassuring because they imply that the compared phenotypes are truly distinct at the molecular level. However, a larger goal of a differential-expression study is to reduce the search space for genes that are most important in driving the phenotype. Lab-based, experimental validation can provide assurances that genes are truly differentially expressed between conditions; however, such procedures do not scale beyond a few genes. Furthermore, the complexity of cancers likely cannot be captured adequately using data from a few genes. Therefore, it is desirable to find a balance between characterizing the complexity of cancers and identifying a gene set that is small enough to be analyzed and interpreted by humans and that can be explored feasibly in subsequent, mechanistic studies. Our novel computational method addresses these goals, while also showing that most of our identified gene sets are more relevant than randomly selected genes. Moreover, it provides insight into the potential to use these genes to predict phenotypes. A limitation of differential-expression studies is that commonly used statistical methods consider evidence from one gene at a time, whereas our classification approach constructs models from many genes simultaneously. We believe this method has potential to be applied in many other contexts. Further validation will be necessary.

Conclusion

Findings from our meta-analyses elucidate gene expression differences between racial groups and clinically relevant breast cancer subtypes, highlighting the roles of diverse cellular processes. They provide additional evidence about mechanisms that may drive breast cancer biology. Such studies are necessary to pave the way for developing targeted diagnostics and therapies in a personalized-medicine context. Further research is warranted to explore the mechanistic pathways of these genes and their potential as biomarkers and therapeutic targets in diverse patient populations.

List of Abbreviations

ER : estrogen receptor

ESR1 : estrogen receptor

HER2 : human epidermal growth factor receptor 2

IGF1 : insulin-like growth factor 1

IHC : immunohistochemistry

MSigDB : Molecular Signatures Database

OSF : Open Science Framework

PGR : Progesterone receptor

PR : progesterone receptor

SCAN-B : The Sweden Cancerome Analysis Network - Breast

TCGA : The Cancer Genome Atlas

TNBC : Triple-negative breast cancers

Additional Data Files

Additional File 1

Format: xlsx

Title: Identifiers for datasets used in the meta-analyses.

Description: This spreadsheet indicates the accession number or other unique identifiers for each dataset analyzed in this study.

Additional File 2

Format: xlsx

Title: Genes that were significantly differentially expressed between individuals of African or European ancestry.

Additional File 3

Format: xlsx

Title: Genes that were significantly differentially expressed between ER-positive and ER-negative breast tumors.

Additional File 4

Format: xlsx

Title: Genes that were significantly differentially expressed between PR-positive and PR-negative breast tumors.

Additional File 5

Format: xlsx

Title: Genes that were significantly differentially expressed between HER2-positive and HER2-negative breast tumors.

Additional File 6

Format: xlsx

Title: Genes that were significantly differentially expressed between TNBC versus non-TNBC breast tumors.

Additional File 7

Format: xlsx

Title: Genes that were significantly differentially expressed in TNBC tumors between individuals of African or European ancestry.

Additional File 8

Format: xlsx

Title: Gene-set overlap results for the top-ranked genes identified in the race meta-analysis.

Additional File 9

Format: xlsx

Title: Gene-set overlap results for the top-ranked genes identified in the ER status meta-analysis.

Additional File 10

Format: xlsx

Title: Gene-set overlap results for the top-ranked genes identified in the PR status meta-analysis.

Additional File 11

Format: xlsx

Title: Gene-set overlap results for the top-ranked genes identified in the HER2 status meta-analysis.

Additional File 12.

Format: xlsx

Title: Gene-set overlap results for the top-ranked genes identified in the TNBC status meta-analysis.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All source code, gene expression data, and metadata are available in a publicly accessible Open Science Framework (OSF)[159] repository (https://osf.io/mr8sb/). The code is packaged and executable using Docker[160] software containers for easier reproducibility.

Competing interests

The authors declare that they have no competing interests

Funding

The authors received no external funding for this research.

Authors' contributions

ION prepared and analyzed data. Both authors designed the study and interpreted the results. SRP supervised the work. Both authors wrote the manuscript and approved the final manuscript.

Acknowledgements

Not applicable

References

1. Vogelstein B, Kinzler KW. Cancer genes and the pathways they control. Nat Med. 2004 Aug;10(8):789–99.

2. Ali MA, Sjöblom T. Molecular pathways in tumor progression: from discovery to functional understanding. Mol Biosyst. 2009 Sep;5(9):902–8.

3. Futreal PA, Coin L, Marshall M, Down T, Hubbard T, Wooster R, et al. A census of human cancer genes. Nat Rev Cancer. 2004 Mar;4(3):177–83.

4. Cosmic. Cancer Gene Census [Internet]. 2018 [cited 2018 Dec 7]. Available from: https://cancer.sanger.ac.uk/census

5. Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA Jr, Kinzler KW. Cancer genome landscapes. Science. 2013 Mar 29;339(6127):1546–58.

6. Pennisi E. Human Genetic Variation. Science. 2007 Dec 21;318(5858):1842–3.

7. Breast cancer facts and statistics 2023 [Internet]. [cited 2023 Sep 5]. Available from: https://www.breastcancer.org/facts-statistics

8. CDCBreastCancer. Breast cancer statistics [Internet]. Centers for Disease Control and Prevention. 2023 [cited 2023 Sep 5]. Available from: https://www.cdc.gov/cancer/breast/statistics/index.htm

9. Breast Cancer Statistics [Internet]. [cited 2023 Sep 6]. Available from: https://www.cancer.org/cancer/types/breast-cancer/about/how-common-is-breast-cancer.html

10. Arranz EE, Vara JÁF, Gámez-Pozo A, Zamora P. Gene signatures in breast cancer: current and future uses. Transl Oncol. 2012 Dec;5(6):398–403.

11. Berry DA, Cronin KA, Plevritis SK, Fryback DG, Clarke L, Zelen M, et al. Effect of

screening and adjuvant therapy on mortality from breast cancer. N Engl J Med. 2005 Oct 27;353(17):1784–92.

12. Lal S, McCart Reed AE, de Luca XM, Simpson PT. Molecular signatures in breast cancer [Internet]. Vol. 131, Methods. 2017. p. 135–46. Available from: http://dx.doi.org/10.1016/j.ymeth.2017.06.032

13. Ringnér M, Fredlund E, Häkkinen J, Borg Å, Staaf J. GOBO: gene expression-based outcome for breast cancer online. PLoS One. 2011 Mar 21;6(3):e17911.

14. Guo L, Kong D, Liu J, Zhan L, Luo L, Zheng W, et al. Breast cancer heterogeneity and its implication in personalized precision therapy. Exp Hematol Oncol. 2023 Jan 9;12(1):3.

15. Melchor L, Benítez J. The complex genetic landscape of familial breast cancer. Hum Genet. 2013 Aug;132(8):845–63.

16. Keenan T, Moy B, Mroz EA, Ross K, Niemierko A, Rocco JW, et al. Comparison of the Genomic Landscape Between Primary Breast Cancer in African American Versus White Women and the Association of Racial Differences With Tumor Recurrence [Internet]. Vol. 33, Journal of Clinical Oncology. 2015. p. 3621–7. Available from: http://dx.doi.org/10.1200/jco.2015.62.2126

17. American Cancer Society. Cancer Facts & Figures for African Americans 2016-2018. American Cancer Society [Internet]. 2016; Available from: https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/cancer-facts-and-figures-for-african-americans/cancer-facts-and-figures-for-african-americans-2016-2018.pdf

18. Lowder D, Rizwan K, McColl C, Paparella A, Ittmann M, Mitsiades N, et al. Racial disparities in prostate cancer: A complex interplay between socioeconomic inequities and genomics. Cancer Lett. 2022 Apr 10;531:71–82.

19. Ryan BM. Lung cancer health disparities. Carcinogenesis. 2018 May 28;39(6):741–51.

20. Tannenbaum SL, Koru-Sengul T, Miao F, Byrne MM. Disparities in survival after female breast cancer diagnosis: a population-based study. Cancer Causes Control. 2013 Sep;24(9):1705–15.

21. Nurgalieva ZZ, Franzini L, Morgan RO, Vernon SW, Liu CC, Du XL. Impact of timing of adjuvant chemotherapy initiation and completion after surgery on racial disparities in survival among women with breast cancer. Med Oncol. 2013 Mar;30(1):419.

22. Polite BN, Dignam JJ, Olopade OI. Colorectal cancer model of health disparities: understanding mortality differences in minority populations. J Clin Oncol. 2006 May 10;24(14):2179–87.

23. Smith EC, Ziogas A, Anton-Culver H. Delay in surgical treatment and survival after breast cancer diagnosis in young women by race/ethnicity. JAMA Surg. 2013 Jun;148(6):516–23.

24. Parise CA, Caggiano V. Disparities in race/ethnicity and socioeconomic status: risk of mortality of breast cancer patients in the California Cancer Registry, 2000--2010. BMC Cancer. 2013;13(1):449.

25. Goodwin S, McPherson JD, McCombie WR. Coming of age: ten years of next-generation sequencing technologies. Nat Rev Genet. 2016 May 17;17(6):333–51.

26. Hansen AR, Bedard PL. Clinical application of high-throughput genomic technologies for treatment selection in breast cancer. Breast Cancer Res. 2013;15(5):R97.

27. Mardis ER. The impact of next-generation sequencing technology on genetics. Trends Genet. 2008 Mar;24(3):133–41.

28. Parker JS, Mullins M, Cheang MCU, Leung S, Voduc D, Vickery T, et al. Supervised risk predictor of breast cancer based on intrinsic subtypes. J Clin Oncol. 2009 Mar

10;27(8):1160–7.

29. Sørlie T, Wang Y, Xiao C, Johnsen H, Naume B, Samaha RR, et al. Distinct molecular mechanisms underlying clinically relevant subtypes of breast cancer: gene expression analyses across three different platforms. BMC Genomics. 2006 May 26;7:127.

30. Al-Ejeh F, Simpson PT, Sanus JM, Klein K, Kalimutho M, Shi W, et al. Meta-analysis of the global gene expression profile of triple-negative breast cancer identifies genes for the prognostication and treatment of aggressive breast cancer. Oncogenesis. 2014 Apr 21;3:e100.

31. Sturtz LA, Melley J, Mamula K, Shriver CD, Ellsworth RE. Outcome disparities in African American women with triple negative breast cancer: a comparison of epidemiological and molecular factors between African American and Caucasian women with triple negative breast cancer. BMC Cancer. 2014;14(1):62.

32. D'Arcy M, Fleming J, Robinson WR, Kirk EL, Perou CM, Troester MA. Race-associated biological differences among Luminal A breast tumors [Internet]. Vol. 152, Breast Cancer Research and Treatment. 2015. p. 437–48. Available from: http://dx.doi.org/10.1007/s10549-015-3474-4

33. Field LA, Love B, Deyarmin B, Hooke JA, Shriver CD, Ellsworth RE. Identification of differentially expressed genes in breast tumors from African American compared with Caucasian women. Cancer. 2012 Mar 1;118(5):1334–44.

34. Stewart PA, Luks J, Roycik MD, Sang QXA, Zhang J. Differentially expressed transcripts and dysregulated signaling pathways and networks in African American breast cancer. PLoS One. 2013 Dec 4;8(12):e82460.

35. Byun JS, Singhal SK, Park S, Yi DI, Yan T, Caban A, et al. Racial Differences in the Association Between Luminal Master Regulator Gene Expression Levels and Breast Cancer Survival. Clin Cancer Res. 2020 Apr 15;26(8):1905–14.

36. Chavez-MacGregor M, Liu S, De Melo-Gagliato D, Chen H, Do KA, Pusztai L, et al. Differences in gene and protein expression and the effects of race/ethnicity on breast cancer subtypes. Cancer Epidemiol Biomarkers Prev. 2014 Feb 1;23(2):316–23.

37. Howlader N, Altekruse SF, Li CI, Chen VW, Clarke CA, Ries LAG, et al. US incidence of breast cancer subtypes defined by joint hormone receptor and HER2 status. J Natl Cancer Inst [Internet]. 2014 Apr 28;106(5). Available from: http://dx.doi.org/10.1093/jnci/dju055

38. Wright JL, Reis IM, Zhao W, Panoff JE, Takita C, Sujoy V, et al. Racial disparity in estrogen receptor positive breast cancer patients receiving trimodality therapy. Breast. 2012 Jun;21(3):276–83.

39. Dent R, Trudeau M, Pritchard KI, Hanna WM, Kahn HK, Sawka CA, et al. Triple-negative breast cancer: Clinical features and patterns of recurrence. Clin Cancer Res. 2007 Aug 1;13(15):4429–34.

40. Perez CA, Zumsteg ZS, Gupta G, Morrow M, Arnold B, Patil SM, et al. Black race as a prognostic factor in triple-negative breast cancer patients treated with breast-conserving therapy: a large, single-institution retrospective analysis. Breast Cancer Res Treat. 2013 Jun 1;139(2):497–506.

41. Gleason MX, Mdzinarishvili T, Sherman S. Breast cancer incidence in black and white women stratified by estrogen and progesterone receptor statuses. PLoS One. 2012 Nov 14;7(11):e49359.

42. Martin DN, Boersma BJ, Yi M, Reimers M, Howe TM, Yfantis HG, et al. Differences in the tumor microenvironment between African-American and European-American breast cancer patients. PLoS One. 2009 Feb 19;4(2):e4531.

43. Lindner R, Sullivan C, Offor O, Lezon-Geyda K, Halligan K, Fischbach N, et al. Molecular phenotypes in triple negative breast cancer from African American patients

suggest targets for therapy. PLoS One. 2013 Nov 18;8(11):e71915.

44. Nwosu IO, Tabler DD, Chipman G, Piccolo SR. Annotated Compendium of 102 Breast Cancer Gene-Expression Datasets [Internet]. bioRxiv. 2023 [cited 2023 Oct 2]. p. 2023.09.22.559045. Available from: https://www.biorxiv.org/content/10.1101/2023.09.22.559045

45. Mikolajewicz N, Komarova SV. Meta-Analytic Methodology for Basic Research: A Practical Guide. Front Physiol. 2019 Mar 27;10:203.

46. Edgar R, Domrachev M, Lash AE. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. Nucleic Acids Res. 2002 Jan 1;30(1):207–10.

47. Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, et al. NCBI GEO: archive for functional genomics data sets--update. Nucleic Acids Res. 2013 Jan;41(Database issue):D991–5.

48. Parkinson H, Kapushesky M, Shojatalab M, Abeygunawardena N, Coulson R, Farne A, et al. ArrayExpress--a public database of microarray experiments and gene expression profiles. Nucleic Acids Res. 2007 Jan;35(Database issue):D747–50.

49. Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, et al. The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. Nature. 2012 Apr 18;486(7403):346–52.

50. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. Nature. 2012 Oct 4;490(7418):61–70.

51. Saal LH, Vallon-Christersson J, Häkkinen J, Hegardt C, Grabau D, Winter C, et al. The Sweden Cancerome Analysis Network - Breast (SCAN-B) Initiative: a large-scale multicenter infrastructure towards implementation of breast cancer genomic analyses in the clinical routine. Genome Med. 2015 Feb 2;7(1):20.

52. Staaf J, Häkkinen J, Hegardt C, Saal LH, Kimbung S, Hedenfalk I, et al. RNA sequencing-based single sample predictors of molecular subtype and risk of recurrence for clinical assessment of early-stage breast cancer. NPJ Breast Cancer. 2022 Aug 16;8(1):94.

53. Villatoro-García JA, Martorell-Marugán J, Toro-Domínguez D, Román-Montoya Y, Femia P, Carmona-Sáez P. DExMA: An R Package for Performing Gene Expression Meta-Analysis with Missing Genes. Sci China Ser A Math. 2022 Sep 17;10(18):3376.

54. Core Team RR. No title [Internet]. [cited 2024 Feb 27]. Available from: https://apps.dtic.mil/sti/citations/AD1039033

55. Venet D, Dumont JE, Detours V. Most random gene expression signatures are significantly associated with breast cancer outcome. PLoS Comput Biol [Internet]. 2011;7(10). Available from: http://dx.doi.org/10.1371/journal.pcbi.1002240

56. Refaeilzadeh P, Tang L, Liu H. Cross-Validation. In: Liu L, Özsu MT, editors. Encyclopedia of Database Systems. Boston, MA: Springer US; 2009. p. 532–8.

57. Lei J. Cross-Validation With Confidence. J Am Stat Assoc. 2020 Dec 11;115(532):1978–97.

58. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn: Machine learning in Python. the Journal of machine Learning research. 2011;12:2825–30.

59. Breiman L. Random Forests. Mach Learn. 2001 Oct 1;45(1):5–32.

60. Piccolo SR, Mecham A, Golightly NP, Johnson JL, Miller DB. The ability to classify patients based on gene-expression data varies by algorithm and performance metric. PLoS Comput Biol. 2022 Mar;18(3):e1009926.

61. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene

set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A. 2005 Oct 25;102(43):15545–50.

62. Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov JP. Molecular signatures database (MSigDB) 3.0. Bioinformatics. 2011 Jun 15;27(12):1739–40.

63. Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell Syst. 2015 Dec 23;1(6):417–25.

64. Bose R, Kavuri SM, Searleman AC, Shen W, Shen D, Koboldt DC, et al. Activating HER2 mutations in HER2 gene amplification negative breast cancer. Cancer Discov. 2013 Feb;3(2):224–37.

65. Baselga J. Why the epidermal growth factor receptor? The rationale for cancer therapy. Oncologist. 2002;7 Suppl 4:2–8.

66. Godoy-Ortiz A, Sanchez-Muñoz A, Chica Parrado MR, Álvarez M, Ribelles N, Rueda Dominguez A, et al. Deciphering HER2 Breast Cancer Disease: Biological and Clinical Implications. Front Oncol. 2019 Oct 29;9:1124.

67. Lanari C, Molinolo AA. Progesterone receptors - animal models and cell signaling in breast cancer Diverse activation pathways for the progesterone receptor - possible implications for breast biology and cancer. Breast Cancer Res. 2002 Dec 1;4(6):1–4.

68. Hua H, Zhang H, Kong Q, Jiang Y. Mechanisms for estrogen receptor expression in human cancer. Exp Hematol Oncol. 2018 Sep 19;7:24.

69. Navasa N, Martin-Ruiz I, Atondo E, Sutherland JD, Angel Pascual-Itoiz M, Carreras-González A, et al. Ikaros mediates the DNA methylation-independent silencing of MCJ/DNAJC15 gene expression in macrophages. Sci Rep. 2015 Sep 30;5:14692.

70. Sinha D, D'Silva P. Chaperoning mitochondrial permeability transition: regulation of transition pore complex by a J-protein, DnaJC15. Cell Death Dis. 2014 Mar 6;5(3):e1101.

71. Jarrous N, Gopalan V. Archaeal/eukaryal RNase P: subunits, functions and RNA diversification. Nucleic Acids Res. 2010 Dec;38(22):7885–94.

72. Abu-Zhayia ER, Khoury-Haddad H, Guttmann-Raviv N, Serruya R, Jarrous N, Ayoub N. A role of human RNase P subunits, Rpp29 and Rpp21, in homology directed-repair of double-strand breaks. Sci Rep. 2017 Apr 21;7(1):1002.

73. Guyon A. CXCL12 chemokine and its receptors as major players in the interactions between immune and nervous systems. Front Cell Neurosci. 2014 Mar 6;8:65.

74. Meng W, Xue S, Chen Y. The role of CXCL12 in tumor microenvironment. Gene. 2018 Jan 30;641:105–10.

75. Ning L, Wang L, Zhang H, Jiao X, Chen D. Eukaryotic translation initiation factor 5A in the pathogenesis of cancers. Oncol Lett. 2020 Oct;20(4):81.

76. Nakanishi S, Cleveland JL. Targeting the polyamine-hypusine circuit for the prevention and treatment of cancer. Amino Acids. 2016 Oct;48(10):2353–62.

77. Bauer MF, Sirrenberg C, Neupert W, Brunner M. Role of Tim23 as voltage sensor and presequence receptor in protein import into mitochondria. Cell. 1996 Oct 4;87(1):33–41.

78. Elvidge GP, Glenny L, Appelhoff RJ, Ratcliffe PJ, Ragoussis J, Gleadle JM. Concordant regulation of gene expression by hypoxia and 2-oxoglutarate-dependent dioxygenase inhibition: the role of HIF-1alpha, HIF-2alpha, and other pathways. J Biol Chem. 2006 Jun 2;281(22):15215–26.

79. Miziak P, Baran M, Błaszczak E, Przybyszewska-Podstawka A, Kałafut J, Smok-Kalwat J, et al. Estrogen Receptor Signaling in Breast Cancer. Cancers [Internet]. 2023 Sep

23;15(19). Available from: http://dx.doi.org/10.3390/cancers15194689

80. Takaku M, Grimm SA, Roberts JD, Chrysovergis K, Bennett BD, Myers P, et al. GATA3 zinc finger 2 mutations reprogram the breast cancer transcriptional network. Nat Commun. 2018 Mar 13;9(1):1059.

81. Theodorou V, Stark R, Menon S, Carroll JS. GATA3 acts upstream of FOXA1 in mediating ESR1 binding by shaping enhancer accessibility. Genome Res. 2013 Jan;23(1):12–22.

82. Badve S, Turbin D, Thorat MA, Morimiya A, Nielsen TO, Perou CM, et al. FOXA1 expression in breast cancer--correlation with luminal subtype A and survival. Clin Cancer Res. 2007 Aug 1;13(15 Pt 1):4415–21.

83. Wilson BJ, Giguère V. Meta-analysis of human cancer microarrays reveals GATA3 is integral to the estrogen receptor alpha pathway. Mol Cancer. 2008 Jun 4;7:49.

84. Li Y, Lei B, Zou J, Wang W, Chen A, Zhang J, et al. High expression of carbonic anhydrase 12 (CA12) is associated with good prognosis in breast cancer. Neoplasma. 2019 May 23;66(3):420–6.

85. Hurtado A, Holmes KA, Ross-Innes CS, Schmidt D, Carroll JS. FOXA1 is a key determinant of estrogen receptor function and endocrine response. Nat Genet. 2011 Jan;43(1):27–33.

86. Seachrist DD, Anstine LJ, Keri RA. FOXA1: A Pioneer of Nuclear Receptor Action in Breast Cancer. Cancers [Internet]. 2021 Oct 17;13(20). Available from: http://dx.doi.org/10.3390/cancers13205205

87. Kothari C, Clemenceau A, Ouellette G, Ennour-Idrissi K, Michaud A, C-Gaudreault R, et al. TBC1D9: An Important Modulator of Tumorigenesis in Breast Cancer. Cancers [Internet]. 2021 Jul 16;13(14). Available from:

http://dx.doi.org/10.3390/cancers13143557

88. Wu JR, Zhao Y, Zhou XP, Qin X. Estrogen receptor 1 and progesterone receptor are distinct biomarkers and prognostic factors in estrogen receptor-positive breast cancer: Evidence from a bioinformatic analysis. Biomed Pharmacother. 2020 Jan;121:109647.

89. Kurozumi S, Matsumoto H, Hayashi Y, Tozuka K, Inoue K, Horiguchi J, et al. Power of PgR expression as a prognostic factor for ER-positive/HER2-negative breast cancer patients at intermediate risk classified by the Ki67 labeling index. BMC Cancer. 2017 May 22;17(1):354.

90. Hao N, Li M, Wang J, Song Y, Zhao Y, Zhang L, et al. High PGAP3 expression is associated with lymph node metastasis and low CD8+T cell in patients with HER2+ breast cancer. Pathol Res Pract. 2023 Nov;251:154890.

91. Fararjeh AFS, Al Khader A, Kaddumi E, Obeidat M, Al-Fawares O 'la. Differential Expression and Prognostic Significance of STARD3 Gene in Breast Carcinoma. Int J Mol Cell Med. 2021 May 22;10(1):34–41.

92. Kauraniemi P, Kallioniemi A. Activation of multiple cancer-associated genes at the ERBB2 amplicon in breast cancer. Endocr Relat Cancer. 2006 Mar;13(1):39–49.

93. Vermehren-Schmaedick A, Mhawech-Fauceglia P, Park BS, Pejovic T, Luoh SW. The prognostic significance of GRB7 protein expression and localization in human breast and ovarian cancers. Oncotarget. 2020 Jun 16;11(24):2273–89.

94. Bivin WW, Yergiyev O, Bunker ML, Silverman JF, Krishnamurti U. GRB7 Expression and Correlation With HER2 Amplification in Invasive Breast Carcinoma. Appl Immunohistochem Mol Morphol. 2017 Sep;25(8):553–8.

95. Choi SH, Martinez TF, Kim S, Donaldson C, Shokhirev MN, Saghatelian A, et al. CDK12 phosphorylates 4E-BP1 to enable mTORC1-dependent translation and mitotic genome

stability. Genes Dev. 2019 Apr 1;33(7-8):418–35.

96. Lei T, Zhang P, Zhang X, Xiao X, Zhang J, Qiu T, et al. Cyclin K regulates prereplicative complex assembly to promote mammalian cell proliferation. Nat Commun. 2018 May 14;9(1):1876.

97. Malumbres M, Barbacid M. Cell cycle, CDKs and cancer: a changing paradigm. Nat Rev Cancer. 2009 Mar;9(3):153–66.

98. Liang S, Hu L, Wu Z, Chen Z, Liu S, Xu X, et al. CDK12: A Potent Target and Biomarker for Human Cancer Therapy. Cells [Internet]. 2020 Jun 18;9(6). Available from: http://dx.doi.org/10.3390/cells9061483

99. Fararjeh AS, Chen LC, Ho YS, Cheng TC, Liu YR, Chang HL, et al. Proteasome 26S Subunit, non-ATPase 3 (PSMD3) Regulates Breast Cancer by Stabilizing HER2 from Degradation. Cancers [Internet]. 2019 Apr 12;11(4). Available from: http://dx.doi.org/10.3390/cancers11040527

100. Lodi M, Voilquin L, Alpy F, Molière S, Reix N, Mathelin C, et al. STARD3: A New Biomarker in HER2-Positive Breast Cancer. Cancers [Internet]. 2023 Jan 5;15(2). Available from: http://dx.doi.org/10.3390/cancers15020362

101. Park HK, Kwak MK, Kim HJ, Ahima RS. Linking resistin, inflammation, and cardiometabolic diseases. Korean J Intern Med. 2017 Mar;32(2):239–47.

102. Steppan CM, Bailey ST, Bhat S, Brown EJ, Banerjee RR, Wright CM, et al. The hormone resistin links obesity to diabetes. Nature. 2001 Jan 18;409(6818):307–12.

103. Dalamaga M. Resistin as a biomarker linking obesity and inflammation to cancer: potential clinical perspectives. Biomark Med. 2014;8(1):107–18.

104. Lee JO, Kim N, Lee HJ, Lee YW, Kim SJ, Park SH, et al. Resistin, a fat-derived secretory factor, promotes metastasis of MDA-MB-231 human breast cancer cells

through ERM activation. Sci Rep. 2016 Jan 5;6:18923.

105.   Deb A, Deshmukh B, Ramteke P, Bhati FK, Bhat MK. Resistin: A journey from metabolism to cancer. Transl Oncol. 2021 Oct;14(10):101178.

106.   Kumar S, Rao N, Ge R. Emerging Roles of ADAMTSs in Angiogenesis and Cancer. Cancers . 2012 Nov 29;4(4):1252–99.

107.   Redondo-García S, Peris-Torres C, Caracuel-Peramos R, Rodríguez-Manzaneque JC. ADAMTS proteases and the tumor immune microenvironment: Lessons from substrates and pathologies. Matrix Biol Plus. 2021 Feb;9:100054.

108.   Van Veldhoven PP. Biochemistry and genetics of inherited disorders of peroxisomal fatty acid metabolism. J Lipid Res. 2010 Oct;51(10):2863–95.

109.   O'Hurley G, Busch C, Fagerberg L, Hallström BM, Stadler C, Tolf A, et al. Analysis of the Human Prostate-Specific Proteome Defined by Transcriptomics and Antibody-Based Profiling Identifies TMEM79 and ACOXL as Two Putative, Diagnostic Markers in Prostate Cancer. PLoS One. 2015 Aug 3;10(8):e0133449.

110.   Wang C, Chi Y, Li J, Miao Y, Li S, Su W, et al. FAM3A activates PI3K p110α/Akt signaling to ameliorate hepatic gluconeogenesis and lipogenesis. Hepatology. 2014 May;59(5):1779–90.

111.   Yan H, Meng Y, Li X, Xiang R, Hou S, Wang J, et al. FAM3A maintains metabolic homeostasis by interacting with F1-ATP synthase to regulate the activity and assembly of ATP synthase. Metabolism. 2023 Feb;139:155372.

112.   Dong QT, Ma DD, Gong Q, Lin ZY, Li ZH, Ye JX, et al. FAM3 family genes are associated with prognostic value of human cancer: a pan-cancer analysis. Sci Rep. 2023 Sep 13;13(1):15144.

113.   Takkar S, Sharma G, Kaushal JB, Abdullah KM, Batra SK, Siddiqui JA. From orphan

to oncogene: The role of GPR35 in cancer and immune modulation. Cytokine Growth Factor Rev [Internet]. 2024 Mar 19; Available from: http://dx.doi.org/10.1016/j.cytogfr.2024.03.004

114.   Milligan G. GPR35: from enigma to therapeutic target. Trends Pharmacol Sci. 2023 May;44(5):263–73.

115.   Yashin DV, Sashchenko LP, Georgiev GP. Mechanisms of Action of the PGLYRP1/Tag7 Protein in Innate and Acquired Immunity. Acta Naturae. 2021 Jan-Mar;13(1):91–101.

116.   Bhusal A, Kim JH, Kim SC, Hwang EM, Ryu H, Ali MS, et al. The microglial innate immune protein PGLYRP1 mediates neuroinflammation and consequent behavioral changes. Cell Rep. 2024 Mar 26;43(3):113813.

117.   Schnell A, Huang L, Regan BML, Singh V, Vonficht D, Bollhagen A, et al. Targeting PGLYRP1 promotes antitumor immunity while inhibiting autoimmune neuroinflammation. Nat Immunol. 2023 Nov;24(11):1908–20.

118.   Nwosu IO, Piccolo SR. A systematic review of datasets that can help elucidate relationships among gene expression, race, and immunohistochemistry-defined subtypes in breast cancer. Cancer Biol Ther. 2021 Sep 2;22(7-9):417–29.

119.   Mehra R, Varambally S, Ding L, Shen R, Sabel MS, Ghosh D, et al. Identification of GATA3 as a breast cancer prognostic marker by global gene expression meta-analysis. Cancer Res. 2005 Dec 15;65(24):11259–64.

120.   Goldhirsch A, Winer EP, Coates AS, Gelber RD, Piccart-Gebhart M, Thürlimann B, et al. Personalizing the treatment of women with early breast cancer: highlights of the St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2013. Ann Oncol. 2013 Sep;24(9):2206–23.

121.   Dowsett M, Smith IE, Ebbs SR, Dixon JM, Skene A, A'Hern R, et al. Prognostic value of Ki67 expression after short-term presurgical endocrine therapy for primary breast cancer. J Natl Cancer Inst. 2007 Jan 17;99(2):167–70.

122.   Stover DG, Coloff JL, Barry WT, Brugge JS, Winer EP, Selfors LM. The Role of Proliferation in Determining Response to Neoadjuvant Chemotherapy in Breast Cancer: A Gene Expression-Based Meta-Analysis. Clin Cancer Res. 2016 Dec 15;22(24):6039–50.

123.   Julka PK, Chacko RT, Nag S, Parshad R, Nair A, Oh DS, et al. A phase II study of sequential neoadjuvant gemcitabine plus doxorubicin followed by gemcitabine plus cisplatin in patients with operable breast cancer: prediction of response using molecular profiling. Br J Cancer. 2008 Apr 22;98(8):1327–35.

124.   Chang JC, Wooten EC, Tsimelzon A, Hilsenbeck SG, Gutierrez MC, Elledge R, et al. Gene expression profiling for the prediction of therapeutic response to docetaxel in patients with breast cancer. Lancet. 2003 Aug 2;362(9381):362–9.

125.   Burstein MD, Tsimelzon A, Poage GM, Covington KR, Contreras A, Fuqua SAW, et al. Comprehensive genomic analysis identifies novel subtypes and targets of triple-negative breast cancer. Clin Cancer Res. 2015 Apr 1;21(7):1688–98.

126.   Ellsworth RE, Seebach J, Field LA, Heckman C, Kane J, Hooke JA, et al. A gene expression signature that defines breast cancer metastases. Clin Exp Metastasis. 2009;26(3):205–13.

127.   Loi S, Sotiriou C, Haibe-Kains B, Lallemand F, Conus NM, Piccart MJ, et al. Gene expression profiling identifies activated growth factor signaling in poor prognosis (Luminal-B) estrogen receptor positive breast cancer. BMC Med Genomics. 2009 Jun 24;2:37.

128.   Schneider J, Ruschhaupt M, Buness A, Asslaber M, Regitnig P, Zatloukal K, et al.

Identification and meta-analysis of a small gene expression signature for the diagnosis of estrogen receptor status in invasive ductal breast cancer. Int J Cancer. 2006 Dec 15;119(12):2974–9.

129.   McKinley KL, Cheeseman IM. The molecular basis for centromere identity and function. Nat Rev Mol Cell Biol. 2016 Jan;17(1):16–29.

130.   Bauer NG, Richter-Landsberg C, Ffrench-Constant C. Role of the oligodendroglial cytoskeleton in differentiation and myelination. Glia. 2009 Dec;57(16):1691–705.

131.   Neefjes J, Jongsma MLM, Paul P, Bakke O. Towards a systems understanding of MHC class I and MHC class II antigen presentation. Nat Rev Immunol. 2011 Nov 11;11(12):823–36.

132.   Matsuda T, Sasaki Y, Kagohashi K, Saitoh K, Sekine Y, Kashiwakura JI, et al. Signal-transducing adaptor protein-2 modulates T-cell functions. Explor Immunol. 2022 Dec 27;771–82.

133.   He XY, Isaacs C, Yang SY. Roles of Mitochondrial 17β-Hydroxysteroid Dehydrogenase Type 10 in Alzheimer's Disease. J Alzheimers Dis. 2018;62(2):665–73.

134.   Amelio I, Cutruzzolá F, Antonov A, Agostini M, Melino G. Serine and glycine metabolism in cancer. Trends Biochem Sci. 2014 Apr;39(4):191–8.

135.   Phang JM, Liu W, Hancock C, Christian KJ. The proline regulatory axis and cancer. Front Oncol. 2012 Jun 21;2:60.

136.   Siddharth S, Sharma D. Racial disparity and triple-negative breast cancer in African-American women: A multifaceted affair between obesity, biology, and socioeconomic determinants. Cancers [Internet]. 2018 Dec 1;10(12). Available from: http://dx.doi.org/10.3390/cancers10120514

137.   Dietze EC, Sistrunk C, Miranda-Carboni G, O'Regan R, Seewaldt VL. Triple-negative

breast cancer in African-American women: disparities versus biology. Nat Rev Cancer. 2015 Apr;15(4):248–54.

138.   Zheng L, Cao L, Zheng XL. ADAMTS13 protease or lack of von Willebrand factor protects irradiation and melanoma-induced thrombotic microangiopathy in zebrafish. J Thromb Haemost. 2022 Oct;20(10):2270–83.

139.   López-Gil JC, García-Silva S, Ruiz-Cañas L, Navarro D, Palencia-Campos A, Giráldez-Trujillo A, et al. The Peptidoglycan Recognition Protein 1 confers immune evasive properties on pancreatic cancer stem cells. Gut [Internet]. 2024 May 16; Available from: http://dx.doi.org/10.1136/gutjnl-2023-330995

140.   Cui X, Chang Z, Dang T, Meng J, Wang P, Wu J, et al. TNF upregulates peptidoglycan recognition protein 1 in esophageal cancer cells to clear the path to its signaling: Making the "enemy" a friend. Arch Biochem Biophys. 2022 Jun 15;722:109192.

141.   Quon T, Lin LC, Ganguly A, Tobin AB, Milligan G. Therapeutic Opportunities and Challenges in Targeting the Orphan G Protein-Coupled Receptor GPR35. ACS Pharmacol Transl Sci. 2020 Oct 9;3(5):801–12.

142.   Guo YJ, Zhou YJ, Yang XL, Shao ZM, Ou ZL. The role and clinical significance of the CXCL17-CXCR8 (GPR35) axis in breast cancer. Biochem Biophys Res Commun. 2017 Nov 25;493(3):1159–67.

143.   Pitteri SJ, Kelly-Spratt KS, Gurley KE, Kennedy J, Buson TB, Chin A, et al. Tumor microenvironment-derived proteins dominate the plasma proteome response during breast cancer induction and progression. Cancer Res. 2011 Aug 1;71(15):5090–100.

144.   Wang CQ, Tang CH, Tzeng HE, Jin L, Zhao J, Kang L, et al. Impacts of RETN genetic polymorphism on breast cancer development. J Cancer. 2020 Feb 20;11(10):2769–77.

145. Aziz MA, Akter T, Sarwar MS, Islam MS. The first combined meta‑analytic approach for elucidating the relationship of circulating resistin levels and RETN gene polymorphisms with colorectal and breast cancer. Egyptian Journal of Medical Human Genetics. 2022 Mar 1;23(1):1–19.

146. Smedley A. "race" and the construction of human identity. Am Anthropol. 1998 Sep;100(3):690–702.

147. Rusidzé M, Adlanmérini M, Chantalat E, Raymond-Letron I, Cayre S, Arnal JF, et al. Estrogen receptor-α signaling in post-natal mammary development and breast cancers. Cell Mol Life Sci. 2021 Aug;78(15):5681–705.

148. Asselin-Labat ML, Sutherland KD, Barker H, Thomas R, Shackleton M, Forrest NC, et al. Gata-3 is an essential regulator of mammary-gland morphogenesis and luminal-cell differentiation. Nat Cell Biol. 2007 Feb;9(2):201–9.

149. Watson PH, Chia SK, Wykoff CC, Han C, Leek RD, Sly WS, et al. Carbonic anhydrase XII is a marker of good prognosis in invasive breast carcinoma. Br J Cancer. 2003 Apr 7;88(7):1065–70.

150. Allison KH, Hammond MEH, Dowsett M, McKernin SE, Carey LA, Fitzgibbons PL, et al. Estrogen and Progesterone Receptor Testing in Breast Cancer: American Society of Clinical Oncology/College of American Pathologists Guideline Update. Arch Pathol Lab Med. 2020 May;144(5):545–63.

151. Scabia V, Ayyanan A, De Martino F, Agnoletto A, Battista L, Laszlo C, et al. Estrogen receptor positive breast cancers have patient specific hormone sensitivities and rely on progesterone receptor. Nat Commun. 2022 Jun 6;13(1):3127.

152. Swain SM, Shastry M, Hamilton E. Targeting HER2-positive breast cancer: advances and future directions. Nat Rev Drug Discov. 2023 Feb;22(2):101–26.

153.   Madrid-Paredes A, Cañadas-Garre M, Sánchez-Pozo A, Calleja-Hernández MÁ. Non-HER2 signaling pathways activated in resistance to anti-HER2 therapy in breast cancer. Breast Cancer Res Treat. 2015 Oct;153(3):493–505.

154.   Sahlberg KK, Hongisto V, Edgren H, Mäkelä R, Hellström K, Due EU, et al. The HER2 amplicon includes several genes required for the growth and survival of HER2 positive breast cancer cells. Mol Oncol. 2013 Jun;7(3):392–401.

155.   Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y, et al. Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. J Clin Invest. 2011 Jul;121(7):2750–67.

156.   Pegram M, Jackisch C, Johnston SRD. Estrogen/HER2 receptor crosstalk in breast cancer: combination therapies to improve outcomes for patients with hormone receptor-positive/HER2-positive breast cancer. NPJ Breast Cancer. 2023 May 31;9(1):45.

157.   Burguin A, Diorio C, Durocher F. Breast Cancer Treatments: Updates and New Challenges. J Pers Med [Internet]. 2021 Aug 19;11(8). Available from: http://dx.doi.org/10.3390/jpm11080808

158.   Mazumder A, Shiao S, Haricharan S. HER2 Activation and Endocrine Treatment Resistance in HER2-negative Breast Cancer. Endocrinology [Internet]. 2021 Oct 1;162(10). Available from: http://dx.doi.org/10.1210/endocr/bqab153

159.   Foster ED, Deardorff A. Open Science Framework (OSF). J Med Libr Assoc. 2017 Apr;105(2):203.

160.   Merkel D. Docker: lightweight Linux containers for consistent development and deployment. Linux J. 2014 Mar 1;2014:2.