Title: Evolutionary feedbacks for *Drosophila* aggression revealed through experimental evolution

Authors: Anna R. Girardeau[1], Grace E. Enochs[1], Julia B. Saltz[1]*

Affiliations:

[1]Biosciences, Rice University; Houston TX 77005, USA

*Corresponding author. Email: Julia.b.saltz@rice.edu

Preprint available: https://www.biorxiv.org/content/10.1101/2024.08.30.610518v1

Classification: Biological Sciences -- Evolution

Keywords: Indirect genetic effects, aggression, Drosophila melanogaster, evolutionary feedbacks, experimental evolution

Abstract: Evolutionary feedbacks occur when evolution in one generation alters the
environment experienced by subsequent generations, and are an expected result of Indirect
Genetic Effects (IGEs). Hypotheses abound for the role of evolutionary feedbacks in climate
change, agriculture, community dynamics, population persistence, social interactions, the genetic
basis of evolution, and more; but evolutionary feedbacks have rarely been directly measured
experimentally, leaving open questions about how feedbacks influence evolution. Using
experimental evolution, we manipulated the social environment in which aggression was
expressed and selected in fruit fly (*Drosophila melanogaster*) populations to allow or limit
feedbacks. We selected for increased male-male aggression while allowing either positive,
negative, or no feedbacks, alongside unselected controls. We show that populations undergoing
negative feedbacks had the weakest evolutionary changes in aggression, while populations
undergoing positive evolutionary feedbacks evolved supernormal aggression. Further, the
underlying social dynamics evolved only in the negative feedbacks treatment. Our results
demonstrate that IGE-mediated evolutionary feedbacks can alter the rate and pattern of
behavioral evolution.

Significance: Evolutionary feedbacks occur when evolution in one generation alters the
environment experienced by subsequent generations, thereby influencing subsequent evolution.
While feedbacks are thought to be important, they have rarely been directly observed because
they happen over many generations and alongside other confounding processes. Here, we studied
fruit flies, *Drosophila melanogaster*. In each population, we selected the most aggressive males
in experimental treatments expected to produce positive, negative, or no feedbacks. Because flies
have short lifespans, we could watch evolution as it happened in the lab over 24 generations. As
predicted, negative feedbacks slowed down behavioral change over generations, while positive
feedbacks accelerated it. Fly "boxing" was especially affected by selection under positive
feedbacks. Our results highlight how small decisions, such as retaliation, can have profound
effects on evolution itself.

Main Text:
Evolutionary feedbacks occur when trait evolution in one generation alters evolutionary
processes in subsequent generations (1–3). Traditional behavioral biology focused on how the
environment affects behavior, often ignoring reciprocal effects whereby behaviors shape the
environment (4). Yet, organisms typically—perhaps universally—alter their environments (5–7),
e.g., by excreting waste, eating other organisms, and behaving socially. This interplay has
important implications for evolution, because the environment mediates the genetic basis and
fitness consequences of variation in traits. Consequently, the potential for evolutionary feedbacks
between organisms and their environments has come into focus as a major, but largely
unexplored, force driving phenotypic evolution (1, 8, 9), even on short timescales (1, 10).
Hypotheses abound for the role of feedbacks in climate change (11, 12), community dynamics
(13), population persistence (10), social interactions (14), the genetic basis of evolution (15–17),
and more.
95
For social behaviors, the "environment" in which behaviors are expressed is formed by the
behaviors of interacting conspecifics. Indirect genetic effects (IGEs) explicitly consider how
genes expressed in these social partners may influence a focal individual's behavior (18–21).
Note that "indirect" refers here to the effect of one individual's genes on another individual's
behaviors, and is distinct from other "indirect effects" in ecology (12). IGEs stand in contrast to
"direct" genetic effects, DGEs, whereby an individual's own genotype affects its own behavior.
For example, guppies (*Poecilia reticulata*) are more willing to inspect a predator if they are
paired with a partner who is also willing to inspect; and these behaviors are heritable (22). This
example illustrates how social interactions may be determined by the genotypes of the
participants. Indirect genetic effects are widespread, and have been identified for a wide range of
traits and taxa (23–26).
107
Under indirect genetic effects, the environment itself can evolve: evolutionary changes in
population-mean behavior also represent changes in the environment in which the behavior is
expressed (18, 27). Indeed, many familiar evolutionary feedbacks are examples of evolutionary
outcomes of indirect genetic effects for social behaviors, such as Fisherian runaway selection and
arms races (8). The existence of these feedbacks highlights the challenge of predicting behavioral
evolution over more than one generation (28–31). Yet, empirical demonstrations of such
feedbacks are scarce (29, 32–34), likely because isolating the effects of evolutionary feedbacks –
if any – is nearly impossible for wild populations (35).
116
Furthermore, theoretical predictions about how IGEs will govern evolution are most relevant to
situations where the IGEs themselves remain relatively stable over generations (36). As with any
evolutionary process, if the underlying mechanisms – here, ways in which the behaviors of social
partners alter the behavior of a focal individual– themselves evolve, then these changes may alter
future evolution. Previous studies have shown that the magnitude and direction of IGEs can
readily evolve, in both the lab and the field (36, 37).
123
To directly identify the effect(s) of evolutionary feedbacks, we founded 12 replicate *Drosophila*
melanogaster populations from the same wild base population and subjected them to 24
generations of selection for increased aggression, while allowing or limiting the opportunity for
different evolutionary feedbacks. Experimental evolution is an important tool for understanding
evolutionary processes, because putative causal factors may be directly manipulated under

controlled conditions, and the resulting evolutionary trajectories can be carefully monitored (38–
41). In experimental evolution studies, the goal is to manipulate the presence, type, or magnitude
of a *single* evolutionary process to identify its effects on evolutionary outcomes. Such studies
thus provide an important complement to field studies, in which multiple evolutionary processes
may be acting simultaneously.

IGEs have been well-established for aggressive behavior in flies (42, 43) and other species (44,
45). Aggression is inherently a highly interactive trait, because animals plastically adjust their
fighting behavior based on the responses of their opponents. Thus, the aggressive behavior of
one individual is an environment that influences the aggressiveness of opponents. This plasticity
occurs on two distinct timescales. On short timescales, i.e., during fights, aggression by one
individual typically provokes retaliatory aggression by its opponent (in flies: (43, 46) across taxa:
(45, 47)). Such **retaliation** represents a social environment that augments aggression: the more
aggressive your opponent is, the more aggressive you are in return. Over longer timescales, i.e.,
across fights, flies who have previously experienced aggressive attacks show reduced
aggressiveness in subsequent encounters (43, 48–51). **Prior experience effects** thus represent a
social environment that reduces aggression: the more aggressive your first opponent is towards
you, the *less* aggressive you will be in your next encounter.

These social dynamics -- retaliation and prior experience effects -- are predicted to produce
potent and opposite evolutionary feedbacks (45). Selection for increased aggressiveness means
that future generations will experience a highly-aggressive social environment—an environment
in which retaliation will augment aggression even further. This process represents a positive
feedback because the direct effect of selection and the indirect effects of changing the
environment in which the behavior is expressed act in the same direction—i.e., to increase
aggression, augmenting the expected evolutionary response.

155
156

157
158

Figure 1. Study design and predictions. Our selection treatments manipulated the opportunity for different evolutionary feedbacks. Left: during selection (generations 1-24), 100 male flies per generation were measured for aggression in their treatment-specific social environments and times. Red boxes indicate the social environment and timing in which the 50 most aggressive males were chosen to become fathers for the next generation. Right: at the end of the experiment, all populations were measured for prior experience effects to test whether these social dynamics had themselves evolved.

To test this prediction, we had a **Positive Feedbacks (P) treatment.** To permit positive feedbacks, two naïve males from the same experimental population were paired each generation, for a total of 100 males in 50 pairs. Total aggression of each male was measured, and the 50 most aggressive individuals were selected (Figure 1). This treatment is expected to produce positive feedbacks because, as aggression evolves (i.e., the population responds to selection), males will face increasingly-aggressive opponents that will stimulate increased retaliatory responses, exaggerating the population's response to selection.

In contrast, with prior experience effects, as the population-mean aggressiveness increases (i.e., the population responds to selection), more individuals will experience aggressive opponents in their first aggressive encounter, resulting in increasingly suppressed aggressiveness in subsequent encounters. This process represents a negative feedback because the direct effect of selection and the indirect effect of changing the environment in which the behavior is expressed are in opposite directions—selection acts to increase aggression, provoking more potent prior experience effects that reduce aggression, dampening the expected evolutionary response.

To test this prediction, we had a **Negative Feedbacks (N) treatment.** In this treatment, males were subjected to 2 days of behavioral analysis (Figure 1). On Day 1, two naïve males from the same experimental population were paired, for a total of 100 males in 50 pairs; this experience provides the opportunity for prior experience. The next day, Day 2, the same males were then paired with a standard opponent and aggressiveness was again measured. These standard opponents were males from the DGRP-208 inbred genotype, an inbred line whose progenitors were wild-collected and then highly inbred over many generations. Importantly, the inbred nature of these opponents mean that every male standard opponent was genetically identical to every other standard opponent in every generation. Selection was based only on males' day 2 aggressiveness: males from the experimental population were compared for total aggression on Day 2, and the 50 most aggressive males were selected.

It may seem intuitive that "negative" feedbacks should involve selection for reduced aggression—however, this is not the case. Negative feedbacks occur when behavioral plasticity and evolutionary changes in behavior act in **opposite** directions, i.e., there is a negative covariance between behavior of opponents (18). Here, this occurs because selection for increased aggression should result in behavioral plasticity (prior experience effects) that reduces aggression.

To quantitatively identify the effects of feedbacks, we compared behavioral evolution in the **Positive feedbacks** and **Negative feedbacks** treatments to a **No feedbacks (0) treatment.** To limit opportunities for feedbacks, 100 males from the experimental population were paired with standard opponents – the same genotype of standard opponents as in the **N** treatment on day 2 (i.e., DGRP-208 inbred genotype). Males from the experimental population were compared for

207 total aggression, and the 50 most aggressive males were selected. This treatment abrogates the
208 opportunity for IGE-mediated evolutionary feedbacks because the standard genotype was
209 genetically identical in every generation, meaning the social environment was not allowed to
210 evolve.
211
212 Finally, we had **Control (C)** populations that did not undergo selection.
213
Each replicate was founded by collecting 4 samples of the base population, with each sample
consisting of 100 males and 100 females. These 4 samples became the 4 populations from this
replicate, and each one was subjected to a different experimental treatment: one population from
each of the 3 selection treatments and one unselected control population. In generation 0, 100
males from each population were paired together and measured for baseline aggression to ensure
similar starting points. In generations 1-24, each population was subjected to evolution for
increased aggression (except unselected control populations). Each population was measured for
aggression every generation as part of the selection protocol (see below), except control
populations, which were measured at generation 0 and again at the midway point and at the end
of the experiment. Populations within each replicate were carefully matched demographically
(see Materials and Methods and Table S1) to rule out any confounding effects of population
history on evolutionary change. In the final generation, samples from all populations were
measured for prior experience effects so that we could test whether prior experience effects
evolved (Figure 1, right), and the experiment was concluded.
228
229 By measuring 39,018 males over 25 generations in 12 initially-identical populations, our design
230 allowed us to directly identify how evolutionary feedbacks shape changes in aggressive
231 behavior.
232
Results:
In every generation, 100 males from each population (except for unselected controls) were scan-
sampled (52) for aggression 24 times in one day during peak activity periods. For each
observation, trained observers recorded the presence or absence of 3 behavioral components of
aggression: fencing - both flies extending and engaging with each other's legs by tapping and/or
pushing; lunging - one fly rearing on his back legs and slamming his foretarsi onto his opponent;
and boxing - both flies rearing on their hindlegs, "kangaroo style," and engaging each other with
their front legs by pushing and/or hitting (46). When fencing and lunging occurred, observers
noted which individual was the aggressor. Boxing can only occur when both individuals are
engaged. These measures are considered behavioral components of aggression (53), rather than
independent behaviors, as they very often occur in a sequence during aggressive encounters (46).
Finally, observers noted whether either of the flies appeared to be temporarily stuck on the food
patch, as occasionally occurred.
246
247 We quantified total aggression as the unweighted sum of these behavioral components. In the
248 selection treatments (Positive feedbacks, Negative feedbacks, and No feedbacks), the 50 males
249 with the highest total aggression (in their treatment-specific social environment) were chosen to
250 become fathers to the next generation alongside randomly-chosen, unmated females from their
251 population.
252
253 After confirming that populations showed indistinguishable aggression at the beginning of the
254 experiment (Table S3), we compared the evolutionary trajectories of total aggression across the

three selection treatments and the controls using Generalized Linear Mixed Models (GLMM).
Our analysis revealed a significant interaction between treatment and generation, indicating that
treatments differed in how aggression changed over generations (Figure 2A, Table 1: Model 1).
All 3 selection treatments evolved increased aggression, but treatments differed in the rate of
behavioral change: the Negative feedbacks treatment showed slower evolutionary changes in
aggression compared to the Positive feedbacks and No feedbacks treatments, while the Positive
feedbacks and No feedbacks treatments showed similar rates of change, with a non-significant
trend towards faster evolution in the Positive feedbacks treatment (Figure 2A, Table 2: total
aggression). In contrast, unselected control populations differed significantly from all other
treatments and did not evolve increased aggression (Figure 2A, Table 2: total aggression).

Figure 2. Evolutionary response to selection. **A.** Y axis is least-squares means of total
aggression adjusted for other terms in the model, estimated every generation from model 1. Grey
error bars indicate standard errors estimated from the model. **B.** Y axis is least-squares means
for each behavioral component of aggression, adjusted for other terms in the model, estimated
every generation from model 2. Note that for Boxing, the slopes for the 0 and N treatments are so

272 low and similar that they cannot be distinguished visually on the plot. 0 indicates the No
273 feedbacks treatment, N indicates the Negative feedbacks treatment, P indicates the Positive
274 feedbacks treatment, and C indicates the unselected controls. N = 3 populations per treatment.
275
During the experiment, it was apparent even through casual observation that the Positive
feedbacks populations showed extreme levels of boxing compared to the other treatments (Figure
2B). Thus, we used a trait covariance approach to analyze how each component of aggression
evolved. Lunging, fencing, and boxing were treated as repeated measures of each individual
male. The resulting GLMM describing evolution in the selected populations (Model 2) revealed
a significant 3-way interaction between generation, treatment, and component (Figure 2B, Table
1: Model 2), meaning that treatment differences in evolutionary change differed between
fencing, lunging, and boxing. For fencing, we found that all three selection treatments evolved
increased fencing; P and 0 had indistinguishable slopes, with the Negative feedbacks treatment
evolving slowest (Table 2: fencing). For lunging, only the No feedbacks treatment showed
significant evolutionary changes (Table 2: lunging). For boxing, we found that the Positive and
Negative Feedbacks treatments showed significant evolution of boxing, but not the No feedbacks
treatment (Table 2: boxing). The Positive feedbacks treatment in particular showed dramatically
faster evolution of boxing compared to the other treatments; indeed, on the scale of the original
data (i.e., not the latent scale), the estimated slope for the P treatment was more than 35 times
greater than the corresponding estimate for the N treatment (Figure 2B, Table 2: boxing).
Together, results from Models 1 and 2 demonstrate that evolutionary feedbacks alter
populations' phenotypic evolution.
294

model	results				
	term	parameter estimate (SE)	chisq	df	p
model 1: did feedbacks influence the evolution of aggression?	Generation	-0.02144 (0.01117)	3.6831	1	0.05496
	Treatment	**.**	**385.1448**	**3**	**< 2.2e-16**
	focal.stuck	**-0.08493 (0.01579)**	**28.9475**	**1**	**7.437E-08**
	Generation:Treatment	**.**	**44.6871**	**3**	**1.078E-09**

model	results				
	term	parameter estimate (SE)	chisq	df	p
model 2: how did each component of aggression evolve?	**Generation**	**0.094761 (0.034816)**	**7.4078**	**1**	**0.00649**
	Treatment	**.**	**104.4534**	**2**	**< 2.2e-16**
	component	**.**	**222.8693**	**2**	**< 2.2e-16**
	focal.stuck	**-0.087871 (0.015889)**	**30.5832**	**1**	**3.198E-08**
	Generation:Treatment	.	1.2900	2	0.52466

Generation:component	.	15.9748	2	0.00034
Treatment:component	.	69.3866	4	3.058E-14
Generation:Treatment:component	.	23.1491	4	0.00012

model	results			
	non-linear term	edf (measure of "wiggliness")	chisq	reference df	p
model 3: did prior experience effects evolve over generations?	day 1 opponent aggression	4.61099	873.724	5.489	< 2e-16
	day 2 opponent aggression	2.78316	261.970	3.392	< 2e-16
	Generation:day 1 opponent aggression	2.24636	14.948	2.638	0.00381
	Generation:day 2 opponent aggression	1.01328	0.159	1.026	0.69746

model	results			
	non-linear term	edf (measure of "wiggliness")	chisq	reference df	p
model 4: did prior experience effects differ across treatments at the end of the experiment?	effect of day 1 opponent behavior in control populations	1.0000937	12.656	1	0.000374
	effect of day 2 opponent behavior in control (C) populations	1.0000411	2.256	1	0.133111
	difference in effect of day 1 opponent behavior between C and 0 treatment	1.0000432	1.042	1	0.307341
	difference in effect of day 1 opponent behavior between C and P treatment	1.000115	0.028	1	0.867075
	difference in effect of day 1 opponent behavior between C and N treatment	1.0000181	3.931	1	0.047411
	difference in effect of day 2 opponent behavior between C and 0 treatment	1.0000268	0.039	1	0.843981
	difference in effect of day 2 opponent behavior between C and P treatment	1.0000804	0.002	1	0.964223
	difference in effect of day 2 opponent behavior between C and N treatment	1.000015	0.517	1	0.472197

Table 1. Evolutionary feedbacks influenced the evolution of aggression and prior experience effects. Important results from each model are presented. Models 1 was a GLMM describing the evolution of aggression (generations 1-24). In Model 2, we fitted a GLMM to the selected treatments only (not controls) analyzing aggression by behavioral component -- lunging, fencing, and boxing -- while accounting for repeated measurements of the same male. Model 3 was a Generalized Additive Mixed Model (GAMM) using only data from the N treatment, which is the only treatment for which prior experience effects were measured during generations 1-24. We tested whether changes in aggressive behavior over days varied across generations and due to opponent aggression on days 1 and 2 (i.e., prior experience effects). EDF describes the degree of non-linearity estimated for each term. In Model 4, we tested whether prior experience effects differed across treatments at the end of the experiment. We fitted a GAMM describing the effects of opponent aggression on each day in the Control treatment as the "reference smooth;" the other terms represent "difference smooths" that estimate differences between each treatment's prior experience effects and those of the Control populations. For details please see Materials and Methods.

311
312

model 1: total aggression	selection treatment	estimated trend over generations (SE)	is the slope greater than 0?	differences among treatments
evolution of total aggression	C	-0.0122 (0.00663)	no (z ratio = -1.840, p = 0.0657)	a
	O	0.0422 (0.01009)	**yes** (z ratio = 4.183, p < 0.001)	c
	N	0.0205 (0.00536)	**yes** (z ratio = 3.813, p = 0.0001)	b
	P	0.0547 (0.01345)	**yes** (z ratio = 4.066, p < 0.0001)	c
model 2: behavioral components	selection treatment	estimated trend over generations (SE)	is the slope greater than 0?	differences among treatments
evolution of fencing	O	2.22e-02 (3.64e-03)	**yes** (z ratio = 6.094 , p <.0001)	a
	N	1.15e-02 (2.12e-03)	**yes** (z ratio = 5.425, p <.0001)	b
	P	2.08e-02 (4.11e-03)	**yes** (z ratio = 5.057, p <.0001)	a
evolution of lunging	O	3.10e-03 (5.82e-04)	**yes** (z ratio = 5.333, p <.0001)	a
	N	6.73e-05 (2.07e-04)	no (z ratio = 0.325, p = 0.7453)	b

	selection treatment	estimated effect of day 1 experience in gen 25 (SE)	is the slope greater than 0?	differences among treatments
	P	-6.92e-05 (4.99e-04)	no (z ratio = -0.138, p = 0.8899)	b
evolution of boxing	O	7.49e-05 (4.95e-05)	no (z ratio = 1.513, p = 0.1303)	b
	N	1.21e-04 (1.21e-04)	yes (z ratio = 2.877, p = 0.0040)	b
	P	6.70e-03 (1.10e-03)	yes (z ratio = 6.071, p <.0001)	a
model 4: prior experience effects	**selection treatment**	**estimated effect of day 1 experience in gen 25 (SE)**	**is the slope greater than 0?**	**differences among treatments**
	C	-0.259 (0.0727)	yes (t ratio = -3.558, p = 0.0004)	ab
change in aggression over days	O	-0.161 (0.0632)	yes (t ratio = -2.549, p = 0.0110)	a
	N	-0.448 (0.0636)	yes (t ratio = -7.041, p <.0001)	b
	P	-0.273 (0.0508)	yes (t ratio = -5.384, p <.0001)	ab

Table 2. Evolutionary effects of feedbacks. Estimated marginal means of linear trends extracted from the indicated model . Slopes are presented on the scale of the original data. We tested whether each slope differed from 0 using t-tests, and we tested for slope differences between treatments. Comparisons between slopes were adjusted for multiple testing using the tukey method.

Feedbacks for social behavior depend on the underlying social dynamics – here, retaliation and prior experience effects. To test whether these social dynamics changed during our experiments, we fitted additive mixed models (GAMM) that can account for non-linear effects of opponent behavior on focal male aggression. For retaliation, we tested whether the magnitude of the covariance between interacting males changed directionally over generations. While retaliation was present in all populations tested, none of the treatments showed evidence that the magnitude or form of retaliation changed over generations (Table S4).

To test for evolution of prior experience effects, we fitted a GAMM to data from the Negative feedbacks treatment (Model 3). We measured prior experience effects for each male as the change in aggression across days (i.e., focal male Day 2 aggression – focal male Day 1 aggression, (54, 55)). Our model uncovered a significant interaction between Day 1 opponent behavior and Generation (Table 1: model 3), indicating that the effects of social experience on male behavioral plasticity evolved over generations (Figure 3A – compare Generation 1 (right) to Generation 12 (center) to Generation 24 (right)). To test whether this evolutionary change in prior experience effects was unique to the Negative feedbacks treatment, we compared prior experience effects between the Positive feedbacks, Negative feedbacks, No feedbacks, and Control populations at the end of each replicate (Model 4). Our analysis (Table 1: model 4)

338 revealed that prior experience effects differed significantly from the unselected Control
339 populations only in the Negative feedbacks treatment (Figure 3B, Table 2: model 4).
340
341

Figure 3. Prior experience effects evolved in the Negative Feedbacks treatment. A: prior experience effects over 24 generations in the negative feedbacks treatment. Y axis is the change in behavior over days, where positive values mean that focal male aggression was enhanced after day 1 social experience, negative values mean that focal male aggression was suppressed after day 1 social experience, and zero indicates no effect of prior experience. The x-axis is the aggression of the focal males' day 1 opponent, and the line represents the cubic spline from Model 2 with associated standard error. Comparison of Generation 1 (A, left) with Generation 12 (A, center) and Generation 24 (A, right) illustrates that prior experience effects evolved. B: comparison of prior experience effects at the end of the experiment for each treatment. The x-axis is the aggression of the focal males' day 1 opponent, and the line represents the cubic spline from Model 4 with associated standard error. Only the N treatment showed evolutionary changes in prior experience effects.

Discussion:

Evolutionary feedbacks are widely hypothesized to be both common and important, but quantifying the effects of evolutionary feedbacks in the midst of other population-genetic processes is challenging, leaving open questions about how evolutionary feedbacks play out in populations over time. Here, we manipulated the social environment in which aggression was expressed and selected to allow or limit the opportunity for indirect genetic effect (IGE)-mediated evolutionary feedbacks to contribute to evolution (Figure 1). We tracked the resulting evolutionary dynamics over 24 generations in 12 populations, requiring behavioral measurement of tens of thousands of males. Our results broadly support predictions (Figure 2, Tables 1-2), i.e., that positive feedbacks should accelerate phenotypic evolution while negative feedbacks should slow it (18, 19, 27, 31). As unselected control populations did not evolve increased aggression, we can be confident that these evolutionary changes are due to our experimental selection treatments. Thus, our results demonstrate that evolutionary feedbacks alter populations' phenotypic evolution.

Not every component of aggression evolved the same way; in particular, the evolutionary changes in the Positive feedbacks treatment were dominated by the very striking evolution of supernormal boxing behavior (Figure 2B: boxing). Boxing is a mutually-expressed high-level component of aggression in which active participation by both males is required. Thus, small increases in aggressiveness over generations may have been magnified in the Positive feedbacks treatment through retaliation, making mutually-expressed, high-intensity fights substantially more common.

Evolutionary feedbacks are predicted to alter both the magnitude of selection and realized heritability (27, 31) to shape phenotypic change over time. Existing statistical approaches to disentangle these processes are focused on normally-distributed traits (56) limiting our ability to piece apart these processes in our data. We consider it most likely that IGE-mediated effects on both selection strength and realized heritability played roles in our observed results. A recent meta-analysis provides strong evidence that IGEs can alter trait heritability, particularly for behaviors, as measured through standing variation (i.e., heritability, not realized heritability, (57)). On the selection side, we performed truncation selection with the most aggressive 50% of males selected to reproduce each generation. As the populations evolved increased average

aggression, the absolute number of aggressive behaviors required of a male to qualify for the selected fraction also increased. Studying evolutionary feedbacks under alternative forms of selection would be an exciting future direction.

Another key dimension of evolution under evolutionary feedbacks is that the behavioral dynamics that produce feedbacks – here, retaliation and prior experience effects – may themselves evolve. While we found no evidence for evolutionary changes in retaliation, prior experience effects showed evolutionary changes in the Negative feedbacks treatment only (Figure 3). Inspection of the reaction norms describing the effect of prior experience on later aggression, both across generations of the Negative feedbacks treatment (Figure 3A) and across treatments at the end of the experiment (Figure 3B, table 2) indicate that prior experience effects evolved to be more pronounced: being attacked by an opponent male on Day 1 produced more severe depression of aggression on Day 2, compared to generation 1 and to other treatments (Table 2, model 4). These findings suggest that negative feedbacks and their evolutionary consequences would be exacerbated under continued selection. Predicting and testing the evolution of social dynamics as indirect targets of selection is a critical future direction for understanding evolutionary feedbacks (36, 37).

Conclusions:
The existence and potency of IGE-mediated evolutionary feedbacks highlight how simple behavioral choices – such as responding to an aggressive attack – can scale up to impact the entire evolutionary process. Our results provide some of the clearest evidence to date that evolutionary changes can alter the environment to produce evolutionary feedbacks that, over time, influence evolutionary change. Together, these findings indicate that evolutionary feedbacks may not be a unique or unusual feature of evolution in specific cases, but rather, a widespread outcome of the ways that organisms interact with their environments. Thus, evolutionary feedbacks should be considered in a broad range of evolutionary studies, even when IGEs and related processes are not the explicit focus.

Methods:
Experimental Overview: *Replicate populations were selected for increased aggression under different social conditions, alongside unselected controls*
We had 3 selection treatments representing selection for increased aggression under positive, negative, or no evolutionary feedbacks, as described below. In addition, we had control populations that did not undergo selection.

In generation 0, 100 males from each population were paired together and measured for baseline aggression to ensure similar starting points. In generations 1-24, each population was subjected to evolution for increased aggression, or not, corresponding to their assigned treatments. Each population was measured for aggression every generation as part of the selection protocol (see below), except control populations, which were measured at generation 0 and again at the midway point (generation 12 or 13). In the final generation (generation 25 for replicates 1 and 2, generation 26 for replicate 3), samples from all populations were measured for prior experience effects so that we could test whether prior experience effects evolved, and the experiment was concluded.

Replicate 1 was founded in July 2021, Replicate 2 was founded in September 2021, and Replicate 3 was founded in February 2022.

Experimental details: flies and rearing conditions.
Base population: As our base population, we used an outbred population of flies originally
collected from Coral Gables, FL, by the de Bivort lab. Mated females were collected from the
same wild population over a period of June-November 2018 and their descendants were kept at
large population sizes in the de Bivort lab (61).

Standard opponents. We used standard opponents in several of the treatments as explained
below. These were males from the DGRP-208 inbred genotype, an inbred line whose progenitors
were wild-collected and then highly inbred over many generations. DGRP-208 was chosen
because it showed intermediate aggression in prior studies (Hutchins et al 2024). Importantly, the
inbred nature of these opponents mean that every male standard opponent was genetically
identical to every other standard opponent in every generation.

Standard opponents were reared for trials by adding 10 unmated DGRP-208 females and 10
DGRP-208 males to a vial containing standard fly food. This approach minimizes variation in
rearing conditions across vials. Male progeny were collected within 12 hours of eclosion and
housed with other standard opponent males for 5-7 days prior to testing.

Rearing and preparation for aggression experiments
Experimental populations were reared on high-protein food (consisting of standard fly food
supplemented with additional dead yeast to produce a 4:1 protein:carbohydrate ratio; (62)) in 6oz
plastic bottles. Each bottle was founded by 25 males and 25 unmated females. The high protein
food ensured that sufficiently large numbers of males eclosed around the same time, allowing us
to conduct the experiment. Newly-eclosed unmated females were housed in same-sex groups.
Newly-eclosed experimental males were marked with a small dot of yellow or blue acrylic paint
under anesthesia so that we could distinguish individuals during behavioral trials. After painting,
males were isolated for 5-7 days. This approach ensured that males were recovered from
anesthesia, but had no adult social experience prior to behavioral trials.

Selection on aggressive behavior
Aggression arenas
Arenas were composed of two small Petri dishes (3 cm x 1 cm) taped together. The floor of the
arena was filled with standard fly food medium and a dot of active yeast. This is a typical arena
design for measuring aggression in flies (e.g., (43)).

Measuring aggression
Because of the large number of males that needed to be measured simultaneously, we used an
instantaneous scan sampling approach to measure aggression (52). Flies were added to arenas in
the morning, at approximately 9:00am. Every 5 minutes during the following 60 minutes, we
noted whether flies in each arena were engaged in any of 3 aggressive behaviors. A second 60-
minute observation period occurred from 3:00-4:00 pm. Thus, we collected 24 total observations
of each fly.

For each observation, trained observers recorded the presence or absence of 3 components of
aggressive behavior: fencing - both flies extending and engaging with each other's legs by
tapping and/or pushing; lunging - one fly rearing on his back legs and slamming his foretarsi
onto his opponent; and boxing - both flies rearing on their hindlegs, "kangaroo style," and

engaging each other with their front legs by pushing and/or hitting (46, 63). When lunging occurred, observers noted which individual was the aggressor. Fencing and boxing are mutually performed behaviors so both individuals were counted as engaging in each instance of these. Finally, observers noted whether either of the flies appeared to be temporarily stuck on the food patch, as occasionally occurred. Observers did not know which flies were from which treatments while measuring aggression. To choose flies to become fathers of the next generation (in the selection treatments), we computed total aggression, which was the unweighted sum of fencing, lunging, and boxing instances observed for each individual male. For each selected population, the 50 males with the highest total aggression were selected each generation. If fewer than 50 males showed non-zero aggression, then all the males that showed any aggression were selected alongside randomly-chosen additional males until we reached 50 total males. For example, if only 45 males showed aggression, 5 males that did not show aggression (during our sampling periods) would be randomly chosen, resulting in 50 fathers for the next generation.

Founding the next generation. After selection, the 50 chosen males were placed into food bottles with 50 randomly chosen unmated females from their own population. The progeny of these flies became the next generation.

Deviations from the protocol. Due to unavoidable variability in fly eclosion rates, there were some generations in which fewer than 100 males were available from one or more of the populations. In these cases, the other populations in that replicate were subsampled to produce equal numbers of males considered for selection. For example, if one population only had 70 males available for testing, we measured those 70 males alongside 100 males from the *other* populations in that replicate. Before selection, we randomly discarded data from 30 males from the other populations, and chose the 50 most aggressive males from the remaining 70 data points.

For replicate 3 only, we had 3 generations where no selection was possible. In Generation 15, flies eclosed almost a week later than usual, possibly due to unusual weather in Houston TX. Since we were unsure of whether these flies were healthy and representative of the evolving population, we chose to allow random parents to found the next generation rather than conducting selection. Later, the standard opponent colonies became contaminated with bacteria resulting in insufficient standard opponents in generations 22, 23, and 25 (for Replicate 3 only). To ensure identical demographic histories among Replicate 3 populations, we therefore chose random flies to propagate all the populations during these "lost" generations. We concluded the experiment in Generation 26, in which we measured prior experience effects. Note that all results are adjusted for these and any other differences between Replicates.

In sum, this approach ensured identical demographic histories among populations from the same replicate. A complete record of the number of males measured for each population in each replicate is available in Table S1 in supplementary materials.

Final generation: test for prior experience effects. In the final generation, males from selected and control populations were reared as normal. But instead of their normal testing protocol, males from all populations were tested for prior experience effects: males from each population were paired together on Day 1 and aggression was measured, then those same males were paired with a standard opponent the next day and their aggression was measured again (i.e., as in (43, 51)).

Analysis Methods

Overview. We fit a series of models using data from generations 1-24 to test for evolutionary changes in aggression, retaliation, and prior experience effects, and differences in these slopes among treatments. Finally, we analyzed prior experience effect differences in the final generation of each replicate. All models included a random effect of Replicate to account for differences among replicates. A full summary of each model, the question that it answers, and its structure, is available in table S2.

In initial models, we found that the relationship between focal male aggression and opponent aggression was highly non-linear. Thus, analyses that included opponent aggression as a predictor were fit as generalized additive mixed models (GAMMs) in the package mcgv, enabling us to capture these nonlinear effects (64). Opponent aggression and its interactions were fit using thin-plate splines, which include penalties for overfitting (64). We allowed the model to fit up to 10 "knots" for each spline. Models not including the effects of opponent behavior were fit as generalized linear mixed models (GLMMs) using the package glmmTMB, enabling us to include zero-inflation terms as needed (65).

Our measure of prior experience effects focused on changes in aggression across days and was calculated as: focal male Day 2 aggression – focal male Day 1 aggression, for each male (54, 55). For these models, we used the scaled t error distribution, which is appropriate for errors that have a heavy-tailed gaussian-type distribution. For models where aggression itself was the response variable, we used a negative binomial error distribution, which is appropriate for overdispersed count data.

Preliminary models showed that fly age (i.e., whether the flies were 5, 6, or 7 days post-eclosion) was never significant, so we did not include this term in final models.

Model fitting and inference. Residual plots from each model were examined to ensure adequate model fit. For GLMMs, residual plots were produced by simulation using the package DHARMa (66); for GAMMs, we used the gam.check function in mcgv. For GLMMs, we also used AIC analysis to decide what zero-inflation terms to include, if any.

In the final models, we used Wald type III tests (implemented in the 'car' package for GLMMs (67), and F tests in the "summary.gam" function in mcgv for GAMMs) to test the significance of fixed effects. We did not conduct inference on random effects.

We estimated slopes, i.e., estimated marginal means of linear trends, representing change in behavior over generations. Slopes were de-transformed to the scale of the original data, which is appropriate for interpreting evolutionary patterns (68). We tested whether each slope differed from 0 using t-tests, and we performed planned contrasts to compare slopes from different treatments. Comparisons between slopes were adjusted for multiple testing using the tukey method. These computations were implemented in the emmeans package (69).

Model details

Model 1: How did aggression evolve? To test how aggression evolved, we modeled total aggression (i.e., the unweighted sum of all aggression components) across generations. This model included data from the selection treatments as well as the controls, starting in generation 0. Fixed predictors were treatment, generation, and their interaction, with a fixed covariate

indicating the number of times the fly appeared stuck on the food patch. We included terms for
replicate, the date the fly was measured, the identities of the trained observers.

Model 2: how did each component of aggression evolve? We used a trait covariance approach
(similar to MANOVA) to analyze how total aggression and each component of aggression
evolved in our experiments. For selection treatments only, we modeled "focal male aggression"
as our response variable, and each individual was represented by three measurements –lunging,
fencing, and boxing-- with a variable, "component," indicating which observation corresponded
to which component of aggression. The model included fixed effects of generation, treatment,
component, and all possible interactions among these variables. We also included a fixed
covariate indicating the number of times the fly appeared stuck on the food patch. We included
the same random effects as in Model 1, plus random effect of individual ID to account for the
non-independence of the 3 measurements from each individual.

Model 3: did prior experience effects evolve in the N treatment? In this model, we used only data
from the N treatment. We tested whether within-individual changes in aggressive behavior over
the 2 days of trials (measured as Day 2 aggression –Day 1 aggression for each male, see above)
varied over generations and due to opponent aggression on days 1 and 2. Generation, opponent
aggression on each day, and 2-way interactions between generation and opponent aggression on
each day were included as fixed effects, with other terms in the model as above. In this model, an
effect of a generation x opponent aggression interaction (on either day) would indicate an
evolutionary change in how males responded to opponent behavior. We included the same
random effects as in Model 1.

Model 4: did prior experience effects evolve in any of the treatments? To compare the magnitude
of prior experience effects across treatments, we tested whether changes in aggressive behavior
across days (measured as Day 2 aggression – Day 1 aggression for each male, see main text)
differed across treatments in Generation 25. Using GAMMs, we fitted the effects of opponent
aggression on each day in the Control treatment as the "reference smooth," and fitted effects that
estimated differences between Control and each of the other treatments for these relationships as
"difference smooths" (70). Significant effects of each difference smooth would indicate
significant differences between this treatment's prior experience effects and those of the control
populations. We included the same random effects as in Model 1.

Model S1: checking assumptions. In this model we confirmed that populations had
indistinguishable levels of aggression at the beginning of the experiment (generation 0) See
Table S3.

Models S2.0, S2.N, and S2.P: did retaliation evolve? In this series of models, we tested whether
retaliation changed over generations in each selection treatment. See Table S4.

Acknowledgments: We thank Charlotte Hovland, Autumn Hildebrand, Philip DuBose, Samitha
Nemirajaiah, Vivian Ha, Jose Ramirez, Maggie Bao, Priya Trakru, Daniela Vodo, and
Brooke Ermias, for logistical assistance. We thank the de Bivort lab, especially Jamilla
Akhund-Zade, for supplying the outbred population.

Funding: This work was supported by the National Science Foundation (Award ID #1856577).

Author contributions:

631   Conceptualization: JBS

632   Methodology: ARG, GEE, JBS

633   Investigation: ARG, GEE, JBS

634   Visualization: JBS

635   Funding acquisition: JBS

636   Project administration: ARG, GEE, JBS

637   Supervision: ARG, GEE, JBS

638   Writing – original draft: JBS

639   Writing – review & editing: ARG, GEE, JBS

641 **Competing interests:** Authors declare that they have no competing interests

642 **Reproducible research:** Data and code needed to reproduce the analyses presented are included
643   as supplemental files.

References

1.   D. M. Post, E. P. Palkovacs, Eco-evolutionary feedbacks in community and ecosystem ecology: interactions
between the ecological theatre and the evolutionary play. *Philosophical Transactions of the Royal Society B:*
Biological Sciences **364**, 1629–1640 (2009).

2.   K. N. Laland, F. J. Olding-Smee, M. W. Feldman, Evolutionary consequences of niche construction and their
implications for ecology. *Proceedings of the National Academy of Sciences of the United States of America*
96, 10242–10247 (1999).

3.   J. N. Thompson, Rapid evolution as an ecological process. *Trends in Ecology & Evolution* **13**, 329–32 (1998).

4.   F. Odling-Smee, K. Laland, M. Feldman, Niche construction. *American Naturalist* **147**, 641–648 (1996).

5.   T. G. Whitham, *et al.*, A framework for community and ecosystem genetics: from genes to ecosystems.
Nature Reviews Genetics **7**, 510–23 (2006).

6.   K. Donohue, *et al.*, Niche construction through germination cueing: life-history responses to timing of
germination in Arabidopsis thaliana. *Evolution* **59**, 771–85 (2005).

7.   J. B. Saltz, A. P. Geiger, R. Anderson, B. Johnson, R. Marren, What, if anything, is a social niche?
Evolutionary Ecology **30**, 349–364 (2016).

8.   B. J. Crespi, Vicious circles: Positive feedback in major evolutionary and ecological transitions. *Trends in*
Ecology & Evolution **19**, 627–633 (2004).

9.   F. Pelletier, D. Garant, A. P. Hendry, Eco-evolutionary dynamics. *Philosophical Transactions of the Royal*
Society B: Biological Sciences **364**, 1483–1489 (2009).

10. M. T. Kinnison, N. G. Hairston, Eco-evolutionary conservation biology: Contemporary evolution and the
dynamics of persistence. *Functional Ecology* **21**, 444–454 (2007).

11. R. Bonduriansky, A. J. Crean, T. Day, The implications of nongenetic inheritance for evolution in changing
environments. *Evolutionary Applications* **5**, 192–201 (2012).

12. J. K. Bailey, *et al.*, Indirect genetic effects: an evolutionary mechanism linking feedbacks, genotypic diversity
and coadaptation in a climate change context. *Functional Ecology* **28**, 87–95 (2014).

13. G. F. Fussmann, M. Loreau, P. A. Abrams, Eco-evolutionary dynamics of communities and ecosystems.
Functional Ecology **21**, 465–477 (2007).

14. J. B. Saltz, B. R. Foley, Natural Genetic Variation in Social Niche Construction: Social Effects of Aggression
Drive Disruptive Sexual Selection in Drosophila melanogaster. *The American Naturalist* **177**, 645–54 (2011).

15. L. M. Meffert, S. K. Hicks, J. L. Regan, Nonadditive genetic effects in animal behavior. *The American
Naturalist* **160 Suppl**, S198-213 (2002).

16. G. C. K. Chiang, *et al.*, PLEIOTROPY IN THE WILD: THE DORMANCY GENE DOG1 EXERTS
CASCADING CONTROL ON LIFE CYCLES. *Evolution* **67**, 883–893 (2012).

17. J. B. Saltz, S. V. Nuzhdin, Genetic variation in niche construction: Implications for development and
evolutionary genetics. *Trends in Ecology and Evolution* **29** (2014).

18. J. Wolf, E. D. Brodie III, J. J. M. Cheverud, A. J. Moore, M. J. Wade, Evolutionary consequences of indirect
genetic effects. *Trends in Ecology & Evolution* **13**, 64–69 (1998).

19. A. J. Moore, E. D. Brodie III, J. B. Wolf, Interacting Phenotypes and the Evolutionary Process: I. Direct and
Indirect Genetic Effects of Social Interactions. *Evolution* **51**, 1352–1362 (1997).

20. P. Bijma, Multilevel selection 4: modeling the relationship of indirect genetic effects and group size. *Genetics*
186, 1029–31 (2010).

21. K. Donohue, Setting the stage: phenotypic plasticity as habitat selection. *International Journal of Plant
Sciences* **164**, S79–S92 (2003).

22. B. H. Bleakley, E. D. Brodie III, Indirect genetic effects influence antipredator behavior in guppies: estimates
of the coefficient of interaction psi and the inheritance of reciprocity. *Evolution* **63**, 1796–806 (2009).

23. J. J. Mutic, J. B. Wolf, Indirect genetic effects from ecological interactions in Arabidopsis thaliana. *Molecular
Ecology* **16**, 2371–81 (2007).

24. B. H. Bleakley, D. J. Parker, E. D. Brodie III, Nonadditive effects of group membership can lead to additive
group phenotypes for anti-predator behaviour of guppies, Poecilia reticulata. *Journal of Evolutionary Biology*
20, 1375–1384 (2007).

25. T. A. Linksvayer, Direct, maternal, and sibsocial genetic effects on individual and colony traits in an ant.
Evolution **60**, 2552–61 (2006).

26. D. Petfield, S. F. Chenoweth, H. D. Rundle, M. W. Blows, Genetic variance in female condition predicts
indirect genetic variance in male sexual display traits. *Proceedings of the National Academy of Sciences of the
United States of America* **102**, 6045–50 (2005).

27. J. W. McGlothlin, A. J. Moore, J. B. Wolf, E. D. Brodie III, Interacting Phenotypes and the Evolutionary
Process. III. Social Evolution. *Evolution* **64–9**, 2558–2574 (2010).

28.   E. Kazancıoğlu, H. Klug, S. H. Alonzo, The evolution of social interactions changes predictions about
704   interacting phenotypes. *Evolution* **66**, 2056–64 (2012).

705   29.   M. D. Greenfield, S. Alem, D. Limousin, N. W. Bailey, The dilemma of Fisherian sexual selection: Mate
706   choice for indirect benefits despite rarity and overall weakness of trait-preference genetic correlation.
707   *Evolution* **68**, 3524–3536 (2014).

708   30.   A. F. Agrawal, E. D. Brodie III, M. J. Wade, On indirect genetic effects in structured populations. *The
709   American Naturalist* **158**, 308–23 (2001).

710   31.   N. W. Bailey, L. Marie-Orleach, A. J. Moore, Indirect genetic effects in behavioral ecology: does behavior
711   play a special role in evolution? *Behavioral Ecology* **29**, 1–11 (2018).

712   32.   S. H. Alonzo, Social and coevolutionary feedbacks between mating and parental investment. *Trends in
713   Ecology & Evolution* **25**, 99–108 (2010).

714   33.   M. Cantor, *et al.*, The importance of individual-to-society feedbacks in animal ecology and evolution. *J Anim
715   Ecol* **90**, 27–44 (2021).

716   34.   Y. G. Araya-Ajoy, D. F. Westneat, J. Wright, Pathways to social evolution and their evolutionary feedbacks.
717   *Evolution* **74**, 1894–1907 (2020).

718   35.   R. A. Duckworth, S. M. Aguillon, Eco-evolutionary dynamics: investigating multiple causal pathways linking
719   changes in behavior, population density and natural selection. *Journal of Ornithology* **156**, 115–124 (2015).

720   36.   S. F. Chenoweth, H. D. Rundle, M. W. Blows, EXPERIMENTAL EVIDENCE FOR THE EVOLUTION OF
721   INDIRECT GENETIC EFFECTS: CHANGES IN THE INTERACTION EFFECT COEFFICIENT, PSI (Ψ),
722   DUE TO SEXUAL SELECTION. *Evolution* **64**, 1849–1856 (2010).

723   37.   N. W. Bailey, M. Zuk, Socially flexible female choice differs among populations of the Pacific field cricket:
724   geographical variation in the interaction coefficient psi ( ). *Proceedings of the Royal Society B: Biological
725   Sciences* **279**, 3589–3596 (2012).

726   38.   R. E. Lenski, M. Travisano, Dynamics of adaptation and diversification: a 10,000-generation experiment with
727   bacterial populations. *Proceedings of the National Academy of Sciences of the United States of America* **91**,
728   6808–6814 (1994).

729   39.   P. M. P. Brakefield, Artificial selection and the development of ecologically relevant phenotypes. *Ecology* **84**,
730   1661–1671 (2003).

731   40.   R. C. Fuller, C. F. Baer, J. Travis, How and when selection experiments might actually be useful. *Integrative
732   and Comparative Biology* **45**, 391–404 (2005).

733   41.   T. J. Kawecki, *et al.*, Experimental evolution. *Trends in Ecology & Evolution* **27**, 547–60 (2012).

734   42.   J. B. Saltz, Genetic composition of social groups influences male aggressive behaviour and fitness in natural
735   genotypes of Drosophila melanogaster. *Proceedings of the Royal Society B: Biological Sciences* **280**, 31926
736   (2013).

737   43.   M. Hutchins, T. Douglas, L. Pollack, J. B. Saltz, Genetic Variation in Male Aggression Is Influenced by
738   Genotype of Prior Social Partners in *Drosophila melanogaster*. *The American Naturalist* **203**, 551–561
739   (2024).

740   44.   I. Camerlink, W. W. Ursinus, P. Bijma, B. Kemp, J. E. Bolhuis, Indirect Genetic Effects for Growth Rate in
741   Domestic Pigs Alter Aggressive and Manipulative Biting Behaviour. *Behav Genet* **45**, 117–126 (2015).

45. A. J. Wilson, U. Gelin, M.-C. Perron, D. Réale, Indirect genetic effects and the evolution of aggression in a
vertebrate system. *Proc. R. Soc. B.* **276**, 533–541 (2009).

46. S. P. Nilsen, Y.-B. Chan, R. Huber, E. A. Kravitz, Gender-selective patterns of aggressive behavior in
Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America*
101, 12342–7 (2004).

47. C. Rutte, M. Taborsky, M. W. G. Brinkhof, What sets the odds of winning and losing? *Trends in Ecology &*
Evolution **21**, 16–21 (2006).

48. A. Yurkovic, O. Wang, A. C. Basu, E. A. Kravitz, Learning and memory associated with aggression in
Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America*
103, 17519–24 (2006).

49. J. K. M. Penn, M. F. Zito, E. A. Kravitz, A single social defeat reduces aggression in a highly aggressive
strain of Drosophila. *Proceedings of the National Academy of Sciences of the United States of America* **107**,
12682–12686 (2010).

50. S. Trannoy, J. Penn, K. Lucey, D. Popovic, E. A. Kravitz, Short and long-lasting behavioral consequences of
agonistic encounters between male Drosophila melanogaster. *Proceedings of the National Academy of*
Sciences of the United States of America **113**, 4818–4823 (2016).

51. J. B. Saltz, Genetic variation in social environment construction influences the development of aggressive
behavior in Drosophila melanogaster. *Heredity (Edinb)* **118**, 340–347 (2017).

52. J. Altmann, Observational Study of Behavior: Sampling Methods. *Behav* **49**, 227–266 (1974).

53. E. Vrontou, S. P. Nilsen, E. Demir, E. A. Kravitz, B. J. Dickson, fruitless regulates aggression and dominance
in Drosophila. *Nature Neuroscience* **9**, 1469–1471 (2006).

54. J. A. Stamps, P. A. Biro, D. J. Mitchell, J. B. Saltz, Bayesian updating during development predicts genotypic
differences in plasticity: PREDICTING GENOTYPIC DIFFERENCES IN PLASTICITY. *Evolution* (2018).
https://doi.org/10.1111/evo.13585.

55. J. B. Saltz, S. Lymer, J. Gabrielian, S. V. Nuzhdin, Genetic Correlations among Developmental and
Contextual Behavioral Plasticity in Drosophila melanogaster. *Am. Nat.* **190**, 61–72 (2017).

56. D. S. Falconer, T. F. C. Mackay, *Introduction to quantitative genetics*, 4. ed., [16. print.] (Pearson, Prentice
Hall, 2009).

57. F. Santostefano, M. Moiron, A. Sánchez-Tójar, D. N. Fisher, Indirect genetic effects increase the heritable
variation available to selection and are largest for behaviors: a meta-analysis. *Evolution Letters* qrae051
(2024). https://doi.org/10.1093/evlett/qrae051.

58. A. Baud, S. McPeek, N. Chen, K. A. Hughes, Indirect Genetic Effects: A Cross-disciplinary Perspective on
Empirical Studies. *Journal of Heredity* esab059 (2021). https://doi.org/10.1093/jhered/esab059.

59. M. Murray, J. Wright, Y. G. Araya-Ajoy, Evolutionary rescue from climate change: male indirect genetic
effects on lay-dates and their consequences for population persistence. *Evolution Letters* **8**, 137–148 (2024).

60. D. N. Fisher, *et al.*, Anticipated effects of abiotic environmental change on intraspecific social interactions.
Biol Rev **96**, 2661–2693 (2021).

61. J. Akhund-Zade, *et al.*, "Wild flies hedge their thermal preference bets in response to seasonal fluctuations"
(Evolutionary Biology, 2020).

62. E. W. Wice, J. B. Saltz, Selection on heritable social network positions is context-dependent in Drosophila
melanogaster. *Nat Commun* **12**, 3357 (2021).

63. S. Chen, A. Y. Lee, N. M. Bowens, R. Huber, E. A. Kravitz, Fighting fruit flies: A model system for the study
of aggression. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 5664–5668 (2002).

64. S. N. Wood, *Generalized Additive Models: An Introduction with R*, 2nd Ed. (Chapman and Hall/CRC, 2017).

65. M. E. Brooks, *et al.*, glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated
Generalized Linear Mixed Modeling. *The R Journal* **9**, 378–400 (2017).

66. F. Hartig, DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models.
(2020). Deposited 2020.

67. J. Fox, S. Weisberg, *An {R} companion to applied regression, second edition*, Second (Sage, 2011).

68. P. De Villemereuil, H. Schielzeth, S. Nakagawa, M. Morrissey, General Methods for Evolutionary
Quantitative Genetic Inference from Generalized Mixed Models. *Genetics* **204**, 1281–1294 (2016).

69. R. Lenth, emmeans: Estimated Marginal Means, aka Least-Squares Means. (2019). Deposited 2019.

70. M. Sóskuthy, Generalised additive mixed models for dynamic analysis in linguistics: a practical introduction.
[Preprint] (2017). Available at: https://arxiv.org/abs/1703.05339 [Accessed 24 July 2024].

Supplementary Materials

Supplemental Tables S1 to S4

Data files

Readme for data files

R Code

SUPPLEMENTAL TABLES
Table of contents

813 Table S1. Number of males measured for aggression every generation for every population. In cases where less than
814 100 were measured, we matched the demographic histories of all populations in that replicate. See Materials and
815 Methods for details.

Replicate		Treatment	Generation	n.males
	1	C	0	98
	1	N	0	100
	1	P	0	99

1	Z	0	100
1	N	1	95
1	P	1	100
1	Z	1	99
1	N	2	101
1	P	2	139
1	Z	2	142
1	N	3	105
1	P	3	90
1	Z	3	105
1	N	4	65
1	P	4	62
1	Z	4	67
1	N	5	97
1	P	5	100
1	Z	5	98
1	N	6	99
1	P	6	100
1	Z	6	100
1	N	7	95
1	P	7	100
1	Z	7	99
1	N	8	96
1	P	8	100
1	Z	8	99
1	N	9	90
1	P	9	100
1	Z	9	99
1	N	10	98
1	P	10	100
1	Z	10	100
1	N	11	98
1	P	11	94
1	Z	11	97
1	C	12	87
1	N	12	99
1	P	12	100
1	Z	12	99
1	N	13	98
1	P	13	100
1	Z	13	99

1	N	14	95
1	P	14	100
1	Z	14	99
1	N	15	93
1	P	15	100
1	Z	15	98
1	N	16	93
1	P	16	100
1	Z	16	98
1	N	17	127
1	P	17	132
1	Z	17	130
1	N	18	67
1	P	18	68
1	Z	18	66
1	N	19	97
1	P	19	100
1	Z	19	84
1	N	20	100
1	P	20	100
1	Z	20	100
1	N	21	99
1	P	21	100
1	Z	21	100
1	N	22	97
1	P	22	100
1	Z	22	100
1	N	23	97
1	P	23	97
1	Z	23	100
1	N	24	96
1	P	24	100
1	Z	24	97
1	C	25	70
1	N	25	72
1	P	25	122
1	Z	25	72
2	C	0	95
2	N	0	96
2	P	0	94
2	Z	0	96

2	N	1	98
2	P	1	100
2	Z	1	100
2	N	2	97
2	P	2	100
2	Z	2	100
2	N	3	95
2	P	3	100
2	Z	3	100
2	N	4	96
2	P	4	100
2	Z	4	100
2	N	5	96
2	P	5	100
2	Z	5	99
2	N	6	100
2	P	6	100
2	Z	6	100
2	N	7	96
2	P	7	100
2	Z	7	100
2	N	8	96
2	P	8	100
2	Z	8	100
2	N	9	91
2	P	9	100
2	Z	9	96
2	N	10	95
2	P	10	98
2	Z	10	99
2	N	11	98
2	P	11	100
2	Z	11	98
2	C	12	41
2	N	12	98
2	P	12	100
2	Z	12	99
2	N	13	91
2	P	13	97
2	Z	13	98
2	N	14	93

2	P	14	100
2	Z	14	100
2	N	15	97
2	P	15	100
2	Z	15	98
2	N	16	98
2	P	16	100
2	Z	16	100
2	N	17	100
2	P	17	100
2	Z	17	100
2	N	18	98
2	P	18	100
2	Z	18	100
2	N	19	99
2	P	19	100
2	Z	19	99
2	N	20	98
2	P	20	100
2	Z	20	98
2	N	21	97
2	P	21	100
2	Z	21	95
2	N	22	98
2	P	22	100
2	Z	22	99
2	N	23	99
2	P	23	100
2	Z	23	100
2	N	24	79
2	P	24	100
2	Z	24	100
2	C	25	124
2	N	25	126
2	P	25	131
2	Z	25	130
3	C	0	100
3	N	0	99
3	P	0	94
3	Z	0	99
3	N	1	96

3	P	1	98
3	Z	1	100
3	N	2	94
3	P	2	100
3	Z	2	100
3	N	3	99
3	P	3	100
3	Z	3	99
3	N	4	97
3	P	4	100
3	Z	4	99
3	N	5	94
3	P	5	100
3	Z	5	96
3	N	6	92
3	P	6	100
3	Z	6	99
3	N	7	93
3	P	7	98
3	Z	7	97
3	N	8	97
3	P	8	90
3	Z	8	98
3	N	9	96
3	P	9	100
3	Z	9	100
3	N	10	96
3	P	10	100
3	Z	10	99
3	N	11	97
3	P	11	98
3	Z	11	100
3	N	12	96
3	P	12	98
3	Z	12	99
3	C	13	95
3	N	13	91
3	P	13	94
3	Z	13	97
3	N	14	97
3	P	14	100

3	Z	14	98
3	N	15	0
3	P	15	0
3	Z	15	0
3	N	16	93
3	P	16	100
3	Z	16	99
3	N	17	96
3	P	17	100
3	Z	17	92
3	N	18	97
3	P	18	98
3	Z	18	99
3	N	19	85
3	P	19	100
3	Z	19	97
3	N	20	75
3	P	20	78
3	Z	20	92
3	N	21	86
3	P	21	96
3	Z	21	93
3	N	22	0
3	P	22	0
3	Z	22	0
3	N	23	0
3	P	23	0
3	Z	23	0
3	N	24	66
3	P	24	85
3	Z	24	73
3	N	25	0
3	P	25	0
3	Z	25	0
3	C	26	66
3	N	26	70
3	P	26	125
3	Z	26	68

817
818
819
820   Table S2. Full model structures for main and supplemental models

model name	question	generations included	treatments included	response variable	fixed effects	random effects	error distribution	model type
model 1	how did aggression evolve?	0 to 24	C, 0, N, P	total aggression	Generation*Treatment+focal.stuck	Replicate + Date + observer identities	negative binomial	GLMM
model 2	how did each component of aggression evolve?	1 to 24	0, N, P	each component of aggressive behavior	Generation*Treatment*component+ focal.stuck	Replicate + Date + Individual.ID + observer identities	negative binomial	GLMM
model 3	did prior experience effects evolve?	1 to 24	N	change in aggression: total aggression on day 2 - total aggression on day 1	Generation + day1 opponent aggression*Generation + day2 opponent aggression*Generation + focal.stuck	Replicate + Date + observer identities	scaled *t*	GAMM -- day1 and day2 opponent aggression and their interactions fit as thin plate splines
model 4	did prior experience effects differ across treatments at the end of the experiment?	25	C, 0, N, P	change in aggression: total aggression on day 2 - total aggression on day 1	Treatment + day1 opponent aggression*Treatment + day2 opponent aggression*Treatment + focal.stuck	Replicate + Date	negative binomial	GAMM -- opponent aggression and its interactions fit as thin plate splines
model S1	were populations the same at the beginning?	0	C, 0, N, P	each component of aggressive behavior	Treatment*component + focal.stuck	Replicate + Date + Individual.ID + observer identities	negative binomial	GLMM
model S2.0, S2.N, S2.P	did retaliation evolve?	1 to 24	0, N, P in separate models	total aggression	total aggression ~ opponent aggression*Generation + focal.stuck	Replicate + Date + observer identities	negative binomial	GAMM -- opponent aggression and its interactions fit as thin plate splines

Table S3. Model S1 Results.

model S1: were populations the same at the beginning?				
term	parameter estimate (se)	Chisq	Df	p
Treatment	.	0.1267	3	0.98846
focal.stuck	-0.13 (0.078)	2.8529	1	0.09121

Table S4. Results from Models S2.0, S2.N, and S2.P.

model	results				
Model 2.N	non-linear term	edf (measure of "wiggliness")	chisq	reference df	p
	opponent aggression	3.429	442.508	4.128	< 2e-16
	Generation: opponent aggression	1.013	0.122	1.026	0.739
Model 2.P	non-linear term	edf (measure of "wiggliness")	chisq	reference df	p
	opponent aggression	6.298	785.013	7.115	< 2e-16
	Generation: opponent aggression	2.117	4.845	2.577	0.11528

	non-linear term	edf (measure of "wiggliness")	chisq	reference df	p
Model 2.0	**opponent aggression**	**1**	**207.259**	**1**	**<2e-16**
	Generation: opponent aggression	1.152	3.458	1.284	0.0682