An alveolus lung-on-a-chip model of *Mycobacterium fortuitum* lung infection

Victoria Ektnitphong[1], Beatriz R.S. Dias[1], Priscila C. Campos[1] and Michael U. Shiloh[1,2]

[1]Department of Internal Medicine and [2]Department of Microbiology

University of Texas Southwestern Medical Center

5323 Harry Hines Blvd., Y9.308

Dallas, TX 75390-9113

Corresponding author: Michael.shiloh@utsouthwestern.edu

ABSTRACT

Lung disease due to non-tuberculous mycobacteria (NTM) is rising in incidence. While both two

dimensional cell culture and animal models exist for NTM infections, a major knowledge gap is

the early responses of human alveolar and innate immune cells to NTM within the human alveolar

microenvironment. Here we describe development of a humanized, three-dimensional, alveolus

lung-on-a-chip (ALoC) model of *Mycobacterium fortuitum* lung infection that incorporates only

primary human cells such as pulmonary vascular endothelial cells in a vascular channel, and type

I and II alveolar cells and monocyte-derived macrophages in an alveolar channel along an air-

liquid interface. *M. fortuitum* introduced into the alveolar channel primarily infected macrophages,

with rare bacteria inside alveolar cells. Bulk-RNA sequencing of infected chips revealed marked

upregulation of transcripts for cytokines, chemokines and secreted protease inhibitors

(SERPINs). Our results demonstrate how a humanized ALoC system can identify critical early

immune and epithelial responses to *M. fortuitum* infection. We envision potential application of

the ALoC to other NTM and for studies of new antibiotics.

INTRODUCTION

Nontuberculous mycobacteria (NTM) represent a diverse group of environmental organisms

found in varied natural habitats such as soil, water, and vegetation (Mercaldo et al., 2023).

Traditionally considered opportunistic pathogens of people with structural or immune deficits, lung

disease secondary to NTM is increasing in incidence in women and people over 65 years of age

(Winthrop et al., 2020). Among the NTM, *M. fortuitum* is the second most frequently encountered

rapidly-growing NTM after *Mycobacterium abscessus*, and can cause a spectrum of infections

including pulmonary, skin and soft tissue, and disseminated infections (Roquet-Baneres et al.,

2023, Kim et al., 2023). NTM lung infections are more frequent and more severe in

immunocompromised individuals and individuals with pre-existing conditions such as

bronchiectasis, severe chronic obstructive pulmonary disease (COPD), cystic fibrosis, organ

transplant and cancer (Roquet-Baneres et al., 2023, Kim et al., 2023). Treatment of NTM lung

disease can be complex due to the presence of antibiotic resistance, medication side effects and

extended duration of drug treatment. More efforts are needed to develop new, safe and efficacious

therapeutics and to shorten treatment duration (Johansen and Kremer, 2020, Orme and Ordway,

2014, Kremer, 2020). In addition, the rising incidence of NTM lung infections among individuals

without defined predisposing conditions indicates that a greater understanding of the immune

response to NTM lung infection is needed.

Expanding the scope of knowledge of the dynamic interactions between humans and NTM is

essential for developing effective prevention and treatment strategies for NTM lung disease, but

gaining such knowledge has been hindered by the lack of suitable *in vitro* and *in vivo* models of

NTM lung infection (Baldwin et al., 2019). Current *in vitro* models typically use traditional two-

dimensional cell culture methods with primary human CD14[+] peripheral blood-derived

macrophages or immortalized monocytic cell lines THP-1 and U937 (Kilinc et al., 2022), and

alveolar epithelial cell lines such as A549 (Rampacci et al., 2020). Although immortalized cell lines

allow for large-scale screening and assay reproducibility, a major disadvantage is that they may

not completely replicate myeloid cell responses in humans, especially in the unique alveolar

microenvironment. For example, the use of PMA to differentiate THP-1 and U937 cells into

adherent macrophage-like cells alters monocytic surface markers, transcriptional activities, and

cytokine production (Rampacci et al., 2020). Animal models have also been established to study

host immune responses to NTM and for testing potential antimicrobial compounds and vaccines.

Most animal models of NTM lung infection use mice as the host species including for *M.*

abscessus (Caverly et al., 2015, De Groote et al., 2014) and *Mycobacterium avium* (Gonzalez-

Perez et al., 2013, Gangadharam, 1995). However, the variable virulence of NTM in model

organisms makes establishing consistent animal models challenging. For example, *M. avium* can

establish a productive lung infection in immunocompetent mice, whereas *M. abscessus* is typically

rapidly cleared and requires immunocompromised mice to achieve a similar outcome (Chan et

al., 2016). Animal models for other NTM species such as *M. fortuitum* are lacking, delaying

progress towards understanding shared and unique molecular mechanisms of pathogenicity for

various NTM. Thus, developing *in vitro* models that can recapitulate the human lung

microenvironment using primary human cells has the potential to advance the study of pulmonary

NTM infections, and is a critical step towards developing improved prevention, vaccine and

treatment strategies.

The alveolus lung-on-a-chip (ALoC) is an innovative microfluid device composed of

microfabricated channels and chambers that mimic the architecture and cellular composition of

the lung alveolus, the initial site of interaction for pulmonary NTM infection (Bethencourt et al.,

2024) . The "chip" contains two overlapping channels, parallel to each other, that are separated

by a porous membrane. It features alveolar epithelial cells lining one channel and pulmonary

microvascular endothelial cells (HMVECs) lining the other to closely replicate the alveolar-

capillary interface of the human lung (Dasgupta et al., 2023). Because the cell types grow in

distinct channels, media or air flow through the channels can be independently controlled to create

distinct environments for the cells. Additional advantages of the ALoC include the ability to

generate an air-liquid interphase (ALI) that is hallmark of the alveolus and application of

mechanical stretch that is essential for the proper differentiation of alveolar epithelial cells into

type I (AT1) and type II (AT2) pneumocytes. In that respect, the ALoC helps overcome one of the

major obstacles of *in vitro* studies of alveolar epithelial cells, namely, simultaneously differentiating

both AT1 and AT2 cells in culture by leveraging stretch and growth factor addition (Li et al., 2018).

Thus, compared to traditional tissue culture methods, the ALoC system provides a more

physiologically relevant environment to study cells and pathogens of the human lung and has

been applied to several airway diseases and pathogens including influenza A virus (Bai et al.,

2022, Haiqing Bai et al., 2022), *Mycobacterium tuberculosis* (Thacker et al., 2020), and acute

radiation induced lung injury (RILI) (Dasgupta et al., 2023).

The complex interaction between *M. fortuitum* and the human alveolar epithelial and immune cells

remains poorly understood. Here, we develop and use a fully humanized ALoC model as a

platform to study the host-pathogen interactions of *M. fortuitum* within the alveolar

microenvironment. By recapitulating key features of the alveolar interface, including cellular

composition, mechanical forces, and biochemical gradients, this model offers a unique

opportunity to investigate the dynamics of *M. fortuitum* infection in a controlled and reproducible

manner. Through integration of advanced imaging techniques and transcriptomic analyses, we

identify macrophages as the primary cell type infected in the alveolus and upregulation of several

inflammatory pathways after *M. fortuitum* infection of the humanized ALoC. We propose

application of the humanized alveolus lung-on-a-chip as a new model to study NTM infections.

RESULTS

Two types of lung chips have been developed to date, one that approximates a bronchial airway

(Benam et al., 2015, Benam et al., 2016) and another more like an alveolus (**Figure 1A**) (Huh et

al., 2012a, Huh et al., 2010, Huh et al., 2012b). Since early NTM infection occurs in the alveolus,

we chose the alveolus lung-on-a-chip (ALoC) as our working model for NTM infection (**Figure**

1B), as previously demonstrated with *E. coli* (Huh et al., 2010). We initially decided to use *M.*

fortuitum to model airway NTM infection using the ALoC model both because it is rapidly-growing,

with a doubling time of 2-3 hours (Qin et al., 1999), and because amongst the rapid growers it

has one of the highest age-adjusted incidence of disease (Donohue, 2021).

Constructing a functional ALoC requires meticulously following a stepwise protocol

(**Supplemental Figure 1A**). Briefly, the membranes of individual chips are chemically activated,

coated with extracellular matrix and then human alveolar epithelial cells are added to the air (or

apical) channel. After three days, human primary lung microvascular endothelial cells are

introduced to the vascular (or basal) channel. Once cells are confluent (~1 day), the alveolus-on-

a-chip is connected to regulated media flow (30 uL/hr) in both channels. The next day, the apical

channel media is removed, generating an air-liquid interface (ALI), and after 2 days of ALI, stretch

is introduced to mimic breathing (5% stretch intensity, 0.2 Hz) and induce surfactant production

(Thacker et al., 2020). After two days of stretch we add differentiated human CD14[+] peripheral

blood-derived macrophages (hereafter called 'macrophages') from anonymous donors to the top

channel (**Figure 1C**). To assess the donor compatibility of the alveolar epithelial cells and

macrophages in the apical channel, we used immunofluorescence to compare AT1 and AT2 cell

differentiation, cell morphology and distribution of cells along the apical channel of ALoCs with

and without macrophages. In ALoCs without macrophages, a monolayer of both AT1 and AT2 cell

types was distributed equally across the apical channel (**Figure 1D, Supplemental Figure 1B**).

High resolution images of ALoCs without macrophages highlighted distinct AT1 and AT2 cell types

after 15 days in culture on the ALoC (**Figure 1E**). Three-dimensional reconstruction of the ALoC

contrasted the cuboidal, cobblestone morphology of AT2 cells to the flat, squamous morphology

of AT1 cells (**Figure 1F**). Cross sectional imaging of the airway channel further highlighted the

nuclei in the basal region of the apical channel with interspersed AT1 and AT2 cells (**Figure 1G**).

In ALoCs with macrophages, we also observed a monolayer of differentiated AT1 and AT2 cells

with seeded macrophages residing on top of the alveolar epithelial monolayer (**Figure 1H-J**).

Since macrophages were added 2 days after 100% confluency of the apical channel, it is expected

that they would be found loosely attached to the alveolar epithelial layer rather than within the

alveolar epithelial layer. Although we introduced the same number of macrophages as alveolar

epithelial cells, we observed many fewer macrophages overall, suggesting that the majority failed

to adhere to the epithelial layer and were washed away or clumped together (**Supplemental**

Figure 1C). Stitched images of ALoCs containing macrophages showed that, despite losing many

macrophages, the remaining macrophages were sufficient to evenly distribute across the entire

apical channel (**Supplemental Figure 1D**). We also observed that despite the addition of

macrophages, AT cell distribution was comparable to uninfected ALoCs without macrophages. At

higher magnification, we observed rounded macrophages dispersed on top of the cuboidal

shaped AT2 cells (**Figure 1I**). Three-dimensional reconstruction of the apical channel also

demonstrated the same cuboidal morphology of AT2 cells and flat, squamous AT1 cells that we

observed in ALoCs without macrophages (**Figure 1J**). Cross-sectional imaging showed

macrophages residing predominantly on top of the alveolar epithelial monolayer (**Figure 1K**).

From these experimental observations, namely that AT1 and AT2 cell differentiation was

maintained after the addition of primary human CD14^{+} peripheral blood macrophages, we

concluded that macrophages could safely be added to the human ALoC model without

dramatically altering epithelial biology.

After establishing the ALoC model with human macrophages, we next incorporated mCherry-

expressing *M. fortuitum* to the system. We introduced mCherry-expressing *M. fortuitum* at an

estimated multiplicity of infection (MOI) of ~1 for the approximate number of AT1 and AT2 cells on

the chip (~3.0 x 10^{4}) (**Figure 2A**). Because we used immunofluorescence to monitor different cell

types on the ALoC, uninfected ALoCs from the same experiment were used to confirm AT1 and

AT2 differentiation (**Supplemental Figure 2A**) and we observed similar trends of AT1 and AT2

differentiation. When ALoCs without macrophages were infected with *M. fortuitum*, we observed

appropriate AT2 cell distribution across the chip in addition to large clumps of bacteria scattered

across the ALoC (**Figure 2B**). Along the apical channel in *M. fortuitum*-infected ALoC, we

observed areas that had little to no immunofluorescence signal throughout the z-stack that we did

not observe in uninfected ALoCs with or without macrophages. In the areas on the ALoCs

containing "dark patches" lacking AT1 or AT2 immunofluorescence, we could not determine at

lower resolution if those areas were due to cell death, but higher resolution images of ALoCs

without macrophages after 24 hours of mCherry *M. fortuitum* infection suggested loss of viability

of AT1 and AT2 cells in areas adjacent to bacteria (**Figure 2C**). Three-dimensional reconstruction

of *M. fortuitum*-infected ALoCs without macrophages demonstrated many AT2 cells with a

cuboidal morphology but interestingly we also observed a more "flattened" AT2 morphology

(**Figure 2D**). Whether these are AT1 cells with HT2 signal or AT2 cells with flattened morphology

could not be determined. HT1 labeling was very diffuse in *M. fortuitum*-infected ALoCs and we

also observed co-staining of HT1 and HT2 in both alveolar epithelial cell types in *M. fortuitum*-

infected ALoCs without macrophages compared to uninfected ALoCs without macrophages

(**Supplemental Figure 2B-D**). High-resolution cross-sectional analysis bacterial clumps on the

M. fortuitum-infected ALoC without macrophages demonstrated bacteria interspersed amongst

AT1 and AT2 cells (**Figure 2E**). No *M. fortuitum* cord-like structures were observed within the

initial 24 hours of infection. Taken together, introducing *M. fortuitum* to the ALoC model at an

approximate MOI of 1 generated a productive infection without overwhelming the alveolar

channel.

We next wanted to determine how *M. fortuitum* grows on ALoC in the presence of human

macrophages. Twenty-four hours after seeding human macrophages onto the ALoC, we infected

ALoCs with macrophages with *M. fortuitum* as described above, using the same donors for

alveolar epithelial and pulmonary microvascular endothelial cell as for ALoCs without

macrophages (**Figure 3A**). On ALoCs infected with *M. fortuitum* 24 hours after macrophage

seeding, there was suitable distribution of AT2 cells across the chips (comparable to infected

ALoCs without macrophages). In contrast to ALoC lacking macrophages infected with *M.*

fortuitum, inclusion of macrophages resulted in most bacteria identified intracellularly within

macrophages with no apparent bacterial clumps (**Figure 3B**). The AT2 cell morphology appeared

more rounded in shape in areas adjacent to infected macrophages and AT2 cells remained intact

with no areas of apparent cell death as we observed in *M. fortuitum*-infected ALoCs without

macrophages (**Figure 3C**). Similarly to *M. fortuitum*-infected ALoCs without macrophages, we

again did not observe *M. fortuitum* cord-like structures. Of note, we were unable to image AT1

cells directly in infected ALoCs with macrophages due to limitations in the number of fluorophores

available in one experiment. Three-dimensional reconstruction highlighted the expected cuboidal

morphology of AT2 cells and demonstrated macrophage clusters near the chip edges (**Figure**

3D). We observed less "flattening" of the AT2 cells in ALoCs with macrophages but were not able

to directly assess AT1 cells in the infected ALoCs with macrophages to determine if they

demonstrated the same trends of AT1 and AT2 co-staining and AT1 signal diffusion. Cross-

sectional analysis of the ALoC with macrophages also highlighted that bacteria were intracellular

in macrophages and not residing on top of the macrophages (**Figure 3E**). In addition, by removing

the AT2 signal, we observed *M. fortuitum* within macrophages in a three-dimensional

reconstruction of the ALoC (**Figure 3F**). To quantify bacteria within either AT2 or CD14$^+$ cells, we

analyzed 10 z-stacks from 3 independent ALoCs containing macrophages. We used three-

dimensional reconstruction and ortho-projections to assign bacteria within AT2 cells (HT2$^+$) or

macrophages (CD14$^+$) and determined that bacteria were found more often in macrophages than

AT2 cells (**Figure 3G**). Taken together, we conclude that *M. fortuitum* infection of ALoC containing

macrophages results in a productive infection, and that ingestion of bacteria by macrophages may

protect adjacent epithelia from infection-induced cell death.

To determine the alveolar response to *M. fortuitum* infection using the human ALoC, we

determined the impact of *M. fortuitum* infection on ALoC with macrophages compared to

uninfected ALoC with macrophages using bulk RNA sequencing. After infection at an MOI of 1 for

24 hours, we collected total RNA from the apical channel containing AT1 cells, AT2 cells and

macrophages for sequencing. While there was some heterogeneity in the response, as would be

expected in a primary infection, we observed 1454 genes were more than 2-fold upregulated,

while 588 were downregulated more than 2-fold (**Figure 4A, B**). Amongst the upregulated genes,

we observed significant upregulation of key cytokines such as tumor necrosis factor (*TNF*),

granulocyte-macrophage colony-stimulating factor (*GM-CSF* or *CSF2*), macrophage colony-

stimulating factor (*M-CSF* or *CSF3*), interleukin 1A (*IL1A*), interleukin 1B (*IL1B*), interleukin 6 (*IL6*),

and interleukin 8 (*IL8*), along with the alarmin, calprotectin (*S100A8* and *S100A9*) (**Figure 4C**).

We also observed several chemokines upregulated including the chemokine (C-X-C motif) ligand

family members *CXCL1, CXCL2, CXCL3, CXCL5, CXCL6, CXCL10* and *CXCL11* and the C-C

motif chemokine ligand family members *CCL2, CCL5, CCL20* and *CCL28* (**Figure 4C**). Finally,

we also noted that many secreted serine protease inhibitor (SERPIN) genes were also

upregulated by *M. fortuitum* infection (**Figure 4D**). Using pathway analysis (Ge et al., 2018), we

found that the most significantly upregulated processes were involved in host defense including

"response to bacterium", "defense response to bacterium", "antimicrobial humoral response",

"response to molecule of bacterial origin" and "response to lipopolysaccharide" (**Table 1**).

Likewise, upregulated molecular functions included several annotated as involved in "signaling"

in addition to "cytokine" and various channel functions (**Table 2**). Taken together, our data highlight

the acute inflammatory response of humanized ALoC containing human macrophages to *M.*

fortuitum infection.

DISCUSSION

We report the development of a fully humanized ALoC model for alveolar NTM infection. This

model develops features consistent with initial primary alveolar infection, including infection of

both airway macrophages and occasional alveolar cells. Moreover, bulk RNA sequencing reveals

activation of an acute inflammatory signaling cascade, including induction of cytokines,

chemokines and alarmins like serpins and calprotectin, with potential roles in recruitment of

myeloid and lymphoid cells from the vasculature and in triggering a robust immune response to

restrict bacterial growth. These findings highlight the value of using the humanized lung on a chip

to model a primarily human infectious disease.

Prior studies using the alveolus-on-a-chip have included infection models, such as infection with

E. coli (Huh et al., 2010), *M. tuberculosis* (Thacker et al., 2020), *Aspergillus fumigatus* (Hoang et

al., 2022), *Staphylococcus aureus* (Bai et al., 2022, Deinhardt-Emmer et al., 2020), and influenza

virus (Bai et al., 2022, Deinhardt-Emmer et al., 2020). The alveolus-on-a-chip has also been used

to determine the effect of mechanical strain on inflammation and particle uptake (Huh et al., 2010),

pulmonary thrombosis (Jain et al., 2018), and mechanotransduction and pulmonary edema (Li et

al., 2019). In the context of *M. tuberculosis* and *A. fumigatus* infection models, addition of human

monocytes/macrophages impacts pathogenesis (Thacker et al., 2020, Hoang et al., 2022). With

M. tuberculosis, cord-like structures are observed compressing macrophage nuclei (Mishra et al.,

2023). We did not observe a similar phenomenon with *M. fortuitum* infection of human

macrophages, which might be due to differences in the cell walls of the organisms or other factors

that impact the growth characteristics of *M. tuberculosis* (slow) as compared to *M. fortuitum*

(rapid). Alternatively, differences in the time after infection used for analysis (24 hours in this study

versus 72 hours (Mishra et al., 2023)) or imaging techniques may also account for these

differences.

We identified a transcriptional signature of early human alveolar infection with *M. fortuitum*. Some

notable findings include marked upregulation of important cytokines and chemokines that may be

involved in stimulating an early innate immune response to airway infection. For example, both

IL6 and IL8 are key cytokines involved in neutrophil recruitment to the airway (Fu and Harrison,

2021), and such recruited cells could have both beneficial (contributing to NTM killing) and

detrimental (involvement in tissue damage and bronchiectasis) effects (Alkarni et al., 2023).

While there have been previous studies of human peripheral blood transcriptional responses in

the context of pulmonary NTM infection (Wang et al., 2024, Cowman et al., 2018, Lindestam

Arlehamn et al., 2022) or tuberculosis (Tabone et al., 2021, Berry et al., 2010), as well as efforts

to determine transcriptional responses in human lung tissue from tuberculosis patients (Wang et

al., 2023a), to our knowledge this is the first analysis of the early interaction of an NTM with a

humanized ALoC. Interestingly, in the RNA sequencing analysis of PBMCs from individuals

infected with *M. avium*, TNF signaling was enriched, and individual genes such as those for

SERPINA1 and calprotectin were modestly upregulated (Lindestam Arlehamn et al., 2022).

Likewise, in a proteomics analysis of serum exosomes in humans with NTM infection, a variety of

serpins including SERPINA1 and SERPINA5 were enriched in patients with *M. abscessus* or *M.*

avium infection (Wang et al., 2023b). Recent work has found the induction of cytosolic SERPINs

may protect *M. marinum* infected macrophages from cathepsin B mediated lysosomal rupture and

cell death (Nobs et al., 2024), a process called lysoptosis that is evolutionary conserved from *C.*

elegans to mammals (Luke et al., 2022). Thus, SERPIN induction by *M. fortuitum* infection,

potentially within infected macrophages, may act to protect infected cells from lysoptosis. Of note,

because we used bulk RNA sequencing, we were unable to identify the precise cellular source of

transcriptional changes identified in our analysis. Future single cell sequencing studies will be

necessary to determine the responses of individual cell types within the model system, and how

such responses impact intercellular communication. Taken together, our bulk RNAseq analysis

has identified unique features of early alveolar NTM infection.

There are several limitations to our model. First, because our model uses all primary human cells,

genetic manipulation of the various cellular constituents is impractical, though primary human

monocytes can be transduced with lentivirus to abrogate gene expression (Campbell and Spector,

2012, Franco et al., 2017). Alternatively, use of genetically modified human or mouse cell lines

can overcome this limitation, though immortalized cell lines have their own inherent drawbacks.

Second, because primary cells, whether pulmonary epithelial cells, pulmonary vascular cells or

PBMCs, are obtained from unique individuals, genetic variation generates greater complexity.

Finally, though our model does allow multiple cell types (epithelia, endothelia and myeloid cells)

to interact with each other and the NTM in the alveolar microenvironment, the interactions are

limited to the cells added to the chamber, and thus do not include all the possible myeloid and

lymphoid cells that could be recruited to the airway in the setting of pulmonary mycobacterial

disease, such as neutrophils (Kimmey et al., 2015), eosinophils (Bohrer et al., 2021) or innate

lymphoid cells (Ardain et al., 2019).

In conclusion, we have developed a fully humanized alveolus on a chip model of pulmonary NTM

infection. By capturing the complexity of NTM in three dimensions and with multiple cell types, as

would be found in a human lung, this model creates an opportunity to better characterize the

cellular and molecular mechanisms that mediate the outcome of human airway NTM infections.

Because NTM infections are on the rise, particularly in individuals with cystic fibrosis,

bronchiectasis and people with HIV, it may be useful to apply this technology to determine the

relative impact of genetic risk alleles (i.e. for those with cystic fibrosis) or coinfection with HIV on

the alveolar transcriptional responses or bacterial survival in the alveolus over time. Furthermore,

by leveraging this system to directly compare responses to other common human pulmonary NTM

infections, such as *M. avium* (Cristancho-Rojas et al., 2023, Varley and Winthrop, 2022, Donohue,

2021), *M. kansasii* and *M. abscessus* (Toure et al., 2023, Lagune et al., 2023, Cristancho-Rojas

et al., 2023, Ordway et al., 2008, Klever et al., 2023) future studies could identify shared and

unique survival strategies used by each organism, and establish distinct immune pathways

responsive to each pathogen which could function to limit or exacerbate infection, or be used

clinically as biomarkers of disease. Finally, as new drugs are developed to treat pulmonary NTM

infection, we envision applying this model for the simultaneous assessment of efficacy (i.e.

reduction in bacterial load) and cellular toxicity via infusion of test compounds into the vascular

channel.

Materials and Methods

Bacteria

Mycobacterium fortuitum subsp. *fortuitum* strain was obtained from ATCC (strain TMC 1529). We

transformed *M. fortuitum* with a vector containing *mCherry* driven by the GroEL constitutive

promoter.

Primary Cell Culture

Primary human alveolar epithelial cells (ATs) were obtained from Cell Biologics (H-6053). Prior to

seeding on Alveolus Lung-on-Chips (ALoCs), ATs were expanded *in vitro* in a T25 flask coated

with a 1% gelatin-based coating solution (Cell Biologics, catalog #6950) and complete medium

comprising of base media and supplements (Lonza, CC-3118). Medium was prepared according

to manufacturer's instructions using all supplements except GA-1000 where 1% Pen-Strep

solution (Gibco 15140-122) was added instead along with 5% FBS and hereby called Small

Airway Growth Medium (SAGM). Primary human microvascular endothelial cells were purchased

from Lonza (CC-2527). Human Lung Microvascular Endothelial cells (HMVECs) were expanded

in a T75 in complete medium comprising of base media and supplements (Lonza, CC-3202).

Medium was prepared according to manufacturer's instructions using all supplements except GA-

1000 where 1% Pen-Strep solution was used instead along with 5% FBS and hereby called EGM-

2MV. Both primary cell lines were cultured at 37°C in 5% CO_2 until ~80% confluency before

detachment with TrypLE Express (Gibco 12604013) and use on the ALoC.

One week prior to seeding the macrophages on the ALoC, peripheral blood mononuclear cells

(PBMCs) were obtained from buffy coat purchased from anonymous donors (Carter BloodCare).

PBMCs were isolated using Ficoll (Cytivia, 17144003) and SepMate50 tubes from StemCell

Technologies (85450). $CD14^+$ monocytes were positively selected from the PBMCs using CD14

Microbeads (Miltenyi, 130-050-201), and seeded onto uncoated petri dishes. The monocytes

were cultured overnight in RPMI medium (Gibco 11875-093) supplemented with 10% heat-

inactivated human serum obtained from the buffy coat of the donor, 1% HEPES buffered solution

(Lonza CC-5022), 1% sodium pyruvate (Gibco 11360-070), and 50 ng/mL human granulocyte-

macrophage colony-stimulating factor (GM-CSF) (Peprotech 300-03-100UG) and is hereby called

macrophage medium. The next day, the serum in the medium was changed to 10% FBS for the

remainder of the culturing. $CD14^+$ monocytes were differentiated to human monocyte-derived

macrophages (HMDMs) for 7 days at 37°C in 5% CO_2 until use on the ALoCs. GM-CSF was

added to medium for the first 4 days to differentiate the $CD14^+$ monocytes.

Human Alveolus LoC Model

ALoCs fabricated with polydimethylsiloxane (PDMS) were purchased from Emulate. The chips

were activated using a 0.5 mg/mL solution comprised of ER-1 and ER-2 (Emulate) that was

protected from light during the activation process. Working in a dark biosafety cabinet (BSC), both

the top and bottom channel were filled completely with ER-1/ER-2 and placed under a UV light

for 10 min, inspected for bubbles with a brightfield microscope and placed under the UV light for

an additional 10 min. After activation of all chips, the top and bottom channels were coated with

an extracellular matrix (ECM) solution specific to the cell type. The top channel (ATs) was coated

with an ECM solution containing Collagen IV at 200 µg/mL (Sigma C5533), Fibronectin at 30

µg/mL (Millipore Sigma F2006) and Laminin at 5µg/mL (Sigma L6274). The bottom channel

(endothelial cells) was coated with an ECM solution containing Collagen IV at 200 µg/mL and

Fibronectin at 30 µg/mL. All ECM components and solutions were kept on ice and prepared by

manufacturer's recommendations. With an empty 200 µL pipette tip plugged into each outlet port,

100 uL of ECM solution were added to the respective channels by expelling ECM solution from

pipette tip and plugging the inlet ports with the tip. The ECM-coated ALoCs were incubated at 4°C

overnight and the next day each channel was washed twice with 200 µL of respective medium.

Respective mediums were left in each channel prior to cell seeding and ALoCs were stored at

4°C until cell seeding.

ATs were seeded first in the top channel of all ALoCs. After detaching the cells from the T75 flasks,

they were adjusted to a concentration of 1×10^6 cells/mL in SAGM medium described above. The

bottom channels were filled with SAGM medium during epithelial cell seeding. 50 µL of the AT cell

suspension was pipetted into the top channel rapidly and checked under a microscope to ensure

correct seeding density and cell homogeneity. Of note, the top channel volume is ~28 µL total,

but to ensure no bubbles are introduced, the channels are overfilled and the flow-through gently

aspirated. Once all chips were seeded, they were placed in a chip cradle (Emulate) and incubated

at 37°C for at least 2 hours. After confirming all cells had attached, each top and bottom channel

were gently washed with 200 µL of warm SAGM medium and incubated overnight at 37°C.

HMVECs were seeded 2-3 days later in the bottom channels of all ALoCs. Prior to HMVEC cell

seeding, media was replenished daily for the ATs seeded on the ALoCs. The SAGM medium used

for maintenance of the ATs on the chips (hereby called AT Maintenance Medium) was

supplemented with Dexamethasone at 100 nM (Sigma D4902), Keratinocyte growth factor (KGF)

at 5 ng/mL (Thermo Fisher PHG0094), 8-Br-cAMP at 50 µM (Sigma B7880), and Isobutyl

methylxanthine (IBMX) at 25 µM (Sigma I7018). When HMVECs reached ~80% confluency, they

were detached and adjusted to a concentration of 5×10^6 cells/mL in EGM-2MV medium

(described above). The bottom channels of all ALoCs were filled with EGM-2MV medium prior to

seeding. 20 µL of the HMVEC cell suspension was pipetted rapidly into the bottom channel and

checked under the microscope for correct seeding density and cell homogeneity. The bottom

channel volume is ~6 µL total. Similarly to the top channel, the bottom channel was overfilled to

prevent the introduction of bubbles to the channel. The chip was then immediately flipped upside

down and placed in the chip cradle to ensure cells were seeded on the porous membrane in the

bottom channel. Once all ALoCs were seeded and flipped upside down, they were incubated at

37°C for at least 2 hours until all cells were attached. After all cells had attached, the bottom

channel was washed with 200 µL of warm EGM-2MV medium and incubated overnight at 37°C.

Prior to attaching ALoCs to Pods, warm EGM-2MV and AT Maintenance medium were degassed

using a Steriflip device for 5 min each to prevent air bubbles being trapped in the microfluidic lines

of the Pod. 3 mL of each medium were pipetted into its respective reservoir within the Pod. Using

Emulate's Zoe, the Pods were primed with medium to prevent bubbles from impeding medium

flow to the ALoCs. Once primed, each ALoC was snapped into each respective Pod. To further

help prevent bubble formation within the microfluidic lines, a Regulate Cycle was performed on

each chip via the Zoe. The ALoCs from this point on were harbored within the Pods and fresh

medium was exchanged through the chip at a flow rate of 30 µL/hr supplied by the Zoe.

After 24 hours of continuous media flow on all ALoCs, air-liquid interface (ALI) was introduced to

all ALoCs by removing all medium from the top channel and changing the top channel from liquid

media flow to air. The bottom channel medium was changed to ALI medium which comprised of

a Medium 199 base (Thermo Fisher 11043023) supplemented with 10 ng/mL Human Epidermal

Growth Factor (Peprotech AF-100-15), 3 ng/mL Human Basic Fibroblast Growth Factor

(Peprotech AF-100-18B), 0.125 ng/mL Human Vascular Endothelial Growth Factor (Peprotech

AF-100-20), 1 µg/mL Hydrocortisone (Sigma H0135), 10 µg/mL Heparin (Sigma H3149), 80 µM

di-butyryl cAMP (Sigma B7880), 1mM L-Glutamax (Gibco 35050-061), 20 nM Dexamethasone

(Sigma D4902), 1% Pen-Srep solution (Gibco 125140-122), and 2% FBS. Bottom channel liquid

flow rate was set to 30 µL/hr and fresh ALI medium was replenished in the bottom channel

reservoir every 2-3 days. After 48 hours of ALI, mechanical stretch (5%, 0.20 Hz) was initiated

using the Zoe. All ALoCs were maintained in this way until seeding of macrophages and infection

with *M. fortuitum*.

Macrophage Seeding on ALoCs

After 7 days of differentiation with M-CSF to HMDMs, macrophages were detached from petri

dishes using ice-cold 5 mM EDTA in PBS and gentle scraping. HMDMs were centrifuged at 300xg

for 5 min and resuspended in a 1:1 ratio of AT Maintenance Medium (without dexamethasone)

and Macrophage Medium to a concentration of 1×10^6 cells/mL. ALoCs were removed from the

Zoe and detached from the Pods, and top channel was washed with 200 µL of HPAEC

Maintenance Medium. With medium still in the top channel, 50 µL of HMDM cell suspension was

rapidly pipetted through the top channel. ALoCs were placed in square petri dishes and incubated

at 37°C with 5% CO_2 for 3-4 hours to allow attachment to the top channel. Once attached, ALoCs

were attached back to Pods after priming and after 2 hours of a Regulate Cycle were returned to

ALI.

***M. fortuitum* Infection on Human ALoC**

Upon returning ALoCs to ALI conditions after macrophage attachment to the ALoCs, ALI medium

was replaced with ALI medium without penicillin/streptomycin, to avoid impacting subsequent

bacterial infection. Twenty-four hours after returning to ALI conditions, the ALoCs were infected

with *M. fortuitum*. mCherry-expressing *M. fortuitum* was cultured at 37°C in liquid 7H9 medium

(BD Difco 271310) supplemented with 10% OADC enrichment (BD 212351), 50 µg/mL

Kanamycin (Sigma 60615) and 0.05% Tyloxapol (Sigma T8761) until it reached an OD_{600} 0.5 –

0.7. The culture was pelleted by centrifugation at 3500 RPM for 10 min. The pellet was washed

418 three times with 50 mL of 1X PBS (-Ca$^+$/-Mg$^+$) by centrifugation at 3500 RPM for 10 min, and

419 resuspended in 5 mL PBS for a slow spin to remove cell debris at 500 RPM for 5 min. The

420 supernatant was collected and passed through a 26-gauge needle three times to generate a

421 single-cell suspension. The bacterial suspension was adjusted to an MOI of 1 with respect to the

422 epithelial cells in HPAEC Maintenance Medium, and 50 µL of the *M. fortuitum* suspension was

423 pipetted rapidly through the top channel. The ALoCs were incubated at 37°C for 1 hr. under static

424 conditions, and then washed three times with epithelial medium before returning ALoCs back to

425 Pods and into the Zoe for another Regulate Cycle. ALI was initiated again and the ALoCs were

426 infected for a total of 24 hours.

RNA Extraction from ALoCs

428 ALoCs were removed from the Zoe and Pods. Top and bottom channels were washed three times

429 with 200 µL of ice-cold 1X DPBS (-Ca$^+$/-Mg$^+$). An empty p200 filtered tip was inserted into the

430 bottom inlet, bottom outlet, and top outlet ports. 200 µL of TRIzol reagent (Invitrogen #15596026)

431 was pipetted in each ALoC by rapidly pressing and releasing the plunger three times before

432 collecting the supernatant in an RNase-free 1.5 mL tube. This process was repeated once more

433 for a total of ~400 µL of supernatant. TRIzol was added to each tube of RNA to a final volume of

434 1 mL. The RNA was stored at -80°C before extraction using the Qiagen RNeasy Mini Columns

435 (74106). AT and macrophage lysates were thawed on ice and incubated at room temperature for

436 5 min. 200 µL of chloroform per 1 mL of lysates was added to each tube, shaken vigorously for

437 15 sec., incubated at room temperature for 2-3 min, and centrifuged for 5 min at 12,000xg at 4°C.

438 The aqueous phase was transferred to fresh RNase-free tubes and the rest of the RNA extraction

439 was performed according to the Qiagen RNeasy Mini Column Kit. Extracted RNA was stored at -

440 80°C until use for library preparation and RNA-sequencing.

Library Preparation and RNA-sequencing

Samples were analyzed on an Agilent Tapestation 4200 to determine level of degradation to

ensure that only high quality RNA was used (RIN Score 8 or higher). We used a Qubit 4.0

Fluorimeter (ThermoFisher) to determine RNA concentration prior to starting library prep. One

microgram of total DNAse treated RNA was then prepared with the TruSeq Stranded mRNA

Library Prep Kit (Illumina). Poly-A RNA was purified and fragmented before strand-specific cDNA

synthesis. cDNA was then A-tailed and indexed adapters ligated. After adapter ligation, samples

were PCR amplified and purified with AmpureXP beads, then validated again on the Agilent

Tapestation 4200. Before being normalized and pooled, samples were quantified by Qubit then

sequenced on the Illumina NextSeq 2000 using a P2-100 flowcell.

Bulk RNA-sequencing Analysis

All RNA-sequencing analyses were performed using integrated Differential Expression and

Pathways Analysis, iDEP 2.01 (Ge et al., 2018). Genes expressed at extremely low levels were

filtered out of the gene set by removing any genes with less than 0.4 counts per million (CPM) in

at least one sample (n=1) (**Supplemental Figure 3A**). 13,934 filtered genes were converted to

Ensembl gene IDs, normalized in edgeR and transformed via rlog. Hierarchical clustering of the

samples and the top 2000 genes were performed (**Supplemental Figure 3B**). We then performed

principal component analysis (PCA) using the 2000 most variable genes. Using DESeq2 and the

Wald test (FDR cutoff = 0.1, minimum fold change = 1.5), we generated adjusted p-values and

log_2 fold changes. Gene expression was then compared between the "infected" AloCs and

"control_uninfected" ALoCs. A total of 1,454 genes were found to be downregulated (log_2 fold

change <1) and 588 genes were upregulated (log_2 fold change >1). Using analysis of variance

(ANOVA) and post-hoc pairwise t-tests, we evaluated between-group differences for each cluster

using these scaled expression values. We performed functional enrichment analysis using the

hypergeometric test in hypeR for enriched pathways within the differentially expressed gene set.

Enrichment analysis was run using significantly upregulated and significantly downregulated

genes. Gene categories with fewer than 15 genes were excluded. Gene categories were

considered significant if they had Benjamini-Hochberg adjusted *p*-values less than 0.1.

Immunofluorescence

ALoCs were washed three times with 200 µL of 1X DPBS and 200 uL of 4% PFA in PBS were

added to each channel of each ALoC for 20 min. at room temperature. ALoCs were washed three

times with 200 µL of 1X DPBS and incubated for 30 min. at room temperature with 200 µL of 0.1%

Triton X-100 and 2% saponin. The ALoCs were then washed three times with 200 µL of 1X DPBS

and incubated overnight at 4°C with primary antibody (1:100) (**Supplemental Table 1**). ALoCs

were washed again three times with 200 µL of DPBS and incubated with secondary antibody

(1:1000) (**Supplemental Table 1**) for 2 hours at room temperature, protected from light. ALoCs

were washed three times with 200 µL of 1X DPBS and incubated with DAPI for 10 min at room

temperature. They were washed again three times with 200 µL of 1X DPBS and stored at 4°C

with both channels filled with 1X DPBS. Confocal images were acquired on a Nikon CSU W1

spinning disk confocal microscope with an CFI S Plan Fluor ELWD 10X objective.

Statistical analyses

All statistical analyses were performed using GraphPad Prism Software (version 9). For *in vitro*

studies, data was analyzed using unpaired two-tailed *t*-test. For RNA sequencing analysis,

analysis of variance (ANOVA) and post-hoc pairwise t-tests were used to evaluate between-group

differences for each cluster using scaled expression values.

ACKNOWLEDGEMENTS

The authors thank Gautam Mahajan and Ben Swenor from Emulate Inc. for providing protocols

and support. The authors also thank core facilities at UT Southwestern Medical Center for their

important contributions to this work including the UT Southwestern McDermott Center Next

Generation Sequencing (NGS) Core and the Quantitative Light Microscopy Core, particularly the

core director Marcel Mettlen, a Shared Resource of the Harold C. Simmons Cancer Center,

supported in part by an NCI Cancer Center Support Grant, 1P30 CA142543-01.

COMPETING INTERESTS

All authors declare that they have no competing interests.

FUNDING

This work was supported by the National Institutes of Health U19 AI142784, R01 AI184584, R01

AI158688 and P01 AI159402 to M.U.S.

DATA AVAILABILITY

Bulk RNA sequencing data has been deposited at the NCBI GEO database, GSE276053, and

will be made publicly available at the time of publication.

AUTHOR CONTRIBUTIONS

Conceptualization, V.E., M.U.S.; Formal analysis, V.E., M.U.S.; Funding acquisition, M.U.S.;

Investigation, V.E., B.R., P.C.; Project administration, M.U.S.; Supervision, M.U.S.; Visualization,

M.U.S.; Writing – original draft, V.E., M.U.S.; Writing – review and editing, All authors.

References

ALKARNI, M., LIPMAN, M. and LOWE, D. M. 2023. The roles of neutrophils in non-tuberculous
mycobacterial pulmonary disease. *Ann Clin Microbiol Antimicrob,* 22**,** 14.

ARDAIN, A., DOMINGO-GONZALEZ, R., DAS, S., KAZER, S. W., HOWARD, N. C., SINGH, A.,
AHMED, M., NHAMOYEBONDE, S., RANGEL-MORENO, J., OGONGO, P., LU, L.,
RAMSURAN, D., DE LA LUZ GARCIA-HERNANDEZ, M., T, K. U., DARBY, M., PARK, E.,
KARIM, F., MELOCCHI, L., MADANSEIN, R., DULLABH, K. J., DUNLAP, M., MARIN-
AGUDELO, N., EBIHARA, T., NDUNG'U, T., KAUSHAL, D., PYM, A. S., KOLLS, J. K.,
STEYN, A., ZUNIGA, J., HORSNELL, W., YOKOYAMA, W. M., SHALEK, A. K.,
KLOVERPRIS, H. N., COLONNA, M., LESLIE, A. and KHADER, S. A. 2019. Group 3 innate
lymphoid cells mediate early protective immunity against tuberculosis. *Nature,* 570**,** 528-532.

BAI, H., SI, L., JIANG, A., BELGUR, C., ZHAI, Y., PLEBANI, R., OH, C. Y., RODAS, M., PATIL,
A., NURANI, A., GILPIN, S. E., POWERS, R. K., GOYAL, G., PRANTIL-BAUN, R. and
INGBER, D. E. 2022. Mechanical control of innate immune responses against viral infection
revealed in a human lung alveolus chip. *Nat Commun,* 13**,** 1928.

BALDWIN, S. L., LARSEN, S. E., ORDWAY, D., CASSELL, G. and COLER, R. N. 2019. The
complexities and challenges of preventing and treating nontuberculous mycobacterial
diseases. *PLoS Negl Trop Dis,* 13**,** e0007083.

BENAM, K. H., DAUTH, S., HASSELL, B., HERLAND, A., JAIN, A., JANG, K. J., KARALIS, K.,
KIM, H. J., MACQUEEN, L., MAHMOODIAN, R., MUSAH, S., TORISAWA, Y. S., VAN DER
MEER, A. D., VILLENAVE, R., YADID, M., PARKER, K. K. and INGBER, D. E. 2015.
Engineered in vitro disease models. *Annu Rev Pathol,* 10**,** 195-262.

BENAM, K. H., VILLENAVE, R., LUCCHESI, C., VARONE, A., HUBEAU, C., LEE, H. H.,
ALVES, S. E., SALMON, M., FERRANTE, T. C., WEAVER, J. C., BAHINSKI, A., HAMILTON,
G. A. and INGBER, D. E. 2016. Small airway-on-a-chip enables analysis of human lung
inflammation and drug responses in vitro. *Nat Methods,* 13**,** 151-7.

BERRY, M. P., GRAHAM, C. M., MCNAB, F. W., XU, Z., BLOCH, S. A., ONI, T., WILKINSON, K.
A., BANCHEREAU, R., SKINNER, J., WILKINSON, R. J., QUINN, C., BLANKENSHIP, D.,
DHAWAN, R., CUSH, J. J., MEJIAS, A., RAMILO, O., KON, O. M., PASCUAL, V.,
BANCHEREAU, J., CHAUSSABEL, D. and O'GARRA, A. 2010. An interferon-inducible
neutrophil-driven blood transcriptional signature in human tuberculosis. *Nature,* 466**,** 973-7.

BETHENCOURT, M. A., NGUYEN, A. D. and FERRER, G. 2024. Lung Nontuberculous
Mycobacterial Infections. *StatPearls.* Treasure Island (FL).

BOHRER, A. C., CASTRO, E., HU, Z., QUEIROZ, A. T. L., TOCHENY, C. E., ASSMANN, M.,
SAKAI, S., NELSON, C., BAKER, P. J., MA, H., WANG, L., ZILU, W., DU BRUYN, E., RIOU,
C., KAUFFMAN, K. D., TUBERCULOSIS IMAGING, P., MOORE, I. N., DEL NONNO, F.,
PETRONE, L., GOLETTI, D., MARTINEAU, A. R., LOWE, D. M., CRONAN, M. R.,
WILKINSON, R. J., BARRY, C. E., VIA, L. E., BARBER, D. L., KLION, A. D., ANDRADE, B.
B., SONG, Y., WONG, K. W. and MAYER-BARBER, K. D. 2021. Eosinophils are part of the
granulocyte response in tuberculosis and promote host resistance in mice. *J Exp Med,* 218.

CAMPBELL, G. R. and SPECTOR, S. A. 2012. Vitamin D inhibits human immunodeficiency
virus type 1 and Mycobacterium tuberculosis infection in macrophages through the induction
of autophagy. *PLoS Pathog,* 8**,** e1002689.

CAVERLY, L. J., CACERES, S. M., FRATELLI, C., HAPPOLDT, C., KIDWELL, K. M.,
MALCOLM, K. C., NICK, J. A. and NICHOLS, D. P. 2015. Mycobacterium abscessus
morphotype comparison in a murine model. *PLoS One,* 10**,** e0117657.

CHAN, E. D., BAI, X., ORDWAY, D. J. and VERMA, D. 2016. Animal Models of Non-
Tuberculous Mycobacterial Infections. *Mycobacterial Diseases,* 6.

COWMAN, S. A., JACOB, J., HANSELL, D. M., KELLEHER, P., WILSON, R., COOKSON, W.
O. C., MOFFATT, M. F. and LOEBINGER, M. R. 2018. Whole-Blood Gene Expression in
Pulmonary Nontuberculous Mycobacterial Infection. *Am J Respir Cell Mol Biol,* 58**,** 510-518.

CRISTANCHO-ROJAS, C., VARLEY, C. D., LARA, S. C., KHERABI, Y., HENKLE, E. and
WINTHROP, K. L. 2023. Epidemiology of Mycobacterium abscessus. *Clin Microbiol Infect*.

DASGUPTA, Q., JIANG, A., WEN, A. M., MANNIX, R. J., MAN, Y., HALL, S., JAVORSKY, E.
and INGBER, D. E. 2023. A human lung alveolus-on-a-chip model of acute radiation-induced
lung injury. *Nat Commun,* 14**,** 6506.

DE GROOTE, M. A., JOHNSON, L., PODELL, B., BROOKS, E., BASARABA, R. and
GONZALEZ-JUARRERO, M. 2014. GM-CSF knockout mice for preclinical testing of agents
with antimicrobial activity against Mycobacterium abscessus. *J Antimicrob Chemother,* 69**,**
1057-64.

DEINHARDT-EMMER, S., RENNERT, K., SCHICKE, E., CSERESNYES, Z., WINDOLPH, M.,
NIETZSCHE, S., HELLER, R., SIWCZAK, F., HAUPT, K. F., CARLSTEDT, S., SCHACKE, M.,
FIGGE, M. T., EHRHARDT, C., LOFFLER, B. and MOSIG, A. S. 2020. Co-infection with
Staphylococcus aureus after primary influenza virus infection leads to damage of the
endothelium in a human alveolus-on-a-chip model. *Biofabrication,* 12**,** 025012.

DONOHUE, M. J. 2021. Epidemiological risk factors and the geographical distribution of eight
Mycobacterium species. *BMC Infect Dis,* 21**,** 258.

FRANCO, L. H., NAIR, V. R., SCHARN, C. R., XAVIER, R. J., TORREALBA, J. R., SHILOH, M.
U. and LEVINE, B. 2017. The Ubiquitin Ligase Smurf1 Functions in Selective Autophagy of
Mycobacterium tuberculosis and Anti-tuberculous Host Defense. *Cell Host Microbe,* 21**,** 59-
72.

FU, Y. L. and HARRISON, R. E. 2021. Microbial Phagocytic Receptors and Their Potential
Involvement in Cytokine Induction in Macrophages. *Front Immunol,* 12**,** 662063.

GANGADHARAM, P. R. 1995. Beige mouse model for Mycobacterium avium complex disease.
Antimicrob Agents Chemother, 39**,** 1647-54.

GE, S. X., SON, E. W. and YAO, R. 2018. iDEP: an integrated web application for differential
expression and pathway analysis of RNA-Seq data. *BMC Bioinformatics,* 19**,** 534.

GONZALEZ-PEREZ, M., MARINO-RAMIREZ, L., PARRA-LOPEZ, C. A., MURCIA, M. I., MARQUINA, B., MATA-ESPINOZA, D., RODRIGUEZ-MIGUEZ, Y., BAAY-GUZMAN, G. J., HUERTA-YEPEZ, S. and HERNANDEZ-PANDO, R. 2013. Virulence and immune response induced by Mycobacterium avium complex strains in a model of progressive pulmonary tuberculosis and subcutaneous infection in BALB/c mice. *Infect Immun,* 81**,** 4001-12.

HAIQING BAI, L. S., AMANDA JIANG, CHAITRA BELGUR, YUNHAO ZHAI, ROBERTO PLEBANI,, CRYSTAL YURI OH, M. R., ADITYA PATIL, ATIQ NURANI , SARAH E. GILPIN, RANI K. POWERS, GIRIJA GOYAL, and INGBER, R. P.-B. D. E. 2022. Mechanical control of innate immune responses against viral infection revealed in a human lung alveolus chip. *Nature Communications,* 13.

HOANG, T. N. M., CSERESNYES, Z., HARTUNG, S., BLICKENSDORF, M., SAFFER, C., RENNERT, K., MOSIG, A. S., VON LILIENFELD-TOAL, M. and FIGGE, M. T. 2022. Invasive aspergillosis-on-chip: A quantitative treatment study of human Aspergillus fumigatus infection. *Biomaterials,* 283**,** 121420.

HUH, D., LESLIE, D. C., MATTHEWS, B. D., FRASER, J. P., JUREK, S., HAMILTON, G. A., THORNELOE, K. S., MCALEXANDER, M. A. and INGBER, D. E. 2012a. A human disease model of drug toxicity-induced pulmonary edema in a lung-on-a-chip microdevice. *Sci Transl Med,* 4**,** 159ra147.

HUH, D., MATTHEWS, B. D., MAMMOTO, A., MONTOYA-ZAVALA, M., HSIN, H. Y. and INGBER, D. E. 2010. Reconstituting organ-level lung functions on a chip. *Science,* 328**,** 1662-8.

HUH, D., TORISAWA, Y. S., HAMILTON, G. A., KIM, H. J. and INGBER, D. E. 2012b. Microengineered physiological biomimicry: organs-on-chips. *Lab Chip,* 12**,** 2156-64.

JAIN, A., BARRILE, R., VAN DER MEER, A. D., MAMMOTO, A., MAMMOTO, T., DE CEUNYNCK, K., AISIKU, O., OTIENO, M. A., LOUDEN, C. S., HAMILTON, G. A., FLAUMENHAFT, R. and INGBER, D. E. 2018. Primary Human Lung Alveolus-on-a-chip Model of Intravascular Thrombosis for Assessment of Therapeutics. *Clin Pharmacol Ther,* 103**,** 332-340.

JOHANSEN, M. D. and KREMER, L. 2020. CFTR Depletion Confers Hypersusceptibility to Mycobacterium fortuitum in a Zebrafish Model. *Front Cell Infect Microbiol,* 10**,** 357.

KILINC, G., WALBURG, K. V., FRANKEN, K., VALKENBURG, M. L., AUBRY, A., HAKS, M. C., SARIS, A. and OTTENHOFF, T. H. M. 2022. Development of Human Cell-Based In Vitro Infection Models to Determine the Intracellular Survival of Mycobacterium avium. *Front Cell Infect Microbiol,* 12**,** 872361.

KIM, S. R., YU, J. Y., KIM, S. Y., KIM, D. H. and JHUN, B. W. 2023. Clinical Characteristics and Treatment Outcomes of Mycobacterium fortuitum Pulmonary Disease. *Microbiol Spectr,* 11**,** e0205123.

KIMMEY, J. M., HUYNH, J. P., WEISS, L. A., PARK, S., KAMBAL, A., DEBNATH, J., VIRGIN, H. W. and STALLINGS, C. L. 2015. Unique role for ATG5 in neutrophil-mediated immunopathology during M. tuberculosis infection. *Nature,* 528**,** 565-9.

KLEVER, A. M., ALEXANDER, K. A., ALMEIDA, D., ANDERSON, M. Z., BALL, R. L., BEAMER, G., BOGGIATTO, P., BUIKSTRA, J. E., CHANDLER, B., CLAEYS, T. A., CONCHA, A. E., CONVERSE, P. J., DERBYSHIRE, K. M., DOBOS, K. M., DUPNIK, K. M., ENDSLEY, J. J., ENDSLEY, M. A., FENNELLY, K., FRANCO-PAREDES, C., HAGGE, D. A., HALL-STOODLEY, L., HAYES, D., JR., HIRSCHFELD, K., HOFMAN, C. A., HONDA, J. R., HULL, N. M., KRAMNIK, I., LACOURCIERE, K., LAHIRI, R., LAMONT, E. A., LARSEN, M. H., LEMAIRE, T., LESELLIER, S., LEE, N. R., LOWRY, C. A., MAHFOOZ, N. S., MCMICHAEL, T. M., MERLING, M. R., MILLER, M. A., NAGAJYOTHI, J. F., NELSON, E., NUERMBERGER, E. L., PENA, M. T., PEREA, C., PODELL, B. K., PYLE, C. J., QUINN, F. D., RAJARAM, M. V. S., MEJIA, O. R., ROTHOFF, M., SAGO, S. A., SALVADOR, L. C. M., SIMONSON, A. W., SPENCER, J. S., SREEVATSAN, S., SUBBIAN, S., SUNSTRUM, J., TOBIN, D. M., VIJAYAN, K. K. V., WRIGHT, C. T. O. and ROBINSON, R. T. 2023. The Many Hosts of Mycobacteria 9 (MHM9): A conference report. *Tuberculosis (Edinb),* 142**,** 102377.

KREMER, M. D. J. A. L. 2020. CFTR Depletion Confers Hypersusceptibility to Mycobacterium fortuitum in a Zebrafish Model. *Frontiers in Cellular and Infection Microbiology,* 10.

LAGUNE, M., KREMER, L. and HERRMANN, J. L. 2023. Mycobacterium abscessus, a complex of three fast-growing subspecies sharing virulence traits with slow-growing mycobacteria. *Clin Microbiol Infect*.

LI, J., WANG, Z., CHU, Q., JIANG, K., LI, J. and TANG, N. 2018. The Strength of Mechanical Forces Determines the Differentiation of Alveolar Epithelial Cells. *Dev Cell,* 44**,** 297-312 e5.

LI, J., WEN, A. M., POTLA, R., BENSHIRIM, E., SEEBARRAN, A., BENZ, M. A., HENRY, O. Y. F., MATTHEWS, B. D., PRANTIL-BAUN, R., GILPIN, S. E., LEVY, O. and INGBER, D. E. 2019. AAV-mediated gene therapy targeting TRPV4 mechanotransduction for inhibition of pulmonary vascular leakage. *APL Bioeng,* 3**,** 046103.

LINDESTAM ARLEHAMN, C. S., BENSON, B., KUAN, R., DILL-MCFARLAND, K. A., PETERSON, G. J., PAUL, S., NGUYEN, F. K., GILMAN, R. H., SAITO, M., TAPLITZ, R., ARENTZ, M., GOSS, C. H., AITKEN, M. L., HORNE, D. J., SHAH, J. A., SETTE, A. and HAWN, T. R. 2022. T-cell deficiency and hyperinflammatory monocyte responses associate with Mycobacterium avium complex lung disease. *Front Immunol,* 13**,** 1016038.

LUKE, C. J., MARKOVINA, S., GOOD, M., WIGHT, I. E., THOMAS, B. J., LINNEMAN, J. M., LANIK, W. E., KOROLEVA, O., COFFMAN, M. R., MIEDEL, M. T., GONG, Q., ANDRESS, A., CAMPOS GUERRERO, M., WANG, S., CHEN, L., BEATTY, W. L., HAUSMANN, K. N., WHITE, F. V., FITZPATRICK, J. A. J., ORVEDAHL, A., PAK, S. C. and SILVERMAN, G. A. 2022. Lysoptosis is an evolutionarily conserved cell death pathway moderated by intracellular serpins. *Commun Biol*, 5**,** 47.

MERCALDO, R. A., MARSHALL, J. E., CANGELOSI, G. A., DONOHUE, M., FALKINHAM, J. O., 3RD, FIERER, N., FRENCH, J. P., GEBERT, M. J., HONDA, J. R., LIPNER, E. M., MARRAS, T. K., MORIMOTO, K., SALFINGER, M., STOUT, J., THOMSON, R. and PREVOTS, D. R. 2023. Environmental risk of nontuberculous mycobacterial infection: Strategies for advancing methodology. *Tuberculosis (Edinb),* 139**,** 102305.

MISHRA, R., HANNEBELLE, M., PATIL, V. P., DUBOIS, A., GARCIA-MOUTON, C., KIRSCH, G. M., JAN, M., SHARMA, K., GUEX, N., SORDET-DESSIMOZ, J., PEREZ-GIL, J., PRAKASH, M., KNOTT, G. W., DHAR, N., MCKINNEY, J. D. and THACKER, V. V. 2023. Mechanopathology of biofilm-like Mycobacterium tuberculosis cords. *Cell,* 186**,** 5135-5150 e28.

NOBS, E., LASCHANZKY, K., MUNKE, K., MOVERT, E., VALFRIDSSON, C. and CARLSSON, F. 2024. Cytosolic serpins act in a cytoprotective feedback loop that limits ESX-1-dependent death of Mycobacterium marinum-infected macrophages. *mBio***,** e0038424.

ORDWAY, D., HENAO-TAMAYO, M., SMITH, E., SHANLEY, C., HARTON, M., TROUDT, J., BAI, X., BASARABA, R. J., ORME, I. M. and CHAN, E. D. 2008. Animal model of Mycobacterium abscessus lung infection. *J Leukoc Biol,* 83**,** 1502-11.

ORME, I. M. and ORDWAY, D. J. 2014. Host response to nontuberculous mycobacterial infections of current clinical importance. *Infect Immun,* 82**,** 3516-22.

QIN, M. H., MADIRAJU, M. V. and RAJAGOPALAN, M. 1999. Characterization of the functional replication origin of Mycobacterium tuberculosis. *Gene,* 233**,** 121-30.

RAMPACCI, E., STEFANETTI, V., PASSAMONTI, F. and HENAO-TAMAYO, M. 2020. Preclinical Models of Nontuberculous Mycobacteria Infection for Early Drug Discovery and Vaccine Research. *Pathogens,* 9.

ROQUET-BANERES, F., ALCARAZ, M., HAMELA, C., ABENDROTH, J., EDWARDS, T. E. and KREMER, L. 2023. In Vitro and In Vivo Efficacy of NITD-916 against Mycobacterium fortuitum. *Antimicrob Agents Chemother,* 67**,** e0160722.

TABONE, O., VERMA, R., SINGHANIA, A., CHAKRAVARTY, P., BRANCHETT, W. J., GRAHAM, C. M., LEE, J., TRANG, T., REYNIER, F., LEISSNER, P., KAISER, K., RODRIGUE, M.,

WOLTMANN, G., HALDAR, P. and O'GARRA, A. 2021. Blood transcriptomics reveal the
evolution and resolution of the immune response in tuberculosis. *J Exp Med,* 218.

THACKER, V. V., DHAR, N., SHARMA, K., BARRILE, R., KARALIS, K. and MCKINNEY, J. D.
2020. A lung-on-chip model of early Mycobacterium tuberculosis infection reveals an
essential role for alveolar epithelial cells in controlling bacterial growth. *Elife,* 9.

TOURE, H., GALINDO, L. A., LAGUNE, M., GLATIGNY, S., WATERHOUSE, R. M., GUENAL, I.,
HERRMANN, J. L., GIRARD-MISGUICH, F. and SZUPLEWSKI, S. 2023. Mycobacterium
abscessus resists the innate cellular response by surviving cell lysis of infected phagocytes.
PLoS Pathog, 19**,** e1011257.

VARLEY, C. D. and WINTHROP, K. L. 2022. Nontuberculous Mycobacteria: Diagnosis and
Therapy. *Clin Chest Med,* 43**,** 89-98.

WANG, L., MA, H., WEN, Z., NIU, L., CHEN, X., LIU, H., ZHANG, S., XU, J., ZHU, Y., LI, H.,
CHEN, H., SHI, L., WAN, L., LI, L., LI, M., WONG, K. W. and SONG, Y. 2023a. Single-cell
RNA-sequencing reveals heterogeneity and intercellular crosstalk in human tuberculosis
lung. *J Infect,* 87**,** 373-384.

WANG, L., ZHENG, X., MA, J., GU, J. and SHA, W. 2023b. Comparative Proteomic Analysis of
Exosomes Derived from Patients Infected with Non-Tuberculous Mycobacterium and
Mycobacterium tuberculosis. *Microorganisms,* 11.

WANG, P., YANG, G. L., HE, Y. F., SHEN, Y. H., HAO, X. H., LIU, H. P., SHEN, H. B., WANG, L.
and SHA, W. 2024. Single-cell transcriptomics of blood identified IFIT1(+) neutrophil
subcluster expansion in NTM-PD patients. *Int Immunopharmacol,* 137**,** 112412.

WINTHROP, K. L., MARRAS, T. K., ADJEMIAN, J., ZHANG, H., WANG, P. and ZHANG, Q.
2020. Incidence and Prevalence of Nontuberculous Mycobacterial Lung Disease in a Large
U.S. Managed Care Health Plan, 2008-2015. *Ann Am Thorac Soc,* 17**,** 178-185.

 Figure legends

Figure 1. **Establishment of a fully humanized alveolus lung-on-a-chip (ALoC) model system**

A. Schematic of a human alveolus and the surrounding vasculature, highlighting alveolar epithelial

type I (ATI), alveolar epithelial type II (AT2) cells, and alveolar macrophages in the alveolar sac,

with pulmonary microvascular endothelial cells lining the vasculature. **B.** Cross-section of the

ALoC model for *M. fortuitum* infection. The apical channel contains primary human alveolar

epithelial cells (ATI and ATII) with or without human CD14$^+$ peripheral blood-derived

macrophages, while the basal channel is lined by human primary pulmonary microvascular

endothelial cells (HMVECs). **C.** Schematic of the addition of macrophages to the apical channel.

D. Stitched image of the apical channel of the ALoC at 15 days of culture highlighting AT1 cells

(HT1$^+$, red) and AT2 cells (HT2$^+$, green). Nuclei are labelled with DAPI (blue). Scale bar = 1000

μm. **E.** Epifluorescent image of differentiated AT1 and AT2 cells on the ALoC. Scale bar = 200

μm. **F.** Three-dimensional immunofluorescence reconstruction of ALoC after 15 days of culture,

highlighting differentiated AT2 cells and AT1 cells. Scale bar = 500 μm. **G.** Cross-sectional image

of the ALoC highlighting AT1 and AT2 cells in a monolayer. Scale bar = 200 μm. **H.** Stitched

image of the apical epithelial channel of the ALoC after the addition of HMDMs (CD14$^+$, yellow)

on the ALoC. Nuclei are labelled with DAPI. Scale bar = 1,000 μm. **I.** Magnified view of the

outlined area in panel H of AT1 cells, AT2 cells, and macrophages. Scale bar = 200 μm. **J.** Three-

dimensional reconstruction of ALoC highlighting the addition of macrophages amongst

differentiated AT1 and AT2 cells. Scale bar = 200 μm. **K.** Cross-sectional image of the ALoC

showing AT1 cells (HT1$^+$, red), AT2 cells (HT2$^+$, green), and HMDMs (CD14$^+$, yellow). Scale bar

= 200 μm. Images are representative of at least 3 independent chips from 2 independent

experiments.

Figure 2. *Mycobacterium fortuitum* **infection on an Alveolus Lung-on-a-Chip without**

macrophages A. Schematic of *M. fortuitum* addition to the ALoC. *M. fortuitum* was added to the

ALoC at an MOI of 1 with respect to the alveolar epithelial cells. After 1 hour of static incubation

at 37°C, ALoCs were washed and returned to ALI. **B.** Stitched image of mCherry-expressing *M.*

fortuitum (red) on ALoCs with AT1 (HT1$^+$, purple) and AT2 (HT2$^+$, green) cells. Nuclei labelled with

DAPI (Blue). Scale bar = 500 µm**. C.** Immunofluorescence image highlighting alveolar epithelial

cell damage. Scale bar = 200 µm. **D.** Three-dimensional reconstruction of an infected ALoC

without macrophages. AT2 cells with "flattened" morphology highlighted within normal cuboidal

AT2 morphology. Scale bar = 200 µm**. E.** Cross-sectional image of *M. fortuitum* (mCherry, red)

infection on ALoC with AT1 and AT2 cells. Scale bar = 200 µm**.** Images are representative of at

least 3 independent chips from 2 independent experiments.

Figure 3. ***Mycobacterium fortuitum* infection on an Alveolus Lung-on-a-Chip containing**

macrophages A. Schematic of *M. fortuitum* infection on ALoC containing macrophages. 24 hours

after the addition macrophages, *M. fortuitum* was added to the ALoC at an MOI of 1 with respect

to the alveolar epithelial cells, incubated at 37°C for 1 hour, washed, and returned to ALI. **B**.

Stitched image of the apical channel, highlighting macrophages (CD14$^+$, grey) and AT2 cells

(HT2$^+$, green) 24 hours after infection with mCherry-expressing *M. fortuitum* (red). Nuclei are

labelled by DAPI. Scale bar = 500 µm. **C.** Immunofluorescence image highlighting *M. fortuitum*

(mCherry, red)-infected macrophages (CD14$^+$, grey) and AT2 cells (HT2$^+$, green). Scale bar = 200

µm. **D.** Three-dimensional reconstructed image of infected macrophages (CD14, grey) along the

edge of an ALoC. Scale bar = 350 µm. **E.** Cross-sectional image of *M. fortuitum* infection on ALoC

with the addition of macrophages. **F.** Three-dimensional reconstruction of *M. fortuitum* bacilli and

macrophages on ALoCs. Scale bar = 200 µm. **G.** Quantification of *M. fortuitum* bacilli within

macrophages or AT2 cell. Each point represents the cumulative number of *M. fortuitum* bacilli

counted in each cell type for 10 fields within a single chip (N=3 independent chips, $p < 0.05$ by

unpaired Student's t-test). Images are representative of at least 3 independent chips from 2

independent experiments.

Figure 4. **Bulk RNA-seq Analysis of *M. fortuitum*-infected ALoCs with macrophages A.**

Heat-map of the top 2000 genes expressed in uninfected and *M. fortuitum*-infected ALoCs (24

hours post-infection) (N=3 per group) using Pearson distance and average linkage. **B.** The -log$_{10}$

(Adjusted p-Value) was plotted against log$_2$ fold change to create a volcano plot representing the

top 25 differentially expressed genes (DEGs). Genes with an adjusted p-value<25 and log$_2$ fold

change<2 were filtered from this analysis. **C.** Heat map comparing the log$_2$ fold change of the

most differentially expressed chemokines and cytokines in ALoCs infected with *M. fortuitum*

versus uninfected ALoCs after 24 hours of infection. **D.** Heat map comparing the log$_2$ fold change

of the most differentially expressed SERPINs in ALoCS infected with *M. fortuitum* versus

uninfected ALoCs after 24 hours of infection.

Supplemental Figure Legends

Supplemental Figure 1. **Establishment of the Alveolus Lung-on-a-Chip Model**. **A.** Typical timeline of a single alveolus lung-on-a-chip (ALoC) experiment. **B.** Three stitched images, representative of the entire apical channel containing AT1 (HT1+, red) and AT2 cells (HT2+, green). Nuclei are labeled with DAPI (blue). Scale bars = 500 µm. **C. left** Brightfield image of ALoCs immediately after the addition human $CD14^+$ peripheral blood-derived macrophages (macrophages) on top of the alveolar epithelial cells. **C. right** Brightfield image of ALoCs 24 hours after the addition of macrophages. **D.** Stitched image of the entire apical channel of an ALoC containing macrophages (CD14+), AT1 cells (HT1+), and AT2 cells (HT2+). Nuclei are labelled DAPI (blue). Scalebar = 500 µm.

Supplemental Figure 2. **Alveolar epithelial type I and II co-staining on ALoCs infected with *M. fortuitum*.** **A.** Stitched image of an uninfected ALoC control containing AT1 (HT1+, red) and AT2 (HT2+, green) cells. Nuclei are labelled with DAPI (blue). **B.** Confocal image of *M. fortuitum*-infected ALoCs after 24 hours of infection, highlighting AT1 cells (HT1+, purple). Adherens junctions are labelled with E-cadherin (yellow) and nuclei are labelled with DAPI. Scale bar = 500 µm. **C.** Confocal image highlighting AT2 cells (HT2+, green), adherens junctions (E-cadherin, yellow) and nuclei (DAPI, blue). Scale bar = 500 µm. **D.** Confocal image showing the merged channels of panels B and C, highlighting the co-staining of alveolar epithelial cells observed in *M. fortuitum*-infected ALoCs after 24 hours of infection. Scale bar = 500 µm.

Supplemental Figure 3. **Bulk RNA-sequencing analysis of *M. fortuitum*-infected ALoCs A.** Total raw read counts of uninfected (n=3) and *M. fortuitum*-infected (n=3) ALoCs. **B.** Pearson (average linkage) clustering of *M. fortuitum*-infected ALoCs and uninfected ALoCs.

Figure 2

Figure 3

Figure 4

Table 1. The most significant up-regulated and down-regulated GO Biological Process pathways from M. fortuitum-infected ALoCs containing macrophages

Direction	GO Biological Process	Statistic	Genes	adj.Pval
Up	Response to bacterium	7.3996	438	1.1e-09
	Humoral immune response	6.5518	138	6.0e-07
	Defense response to bacterium	6.3097	169	1.1e-06
	Antimicrobial humoral response	6.0602	65	1.9e-05
	Response to molecule of bacterial origin	5.5061	251	3.7e-05
	Chemotaxis	5.2453	444	8.2e-05
	Taxis	5.2402	445	8.2e-05
	Response to lipopolysaccharide	5.2654	240	8.2e-05
	Antimicrobial humoral immune response mediated by antimicrobial peptide	5.6504	41	1.4e-04
Down	MRNA processing	-6.8939	402	4.4e-08
	RNA splicing	-6.5273	385	2.3e-07
	Histone modification	-5.7112	429	1.6e-05
	Proteasome-mediated ubiquitin-dependent protein catabolic process	-5.5755	410	2.5e-05
	Proteasomal protein catabolic process	-5.534	470	2.5e-05
	Chromatin organization	-5.4523	478	3.2e-05
	NcRNA metabolic process	-5.3774	477	4.2e-05
	Peptidyl-lysine modification	-5.128	343	1.5e-04
	RNA splicing via transesterification reactions with bulged adenosine as nucleophile	-5.1146	265	1.5e-04
	MRNA splicing via spliceosome	-5.1146	265	1.5e-04
	RNA splicing via transesterification reactions	-5.097	269	1.5e-04

Table 2. The most significant up-regulated and down-regulated GO Molecular Function pathways from M. fortuitum-infected ALoCs containing macrophages

Direction	GO Molecular Function	Statistic	Genes	adj.Pval
Up	Transmembrane signaling receptor activity	8.0205	486	1.9e-12
	Receptor ligand activity	6.4443	275	6.9e-08
	Signaling receptor regulator activity	6.3634	295	6.9e-08
	Signaling receptor activator activity	6.3463	281	6.9e-08
	Cytokine activity	5.657	148	4.8e-06
	Channel activity	5.1005	274	3.6e-05
	Passive transmembrane transporter activity	5.1005	274	3.6e-05
	Glycosaminoglycan binding	5.1296	151	3.7e-05
	Ion channel activity	4.8906	248	8.2e-05
	Calcium ion binding	4.8317	497	8.2e-05
	G protein-coupled receptor activity	4.872	193	8.3e-05
	Gated channel activity	4.7825	188	1.1e-04
	Extracellular matrix structural constituent	4.7767	119	1.3e-04
	Cation channel activity	4.7291	184	1.3e-04
	Inorganic molecular entity transmembrane transporter activity	4.6534	455	1.3e-04
	Heparin binding	4.4603	110	4.5e-04
	Endopeptidase regulator activity	4.3733	123	6.1e-04
	Inorganic cation transmembrane transporter activity	4.2753	363	6.3e-04
Down	Catalytic activity acting on RNA	-5.0214	311	3.8e-04
	Histone binding	-4.8025	216	6.4e-04

Supplemental Table 1. Antibodies used on ALoC for immunofluorescence

Cell Type	Target	Primary Antibody	Secondary Antibody
AT1	HT1	Anti HT1-56 (Terrace Biotech, TB-29AHT1-56)	Alpaca anti-Mouse IgG1 Nano (VHH) Recombinant Secondary Antibody conjugated with Alexa Fluor 647 (Thermo Fisher, SA5-10333)
AT1	PDPN	Anti-Podoplanin/gp36 antibody (18H5) (abcam, ab10288)	Alpaca anti-Mouse IgG1 Nano (VHH) Recombinant Secondary Antibody conjugated with Alexa Fluor 647 (Thermo Fisher, SA5-10333)
AT2	HT2	Anti HT2-280 (Terrace Biotech, TB-27AHT2-280)	Goat anti-Mouse IgG2a Cross-Adsorbed Secondary Antibody, Alexa Fluor 488 (Thermo Fisher, A21131)
Macrophage	CD14	CD14 Recombinant Rabbit Monoclonal Antibody (SC69-02) (Thermo Fisher, MA5-32248)	Donkey anti-Rabbit IgG (H+L), highly cross-adsorbed Secondary Antibody with Alexa Fluor 555 Plus
Epithelial cell	E-cadherin	E-Cadherin Recombinant Rabbit Monoclonal Antibody (5H6L18) (Invitrogen, 701134)	Donkey anti-Rabbit IgG (H+L), highly cross-adsorbed Secondary Antibody with Alexa Fluor 555 Plus
n/a	Nuclei	NucBlue™ Fixed Cell ReadyProbes™ Reagent (DAPI) (Thermo Fisher, R37606)	n/a

A

Lung Alveolus-Chip Stabilization, ECM coating, Seeding, Flow	ALI

Infection

Day -3 and -1 — Day 0 — Day 3 — Day 4 — Day 5 — Day 7 — Day 9 — Day 10 — Day 11 — Day 13

Cell Expansion

Top Channel: Human Primary Alveolar Epithelial Cells (HPAEC)

Bottom Channel: Human Microvascular Endothelial Cells (HMVEC)

Flow

ALI

Stretch

Top Channel: Human Primary CD14+ Monocytes

Top Channel: M. fortuitum infection

Endpoint:
- CFU
- Fixation for IF
- Cell lysing for RNA

B HT1 HT2 DAPI

C

D HT1HT2DAPI

A

B