Fe deficiency causes transcriptional shift in roots leading to disruption of drought tolerance in soybean

Md Rokibul Hasan[1], Asha Thapa[1], Mohammad Golam Mostofa[2], Ahmad H. Kabir[1*]

[1]School of Sciences, University of Louisiana at Monroe, Monroe, LA 71209, USA

[2]Department of Chemistry, State University of New York, College of Environmental Science and Forestry, Syracuse, NY, USA

[*]e-mail: kabir@ulm.edu

Abstract

Iron (Fe) deficiency in alkaline soils, exacerbated by drought, collectively affects soybean health. However, the availability of Fe and the transcriptional changes associated with drought tolerance remain unclear in soybean. This study aimed to evaluate the physiological and transcriptional changes in Fiskeby IV, a drought-tolerant genotype that loses its resilience under simultaneous Fe deficiency and drought. In this growth incubator study, Fe deficiency and drought stress led to significant reductions in plant biomass, photosynthetic efficiency, and nutrient uptake in Fiskeby IV. Despite these disruptions, the photochemical efficiency of photosystem II remained stable, suggesting the activation of protective mechanisms to maintain essential photosynthetic functions. RNA-seq analysis highlighted a complex response, showing the upregulation of ethylene-responsive genes (*Ethylene-response sensor 2*, *Ethylene-responsive TF018*, *Ethylene-responsive TF5*) as well as the genes related to rhizosphere acidification (*ATPase 1*) and redox homeostasis (*Glutaredoxin-3*). It suggests that ethylene signaling and rhizosphere acidification may be responsive in coordinating Fe homeostasis and drought adaptation in soybean. On the flip side, combined stresses caused the downregulation of several genes related to nutrient uptake (*nicotianamine transporter YSL1*, *ammonium transporter 2*, *sulfate transporter 3.4*, and *major facilitator family protein*). In a targeted study, soybean plants supplemented with 1-aminocyclopropane-1-carboxylic acid (an ethylene precursor) showed significant improvements in morpho-physiological traits and Fe status under combined stress conditions. Furthermore, the ethylene precursor enhanced root flavonoid and rhizosphere siderophore levels, along with restoring the overall abundance of bacterial and fungal microbial cells under Fe deficiency and drought. It implies that elevated ethylene may modulate root exudates, enhance microbial recruitment, and promote a stress-resilient rhizosphere environment to help soybean plants cope with combined stresses. It underscores the role of targeting ethylene signaling that may facilitate Fe mobilization and microbial interactions to induce soybean tolerance to concurrent Fe deficiency and drought. This is the first report on the transcriptional response and Fe status requirement underlying drought tolerance in soybean, potentially guiding future strategies to improve combined stress resilience in legume crops.

Keywords: Combined stress, ethylene signaling, legume, siderophore, symbiosis.

1 Introduction

Global agricultural sustainability faces increasing challenges due to climate change, which exacerbates abiotic stressors such as drought and nutrient deficiencies. Drought stress severely limits crop productivity by disrupting physiological processes essential for growth and development (Barnabas et al., 2008). The lack of soil moisture causes drought stress, limiting the availability and uptake of essential minerals by plant roots (Li et al., 2023). In plants, drought response involves intricate regulatory networks that coordinate responses at molecular, physiological, and morphological levels (Geng et al., 2024; Seleiman et al., 2021). However, recent research highlights that the presence of additional stress factors, such as Fe deficiency, can significantly alter these mechanisms, potentially negating the benefits of drought-tolerant plant species/cultivars (Afshar et al., 2012; Elanz et al., 2011). However, Fe bioavailability in the soil is often limited, especially in alkaline or calcareous soils common to many agricultural regions (Kabir et al., 2012; 2015; Li and Lan, 2017). Fe deficiency is a common nutritional disorder affecting plants, characterized by impaired chlorophyll synthesis and reduced photosynthetic efficiency (Li and Lan, 2017). Furthermore, drought stress reduces soil moisture and restricts Fe solubility, further exacerbating Fe deficiency and negatively impacting crop yield (García et al., 2012). Beyond its direct impacts on plant health, Fe deficiency interacts with other stressors, altering gene expression patterns and metabolic pathways critical for stress adaptation (Liang, 2022). Especially in legume crops, Fe is essential for nitrogen fixation and yield productivity (Briat et al., 2015). Therefore, understanding how Fe deficiency modulates drought response in plants is pivotal for devising strategies to enhance crop resilience under multifaceted stress environments.

In plants, two primary strategies are used to absorb Fe from the soil: Strategy I and Strategy II. Strategy I is utilized by non-graminaceous (non-grass) plants, such as dicots (e.g., soybean, pea) and most non-grass monocots (Liang, 2022). Plants employing Strategy I responses to Fe deficiency enhance Fe uptake through several mechanisms to help manage and mitigate the deficiency. They increase proton extrusion via *ATPase* 1 to acidify the rhizosphere, improving Fe solubility (Kabir et al., 2012). Additionally, they boost the activity of ferric-chelate reductase enzymes to reduce Fe(III) to the more readily absorbed Fe(II) form (Kobayashi and Nishizawa, 2012; Morrissey and Guerinot, 2009). Plants also enhance the expression of Fe transporter proteins to facilitate Fe uptake into root cells (Kroh and Pilon, 2019). Furthermore, ethylene signaling influences the plant's ability to cope with low Fe availability by modulating the expression of genes involved in Fe uptake and utilization (Kabir et al., 2012; Lucena et al., 2015). In addition, microbes also play a crucial role in stress resilience in plants. Flavonoids released by plants have been identified as pre-symbiotic signaling molecules in microbial-plant symbiosis (Lidoy et al., 2023; Schliemann et al., 2008). When plants are deficient in Fe, there is typically an increase in siderophores in the rhizosphere, secreted by both bacterial and fungal communities associated with host plants in a symbiotic relationship, to chelate Fe (Jin et al., 2010; Crowley et al., 2006).

In response to drought, plants possess a diverse array of adaptive strategies aimed at maintaining cellular hydration, sustaining metabolic processes, and minimizing damage during water deficit conditions. These strategies include morphological adjustments such as root architecture

modification to enhance water uptake and leaf rolling to reduce water loss, alongside
physiological mechanisms like osmotic adjustment, where compatible solutes accumulate to
protect cellular structures (Volaire, 2018; Kumar et al., 2021; Khan et al., 2023; D'Oria et al.,
2022). On a molecular level, plants undergo dynamic changes involving the activation of stress-
responsive genes and signaling pathways. Among these, abscisic acid (ABA) signaling plays a
central role by coordinating responses to water stress; ABA mediates stomatal closure, conserves
water, and activates a cascade of stress-responsive genes to help the plant acclimate to drought
conditions (Chaves et al., 2003; Ali et al., 2020). Additionally, ethylene contributes significantly
to drought tolerance, not only by regulating drought-responsive genes but also by influencing
cell wall remodeling and antioxidant responses, which are essential for minimizing oxidative
damage under stress (Fatma et al., 2022). Ethylene's role is further highlighted in conditions of
Fe deficiency, where it activates pathways that modify root architecture to improve nutrient
acquisition, underscoring its multifaceted involvement in stress tolerance mechanisms (Sánchez-
Rodríguez et al., 2011; Barnaby et al., 2019).
107
Among staple crops, soybeans (*Glycine max*) play a crucial role in global food security and
industrial applications (Kulkarni et al., 2018). Efforts to enhance soybean resilience to drought
have yielded promising cultivars, yet their efficacy can be compromised under concurrent stress
conditions, such as Fe deficiency. While the individual roles of Fe deficiency and drought stress
in limiting plant growth are well-documented, the interplay between Fe availability and drought
tolerance remains underexplored. Particularly, the modulation of transcriptional changes and
drought tolerance due to differential Fe availability in soybeans is still not well understood.
Therefore, this study aims to unravel the physiological responses and transcriptome governing
the interplay between Fe deficiency and drought tolerance in soybean. Furthermore, we
examined the role of ethylene signaling and potential crosstalk with other hormones, providing
insights into adaptive responses that may improve soybean resilience in response to combined Fe
deficiency and drought in soybeans. These insights may optimize breeding strategies for
developing resilient crop varieties of soybeans capable of withstanding complex environmental
challenges.
122
2 Materials and methods:
2.1 Plant cultivation and growth conditions
The seeds of soybean (var. Fiskeby IV: drought tolerant) were collected from USDA-GRIN.
Before germination, seeds were surface sterilized with 4% sodium hypochlorite solution for 5
min, followed by three washes with sterile water. Subsequently, seeds were placed in a tray and
incubated at 25°C in the dark. Once uniform and healthy seedlings emerged in 3d, they were
transplanted into soil pots (500 g soil), consisting of a mixture of natural field soil and
commercial potting mix in a 1:2 ratio, for two distinct treatments: control (soil without lime, pH
~6.5) and Fe- drought+ (soil amended with 4.5 g $NaHCO_3$ and 3.0 g $CaCO_3$, pH ~7.8).
Throughout the cultivation period, 15 mM $NaHCO_3$ was added weekly to the Fe-deficient soil to
maintain its pH to 7.8 to make the Fe unavailable for plants (Kabir et al., 2012). Drought
conditions were initiated by reducing water availability to 75% compared to the controls (well-
watered) two weeks after transplanting the seedlings into the soil pots. Daily monitoring of soil
moisture content was conducted to maintain 75% and 25% soil moisture levels for control and
drought-stressed plants, respectively. Plants were cultivated following a randomized complete
block design, with 9 replications (3 plants/pot) per treatment in a growth chamber (10 h light/14

h darkness, 250 µmol m^{-2} s^{-1}) at 25°C up to 4 weeks before data measurement. In a targeted
experiment, a precursor, 1-aminocyclopropane-1-carboxylic acid (ACC) of the ethylene
synthesis (McDonnell et al., 2009), was supplemented in soil (10 mM) to validate whether it can
induce tolerance in 5-week-old soybean plants in response to combined Fe deficiency and
drought.
144
2.2 Determination of morphological and photosynthetic parameters
The height of each plant's shoot and the length of the longest root were measured using a digital
caliper. The shoots and roots were then dried for three days at 75°C in an electric oven before
their dry weights were recorded. Furthermore, chlorophyll scoring and chlorophyll fluorescence
kinetics (OJIP) measurements, including Fv/Fm (maximal photochemical efficiency of PSII) and
Pi_ABS (photosynthesis performance index) were conducted on the uppermost fully developed
leaves at three different locations using a handheld chlorophyll meter (AMTAST, United States)
and FluorPen FP 110 (Photon Systems Instruments, Czech Republic), respectively. For OJIP
analysis, the leaves were acclimated to darkness for 1 hour before data collection. Relative water
content (RWC) of leaves was measured on the fully expanded trifoliate leaves at the tip of the
main stem as previously described (Schonfeld et al., 1988). Briefly, the fresh weight (FW) of
leaves from each treatment was recorded immediately after removal. The turgid weight (TW)
was measured after soaking the leaves in distilled water in beakers for 24 h at room temperature.
Following the soaking period, the leaves were quickly and carefully blotted dry with tissue paper
to prepare for turgid weight determination. The dry weight (DW) of the leaves was obtained by
oven-drying the leaf samples for 72 h at 75ºC. RWC was calculated using the formula: RWC (%)
= (FW – DW) / (TW – DW) x 100.
162
2.3 RNA-seq analysis in roots
RNA-seq analysis was performed on root samples collected from 4-week-old plants. The
soybean roots were thoroughly cleaned by washing twice in sterile phosphate-buffered saline
with 10 seconds of vortexing, followed by two rinses with sterile water. The cleaned roots were
then ground into a powder with a pre-cooled pestle and mortar using liquid nitrogen. RNA was
extracted from the powdered material using the SV Total RNA Isolation System (Promega,
USA). For each sample, 1 µg of RNA with an RIN value above 8 was used for RNA-seq library
preparations. The libraries were created with the KAPA HiFi HotStart Library Amplification Kit
(Kapa Biosystems, USA) and sequenced on an Illumina NovaSeq 6000 instrument at the RTSF
Genomics Core at Michigan State University. A total of 92.5% of reads passed filters with a
quality score of ≥ Q30.
174
Bioinformatics analysis of raw FastQ files was performed using Partek Flow genomic analysis
software (Partek, St. Louis, MO, USA). Adaptor sequences and low-quality reads were removed
with Trimmomatic (Bolger et al., 2014) to obtain high-quality clean reads. The clean reads were
aligned to the soybean reference genome, downloaded from HISAT2 (v2.0.5) (Yates et al., 2022;
Kim et al., 2015). Gene read counts were determined using HTSeq (Anders et al., 2015).
Differentially expressed genes (DEGs) were identified using the limma v.3.19 R package
(Ritchie et al., 2015), with FPKM (fragments per kilobase of exon per million mapped
fragments) calculations. DEGs were defined as those with an adjusted P-value < 0.05 and a fold
change ≥ 2, using the false discovery rate (FDR) method. Venn diagrams and heatmaps were
created with VennDetail (Hinder et al., 2017) and pheatmap (Hu et al., 2021) in R, respectively.

Gene ontology and enrichment analyses of soybean DEGs were conducted using ShinyGO 0.81 (Ge et al., 2020).

2.4 Determination of tissue nutrient elements

Nutrient analysis was conducted on the roots and young leaves harvested from 4-week-old plants. The roots were detached, washed with running tap water, soaked in a 0.1 mM $CaSO_4$ solution for 10 min, and then rinsed with deionized water. The young leaves were washed separately with deionized water. All samples were placed in envelopes and dried in an electric oven at 75°C for three days. Elemental concentrations were measured using inductively coupled plasma mass spectrometry (ICP-MS) at the Agricultural and Environmental Services Laboratories at the University of Georgia. The concentrations were determined by comparing the intensity of the ion peaks in the spectrum to calibration standards.

2.5 Estimation of flavonoid content in roots

The total flavonoid content in soybean roots was measured as described by Piyanete et al., (2009). In brief, 0.5 mL of 70% ethanol extract from the roots was mixed with 2 mL of distilled water and 0.15 mL of a 5% $NaNO_2$ solution. After incubating for 6 min, 0.15 mL of a 10% $AlCl_3$ solution was added and left to stand for another 6 min. Then, 2 mL of a 4% NaOH solution was added, and the mixture was diluted to a final volume of 5 mL with methanol and thoroughly mixed. After a 15-min incubation, the absorbance was measured at 510 nm using a spectrophotometer against a blank. The total flavonoid concentration was expressed in milligrams per gram of dry weight, calculated using a quercetin standard curve.

2.6 Determination of siderophore in the rhizosphere

We quantified the siderophore content in the rhizosphere soil using a chrome azurol S (CAS) assay, following the procedure outlined by Himpsl and Mobley (2019). Briefly, the rhizosphere soil was collected by washing off the soil tightly attached to the root surface. The rhizosphere soil was then homogenized with 80% methanol and subsequently centrifuged for 15 min at 10,000 rpm. The resulting supernatant (500 μl) was mixed with 500 μl of CAS solution and allowed to incubate at room temperature for 5 min. The absorbance of each reaction was then measured at 630 nm against a reference containing 1 ml of CAS reagent alone. The siderophore level per gram of soil sample was calculated using the formula: % siderophore unit = [(Ar - As) / Ar] x 100, where Ar represents the absorbance at 630 nm of the reference (CAS reagent only), and As represents the absorbance at 630 nm of the sample (CAS reagent + soil sample).

2.7 Determination of relative abundance of microbial cells

Harvested roots were cleaned by vortexing in sterile phosphate-buffered saline (PBS) for 10 sec, followed by two rinses with sterile water to remove surface contaminants. Approximately 0.2 g aliquots of cleaned root samples were used for DNA extraction following the procedure outlined by Clarke (2009), along with RNase and Proteinase K treatments. The DNA samples were then quantified and diluted before performing qPCR in a CFX96 Real-Time PCR Detection System (Bio-Rad, USA) using specific primers for 16S bacteria (16S-341_F: CCTACGGGNGGCWGCAG, 16S-341_R: GACTACHVGGGTATCTAATCC) and ITS fungi (ITS1-F: TCCGTAGGTGAACCTGCGG, ITS4-R: TCCTCCGCTTATTGATATGC). The

231 soybean *Beta-tubulin* gene (Fw: GCTCCCAGCAGTACAGGACTCT, Rv:
232 TGGCATCCCACATTTGTTGA) served as the internal control for relative quantification.
233
234
235
236 **2.8 Data analysis**
237 All data were collected from three independent biological replicates for each treatment. The
238 statistical significance of each variable was assessed using Student's *t*-test and one-way ANOVA,
239 followed by Duncan's Multiple Range Test (DMRT) where applicable. Graphical presentations
240 were prepared using ggplot2 in the R program.
241
3 Results
3.1 Changes in morphological traits and photosynthetic features
At weeks 3 and 4, we measured the morphological and photosynthetic attributes to study the
responses of plants exposed to combined Fe deficiency and drought. The plants grown under Fe
deficiency and drought showed a significant decrease in root length, root dry weight, shoot
height, and shoot dry weight compared to the untreated controls (Figure 1A–1D) at both 3 weeks
and 4 weeks. We observed a significant decrease in leaf SPAD score (an indicator of chlorophyll
content) under Fe deficiency combined with drought compared to the controls (Figure 2A) at
both timepoints. Fv/Fm remained the same in the Fe-deficient and drought-stressed plants
compared to the controls (Figure 2B). Interestingly, Pi_ABS showed no significant changes at
week 3 but was significantly decreased under Fe deficiency combined with drought compared to
the controls at week 4 (Figure 2C). We did not observe any significant changes in the relative
water content (%) in the leaves of plants exposed to combined Fe deficiency and drought
compared to the control at either 3 weeks or 4 weeks (Figure 2D).
256
3.2 Elemental changes in plant tissues
The ICP-MS analysis of soybean roots and shoots under control and Fe-drought+ conditions
revealed significant differences in nutrient concentrations (Table 1). Fe and N levels significantly
decreased in both root and shoot under Fe-drought+ conditions compared to control. Zn, Mn, and
Ca levels showed no significant changes in the roots due to Fe-drought+ conditions but these
minerals significantly declined in the shoot in such conditions relative to controls (Table 1). S
and K levels significantly declined in the roots of soybean cultivated in response to combined Fe
deficiency and drought compared to controls, while these minerals showed no significant
changes in the shoot in such conditions. However, Mg levels showed no significant changes
either in the root or shoot under Fe-drought+ conditions relative to untreated controls (Table 1).
267
3.3 RNA-seq analysis of differentially expressed genes
We conducted RNA-seq analysis to examine the changes in the transcriptome of soybean roots
harvested from 4-week-old plants exposed to combined Fe deficiency and drought. Principal
component analysis (PCA) using three biological replicates revealed that the first two principal
components, PC1 and PC2, accounted for 53.4% and 23.6% of the variance, respectively (Figure
3A). The separation between PC1 and PC2 indicates distinct grouping patterns among the
samples, suggesting that treatment groups led to substantial differences in gene expression.
Furthermore, the Venn diagram highlighted the genes that were significantly altered in response
to the combined Fe deficiency and drought compared to non-treated controls (Figure 3B).

Specifically, we identified 705 upregulated and 1,297 downregulated genes differentially expressed under Fe-deficient and drought conditions compared to the control. Additionally, the MA plot showed a higher number of downregulated genes than upregulated ones in response to dual stress (Figure 3C).

3.4 Heatmap analysis of top 15 differentially upregulated and downregulated genes in roots
From the RNA-seq analysis, we present the top 15 upregulated and downregulated genes that are differentially expressed (> Log2 fold, < P value 0.05) and associated with plant growth promotion and stress responses (Figure 4A, 4B). The expression of the top 15 key genes that were significantly upregulated in the roots of soybean treated with Fe deficiency and drought relative to controls includes: *ethylene-responsive TF5*, *ABC transporter C family 10*, *ATPase 1*, dehydration-responsive binding protein 2A, *calcium-dependent protein kinase 6*, *copper ion transporter*, *sulfate transporter 1.1*, *ferredoxin*, MATE efflux family protein, *ethylene-responsive TF018*, *BZIP transcription factor ATB2*, *ABA-GTase*, *glutaredoxin-3*, *ethylene-response sensor 2*, and heavy-metal-associated protein 26 (Figure 4A). In contrast, the top 15 downregulated genes include *nicotianamine transporter YSL1*, *multi-copper oxidase type I*, *anion transporter 6*, *sulfate transporter 3.4*, heme-binding protein-related, *auxin efflux carrier component 6*, calcium transporting *ATPase 4*, jasmonic acid-amido synthesis *JAR1*, *ammonium transporter 2*, *molybdate transporter 1*, *neoxanthin synthase*, solute carrier family 40, major facilitator family protein, heavy metal transport protein, and *peroxidase 66* (Figure 4B).

3.5 Gene enrichment analysis
Gene enrichment analysis revealed the enrichment of several pathways, categorized into three groups: biological processes and molecular functions (Figure 5A-5B). The upregulated genes showed significant enrichment in major biological processes, including sequestering of actin monomers, cell redox homeostasis, negative regulation of actin filament polymerization, cellular homeostasis, and the oxidation-reduction process (Figure 5A). Among the molecular functions, oxidoreductase activity exhibited the highest fold enrichment, followed by actin monomer binding, protein disulfide isomerase activity, phosphorelay sensor kinase activity, and disulfide oxidoreductase activity (Figure 5A). Conversely, the top enriched biological processes among downregulated genes included defense response to insects, which showed the highest fold enrichment, followed by regulation of vacuole fusion, DNA replication, and cell redox homeostasis (Figure 5B). Additionally, the top enriched molecular functions among downregulated genes included beta-galactosidase activity, beta-amylase activity, galactosidase activity, amylopectin maltohydrolase activity, and oxidoreductase activity (Figure 5B).

3.6 ACC (ethylene precursor) on promoting dual stress tolerance
Given that we observed the upregulation of several ethylene-responsive genes in RNA-seq, we further investigated whether ACC (ethylene precursor) has significant biological effects on recovering dual stress in soybean cultivated for 6 weeks. In this study, the combined Fe deficiency and drought exhibited a significant decrease in chlorophyll score, plant height, plant dry weight, stem diameter, and leaf RWC (%) in soybeans relative to untreated controls (Figure 6A-6F). ACC supplementation resulted in significant improvements in these traits compared to the Fe-Drought+ conditions without ACC (Figure 6A-6F). ICP-MS analysis showed a significant decrease in leaf Fe concentration due to Fe- drought+ conditions compared to controls. However, the addition of ACC under Fe- drought+ conditions caused a significant increase in Fe levels in

the leaf compared to the plants in such conditions without ACC (Figure 6G). We found that Fe-drought+ conditions showed no significant changes in flavonoid content in the roots of soybean relative to controls. However, the addition of ACC under Fe-drought+ caused a significant increase in flavonoid content compared to those plants exposed to combined Fe deficiency and drought without ACC (Figure 6H). Interestingly, siderophore levels in the rhizosphere of soybean exposed to Fe-drought+ conditions showed a significant increase relative to controls (Figure 6I). Soybean plants exposed to combined Fe deficiency and drought supplemented with ACC showed a significant increase in rhizosphere siderophore compared to those plants exposed to Fe-drought+ conditions without ACC supplementation (Figure 6I).

We further determined the relative of 16S and ITS microbial cells in roots by DNA-based qPCR. We found that combined Fe deficiency and drought caused a significant decline in 16S bacterial abundance in roots relative to controls. However, the addition of ACC in Fe-Drought+ conditions showed a significant increase in 16S, which is similar to controls (Figure 7A). Similarly, the abundance of ITS fungal cells significantly decreased in the roots due to Fe-Drought+ conditions in contrast to untreated controls. Interestingly, ITS fungal abundance showed a significant increase when ACC was added to soil exposed to Fe deficiency and drought. Also, the increase in ITS abundance was relatively higher than the plants cultivated in control conditions (Figure 7B).

4 Discussion
This study reveals that drought tolerant cultivar (Fiskeby IV) exposed to combined Fe deficiency and drought stress experience significant reductions in morphological and photosynthetic efficiency. Plants are constantly exposed to combined abiotic stresses, such as mineral deficiency and drought, in their natural environments. These simultaneous challenges can lead to an intensified effect, making it harder for plants to adapt and survive (Teshome et al., 2020). This is the first growth incubator study to reveal transcriptional changes, with many genes related to growth and stress responses being differentially expressed. We also observed the requirement for ethylene-mediated Fe homeostasis and possible indications of symbiosis that may be necessary for combined stress tolerance in soybean.

The significant reduction in plant biomass subjected to Fe deficiency and drought stress, as observed in our study, highlights the detrimental impact of these combined abiotic stresses on the drought-tolerant soybean genotype. These morphological changes are likely due to impaired cellular processes and metabolic activities resulting from inadequate Fe availability and water stress. Fe is a critical micronutrient involved in various physiological processes, including chlorophyll synthesis and electron transport in photosynthesis (Samira et al., 2013). The observed decrease in chlorophyll content and photosynthetic performance index under Fe deficiency and drought stress conditions underscores the compromised photosynthetic capacity of the plants. Interestingly, the maximum quantum efficiency of photosystem II (Fv/Fm) remained unaffected under Fe deficiency and drought stress suggesting that while the overall photosynthetic apparatus is impacted, the core photochemical efficiency of PSII remains intact. It is possible that the non-photochemical quenching helps dissipate excess light energy as heat to prevent damage to PSII reaction centers (Ruban et al., 2016; Demmig-Adams et al., 2014). Furthermore, enhanced scavenging of reactive oxygen species (ROS) may also contribute to the stability of PSII by neutralizing ROS generated under stress (Foyer & Noctor, 2011). We

369 observed the upregulation of glutaredoxin-3 in the roots of soybean. Glutaredoxin-3 is involved
370 in the assembly and maintenance of Fe-S clusters, which are crucial for the function of many
371 enzymes, including those involved in ROS detoxification and other metabolic processes
372 (Couturier et al., 2015).
373
The elemental analysis of soybean tissues revealed significant alterations in nutrient
concentrations under Fe deficiency and drought stress. The marked decrease in Fe and N levels
in both roots and shoots highlights the impaired nutrient uptake and transport mechanisms under
these stress conditions. Also, the decline in these nutrients in the roots indicates a potential trade-
off between root and shoot nutrient allocation to mitigate stress effects. However, the unchanged
levels of Zn, Mn, and Ca in the roots, despite significant declines in the shoots under Fe
deficiency and drought stress, suggest differential regulation of nutrient allocation and
homeostasis between root and shoot tissues. The stability of these elements in the roots might be
attributed to their relatively lower mobility and the plant's efforts to prioritize root functionality
under stress. Furthermore, the significant reduction in S and K levels in the roots, coupled with
no significant changes in the shoots, further underscores the differential nutrient management
strategies adopted by the plants under Fe deficiency and drought stress. Sulfur is vital for the
synthesis of amino acids and proteins (Ren et al., 2022; Künstler et al., 2020). Furthermore,
potassium regulates stomatal exposure, helping to reduce water loss through transpiration and
maintain turgor pressure under drought conditions (Wang et al., 2013). K also activates
numerous enzymes, including those involved in photosynthesis, protein synthesis, and
carbohydrate metabolism, helping plants sustain metabolic functions under stress (Ho et al.,
2022; Jin et al., 2011). Additionally, K is critical for maintaining ion homeostasis and enhancing
the plant's ability to uptake other essential nutrients (Lian et al., 2023). Hence, the retention of K
levels in the shoot may therefore reflect the soybean plants' attempt to optimize water usage and
nutrient uptake under combined Fe deficiency and drought stress. Furthermore, the retention of
sulfur levels in the shoot aligns with the gene enrichment in oxidoreductase activity, indicating a
significant role in redox reactions and sulfur metabolism in soybeans exposed to the combined
effects of Fe deficiency and drought. Several studies demonstrated that redox reactions and
sulfur metabolism play a critical role in responses to abiotic stress plants (Mittler, 2002; Kopriva
& Rennenberg, 2004).
400
The transcriptomic analysis reveals a complex network of gene expression changes in response
to combined Fe deficiency and drought stress in soybean roots. We found that upregulated genes
are primarily associated with stress responses and adaptive mechanisms. In Strategy I plants,
ATPase 1, a plasma membrane proton pump, plays a significant role in Fe uptake (Zhang et al.,
2017). *ATPase 1* contributes to this by creating a proton gradient across the root plasma
membrane, which is essential for the acidification of the rhizosphere (Michalak et al., 2022).
This acidification helps solubilize Fe^{3+} making it more accessible for uptake by the plant (Kabir
et al., 2012). Thus, the upregulation of *ATPase 1* in roots suggests an enhanced effort to regulate
rhizosphere acidification which helps plants mobilize Fe under combined Fe deficiency and
drought in soybean. In addition, the induction of ferredoxin among the upregulated genes also
points to the importance of redox homeostasis in stress adaptation. Ferredoxin is involved in
maintaining cellular redox homeostasis by ROS, particularly under drought, salinity, and high
light-intensity stresses in plants (Huang et al., 2020; Lehtimäki et al., 2010). Our gene
enrichment analysis also indicates an upregulation of cell redox homeostasis, highlighting the

importance of redox regulation required for soybean plants to cope with combined Fe deficiency and drought. Furthermore, soybean plants exposed to concurrent Fe deficiency and drought showed upregulation of the *ABA-Gtase* gene. It serves as an ABA receptor and is also a key component of the ABA signaling pathway in plants (Xu et al., 2002). Although ABA is a well-known stress hormone, particularly involved in drought responses, it can also induce the expression of Fe-deficiency-responsive genes, suggesting its role in regulating Fe uptake and transport under Fe-limiting conditions (Lei et al., 2014; García et al., 2011). This crosstalk between ABA and Fe deficiency is particularly important in drought conditions, where the plant's need for Fe might increase due to compromised root functionality.

Interestingly, we found the upregulation of several ethylene-responsive genes, such as ethylene-responsive transcription factors (TF5 and TF018) and dehydration-responsive binding protein 2A, in response to combined Fe deficiency and drought in soybean. This may imply the activation of stress-related signaling pathways and the role of ethylene signaling in soybean's adaptive response to combined Fe deficiency and drought. Under Fe deficiency, ethylene-responsive genes are induced to enhance the plant's ability to acquire and utilize Fe and other nutrients efficiently (Müller and Munné-Bosch, 2015; Li and Lan, 2017; Wu et al., 2011). This is correlated with the upregulation of sulfate transporter 1.1, heavy-metal-associated protein 26, and copper ion transporter, indicating their efforts in maintaining cellular ion balance under stress conditions. Interestingly, ethylene may interact with Fe deficiency signaling pathways to enhance ATPase 1 activity (Waters et al., 2007), a finding that aligns with our RNA-seq results. Additionally, ethylene biosynthesis may induce drought-responsive genes that help plants conserve water and maintain cellular homeostasis (Mohorović et al., 2023; Arraes et al., 2015). Naing et al., (2022) showed that ethylene-insensitive mutants exhibited increased sensitivity to drought stress, indicating that ethylene-responsive genes contribute to drought adaptation by modulating stomatal conductance and enhancing root growth. Ethylene signaling may facilitate ROS scavenging enzymes, helping mitigate oxidative stress, which is common under both Fe deficiency and drought conditions (Zhang et al., 2016; Kabir et al., 2012), and aligns with the upregulation of glutaredoxin-3 in the roots of soybean. In most cases, ethylene, whose production increases under nutrient deficiency in genotypically efficient genotypes, acts as an activator of stress responses in plants (Dubois et al., 2018). Precious studies indicating the upregulation of *1-aminocyclopropane-1-carboxylic acid synthase (ACS)*, and *1-aminocyclopropane-1-carboxylic acid oxidade* genes involved in ethylene synthesis further support the increased ethylene production as part of the adaptive responses under Fe deficiency in plants (García et al., 2015; Sauter et al., 2013). Thus, it appears that ethylene-related or Fe-uptake genes could be potential targets for improving stress tolerance in soybean exposed to Fe deficiency and drought.

To further elucidate the role of ethylene, we conducted a targeted study using an ethylene precursor (ACC) to determine if it could eliminate the impairment in drought-tolerant Fiskeby IV exposed to combined Fe deficiency and drought. In this study, supplementation with ACC led to significant improvements in these morpho-physiological traits, suggesting that ethylene signaling helps restore key growth parameters along with leaf Fe status disrupted by combined stress conditions. Interestingly, we observed a correlation between root flavonoid content and rhizosphere siderophore levels induced by the presence of an ACC, further underscoring the significance of microbial associations in soybean subjected to combined Fe deficiency and drought. While 'Fe- drought+' stress conditions alone did not affect root flavonoid content, ACC

supplementation significantly increased flavonoid levels. This enhancement in flavonoids might
contribute to improved stress resilience in soybean as flavonoids are known to play roles in
symbiotic associations (Wang et al., 2022). Flavonoids play an essential role in rhizobium-
legume symbiosis as chemoattractants and *Nod* gene inducers, and they are also believed to
stimulate fungal spore germination and hyphal growth (Abdel-Lateif et al., 2012; Hassan et al.,
2012). Similarly, siderophore levels in the rhizosphere were elevated under 'Fe-drought stress',
and ACC further boosted siderophore production. This indicates that ACC not only improves
plant tolerance to concurrent Fe deficiency and drought but also potentially aids in Fe acquisition
by increasing siderophore-mediated Fe mobilization in the rhizosphere. Microbial siderophores
are compounds secreted by both bacterial and fungal microbial communities that bind and
solubilize Fe, making them more accessible for uptake by plants (Singh et al., 2022). These
siderophores can enhance plant Fe acquisition, particularly in Fe-deficient soils, by facilitating
the transfer of Fe from the soil to plant roots (Kabir and Bennetzen, 2024; Radzki et al., 2013). In
this study, the decline in microbial cells due to Fe deficiency and drought could be attributed to
the combined stress effects on soil and plant environments. Fe deficiency reduces the availability
of essential nutrients and impacts plant root exudation, which serves as a primary energy source
for microbes (Chen and Liu, 2024; Pascale et al., 2020). Additionally, drought conditions limit
soil moisture, further restricting microbial activity and growth due to reduced mobility and
access to nutrients. Furthermore, drought conditions may limit soil moisture, further restricting
microbial activity and growth due to reduced mobility and access to nutrients (Qu et al., 2024).
Interestingly, we observed the restoration of the overall 16S bacterial and ITS fungal abundance
when ACC was augmented in soybeans exposed to dual stresses, compared to plants exposed to
Fe-drought+ conditions. Thus, it appears that ethylene signaling may indirectly promote
microbial symbiosis, thereby recovering Fe status and drought tolerance in Fiskeby IV exposed
to combined Fe deficiency and drought. Previous studies demonstrated that ethylene plays a
complex yet crucial role in enhancing microbial symbiosis in plants by modulating defense
responses and facilitating beneficial interactions (Shekhawat et al., 2022; Ravanbakhsh et al.,
2018). In legume-rhizobia symbiosis, ethylene levels influence root nodule formation essential
for nitrogen fixation (Guinel, 2015). Similarly, in mycorrhizal associations, ethylene can
modulate root architecture and colonization (Khatabi et al., 2012). Also, plants exposed to abiotic
stresses often produce elevated levels of ethylene to influence root architecture and the
composition of root exudates (Glick, 2014). These exudates, which include organic acids, amino
acids, and secondary metabolites, act as signaling molecules and attractants for beneficial
microbes, such as rhizobacteria and mycorrhizal fungi (Vacheron et al., 2013). Thus, our findings
imply that the induction of ethylene-responsive genes may act as a central regulator, integrating
signals from both Fe deficiency and drought stress to orchestrate an adaptive response in
soybean. This study may inspire future research on the composition of microbial dynamics,
including rhizobia, to uncover the interactions between ethylene signaling and microbiome to
possess plant growth-promoting and stress resilience functions.
501
Our RNA-seq analysis exhibited that several downregulated genes are associated with various
physiological processes, including nutrient transport, hormone signaling, and stress responses
due to combined Fe deficiency and drought. Fe deficiency and/or drought can cause significant
nutritional imbalances in plants, disrupting essential metabolic processes and leading to stunted
growth and poor health (Kumari et al., 2022). In this study, the downregulation of nicotianamine
transporter *YSL1* and anion transporter 6 reflects a possible disruption in nutrient uptake and

translocation in soybean (Fiskeby IV) exposed to Fe deficiency and drought. Furthermore, the
decreased expression of jasmonic acid-amido synthetase *JAR1* and auxin efflux carrier
component 6 suggests an alteration in hormonal balance, which could be one of the factors
disrupting the drought tolerance ability of the genotype Fiskeby IV to combat dual stresses on
soybean. Decreased *JAR1* expression can impair drought resilience by weakening the jasmonic
acid signaling pathways that regulate stomatal closure and stress-responsive gene activation (Ali
et al., 2020). Another study demonstrates that over-expression of *JAR1*) enhances drought
tolerance in *Arabidopsis thaliana* by increasing jasmonyl-L-isoleucine production, which leads
to better water retention (Mahmud et al., 2022). Also, JA has been shown to promote root hair
development, which helps increase the plant's ability to acquire Fe (Kobayashi et al., 2016.
Kobayashi and Nishizawa, 2012). Although Fe is not a direct cofactor for the enzymes specific to
jasmonic acid signaling, it is crucial for the biosynthesis of JA, as it is required for the
functioning of lipoxygenases and other enzymes that convert fatty acids into jasmonic acid
(Singh et al., 2022; Oliw 2022). In contrast, jasmonate-deficient *cpm2* mutant revealed that
jasmonates inhibit the expression of several Fe deficiency-inducible genes associated with
Feuptake in rice (Kobayashi et al., 2016). Particularly, auxin plays a critical role in mediating
adaptive responses to improve Fe acquisition and drought tolerance in plants (Liu et al., 2024;
Iqbal et al., 2022; Kabir et al., 2012). In this study, we also found the downregulation of
neoxanthin synthase and solute carrier family 40 highlights disruptions in carotenoid
biosynthesis and solute transport, which is consistent with the decreased photosynthetic
efficiency and nutrient uptake in Fiskeby IV exposed to concurrent Fe deficiency and drought.
Under abiotic stress, such as drought or nutrient deficiency, plants often experience disruptions
in chlorophyll and carotenoid biosynthesis resulting in decreased energy production and plant
growth (Muhammad et al., 2021; Dong et al., 2007). Carotenoid synthesis may also be affected,
impacting plant protection against oxidative damage and influencing overall stress tolerance
(Giani and Martínez-Espinosa 2020). Thus, phytohormone and other downregulated genes can be
targeted for genetic engineering or breeding strategies aimed at enhancing stress resilience in
soybean.

5 Conclusion
Fe deficiency, intensified by drought in alkaline soils, compromises soybean health by reducing
biomass, photosynthetic efficiency, and nutrient uptake. Despite these stresses, Fiskeby IV
maintained stable photosystem II efficiency, indicating protective adaptations. RNA-seq revealed
upregulation of ethylene-responsive genes, rhizosphere acidification, and redox homeostasis
factors, suggesting these mechanisms aid in coordinating Fe and drought resilience (Figure 8).
Conversely, key nutrient transporters were downregulated under combined stress.
Supplementation with an ethylene precursor improved physiological traits, Fe status, root
flavonoid, and siderophore levels. Supplementation with an ethylene precursor, improved
morpho-physiological traits and Fe status under combined stress, enhanced root flavonoid
content, and increased rhizosphere siderophore levels and microbial abundance. These findings
reveal the potential of targeting ethylene signaling and microbial interactions to enhance soybean
tolerance to concurrent Fe deficiency and drought. Future research should focus on these
mechanisms to develop strategies that improve stress resilience and sustainability in soybean
crops.

Data availability statement

Raw RNA-seq data was submitted to NCBI under the following accession number: PRJNA1140285.

Author contributions
MRH cultivated the plants, conducted most of the experiments, performed RNA-seq analysis, and prepared the draft manuscript. AT extracted RNA from root samples and assisted with plant phenotyping. MGM helped with data interpretations and critically revised the manuscript. AHK conceived the idea, supervised the whole work, and revised the manuscript.

Acknowledgments
This research was supported by a startup grant (5SFAES-293007) and Elizabeth and Hayden Cutler's Endowed Professorship in Biotechnology from the University of Louisiana at Monroe.

Conflict of Interest Statement
We have no conflict of interest.

References
Abdel-Lateif, K., Bogusz, D., and Hocher, V. (2012). The role of flavonoids in the establishment of plant roots endosymbioses with arbuscular mycorrhiza fungi, rhizobia and Frankia bacteria. *Plant Signal Behav.* 7(6), 636–641.

Afshar, R. M., Hadi, H., and Pirzad, A. (2012). Effect of Nano-Fe foliar application on qualitative and quantitative characteristics of cowpea, under end season drought stress. *Int. Res. J. Appl. Basic Sci.* 3, 1709–1717.

Ali, M. S., and Baek, K. H. (2020). Jasmonic Acid Signaling Pathway in Response to Abiotic Stresses in Plants. *Int. Journal Mol. Sci.* 21(2), 621.

Ali, S., Hayat, K., Iqbal, A., and Xie, L. (2020) Implications of Abscisic Acid in the Drought Stress Tolerance of Plants. *Agronomy.* 10, 1323.

Arraes, F. B., Beneventi, M. A., Lisei de Sa, M. E., Paixao, J. F., Albuquerque, E. V., Marin, S. R., Purgatto, E., Nepomuceno, A. L., and Grossi-de-Sa, M. F. (2015). Implications of ethylene biosynthesis and signaling in soybean drought stress tolerance. *BMC Plant Biol.* 15, 213.

Barnabas, B., Jäger, K., and Fehér, A. (2008). The effect of drought and heat stress on reproductive processes in cereals. *Plant Cell EnvFe.* 31, 11–38.

Barnaby, J. Y., Bauerle, T. L., Edwards, N. T., & DeLucia, E. H. (2019). Ethylene synthesis and downstream signaling interact to regulate root growth under iron-deficient conditions in soybean. Frontiers in Plant Science, 10, 783.

Briat, J. F., Dubos, C., & Gaymard, F. (2015). Iron nutrition, biomass production, and plant product quality. *Trends Plant Sci.* 20(1), 33–40.

Chaves M.M., Maroco J.P., and Pereira J.S. (2003) Understanding plant responses to drought- from genes to the whole plant. *Funct. Plant Biol.* 30, 239–264.

Chen, L., & Liu, Y. (2024). The Function of Root Exudates in the Root Colonization by Beneficial Soil Rhizobacteria. Biology, 13(2), 95.

Couturier, J., Przybyla-Toscano, J., Roret, T., Didierjean, C., and Rouhier, N. (2015). The roles of glutaredoxins ligating Fe-S clusters: Sensing, transfer or repair functions? *Biochimica et Biophysica Act.* 1853(6), 1513–1527.

Crowley D.E. Microbial siderophores in the plant rhizosphere. In: Barton L.L., Abadia J., editors. Fe Nutrition in Plants and Rhizospheric Microorganisms. Springer; Dordrecht, The Netherlands: 2006. pp. 169–198.

Demmig-Adams, B., Stewart, J. J., Burch, T. A., & Adams, W. W., 3rd (2014). Insights from Placing Photosynthetic Light Harvesting into Context. *J. Physical Chemistry Lett.* 5(16), 2880–2889.

Dong, H., Deng, Y., Mu, J., Lu, Q., Wang, Y., Xu, Y., Chu, C., Chong, K., Lu, C., and Zuo, J. (2007). The Arabidopsis Spontaneous Cell Death1 gene, encoding a zeta-carotene desaturase essential for carotenoid biosynthesis, is involved in chloroplast development, photoprotection and retrograde signalling. *Cell Res.* 17(5), 458–470.

D'Oria, A., Courbet, G., Billiot, B., Jing, L., Pluchon, S., Arkoun, M., Maillard, A., Roux, C. P., Trouverie, J., Etienne, P., Diquélou, S., and Ourry, A. (2022). Drought specifically downregulates mineral nutrition: Plant ionomic content and associated gene expression. *Plant Direct* 6(8), e402.

Dubois, M., Van den Broeck, L., and Inzé, D. (2018). The Pivotal Role of Ethylene in Plant Growth. *Trends Plant Sci.* 23(4), 311–323.

Elanz, E., Mohsen, R., and Ahmed, B. (2011). Influence of salt stress on cations concentration quality and quantity of sunflower cultivars. *J. Food Agric. EnvFe.* 2, 469–476.

Fatma, M., Asgher, M., Iqbal, N., Rasheed, F., Sehar, Z., Sofo, A., and Khan, N. A. (2022). Ethylene Signaling under Stressful EnvFements: Analyzing Collaborative Knowledge. Plants (Basel, Switzerland), 11(17), 2211.

Fatma, M., Iqbal, N., Gautam, H., Sehar, Z., Sofo, A., D'Ippolito, I., & Khan, N. A. (2021). Ethylene and Sulfur Coordinately Modulate the Antioxidant System and ABA Accumulation in Mustard Plants under Salt Stress. Plants (Basel, Switzerland), 10(1), 180.

García, M. J., Romera, F. J., Lucena, C., Alcántara, E., and Pérez-Vicente, R. (2015). Ethylene and the Regulation of Physiological and Morphological Responses to Nutrient Deficiencies. Plant physiology, 169(1), 51–60.

García, M. J., Suárez, V., Romera, F. J., Alcántara, E., & Pérez-Vicente, R. (2011). A new model involving ethylene, nitric oxide and Fe to explain the regulation of Fe-acquisition genes in Strategy I plants. *Plant Physiol. Biochem.* 49(5), 537–544.

Ge, S.X., Jung, D., Yao, R. (2020). Bioinformatics 36:2628–2629.

Geng, A., Lian, W., Wang, Y., Liu, M., Zhang, Y., Wang, X., and Chen, G. (2024). Molecular Mechanisms and Regulatory Pathways Underlying Drought Stress Response in Rice. *Int. J Mol. Sci.* 25(2), 1185.

Giani, M., and Martínez-Espinosa, R. M. (2020). Carotenoids as a Protection Mechanism against Oxidative Stress in *Haloferax mediterranei*. *Antioxidants* 9(11), 1060.

Glick B. R. (2014). Bacteria with ACC deaminase can promote plant growth and help to feed the world. Microbiological research, 169(1), 30–39.

Guinel, F.C. (2015) Ethylene, a Hormone at the Center-Stage of Nodulation. *Front. Plant Sci.* 6, 1121.

Hassan, S., and Mathesius, U. (2012). The role of flavonoids in root-rhizosphere signalling: opportunities and challenges for improving plant-microbe interactions. *J. Exp. Bot.* 63(9), 3429–3444.

Ho, L. H., Rode, R., Siegel, M., Reinhardt, F., Neuhaus, H. E., Yvin, J. C., Pluchon, S., Hosseini, S. A., & Pommerrenig, B. (2020). Potassium application boosts photosynthesis and sorbitol biosynthesis and accelerates cold acclimation of common Plantain (*Plantago major* L.). *Plants* (Basel, Switzerland), 9(10), 1259.

Huang, H. E., Ho, M. H., Chang, H., Chao, H. Y., and Ger, M. J. (2020). Overexpression of plant ferredoxin-like protein promotes salinity tolerance in rice (*Oryza sativa*). *Plant Physiol. Biochem.* 155, 136–146.

Iqbal, S., Wang, X., Mubeen, I., Kamran, M., Kanwal, I., Díaz, G. A., Abbas, A., Parveen, A., Atiq, M. N., Alshaya, H., Zin El-Abedin, T. K., and Fahad, S. (2022). Phytohormones Trigger Drought Tolerance in Crop Plants: Outlook and Future Perspectives. *Front. Plant Sci.* 12, 799318.

Jin, C. W., Li, G. X., Yu, X. H., and Zheng, S. J. (2010). Plant Fe status affects the composition of siderophore-secreting microbes in the rhizosphere. *Ann Bot.* 105(5), 835–841.

Jin, S. H., Huang, J. Q., Li, X. Q., Zheng, B. S., Wu, J. S., Wang, Z. J., Liu, G. H., & Chen, M. (2011). Effects of potassium supply on limitations of photosynthesis by mesophyll diffusion conductance in Carya cathayensis. *Tree physiology*, 31(10), 1142–1151.

Kabir, A. H., and Bennetzen, J. L. (2024). Molecular insights into the mutualism that induces Fe deficiency tolerance in sorghum inoculated with *Trichoderma harzianum. Microbiol. Res.* 281, 127630.

Kabir, A. H., Paltridge, N. G., Able, A. J., Paull, J. G., and Stangoulis, J. C. (2012). Natural variation for Fe-efficiency is associated with upregulation of Strategy I mechanisms and enhanced citrate and ethylene synthesis in Pisum sativum L. Planta, 235(6), 1409–1419.

Khan, M. I. R., Nazir, F., Maheshwari, C., Chopra, P., Chhillar, H., and Sreenivasulu, N. (2023). Mineral nutrients in plants under changing envFements: A road to future food and nutrition security. *Plant Genome* 16(4), e20362.

Khatabi, B., Molitor, A., Lindermayr, C., Pfiffi, S., Durner, J., von Wettstein, D., Kogel, K. H., and Schäfer, P. (2012). Ethylene supports colonization of plant roots by the mutualistic fungus *Piriformospora indica. PloS one* 7(4), e35502.

Kobayashi, T., and Nishizawa, N. K. (2012). Fe uptake, translocation, and regulation in higher plants. *Ann. Rev. Plant Biol.* 63, 131–152.

Kobayashi, T., Itai, R. N., Senoura, T., Oikawa, T., Ishimaru, Y., Ueda, M., Nakanishi, H., & Nishizawa, N. K. (2016). Jasmonate signaling is activated in the very early stages of iron deficiency responses in rice roots. *Plant Mol. Biol.* 91(4-5), 533–547.

Kobayashi, T., Itai, R. N., Senoura, T., Oikawa, T., Ishimaru, Y., Ueda, M., Nakanishi, H., and Nishizawa, N. K. (2016). Jasmonate signaling is activated in the very early stages of Fe deficiency responses in rice roots. *Plant Mol. Biol.* 91(4-5), 533–547.

Kopriva, S., & Rennenberg, H. (2004). Control of sulphate assimilation and glutathione synthesis: interaction with N and C metabolism. Journal of Experimental Botany, 55(404), 1831-1842.

Kroh, G.E., and Pilon, M. (2019). Connecting the negatives and positives of plant Fe homeostasis. *New Phytol.* 223(3), 1052-1055.

Kulkarni, K. P., Tayade, R., Asekova, S., Song, J. T., Shannon, J. G., and Lee, J. D. (2018). Harnessing the Potential of Forage Legumes, Alfalfa, Soybean, and Cowpea for Sustainable Agriculture and Global Food Security. *Front. Plant Sci.* 9, 1314.

Kumar, M., Kumar Patel, M., Kumar, N., Bajpai, A. B., and Siddique, K. H. M. (2021). Metabolomics and Molecular Approaches Reveal Drought Stress Tolerance in Plants. *Int. J. Mol. Sci.* 22(17), 9108.

Kumari, V. V., Banerjee, P., Verma, V. C., Sukumaran, S., Chandran, M. A. S., Gopinath, K. A., Venkatesh, G., Yadav, S. K., Singh, V. K., and Awasthi, N. K. (2022). Plant Nutrition: An Effective Way to Alleviate Abiotic Stress in Agricultural Crops. *Int. J. Mol. Sci.* 23(15), 8519.

Künstler, A., Gullner, G., Ádám, A. L., Kolozsváriné Nagy, J., and Király, L. (2020). The Versatile Roles of Sulfur-Containing Biomolecules in Plant Defense-A Road to Disease Resistance. *Plants* 9(12), 1705.

Lehtimäki, N., Lintala, M., Allahverdiyeva, Y., Aro, E. M., and Mulo, P. (2010). Drought stress-induced upregulation of components involved in ferredoxin-dependent cyclic electron transfer. *J Plant Phys.* 167(12), 1018–1022.

Lei, G. J., Zhu, X. F., Wang, Z. W., Dong, F., Dong, N. Y., & Zheng, S. J. (2014). Abscisic acid alleviates iron deficiency by promoting root iron reutilization and transport from root to shoot in Arabidopsis. *Plant Cell Environ.* 37(4), 852–863.

Li, S., Yang, L., Huang, X., Zou, Z., Zhang, M., Guo, W., Addo-Danso, S. D., and Zhou, L. (2023). Mineral Nutrient Uptake, Accumulation, and Distribution in *Cunninghamia lanceolata* in Response to Drought Stress. *Plants* 12(11), 2140.

Li, W., and Lan, P. (2017). The Understanding of the Plant Fe Deficiency Responses in Strategy I Plants and the Role of Ethylene in This Process by Omic Approaches. *Front Plant Sci.* 8, 40.

Lian, W., Geng, A., Wang, Y., Liu, M., Zhang, Y., Wang, X., & Chen, G. (2023). The Molecular Mechanism of Potassium Absorption, Transport, and Utilization in Rice. *Int. J. Mol Sci.* 24(23), 16682.

Liang G. (2022). Fe uptake, signaling, and sensing in plants. *Plant Comm.* 3(5), 100349.

Lidoy J, Berrio E, García M, España-Luque L, Pozo MJ, López-Ráez JA. (2023). Flavonoids promote *Rhizophagus irregularis* spore germination and tomato root colonization: A target for sustainable agriculture. *Front Plant Sci.* 5, 1094194.

Liu, L., Yahaya, B. S., Li, J., and Wu, F. (2024). Enigmatic role of auxin response factors in plant growth and stress tolerance. *Front Plant Sci.* 15, 1398818.

Lucena, C., Romera, F. J., García, M. J., Alcántara, E., and Pérez-Vicente, R. (2015). Ethylene Participates in the Regulation of Fe Deficiency Responses in Strategy I Plants and in Rice. *Front Plant Sci.* 6, 1056.

Mahmud, S., Ullah, C., Kortz, A., Bhattacharyya, S., Yu, P., Gershenzon, J., and Vothknecht, U. C. (2022). Constitutive expression of Jasmonate resistant 1 induces molecular changes that prime the plants to better withstand drought. *Plant Cell Env.* 45(10), 2906–2922.

McDonnell, L., Plett, J. M., Andersson-Gunnerås, S., Kozela, C., Dugardeyn, J., Van Der Straeten, D., Glick, B. R., Sundberg, B., and Regan, S. (2009). Ethylene levels are regulated by a plant encoded 1-aminocyclopropane-1-carboxylic acid deaminase. *Physiol Plant.* 136(1), 94-109.

Michalak, A., Wdowikowska, A., and Janicka, M. (2022). Plant Plasma Membrane Proton Pump: One Protein with Multiple Functions. *Cells* 11(24), 4052.

Mittler, R. (2002). Oxidative stress, antioxidants and stress tolerance. Trends in Plant Science, 7(9), 405-410.

Mohorović, P., Geldhof, B., Holsteens, K., Rinia, M., Ceusters, J., and Van de Poel, B. (2023). Effect of ethylene pretreatment on tomato plant responses to salt, drought, and waterlogging stress. *Plant Direct* 7(11), e548.

Morrissey, J., and Guerinot, M. L. (2009). Fe uptake and transport in plants: the good, the bad, and the ionome. *Chem Rev.* 109(10), 4553–4567.

Muhammad, I., Shalmani, A., Ali, M., Yang, Q. H., Ahmad, H., and Li, F. B. (2021). Mechanisms Regulating the Dynamics of Photosynthesis Under Abiotic Stresses. *Front Plant Sci.* 11, 615942.

Müller, M., and Munné-Bosch, S. (2015). Ethylene Response Factors: A Key Regulatory Hub in Hormone and Stress Signaling. *Plant Physiol.* 169(1), 32–41.

Naing, A. H., Campol, J. R., Kang, H., Xu, J., Chung, M. Y., and Kim, C. K. (2022). Role of Ethylene Biosynthesis Genes in the Regulation of Salt Stress and Drought Stress Tolerance in Petunia. *Front Plant Sci.* 13, 844449.

Oliw E. H. (2022). Fe and manganese lipoxygenases of plant pathogenic fungi and their role in biosynthesis of jasmonates. *Arch. Biochem. Biophy.* 722, 109169.

Pascale A, Proietti S, Pantelides IS, Stringlis IA. (2022) Modulation of the Root Microbiome by Plant Molecules: The Basis for Targeted Disease Suppression and Plant Growth Promotion. Front Plant Sci. 10:1741.

Qu Q, Wang Z, Gan Q, Liu R, Xu H. (2024) Impact of drought on soil microbial biomass and extracellular enzyme activity [published correction appears in Front Plant Sci. 15, 1393574.

Radzki, W., Gutierrez Mañero, F. J., Algar, E., Lucas García, J. A., García-Villaraco, A., and Ramos Solano, B. (2013). Bacterial siderophores efficiently provide Fe to Fe-starved tomato plants in hydroponics culture. *Antonie van Leeuwenhoek* 104(3), 321–330.

Ravanbakhsh, M., Sasidharan, R., Voesenek, L. A. C. J., Kowalchuk, G. A., & Jousset, A. (2018). Microbial modulation of plant ethylene signaling: ecological and evolutionary consequences. *Microbiome* 6(1), 52.

Ren, Z., Wang, R.Y., Huang, X.Y., and Wang Y. (2022) Sulfur Compounds in Regulation of Stomatal Movement. *Front. Plant Sci.* 13, 846518.

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., and Smyth, G. K. (2015). limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* 43(7), e47.

Ruban AV. (2016) Nonphotochemical chlorophyll fluorescence quenching: mechanism and effectiveness in protecting plants from photodamage. *Plant Physiol.* 170(4), 1903-1916.

Samira, R., Stallmann, A., Massenburg, L.N., Long, T.A. (2013) Feing out the issues: integrated approaches to understanding Fe homeostasis in plants. *Plant Sci.* 210, 250-259.

Sánchez-Rodríguez, E., Del Mar Rubio-Wilhelmi, M., Blasco, B., Leyva, R., Romero, L., & Ruiz, J. M. (2011). Antioxidant response in lemon plants subjected to boron and drought stress. Plant Physiology and Biochemistry, 49(7), 774–782.

Sauter M, Moffatt B, Saechao MC, Hell R, Wirtz M (2013) Methionine salvage and S-adenosylmethionine: essential links between sulfur, ethylene and polyamine biosynthesis. *Biochem J.* 451, 145–154.

Schliemann, W., Ammer, C., and Strack, D. (2008). Metabolite profiling of mycorrhizal roots of *Medicago truncatula*. *Phytochem.* 69(1), 112–146.

Seleiman, M. F., Al-Suhaibani, N., Ali, N., Akmal, M., Alotaibi, M., Refay, Y., Dindaroglu, T., Abdul-Wajid, H. H., and Battaglia, M. L. (2021). Drought Stress Impacts on Plants and Different Approaches to Alleviate Its Adverse Effects. *Plants* 10(2), 259.

Shekhawat, K., Fröhlich, K., García-Ramírez, G. X., Trapp, M. A., and Hirt, H. (2022). Ethylene: A Master Regulator of Plant-Microbe Interactions under Abiotic Stresses. *Cells* 12(1), 31.

Singh, P., Arif, Y., Miszczuk, E., Bajguz, A., and Hayat, S. (2022). Specific Roles of Lipoxygenases in Development and Responses to Stress in Plants. *Plants* 11(7), 979.

Singh, P., Chauhan, P. K., Upadhyay, S. K., Singh, R. K., Dwivedi, P., Wang, J., Jain, D., and Jiang, M. (2022). Mechanistic Insights and Potential Use of Siderophores Producing Microbes in Rhizosphere for Mitigation of Stress in Plants Grown in Degraded Land. *Front. Microbiol.* 13, 898979.

Teshome, D. T., Zharare, G. E., and Naidoo, S. (2020). The Threat of the Combined Effect of Biotic and Abiotic Stress Factors in Forestry Under a Changing Climate. *Front. Plant Sci.* 11, 601009.

Vacheron, J., Desbrosses, G., Bouffaud, M. L., Touraine, B., Moënne-Loccoz, Y., Muller, D., Legendre, L., Wisniewski-Dyé, F., & Prigent-Combaret, C. (2013). Plant growth-promoting rhizobacteria and root system functioning. Frontiers in plant science, 4, 356.

Volaire F. (2018). A unified framework of plant adaptive strategies to drought: Crossing scales and disciplines. *Global Change Biol.* 24(7), 2929–2938.

Wang M, Zheng Q, Shen Q, Guo S. The critical role of potassium in plant stress response. (2013) *Int J Mol Sci.* 14(4), 7370-7390.

Wang, L., Chen, M., Lam, P. Y., Dini-Andreote, F., Dai, L., and Wei, Z. (2022). Multifaceted roles of flavonoids mediating plant-microbe interactions. *Microbiome* 10(1), 233.

Waters, B. M., Lucena, C., Romera, F. J., Jester, G. G., Wynn, A. N., Rojas, C. L., Alcántara, E., and Pérez-Vicente, R. (2007). Ethylene involvement in the regulation of the H(+)-ATPase *CsHA1* gene and of the new isolated ferric reductase *CsFRO1* and Fe transporter *CsIRT1* genes in cucumber plants. *Plant Physiol. Biochem.* 45(5), 293–301.

Wu, J., Wang, C., Zheng, L., Wang, L., Chen, Y., Whelan, J., and Shou, H. (2011). Ethylene is involved in the regulation of Fe homeostasis by regulating the expression of Fe-acquisition-related genes in Oryza sativa. *J. Exp. Bot.* 62(2), 667–674.

Xu ZJ, Nakajima M, Suzuki Y, Yamaguchi I. (2002). Cloning and characterization of the abscisic acid-specific glucosyltransferase gene from adzuki bean seedlings. *Plant Physiol.* 129(3), 1285-1295.

Zhang, H., Li, A., Zhang, Z., Huang, Z., Lu, P., Zhang, D., Liu, X., Zhang, Z. F., and Huang, R. (2016). Ethylene Response Factor *TERF1*, Regulated by ETHYLENE-INSENSITIVE3-like Factors, Functions in Reactive Oxygen Species (ROS) Scavenging in Tobacco (Nicotiana tabacum L.). *Sci. Rep.* 6, 29948.

Zhang, J., Wei, J., Li, D., Kong, X., Rengel, Z., Chen, L., Yang, Y., Cui, X., and Chen, Q. (2017). The Role of the Plasma Membrane H+-ATPase in Plant Responses to Aluminum Toxicity. *Front. Plant Sci.* 8, 1757.

Table 1. ICP-MS analysis of nutrients in the root and shoot of soybean grown in control and Fe-drought+ conditions. Different letters indicate statistically significant differences between the treatment groups, as determined by a *t*-test at the $P < 0.05$ level. The data represents the means ± SD of three independent biological samples ($n = 3$).

Elements	Root		Shoot	
	Control	Fe- Drought+	Control	Fe- Drought+
Fe (ppm)	460.8±32.5a	339.9±28.8b	95.3±8.6a	66.7±4.9b
Zn (ppm)	34.4±1.36a	34.2±1.64a	40.5±6.0a	28.6±3.1b

Mn (ppm)	24.1±4.0a	22.1±2.9a	49.2±8.5a	25.9±4.4b
N (%)	2.48±0.19a	1.77±0.19b	3.79±0.01a	3.45±0.07b
S (%)	0.23±0.02a	0.14±0.04b	0.25±0.01a	0.25±0.03a
K (%)	0.95±0.11a	0.70±0.05b	3.50±0.13a	3.51±0.21a
Ca (%)	0.64±0.06a	0.65±0.02a	1.02±0.17a	0.56±0.02b
Mg (%)	0.59±0.13a	0.40±0.07a	0.36±0.03a	0.40±0.04a

Figure 1. Changes in root length (A), root dry weight (B), shoot height (C), and shoot dry weight (D) in soybean cultivated for 3-week and 4-week in 'control' and 'Fe-drought+' conditions. Statistical significance between treatments was determined using a two-tailed t-test (* indicates significant differences, $p < 0.05$) The data represents the means ± SD of three independent biological samples ($n = 3$).

829
Figure 2. Changes in chlorophyll score (A), Fv/Fm: maximal quantum yield of PS II (B), and
Pi_ABS: photosynthesis performance index (C) in the leaves of soybean cultivated for 3-week
and 4-week in 'control' and 'Fe-drought+' conditions. Statistical significance between treatments
was determined using a two-tailed t-test (**$p < 0.01$, *$p < 0.05$, ***$p < 0.001$, and "ns" indicates
no significant difference). The data represents the means ± SD of three independent biological
samples ($n = 3$).
836

837
Figure 3. Analysis of RNA-sequencing data in roots of soybean cultivated for 4-week for
differentially expressed genes (DEGs): principal component analysis of DEGs (A), vein diagram
of upregulated and downregulated genes (>Log2 fold, $P < 0.05$) grown in 'Fe-drought+' vs.
'control' (B) and MA plot of genes expressed in 'Fe-drought+' vs. 'control' conditions (C).

857 **Figure 4.** Heatmap of the top 15 differentially expressed genes (Log2 fold change > 1, $P < 0.05$)
858 in the roots of Fiskeby IV cultivated for 4-week to combined Fe deficiency and drought relative
859 to untreated controls. (A) Top 15 upregulated genes. (B) Top 15 downregulated genes.

866
867 **Figure 5.** Gene enrichment analysis for the upregulated (A) and downregulated genes (B) in
868 roots of Fiskeby IV cultivated in 'Fe-drought+' and 'control' conditions for biological processes,
869 and molecular functions. The size and color of the data points indicate the number of genes
870 involved and the significance level, respectively, with larger points and deeper colors
871 representing more substantial enrichment and higher significance.
872
873

Figure 6. Effect of ACC on plant phenotype (A), chlorophyll score (B), plant height (C), plant dry weight (D), stem diameter (E), relative water content (F), leaf Fe (G), flavonoid content (H) and rhizosphere siderophore (I) in soybean exposed to Fe-drought cultivated for 4 weeks. Different letters above the bar indicate significant differences between treatment groups at a *P* <0.05 level, where applicable. The data represents the means ± SD of three independent biological samples (*n* = 3).

888
889 **Figure 7.** Effect of ACC on the relative abundance of 16S (bacteria) and ITS (fungi) cells in
890 roots of Fiskeby IV exposed to combined Fe deficiency and drought for 4 weeks. Different
891 letters above the bar indicate significant differences between treatment groups at a P <0.05 level,
892 where applicable. The data represents the means ± SD of three independent biological samples (*n*
893 = 3).

894
895

Figure 8. Physiological and transcriptional responses of soybean plants exposed to concurrent Fe deficiency and drought. The combined Fe deficiency and drought caused the upregulation of genes related to ethylene signaling (*Ethylene-response sensor 2*, *Ethylene-responsive TF018*, *Ethylene-responsive TF5*), proton extrusion (*ATPase 1*) and redox homeostasis (*Glutaredoxin-3*) while the expression of genes associated with peroxidase activity (*Peroxidase 66*), phytohormone synthesis (*Auxin efflux carrier component-6*, *Jasmonic acid-amido synthetase*) and nutrient transport (*Sulfate transporter 3.4*, *Ammonium transporter 2*) showed downregulation. 'Fe-Drought+' conditions caused a decline in mineral elements in both root and leaf. Interestingly, ethylene precursor (ACC) improved Fe homeostasis leading to drought resilience either directly or indirectly through root flavonoid and rhizosphere siderophore to induce drought tolerance.