Fe deficiency causes transcriptional shift in roots leading to disruption of drought tolerance in soybean

Md Rokibul Hasan[1], Asha Thapa[1], Mohammad Golam Mostofa[2], Ahmad H. Kabir[1*]

[1]School of Sciences, University of Louisiana at Monroe, Monroe, LA 71209, USA

[2]Department of Chemistry, State University of New York, College of Environmental Science and Forestry, Syracuse, NY, USA

[*]e-mail: kabir@ulm.edu

Abstract

Iron (Fe) deficiency in alkaline soils, exacerbated by drought, collectively affects soybean health. This study aimed to evaluate the physiological and transcriptional changes in Fiskeby IV, a drought-tolerant genotype that loses its tolerance when exposed to simultaneous Fe deficiency and drought. In this growth incubator study, Fe deficiency and drought stress resulted in substantial reductions in plant biomass, photosynthetic efficiency, and nutrient uptake in Fiskeby IV. Despite these disruptions, the photochemical efficiency of photosystem II remained stable, suggesting the activation of protective mechanisms to maintain essential photosynthetic functions. RNA-seq analysis highlighted a complex response, showing the upregulation of ethylene-responsive genes (*Ethylene-response sensor 2*, *Ethylene-responsive TF018*, *Ethylene-responsive TF5*) as well as the genes related to rhizosphere acidification (*ATPase 1*) and redox homeostasis (*Glutaredoxin-3*). It suggests that ethylene signaling and rhizosphere acidification may be responsive in coordinating Fe homeostasis and drought adaptation in soybean. On the flip side, combined stresses caused the downregulation of several genes related to nutrient uptake (*nicotianamine transporter YSL1*, *ammonium transporter 2*, *sulfate transporter 3.4*, and *major facilitator family protein*). In a targeted study, supplementation with 1-aminocyclopropane-1-carboxylic acid (ACC), an ethylene precursor, led to substantial improvements in morpho-physiological traits and Fe status under combined stress conditions. This ACC treatment enhanced root flavonoid content and rhizosphere siderophore levels accompanied by restoration of 16S and ITS microbial community under Fe deficiency and drought. It underscores the potential of targeting ethylene signaling that may facilitate Fe mobilization and microbial interactions to enhance soybean tolerance to concurrent Fe deficiency and drought. This is the first report on the transcriptional response and requirement of Fe status underlying drought tolerance, potentially guiding future strategies for improving combined stress resilience in soybean.

Keywords: Combined stress, ethylene signaling, legume, siderophore, symbiosis.

1 Introduction

Global agricultural sustainability faces increasing challenges due to climate change, which exacerbates abiotic stressors such as drought and nutrient deficiencies. Drought stress severely limits crop productivity by disrupting physiological processes essential for growth and development (Barnabas et al., 2008). Drought stress reduces soil moisture, limiting the availability and uptake of essential minerals by plant roots (Li et al., 2023). In plants, drought response involves intricate regulatory networks that coordinate responses at molecular, physiological, and morphological levels (Geng et al., 2024; Seleiman et al., 2021). However, recent research highlights that the presence of additional stress factors, such as Fe (Fe) deficiency, can significantly alter these mechanisms, potentially negating the benefits of drought-tolerant plant species/cultivars (Afshar et al., 2012; Elanz et al., 2011). Fe deficiency is a common nutritional disorder affecting plants, characterized by impaired chlorophyll synthesis and reduced photosynthetic efficiency (Li and Lan, 2017). Beyond its direct impacts on plant health, Fe deficiency interacts with other stressors, altering gene expression patterns and metabolic pathways critical for stress adaptation (Liang, 2022). Understanding how Fe deficiency modulates drought tolerance mechanisms in plants is therefore pivotal for devising strategies to enhance crop resilience under multifaceted stress environments.

Plants employing Strategy I responses to Fe deficiency enhance Fe uptake through several mechanisms to help manage and mitigate the deficiency. They increase proton extrusion via *ATPase* 1 to acidify the rhizosphere, improving Fe solubility (Kabir et al. 2012). Additionally, they boost the activity of ferric-chelate reductase enzymes to reduce Fe(III) to the more readily absorbed Fe(II) form (Morrissey and Guerinot, 2009). Plants also enhance the expression of Fe transporter proteins to facilitate Fe uptake into root cells (Kroh and Pilon, 2019). In addition, drought tolerance in plants encompasses a diverse array of adaptive strategies aimed at maintaining cellular hydration, sustaining metabolic processes, and minimizing damage during water deficit conditions (Volaire, 2018; Kumar et al., 2021). Furthermore, plants exhibit dynamic changes at the molecular level, involving the activation of stress-responsive genes and signaling pathways such as abscisic acid (ABA) signaling, which coordinates responses to water stress by regulating stomatal closure and promoting stress-responsive gene expression (Chaves et al., 2003; Ali et al., 2020). Also, ethylene plays a pivotal role in mediating plant responses to both drought stress and Fe deficiency, highlighting its multifaceted involvement in stress tolerance mechanisms (Fatma et al., 2022). Ethylene signaling influences the plant's ability to cope with low Fe availability by modulating the expression of genes involved in Fe uptake and utilization (Kabir et al., 2012; Lucena et al., 2015). Furthermore, plants employ a variety of strategies to cope with drought stress, among which the Strategy I response in dicot plants plays a crucial role in ensuring adequate mineral nutrition, particularly under water-deficit conditions (Khan et al., 2023; D'Oria et al., 2022). Strategy I refers to the mechanism by which plants acquire Fe from the soil through the reduction of Fe(III) to Fe(II) at the root surface, facilitated by proton extrusion and subsequent uptake into root cells via specific transporters (Kabir et al., 2012; Kobayashi and Nishizawa, 2012). These adjustments are crucial for ensuring sufficient Fe acquisition despite limited water availability, thereby safeguarding essential metabolic functions and overall plant growth and productivity. Flavonoids released by plants have been identified as pre-symbiotic signaling molecules in microbial-plant symbiosis (Lidoy et al., 2023; Schliemann et al., 2008). When plants are deficient in Fe, there is

typically an increase in siderophores in the rhizosphere, secreted by both bacterial and fungal communities associated with host plants in a symbiotic relationship, to chelate Fe (Jin et al., 2010; Crowley et al., 2006). However, the role of microbial association and siderophores in the rhizosphere of legume crops exposed to combined abiotic stresses is understudied.

Among staple crops, soybeans (*Glycine max*) play a crucial role in global food security and industrial applications (Kulkarni et al., 2018). Efforts to enhance soybean resilience to drought have yielded promising cultivars, yet their efficacy can be compromised under concurrent stress conditions, such as Fe deficiency. However, the modulation of drought tolerance due to differential Fe availability in soybeans is still not well understood. Therefore, this study aims to unravel the physiological responses and transcriptome governing the interplay between Fe deficiency and drought tolerance in soybean. Furthermore, we studied the potential role of an ethylene precursor in mitigating the detrimental impacts of Fe deficiency and drought in soybeans. These insights may optimize breeding strategies for developing resilient crop varieties of soybeans capable of withstanding complex environmental challenges.

2 Materials and methods:
2.1 Plant cultivation and growth conditions
The seeds of soybean (var. Fiskeby IV: drought tolerant) were collected from USDA-GRIN. Before germination, seeds were surface sterilized with 4% sodium hypochlorite solution for 5 min, followed by three washes with sterile water. Subsequently, seeds were placed in a tray and incubated at 25°C in the dark. Once uniform and healthy seedlings emerged in 3d, they were transplanted into soil pots (500 g soil), consisting of a mixture of natural field soil and commercial potting mix in a 1:2 ratio, for two distinct treatments: control (soil without lime, pH ~6.5) and Fe-drought+ (soil amended with 4.5 g $NaHCO_3$ and 3.0 g $CaCO_3$, pH ~7.8). Throughout the cultivation period, 15 mM $NaHCO_3$ was added weekly to the Fe-deficient soil to maintain its pH to 7.8 to make the Fe unavailable for plants (Kabir et al., 2012). Drought conditions were initiated by reducing water availability to 75% compared to the controls (well-watered) two weeks after transplanting the seedlings into the soil pots. Daily monitoring of soil moisture content was conducted to maintain 75% and 25% soil moisture levels for control and drought-stressed plants, respectively. Plants were cultivated following a randomized complete block design, with 9 replications (3 plants/pot) per treatment in a growth chamber (10 h light/14 h darkness, 250 μmol $m^{-2} s^{-1}$) at 25°C for 5 weeks before data measurement. In a targeted experiment, a precursor, 1-aminocyclopropane-1-carboxylic acid (ACC) of the ethylene synthesis (McDonnell et al., 2009), was supplemented in soil (10 mM) to validate whether it can induce tolerance in 6-week-old plants in response to combined Fe deficiency and drought.

2.2 Determination of morphological and photosynthetic parameters
The height of each plant's shoot and longest root were measured. The shoots and roots were then dried for three days at 70°C in an electric oven before their dry weights were recorded. Furthermore, chlorophyll scoring and chlorophyll fluorescence kinetics (OJIP) measurements, including Fv/Fm (maximal photochemical efficiency of PSII) and Pi_ABS (photosynthesis performance index) were conducted on the uppermost fully developed leaves at three different locations using a handheld chlorophyll meter (AMTAST, United States) and FluorPen FP 110 (Photon Systems Instruments, Czech Republic), respectively. For OJIP analysis, the leaves were acclimated to darkness for 1 hour before data collection. Relative water content (RWC) of leaves

was measured on the fully expanded trifoliate leaves at the tip of the main stem as previously described (Schonfeld et al., 1988). Briefly, the fresh weight (FW) of leaves from each treatment was recorded immediately after removal. The turgid weight (TW) was measured after soaking the leaves in distilled water in beakers for 24 h at room temperature. Following the soaking period, the leaves were quickly and carefully blotted dry with tissue paper to prepare for turgid weight determination. The dry weight (DW) of the leaves was obtained by oven-drying the leaf samples for 72 h at 70ºC. RWC was calculated using the formula: RWC (%) = (FW – DW) / (TW – DW) x 100.

2.3 RNA-seq analysis in roots

RNA-seq analysis was performed on root samples. The soybean roots were thoroughly cleaned by washing twice in sterile phosphate-buffered saline with 10 seconds of vortexing, followed by two rinses with sterile water. The cleaned roots were then ground into a powder with a pre-cooled pestle and mortar using liquid nitrogen. RNA was extracted from the powdered material using the SV Total RNA Isolation System (Promega, USA). For each sample, 1 µg of RNA with an RIN value above 8 was used for RNA-seq library preparations. The libraries were created with the KAPA HiFi HotStart Library Amplification Kit (Kapa Biosystems, USA) and sequenced on an Illumina NovaSeq 6000 instrument at the RTSF Genomics Core at Michigan State University. A total of 92.5% of reads passed filters with a quality score of $\geq$ Q30.

Bioinformatics analysis of raw FastQ files was performed using Partek Flow genomic analysis software (Partek, St. Louis, MO, USA). Adaptor sequences and low-quality reads were removed with Trimmomatic (Bolger et al., 2014) to obtain high-quality clean reads. The clean reads were aligned to the soybean reference genome, downloaded from HISAT2 (v2.0.5) (Yates et al., 2022; Kim et al., 2015). Gene read counts were determined using HTSeq (Anders et al., 2015). Differentially expressed genes (DEGs) were identified using the limma v.3.19 R package (Ritchie et al., 2015), with FPKM (fragments per kilobase of exon per million mapped fragments) calculations. DEGs were defined as those with an adjusted P-value < 0.05 and a fold change $\geq$ 2, using the false discovery rate (FDR) method. Venn diagrams and heatmaps were created with VennDetail (Hinder et al., 2017) and pheatmap (Hu et al., 2021) in R, respectively. Gene ontology and enrichment analyses of soybean DEGs were conducted using the SoyBase website developed by the USDA-ARS SoyBase and Legume Clade Database group at Iowa State University (Ames, IA, United States).

2.4 Determination of tissue nutrient elements.

Nutrient analysis was conducted on the roots and young leaves. The roots were detached, washed with running tap water, soaked in a 0.1 mM $CaSO_4$ solution for 10 min, and then rinsed with deionized water. The young leaves were washed separately with deionized water. All samples were placed in envelopes and dried in an electric oven at 75°C for three days. Elemental concentrations were measured using inductively coupled plasma mass spectrometry (ICP-MS) at the Agricultural and Environmental Services Laboratories at the University of Georgia. The concentrations were determined by comparing the intensity of the ion peaks in the spectrum to calibration standards.

2.5 Estimation of flavonoid content in roots

The total flavonoid content in soybean roots was measured as described by Piyanete et al., (2009). In brief, 0.5 mL of 70% ethanol extract from the roots was mixed with 2 mL of distilled water and

0.15 mL of a 5% NaNO$_2$ solution. After incubating for 6 min, 0.15 mL of a 10% AlCl$_3$ solution
was added and left to stand for another 6 min. Then, 2 mL of a 4% NaOH solution was added, and
the mixture was diluted to a final volume of 5 mL with methanol and thoroughly mixed. After a
15-minute incubation period, the absorbance was measured at 510 nm using a spectrophotometer
against a blank. The total flavonoid concentration was expressed in milligrams per gram of dry
weight, calculated using a quercetin standard curve.
191
2.6 Determination of siderophore in the rhizosphere
We quantified the siderophore content in the rhizosphere soil using a chrome azurol S (CAS) assay,
following the procedure outlined by Himpsl and Mobley (2019). In summary, rhizosphere soil was
homogenized with 80% methanol and subsequently centrifuged for 15 min at 10,000 rpm. The
resulting supernatant (500 µl) was mixed with 500 µl of CAS solution and allowed to incubate at
room temperature for 5 min. The absorbance of each reaction was then measured at 630 nm against
a reference containing 1 ml of CAS reagent alone. The siderophore level per gram of soil sample
was calculated using the formula: % siderophore unit = [(Ar - As) / Ar] x 100, where Ar represents
the absorbance at 630 nm of the reference (CAS reagent only), and As represents the absorbance
at 630 nm of the sample (CAS reagent + soil sample).
202
2.7 Determination of relative abundance of microbial community
Harvested roots were cleaned by vortexing in sterile phosphate-buffered saline (PBS) for 10 sec,
followed by two rinses with sterile water to remove surface contaminants. Approximately 0.2 g
aliquots of cleaned root samples were used for DNA extraction following the procedure outlined
by Clarke (2009), along with RNase and Proteinase K treatments. The DNA samples were then
quantified and diluted before performing qPCR in a CFX96 Real-Time PCR Detection System
(Bio-Rad, USA) using specific primers for bacterial (16S-341_F: CCTACGGGNGGCWGCAG,
16S-341_R: GACTACHVGGGTATCTAATCC)   and   ITS   fungal   (ITS1-F:
TCCGTAGGTGAACCTGCGG, ITS4-R: TCCTCCGCTTATTGATATGC) community. The
soybean *Tubulin* gene (Fw: GCTCCCAGCAGTACAGGACTCT,   Rv:
TGGCATCCCACATTTGTTGA) served as the internal control for relative quantification.
214
2.8 Data analysis
216 All data were collected from three independent biological replicates for each treatment. The
217 statistical significance of each variable was assessed through Student's t-test. Graphical
218 presentations were prepared using ggplot2 in the R program.
219
3 Results
3.1 Changes in morphological and traits and photosynthetic features
222 The plants grown under Fe deficiency and drought showed a significant decrease in root length,
223 root dry, shoot height and shoot dry weight compared to the untreated controls (Figure 1A-1D).
224 We observed a significant decrease in chlorophyll scores and Pi_ABS under Fe deficiency
225 combined with drought compared to the controls (Figure 2A, 2C). However, Fv/Fm remained the
226 same in the Fe-deficient and drought-stressed plants compared to the controls (Figure 2B).
227
3.2 Elemental changes in plant tissues
229 The ICP-MS analysis of soybean roots and shoots under control and Fe-drought+ conditions
230 revealed significant differences in nutrient concentrations (Table 1). Fe and N levels significantly

decreased in both root and shoot under Fe-drought+ conditions compared to control. Zn, Mn and Ca levels showed no significant changes in the roots due to Fe-drought+ conditions but these minerals significantly declined in the shoot in such conditions relative to controls (Table 1). S and K levels significantly declined in the roots of soybean cultivated in response to combined Fe deficiency and drought compared to controls, while these minerals showed no significant changes in the shoot in such conditions. However, Mg levels showed no significant changes either in the root or shoot under Fe-drought+ conditions relative to untreated controls (Table 1).

3.3 RNA-seq analysis of differentially expressed genes

We conducted RNA-seq analysis to examine the changes in the transcriptome of soybean roots exposed to a combination of Fe (Fe) deficiency and drought. Principal component analysis (PCA) using three biological replicates revealed that the first two principal components, PC1 and PC2, accounted for 53.4% and 23.6% of the variance, respectively (Figure 3A). The PCA results indicated significant differences among the sample groups. Furthermore, the Venn diagram highlighted the genes that were significantly altered in response to the combined Fe deficiency and drought compared to non-treated controls (Figure 3B). Specifically, we identified 705 upregulated and 1,297 downregulated genes differentially expressed under Fe-deficient and drought conditions compared to the control. Additionally, the MA plot showed a higher number of downregulated genes than upregulated ones in response to dual stress (Figure 3C).

3.4 Heatmap analysis of top 15 differentially upregulated and downregulated genes in roots

From the RNA-seq analysis, we present the top 15 upregulated and downregulated genes that are differentially expressed (> Log2 fold, < P value 0.05) and associated with plant growth promotion and stress responses (Figure 4A, 4B). The expression of the top 15 key genes that were significantly upregulated in the roots of soybean treated with Fe deficiency and drought relative to controls includes: ethylene-responsive TF5, ABC transporter C family 10, ATPase 1, dehydration-responsive binding protein 2A, calcium-dependent protein kinase 6, copper ion transporter, sulfate transporter 1.1, ferredoxin, MATE efflux family protein, ethylene-responsive TF018, BZIP transcription factor ATB2, ABA-GTase, glutaredoxin-3, ethylene-response sensor 2, and heavy-metal-associated protein 26 (Figure 4A, Table 2). In contrast, the top 15 downregulated genes include nicotianamine transporter YSL1, multi-copper oxidase type I, anion transporter 6, sulfate transporter 3.4, heme-binding protein-related, auxin efflux carrier component 6, calcium transporting *ATPase 4*, jasmonic acid-amido synthesis *JAR1*, ammonium transporter 2, molybdate transporter 1, neoxanthin synthase, solute carrier family 40, major facilitator family protein, heavy metal transport protein, and peroxidase 66 (Figure 4B, Table 2).

3.5 Gene enrichment analysis

Gene enrichment analysis revealed the enrichment of several pathways, categorized into three groups: biological processes, cellular components, and molecular functions (Figure 5A-5B). The biological process category showed significant upregulation in translation, signal transduction, biosynthesis, cell cycle, transport, embryo development, flower development, cell differentiation, and cellular metabolic processes. In contrast, the cellular component category indicated significant upregulation of the nucleus, cytoplasm, mitochondria, membranes, plasma membrane, cytoskeleton, endoplasmic reticulum, and Golgi apparatus. Additionally, the molecular function category showed significant upregulation of protein binding, nucleic acid binding, RNA binding,

catalytic activity, sequence-specific DNA binding transcription factor activity, nucleotide binding, kinase activity, and transferase activity - specifically transferring phosphorus-containing groups. We also observed downregulation in several pathways compared to the control. In the biological process category, there was significant downregulation in signal transduction, translation, lipid metabolic processes, flower development, cell death, transport, carbohydrate metabolic processes, cell differentiation, protein modification processes, and reproductive structure development. The cellular component category showed significant downregulation of the nucleus, mitochondria, plasma membrane, thylakoid, cytoplasm, extracellular region, golgi apparatus, cytosol, and vacuole. Furthermore, the molecular function category exhibited significant downregulation of protein binding, catalytic activity, nucleic acid binding, RNA binding, DNA binding, sequence-specific DNA binding transcription factor activity, nucleotide binding, kinase activity, and chromatin binding compared to controls under Fe deficiency and drought conditions.

3.6 ACC (ethylene precursor) on promoting dual stress tolerance

Given that we observed the upregulation of several ethylene-responsive genes in RNA-seq, we further investigated whether ACC (an ethylene precursor) has significant biological effects on recovering dual stress in soybean cultivated for 6 weeks. In this study, the combined Fe deficiency and drought exhibited a significant decrease in chlorophyll score, plant height, plant dry weight, stem diameter and leaf RWC (%) in soybeans relative to untreated controls (Figure 6A-6F). ACC supplementation resulted in significant improvements in these traits compared to the Fe-Drought+ conditions without ACC (Figure 6A-6F). ICP-MS analysis showed a significant decrease in leaf Fe concentration due to Fe- drought+ conditions compared to controls. However, the addition of ACC under Fe- drought+ conditions caused a significant increase in Fe levels in the leaf compared to the plants in such conditions without ACC (Figure 6G). We found that Fe-drought+ conditions showed no significant changes in flavonoid content in the roots of soybean relative to controls. However, the addition of ACC under Fe-drought+ caused a significant increase in flavonoid content compared to those plants exposed to combined Fe deficiency and drought without ACC (Figure 6H). Interestingly, siderophore levels in the rhizosphere of soybean exposed to Fe-drought+ conditions showed a significant increase relative to controls (Figure 6I). Soybean plants exposed to combined Fe deficiency and drought supplemented with ACC showed a significant increase in rhizosphere siderophore compared to those plants exposed to Fe-drought+ conditions without ACC supplementation (Figure 6I).

We further determined the relative of 16S and ITS microbial communities in roots by DNA-based qPCR. We found that combined Fe deficiency and drought caused a significant decline in 16S bacterial abundance in roots relative to controls. However, the addition of ACC in Fe-Drought+ conditions showed a significant increase in 16S, which is similar to controls (Figure 7A). Similarly, the relative abundance of ITS fungal community significantly decreased in the roots due to Fe-Drought+ conditions in contrast to untreated controls. Interestingly, ITS fungal abundance showed a significant increase when ACC was added to soil exposed to Fe deficiency and drought. Also, the increase in ITS abundance was relatively higher than the plants cultivated in control conditions (Figure7B).

4 Discussion

This study reveals that drought tolerant cultivar (Fiskeby IV) exposed to combined Fe deficiency and drought stress experience significant reductions in morphological and photosynthetic

efficiency. Plants are constantly exposed to combined abiotic stresses, such as mineral deficiency
and drought, in their natural environments. These simultaneous challenges can lead to an
intensified effect, making it harder for plants to adapt and survive (Teshome et al., 2020). This is
the first growth incubator study to reveal transcriptional changes, with many genes related to
growth and stress responses being differentially expressed. We also observed the requirement for
ethylene-mediated Fe homeostasis and possible indications of symbiosis that may be necessary for
combined stress tolerance in soybean.
329
Morphological and physiological responses to combined stresses
The significant reduction in plant biomass subjected to Fe deficiency and drought stress, as
observed in our study, highlights the detrimental impact of these combined abiotic stresses on
soybean growth. These morphological changes are likely due to impaired cellular processes and
metabolic activities resulting from inadequate Fe availability and water stress. Fe is a critical
micronutrient involved in various physiological processes, including chlorophyll synthesis and
electron transport in photosynthesis (Samira et al., 2013). The observed decrease in chlorophyll
content and photosynthetic performance index under Fe deficiency and drought stress conditions
underscores the compromised photosynthetic capacity of the plants. Interestingly, the maximum
quantum efficiency of photosystem II (Fv/Fm) remained unaffected under Fe deficiency and
drought stress suggesting that while the overall photosynthetic apparatus is impacted, the core
photochemical efficiency of PSII remains intact. It is possible that the plants activate protective
mechanisms to maintain the functional integrity of PSII under combined stress conditions, even
though the overall photosynthetic capacity is reduced.
344
The elemental analysis of soybean tissues revealed significant alterations in nutrient concentrations
under Fe deficiency and drought stress. The marked decrease in Fe and N levels in both roots and
shoots highlights the impaired nutrient uptake and transport mechanisms under these stress
conditions. The unchanged levels of Zn, Mn, and Ca in the roots, despite significant declines in
the shoots under Fe deficiency and drought stress, suggest differential regulation of nutrient
allocation and homeostasis between root and shoot tissues. The stability of these elements in the
roots might be attributed to their relatively lower mobility and the plant's efforts to prioritize root
functionality under stress. Furthermore, the significant reduction in S and K levels in the roots,
coupled with no significant changes in the shoots, further underscores the differential nutrient
management strategies adopted by the plants under Fe deficiency and drought stress. Sulfur is vital
for the synthesis of amino acids and proteins, while potassium plays a crucial role in osmotic
regulation and enzyme activation (Ren et al., 2022; Künstler et al., 2020). The observed decline in
these nutrients in the roots indicates a potential trade-off between root and shoot nutrient allocation
to mitigate stress effects.
359
Transcriptomic responses to combined Fe deficiency and drought stress
The transcriptomic analysis reveals a complex network of gene expression changes in response to
combined Fe deficiency and drought stress in soybean roots. We found that upregulated genes are
primarily associated with stress responses and adaptive mechanisms. Particularly, the upregulation
of ethylene-responsive transcription factors (TF5 and TF018) and dehydration-responsive binding
protein 2A highlights the activation of stress-related signaling pathways. The induction of several
ethylene-responsive genes suggests a connection of ethylene signaling as part of the adaptive
responses to combined Fe deficiency and drought stress in soybean. Ethylene is a critical plant

hormone involved in regulating various physiological and developmental processes, including
responses to biotic and abiotic stresses (Chen et al., 2021; Dubois et al., 2018). Ethylene has been
shown to be involved in the regulation of Fe homeostasis (Wu et al., 2011). Under Fe deficiency,
ethylene-responsive genes are induced to enhance the plant's ability to acquire and utilize Fe and
other nutrients efficiently (Müller and Munné-Bosch, 2015; Li and Lan, 2017). This is correlated
with the upregulation of sulfate transporter 1.1, heavy-metal-associated protein 26, and copper ion
transporter, indicating their efforts in maintaining cellular ion balance under stress conditions.
Ethylene has also been shown to interact with Fe deficiency signaling pathways, enhancing the
expression of Fe acquisition genes, including *ATPase 1* (Waters et al., 2007). In Strategy I plants,
ATPase 1, a plasma membrane proton pump, plays a significant role in Fe uptake (Zhang et al.,
2017). *ATPase 1* contributes to this by creating a proton gradient across the root plasma membrane,
which is essential for the acidification of the rhizosphere (Michalak et al., 2022). This acidification
helps solubilize Fe^{3+} making it more accessible for uptake by the plant (Kabir et al., 2012). Thus,
the upregulation of *ATPase 1* in roots suggests an enhanced effort to regulate rhizosphere
acidification which helps plants mobilize Fe under combined Fe deficiency and drought in
soybean.

Furthermore, studies suggest that ethylene biosynthesis is often upregulated, leading to changes in
the expression of drought-responsive genes that help plants conserve water and maintain cellular
homeostasis (Mohorović et al., 2023; Arraes et al., 2015). Naing et al., (2022) showed that
ethylene-insensitive mutants exhibited increased sensitivity to drought stress, indicating that
ethylene-responsive genes contribute to drought adaptation by modulating stomatal conductance
and enhancing root growth. Thus, our findings imply that the induction of ethylene-responsive
genes may act as a central regulator, integrating signals from both Fe deficiency and drought stress
to orchestrate an adaptive response in soybean. Ethylene signaling may facilitate crosstalk between
Fe deficiency and drought stress pathways. This crosstalk can regulate the reactive oxygen species
(ROS) scavenging enzymes can help mitigate oxidative stress, which is common under both Fe
deficiency and drought conditions (Zhang et al., 2016; Kabir et al., 2012). In support of this
possible phenomenon, we observed the upregulation of glutaredoxin-3 in the roots of soybean.
Glutaredoxin-3 is involved in the assembly and maintenance of Fe-S clusters, which are crucial
for the function of many enzymes, including those involved in ROS detoxification and other
metabolic processes (Couturier et al., 2015). In addition, the induction of ferredoxin among the
upregulated genes also points to the importance of redox homeostasis in stress adaptation.
Ferredoxin is involved in maintaining cellular redox homeostasis by ROS, particularly under
drought, salinity, and high light-intensity stresses in plants (Huang et al., 2020; Lehtimäki et al.,
2010). Taken together, the top upregulated genes found in this study can be leveraged to develop
soybean varieties with enhanced resilience to concurrent abiotic stresses, potentially leading to
improved crop yields and sustainability.

Our RNA-seq analysis exhibited that several downregulated genes are associated with various
physiological processes, including nutrient transport, hormone signaling, and stress responses due
to combined Fe deficiency and drought. Fe deficiency and/or drought can cause significant
nutritional imbalances in plants, disrupting essential metabolic processes and leading to stunted
growth and poor health (Kumari et al., 2022). In this study, the downregulation of nicotianamine
transporter *YSL1* and anion transporter 6 reflects a possible disruption in nutrient uptake and
translocation in soybean (Fiskeby IV) exposed to Fe deficiency and drought. Furthermore, the

decreased expression of jasmonic acid-amido synthetase *JAR1* and auxin efflux carrier component 6 suggests an alteration in hormonal balance, which could be one of the factors disrupting the drought tolerance ability of the genotype Fiskeby IV to combat dual stresses on soybean. Decreased *JAR1* expression can impair drought resilience by weakening the jasmonic acid signaling pathways that regulate stomatal closure and stress-responsive gene activation (Ali et al., 2020). Another study demonstrates that over-expression of *JAR1)* enhances drought tolerance in *Arabidopsis thaliana* by increasing jasmonyl-L-isoleucine production, which leads to better water retention (Mahmud et al., 2022). Although Fe is not a direct cofactor for the enzymes specific to jasmonic acid signaling, it is crucial for the biosynthesis of JA, as it is required for the functioning of lipoxygenases and other enzymes that convert fatty acids into jasmonic acid (Singh et al., 2022; Oliw 2022). In contrast, jasmonate-deficient *cpm2* mutant revealed that jasmonates inhibit the expression of several Fe deficiency-inducible genes associated with Feuptake in rice (Kobayashi et al., 2016). Particularly, auxin plays a critical role in mediating adaptive responses to improve Fe acquisition and drought tolerance in plants (Liu et al., 2024; Iqbal et al., 2022; Kabir et al., 2012). In this study, we also found the downregulation of neoxanthin synthase and solute carrier family 40 highlights disruptions in carotenoid biosynthesis and solute transport, which is consistent with the decreased photosynthetic efficiency and nutrient uptake in Fiskeby IV exposed to concurrent Fe deficiency and drought. Under abiotic stress, such as drought or nutrient deficiency, plants often experience disruptions in chlorophyll and carotenoid biosynthesis resulting in decreased energy production and plant growth (Muhammad et al., 2021; Dong et al., 2007). Carotenoid synthesis may also be affected, impacting plant protection against oxidative damage and influencing overall stress tolerance (Giani and Martínez-Espinosa 2020). Thus, phytohormone and other downregulated genes can be targeted for genetic engineering or breeding strategies aimed at enhancing stress resilience in soybean.

Ethylene enhances dual stress tolerance in soybean
In light of the RNA-seq analysis revealing upregulation of several ethylene-related genes, we conducted a targeted study using an ethylene precursor to determine if it could eliminate the impairment in drought-tolerant Fiskeby IV exposed to combined Fe deficiency and drought. The results of our study underscore the potential of ACC, an ethylene precursor, in enhancing dual stress tolerance in soybean. In this study, supplementation with ACC led to significant improvements in these morpho-physiological traits, suggesting that ethylene signaling helps restore key growth parameters along with leaf Fe status disrupted by combined stress conditions. Precious studies indicating the up-regulation of genes (i.e. *1-aminocyclopropane-1-carboxylic acid synthase (ACS)*, and *1-aminocyclopropane-1-carboxylic acid oxidade*) involved in ethylene synthesis further support the increased ethylene production as part of the adaptive responses under Fe deficiency in plants (García et al., 2015; Sauter et al., 2013). In most cases, ethylene, whose production increases under nutrient deficiency in genotypically efficient genotypes, acts as an activator of stress responses in plants (Dubois et al., 2018). Thus, it appears that ethylene-related or Fe-uptake genes could be potential targets for improving stress tolerance in soybean exposed to Fe deficiency and drought.

Our extended study, which revealed a correlation between root flavonoid content and rhizosphere siderophore levels induced by the presence of an ethylene precursor, further underscores the significance of microbial associations in soybean subjected to combined Fe deficiency and drought. While 'Fe- drought+' stress conditions alone did not affect root flavonoid content, ACC

supplementation significantly increased flavonoid levels. This enhancement in flavonoids might contribute to improved stress resilience in soybean as flavonoids are known to play roles in symbiotic associations (Wang et al., 2022). Flavonoids play an essential role in rhizobium-legume symbiosis as chemoattractants and *Nod* gene inducers, and they are also believed to stimulate fungal spore germination and hyphal growth (Abdel-Lateif et al., 2012; Hassan et al., 2012). Similarly, siderophore levels in the rhizosphere were elevated under 'Fe-drought stress', and ACC further boosted siderophore production. This indicates that ACC not only improves plant tolerance to concurrent Fe deficiency and drought but also potentially aids in Fe acquisition by increasing siderophore-mediated Fe mobilization in the rhizosphere. Microbial siderophores are compounds secreted by both bacterial and fungal microbial communities that bind and solubilize Fe, making them more accessible for uptake by plants (Singh et al., 2022). These siderophores can enhance plant Fe acquisition, particularly in Fe-deficient soils, by facilitating the transfer of Fe from the soil to plant roots (Kabir and Bennetzen, 2024; Radzki et al., 2013). Our qPCR experiment further demonstrated the restoration of 16S bacterial and ITS fungal community when ACC was augmented in soybeans exposed to dual stresses, compared to plants exposed to Fe-drought+ conditions. Ethylene plays a complex yet crucial role in enhancing microbial symbiosis in plants by modulating defense responses and facilitating beneficial interactions (Shekhawat et al. 2022; Ravanbakhsh et al. 2018). In legume-rhizobia symbiosis, ethylene levels influence root nodule formation essential for nitrogen fixation (Guinel, 2015). Similarly, in mycorrhizal associations, ethylene can modulate root architecture and colonization (Khatabi et al. 2012). Thus, it appears that ethylene signaling may indirectly promote microbial symbiosis, thereby recovering Fe status and drought tolerance in Fiskeby IV exposed to combined Fe deficiency and drought. This study may inspire future research on the composition of microbial dynamics, including rhizobia, to uncover the interactions between ethylene signaling and microbiome to possess plant growth-promoting and stress resilience functions. This could lead to improved stress tolerance in soybeans in response to Fe deficiency and drought.

5 Conclusion

This study evaluated the physiological and transcriptional changes in the drought-tolerant soybean genotype Fiskeby IV, which loses its tolerance when exposed simultaneously to Fe deficiency and drought. Exposure to these combined stresses resulted in significant reductions in plant biomass, photosynthetic efficiency, and nutrient uptake. Despite these disruptions, the core photochemical efficiency of photosystem II remained stable, suggesting protective mechanisms at play. RNA-seq analysis highlighted the upregulation of genes related to ethylene signaling, rhizosphere acidification, and redox homeostasis (Figure 8). This indicates that ethylene is crucial for coordinating Fe homeostasis and drought adaptation pathways. Conversely, combined Fe deficiency and drought led to the downregulation of genes related to nutrient uptake. Supplementation with an ethylene precursor, improved morpho-physiological traits and Fe status under combined stress, enhanced root flavonoid content, and increased rhizosphere siderophore levels and microbial community abundance. These findings underscore the potential of targeting ethylene signaling and microbial interactions to enhance soybean tolerance to concurrent Fe deficiency and drought. Future research should focus on these mechanisms to develop strategies that improve stress resilience and sustainability in soybean crops.

Data availability statement

Raw RNA-seq data was submitted to NCBI under the following accession number: PRJNA1140285.

Author contributions

MRH cultivated the plants, conducted most of the experiments, performed RNA-seq analysis, and prepared the draft manuscript. AT extracted the RNA from root samples. MGM helped with data interpretations and critically revised the manuscript. AHK conceived the idea, supervised the whole work, and revised the manuscript.

Acknowledgments

This research was supported by a startup grant (5SFAES-293007) and Elizabeth and Hayden Cutler's Endowed Professorship in Biotechnology from the University of Louisiana at Monroe.

Conflict of Interest Statement

We have no conflict of interest.

References

Abdel-Lateif, K., Bogusz, D., and Hocher, V. (2012). The role of flavonoids in the establishment of plant roots endosymbioses with arbuscular mycorrhiza fungi, rhizobia and Frankia bacteria. *Plant Signal Behav.* 7(6), 636–641.

Afshar, R. M., Hadi, H., and Pirzad, A. (2012). Effect of Nano-Fe foliar application on qualitative and quantitative characteristics of cowpea, under end season drought stress. *Int. Res. J. Appl. Basic Sci.* 3, 1709–1717.

Ali, S., Hayat, K., Iqbal, A., and Xie, L. (2020) Implications of Abscisic Acid in the Drought Stress Tolerance of Plants. *Agronomy.* 10, 1323.

Ali, M. S., and Baek, K. H. (2020). Jasmonic Acid Signaling Pathway in Response to Abiotic Stresses in Plants. *Int. Journal Mol. Sci.* 21(2), 621.

Arraes, F. B., Beneventi, M. A., Lisei de Sa, M. E., Paixao, J. F., Albuquerque, E. V., Marin, S. R., Purgatto, E., Nepomuceno, A. L., and Grossi-de-Sa, M. F. (2015). Implications of ethylene biosynthesis and signaling in soybean drought stress tolerance. *BMC Plant Biol.* 15, 213.

Barnabas, B., Jäger, K., and Fehér, A. (2008). The effect of drought and heat stress on reproductive processes in cereals. *Plant Cell EnvFe.* 31, 11–38.

Chaves M.M., Maroco J.P., and Pereira J.S. (2003) Understanding plant responses to drought-from genes to the whole plant. *Funct. Plant Biol.* 30, 239–264.

Chen, H., Bullock, D. A., Jr, Alonso, J. M., and Stepanova, A. N. (2021). To Fight or to Grow: The Balancing Role of Ethylene in Plant Abiotic Stress Responses. *Plants* 11(1), 33.

Couturier, J., Przybyla-Toscano, J., Roret, T., Didierjean, C., and Rouhier, N. (2015). The roles of glutaredoxins ligating Fe-S clusters: Sensing, transfer or repair functions? *Biochimica et Biophysica Act.* 1853(6), 1513–1527.

Crowley D.E. Microbial siderophores in the plant rhizosphere. In: Barton L.L., Abadia J., editors. Fe Nutrition in Plants and Rhizospheric Microorganisms. Springer; Dordrecht, The Netherlands: 2006. pp. 169–198.

Dong, H., Deng, Y., Mu, J., Lu, Q., Wang, Y., Xu, Y., Chu, C., Chong, K., Lu, C., and Zuo, J. (2007). The Arabidopsis Spontaneous Cell Death1 gene, encoding a zeta-carotene desaturase

essential for carotenoid biosynthesis, is involved in chloroplast development, photoprotection and retrograde signalling. *Cell Res.* 17(5), 458–470.

D'Oria, A., Courbet, G., Billiot, B., Jing, L., Pluchon, S., Arkoun, M., Maillard, A., Roux, C. P., Trouverie, J., Etienne, P., Diquélou, S., and Ourry, A. (2022). Drought specifically downregulates mineral nutrition: Plant ionomic content and associated gene expression. *Plant Direct* 6(8), e402.

Dubois M, Van den Broeck L, Inzé D. The Pivotal Role of Ethylene in Plant Growth. Trends Plant Sci. 2018;23(4):311-323.

Dubois, M., Van den Broeck, L., and Inzé, D. (2018). The Pivotal Role of Ethylene in Plant Growth. *Trends Plant Sci.* 23(4), 311–323.

Elanz, E., Mohsen, R., and Ahmed, B. (2011). Influence of salt stress on cations concentration quality and quantity of sunflower cultivars. *J. Food Agric. EnvFe.* 2, 469–476.

Fatma, M., Asgher, M., Iqbal, N., Rasheed, F., Sehar, Z., Sofo, A., and Khan, N. A. (2022). Ethylene Signaling under Stressful EnvFements: Analyzing Collaborative Knowledge. Plants (Basel, Switzerland), 11(17), 2211.

García, M. J., Romera, F. J., Lucena, C., Alcántara, E., and Pérez-Vicente, R. (2015). Ethylene and the Regulation of Physiological and Morphological Responses to Nutrient Deficiencies. Plant physiology, 169(1), 51–60.

Geng, A., Lian, W., Wang, Y., Liu, M., Zhang, Y., Wang, X., and Chen, G. (2024). Molecular Mechanisms and Regulatory Pathways Underlying Drought Stress Response in Rice. *Int. J Mol. Sci.* 25(2), 1185.

Giani, M., and Martínez-Espinosa, R. M. (2020). Carotenoids as a Protection Mechanism against Oxidative Stress in *Haloferax mediterranei*. *Antioxidants* 9(11), 1060.

Guinel, F.C. (2015) Ethylene, a Hormone at the Center-Stage of Nodulation. *Front. Plant Sci.* 6, 1121.

Hassan, S., and Mathesius, U. (2012). The role of flavonoids in root-rhizosphere signalling: opportunities and challenges for improving plant-microbe interactions. *J. Exp. Bot.* 63(9), 3429–3444.

Huang, H. E., Ho, M. H., Chang, H., Chao, H. Y., and Ger, M. J. (2020). Overexpression of plant ferredoxin-like protein promotes salinity tolerance in rice (*Oryza sativa*). *Plant Physiol. Biochem.* 155, 136–146.

Iqbal, S., Wang, X., Mubeen, I., Kamran, M., Kanwal, I., Díaz, G. A., Abbas, A., Parveen, A., Atiq, M. N., Alshaya, H., Zin El-Abedin, T. K., and Fahad, S. (2022). Phytohormones Trigger Drought Tolerance in Crop Plants: Outlook and Future Perspectives. *Front. Plant Sci.* 12, 799318.

Jin, C. W., Li, G. X., Yu, X. H., and Zheng, S. J. (2010). Plant Fe status affects the composition of siderophore-secreting microbes in the rhizosphere. *Ann Bot.* 105(5), 835–841.

Kabir, A. H., Paltridge, N. G., Able, A. J., Paull, J. G., and Stangoulis, J. C. (2012). Natural variation for Fe-efficiency is associated with upregulation of Strategy I mechanisms and enhanced citrate and ethylene synthesis in Pisum sativum L. Planta, 235(6), 1409–1419.

Kabir, A. H., and Bennetzen, J. L. (2024). Molecular insights into the mutualism that induces Fe deficiency tolerance in sorghum inoculated with *Trichoderma harzianum*. *Microbiol. Res.* 281, 127630.

Khatabi, B., Molitor, A., Lindermayr, C., Pfiffi, S., Durner, J., von Wettstein, D., Kogel, K. H., and Schäfer, P. (2012). Ethylene supports colonization of plant roots by the mutualistic fungus *Piriformospora indica*. *PloS one* 7(4), e35502.

Khan, M. I. R., Nazir, F., Maheshwari, C., Chopra, P., Chhillar, H., and Sreenivasulu, N. (2023). Mineral nutrients in plants under changing envFements: A road to future food and nutrition security. *Plant Genome* 16(4), e20362.

Kobayashi, T., and Nishizawa, N. K. (2012). Fe uptake, translocation, and regulation in higher plants. *Ann. Rev. Plant Biol.* 63, 131–152.

Kobayashi, T., Itai, R. N., Senoura, T., Oikawa, T., Ishimaru, Y., Ueda, M., Nakanishi, H., and Nishizawa, N. K. (2016). Jasmonate signaling is activated in the very early stages of Fe deficiency responses in rice roots. *Plant Mol. Biol.* 91(4-5), 533–547.

Kroh, G.E., and Pilon, M. (2019). Connecting the negatives and positives of plant Fe homeostasis. *New Phytol.* 223(3), 1052-1055.

Kulkarni, K. P., Tayade, R., Asekova, S., Song, J. T., Shannon, J. G., and Lee, J. D. (2018). Harnessing the Potential of Forage Legumes, Alfalfa, Soybean, and Cowpea for Sustainable Agriculture and Global Food Security. *Front. Plant Sci.* 9, 1314.

Kumar, M., Kumar Patel, M., Kumar, N., Bajpai, A. B., and Siddique, K. H. M. (2021). Metabolomics and Molecular Approaches Reveal Drought Stress Tolerance in Plants. *Int. J. Mol. Sci.* 22(17), 9108.

Kumari, V. V., Banerjee, P., Verma, V. C., Sukumaran, S., Chandran, M. A. S., Gopinath, K. A., Venkatesh, G., Yadav, S. K., Singh, V. K., and Awasthi, N. K. (2022). Plant Nutrition: An Effective Way to Alleviate Abiotic Stress in Agricultural Crops. *Int. J. Mol. Sci.* 23(15), 8519.

Künstler, A., Gullner, G., Ádám, A. L., Kolozsváriné Nagy, J., and Király, L. (2020). The Versatile Roles of Sulfur-Containing Biomolecules in Plant Defense-A Road to Disease Resistance. *Plants* 9(12), 1705.

Lehtimäki, N., Lintala, M., Allahverdiyeva, Y., Aro, E. M., and Mulo, P. (2010). Drought stress-induced upregulation of components involved in ferredoxin-dependent cyclic electron transfer. *J Plant Phys.* 167(12), 1018–1022.

Li, S., Yang, L., Huang, X., Zou, Z., Zhang, M., Guo, W., Addo-Danso, S. D., and Zhou, L. (2023). Mineral Nutrient Uptake, Accumulation, and Distribution in *Cunninghamia lanceolata* in Response to Drought Stress. *Plants* 12(11), 2140.

Li, W., and Lan, P. (2017). The Understanding of the Plant Fe Deficiency Responses in Strategy I Plants and the Role of Ethylene in This Process by Omic Approaches. *Front Plant Sci.* 8, 40.

Liang G. (2022). Fe uptake, signaling, and sensing in plants. *Plant Comm.* 3(5), 100349

Lidoy J, Berrio E, García M, España-Luque L, Pozo MJ, López-Ráez JA. (2023). Flavonoids promote *Rhizophagus irregularis* spore germination and tomato root colonization: A target for sustainable agriculture. *Front Plant Sci.* 5, 1094194.

Liu, L., Yahaya, B. S., Li, J., and Wu, F. (2024). Enigmatic role of auxin response factors in plant growth and stress tolerance. *Front Plant Sci.* 15, 1398818.

Lucena, C., Romera, F. J., García, M. J., Alcántara, E., and Pérez-Vicente, R. (2015). Ethylene Participates in the Regulation of Fe Deficiency Responses in Strategy I Plants and in Rice. *Front Plant Sci.* 6, 1056.

Mahmud, S., Ullah, C., Kortz, A., Bhattacharyya, S., Yu, P., Gershenzon, J., and Vothknecht, U. C. (2022). Constitutive expression of Jasmonate resistant 1 induces molecular changes that prime the plants to better withstand drought. *Plant Cell Env.* 45(10), 2906–2922.

McDonnell, L., Plett, J. M., Andersson-Gunnerås, S., Kozela, C., Dugardeyn, J., Van Der Straeten, D., Glick, B. R., Sundberg, B., and Regan, S. (2009). Ethylene levels are regulated by a plant encoded 1-aminocyclopropane-1-carboxylic acid deaminase. *Physiol Plant.* 136(1), 94-109.

Michalak, A., Wdowikowska, A., and Janicka, M. (2022). Plant Plasma Membrane Proton Pump: One Protein with Multiple Functions. *Cells* 11(24), 4052.

Mohorović, P., Geldhof, B., Holsteens, K., Rinia, M., Ceusters, J., and Van de Poel, B. (2023). Effect of ethylene pretreatment on tomato plant responses to salt, drought, and waterlogging stress. *Plant Direct* 7(11), e548.

Morrissey, J., and Guerinot, M. L. (2009). Fe uptake and transport in plants: the good, the bad, and the ionome. *Chem Rev.* 109(10), 4553–4567.

Muhammad, I., Shalmani, A., Ali, M., Yang, Q. H., Ahmad, H., and Li, F. B. (2021). Mechanisms Regulating the Dynamics of Photosynthesis Under Abiotic Stresses. *Front Plant Sci.* 11, 615942.

Müller, M., and Munné-Bosch, S. (2015). Ethylene Response Factors: A Key Regulatory Hub in Hormone and Stress Signaling. *Plant Physiol.* 169(1), 32–41.

Naing, A. H., Campol, J. R., Kang, H., Xu, J., Chung, M. Y., and Kim, C. K. (2022). Role of Ethylene Biosynthesis Genes in the Regulation of Salt Stress and Drought Stress Tolerance in Petunia. *Front Plant Sci.* 13, 844449.

Oliw E. H. (2022). Fe and manganese lipoxygenases of plant pathogenic fungi and their role in biosynthesis of jasmonates. *Arch. Biochem. Biophy.* 722, 109169.

Radzki, W., Gutierrez Mañero, F. J., Algar, E., Lucas García, J. A., García-Villaraco, A., and Ramos Solano, B. (2013). Bacterial siderophores efficiently provide Fe to Fe-starved tomato plants in hydroponics culture. *Antonie van Leeuwenhoek* 104(3), 321–330.

Ravanbakhsh, M., Sasidharan, R., Voesenek, L. A. C. J., Kowalchuk, G. A., & Jousset, A. (2018). Microbial modulation of plant ethylene signaling: ecological and evolutionary consequences. *Microbiome* 6(1), 52.

Ren, Z., Wang, R.Y., Huang, X.Y., and Wang Y. (2022) Sulfur Compounds in Regulation of Stomatal Movement. *Front. Plant Sci.* 13, 846518.

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., and Smyth, G. K. (2015). limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* 43(7), e47.

Samira, R., Stallmann, A., Massenburg, L.N., Long, T.A. (2013) Feing out the issues: integrated approaches to understanding Fe homeostasis in plants. *Plant Sci.* 210, 250-259.

Sauter M, Moffatt B, Saechao MC, Hell R, Wirtz M (2013) Methionine salvage and S-adenosylmethionine: essential links between sulfur, ethylene and polyamine biosynthesis. *Biochem J.* 451, 145–154.

Schliemann, W., Ammer, C., and Strack, D. (2008). Metabolite profiling of mycorrhizal roots of *Medicago truncatula*. *Phytochem.* 69(1), 112–146.

Seleiman, M. F., Al-Suhaibani, N., Ali, N., Akmal, M., Alotaibi, M., Refay, Y., Dindaroglu, T., Abdul-Wajid, H. H., and Battaglia, M. L. (2021). Drought Stress Impacts on Plants and Different Approaches to Alleviate Its Adverse Effects. *Plants* 10(2), 259.

Shekhawat, K., Fröhlich, K., García-Ramírez, G. X., Trapp, M. A., and Hirt, H. (2022). Ethylene: A Master Regulator of Plant-Microbe Interactions under Abiotic Stresses. *Cells* 12(1), 31.

Singh, P., Arif, Y., Miszczuk, E., Bajguz, A., and Hayat, S. (2022). Specific Roles of Lipoxygenases in Development and Responses to Stress in Plants. *Plants* 11(7), 979.

Singh, P., Chauhan, P. K., Upadhyay, S. K., Singh, R. K., Dwivedi, P., Wang, J., Jain, D., and Jiang, M. (2022). Mechanistic Insights and Potential Use of Siderophores Producing Microbes in Rhizosphere for Mitigation of Stress in Plants Grown in Degraded Land. *Front. Microbiol.* 13, 898979.

Teshome, D. T., Zharare, G. E., and Naidoo, S. (2020). The Threat of the Combined Effect of Biotic and Abiotic Stress Factors in Forestry Under a Changing Climate. *Front. Plant Sci.* 11, 601009.

Volaire F. (2018). A unified framework of plant adaptive strategies to drought: Crossing scales and disciplines. *Global Change Biol.* 24(7), 2929–2938.

Wang, L., Chen, M., Lam, P. Y., Dini-Andreote, F., Dai, L., and Wei, Z. (2022). Multifaceted roles of flavonoids mediating plant-microbe interactions. *Microbiome* 10(1), 233.

Waters, B. M., Lucena, C., Romera, F. J., Jester, G. G., Wynn, A. N., Rojas, C. L., Alcántara, E., and Pérez-Vicente, R. (2007). Ethylene involvement in the regulation of the H(+)-ATPase *CsHA1* gene and of the new isolated ferric reductase *CsFRO1* and Fe transporter *CsIRT1* genes in cucumber plants. *Plant Physiol. Biochem.* 45(5), 293–301.

Wu, J., Wang, C., Zheng, L., Wang, L., Chen, Y., Whelan, J., and Shou, H. (2011). Ethylene is involved in the regulation of Fe homeostasis by regulating the expression of Fe-acquisition-related genes in Oryza sativa. *J. Exp. Bot.* 62(2), 667–674.

Zhang, H., Li, A., Zhang, Z., Huang, Z., Lu, P., Zhang, D., Liu, X., Zhang, Z. F., and Huang, R. (2016). Ethylene Response Factor *TERF1*, Regulated by ETHYLENE-INSENSITIVE3-like Factors, Functions in Reactive Oxygen Species (ROS) Scavenging in Tobacco (Nicotiana tabacum L.). *Sci. Rep.* 6, 29948.

Zhang, J., Wei, J., Li, D., Kong, X., Rengel, Z., Chen, L., Yang, Y., Cui, X., and Chen, Q. (2017). The Role of the Plasma Membrane H+-ATPase in Plant Responses to Aluminum Toxicity. *Front. Plant Sci.* 8, 1757.

Table 1. ICP-MS analysis of nutrients in the root and shoot of soybean grown in control and Fe-drought+ conditions. Different letters indicate significant differences between the two treatment groups at a P <0.05 level. The data represents the means ± SD of three independent biological samples (*n* = 3).

Elements	Root		Shoot	
	Control	Fe- Drought+	Control	Fe- Drought+
Fe (ppm)	460.8±32.5a	339.9±28.8b	95.3±8.6a	66.7±4.9b
Zn (ppm)	34.4±1.36a	34.2±1.64a	40.5±6.0a	28.6±3.1b
Mn (ppm)	24.1±4.0a	22.1±2.9a	49.2±8.5a	25.9±4.4b
N (%)	2.48±0.19a	1.77±0.19b	3.79±0.01a	3.45±0.07b
S (%)	0.23±0.02a	0.14±0.04b	0.25±0.01a	0.25±0.03a
K (%)	0.95±0.11a	0.70±0.05b	3.50±0.13a	3.51±0.21a
Ca (%)	0.64±0.06a	0.065±0.02a	1.02±0.17a	0.56±0.02b
Mg (%)	0.59±0.13a	0.40±0.07a	0.36±0.03a	0.40±0.04a

725 **Table 2.** The top 15 differentially expressed genes in roots of soybean exposed to combined Fe
726 deficiency and drought relative to untreated controls. Values marked with an asterisk are significant
727 (> Log2 fold, < P value 0.05).

Upregulated genes	Fe- Drought+ vs Control (fold change)
Glyma.04G089000 (MATE efflux family protein)	2.0
Glyma.20G202200 (Ethylene-response sensor 2)	2.1
Glyma.07G122500 (Copper ion transporter)	2.0
Glyma.15G180000 (Ethylene-responsive TF018)	2.5
Glyma.11G091400 (Heavy-metal-associated protein 26)	2.4
Glyma.13G087200 (Sulfate transporter 1.1)	2.1
Glyma.19G130800 (Ferredoxin)	2.4
Glyma.07G215500 (ABA-GTase)	4.8
Glyma.10G227400 (ABC transporter C family 10)	3.0
Glyma.13G088100 (Ethylene-responsive TF5)	3.1
Glyma.03G262600 (ATPase 1)	2.1
Glyma.14G071400 (BZIP transcription factor ATB2)	2.2
Glyma.02G261700 (Dehydration-responsive binding protein 2A)	3.0
Glyma.14G000800 (Calcium-dependent protein kinase 6)	2.4
Glyma.13G180400 (Glutaredoxin-3)	2.3
Downregulated genes	
Glyma.03G134200 (Calcium transporting ATPase 4)	-3.0
Glyma.17G203500 (Molybdate transporter 1)	-4.4
Glyma.03G212400 (heavy metal transport protein)	-2.6
Glyma.10G240300 (Peroxidase 66)	-4.1
Glyma.17G087500 (Heme-binding protein-related)	-3.1
Glyma.04G019500 (Multi-copper oxidase type I)	-2.7
Glyma.07G057900 (Jasmonic acid-amido synthetase JAR1)	-3.1
Glyma.13G162900 (Anion transporter 6)	-2.7
Glyma.07G006500 (Sulfate transporter 3.4)	-2.5
Glyma.10G172900 (Nicotianamine transporter YSL1)	-4.2
Glyma.13G038300 (Auxin efflux carrier component-6)	-3.4
Glyma.18G204800 (Ammonium transporter 2)	-3.0
Glyma.10G146600 (Solute carrier family 40)	-4.0
Glyma.13G315200 (Neoxanthin synthase)	-3.9
Glyma.05G211300 (Major facilitator family protein)	-4.0

745
746 **Figure 1.** Changes in root length (A), root dry weight (B), shoot height (C), and shoot dry weight
747 (D) in soybean in 'control' and 'Fe-drought+' conditions. Different letters above the bar indicate
748 significant differences between treatment groups at a P <0.05 level. The data represents the means
749 ± SD of three independent biological samples ($n = 3$).

750
751 **Figure 2.** Changes in chlorophyll score (A), Fv/Fm: maximal quantum yield of PS II (B), and
752 Pi_ABS: photosynthesis performance index (C) in the leaves of soybean in 'control' and 'Fe-
753 drought+' conditions. Different letters above the bar indicate significant differences between
754 treatment groups at a P <0.05 level. The data represents the means ± SD of three independent
755 biological samples ($n = 3$).
756

(A)

(B)

(C)

757
758 **Figure 3.** Analysis of RNA-sequencing data for differentially expressed genes (DEGs): principal
759 component analysis of DEGs (A), vein diagram of upregulated and downregulated genes (>Log2
760 fold, p<0.05) grown in 'Fe-drought+' vs. 'control' (B) and MA plot of genes expressed in 'Fe-
761 drought+' vs. 'control' conditions (C).

778
779 **Figure 4.** Heatmap of top 15 differentially expressed (>Log2 fold, p<0.05) upregulated (A) and
780 downregulated (B) genes in roots of Fiskeby IV in 'Fe-drought+' vs 'control' conditions.

Figure 5. Gene enrichment analysis for the upregulated (A) and downregulated genes (B) in roots of Fiskeby IV cultivated in 'Fe-drought+' and 'control' conditions for biological processes, cellular components, and molecular functions.

Figure 6. Effect of ACC on plant phenotype (A), chlorophyll score (B), plant height (C), plant dry weight (D), stem diameter (E), relative water content (F), leaf Fe (G), flavonoid content (H) and rhizosphere siderophore (I) in soybean plants cultivated with or without Fe-drought+ conditions. Different letters above the bar indicate significant differences between treatment groups at a P <0.05 level, where applicable. The data represents the means ± SD of three independent biological samples (*n* = 3).

Figure 7. Effect of ACC on the relative abundance of 16S (bacteria) and ITS (fungi) communities in roots of Fiskeby IV exposed to combined Fe deficiency and drought. Different letters above the bar indicate significant differences between treatment groups at a P <0.05 level, where applicable. The data represents the means ± SD of three independent biological samples ($n = 3$).

Figure 8. Physiological and transcriptional responses of soybean plants exposed to concurrent Fe deficiency and drought. The combined Fe deficiency and drought caused the upregulation of genes related to ethylene signaling (*Ethylene-response sensor 2*, *Ethylene-responsive TF018*, *Ethylene-responsive TF5*), proton extrusion (*ATPase 1*) and redox homeostasis (*Glutaredoxin-3*) while the expression of genes associated with peroxidase activity (*Peroxidase 66*), phytohormone synthesis (*Auxin efflux carrier component-6*, *Jasmonic acid-amido synthetase*) and nutrient transport (*Sulfate transporter 3.4*, *Ammonium transporter 2*) showed downregulation. 'Fe-Drought+' conditions caused a decline in mineral elements in both root and leaf. Interestingly, ethylene precursor (ACC) improved Fe homeostasis either directly or indirectly through root flavonoid and rhizosphere siderophore to induce drought tolerance.