Pain distribution can be determined by classical conditioning

Jakub Nastaj[1*], Jacek Skalski[1], Daria Nowak[1], Natalia Kruszyna[1], Przemysław Bąbel[4], Tibor Szikszay[2], Kerstin Luedtke[2], Rafał Gnat[3], & Wacław M. Adamczyk[1,2]

[1]Laboratory of Pain Research, Institute of Health Sciences, Academy of Physical Education, Katowice, Poland

[2]Institute of Health Sciences, Department of Physiotherapy, Pain & Exercise Research Luebeck (P.E.R.L.), University of Luebeck, Lübeck, German

[3]Motion Analysis Laboratory, Department of Physiotherapy, Academy of Physical Education, Katowice, Poland

[4]Pain Research Group, Institute of Psychology, Jagiellonian University, Kraków, Poland

*Corresponding author: Laboratory of Pain Research, Institute of Health Sciences, Academy of Physical Education, Katowice, Poland

Number of figures: 3

Number of tables: 2

Number of pages: 22

Number of Appendixes: 1

Disclosure: The authors declare no conflict of interest.

Funding: Polish National Science Center (2020/37/B/HS6/04196)

                                                                  ABSTRACT

Chronic widespread pain (CWP) - as many other clinical presentations - manifests in ongoing pain without identifiable

structural cause, with pain that spreads over multiple body areas. The development and maintenance of symptoms may

involve learning mechanisms. The authors have hypothesized that pain distribution can be learned through classical

conditioning or elicited by verbal suggestion. Ninety-four healthy volunteers participated in this study and were

randomly distributed to four groups. In the classical conditioning combined with verbal suggestion group, US- (small

pain distribution) and US+ (large pain distribution) were paired with visual stimuli (CS+ and CS-) and participants

were told about this association. In the verbal suggestion group, the conditioning was not performed, whereas in

classical conditioning only group, learning was not combined with suggestion. In the control group, conditioning and

suggestion did not take place. Ratings of perceived pain distribution (PD) were collected after each trial and ratings of

pain intensity after each block of trials. During the testing phase, participants were exposed to electrocutaneous stimuli

corresponding to only the small (US-) pain distribution. Results showed significant differences between CS+ and CS-

pain distribution ratings across the experimental groups: conditioning + verbal suggestion ($p<0.01$), conditioning only

group ($p<0.05$) and verbal suggestion only group ($p<0.05$), but not in the control group ($p>0.05$). Furthermore,

significant differences in the perceived pain distribution were found between the control group and all experimental

groups. This result supports our main hypothesis that the perceived pain distribution can be influenced by classical

conditioning as well as verbal suggestion, although the effect is stronger when both are combined.

Key words: pain, classical conditioning, pain distribution, spatial aspect of pain, learning mechanism

1. Introduction

Pain perception requires the integration of spatial and temporal aspects of sensory processing [14,37,45]. Pain can be experienced even in the absence of current or persistent tissue damage [40]. Many clinical presentations manifest themselves in ongoing pain without evident structural pathology or patients continue to experience pain after the completion of healing processes. Examples include chronic non-specific pain states such as chronic widespread pain (CWP) [26], which is characterized by ongoing and spreading musculoskeletal diffuse pain lasting for 3 months or longer [20,26]. To develop effective treatment strategies for CWP, it is crucial to understand the mechanisms that contribute to its onset and persistence, which could involve learning mechanisms.

Acute pain is typically localized to the affected region of the body and is initiated by the activation of primary afferent neurons. This circumstance provides an optimal environment for the mechanisms underlying pain learning, particularly classical conditioning. Evidence supports the modulation of pain through learning processes such as classical conditioning [5,25,32], operant conditioning [1,4] or observational learning [43,44]. Previous studies have only examined the sensory (intensity) and affective (unpleasantness) aspects of pain concerning learning mechanisms, validating that intensity [32] and unpleasantness of pain [39] could be amplified through learning [25]. It is noteworthy, however, that no study has yet investigated whether the distribution of pain can be influenced by learning processes. Given the substantial published data on pain intensity [6,8,21], the authors have hypothesized that pain location could likewise be subject to classical conditioning when visual stimuli are applied as conditioned stimuli (CS). If the distribution of pain indeed originates through learning mechanisms, this can offer new perspectives accounting for CWP and spatial phenomena such as radiation of pain and spatial summation of pain [42].

In most former studies, classical conditioning was paired with verbal suggestion to evoke a nocebo effect. Results of a meta-analysis on nocebo hyperalgesia [35] suggest that this effect is more powerful when it arises from verbal suggestion combined with conditioning, as opposed to either verbal suggestion or conditioning alone. Moreover, generalization has been suggested as one of the mechanisms that could play a substantial role in the onset of chronic pain [33]. So far, stimulus generalization has been shown to contribute to the expansion of the conditioned placebo effect [23]. However, it remains to be explored whether the spatial expansion of pain is similarly prone to stimulus generalization.

The main aim of this study was to investigate whether classical conditioning may be involved in the learning of perceived pain distribution, i.e., to address whether non-nociceptive conditioned stimuli (e.g., visual cues) could

induce pain sensation with different (larger/smaller) perceived pain distribution through classical conditioning. The secondary aim of this experiment was to investigate whether the impact of conditioning is heightened when learning is coupled with a verbal suggestion. The final aim of the experiment was to investigate whether the conditioned pain distribution extends to a perceptually similar but novel stimulus through stimulus generalization.

2. Methods and materials

2.1. Overview of the experiment

The experiment utilized a within- and between-subjects design and involved healthy volunteers assigned to one of four groups: classical conditioning only, verbal suggestion only, conditioning + verbal suggestion, or control group. Each group received the same number and type of stimulation: electrocutaneous stimuli (as unconditioned stimuli, US) inducing pain of small (US-) or large distribution (US+), which were preceded by visual (color) stimuli (as conditioned stimuli, CS). The groups differed from each other in the associations between visual and electrical stimuli. In the conditioning groups, the stimulation inducing pain of large distribution was consistently paired with one specific color presented to participants, whereas in the groups without conditioning, there was no associations and colors were presented randomly. In the suggestion groups, participants were informed that one specific color was associated with pain distributed over a larger area. The main outcome measure in this experiment was the size of the painful area assessed via a pain distribution (PD) rating task.

2.2. General information

The study protocol was approved by the ethics committee at the Academy of Physical Education in Katowice (no. 4/2022) and registered at the Open Science Framework platform (https://osf.io/3qezt) using the AsPredicted.org template. The study was conducted in the certified (ISO) Laboratory of Pain Research. The experiment followed the recommendations of the Declaration of Helsinki [48]. Participants were given written and oral information of the study procedures before informed consent was obtained. Moreover, participants were informed that they can withdraw from the study at any timepoint without any reason and consequences. Each volunteer received financial remuneration for their participation.

2.3. Study population and eligibility

A group of 97 healthy participants (age: M= 21.44, 18-33 years) was recruited (between February and December 2023) (Appendix). The following eligibility criteria were applied: pain free on the day of the study, healthy (self-report), no color vision deficiency, no chronic pain or experienced prolonged pain in the last 3 months, no

pregnancy, no cardiovascular or neurological disease, no chronic medication use, no mental illness or any systemic disease, no electronic devices in or at the body or unremovable metal objects in the area of the non-dominant hand, as well as no skin allergies, skin lesions, tattoos or sensory abnormalities in close proximity to the non-dominant hand.

2.4. Sample size calculation

The required sample size was estimated based on previous conditioning effects observed in analyses on perceived pain intensity [7], which showed that at least n = 15 is needed to detect a within-group effect (pain in CS+ vs. CS- condition) of $d_z = 0.92$ (mean of 0.35 ± 0.38) with 90% of power and α set to 0.05 (G*Power 3.1 software) with t test contrast. However, given the novelty of the procedures and the need to counterbalance the factors such as the color of CS (green, blue) and localization of US (proximal, distal) it was preregistered *a priori* to increase a target sample size to 24 per group. Such sample size also ensured that at least 90% of power will be maintained even with potential dropouts.

2.5. Outcome measures

Main outcome measure in this experiment was the length of PD expressed in centimeters – participants rated the length of the spatial dimension of pain they perceived by drawing a horizontal line on the computer's screen. The longer the line, the greater the PD. The secondary outcome was the rating of pain intensity, which was collected by using the Numerical Rating Scale (NRS), ranging from "0" (no pain) to "10" (worst pain imaginable). Intensity ratings were collected to ensure the stimulation was noxious. Further outcomes collected to characterize the study sample included: age, sex (assigned at birth), handedness, height (cm), and weight (kg). Also, fear of pain as a state was measured on a "0" (not fear at all) to "10" (highest possible fear) NRS scale and as a trait via the Fear of Pain Questionnaire III (FPQ-III) [29].

2.6. Apparatus, stimulation, and experimental setup

During the experiment, participants sat upright at a desk, the non-dominant hand with the electrodes attached was placed on the desktop (with the electrodes facing the desktop). Two planar-concentric ($\varnothing$ 8 mm) electrodes separated by 4cm (WASP electrodes, Brainbox Ltd., Cardiff, United Kingdom) were attached to the ulnar edge of the non-dominant hand such that they were positioned within the C8 dermatome. The dominant hand was used to perform the PD rating task (via a computer mouse) and to assess pain intensity on the NRS scale using the keyboard. Visual stimuli, PD rating task, NRS scale and written instructions were displayed on the screen (E900, BENQ, 1280x1024) placed in front of the participant (distance approximately 60 cm). Visual stimuli used in the study served as

conditioned stimuli (CS) and were in the form of the color of the slide displayed on the screen for two seconds. Slides

could be either blue (RGB: 19, 19, 236), green (RGB: 19, 236, 19) or cyan (RGB: 19, 236, 236). The choice of colors

was driven by the results of a recent study demonstrating that these colors do not impact pain perception [47].

To induce pain, electrocutaneous stimuli were generated via a DS8R constant current stimulator (Digitimer,

Welwyn, Garden City, England) with a capacity of 0 to 100 mA with a maximum voltage of 400 V. Each single

stimulus was formed by a series of five (rectangular) pulses with 100μs duration (interval 100ms) to reduce the

intensity of the current applied to the skin. Stimulation was distributed through the D188 remote electrode selector

(Digitimer, Welwyn, Garden City, England) activating the required electrode (proximal OR distal – depending on

counterbalanced assignment) or pair of electrodes (proximal AND distal). External control of the DS8R and D188 was

ensured through a digital/analogue converter device Labjack U3-LV (LabJack Corporation, Lakewood, CO), which

was controlled using "u3" Python-2 library. Experimental procedures were fully automatic and operated by the

PsychoPy 2.3 software [34].

2.7. Study procedures

140   The experiment started from the briefing and consenting procedure. The examiner verified the inclusion and

141 exclusion criteria and prepared participants for the study procedures. The study procedure consisted of five phases: i.

142 familiarization to electrical stimulation, ii. calibration, iii. scale training, iv. main phase of the study (manipulation

143 depending on the group assignment), v. exit tasks.

2.7.1 Familiarization

145   During the familiarization phase, participants received 5 electrocutaneous stimuli of different intensities in a

146 pseudorandom order (5mA, 20mA, 15mA, 10mA and 25mA), applied using a single electrode, to familiarize them

147 with the sensation of electrocutaneous stimuli. This form of pre-exposure was employed to make the subsequent

148 calibration phase and its results more valid.

2.7.2 Calibration

150   To determine individual levels of noxious stimulus intensity for each participant, a calibration procedure was

151 conducted. Although this procedure does not eliminate individual differences in pain perception [13], it was critical in

152 this experiment: The main outcome measure was distribution of pain, thus, highly insensitive individuals could not

153 participate in the PD rating task and for this reason it was decided to not use stimulation of fixed intensity [3]. Two

154 electrodes were calibrated separately to control for the possible differences in perceived pain intensity due to slightly

different localizations of measurement sites. For each electrode, the method of limits was used [3] to determine the sensory detection threshold (t), pain threshold (T) and moderate pain sensation (T5) that would correspond to pain at the level of 5 on NRS scale. Electrocutaneous stimuli were delivered increasingly by 2mA increments until the participant reported the first innocuous sensation (t) and first painful sensation (T). The procedure was continued until the participant reported a moderate sensation of pain (T5) with a safety cut-off value set to 40 mA (to avoid prolonged skin irritation). Participants were excluded from further participation in the experiment if the stimulus intensity reached 40mA and participants did not report a moderately painful sensation (T5). The final intensity of stimulation (in mA) used in the main phase of the study was the same for each of two electrodes. The final intensity(I_{used}) was determined as average intensity inducing moderate pain (T5) via each of two electrodes: I_{used} = (T5 value for proximal electrode + T5 value for distal electrode) / 2.

2.7.3. Training phase

To familiarize participants with the use of the PD rating task and to train them in how to provide pain intensity ratings via NRS, they took part in a short training phase: Two noxious stimuli were applied separately, preceded by two visual stimuli to first train them in PD rating task and secondly the rating of pain intensity by using the NRS scale.

2.7.4. Main phase

The main phase of the study had two components: "experience" phase and "testing" phase **(Figure 1A)**. Participants in each group were first exposed to three blocks of 22 trials each (5 minutes break between blocks) in which CS displayed to participants were followed by a small or large pain distribution modelled here as activation of one electrode (US-, small distribution of pain) or two electrodes (US+, large distribution of pain). Whether pain was induced via the proximal or distal electrode in the US- condition was counterbalanced across individuals.

Depending on the group allocation, green and blue slides in this "experience" phase were consistently matched with a given US type or not. The sequence of trials was pseudorandom across each block and assignment of the colors to the US stimuli was counterbalanced across participants. For instance, green CS+ was coupled with US+ and blue CS- was coupled with US- in groups with conditioning. In other groups consistent associations were not implemented (control group, suggestion only group). Apart from regular trials with PD rating task, each block ended with one pair of trials in which instead of PD the pain intensity was assessed via NRS.

In the testing phase participants were exposed to 12 trials in which always one electrode was activated (US-) regardless of the color of the CS. Apart from CS+ and CS-, which were always displayed in the "experience" phase, a

new visual stimulus (CS$_{GEN}$), i.e. cyan color, was also presented to investigate the possibility of stimulus

generalization of conditioned PD. The order of CS presentations was pseudorandom.

2.7.4.1. Experimental manipulation

In the control group, there was no association between visual stimuli and pain distribution (CS+ and CS-)

during the three blocks of trials. Activation of one (US-) or two electrodes (US+) was in pseudorandom order and the

order of CS+ and CS- stimuli was also pseudorandom. In the final, testing, block participants were always stimulated

with one electrode (CS-). No suggestion about an association between visual stimuli and distribution of pain was

given to participants.

In the verbal suggestion only group, the procedure was the same as in the control group, but after the first

three blocks, verbal suggestion was given before the testing phase to investigate the effect of suggestion. Associations

between visual stimuli (CS+, CS-) and unconditioned stimuli (US-, US+) were suggested to participants whereas in

fact, always one electrode was activated (US-). The suggestion was: *"During this phase of the experiment, after the*

color (blue/green) you will feel pain in a smaller area, and after the color (blue/green) you will feel pain in a larger

area". Verbal suggestion was given to the participants in written (displayed on the screen) and oral form

simultaneously. The assignment of the colors (blue, green) as CS+ and CS- was counterbalanced across the group.

In this classical conditioning + verbal suggestion group, the first three blocks consisted of trials in which

stimuli leading to small (US-) distribution of pain (US-) were paired with CS- stimuli and stimuli leading to large

distribution of pain (US+) were paired with CS+ stimuli. Verbal suggestion was given before the first of the three

blocks to facilitate learning through classical conditioning and was as follows: *"Before receiving electrical stimuli,*

you will observe a color on the monitor. After the color (blue/green) you will feel pain in a smaller area, and after the

color (blue/green) you will feel pain in a larger area". Verbal suggestion was given to the participants in written

(displayed on the screen) and oral form simultaneously. The assignment of the colors (blue, green) as CS+ and CS-

was counterbalanced across the group.

In the conditioning only group, during the three blocks of trials, small (US-) distribution of pain was paired

with CS- visual stimuli and large (US+) pain distribution was paired with CS+ visual stimuli. The assignment of the

colors (blue, green) to CS+ and CS- was counterbalanced across the group. No suggestion about the association

between visual stimuli and the distribution of pain was given to participants.

2.7.4.2. Single trial and block design

Each trial was formed by four distinct periods which are also illustrated in **Figure 1B**: i. jittered inter-trial-

interval (ITI) period (4 – 8s), ii. CS presentation on the computer screen (color slide), iii. delivery of the US, iv. rating

period. Conditioned stimuli were slides of two colors, i.e. green or blue, which were associated with one of US type,

i.e. electrical stimulus applied to one electrode (US-) to model pain distribution at small area or two electrodes (US+)

to model pain of large distribution. In the rating period (iv), participants were asked to rate the length of their pain as a

proxy measure of the pain size by PD rating task. By using a mouse, they were asked to draw on the screen a digital

line such that the length of the line had to match the length of perceived pain (outcome was stored in cm). Longer line

indicated greater (longer) pain distribution. However, in the last two trials of each block participants were asked to

rate pain intensity by using NRS.

Figure 1. Study structure. (A) The study involved three experimental groups and one control group. Each group received the
same number of nociceptive stimuli and the same number of electrodes activated for each participant (”1” or „2” indicates the
number of activated electrodes). During the "experience" phase, participants underwent three blocks of trials during which
noxious stimuli were delivered to them. Each block consisted of 20 trials where participants rated pain distribution using pain
distribution rating task and 2 additional trials (always the last ones) where they rated pain intensity using a Numeric Rating Scale
(NRS). In the "control" and "suggestion only" groups, the experience blocks featured pseudo-conditioning: the color-based
conditioned stimuli [CS+ (green or blue), CS- (green or blue)] were not systematically associated with the unconditioned stimuli
(US). Specifically, CS+ was not consistently paired with stimulation using two electrodes (US+), nor was CS- consistently paired

with stimulation using one electrode (US-). In the classical conditioning groups, CS+ was always linked to US+ (stimulation with two electrodes), and CS- was always linked to US- (stimulation with one electrode). Of these, only one group received verbal suggestion (indicated as a pictogram) and was informed about the US-CS contingencies. During the testing block, only US- (stimulation with one electrode) was applied regardless of the CS presented. Additionally, a novel stimulus [CS_{GEN} (cyan)], which had not been used during experience blocks was introduced to test for stimulus generalization. **(B)** Each given trial from experience blocks was structured with the following components: an inter-trial interval (ITI), a CS presentation period (2s) linked with US stimulation (lightning), and a rating period. During the rating period, participants either rated the pain distribution (PD) or pain intensity. To rate pain distribution participants were drawing a digital line on the screen to match the length of the pain distribution with the length of the line. Pain intensity was rated on the NRS scale displayed on the screen. **(C)** The trial design from the testing block was similar to the experience block, however, here always only one electrode was activated regardless of CS presentation and one new stimulus (CS_{GEN}) was mixed with CS+ and CS-.

2.7.5. Exit phase

After the main phase of the study, participants were asked to fill the exit questionnaire with five manipulation check questions. Questions were related to study aim, meaning of visual stimuli, contingencies between colors and perceived pain. List of manipulation check questions and results are presented in the **Appendix**.

2.8. Preprocessing and statistical analyses

Data saved in a raw format via PsychoPy software were first preprocessed by using MATLAB. Relevant data from conditioning and testing phases were extracted and re-scaled to real cm values (PD ratings). All analyses were performed using R Statistical Software (v4.1.2; R Core Team 2021) [38] and with $\alpha = 0.05$.

Primary analyses were performed in line with the *a priori* released plan (https://osf.io/3qezt). First, a mixed model ANOVA was performed on the primary outcome "PD" with "group" as a between-subject factor and "condition" (CS+, CS-, CS_{GEN}) as the within-subject factor. In case of significant interaction effects, planned within-subject contrasts were performed to address specified predictions: Paired *t* test contrasts between CS+ and CS- were applied to test outlined hypotheses: effects in groups with conditioning only, and conditioning with verbal suggestion. It was assumed that effect should not be present in the control group and there was no specific hypothesis regarding the group with verbal suggestion only. Similar series of paired *t* tests contrasts between CS- and CS_{GEN} were performed to test for the presence of stimulus generalization effects within each of the group.

Additionally, mean within-subject main effects (differences between CS- and CS+ conditions) as well as generalization effects (differences between CS- and CS_{GEN} conditions) from experimental groups were contrasted with the effects from the control group. Lastly, for the purpose of exploratory analyses, the same set of analyses was performed on pain intensity ratings. To investigate if evoked effects observed in the experimental groups were contaminated by potential spatial summation of pain (higher pain intensity in CS+ vs. CS-), Pearson correlation coefficients were calculated between the effects in PD and effects on pain intensity. Between-group differences in descriptive statistics were explored by using one-way ANOVA with "group" as a between-subjects factor.

3. Results

Two participants were excluded from the study because the required intensity of perceived pain was not reached during the calibration phase (5/10 on NRS scale) within a preset intensity range (40 mA). Data from one participant is not included due to technical problems during the study. A total number of 94 datasets from 94 participants who completed the experiment were used to run statistical analysis. Descriptive characteristics are shown in **Table 1.** No significant between-group differences regarding these variables were found **(Appendix).**

274 **Table 1. Descriptive characteristics of the study population**

Variable	conditioning + vs	conditioning	vs	control
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
Age (years)	22.13 (2.91)	21.09 (2.19)	20.75 (1.57)	21.78 (1.65)
Fear of pain (NRS 0-10)	2.21 (1.79)	1.96 (1.52)	2.46 (1.91)	2.13 (2.12)
FPQ-III: Severe pain	29.21 (6.97)	29.48 (7.22)	32.17 (7.04)	30.91 (8.15)
FPQ-III: Medical pain	23.04 (6.05)	23.13 (7.55)	23.71 (7.19)	24.17 (9.05)
FPQ-III: Minor pain	16.92 (4.91)	16.09 (5.96)	16.08 (3.72)	15.83 (5.46)
FPQ-III total score	69.17 (14.95)	68.70 (15.48)	71.96 (14.64)	70.91 (18.82)
Body mass (kg)	72.08 (12.35)	70.04 (11.85)	70.21 (15.01)	72.63 (14.59)
Height (cm)	173.37 (7.73)	172.39 (9.09)	175.67 (8.58)	174.09 (8.29)
Stimulus Intensity (mA)	19.00 (7.48)	16.74 (5.20)	17.54 (5.35)	19.00 (6.45)
	N (24)	N (23)	N (24)	N (23)
Sex	F = 12; M = 12	F = 13; M = 10	F = 11; M = 13	F = 11; M = 12

275   F- Female, M -Male, R - right, L – left, NRS - numeric rating scale, FPQ-III – Fear of pain questionnaire, SD – standard deviation, vs – verbal suggestion

3.1 Primary analyses: pain distribution

Means and standard deviations for PD and pain intensity in all groups are presented in **Table 2**. The mixed model ANOVA showed a significant effect for the factor "condition" ($F_{(2,180)} = 11.35$, $p < 0.0001$, $\eta_p^2 = 0.11$) and a significant "group" × "condition" interaction was found ($F_{(6,180)} = 3.54$, $p = 0.002$, $\eta_p^2 = 0.11$). A within-group t test comparisons revealed that CS+ condition was, in general, characterized by more extended PD compared to CS- condition (**Figure 2 and 3**). This direction was present in the conditioning + verbal suggestion group ($t_{(23)} = 3.59$, $p < 0.01$, $d_z = 0.73$), conditioning only group ($t_{(22)} = 2.47$, $p < 0.05$, $d_z = 0.52$), verbal suggestion only group ($t_{(24)} = 2.78$, $p < 0.05$, $d_z = 0.56$) but no significant differences between CS+ and CS- were found in the control group ($t_{(21)} = -0.87$, $p = 0.39$, $d_z = -0.19$). Comparison between CS_{GEN} and CS- showed significantly more extended PD in CS_{GEN}, however only in the "verbal suggestion only" group ($t_{(24)} = -2.12$, $p < 0.05$, $d_z = -0.42$). No significant differences between CS_{GEN} and CS- and stimuli were observed in "conditioning only" ($t_{(22)} = -0.51$, $p = 0.61$, $d_z = -0.11$), "conditioning + verbal suggestion" ($t_{(23)} = -1.93$, $p = 0.07$, $d_z = -0.39$) and "control" group ($t_{(21)} = 1.43$, $p = 0.17$, $d_z = 0.31$).

302 **Table 2. Means and standard deviations for pain distribution and pain intensity in all groups**

Outcome	Group	Block 1		Block 2		Block 3		Testing		
		CS+	CS-	CS+	CS-	CS+	CS-	CS+	CS-	CS$_{GEN}$
		Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
PD	conditioning with suggestion	8.05 (4.99)	4.38 (3.51)	8.00 (5.50)	3.64 (3.24)	7.64 (5.17)	3.39 (3.53)	5.58 (4.48)	4.25 (3.83)	4.74 (4.14)
	conditioning	6.63 (4.64)	4.04 (4.09)	6.07 (4.04)	3.65 (3.62)	6.00 (4.01)	3.31 (3.66)	4.26 (4.58)	3.64 (3.91)	3.75 (3.97)
	suggestion	5.91 (4.15)	6.29 (4.38)	6.04 (4.52)	6.18 (4.63)	5.67 (3.94)	5.71 (4.20)	4.32 (3.25)	3.82 (3.00)	4.26 (3.33)
	control	5.96 (3.59)	5.64 (3.54)	6.53 (4.64)	6.56 (4.46)	6.35 (4.58)	6.01 (4.33)	3.38 (3.23)	3.54 (3.66)	3.07 (2.82)
PI	conditioning with suggestion	4.83 (1.69)	3.33 (2.04)	4.92 (1.82)	2.87 (1.62)	4.63 (1.91)	2.50 (1.35)	3.88 (2.09)	3.08 (1.74)	3.37 (2.16)
	conditioning only	4.87 (1.60)	2.78 (1.70)	4.96 (1.64)	2.48 (1.47)	4.61 (1.64)	2.09 (1.51)	3.00 (1.76)	2.48 (1.44)	3.00 (1.65)
	suggestion	3.92 (2.04)	3.92 (1.89)	3.63 (1.95)	3.92 (2.21)	3.54 (2.28)	3.33 (1.71)	3.80 (1.85)	3.40 (1.56)	3.60 (2.12)
	control group	3.22 (1.65)	3.57 (1.75)	3.61 (1.80)	3.52 (1.97)	3.09 (1.62)	3.13 (1.77)	2.77 (1.41)	2.55 (1.30)	2.64 (1.50)

CS+: Mean ratings preceded by visual stimuli associated with large pain distribution (US+) in conditioning only group and conditioning + verbal suggestion group. Mean ratings preceded by
visual stimuli associated by verbal suggestion with large pain distribution (US+) in verbal suggestion only group. CS-: Mean ratings preceded by visual stimuli associated with small pain
distribution (US-) in conditioning only group and conditioning + verbal suggestion group. Mean ratings preceded by visual stimuli associated by verbal suggestion with small pain distribution
(US-) in verbal suggestion only group. CS$_{GEN}$: Mean ratings preceded by novel visual stimuli not previously associated with any pain distribution by conditioning or verbal suggestion. PD-
pain distribution. PI- pain intensity. SD – standard deviation. Block (1,2,3)- three blocks of trials during the experience phase. Testing – trials during testing phase.

Figure 2. Mean ratings of perceived area of pain (pain distribution) preceded by visual stimuli (CS+, CS-) in every trial.
Trials: First block of experimental trials (1-10), Second block of experimental trials (11-20), Third block of experimental trials
(21-30), Testing phase (30-33). CS+: Mean ratings preceded by visual stimuli associated (1-30 trials) with large pain distribution
(US+) in conditioning only group and conditioning + verbal suggestion group. Mean ratings preceded by visual stimuli associated
by verbal suggestion (30-33 trials) with large pain distribution (US+) in verbal suggestion only group. CS-: Mean ratings preceded
by visual stimuli associated (1-30 trials) with small pain distribution (US-) in conditioning only group and conditioning + verbal
suggestion group. Mean ratings preceded by visual stimuli associated by verbal suggestion (30-33 trials) with small pain
distribution (US-) in verbal suggestion only group. Error bars represent the standard error of the mean.

Figure 3. Mean ratings of perceived area and intensity of pain. CS+: Mean ratings preceded by visual stimuli associated with
large pain distribution (US+) in conditioning only group and conditioning + verbal suggestion group. Mean ratings preceded by
visual stimuli associated by verbal suggestion with large pain distribution (US+) in verbal suggestion only group. CS-: Mean
ratings preceded by visual stimuli associated with small pain distribution (US-) in conditioning only group and conditioning +
verbal suggestion group. Mean ratings preceded by visual stimuli associated by verbal suggestion with small pain distribution
(US-) in verbal suggestion only group. CS$_{GEN}$: Mean ratings preceded by novel visual stimuli not previously associated with any

3.1.2 Exploratory analyses

To complement within-group contrasts conducted on PD variable, additional (exploratory) between-group comparisons were performed which showed significantly larger effect sizes (difference between CS+ and CS-) in PD between the conditioning + verbal suggestion group and the control group ($t_{(44)} = 3.51$, $p < 0.01$, $d_z = 1.04$), between the conditioning only group and the control group ($t_{(43)} = 2.50$, $p < 0.05$, $d_z = 0.75$), and also between the suggestion only group and the control group ($t_{(45)} = 2.57$, $p < 0.05$, $d_z = 0.75$).

The mixed model ANOVA performed on pain intensity showed a divergent pattern of results compared to analysis of PD data. Specifically, significant effect was found for the factor "condition" ($F_{(2,180)} = 8.27$, $p < 0.001$, $\eta_p^2 = 0.08$) but not main effect of "group" ($F_{(3,90)} = 1.92$, $p = 0.13$, $\eta_p^2 = 0.06$) nor "condition" × "group" interaction ($F_{(6,180)} = 0.78$, $p = 0.59$, $\eta_p^2 = 0.03$) was observed, indicating that the difference between CS+ and CS- in pain intensity was consistent across all four groups. Due to lack of effects in this outcome, no further exploratory tests were performed.

Correlations between evoked effects in PD (individual differences between CS+ and CS-) and differences in pain intensity were not significant in all groups: conditioning + verbal suggestion ($r = 0.39$, $p = 0.06$), conditioning only ($r = 0.00$, $p > 0.99$), verbal suggestion only ($r = -0.04$, $p = 0.86$), control group ($r = -0.26$, $p = 0.23$).

4. Discussion

The results confirm that PD ratings preceded by CS+ or CS- stimuli were significantly different in each experimental group but not in the control group. Moreover, significant differences in PD between groups were found. Such results show that PD can be learned by classical conditioning and that the perceived PD can be influenced by verbal suggestion. The largest effect size was observed in the conditioning with verbal suggestion group, which is in line with the hypothesis that verbal suggestion can enhance conditioning processes. Only in the verbal suggestion group, a significant effect of generalization was found. No significant differences in pain intensity preceded by CS+ or CS- stimuli were found within groups and no significant correlations were observed between pain distribution and pain intensity ratings.

4.1 Mechanisms of pain distribution

Pain not limited to a given body area is a common feature of CWP. This debilitating condition is often linked with fatigue, sleep disturbances, and cognitive impairment [11] and it has been proposed that peripheral as well as central mechanisms can be attributed to the widespread distribution of reported pain [10,15]. For instance, research suggests central sensitization [11,30], muscular dysfunction [9,19], bifurcation of nociceptive afferents from two different tissues [41] or deficits in endogenous pain-modulation [22,24] as some of the mechanisms explaining CWP. However, experiments on humans and animals also pointed out that experimentally induced (e.g. heat) pain is not precisely represented within the neuroaxis. Spinal cord imaging on rats demonstrated that relatively strong noxious input leads to a widespread rostro-caudal and ipsilateral-contralateral spread of neural activity within the spinal laminae that includes nociceptive-specific neurons and wide-dynamic range (WDR) neurons [12]. Interestingly, this activity spread can manifest as the radiation of pain induced by noxious heat in humans. For instance, in an experiment with two thermal probes attached to the skin and separated by 10 cm, pain was often reported under both probes even though one probe was set to deliver a neutral temperature of 35°C [36]. The most recent report with stimulation of 2.56 cm^2 area of the skin showed that 49°C can produce radiation of pain that is, on average, 12 times greater than the size of stimulation [2]. Thus, these psychophysical findings suggest that the human nervous system has the capacity to construct (and even interpolate) complex patterns of pain based on varied sources of available nociceptive information, including population coding and neuronal recruitment [2,12,36].

It seems that the potential of the human neuroaxis to form and maintain these complex patterns of pain distribution can be shaped by repeated interactions with the internal as well as external environments specific to a given individual. For instance, ongoing nociception from various foci may interact with each other or with neutral environmental signals. However, this hypothetical framework for understanding the formation of pain distribution has largely been overlooked in the literature. The present study is the first to test if classical conditioning could contribute to the acquisition and maintenance of widespread pain, although some translational studies have provided hints that learning can shape pain distribution. Indeed, in the study by Doménech-García et al. [16], participants who reported persistent pain in the shoulder in the past experienced a larger area of referred pain that was triggered by a remote stimulus applied to a previously non-injured area, a pattern not observed in the healthy controls. These results may provide indirect evidence for the involvement of learning mechanisms, as additional pain stimuli could serve as conditioned stimuli eliciting a learned pain response, consistent with the findings of the present study. The engagement of learning mechanisms could account for the differences in pain distribution areas elicited by noxious stimuli of identical intensity and applied to the same region in healthy participants.

4.2 Classical conditioning

The main novelty of the current experiment is a demonstration that enlarged pain distribution can persist as a result of classical conditioning. Results from the primary analysis indicate a significant difference in perceived PD both between and within groups. Results from the conditioning + verbal suggestion and conditioning only group show that participants perceived a wider PD in the testing phase when noxious stimuli delivered by only one electrode were preceded by stimuli previously associated with large PD (CS+), i.e. delivered by two electrodes as compared to noxious stimuli preceded by stimuli previously associated with small PD (CS-), i.e. delivered by one electrode. First, these results indicate that pain distribution elicited by electrocutaneous noxious stimulation was successfully modulated by classical conditioning. Secondly, the size of the stimulated area in the large PD (US+) was twice as large (two electrodes were active) as that of the small PD (US-) (one electrode was active) with a 2:1 proportion, indicating that a conditioned stimulus (CS+), followed by a small unconditioned stimulus (US-), can elicit pain in a larger area than the unconditioned stimuli itself. However, given the implemented method of noxious stimulation by using electrodes, the stimulated body area during the experiment was not as extensive as in many of the possible injuries from which many patients could suffer. Thirdly, differences in perceived PD between CS+ and CS- were higher in the conditioning + verbal suggestion group as compared to the conditioning only group indicating that suggestion could enhance the conditioned response, here pain distribution. Such results are in line with previous studies on pain conditioning, showing a modulation of pain intensity through suggestion [35]. However, differences in the perceived PD between CS+ and CS- were higher in the conditioning only group than in the verbal suggestion only group, suggesting that the learning effect could be more robust in the modulation of PD than suggestion alone, indicating the need for the first-hand experience to observe the conditioning effect.

Previous findings show that pain intensity can be modulated through classical conditioning [5,25,32]. However, no significant difference in perceived pain intensity between the four groups was found in the current study. The lack of a significant difference in perceived pain intensity between the groups could have been caused by the fact that authors did not manipulate pain intensity. The difference in pain intensity between a large pain distribution (US+) and a small pain distribution (US-) was the same for all groups. Furthermore, the stimuli intensity was calibrated individually. This indicates that effect of spatial summation of pain [27] was successfully controlled and did not contaminate the "experience" phase. Moreover, a lack of any significant correlation between pain intensity and pain distribution suggests that pure pain distribution was successfully manipulated and that the results are not biased by pain intensity.

4.3 Generalization

Generalization and discrimination of stimuli and reaction are essential parts of learning through conditioning [28]. Current results do not provide evidence for a significant generalization effect of the conditioned pain distribution, only in verbal suggestion group a significant effect was observed. Therefore, a lack of generalization of the learning of pain distribution seems implausible. However, most previous studies focused on the generalization of a pain related fear reaction [18,31,46] and not on the generalization of pain intensity itself [23]. Thus, more evidence is needed to support the hypothesis that both pain distribution and pain intensity can be generalized as a conditioned reaction. In this experiment, the cyan color (RGB (19, 236, 236)) was chosen as the generalization stimulus because the color is embedded "in-between" blue (RGB (19, 19, 236)) and green (RGB (19, 236, 19)). Generalization depends strongly on the physical properties of similarity of presented novel stimuli to conditioned stimuli, such as the intensity of stimulation (the total activation of receptors in the sense organs), and how a given amount of stimulation is distributed among receptors [17]. Novel stimuli used in the current study could lack crucial physical similarities to conditioned stimuli to elicit a generalized response.

4.5 Limitations and future directions

This study has two main limitations. First, it only used electrocutaneous stimuli, leaving a question open whether other PD modalities can be modulated by learning processes. If PD can be modulated by learning mechanisms using other types of stimuli as extensively as the intensity of pain [25], future research could significantly advance our understanding of pain "chronification". Second, the design with two separate electrodes limits information on the extent of the conditioned responses. Future research should explore the magnitude of the difference between US+ and US- required to condition pain sensation over a larger area, as well as the overall extent of this effect.

4.6 Conclusion

The current results confirm that pain distribution can be learned through classical conditioning and influenced by suggestion. Clinically, these findings contribute to a better understanding of the spatial features of pain, especially when patients experience pain across a more widespread area than would be expected based on the actual site of nociceptive foci. Such understanding is crucial for the development of new pain treatments. Moreover, the role of verbal suggestion in the clinical environment should not be neglected, as it can not only enhance conditioning effects but also contribute to enlarged pain distribution reports as a standalone mechanism.

5. Acknowledgments

The authors declare no conflict of interest. The authors are supported by the National Science Center within grant no. 2020/37/B/HS6/04196.

6. References

[1]   Adamczyk WM, Luedtke K, Buglewicz E, Bąbel P. Pain rewarded: hyperalgesic and allodynic effect of operant conditioning in healthy humans—protocol for a systematic review and meta-analysis. Syst Rev 2018;7:93.

[2]   Adamczyk WM, Ramu V, Jackson C, Schulze G, Goldschneider KR, Kashikar-Zuck S, King CD, Coghill RC. Radiation of pain: Psychophysical evidence for a population coding mechanism. 2024. doi:10.1101/2024.04.02.587666.

[3]   Adamczyk WM, Szikszay TM, Nahman-Averbuch H, Skalski J, Nastaj J, Gouverneur P, Luedtke K. To Calibrate or not to Calibrate? A Methodological Dilemma in Experimental Pain Research. The Journal of Pain 2022;23:1823–1832.

[4]   Adamczyk WM, Wiercioch-Kuzianik K, Bajcar EA, Bąbel P. Rewarded placebo analgesia: A new mechanism of placebo effects based on operant conditioning. European Journal of Pain 2019;23:923–935.

[5]   Bąbel P, Adamczyk W, Świder K, Bajcar EA, Kicman P, Lisińska N. How Classical Conditioning Shapes Placebo Analgesia: Hidden versus Open Conditioning. Pain Medicine 2018;19:1156–1169.

[6]   Bąbel P, Bajcar EA, Adamczyk W, Kicman P, Lisińska N, Świder K, Colloca L. Classical conditioning without verbal suggestions elicits placebo analgesia and nocebo hyperalgesia. PLoS ONE 2017;12:e0181856.

[7]   Bąbel P, Bajcar EA, Adamczyk W, Kicman P, Lisińska N, Świder K, Colloca L. Classical conditioning without verbal suggestions elicits placebo analgesia and nocebo hyperalgesia. PLoS ONE 2017;12:e0181856.

[8]   Bajcar EA, Adamczyk WM, Wiercioch-Kuzianik K, Bąbel P. Nocebo hyperalgesia can be induced by classical conditioning without involvement of expectancy. PLoS ONE 2020;15:e0232108.

[9]   Bengtsson A, Henriksson KG. The muscle in fibromyalgia--a review of Swedish studies. J Rheumatol Suppl 1989;19:144–149.

[10] Clauw DJ. Diagnosing and treating chronic musculoskeletal pain based on the underlying mechanism(s). Best Practice & Research Clinical Rheumatology 2015;29:6–19.

[11] Clauw DJ, Crofford LJ. Chronic widespread pain and fibromyalgia: what we know, and what we need to know. Best Practice & Research Clinical Rheumatology 2003;17:685–701.

[12] Coghill RC. The Distributed Nociceptive System: A Framework for Understanding Pain. Trends in Neurosciences 2020;43:780–794.

[13] Coghill RC, McHaffie JG, Yen Y-F. Neural correlates of interindividual differences in the subjective experience of pain. Proc Natl Acad Sci USA 2003;100:8538–8542.

[14] Defrin R, Urca G. Spatial summation of heat pain: a reassessment. Pain 1996;66:23–29.

[15] DeSantana JM, Sluka KA. Central mechanisms in the maintenance of chronic widespread noninflammatory muscle pain. Current Science Inc 2008;12:338–343.

[16] Doménech-García V, Skuli Palsson T, Boudreau SA, Herrero P, Graven-Nielsen T. Pressure-induced referred pain areas are more expansive in individuals with a recovered fracture. Pain 2018;159:1972–1979.

[17] Ghirlanda S, Enquist M. A century of generalization. Animal Behaviour 2003;66:15–36.

[18] Glogan E, Van Vliet C, Roelandt R, Meulders A. Generalization and Extinction of Concept-BasedPain-Related
Fear. The Journal of Pain 2019;20:325–338.

[19] Henriksson KG. Is fibromyalgia a distinct clinical entity? Pain mechanisms in fibromyalgia syndrome. A
myologist's view. Best Practice & Research Clinical Rheumatology 1999;13:455–461.

[20] Hunt I. The prevalence and associated features of chronic widespread pain in the community using the
"Manchester" definition of chronic widespread pain. Rheumatology 1999;38:275–279.

[21] Jensen KB, Kaptchuk TJ, Kirsch I, Raicek J, Lindstrom KM, Berna C, Gollub RL, Ingvar M, Kong J.
Nonconscious activation of placebo and nocebo pain responses. Proc Natl Acad Sci USA 2012;109:15959–
15964.

[22] Julien N, Goffaux P, Arsenault P, Marchand S. Widespread pain in fibromyalgia is related to a deficit of
endogenous pain inhibition. Pain 2005;114:295–302.

[23] Koban L, Kusko D, Wager TD. Generalization of learned pain modulation depends on explicit learning. Acta
Psychologica 2018;184:75–84.

[24] Kosek E, Ekholm J, Hansson P. Modulation of pressure pain thresholds during and following isometric
contraction in patients with fibromyalgia and in healthy controls. Pain 1996;64:415–423.

[25] Madden VJ, Harvie DS, Parker R, Jensen KB, Vlaeyen JWS, Moseley GL, Stanton TR. Can Pain or
Hyperalgesia Be a Classically Conditioned Response in Humans? A Systematic Review and Meta-Analysis.
Pain Med 2015:pnv044.

[26] Mansfield KE, Sim J, Jordan JL, Jordan KP. A systematic review and meta-analysis of the prevalence of chronic
widespread pain in the general population. Pain 2016;157:55–64.

[27] Marchand S, Arsenault P. Spatial summation for pain perception: interaction of inhibitory and excitatory
mechanisms. Pain 2002;95:201–206.

[28] Mclaren IPL, Mackintosh NJ. Associative learning and elemental representation: II. Generalization and
discrimination. Animal Learning & Behavior 2002;30:177–200.

[29] McNeil DW, Rainwater AJ. Development of the Fear of Pain Questionnaire-III. Journal of Behavioral Medicine
1998;21:389–410.

[30] Meeus M, Nijs J. Central sensitization: a biopsychosocial explanation for chronic widespread pain in patients
with fibromyalgia and chronic fatigue syndrome. Clin Rheumatol 2007;26:465–473.

[31] Meulders A, Jans A, Vlaeyen JWS. Differences in pain-related fear acquisition and generalization: an
experimental study comparing patients with fibromyalgia and healthy controls. Pain 2015;156:108–122.

[32] Miguez G, Laborda MA, Miller RR. Classical conditioning and pain: Conditioned analgesia and hyperalgesia.
Acta Psychologica 2014;145:10–20.

[33] Moseley GL, Vlaeyen JWS. Beyond nociception: the imprecision hypothesis of chronic pain. Pain 2015;156:35–
38.

[34] Peirce JW. PsychoPy—Psychophysics software in Python. Journal of Neuroscience Methods 2007;162:8–13.

[35] Petersen GL, Finnerup NB, Colloca L, Amanzio M, Price DD, Jensen TS, Vase L. The magnitude of nocebo
effects in pain: A meta-analysis. Pain 2014;155:1426–1434.

[36] Quevedo AS, Coghill RC. Filling-In, Spatial Summation, and Radiation of Pain: Evidence for a Neural
Population Code in the Nociceptive System. Journal of Neurophysiology 2009;102:3544–3553.

[37] Quevedo AS, Mørch CD, Andersen OK, Coghill RC. Lateral inhibition during nociceptive processing. Pain 2017;158:1046–1052.

[38] R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing 2021.

[39] Raghuraman N, Wang Y, Schenk LA, Furman AJ, Tricou C, Seminowicz DA, Colloca L. Neural and behavioral changes driven by observationally-induced hypoalgesia. Sci Rep 2019;9:19760.

[40] Raja SN, Carr DB, Cohen M, Finnerup NB, Flor H, Gibson S, Keefe FJ, Mogil JS, Ringkamp M, Sluka KA, Song X-J, Stevens B, Sullivan MD, Tutelman PR, Ushida T, Vader K. The revised International Association for the Study of Pain definition of pain: concepts, challenges, and compromises. Pain 2020;161:1976–1982.

[41] Sinclair, D. C., Weddell, G., & Feindel, W. H. Referred pain and associated phenomena. Brain 1948;71:184–211.

[42] Staud R, Vierck CJ, Robinson ME, Price DD. Spatial summation of heat pain within and across dermatomes in fibromyalgia patients and pain-free subjects. Pain 2004;111:342–350.

[43] Świder K, Bąbel P. The effect of the sex of a model on nocebo hyperalgesia induced by social observational learning. Pain 2013;154:1312–1317.

[44] Świder K, Bąbel P. The Effect of the Type and Colour of Placebo Stimuli on Placebo Effects Induced by Observational Learning. PLoS ONE 2016;11:e0158363.

[45] Szikszay TM, Adamczyk WM, Luedtke K. The Magnitude of Offset Analgesia as a Measure of Endogenous Pain Modulation in Healthy Participants and Patients With Chronic Pain: A Systematic Review and Meta-Analysis. The Clinical Journal of Pain 2019;35:189–204.

[46] Vandael K, Meulders M, Vanden Boer C, Meulders A. The relationship between fear generalization and pain modulation: an investigation in healthy participants. Scandinavian Journal of Pain 2019;20:151–165.

[47] Wiercioch-Kuzianik K, Bąbel P. Color Hurts. The Effect of Color on Pain Perception. Pain Medicine 2019;20:1955–1962.

[48] World Medical Association. World Medical Association Declaration of Helsinki. Ethical principles for medical research involving human subjects. Bull World Health Organ 2001;79:373–374.