Distal super-enhancer drives aberrant *CXCL13* expression in Cancer cells driving growth and p53 dysregulation via CXCR5-CXCL13 axis

Santosh Kumar Gothwal[1,2*], Pieta K. Mattila[3,4,5] Jacqueline H Barlow[6]

[1] Institute of Frontier Science Initiative, Kanazawa University, Japan

[2] Central Research Resource Center, Cancer Research Institute, Kanazawa University,

[3] Institute of Biomedicine and MediCity Research Laboratories, University of Turku, Turku, Finland

[4] InFLAMES Research Flagship, University of Turku, Turku, Finland.

[5] Turku Bioscience, University of Turku and Åbo Akademi University, Turku, Finland.

[6] Department of Microbiology and Molecular Genetics, University of California, Davis, United States

* Corresponding author: skgothwal@staff.kanazawa-u.ac.jp & santoshgothwalbio@gmail.com

Key words- CXCL13, super-enhancer, CXCR5, B-lymphoma, p53, GC-migration

Key Points

1. Super-enhancer near *CCNG2* region interacts with *CXCL13-TSS* driving CXCL13 in cancers.

2. Aberrant CXCL13 prevents CXCR5-mediated migration of B-lymphomas and promotes growth and p53 dysregulation in CXCR5+ cells

3. CXCR5-CXCL13 axis impairs p53 target gene expression and promotes tumor growth

Abstract

The CXCL13 chemokine plays a crucial role in guiding B cell migration to the light zones (LZs)

during the germinal center (GC) reaction. While *CXCL13* expression is absent in most cell types,

aberrant amplification of the CXCR5-CXCL13 signaling is observed in various cancers,

including germinal center-derived B-lymphomas (GCDBL), colorectal adenocarcinoma

(COAD), and liver hepatocellular carcinoma (LIHC). However, the molecular mechanisms

underlying abnormal *CXCL13* transcription in cancer cells and its functional consequences

remain elusive. We identify DNA-CpG methylation binding protein 1 (MBD1) as a suppressor of

CXCL13 expression. Chromosomal conformation capture (3C) analysis reveals a distal super-

enhancer located near *CCNG2* that interacts with the *CXCL13* promoter in GCDBL, suggesting

that enhancer-hijacking drives the aberrant expression. Our functional validation demonstrates

that CXCR5-CXCL13 signaling suppresses p53 and its target genes in GCDBLs, COAD, and

LIHC. Notably, CXCL13 in the GCDBL cell line Raji disrupts CXCR5-mediated migration, a

mechanism essential for (light zone) LZ-entry and affinity maturation of GC B cells. These

findings highlight the dual role of the CXCR5-CXCL13 axis in immune response and cancer

proliferation.

Introduction

Chemokine (C-X-C motif) ligand 13 (CXCL13) binds and exerts signaling through the C-X-C

chemokine receptor type 5 (CXCR5) and G proteins [1-4]. *CXCL13* is primarily expressed in

follicular dendritic cells (FDCs), follicular helper T cells (Tfh), CD4+ T cells, CD8+ T cells,

stromal cells and macrophages and regulates the innate and humoral immune repsonse [1-3, 5-9].

But aberrant *CXCL13* expression is also observed in hematological malignances such as B-cell

chronic lymphocytic leukemia (B-CLL) and multiple myeloma (MM) [10-13]. Among solid

cancers, *CXCL13* expression is observed in non-small cell lung carcinoma (NSCLC), breast

cancers, prostate cancers and *Helicobacter pylori* associated gastric lymphomas [11, 14-18].

Aberrant CXCR5-CXCL13 signaling is evident in autoimmune diseases such as rheumatoid

arthritis, [19, 20] , multiple sclerosis [21], systematic lupus erythematosis [22], primary Sjögren's

syndrome [23] and myasthenia gravis [24, 25].

CXCL13 signaling is known to interfere the chemotherapeutic response and elevated

CXCL13 expression by cancer cells promotes autocrine and paracrine effects in the tumor

microenvironment resulting [26]. CXCL13 secreted by MM cells further induces CXCL13 in

surrounding macrophages by Bruton's tyrosine kinase (BTK) signaling. In turn, macrophages

induce *CXCL13* expression in MM through TGF-β signaling [13]. Benzo(a)pyrene (BaP), a

carcinogen found in tobacco and air pollution, induces *CXCL13* in NSCLC [27]. *CXCL13*

knockdown in MM and NSCLC resulted in slower tumor growth in xenograft experiments **[13,**

27]. The CXCR5-CXCL13 axis exhibits resistance to chemotherapeutic drugs including

bortezomib in MM [28], and 5-flurouracil in colorectal cancer [29] and mantle cell lymphoma

[30]. In breast cancer, CXCR5-CXCL13 axis promotes metastasis by regulating the epithelial to

mesenchymal transition (EMT) [31]. CXCR5-CXCL13 signaling promotes tumorigenesis in

PTEN deficient cancers by Protein kinase C (PKC) signaling [32]. Loss of *PTEN* can induce

CXCL13 expression by NF-kB signaling [32]. *PTEN* is a p53 dependent gene, suggesting that

CXCR5-CXCL13 signaling interferes with p53 regulation, however whether the CXCR5-

CXCL13 axis promotes tumorigenesis by interfering with other p53 target genes is not known.

Despite the widespread expression of *CXCL13* in hematological and solid cancers, the

mechanism(s) driving aberrant *CXCL13* expression in cancer cells is elusive. Little is known about

the factors directly regulating the transcriptional axis of *CXCL13*. TGF-β-induced SOX4 promotes

the *CXCL13* expression during Th2 cells differentiation and retinoic acid and neuronal signaling

induces the CXCL13 expression in murine embryonic stromal cells [7, 33]. Enhancer tethering has

been suggested as a key mechanism for oncogene expression in cancers {Bi, 2020 #127;Dixon,

2012 #121;Roe, 2017 #128. Whether the CXCL13 promoter interacts with a distal enhancer that

drives *CXCL13* expression in cancer cells remains unknown. The chromatin insulating protein

CTCF links chromosome conformation and global gene-expression [34]. CTCF binding to DNA

is driven by local DNA methylation levels and Ten eleven translocation (TET) enzymes [35, 36].

DNA CpG methylation and MBD1 drive transcriptional suppression [37, 38]. Whether MBD1,

TET enzymes, and CTCF regulate *CXCL13* expression remains unclear.

In this study, we identify a super-enhancer, surrounding the *CXCL13* locus

interacting with *CXCL13* locus in cancer cells. Our findings in the human Burkitt's lymphoma cell

line Raji, suggest a role for aberrantly expressed CXCL13 in preventing migration of GCDBL

cells towards external CXCL13, a LZ chemokine. CXCR5-CXCL13 axis suppressed p53 target

genes in B-lymphomas, COAD and LIHC. These findings highlight the precise control of *CXCL13*

expression through epigenetic mechanisms and the regulation of enhancer-promoter interactions.

A loss of these regulatory processes in cancer is associated with the complex roles of CXCR5-

CXCL13 signaling in both hematological and solid tumors. Our study highlights the importance

of CXCL13 regulation in humoral immunity and tumorigenesis.

Material and Methods

Human B lymphoma cultures and treatments: Human B lymphoma Cell lines FL-18, FL-218,

FL-318, FL-518, Raji and Granta-519 were obtained from RIKEN, Japan and Kyoto university

hospital. All B lymphoma cell lines, THP1, LoVo and HC116 were cultured in RPMI (30264-85,

Nakalai) containing 10% FBS, 1% Penicillin, Streptomycin and Amphotericin B solution (Wako#

161-23181) in CO_2 incubator at 37°C under 5% CO_2 concentration. MCF7, HepG2 and 293T

cells were cultured in the DMEM containing 10% FBS, 1% Penicillin, Streptomycin and

Amphotericin B solution (Wako# 161-23181) in CO_2 incubator at 37°C under 5% CO_2

concentration. Treatment of cells with chemotherapeutics drugs Hydroxyurea (Product #085-

06653, Wako, Japan), Gemcitabine (product # S1714, Selleck, Japan) ATR inhibitor

(catalogue#KU55933; Selleck, Japan), NF-κB inhibitor (catalogue## S4902: Selleck, Japan),

AKT inhibitor (catalogue#S1037:Selleck, Japan) and etoposide (product #055-0843, Wako Japan),

bobocat339 (catalogue #SML2611, sigma), Bleomycin (Catalogue# B7116-1ML, Sigma), Nutlin-

3a (Catalogue# S8059, Selleck, Japan) were used as per defined timepoints and concentrations.

Short hairpin knockdown, and lentivirus preparation-

293T cells at nearly 60% confluency were transfected with a mix of *p-PAX2* (3 µg), *pVSVG* (3

µg), and target shRNA plasmids in *pLKO1-puromycin/blasticidin backbone* (3 µg). Plasmids were

mixed in 1 ml of Opti-Mem (Cat# 31985062, Gibco), then 27 µl of Polyethyleneimine (PEI)

solution (1 µg/ml) was added. After 20 minutes at room temperature, the mixture was used for

transfecting 293T cells. Cells were refreshed with DMEM medium after 24 hours, and culture

supernatants were collected 48 hours post-transfection, filtered with a 0.2 µM filter. Lentiviral

medium (1 ml) was used for transduction to 1 million Raji cells, followed by puromycin (1 µg/ml)

or Blasticidin (10 µg/ml) selection for resistance. Puromycin or Blasticidin was refreshed every

three days for one week. For production of CRISPR/Cas9 lentivirus, guideRNA cloned into the

Guide-it™ CRISPR/Cas9 Systems (Takarabio# 632601) were used instead of shRNA plasmids

follwed by the similar procedure and transduction to Raji cells.

B Cell Migration Assay: Transwell (Costar, REF#3421) was filled with RPMI (5% FBS) or RPMI

(5% FBS) medium containing rhCXCL12 (200 ng/ml) or rhCXCL13 (1000 ng/ml) (Biolegend).

Cell suspension of $5*10^5$ cells or $1.5*10^5$ cells for normal human B cells, isolated by institutional

approval and with manufacturers instruction, EasySep™ Human B Cell Isolation Kit

(STEMCELL technologies) were placed in transwell inserts for 6 hours at 37°C in CO_2 chamber.

Collected cells were lysed in 200 μl of CellTiter-Glo® 2.0 Reagent (Promega #G9242) and

allowed to stabilize for 10 minutes to measure luminescence signals. Absorbance was measured

using TECAN luminometer (code # MIC9414). A CXCL13-migration rescue experiment was

conducted in Raji cells, transfected with lentiviral particles containing CRISPR/Cas9 guide RNA

targeting exons 2 and 3 of *CXCL13*. Following transfection, cells were cultured for 4 days before

being utilized in a migration assay.

FACS analysis- Raji cells (1 * 10^6) were surface-stained with isotype controls or antibodies

(Supplementary Table 1) constituted in FACS buffer (PBS from Nakalai Tasque, 0.1% BSA, 0.1%

NaN3). After a 15-minute staining at room temperature, cells were washed, resuspended in 400 μl

FACS buffer, and mixed with 7-AAD before analysis. Flow cytometry was conducted using a

FACS Fortessa Flow cytometer (Becton Dickinson, San Jose, USA), analyzing 10,000 events with

data displayed as two-color plots (SSC-A/APC/PE/BV-421). Regarding CRISPR/Cas9 editing,

Raji cells were transfected with guide RNA targeting *CXCL13* exons 2 and 3, then sorted using a

ZSgreen marker.

Cloning, sequence analysis and overexpression- Full-length CXCR5 transcript amplification

was performed from Raji's cDNA with primers listed in supplementary Table 1. The PCR product

was digested with NheI and BamHI and cloned *pCMV-IRES-Puro vectors*, Sequenced and,

aligned with Uniprot accession (P32302). 5 μg of HpaI-linearized *pCMV-CXCR5-IRES-Puro* was

transfected into 1 million Raji cells via Nepagene 21 electroporation followed by Puromycin

selection (1 μg/ml) for a week to obtain stably transfected cells. Stable integration was confirmed

by PCR, flow cytometry and qPCR. The Nepagene 21 electroporation system was used according

to the instructions for the transient transfection of *pcDNA-MBD1*, *pcDNA-BCL6*, and *pcDNA* into

cell lines.

RNA isolation, cDNA preparation, RT-qPCR analysis and RNA-seq datasets from GEOR-

RNA from of normal Human B and B lymphoma cells was isolated (NucleoSpin® RNA kit,

#740984.50, Takara), cDNA was prepared (FSQ#201, Toyobo) and RT-qPCR was performed with

SYBR qPCR mix (QPS-201, Toyobo). The primer sequence of the analyzed transcripts is provided

(Supplementary table 1). Standard delta-delta Ct method was used for analysis of relative transcript

levels as described [39]. GSE171763 and GSE190319 datasets were used from GEO2R, NCBI.

Accession token (gfkvukuqnpsfvmx) and accession number GSE236240 were utilized fot the

cMYC and p53 ChIP-seq peaks in Raji.

Immunoprecipitation and Immunoblotting–2 million Raji cells were collected, washed with

PBS and lysed in 200 µl of RIPA buffer containing 1X protease inhibitors cocktail

(Sigma#11873580001). Lysed cells were spun at 13,000g for 10 min at 4 °C. The cell lysate from

supernatant was used for IP with 3 µg of CXCL13/IgG antibody (AF801, R&D system) and 20µl

precleared protein A beads (Invitrogen; 10001D, Invitrogen) for 2 hours at room temperatures,

followed by washing of the beads with RIPA buffer up to three times, 5 minutes each. The washed

beads were used for the elution and protein blotting. For Western blotting, cleared cell lysates

from 1 million cells (100 µl of RIPA) was reconstituted in 1X Lamellae buffer (Bio-Rad) with a

final concentration of 10% β-mercaptoethanol, followed by boiling at 95 °C for 5 min. After SDS

SDS-PAGE and transferring to a PVDF membrane using the Transblot® SD Semi-Dry

Electrophoretic Transfer Cell (Product #1703940, Bio-Rad), PVDF membranes were blocked in

5% skim milk for 15 min, washed with 1X TBST, and probed with primary antibody solution

overnight at 4 °C. The following day, the membrane was washed three times with 1X-TBST (10

min each) and then incubated with an HRP-conjugated secondary antibody for 1 hr. at room

temperature. After washing, the membrane was used to develop a chemiluminescent signal using

the Chemi-lumi one ultra (product #11644-40, Nakalai, Japan). The protein qualification was

performed using the ImageJ.

Chromosome Conformation Capture (3C) Assay: The 3C assay was performed as described

[39]. Two million Raji cells (-/+HU) were fixed in 1% PFA/PBS for 10 minutes, quenched with

glycine, and lysed in 500µl buffer (20 mM Tris-HCl, pH-8; 0.5% NP40; 1 mM EDTA, pH-8, 1X

protease inhibitor, Sigma #11873580001) for 30 minutes. 760 µl of 1X CutSmart buffer (NEB

#B6004) and 24µl of 10% SDS buffer were added, incubated at 37°C for 1 hour, followed by

EcoRI-digestion (40µl), overnight. The next day, 80 µl of 10% SDS was added, incubated at 65°C

(20 minutes), then mixed with 9.2 ml of 1X ligation-buffer, 100 µl of Triton X-100, and 1750 units

of Hi-T4™ DNA Ligase (NEB#M2622S). Ligation was performed at 16°C for 5 hours and

addition 16-18 hours at room-temperature. Samples were treated with 10 uL of RNase A (10

mg/ml) and 20 uL of proteinase K (20 mg/ml) before DNA purification. The DNA pellet was

dissolved in 600 µl TE buffer, 1µg of DNA was used for PCR (Takara, HS-Taq#R007A).

Chromatin Immunoprecipitation (ChIP)- ChIP assay was performed as described earlier [39].

Briefly, 2×10^7 cells were crosslinked in 50 ml of PBS with 1% formaldehyde at room temperature,

quenching with 2 ml of 2M glycine, followed by PBS washing, cells lysis (20mM Tris-HCL (pH-

8), 0.5% NP40, 1mM EDTA (pH-8), with 1X protease inhibitors, NaF (20 mM), and Na3Vo4 (1

µM). After brief centrifugation and nuclei collection, they were treated with SDS and 8M urea

buffers, followed by sonication (13 cycles, 30 sec on, 1 min off, Diagenode). Approximately 25

µg of chromatin was immunoprecipitated with 2 µg of antibodies, and 25 ul of Protein A/G Magna

ChIP beads (Sigma). The eluted samples and Input DNA were treated with RNAse and proteinase

K, then purified for qPCR analysis of target promoters. Primers and antibodies for qPCR are shown

in (supplementary table 1 and 2).

Results

Aberrant *CXCL13* expression in hematological malignancies and solid cancers

To identify cancer types exhibiting aberrant *CXCL13* expression, we explored the cancer genome

atlas (TCGA) sorting tumor samples by altered *CXCL13* status including deep deletion, shallow

deletion, diploid, gain, and amplification [40-42]. Genomic Identification of Significant Targets in

Cancer (GISTIC) analysis of *CXCL13* copy number alterations confirmed higher log2 copy

number values in gain and amplified groups compared to deep deletion, shallow deletion, and

diploid groups (Figure 1A). However, the levels of *CXCL13* mRNA in *CXCL13*-gain, *CXCL13*-

amplified, *CXCL13*-deletion and *CXCL13* shallow-deletion groups were not significantly different

from *CXCL13*-diploid control groups (Figure 1B). This suggests that genetic alterations such as

amplification and gains leading to increased copy number of *CXCL13* are not essential

determinants of enhanced *CXCL13* mRNA expression in cancers (Fig 1A-B) and implies that the

normally tight regulation of *CXCL13* expression becomes leaky in specific cancers.

To examine the expression of *CXCL13* mRNA among various cancers, we plotted *CXCR5*

and *CXCL13* mRNA Z-scores normalized with normal samples among cancer types (Figure 1C-

D). For both *CXCR5* and CXC13, mRNA log2 RNA-seq vs RSEM values above log2 cutoff value

of 1 were found in at least 1 sample for all cancers analyzed except for rectum adenocarcinoma

(READ) (Figure 1C,D). These results suggest that both the CXCR5 receptor and CXCL13 ligand

are potentially co-expressed in several cancers and may promote autocrine signaling (Figure 1C-

E). To evaluate whether *CXCL13* and *CXCR5* are upregulated in the same tumor, we randomly

identified seven CHOL samples exhibiting *CXCR5* mRNA log2 value of more than 0

(Supplementary Figure 1A). Four out of seven samples also exhibited *CXCL13* expression

(Supplementary Figure 1A), demonstrating that CHOL exhibits co-expression of *CXCR5* and

CXCL13 and suggests co-upregulation may occur in multiple cancer types (Figure 1C-E,

supplementary Figure 1A). Downstream signaling by the CXCR5-CXCL13 axis in these cancer

types may promote therapy resistance (Figure 1C-E). Of note, the overall survival (OS) was

notably lower among high-CXCL13 expressors compared to those with lower *CXCL13* levels as

seen across Acute Myeloid Leukemia, Glioblastoma, and Uveal Melanoma (Supplementary Figure

1B-D).

Based on widespread *CXCL13* expression in human cancers (Figure 1A- D), we examined

CXCL13 expression in human cell lines derived from hematological and solid malignancies.

CXCL13 mRNA and protein expression was confirmed among Raji (Burkitt's lymphoma), Granta-

519 (mantle cell lymphoma), FL-518 (Follicular Lymphoma), A549, MCF7, HepG2 (LIHC),

HCT116, and LoVo (COADs) MDA-MB231 and THP1 cells (Figure 1F-H). These observations

support the notion of aberrant *CXCL13* expression in hematological and solid cancers.

Induced *CXCL13* expression after chemotherapeutic treatment of cancer cells targeting DNA

replication, DNA-damage and BTK signaling

Chemotherapeutic treatments in MM and NSCLC induce *CXCL13* and therapy resistance [13, 27].

To determine whether chemotherapeutics that induce DNA damage response and inhibit DNA

replication affect *CXCL13* expression in cancer cells, we investigated the effects of hydroxyurea

(HU), gemcitabine, etoposide, and bleomycin on *CXCL13* expression (Figure 2A-D). In Raji and

HCT116 cells, exposure to HU increased CXCL13 expression (Figure 2A, supplementary figure

2A). In contrast, γH2AX, a marker of DNA damage, was decreased in Raji cells treated with 20

mM HU than cells treated with 10 mM HU (Figure 2A, B). Similarly, treatment of Raji cells with

gemcitabine, etoposide and bleomycin increased the CXCL13 levels however no reduction in

γH2AX was observed upon this treatment (Figure 2A-D). Taken together, these results show that

DNA damage induced by various chemotherapeutics induces CXCL13 expression (Figure 2A-D).

It is possible that higher CXCL13 levels may counteract DNA replication stress-induced DNA

damage caused by HU treatment (Figure 2B). These suggest a possibility to check in future
whether CXCR5-CXCL13 axis promotes the γH2AX turnover and DNA-repair.

DNA damage and DNA replication-stress induced signaling pathways activate p53
[43]. Therefore, we tested if p53 activation promotes *CXCL13* expression (Figure 2D). We
examined the effect of Nutlin-3a treatment, an agonist of p53, on *CXCL13* levels in Raji cells
(Figure 2D). Nutlin-3a treatment induced the p53-dependent gene *MDM2*, but at a lower level
than in A549 cells which harbors a wild type p53 (Supplementary Figure 2 A, B), suggesting that
Nutlin-3a partially activates the mutant p53 present in Raji cells (Supplementary Figure 2A, B).
Intriguingly, treatment with either Nutlin-3a or bleomycin further increased *CXCL13* levels as well
as *CXCR5* and *CCR7*, the GPCRs expressed in GC B cells (Figure 2D, E). Surface expression of
CXCR5 in Raji cells was induced upon bleomycin and Nutlin-3a treatment (Figure 2E). These
results suggests that chemotherapeutic treatment also induce CXCR5 expression in cancer cells.
These results also imply a role for mutant p53 in regulation of *CXCL13, CXCR5 and CCR7*
expression (Figure 2D-E). In addition, treatment with Ibrutinib (a BTK inhibitor) and AZ5576 (a
CDK9 inhibitor) led to an upregulation of *CXCL13* expression in OCI-LY3 cells, which are an
activated germinal center B cell-like diffuse large B-cell lymphoma (AGC-DLBCL) cell line, as
well as in VAL cell line (B-lymphoma) (Figure 2F, G) [44]. This suggests that chemotherapeutics
agents inhibiting the BTK signaling and CDK9 inhibitors induce the *CXCL13* and CXCR5
expression in hematological and solid cancers. Cancer cell survival is typically promoted by
increased NF-kB, mTOR1C, AKT and ataxia telangiectasia and Rad3-related protein (ATR)
signaling [45]. To examine if these pathways affect *CXCL13* expression under DNA-replication
stress, we measured *CXCL13* mRNA by qRT-PCR in Raji cells treated with inhibitors against NF-
kB, mTOR1C, AKT, and ATR (Figure 2H) in the HU-treated Raji cells (Figure 2H). The profound
reduction on *CXCL13* mRNA levels was seen in response to ATR inhibition (Figure 2H). Overall
these suggests that chemotherapeutics agents inhibiting the DNA-replication, DNA-damage
response, BTK signaling and CDK9 inhibitors induce the *CXCL13* and CXCR5 expression in
hematological and solid cancers while ATR signaling promotes the CXCL13 signaling.

MBD1 regulates the *CXCL13* transcription independently of CTCF
By *in silico* analysis of a eukaryotic promoter database, we found putative binding sites for CTCF,
retinoic acid receptor alpha (RARA), p53, and MYC in the 1 kilobase upstream of the *CXCL13*

transcriptional start site (TSS) (Figure 3A-D). These factors also exhibited binding on the mouse

CXCL13 promoter (Supplementary figure 3A-C), suggesting an evolutionary conserved regulation

of *CXCL13* by these factors (Figure 3A-D, Supplementary Figure 3A-C). MBD1 physically

interacts with RARA and promyelocytic leukemia protein (PML), leading to histone deacetylase-

3 mediated gene silencing [46]. Given the association among MBD1, RARA and CTCF with

DNA-methylation, we tested the effect of MBD1 and CTCF on *CXCL13* expression (Figure 3E-

F). Short hairpin (shRNA) silencing of *MBD1* and *CTCF* was confirmed in Raji cells (Figure 3F).

Interestingly, shMBD1-Raji cells induced *CXCL13* expression compared to scramble control

(shSCR-Raji) (Figure 3E), while shCTCF-Raji cells exhibited slight reduction in *CXCL13* mRNA

levels (Figure 3E). Compared to shMBD1 alone, shMBD1+shCTCF-Raji cells exhibited a higher

reduction in *CXCL13* expression (Figure 3E). These results suggest a suppressive role of MBD1

on *CXCL13*-expresion but a positive role of CTCF in *CXCL13* expression in MBD1-depleted

conditions (Figure 3E).

Since HU-treatment increases *CXCL13* expression in Raji and HCT116 cells (Figure 2A,

Supplementary Figure 2A, Figure 3E), we next examined whether MBD1 also regulates *CXCL13*

expression under HU treatment. Compared to HU-treated shSCR-Raji cells, HU-treated shMBD1-

Raji cells showed increased *CXCL13* expression (Figure 3E). The mean value of *CXCL13* mRNA

in HU-treated shMBD1-Raji cells was 12.3 than the mean value of 10.6 in non-HU treated

shMBD1 cells (Figure 3E-F). In addition, HU-treated shCTCF-Raji cells exhibited slight reduction

in *CXCL13* levels than the HU-treated shSCR-Raji cells (Figure 3E). Moreover, HU-treated

shCTCF+shMBD1-Raji cells exhibited almost equal levels of *CXCL13* expression as seen in HU-

treated shMBD1-Raji cells, but these levels were greater than HU-treated shCTCF-Raji cells

(Figure 3E). These results suggest that MBD1 plays a more important role in suppressing the

CXCL13 expression under HU-treatment (Figure 3E-F). These results also suggest that CTCF

deficiency does not affect the upregulated levels of *CXCL13* in *MBD1*-deficient cells, suggesting

a downstream role of CTCF in MBD1-dependent CXCL13-regulation (Figure 3E).

Indeed, *MBD1* overexpression in Raji cells led to a stronger reduction in *CXCL13* levels in

response to HU treatment compared to non-HU-treated Raji cells (Figure 3F-G). Again, these

analyses reveal a prominent role of MBD1 in suppression of *CXCL13* expression under the HU

stress (Figure 3E-H). In contrast, overexpression of *BCL6*, a ubiquitously expressed protein in

GC B cells [47], slightly reduced CXCL13 levels in either absence or presence of HU (Figure

3G, Supplementary Figure 3G). The expression of *BCL6* and *MBD1* plasmids was confirmed by

western blotting in HEK293T cells (Supplementary Figure 3D, E).

Given that MBD1 binds to CpG-methylated DNA and promotes gene silencing [48], we next

aimed to clarify the role of DNA methylation in *CXCL13* expression in other B-lymphomas and

colon cancer cells (Figure 3H-I). From published datasets, we explored the effects of 5-

Azacytidine (5-Aza), on *CXCL13* expression [49, 50]. 5-Aza is a chemotherapeutic drug that

inhibits DNA CpG methylation [51]. Our analysis revealed that 5-Aza treatment in HT29 cells, a

colon cancer cell line, induced *CXCL13* expression in the 5 μM and 10 μM 5-Aza-treated groups

compared to the DMSO-treated groups [49] (Figure 3H, I). We also noted that 5-Aza treatment

increased *CXCL13* and *CXCR5* expression in OCI-LY3 cells, a DLBCL cell type [50] (Figure 3I;

GSE190319). These findings suggest the DNA-methylation dependent suppression of *CXCL13*

expression**.** Finally, to test if cMYC, which is highly amplified in Burkitt's lymphoma and the

Raji cells due to translocation of the MYC locus to the Immunoglobulin locus, affects CXCL13

levels in Raji cells, we compared the CXCL13 levels in shSCR-Raji cells to shMYC#1-Raji and

shMYC#2-Raji cells. The shMYC#1-Raji and shMYC#2-Raji cells showed a decrease in

CXCL13 expression than shSCR-Raji cells (Figure 3J), indicating a positive role of c-MYC in

promoting CXCL13 expression in Burkitt's lymphoma cells. These results suggest that DNA

methylation and MBD1 play a regulatory role in *CXCL13* expression, while cMYC positively

regulates CXCL13 expression.

CTCF and RNAPII enrichment on the *CXCL13* promoter upon HU-treatment in Raji cells

Since we observed a slight positive role of CTCF in *CXCL13* expression under non-HU treated

condition (Figure 3E), we next attempted to understand the relationship between the pattern of

CTCF binding on *CXCL13* promoter and its association with *CXCL13* expression. We explored

the CTCF ChIP-seq data from 114 cell lines available on ENCODE (Figure 4A, Supplementary

Figure 4A). We classified the CTCF-binding sites observed on the *CXCL13* locus in these 114

cell lines into three categories: site-1, site-2, and site-3 (Figure 4B). CTCF binding was observed

at site-1 in 47 cell lines including the normal B and prostate cells (Figure 4A). The site-1 is

located about 23 kb upstream of *CXCL13* TSS (Figure 4A, Supplementary Figure 4A). CTCF

binding at site-2, consisting of two closer consecutive CTCF peaks, was observed in 101 out of

114 cell lines, including the normal B and prostate cells, known to be negative for CXCL13

expression (Figure 4A, Supplementary Figure 4A). The consistent pattern of CTCF binding at

site-2 in 101 cell lines suggests a conserved binding of CTCF on the *CXCL13* locus (Figure 4A,

Supplementary Figure 4A). Given that most cell types do not express the *CXCL13* naturally, we

suggest that the conserved CTCF-binding on site-2 might be associated with suppression of

CXCL13 expression (Figure 4A). In contrast, only 8 cell types exhibited CTCF binding at site-3

(Figure 4A, supplementary Figure 4A), which spans between exons 2 and exon 3 of the *CXCL13*

locus (Figure 4A, B). Interestingly, among these 8 cell types, 6 were of primary origin, while 2

cell lines, namely OCI-LY3 (DLBCL) and PC3 cell line (Prostate cancer), have been reported to

exhibit *CXCL13* expression (Figure 4A, Supplementary Figure 4A) [52]. These findings suggest

that CTCF binding at site-3 may be associated with an alternative conformation of the *CXCL13*

promoter that could facilitate *CXCL13* transcription.

To determine if CTCF redistribution occurs at the *CXCL13* promoter, we performed a CTCF-

ChIP assay in HU-treated Raji cells (Figure 4C-D). HU treatment induces *CXCL13* expression

(Figure 2A, 3E, Supplementary Figure 2A); therefore, altered CTCF binding at the *CXCL13*

promoter may be observed under HU-treated conditions. *In silico* analysis predicted dense

binding sites for CTCF between -1500 to -1800 and -597 and -726 on *CXCL13*-TSS (Figure 4I-

J); these were designated as primary pair 1 (PP1) at -1503 bp (Figure 4I-J) and primer pair 2

(PP2) at -754 (Figure 4C-D). CTCF binding was not detected at the PP1 site, but CTCF binding

was induced at the PP2 site in HU-treated Raji cells (Figure 4C-D). These suggests that HU-

treatment leads to CTCF-redistribution on *CXCL13* locus in Raji cells (Figure 4C-D).

Since it is unclear whether CTCF redistribution is associated with *CXCL13* expression, we

hypothesized that CTCF enrichment at the PP2 site might be linked to increased RNA

Polymerase II (RNAPII) occupancy at the *CXCL13* locus (Figure 4E). To test this, we performed

the RNAPII ChIP assay in HU-treated Raji cells (Figure 4E). Interestingly, we observed

increased RNAPII occupancy at the PP2 site (Figure 4E). Based on these findings, we propose

that CTCF recruitment to the *CXCL13* promoter leads to conformational changes of *CXCL13*-

promoter, allowing the RNAPII and transcriptional machinery to access the *CXCL13* promoter

and activate *CXCL13* expression under genotoxic stress.

Finally, considering the association between CTCF binding and TET2 dependent DNA-

demethylation [53], we investigated whether TET1/2 inhibition affect the CXCL13 expression.

Treatment with Bobcat-339, a potent TET1/2 inhibitor, reduced CXCL13 expression in HCT116

cells (Figure 4F). This suggests that TET1/2 enzymes promote CXCL13 expression in HCT116

cells. Although, we did not see an inhibitory effect of Bobocat-339 in Raji cells (Data not

shown), this suggests a cell type specific effect of Bobocat-339 on CXCL13 expression. This

may be possible that cell type specific genetic mutation may determine the TET1/2 activity on

the *CXCL13* promoter. It is noteworthy that HCT116 cells harbors wild type p53 and Raji

harbors two mutant p53. The p53 may interfere with TET1/2 activity on *CXCL13* promoter since

Nultin-3a treatment induced the *CXCL13* transcription in Raji cells (Figure 2D). Overall, these

results suggest a coordinated regulation among CTCF, TET enzymes, DNA methylation in

determining the *CXCL13* expression in cancer and normal cells.

Distal super-enhancer of *CCNG2* locus interacts with the *CXCL13* promoter in B-lymphoma

The human *CXCL13* is located on chromosome 4, and harbors 5 exons (Ensemble ID#

ENST00000286758; CXCL13-201; Figure 5A, Supplementary 4B). Peripheral Blood

Mononuclear Cells (PBMCs), including human B cells do not express *CXCL13* (Supplementary

Figure 5A), suggesting a closed conformation of *CXCL13* promoter in most cell types. To elucidate

the promoter conformation features of *CXCL13* expressing Raji cells, we assessed the binding of

the RNAPII subunit POL2RA along the *CXCL13* locus using ENCODE ChIP-Seq data for Raji,

as well as HepG2 and Ishikawa cells (Figure 5A). Strikingly, POL2RA ChIP-seq peaks were not

observed in near vicinity of *CXCL13* promoter in Raji, HepG2 or Ishikawa cells (Figure 5A). The

closest POL2RA peaks were found at -317,921 bp and -359,311 bp upstream of the *CXCL13* locus

(Figure 5A), while the nearest downstream peak observed in Raji, HepG2, and Ishikawa cells was

at +307,482 bp from the TSS (Figure 5A). Since we did not observe POL2RA peaks close to the

CXCL13-TSS, we hypothesized that a distal enhancer may drive *CXCL13* expression in Cancer

cells. To investigate this hypothesis, we explored the closest upstream and downstream regions

co-occupied with POL2RA, histone H3K4me3, and the super-enhancer specific histone marks, the

Histone H3K27-actyaltion (H3K27-Ac) and Histone H3K4me1 (Figure 5A). We analyzed the

ChIP-seq data available at the ENCODE (UCSD database; Figure 5A-B), and identified three

super-enhancers namely SE1, SE2 and SE3 (Figure 5A). SE1 is located between exon 2 and 3 of

the *CCNG2* locus, while SE2 is downstream of *CCNG2*, and SE3 was present near the *CNOT6L*

TSS (Figure 5A, C). The strongest POL2RA peaks were present at SE1 in Raji, HepG2 and

Ishikawa cells, and the POL2RA peak on SE2 was only found in Raji cells but not in HepG2 or

Ishikawa cells (Figure 5A). POL2RA occupancy was lower at SE2 than SE1 in Raji (Figure 5A).

Given that SE1 harbors the highest enrichment of POL2RA signals (Figure 5A), we hypothesized

that SE1 is a potential super-enhancer driving aberrant *CXCL13* expression in cancer cells. In

mantle cell lymphoma, as demonstrated by HiC interaction data [54], there was an observed

interaction between SE1 and the region flanking outside but nearer to *CXCL13* locus (Figure 5B).

This interaction point further exhibited another interaction with the *CXCL13* locus (Figure 5B).

Also, virtual circular chromosomal conformation capture (4C) from human spleen cells showed

the interaction between region closer to SE1 and the *CXCL13* locus (Figure 5C) [55]. These

suggest a possible interaction between the *CXCL13* promoter and the region near the *SE1* (Figure

5B,C). To test if SE1 interacts with *CXCL13* promoter, we performed chromosome conformation

capture assay (3C) assay using Raji cells. We checked the interaction of SE1 with predicted

CXCL13-TSS, located upstream of exon 1 [31] (Figure 5D-F). The 3C assay revealed an

interaction between the *CXCL13* promoter and SE1 (Figure 5D, E). To confirm the interaction

between SE-1 and the CXCL13 promoter, we performed the nested PCR from the purified PCR

products using the primer FP2 and RP2 (Figure 5E, F). The nested PCR allows amplification from

the first PCR containing 2069 bp product and generates a smaller band of 969 bps (Figure 5E, F).

The presence of smaller band of 969 bps further confirmed the SE1-*CXCL13-TSS* interaction

(Figure 5D-E). These results indicate that *CCNG2* super-enhancer interacts with *CXCL13*-TSS

and induces the *CXCL13* expression in cancer cells. Since the SE1 is enriched with RNAPII, this

interaction allows the access of RNAPII from SE1 to drive *CXCL13* transcription (Figure 5A,D-

E). Moreover, 95 cell lines from ENCODE exhibited DNase-hypersensitivity sites and a higher

value of GeneHancer score at SE1 (Figure 5A), suggesting the nature of open chromatin on SE1.

Multispecies alignment of SE1 and SE3 revealed its conservation among the other vertebrates,

suggesting a role for SE1 in developmentally programmed gene expression among vertebrates and

its leaky access in cancer cells driving he *CXCL13* expression (Figure 5F).

***CXCL13* expression in B-lymphomas perturbs the CXCR5-migration**

FDCs form CXCL13 gradient in the light zone (LZ) of GCs, which attracts CXCR5-positive GC

B cells, and promotes their affinity maturation and selection [1, 9, 56]. Whether aberrant CXCL13

expression in GCDBLs serves as a programmed adaptation, promoting CXCR5-internalization and

deprived LZ-migration is not known. Since peripheral blood mononuclear cell including B cells

do no express *CXCL13* (Supplementary 5A), we hypothesized that *CXCL13* expression in

GCDBLs may lead to CXCR5 receptor-internalization, reducing their migration towards external

CXCL13. To test this, we compared CXCL13-induced migration of normal peripheral B cells

(NPBCs) obtained from healthy donors with GCDBLs, Raji, FL-18, FL-218, and FL-318 (Figure

6A-C). NPBCs efficiently migrated towards recombinant human CXCL13 (rhCXCL13), while the

migration index of multiple GCDBLs was significantly lower (Figure 6A). Strikingly, Raji cells

exhibited robust migration towards recombinant human CXCL12 (rhCXCL12), a dark zone (DZ)

chemokine (Figure 6B, C). This preference of Raji cells for rhCXCL12 over rhCXCL13 suggests

GCDBLs confinement to the DZ and hinderance to LZ entry (Figure 6B, C). To investigate if

internal CXCL13 disrupts CXCL13-mediated migration of GCDBLs towards an external CXCL13

gradient, we examined CXCL13-mediated migration of Raji cells transfected with lentivirus

expressing CRISPR/Cas9 guide RNA (gRNA) targeting *CXCL13* exons 2 and 3 (referred to as

CXCL13-KO Raji cells) (Figure 6D). Interestingly, bulk containing *CXCL13*-KO Raji cells

exhibited a significant rescue in CXCL13-migration compared to Raji cells (Figure 6D). This

suggests that internal CXCL13 disrupts migration of Raji cells and GCDBLs.

A plausible explanation of reduced migration in GCDBLs is CXCR5-internalization from

interaction with internal CXCL13 ligand reduces its affinity towards external CXCL13. To test

this, we performed co-immunoprecipitation (co-IP) of CXCL13 in Raji cells (Figure 6D). Elutes

of CXCL13 co-IP exhibited distinct CXCR5 signals (Figure 6D), indicating a physical interaction

between CXCL13 and CXCR5 in Raji cells (Figure 6D). This suggests that CXCR5-internalization

takes place in Raji, thus B-lymphoma cells (Figure 6D). CXCR5-internalization was also

confirmed by flow cytometry (FC) analysis: as intracellularly stained CXCR5 receptors were seen

in 100% of Raji cells while only 29% of cells were positive for surface-CXCR5 (Figure 6F-G).

These findings indicate internalization of surface-CXCR5 in Raji and possibly other GCDBL. Off

note, to confirm that *CXCR5* coding sequence mutations were not the cause of reduced CXCL13

migration in Raji cells, we aligned Raji *CXCR5* with human *CXCR5* (Uniprot ID# P32302),

revealing three SNPs (Supplementary Fig 6B). Despite these SNPs (Proline 39: CCC, Glycine 43:

GGT, Threonine 338: ACC), however the resulting amino acids remained unchanged, confirming

that Raji's CXCR5 is identical to wild type CXCR5 (Supplementary Figure 6B). Therefore,

alterations in the CXCR5 protein are not the cause of the altered migration of Raji cells towards

CXCL13. Instead, our data suggests that the internalized CXCL13 caused the impaired migration

of B-lymphoma cells (Supplementary Figure 6B).

Accelerated cell cycle progression and suppression of p53-dependent transcripts by the

CXCR5-CXCL13 axis in hematological and solid cancers

The CXCR5-CXCL13 axis activates intracellular signaling and confers chemotherapeutic

resistance in *PTEN*-deficient cancers [32]. Given that *PTEN* is regulated by p53, this suggests a

potential interference of the CXCR5-CXCL13 axis with p53 target genes. Conversely, studies have

shown that p53 suppresses CXCR5 [57], indicating a complex interplay between p53 and the

CXCR5-CXCL13 axis. Thus, the precise impact of the CXCR5-CXCL13 axis on p53 regulation

remains poorly understood. To investigate the effects of CXCR5 overexpression on cancer cell

survival, proliferation, chemotherapeutic resistance, and p53 regulation, we stably introduced the

pCMV-CXCR5-IRES-EGFP plasmid into Raji cells *(Raji-CXCR5^{oe} hereafter,* Supplementary

figure 7A, B). Notably, *Raji-CXCR5^{oe}* cells displayed a marked increase in *CXCL13* levels

(Supplementary Figure 7C), suggesting an activation of a positive feedback loop where CXCR5

signaling enhances *CXCL13* expression through autocrine loop (Supplementary Figure 7A-C).

Despite this, *Raji-CXCR5^{oe}* cells did not exhibit enhanced CXCL13-migration compared to Raji

cells, indicating that increased CXCR5 levels did not drive CXCL13-mediated migration of

GCDBLs, possibly due to increased intracellular CXCL13 levels and receptor internalization

(Supplementary Figure 7D).

To assess the impact of the CXCR5-CXCL13 axis on cellular proliferation, we

synchronized Raji and *Raji-CXCR5^{oe}* cells using HU treatment. After releasing the synchronized

cells (Figure 7A), we monitored the live cell number (Figure 7A, B). Interestingly, *Raji-CXCR5^{oe}*

cells demonstrated significantly higher proliferation rates than Raji cells (Figure 7B). We also

observed elevated *JUN* levels in *Raji-CXCR5^{oe}* cells, a downstream target of CXCR5-CXCL13

signaling (Figure 7C), particularly in the presence of bleomycin or a combination of bleomycin

and Nutlin-3a (Fig 7C) . Compared to control Raji cells, *Raji-CXCR5^{oe}* cells showed increased

levels of *CCR7* (Figure 7C), a GPCR known to promote cancers metastasis [58]. Treatment with

bleomycin, Nutlin-3a, or the combination of both led to increased *CCR7* levels in both Raji and

Raji-CXCR5^oe cells (Figure 7C). However, the levels of *CCR7* were notably higher in *Raji-*

CXCR5^oe cells within their respective treatment groups compared to Raji cells (Figure 7C). In

contrast *Raji-CXCR5^oe* cells exhibited an overall reduction in *TP53* transcript levels compared to

Raji cells (Figure 7D). *TP53* transcripts were also low in individual or simultaneously treated

groups with bleomycin and Nutlin-3a (Figure 7D) in the *Raji-CXCR5^oe* than Raji cells , indicating

a potential interference of the CXCR5-CXCL13 axis in the p53 levels (Figure 7D).

498   The transcriptional targets of p53 play important roles in cell cycle regulation,

therefore we measured mRNA levels of *p53* target genes *CDKN1A* (encodes the p21), *DDB2*

(encodes a DNA-repair factor) and *PTP4A1*(encodes for a protein phosphatase required for mitotic

entry) between Raji and *Raji-CXCR5^oe* cells (Figure 7E). Following treatment with bleomycin,

Nutlin-3a, or their combination to induce p53 activation, we observed an induction of *CDKN1A*

and *DDB2* levels in Raji cells (Figure 7E). Interestingly, *Raji-CXCR5^oe* cells exhibited a lower

induction of *CDKN1A* in response to combined treatment with bleomycin and Nutlin-3a compared

to Raji cells treated with bleomycin and Nutlin-3a (Figure 7E). Conversely, *DDB2* levels were

notably higher in *Raji-CXCR5^oe* cells, particularly in the groups either treated with bleomycin or

co-treated with bleomycin and Nutlin-3a, compared to the corresponding treatment groups of Raji

cells (Figure 7E). Furthermore, the expression of *PTP4A1* was notably increased in *Raji-CXCR5^oe*

cells upon combined treatment with bleomycin and Nutlin-3a, as compared to its decreased

expression in Raji cells with similar treatment conditions (Figure 7E). This indicates that the

CXCR5-CXCL13 axis can impede the p53-mediated regulation of *PTP4A1, DDB2* and *CDKN1A*

(Figure 7D). Since Raji cells exhibited DNA-damage-induced expression of CXCL13 and CXCR5

(Figure 2C, D), we investigated whether this upregulation of CXCR5 and CXCL13 could enhance

PTP4A1 levels without the need for CXCR5 overexpression. Our observations revealed that

treatment of Raji cells with Bleomycin and Nutlin-3a led to nearly a twofold increase in PTP4A1

protein levels (Figure 7F). Moreover, this enhancement of PTP4A1 was further intensified,

reaching up to a fourfold increase, upon combined treatment of Raji cells with Bleomycin and

Nutlin-3a (Figure 7F). These findings confirm that the chemotherapeutic-induced CXCR5-

CXCL13 signaling can stimulate PTP4A1 expression, potentially promoting cell cycle

proliferation and contributing to chemotherapeutic resistance (Figure 7F). While co-treatment of

Raji cells with bleomycin and Nutlin-3a led to a decrease in *PTP4A1* mRNA levels, there was no

significant impact on protein levels. This implies that additional regulatory mechanisms may

regulate PTP4A1 in Raji cells following the combined treatment with bleomycin and Nutlin-3a

(Figure 7F).

Levels of p53 are regulated by ubiquitin ligases MDM2, COP1, ARF1BP1 (HUWE1)

and PIRH2,(Figure 7G) [59, 60]; in turn p53 can be stabilized by HAUSP-mediated de-

ubiquitination of p53 itself, ARF inhibition of ARFBP1 ubiquitin ligase activity, or ARF binding

and sequestration of MDM2 [61]. To assess how CXCR5 regulates p53, we first measured

expression of these ubiquitin ligases in *Raji-CXCR5^oe* cells (Figure 7G, H). Compared to Raji cells,

Raji-CXCR5^oe cells exhibited elevated levels of *PIRH2* (*RCHY1*), while no significant increase

was observed in *COP1* levels (Figure 7H). However, *COP1* levels were induced in Nutlin-3a-

treated *Raji-CXCR5^oe* cells compared to Raji cells treated with Nutlin-3a (Figure 7H). In contrast,

PIRH2 levels were significantly elevated in the DMSO-treated, bleomycin-treated or the

combination of bleomycin and Nutlin-3a treated groups of *Raji-CXCR5^oe* cells compared to

respective groups in Raji cells (Figure 7H).The levels of ARF1 were decreased in *Raji-CXCR5^oe*

cells than Raji cells (Supplementary Figure 7E). Notably, *ARFBP1* and *MDM2* levels were also

reduced in *Raji-CXCR5^oe* cells than Raji cells (Supplementary Figure 7E). The levels of HAUSP,

which stabilizes p53 by promoting its deubiquitylation, were significantly reduced in *Raji-*

CXCR5^oe cells (Figure 7H). In contrast, *Raji-CXCR5^oe* cells treated with bleomycin, Nutlin-3a or

a combination of both showed a significant reduction in the *HAUSP* levels compared to respective

treatment grouped in Raji cells (Figure 7 H). These suggests a potential role for CXCR5-CXCL13

signaling in destabilizing p53 through decreased HAUSP and increased PIRH2 levels (Figure 7H).

These findings suggest that the CXCR5-CXCL13 axis may modulate p53 levels by disrupting the

balance of its ubiquitination and deubiquitinating enzymes (Figure 7H).

Raji cells harbor mutant p53 proteins, so we further analyzed the effect of CXCR5-

CXCL13 axis on p53-target genes using the HCT116 and HepG2 cells, the COAD and LIHC cell

lines respectively (Figure 7I, J). HCT116 and HepG2 harbor wildtype p53, which allows us to

examine the effect of CXCR5-CXCL13 on p53 target genes accurately. HCT116 and HepG2 cells

exhibit *CXCL13* expression, therefore we transiently overexpressed the *CXCR5* in HCT116 and

HepG2 cells to activate the CXCR5-CXCL13 axis (*HCT116-CXCR5^oe* and *HepG2-CXCR5^oe*

hereafter). Interestingly, *HCT116-CXCR5^oe* cells exhibited significant upregulation of *PIRH2*,

MDM2, and *PTP4A1* than HCT116 cells (Figure 7I). Furthermore, *HepG2-CXCR5^oe* cells also

exhibited significant upregulation of *PIRH2* and *MDM2* than HepG2 cells (Figure 7J). The most

induced genes were *PIRH2*, *PTP4A1*, and MDM2 in Raji, HCT116 and HepG2 after CXCR5

overexpression (Figure 7E-J). Collectively, these results suggest that the CXCR5-CXCL13 axis

promotes the expression of p53 specific ubiquitin ligases, *PIRH2* and *PTP4A1* in lymphoid and

solid cancers suggesting a role of CXCR5-CXCL13 axis in destabilizing the p53 levels and

promoting cell cycle progression (Figure 7B, H). Collectively, these results suggest that CXCR5-

CXCL13 axis impairs the p53 axis, promoting the tumor proliferation.

Discussion

Regulation of *CXCL13* expression by DNA methylation, MBD1, CTCF and TET enzymes-

The observed inhibitory effects of 5-Aza treatment and MBD1 on *CXCL13* expression in colorectal

cancer, DLBCL, and Raji cells (Figure 3E-I) are consistent with previous findings highlighting the

suppressive role of the DNA methylation axis on gene expression [62-64]. Furthermore, the

influence of TET enzymes and CTCF on *CXCL13* expression (Figures 3E, 4A-E) suggests a

complex interplay involving DNA methylation, MBD1, TET enzymes, and CTCF in regulating

CXCL13 transcription. However, the stepwise mechanisms by which DNA methylation, MBD1

regulation, TET activity, and CTCF regulation affect *CXCL13* expression require further

investigation. We could clarify that under the DNA-replication stress, MBD1 works upstream of

CTCF in *CXCL13* regulation since the double knockdown of *MBD1* and *CTCF* induced the similar

level of *CXCL13* levels as seen in single knockdown of MBD1 (Figure 3E). On the other hand,

CTCF seems to also affect *CXCL13* expression under normal conditions (Figure 3E), suggesting

a partially non-overlapping role of CTCF and MBD1 on *CXCL13* transcription (Figure 3E). Future

studies should investigate whether TET enzymes affect CTCF distribution and MBD1 regulation

at the *CXCL13* locus in cancer cells, reported previously [53, 65, 66].

ATR-signaling, TET activity and *CXCL13* expression- We observed that DNA damage and

ATR signaling promote CXCL13 expression (Figure 2A, Supplementary Figure 2A, Figure 2H).

This suggests that DNA damage and ATR signaling may alleviate the suppressive effects of DNA

methylation and MBD1 on *CXCL13* transcription. This aligns with ATR's role in TET3-dependent

DNA demethylation and gene expression [67]. TET inhibition reduced CXCL13 expression in

HCT116 cells (Supplementary Figure 2A), suggesting that DNA damage and ATR signaling may

recruit TET enzymes to the *CXCL13* locus, leading to DNA hypomethylation and induction of

CXCL13 transcription. A rational for ATR-dependent *CXCL13* transcription could be in activating

the autocrine signaling in CXCR5+ cancer cells, and accelerated DNA repair. This is supported

by observation that Raji cells treated with 20mM HU, which have higher CXCL13 levels, showed

lower γH2AX signals compared to those treated with 10mM HU (Figure 2A, B). This suggests a

novel link between ATR signaling and TET-dependent CXCL13 expression in DNA repair.

CTCF, enhancer hijacking and RNAPII enrichment on the *CXCL13* promoter

Using 3C assay, we identified that a distal super-enhancer (SE1) at the *CCNG2* locus interacts

with the *CXCL13* transcription start site (TSS) in Raji cells (Figure 5B-D). Evidence from Raji,

HepG2, and Ishikawa cells shows that the *CXCL13*-TSS lacks RNAPII signals at least 300 kb

upstream and downstream (Figure 5A). This suggests that SE1 interaction may facilitate

transcriptional machinery access to *CXCL13*-TSS in cancer cells (Figure 4E, Figure 5A, C). It is

possible that the interaction between SE1 and the *CXCL13* promoter may be regulated by TET

enzymes and CTCF. Supporting this, HU treatment, which induces *CXCL13* expression (Figure

2a, 3E) in Raji cells, leads to CTCF and RNAPII enrichment at the CXCL13 promoter (Figure 4D-

E). This indicates coordinated localization of CTCF and RNAPII under HU stress on *CXCL13*

promoter. Additionally, CTCF peaks at the *CXCL13* promoter site-3 are found in CXCL13 positive

OCI-LY3 and PC3 cells but not in normal B cells or hepatocytes (Figure 4A-D), suggesting CTCF

distribution may influence the inactive state of the *CXCL13* promoter. Further studies are needed

to determine whether the interaction between SE1 and the CXCL13 promoter depends on CTCF

redistribution, potentially altering chromosomal conformation within or beyond the associated

topologically associated domain, thereby facilitating RNAPII access to the *CXCL13* locus. These

findings align with the established interplay among TET enzymes, DNA hypomethylation, CTCF

binding, and chromosomal conformation changes during development [53, 65, 66].

Conserved CTCF binding on *CXCL13* locus and transcriptional suppression-The *CXCL13*

locus exhibits a consistent and conserved pattern of CTCF binding on site-2 in the 101 out of 114

examined cell lines (Figure 4A,B). Based on ENCODE ChIP-seq data sets CTCF binding at site-

2 of the *CXCL13* locus overlaps with RAD21 and SMC3 peaks in the normal hepatocytes. The

CTCF-RAD21-SMC3 may block general transcription elongation, (Nanavaty, 2020, 32333838),

suggesting a suppressive role of site-2 in *CXCL13* expression. It may be possible that cohesin

defective cells may exhibit the *CXCL13* overexpression in few cancer types. ENCODE data

suggest the loss of RAD21 in one sample of HepG2 in site 2, suggesting that RAD21 and

cohesin loss may promote the CXCL13 expression in cancer cells.

Physiological elimination of B-lymphomas during humoral immune response- CXCL13 expression in B-lymphomas may represent a programmed adaptation that restricts their migration to the LZ by CXCR5-internalization and confines them to the DZ. This restriction could potentially limit survival feedback signaling from FDCs, leading to the GCDBLs elimination. The orchestrated *CXCL13* expression via CTCF and altered chromosome conformation (Figure 3A-D, Figure 5A-D) might hinder the LZ-migration and fate of GC B cells. Moreover, Burkitt's lymphomas, and myeloma exhibit translocations between the immunoglobulin heavy and *MYC* loci [68-70], resulting in MYC overexpression. MYC is required for LZ entry of GC B cells [71]. The amplified *MYC*, leading to CXCL13 overexpression, may simultaneously prevent the LZ entry of GCDBLs, suggesting a dual role for MYC in regulating the fate of germinal center B cells. This indicates a unique adaptation within GCDBLs, where MYC and oncogenic factors like BCL6 may paradoxically serve tumor-suppressive roles by preventing the exit of GCDBLs from GCs [72]. These regulations assist the robust immune response by facilitating the elimination of GCDBLs and promoting the generation of high-affinity antibodies. Furthermore, internalized CXCR5 may undergo proteasomal degradation in GCDBLs and needs to be tested in future studies.

Chemotherapeutic resistance - The inducibility of CXCL13 and CXCR5 by DNA damage-inducing chemotherapeutics highlights potential concerns on resistance caused by CXCR5-CXCL13 axis in cancer cells (Figure 2A-D, Supplementary Figure 2A). We show that the CXCR5-CXCL13 axis can counteract p53 levels by increasing expression of the ubiquitin ligase *PIRH2* and reducing the expressions of its de-ubiquitylation enzyme, *HAUSP* (Figure 7H-I). CXCR5-CXCL13 signaling also counteracted the p53 target genes *PTP4A1*, *DDB2* and *CCR7* (Figure 7D-I), which could potentially promote cell cycle progression, DNA repair and metastasis. It will be important to see if *PTP4A1/CCR7/DDB2* knockdown in CXCR5 and CXCL13 co-expressing cancers perturbs tumor growth in xenograft models. *PTP4A1* and *CCR7* could be novel markers exhibiting chemotherapeutic-resistance in the cancer exhibiting the CXCR5-CXCL13 signaling.

Declarations

Ethical Approval – The use of human B cells/ cell lines was approved as per Institutional ethical board.

Competing interests- The authors declare no competing interest financially.

Authors' contributions- SKG designed the original hypothesis, performed experiments, and
analyzed data. PM discussed the results and provided constructive feedback. SKG wrote the
manuscript. JHB critically read the manuscript and provided constructive feedback. All authors
read and agreed to manuscript.

Funding- This work is supported grant number 21K16142 from Japan Science for promotion
sciences (JSPS) to SKG .

Availability of data and materials- All data needed to evaluate the conclusions in the paper are
present in the main text and the supplementary materials.

Acknowledgment- The authors acknowledge TCGA and the ENCODE Consortium for generating
data used in this study. We thank Dr. Katsuya Sakai for anti-goat IgG-HRP antibody. We thank
Dr Susumu Kohno for gifting shMYC plasmids. We thank Dr. Atsuyasu Sato for the discussion.
We thank Dr. Atsushi Hirao for providing the THP1 cells and Dr. Kazuhiro Murakami for useful
discussions and providing the HCT116 and LoVo cell lines, we thank Dr. Y. Endo for HepG2
cells. We thank Dr. Takeshi Suzuki for the supporting SKG's research at Kanazawa University.

Limitations of the study: Limitations of the study: In this study, we have identified and
characterized a potential mechanism that inhibits *CXCL13* expression in adherent cells, as
CXCL13 is typically expressed in certain immune cells. Our findings suggest that the suppression
of *CXCL13* expression involves DNA methylation factors and a redistribution of CTCF on the
CXCL13 promoter, leading to a conformational change that transitions the promoter to a
transcriptionally active state. However, further investigation is required to elucidate whether DNA
methylation and TET activity coordinate these changes, and how the loss of DNA methylation and
CTCF redistribution influences this altered chromosomal conformation and RNAPII enrichment
on *CXCL13* promoter to activate transcription. Additionally, the CTCF boundaries that overlap
with RAD21 and SMC3 seem to play a regulatory role in the polyadenylation of *CXCL13* mRNA,
potentially resulting in various *CXCL13* transcript forms. Therefore, transcriptional elongation
may be perturbed by the RAD21/CTCF/SMC3 sites on intron 2, producing alternatively spliced
CXCL13 transcripts. We have not addressed the alternative splicing of *CXCL13* in this manuscript.
Our study does not conclusively distinguish the forms of the CXCL13 protein reported. This is
also to note that the CXCL13 molecular weight appears higher than the predicted 13 kDa band.
This will also help clarify whether intracellular CXCL13 expression contributes to CXCR5

receiver internalization. Attempts to generate viable CXCL13-knockout cell lines other than in

receptor internalization. Attempts to generate viable CXCL13-knockout cell lines other than in

Raji cells may be necessary to resolve this concern. A secretory form of CXCL13 may arise from

the reported variant, which could impact on CXCR5-CXCL13 signaling in autocrine and paracrine

mode, but its appearance may not be visible on WB due to secretion. These limitations warrant

further investigation in future studies.

Figure legends-

Figure 1 *CXCL13* is expressed in hematological, and solid cancers (A) *CXCL13* putative copy

number values from GISTIC vs log2 copy number values in 10712 samples. p values <0.0001 are

derived from ordinary one-way ANOVA (B) *CXCL13*: mRNA expression z-scores relative to

normal samples (log RNA-Seq V2 RSEM) data are from 6909 samples in both axes. *ns* = not

significant; p value derived from ordinary one-way ANOVA (C) *CXCR5*: mRNA expression z-

scores relative to normal samples (log2 RNA Seq V2 RSEM from 6959 diploid samples *RSEM=*

RNA-seq by expectation-maximization. X axis profile shows TCGA cancer acronyms. Dotted line

in parallel to X axis separates the tumor samples exhibiting cutoff *CXCR5 mRNA* log2 values >1

or log2<1. TCGA Cancer acronyms: rectum adenocarcinoma (READ), bladder urothelial

Carcinoma (BLCA), breast cancer (BRCA), cholangiocarcinoma (CHOL), colon adenocarcinoma

(COAD), esophageal carcinoma (ESCA), head and neck squamous cell carcinoma (HNSC),

kidney chromophobe (KICH), Kidney renal clear cell carcinoma (KIRC), kidney renal papillary

cell carcinoma (KIRP), liver hepatocellular carcinomas (LIHC), lung adenocarcinoma (LUAD),

lung squamous cell carcinoma (LUSC), prostate adenocarcinoma (PRAD), stomach

adenocarcinoma (STAD), thyroid carcinoma (THCA), uterine corpus endometrial carcinoma

(UCEC) (D) *CXCL13*: mRNA expression z-scores relative to normal samples (log RNA Seq V2

RSEM) in 6953 diploid samples. X axis shows TCGA cancer acronyms. Dotted line in parallel to

X axis separates the tumor samples exhibiting cutoff *CXCL13 mRNA* log2 values >1 or log2<1 (E)

Schematic of CXCR5 and CXCL13 overexpressing cancer cells in activating autocrine, paracrine

signaling, and metastasis to distally located CXCL13 chemokine gradient. (F) Ethidium bromide

gel-electrophoresis of *CXCL13* PCR using the cDNA samples from normal human peripheral B

cells, Raji and Granta-519 cells (G) Relative quantitative real-time PCR (qRT-PCR) analysis

comparing the mRNA levels of *CXCL13* versus *ACTB* in Raji, FL-518, Granta-519, A549, and

MCF7 cells. Additionally, quantification of *CXCL13*, *CXCR5*, and *CCR7* mRNA levels in multiple

human B cell lymphoma cell lines (Raji, FL-518, Granta-519), along with breast cancer (MCF7)

and lung cancer (A549) cell lines. *CXCL13* mRNA levels in the other cell lines are normalized to

those in A549 cells. Turkey's multiple comparison test for statistical significance. *p= 0.0486* for

Raji cells vs A549 cells and *p= 0.0430* for Raji cells vs MCF-7 cells. Data are mean ± SEM for

n=3 (H) CXCL13 Western blotting using LoVo (COAD), HCT116 (COAD), HepG2 (LIHC),

MDAMB-231 (BRCA) and THP1 (Acute monocytic leukemia) cell lysates.

Figure 2- DNA damage and Replication stress induce *CXCL13* expression in the Raji cells-
(A) Western blotting of CXCL13 and γH2AX from the cell lysates of Raji cells treated with
DMSO, hydroxyurea (10 mM and 20 mM), Gemcitabine (20μM and 40μM) and Etoposide
(50μM and 100μM) for 24 hours. Values above blot indicate densitometric qualification of
CXCL13 normalized to β-actin levels from a representative blot (B) Correlation of densitometry
quantification of CXCL13 and γH2AX signals in Raji cells treated with HU, Gemcitabine and
Etoposide, normalized with CXCL13/γH2AX levels of DMSO treated Raji cells (C) CXCL13
expression analysis by Western blotting in Raji cells treated with Bleomycin (5μg/ml) for 24 hrs.
Values above blot indicate densitometric qualification of CXCL13 normalized to β-actin levels
from a representative blot (D) qRT-PCR analysis of *CXCL13*, *CXCR5*, *TP53* and CCR7 mRNA
levels in Bleomycin (5μg/ml) treated Raji cells, Nutlin-3a (10μM) treated Raji cells and
Bleomycin (5μg/ml) + Nutlin-3a (10μM) treated Raji cells are shown. ***CXCL13***: p= 0.0409 in
Raji (DMSO vs Bleomycin), p= 0.5163 (DMSO vs Nutlin-3a), p= 0.0002 [bleomycin vs
(Nutlin-3a+Bleomycin)], p <0.0001 for [Nutlin-3a vs (Nutlin-3a+Bleomycin)]. Šídák's multiple
comparisons test. ***CXCR5***; p <0.0001 [DMSO vs (Nutlin-3a+Bleomycin)]. ***CCR7***; p <0.0001
[DMSO vs (Nutlin-3a+Bleomycin)]. Dunnett's multiple comparisons test. Data are presented as
mean ± SEM; n=3; p<0.0001 for DMSO versus Bleomycin treated groups, p<0.0001 for DMSO
versus Nutlin-3a (10μM), and p<0.0001 for DMSO vs co-treatment with Bleomycin (5μg/ml)
and Nutlin-3a (10μM). (E) Surface expression of CXCR5 in Raji cells in Raji cells treated with
Bleomycin (5μg/ml), Nutlin-3a (10μM), and the combination of Bleomycin (5μg/ml) + Nutlin-3a
(10μM). (F) Treatment with Bruton Tyrosine Kinase Inhibitor Ibrutinib to OCI-LY3 for 48 hrs
induced *CXCL13* transcripts per million reads (TPM). Unpaired t-test, *p = 0.0004* (G) Panel of
OCI-LY3 and VAL cell lines treated with AZ-5576, a selective CDK9 inhibitor, exhibiting
induced *CXCL13* TPM (GSE113035) in at least 1/2 independent samples. (H) effect of 24 hr
treatment of NF-κB inhibitor (10mM), AKT inhibitor (10μM), ATR inhibitors (10μM),
mTOR1C (10μM) treatment on *CXCL13* mRNA expression in the Raji cells. Data are mean ±
SEM of three independent experiments. Ordinary one-way ANOVA. *p*= 0.0118 for *HU vs HU +*
ATRi groups.
743
Figure 3- MBD1 suppresses the *CXCL13* expression independently of CTCF (A) *In silico*
binding of RARA, MYC, p53 and CTCF on *CXCL13* promoter predicted by eukaryotic promoter
database. Factors binding to promoter with a cutoff p value > 0.001 are shown. Binding sites of
RARA at -242, -824, -941, -961 from the *CXCL13*-TSS (B) Binding sites of CTCF at -67, -597, -
726 from TSS (C) Binding sites for p53 at -633, and -634 from TSS (D) Binding sites for MYC
at -410, -716 and -717 from TSS (E) qRT-PCR quantification of *CXCL13* mRNA in -/+ HU
treated shSCR-Raji, shMBD1-Raji, shCTCF-Raji and (shMBD1+shCTCF)-Raji cells. Data are
mean ± SEM for three independent replicates. *p* values are derived from ordinary one-way
ANOVA (*p<0.0001=*****) (F) qRT-PCR analysis of the *MBD1* and *CTCF* in shSCR-Raji,
shMBD1-Raji, shCTCF-Raji and (shMBD1+shCTCF)-Raji cells. Data are mean ± SEM for three
intendent replicates. *p* values are derived after Šídák multiple comparison test (G) CXCL13

Western blotting using Raji cell lysates transfected with *pcDNA-3.1*, *pcDNA-MBD1* and *pcDNA-*
BCL6 for 24 hours followed by 10 mM HU treatment for 24 hours. Cell lysates were prepared
48 hrs post-transfection and 24 hr. post HU treatment and then subjected to *CXCL13* western
blotting. Values above blot indicate densitometric qualification of CXCL13 normalized to beta-
actin levels from a representative blot (H) HT29-cells treated with 5-Aza induces the *CXCL13*
expression. HT-29 cells treated with dimethyl sulfoxide (vehicle alone= 0 μM 5-Aza), 5μM 5-
Aza and 10 μM 5-Aza for five days. Normalized transcripts per million (TPM) values of the
CXCL13 is shown. Two treated sample out of three samples exhibited the upregulation of
CXCL13 transcript (I) 5-Azacytidine treatment of OCI-LY3 cells induced *CXCL13* and *CXCR5*
transcript levels (GSE190319). Normalized TPM values are shown in the graph (J) WB of
CXCL13 using the cell lysates from shSCR-Raji, shMYC#1-Raji, shMYC#2-Raji cells lysates.
Data are representative of three independent repeats.
767
Figure 4- Enrichment of CTCF and RNAPII on *CXCL13* promoter in HU-treated Raji cells
(A) CTCF-ChIP-seq and distribution of CTCF binding on site-1, site-2 and site-3 on CXCL13
locus in normal human B cells, prostate cells and the OCI-LY3 and PC3 cell lines. The
additional peak of CTCF on site-3 is located at position, chr4:77606576-77606790. Peak 1
sequence Position: chr4:77578302-77578777. Position of peak 2 from: chr4:77599430-77599754
OCI-LY3 and PC3 cell lines exhibit the *CXCL13* expression (B) Distribution of CTCF's
presence/absence on site-1, site-2 and site-3 in 114 cell types. Data represent the percentage of
either presence or absence of CTCF peaks on site-1, site2 and sire-3 among 114 cell lines tested
for CTCF ChIP-seq peaks. Loss of CTCF binding on either 1 or both peaks on Site-2 is counted
as absence of CTCF binding. Data are derived from transcription factor ChIP-seq peaks from
129 cell lines on ENCODE from the UCSD genome browser (C) Schematic of CTCF putative
binding sites on *CXCL13* locus at PP1 and PP2 sites located at -1503 and -597 upstream of the
TSS respectively (D) CTCF ChIP-signals (%) on PP2 site in -/+HU treated Raji cells. Data are
mean ± SEM for n=3. p= 0.0284 (unpaired t test) (E) RNAPII ChIP-signals (%) on PP2 site in -
/+HU treated Raji cells. Data are mean ± SEM for n=3 p= 0.0058 (unpaired t test) (unpaired t
test) (F) HCT116 cells were treated with DMSO and 10μM, 20μM, 40μM and 80μM
concentrations of Bobocat-339 for 24 hours. After the treatment period, the cells were lysed, and
proteins extracts were tested to assess the expression levels of CXCL13 by western blotting
analysis.
787
Figure 5- Distal super enhancer at *CCNG2* locus interacts with *CXCL13* promoter (A)
(A) Genetic features of the *CXCL13* locus are depicted using the UCSD Genome Browser with
Three super-enhancers surrounding the *CXCL13* locus, namely SE1, SE2, and SE3 are shown in
maroon. Histone marks H3K4me3, H3K27Ac, and H3K4me1 are displayed from seven cell lines
from the UCSD database. Super-enhancers highlighted in maroon and their overlapping with
H3K27Ac and H3K4me1 signals. POL2RA ChIP-seq signals from ENCODE3 near the *CXCL13*
locus are shown for Raji, HepG2, and Ishikawa cells. Comparison maps of micro-C and Hi-C

interactions, GeneHancer regulatory elements, and DNase clusters of the *CXCL13* locus are

presented from the UCSD database (B) HiC interaction data: Mantle cell lymphoma (left) and

normal GC B cells (right). In the mantle cell lymphoma HiC map, the highlighted rectangle

indicates an interaction between the *CCGN2* region and an area outside the *CXCL13* promoter,

which also interacts with *CXCL13* promoter (C) Virtual Circular chromosomal conformation

capture (4C) from spleen exhibiting the genomic interaction of *CXCL13* locus. Assembly hg19,

resolution 40,000 base pairs (D) Schematic representation of the super-enhancer identification

and their possible interaction with the *CXCL13* transcription start site (TSS). SE1, SE2, and SE3

exhibit overlapping signals of POL2RA, H3K27Ac, and H3K4me1. Schematic of SE1, SE2 and

SE3 interaction with the super-enhancer located at *CCNG2* or *CNOT6L* (E) Schematic

representation of 3C assay with EcoR1 digestion. Primer locations are shown for the detection of

the first PCR and nested PCR products using primer pairs (FP1+RP1) and (FP2+RP2),

respectively. The band lengths of the first PCR and nested PCR are 2069 and 1100 bp,

respectively (F) Ethidium bromide staining of 3C assay PCR products. PCR performed with

1000 ng of purified genomic DNA from control Raji or 3C-processed (-/+HU) Raji cells. The

nested PCR product for positive interaction between SE1 and *CXCL13*-TSS is shown with arrow

in right (G) UCSD genome browser-based conservation of SE1, SE3 among 100 vertebrates by

PhastCons and Multiz alignment is shown.

Figure 6- CXCL13 expression is human GCDBLs perturbs migration due to CXCR5-

internalization (A) Migration index of normal Human B and multiple Human B lymphoma lines

Raji, FL-18, FL-518 and Granta-519 cell migration towards rhCXCL13 gradient. Data are mean

± SEM. P values for migration index were calculated using ordinary one-way Anova with

multiple comparison. *p*= 0.002 for normal human B vs. FL-18, *p*<0.0001 for normal Human B

vs FL-218, normal human B vs FL-318 and normal Human B vs Raji cells (B) Raji cells migrate

efficiently towards rhCXCL12 (200 ng/ml). Data are mean ± SEM for n=3. NC= Negative

control exhibiting the cell migrated to empty medium without the rhCXCL12 gradient (C) Raji

cells exhibit enhanced migration towards rhCXCL12 compared to rhCXCL13 (n=3). The

migration index of Raji cells towards rhCXCL12 was significantly higher than towards

rhCXCL13 (p = 0.0006, unpaired Student's t test) (D) Comparison of migration index between

Raji cells and *CXCL13*-KO Raji cells towards rhCXCL13. *CXCL13*-KO Raji cells exhibit

increased migration towards rhCXCL13 compared to control Raji cells (Data presented as mean

± SEM for n=3, p values calculated using unpaired t test) (E) Co-immunoprecipitation of

CXCL13 and CXCR5 in Raji cells. Immunoprecipitation was performed using 100 μg of protein

lysates from Raji cells with CXCL13 antibody (ab) or IgG antibody, followed by elution and

Western blotting for CXCL13 and CXCR5 (F) Intracellular CXCR5 in Raji cells was detected

using a CXCR5 antibody, followed by probing with a PE-conjugated secondary antibody.

Unstained and isotype controls were used for flow cytometry staining. (G) Surface CXCR5

staining in Raji cells was performed with an APC-labelled CXCR5 antibody.

Figure 7. CXCR5 Modulates Cell Cycle Progression and Overrides p53 response

(A) Schematic representation of cell synchronization by HU-treatment of Raji and *Raji-CXCR5^oe*
cells (B) Enhanced proliferation in *Raji-CXCR5^oe* cells compared to Raji cells. Cell proliferation
assessed on days 0, 2, 4, and 6 with data from four replicates, plotted as multiples of $(1*10^5)$
cells on the X-axis (C) Treatment of Raji and *Raji-CXCR5^oe* cells with Bleomycin (5μg/ml),
Nutlin-3a (10μM), or their combination for 24 hours and qRT-PCR quantification of *JUN* and
CCR7 transcript levels in *Raji-CXCR5^oe* cells compared to Raji cells. *JUN* transcript is shown as
downstream positive marker gene induced from CXCR5-CXCL13 signaling. p= 0.0002 for Raji
(Bleomycin) vs *Raji-CXCR5^oe* (Bleomycin): p= 0.0336 for Raji (Nutlin-3a) vs *Raji-CXCR5^oe*
(Nutlin-3a): p= 0.0002 in Raji (Nutlin-3a+bleomycin) vs *Raji-CXCR5^oe* (Nutlin 3a+bleomycin)
groups. Šídák's multiple comparisons test. For *CCR7* transcripts; p= 0.0022 for for Raji (DMSO)
vs *Raji-CXCR5^oe*(DMSO) and p<0.001 for Raji vs *Raji-CXCR5^oe* in all three respective
treatment groups. Šídák's multiple comparisons test. Data presented as mean ± SEM for n=3. (D)
qRT-PCR quantification of *TP53* transcript levels in *Raji-CXCR5^oe* cells compared to Raji cells.
p =0.0025 in Raji (DMSO) vs *Raji-CXCR5^oe*(DMSO) , p=0.0064 Raji (bleomycin) vs *Raji-*
CXCR5^oe(Bleomycin) , p= 0.0114 for Raji (Nutlin-3a) vs *Raji-CXCR5^oe*(Nutlin-3a) and p=
0.0052 for Raji (bleomycin+Nutlin-3a) vs *Raji-CXCR5^oe*(Bleomycin+Nutlin-3a) cells. p values
are derivatives of t test. Data presented as mean ± SEM for n=3 (E) Treatment of Raji/*Raji-*
CXCR5^oe cells with Bleomycin (5μg/ml), Nutlin-3a (10μM), or their combination and mRNA
quantification of p53 target genes, *CDKN1A*, *DDB2* and *PTP4A1*. p <0.0001 for *CDKN1A*,
DDB2, *PTP4A1* in Raji (Bleomycin+Nutlin-3a) vs *Raji-CXCR5^oe*(Bleomycin +Nutlin-3a). p
values are derived with Tukey's multiple comparison test. Data presented as mean ± SEM for
n=3 (F) Treatment of Raji cells with Bleomycin (5μg/ml), Nutlin-3a (10μM) for 24 hr and
Western blotting analysis of PTP4A1 and γH2AX using whole cell lysates. PTP4A1 levels were
increased after bleomycin and Nutlin-3a treatment of Raji cells. Values above blot indicate
densitometric qualification of PTP4A1 normalized to β-actin levels. Data are representative of
three independent replicates (G) Schematic representation of enzymes regulating the
ubiquitination/de-ubiquitylation of p53 (H) Treatment of Raji/*Raji-CXCR5^oe* cells with
Bleomycin (5μg/ml), Nutlin-3a (10μM), or their combination and qRT-PCR quantification of
COP1, *PIRH2* and *HAUSP* in *Raji-CXCR5^oe* cells compared to Raji cells. p< <0.0001 for *COP1*
in Raji (Nutlin-3a) vs *Raji-CXCR5^oe* (Nutlin-3a). p= 0. 0.0118 for *PIRH2* in Raji (DMSO) vs
Raji-CXCR5^oe(DMSO) , p<0.0001for *PIRH2* in Raji (Bleomycin) vs *Raji-CXCR5^oe*(Bleomycin).
p <0.0001 for *HAUSP* in all respective treatment groups of Raji vs *Raji-CXCR5^oe* cells. Tukey's
multiple comparisons test. Data presented as mean ± SEM for n=3 (I) HCT116 and *HCT116-*
CXCR5^oe cells utilized for qRT-PCR analysis for *DDB2*, *MDM2*, *PIRH2*, and *PTP4A1* mRNA
levels. p= 0.276680 (*DDB2*), p= 0.000845 (*PIRH2*), p= 0.001291 (PTP4A1), p= 0.000005
(MDM2) in HCT116 vs *HCT116-CXCR5^oe* cells. Unpaired t test. Data presented as mean ± SEM
for n=3. (J) HepG2 and *HepG2-CXCR5^oe* cells utilized for qRT-PCR analysis of for *DDB2*,
MDM2, *PIRH2*, and *PTP4A1* mRNA levels. p= 0.126974 (*DDB2*), p= 0.00017 (*PIRH2*), p=

874   0.026746 (PTP4A1), p= 0.000404 (MDM2) in HepG2 vs *HepG2-CXCR5^{oe}* cells. Unpaired t test.
875   Data presented as mean ± SEM for n=3.
876
Supplementary Figure 1 (A) Comparison of *CXCR5* and *CXCL13* expression in 7 CHOL
samples revealed co-expression of both *CXCR5* and *CXCL13* in 4 samples. Co-expression
indicates a possible autocrine signaling in these cancers. (B-D) Overall survival analysis from
GEPIA (Gene Expression Profiling Interactive Analysis) indicates that groups with low *CXCL13*
TPM and high *CXCL13* TPM for Uveal Melanoma (UVM), Glioblastoma (GBM), and Acute
Myeloid Leukemia (AML) demonstrate varying survival outcomes. The *CXCL13*-high TPM
groups exhibit reduced survival rates in these cancer types (E) Detection of CXCL13 in cell
lysates of Raji cells, CRISPR/Cas9 transfected Raji cells with guide RNA targeting
MYC/*CXCL13* cells. The THP1 cell line serves as a positive control, expressing CXCL13.
Antibody:Novus#AF801. Cell lysates of CRISPR/Cas9 transfected Raji cells were prepared after
transfection with CRISPR/Cas9 plasmid targeting exon 2 and 3 of *CXCL13* locus, followed by
culture for 2-days prior to sorting of transfected cells. Sorted cells were subsequently cultured
and single cell clones were grown until confluency and viability. The single-cell clones
(*CXCL13^{-/+}*) were then subjected to Western blot analysis. The SDS-PAGE gel concentration
used was 12%, and the transfer was conducted for 20 minutes. Homozygous clones *(CXCL13^{-/-})*
of Raji cells could not grow faster and maintained in passage. The Western blot was conducted
using a bulk of sorted cells that were positively transfected with CRISPR/Cas9, following their
culture until viability was maintained. CXCL13 appeared around 22 kilodaltons (kDa), which
aligns with the weight of CXCL13 in THP1 cells—a positive control cell line expressing
CXCL13 and is reduced in two independent *CXCL13^{+/-}* clones of Raji cells.
897
898
Supplementary Figure 2
(A) Western blotting of CXCL13 and γH2AX from the cell lysates of HCT116 cells treated with
DMSO, hydroxyurea (10 mM and 20 mM), Gemcitabine (20μM and 40μM) and Etoposide
(50μM and 100μM) for 24 hours (B) Western blotting analysis of MDM2, p53, γH2AX and β-
actin in Raji cells treated with Bleomycin (5μg/ml), Nutlin-3a (10μM), or the combination of
Bleomycin (5μg/ml) + Nutlin-3a (10μM). (C) *MDM2* mRNA levels in the Raji/A549 cells
treated with Nutlin-3a (10μM) for 24 hrs. Data are presented as mean ± SEM for n=3 samples.
906
Supplementary Figure 3
CTCF, RARA, Myc, and p53 binding sites are evolutionarily conserved on the *CXCL13*
promoter (A) RARA binding sites are located at -216, -336, -399, and -796 from the
Transcription Start Site (TSS) (B) Predicted CTCF binding sites are at 74, 34, -131, and -749
from the TSS (C) Predicted Myc binding site is at -622 from the *CXCL13*-TSS. (D) Western
blotting of MBD1 in *pcDNA-MBD1* (1 μg) overexpressing H3K293T cells post 24 hr of

transfection (E)Western blotting of BCL6 in *pcDNA-BCL6* (1 μg) overexpressing HEK293T
cells post 24 hours of transfection.
915
Supplementary Figure 4
(A) CTCF-ChIP-seq and distribution of CTCF binding on site-1, site-2 and site-3 on CXCL13
locus in 114 cell lines including of normal human B cells and OCI-LY3. The additional peak of
CTCF on site-3 is located at position, chr4:77606576-77606790. Peak 1 sequence Position:
chr4:77578302-77578777. Position of peak 2 from: chr4:77599430-77599754 (B) Genomic
Locus of human *CXCL13*. Ensemble Genome Browser view of the CXCL13 exon-intron
position (transcript ID# ENST00000286758.4 CXCL13-201).
923
924 **Supplementary Figure 5**
925 (A) Expression status of genes neighboring the *CXCL13* locus—SEP11, CCNI, CCNG2,
926 CNOT6L, and MRPL1—in peripheral blood mononuclear B cells (PBMCs) is displayed, as
927 assessed from PBMC data (Hao et al., 2020).
928
Supplementary Figure 6
Sequencing of CXCR5 coding region in Raji cells (A) Absolute similarity in the peptide sequences
derived from *CXCR5* DNA clones of Raji cells with human *CXCR5*. Uniport alignment of the
CXCR5 protein from Raji cells and the human CXCR5 peptide from Uniprot. Three single
nucleotide polymorphisms (SNPs) in Raji's CXCR5 (Proline 39: CCC, Glycine 43: GGT,
Threonine 338: ACC) did not resulted in amino acids confirming the reduced migration of Raji
towards CXCL13 is not caused by *CXCR5* mutations.
936
Supplementary Figure 7
CXCR5 Overexpression induces *CXCL13* mRNA expression via positive feedback in Raji cells
(A) Flow cytometry analysis of CXCR5 surface staining reveals elevated CXCR5 levels in *Raji-*
CXCR5^{oe} cells compared to normal Raji cells (B, C) Quantification of *CXCR5* and *CXCL13*
transcripts using qRT-PCR in *Raji-CXCR5^{oe}* cells compared to Raji cells. The data represent
mean ± SEM for n=3, with p-values < 0.0001 and < 0.0306, respectively (D) Migration assays
against rhCXCL13 gradient in Raji and *Raji-CXCR5^{oe}* cells. Data presented as mean ± SEM for
n=3 (E) qRT-PCR analysis of the expression levels of *ARF1*, *ARFBP1*, and *MDM2* in Raji and
Raji-CXCR5^{oe} cells after 24 hours treatment with either Bleomycin (5μg/ml), Nutlin-3a (10μM),
or Bleomycin (5μg/ml) + Nutlin-3a (10μM). p= 0.0003 for Raji (DMSO) vs *Raji-CXCR5^{oe}* , p=

0.0176 for Raji (DMSO) vs *Raji-CXCR5^{oe}* and p<0.0001 Raji (DMSO) vs *Raji-CXCR5^{oe}* .

Tukey's multiple comparison test. Data are mean ± SEM for n=3 samples.

Supplementary table 1. Primer sequences

Transcript name	Primer sequences for qPCR (5' to 3')
ACTB-FP	GATGATGATATCGCCGCGCT
ACTB-RP	GAATCCTTCTGACCCATGCC
CXCR5-FP	AGATTCTCTTCGCCAAAGTC
CXCR5-RP	CACCAGCATGGGCAGCAGGAAT
CXCL13-FP	GAAGTTCATCTCGACATCTC
CXCL13-RP	GAATTCGATCAATGAAGCGT
CCR7-FP	GTACTCCATCATTTGTTTCGTG
CCR7-RP	CAGAAGGGAAGGGTCAGGAG
CDKN1A-FP	CTGGAGACTCTCAGGGTCGAAA
CDKN1A-RP	GATTAGGGCTTCCTCTTGGAGAA
DDB2-FP	TCTACTCGCTGCCGCACAGG
DDB2-RP	TCGGGACTGAAACAAGCTGCGT
PTP4A1-FP	TAAGACAAAAGCGGCGTGGAGC
PTP4A1-RP	ACGCAGCCGCATTTTAGGACG
HAUSP-FP	CAGCCTGCTACAGACGTTAT
HAUSP-RP	CCCACCCAAATGACTTTGTTA
MDM2-FP	GAACTTGGTAGTAGTCAATCAGC
MDM2-RP	CTCTCCTTCTAGATGAGGTAG
PIRH2-FP	CCGCTTGTGTCATGATAACAA
PIRH2-RP	CCACAGTTTTCACAGTGATACTGC
ARF1-FP	CTTAAGCTGGGTGAGATCGTGAC
ARF1-RP	CTCTCTGTCATTGCTGTCCAC
ARFBP1-FP	CGGCTTTTCCTGAAGAAGGGAC
ARFBP1-RP	GTTTCCAAAGGCTTCAGAGCAGC
TP53-FP	ACTTTGCGTTCGGGCTGGGA
TP53-RP	GTCTGGCTGCCAATCCAGGGA
COP1-FP	GTCCCAAGTGTAACTATGTTG
COP1-RP	CTTGGTCAGTTCCCAACCAATC
MBD1-FP	CTGTCGAGCACAGAGAATTG
MBD1-FP	CGTTCACACTTGCAGAACAG
CTCF-FP	CCGTTACTGTGATGCTGTGT
CTCF-RP	GAAGGTCTTATCGCAGTGGC
	Primers for shRNA-Knockdown (5' to 3')
shRNA-MBD1	CCGGGAACAGAGAATGTTTAA
shRNA-MYC#1	CCTGAGACAGATCAGCAACAA
shRNA-MYC#2	CAGTTGAAACACAAACTTGAA
shRNA-CTCF	ACGTGTCCACGGCGTTCAAAT
	***CXCL13* guide RNA sequence for CRISPR/Cas9-knockout**
hCXCL13-gRNA1-FP	AGGTCTATTACACAAGCTTG
hCXCL13-gRNA2-RP	CAAGCTTGTGTAATAGACCT
hCXCL13-gRNA2-FP	GAACAAGTCAATTGTGTGTG
hCXCL13-gRNA2-RP	CACACACAATTGACTTGTTC
	Primer sequences for ChIP analysis
Primers	Primer sequence
CTCF-PP1_FP	CAAGCTCATGCGTGTGCAAG
CTCF-PP1_RP	CCTACCAGTCTTTTCCATCTC
CTCF-PP2_FP	GGAGTTGTTCTGAATAACCT
CTCF-PP2_RP	GCTTGTGTTGTGATAAGGGG
	Primers for 3C assay
CCNG2-int-PCR1-FP	GAGATCATGACTCTCACAGAC

CXCL13-int-PCR1-RP	GTTCTGATTTCAGGCATTGCC
CCNG2-nested-FP	ACGTAGATCTCTTTCGCTCCCTTGAATAAATG
CXCL13-nested-RP	GTCCTTCACCTTAAGGAGAG

952 **Supplementary table 2 for antibodies for flow cytometry, western blotting, Immunoprecipitation**
953 **and ChIP assays**

Antibody name	Purpose	Product/Manufacturer
CXCR5-APC	FACS	Biolegend (356907)
CXCR4-PE	FACS	Biolegend (306505)
CXCR3-BV421	FACS	Biolegend (353715)
APC-Isotype control	FACS	Biolegend (400121)
PE-Isotype control	FACS	Biolegend (400111)
BV-421-Isotype control	FACS	Biolegend (400341)
BCL6	WB	GeneTex#GTC101338
β-actin	WB	Biolegend#643801
Histone H3	WB	Biolegend (819411)
γH2AX	ChIP	Abcam #ab20669
CTCF	ChIP, WB	Active Motif #61932
CXCR5	WB	GeneTex#GTX100351
p53	WB	Biolegend# 645805
MDM2	WB	SANTACRUZ#sc-5304
CXCL13	WB, IP	AF801, R&D systems
MBD1	WB	39858, Active motif
RNAPII	ChIP	61082, Active motif

References

1. Cyster, J.G., et al., *Follicular stromal cells and lymphocyte homing to follicles.* Immunol Rev, 2000. 176: p. 181-93.
2. Gunn, M.D., et al., *A B-cell-homing chemokine made in lymphoid follicles activates Burkitt's lymphoma receptor-1.* Nature, 1998. 391(6669): p. 799-803.
3. Legler, D.F., et al., *B cell-attracting chemokine 1, a human CXC chemokine expressed in lymphoid tissues, selectively attracts B lymphocytes via BLR1/CXCR5.* J Exp Med, 1998. 187(4): p. 655-60.
4. El-Haibi, C.P., et al., *Differential G protein subunit expression by prostate cancer cells and their interaction with CXCR5.* Molecular Cancer, 2013. 12(1): p. 64.
5. Rao, D.A., et al., *Pathologically expanded peripheral T helper cell subset drives B cells in rheumatoid arthritis.* Nature, 2017. 542(7639): p. 110-114.
6. Kobayashi, S., et al., *A distinct human CD4+ T cell subset that secretes CXCL13 in rheumatoid synovium.* Arthritis Rheum, 2013. 65(12): p. 3063-72.
7. Yoshitomi, H., et al., *Human Sox4 facilitates the development of CXCL13-producing helper T cells in inflammatory environments.* Nature Communications, 2018. 9(1): p. 3762.
8. Manzo, A., et al., *Mature antigen-experienced T helper cells synthesize and secrete the B cell chemoattractant CXCL13 in the inflammatory environment of the rheumatoid joint.* Arthritis Rheum, 2008. 58(11): p. 3377-87.
9. Förster, R., et al., *A putative chemokine receptor, BLR1, directs B cell migration to defined lymphoid organs and specific anatomic compartments of the spleen.* Cell, 1996. 87(6): p. 1037-47.
10. Hussain, S.K., et al., *Serum levels of the chemokine CXCL13, genetic variation in CXCL13 and its receptor CXCR5, and HIV-associated non-hodgkin B-cell lymphoma risk.* Cancer Epidemiol Biomarkers Prev, 2013. 22(2): p. 295-307.
11. Bürkle, A., et al., *Overexpression of the CXCR5 chemokine receptor, and its ligand, CXCL13 in B-cell chronic lymphocytic leukemia.* Blood, 2007. 110(9): p. 3316-25.
12. Kurtova, A.V., et al., *Mantle cell lymphoma cells express high levels of CXCR4, CXCR5, and VLA-4 (CD49d): importance for interactions with the stromal microenvironment and specific targeting.* Blood, 2009. 113(19): p. 4604-13.
13. Beider, K., et al., *CXCL13 chemokine is a novel player in multiple myeloma osteolytic microenvironment, M2 macrophage polarization, and tumor progression.* Journal of Hematology & Oncology, 2022. 15(1): p. 144.
14. Kazanietz, M.G., M. Durando, and M. Cooke, *CXCL13 and Its Receptor CXCR5 in Cancer: Inflammation, Immune Response, and Beyond.* Frontiers in Endocrinology, 2019. 10(471).
15. Singh, S., et al., *Clinical and biological significance of CXCR5 expressed by prostate cancer specimens and cell lines.* Int J Cancer, 2009. 125(10): p. 2288-95.
16. Panse, J., et al., *Chemokine CXCL13 is overexpressed in the tumour tissue and in the peripheral blood of breast cancer patients.* British Journal of Cancer, 2008. 99(6): p. 930-938.
17. Ammirante, M., et al., *Tissue injury and hypoxia promote malignant progression of prostate cancer by inducing CXCL13 expression in tumor myofibroblasts.* Proc Natl Acad Sci U S A, 2014. 111(41): p. 14776-81.

18.   Zhu, Z., et al., *CXCL13-CXCR5 axis promotes the growth and invasion of colon cancer*
cells via PI3K/AKT pathway. Mol Cell Biochem, 2015. 400(1-2): p. 287-95.

19.   Bugatti, S., et al., *High expression levels of the B cell chemoattractant CXCL13 in*
rheumatoid synovium are a marker of severe disease. Rheumatology (Oxford), 2014.
53(10): p. 1886-95.

20.   Shi, K., et al., *Lymphoid chemokine B cell-attracting chemokine-1 (CXCL13) is expressed*
in germinal center of ectopic lymphoid follicles within the synovium of chronic arthritis
patients. J Immunol, 2001. 166(1): p. 650-5.

21.   Krumbholz, M., et al., *Chemokines in multiple sclerosis: CXCL12 and CXCL13 up-*
regulation is differentially linked to CNS immune cell recruitment. Brain, 2006. 129(Pt
1): p. 200-11.

22.   Lee, H.T., et al., *Serum BLC/CXCL13 concentrations and renal expression of*
CXCL13/CXCR5 in patients with systemic lupus erythematosus and lupus nephritis. J
Rheumatol, 2010. 37(1): p. 45-52.

23.   Barone, F., et al., *Association of CXCL13 and CCL21 expression with the progressive*
organization of lymphoid-like structures in Sjögren's syndrome. Arthritis Rheum, 2005.
52(6): p. 1773-84.

24.   Meraouna, A., et al., *The chemokine CXCL13 is a key molecule in autoimmune*
myasthenia gravis. Blood, 2006. 108(2): p. 432-40.

25.   Weiss, J.M., et al., *Novel CXCL13 transgenic mouse: inflammation drives pathogenic*
effect of CXCL13 in experimental myasthenia gravis. Oncotarget, 2016. 7(7): p. 7550-62.

26.   Dai, S., et al., *Intratumoral CXCL13(+)CD8(+)T cell infiltration determines poor*
clinical outcomes and immunoevasive contexture in patients with clear cell renal cell
carcinoma. J Immunother Cancer, 2021. 9(2).

27.   Wang, G.Z., et al., *The chemokine CXCL13 in lung cancers associated with*
environmental polycyclic aromatic hydrocarbons pollution. Elife, 2015. 4.

28.   Zhang, G., et al., *Mesenchymal stem cells from bone marrow regulate invasion and drug*
resistance of multiple myeloma cells by secreting chemokine CXCL13. Bosn J Basic Med
Sci, 2020. 20(2): p. 209-217.

29.   Zhang, G., et al., *CXCL-13 Regulates Resistance to 5-Fluorouracil in Colorectal Cancer.*
Cancer Res Treat, 2020. 52(2): p. 622-633.

30.   Medina, D.J., et al., *Mesenchymal stromal cells protect mantle cell lymphoma cells from*
spontaneous and drug-induced apoptosis through secretion of B-cell activating factor
and activation of the canonical and non-canonical nuclear factor κB pathways.
Haematologica, 2012. 97(8): p. 1255-63.

31.   Biswas, S., et al., *CXCL13-CXCR5 co-expression regulates epithelial to mesenchymal*
transition of breast cancer cells during lymph node metastasis. Breast Cancer Res Treat,
2014. 143(2): p. 265-76.

32.   Garg, R., et al., *Protein Kinase C Epsilon Cooperates with PTEN Loss for Prostate*
Tumorigenesis through the CXCL13-CXCR5 Pathway. Cell Rep, 2017. 19(2): p. 375-388.

33.   van de Pavert, S.A., et al., *Chemokine CXCL13 is essential for lymph node initiation and*
is induced by retinoic acid and neuronal stimulation. Nat Immunol, 2009. 10(11): p.
1193-9.

34.   Kim, S., N.-K. Yu, and B.-K. Kaang, *CTCF as a multifunctional protein in genome*
regulation and gene expression. Experimental & Molecular Medicine, 2015. 47(6): p.
e166-e166.

35. Lai, A.Y., et al., *DNA methylation prevents CTCF-mediated silencing of the oncogene BCL6 in B cell lymphomas.* Journal of Experimental Medicine, 2010. 207(9): p. 1939-1950.

36. Zhang, X., et al., *Large DNA Methylation Nadirs Anchor Chromatin Loops Maintaining Hematopoietic Stem Cell Identity.* Mol Cell, 2020. 78(3): p. 506-521.e6.

37. Tate, P.H. and A.P. Bird, *Effects of DNA methylation on DNA-binding proteins and gene expression.* Curr Opin Genet Dev, 1993. 3(2): p. 226-31.

38. Fujita, N., et al., *Mechanism of transcriptional regulation by methyl-CpG binding protein MBD1.* Mol Cell Biol, 2000. 20(14): p. 5107-18.

39. Gothwal, S.K., et al., *BRD2 promotes antibody class switch recombination by facilitating DNA repair in collaboration with NIPBL.* Nucleic Acids Res, 2024.

40. Gao, J., et al., *Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal.* Sci Signal, 2013. 6(269): p. pl1.

41. Cerami, E., et al., *The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data.* Cancer Discov, 2012. 2(5): p. 401-4.

42. de Bruijn, I., et al., *Analysis and Visualization of Longitudinal Genomic and Clinical Data from the AACR Project GENIE Biopharma Collaborative in cBioPortal.* Cancer Res, 2023. 83(23): p. 3861-3867.

43. Lakin, N.D. and S.P. Jackson, *Regulation of p53 in response to DNA damage.* Oncogene, 1999. 18(53): p. 7644-55.

44. Bertram, K., et al., *Inhibitors of Bcl-2 and Bruton's tyrosine kinase synergize to abrogate diffuse large B-cell lymphoma growth in vitro and in orthotopic xenotransplantation models.* Leukemia, 2022. 36(4): p. 1035-1047.

45. Dan, H.C., et al., *Akt-dependent regulation of NF-{kappa}B is controlled by mTOR and Raptor in association with IKK.* Genes Dev, 2008. 22(11): p. 1490-500.

46. Villa, R., et al., *The methyl-CpG binding protein MBD1 is required for PML-RARalpha function.* Proc Natl Acad Sci U S A, 2006. 103(5): p. 1400-5.

47. Basso, K. and R. Dalla-Favera, *BCL6: master regulator of the germinal center reaction and key oncogene in B cell lymphomagenesis.* Adv Immunol, 2010. 105: p. 193-210.

48. Fujita, N., et al., *Methyl-CpG binding domain 1 (MBD1) interacts with the Suv39h1-HP1 heterochromatic complex for DNA methylation-based transcriptional repression.* J Biol Chem, 2003. 278(26): p. 24132-8.

49. Xu, X., et al., *Parallel comparison of Illumina RNA-Seq and Affymetrix microarray platforms on transcriptomic profiles generated from 5-aza-deoxy-cytidine treated HT-29 colon cancer cells and simulated datasets.* BMC Bioinformatics, 2013. 14 Suppl 9(Suppl 9): p. S1.

50. Liu, J., et al., *Epigenetic priming improves salvage chemotherapy in diffuse large B-cell lymphoma via endogenous retrovirus-induced cGAS-STING activation.* Clinical Epigenetics, 2023. 15(1): p. 75.

51. Bender, C.M., M.M. Pao, and P.A. Jones, *Inhibition of DNA methylation by 5-aza-2'-deoxycytidine suppresses the growth of human tumor cell lines.* Cancer Res, 1998. 58(1): p. 95-101.

52. Fan, L., et al., *CXCL13 is androgen-responsive and involved in androgen induced prostate cancer cell migration and invasion.* Oncotarget, 2017. 8(32): p. 53244-53261.

53. Nanan, K.K., et al., *TET-Catalyzed 5-Carboxylcytosine Promotes CTCF Binding to Suboptimal Sequences Genome-wide.* iScience, 2019. 19: p. 326-339.

54.   Vilarrasa-Blasi, R., et al., *Dynamics of genome architecture and chromatin function*
during human B cell differentiation and neoplastic transformation. Nature
Communications, 2021. 12(1): p. 651.
55.   Schmitt, A.D., et al., *A Compendium of Chromatin Contact Maps Reveals Spatially*
Active Regions in the Human Genome. Cell Rep, 2016. 17(8): p. 2042-2059.
56.   Liu, Y.J., et al., *Mechanism of antigen-driven selection in germinal centres.* Nature,
1989. 342(6252): p. 929-31.
57.   Mitkin, N.A., et al., *p53-dependent expression of CXCR5 chemokine receptor in MCF-7*
breast cancer cells. Scientific Reports, 2015. 5(1): p. 9330.
58.   Legler, D.F., E. Uetz-von Allmen, and M.A. Hauser, *CCR7: roles in cancer cell*
dissemination, migration and metastasis formation. Int J Biochem Cell Biol, 2014. 54: p.
78-82.
59.   Leng, R.P., et al., *Pirh2, a p53-induced ubiquitin-protein ligase, promotes p53*
degradation. Cell, 2003. 112(6): p. 779-91.
60.   Kubbutat, M.H., S.N. Jones, and K.H. Vousden, *Regulation of p53 stability by Mdm2.*
Nature, 1997. 387(6630): p. 299-303.
61.   Chen, D., et al., *ARF-BP1/Mule is a critical mediator of the ARF tumor suppressor.* Cell,
2005. 121(7): p. 1071-83.
62.   Rasmussen, K.D. and K. Helin, *Role of TET enzymes in DNA methylation, development,*
and cancer. Genes Dev, 2016. 30(7): p. 733-50.
63.   Larsen, F., et al., *CpG islands as gene markers in the human genome.* Genomics, 1992.
13(4): p. 1095-107.
64.   Singh, D.K., et al., *5-Azacytidine- and retinoic-acid-induced reprogramming of DCCs*
into dormancy suppresses metastasis via restored TGF-β-SMAD4 signaling. Cell Rep,
2023. 42(6): p. 112560.
65.   Tian, H., et al., *Tet-mediated DNA methylation dynamics affect chromosome*
organization. Nucleic Acids Research, 2024.
66.   Nanavaty, V., et al., *DNA Methylation Regulates Alternative Polyadenylation via CTCF*
and the Cohesin Complex. Mol Cell, 2020. 78(4): p. 752-764.e6.
67.   Jiang, D., et al., *TET3-mediated DNA oxidation promotes ATR-dependent DNA damage*
response. EMBO reports, 2017. 18(5): p. 781-796-796.
68.   Chiecchio, L., et al., *Frequent upregulation of MYC in plasma cell leukemia.* Genes
Chromosomes Cancer, 2009. 48(7): p. 624-36.
69.   Avet-Loiseau, H., et al., *Rearrangements of the c-myc oncogene are present in 15% of*
primary human multiple myeloma tumors. Blood, 2001. 98(10): p. 3082-6.
70.   Zeidler, R., et al., *Breakpoints of Burkitt's lymphoma t(8;22) translocations map within a*
distance of 300 kb downstream of MYC. Genes Chromosomes Cancer, 1994. 9(4): p. 282-
7.
71.   Dominguez-Sola, D., et al., *The proto-oncogene MYC is required for selection in the*
germinal center and cyclic reentry. Nat Immunol, 2012. 13(11): p. 1083-91.
72.   Huang, C., et al., *Cooperative transcriptional repression by BCL6 and BACH2 in*
germinal center B-cell differentiation. Blood, 2014. 123(7): p. 1012-20.

Paper outline

CXCL13 expression in hematological, solid cancers	CXCL13 regulation by DNA-methylation, MBD1, SE-CXCL13-TSS interaction	Effect of CXCR5-CXCL13 axis on p53 target genes

Highlights:
- **Aberrant *CXCL13* expression in hematological and solid cancers**
- **Chemotherapeutic treatment of cancer cells promotes *CXCL13* and *CXCR5* expression**
- **Distal super-enhancer on *CCNG2* interacts with *CXCL13* promoter**
- **CXCL13 expression in B-lymphomas prevents CXCR5-dependent migration**
- **CXCR5-CXCL13 axis encounters p53 function in hematological and solid cancer cells**

Figure- 1 Aberrant CXCL13 expression hematological and solid cancers

A. **CXCL13: copy number values in *CXCL13* altered groups**

B. *CXCL13* mRNA expression in different cancer types

C. *CXCR5* mRNA normalized to illumina Hiseq_RNAseqV2 in diploid cancers

D. *CXCL13* mRNA normalized to illumina Hiseq_RNAseqV2 in diploid cancer only

E. Schematic of CXCR5 and CXCL13 overexpressing cancer cells activating autocrine/paracrine signaling and metastasis

F. *CXCL13* mRNA levels in B-lymphomas

G. Relative *CXCL13* transcript level in cancer cells

H. CXCL13 expresion in solid cancers clls

Figure 2- DNA damage and Replication stress induce CXCL13 expression in cancer cells

A. CXCL13 expression is induced by DNA-replication stress and DNA damage

B. Corelation between CXCL13 and γH2AX levels

C. Bleomycin and Nutlin-3a treatment increases CXCL13 levels

D. Bleomycin and Nutlin-3a treatment induce *CXCL13* and *CXCR5* transcription

E. Bleomycin and Nutlin-3a treatment increase surface CXCR5 expression

F. BTK inhibitor Ibrutinib treatment induces the CXCL13 expression

G. CDK9 inhibitor induces the *CXCL13* expression

H. ATR inhibition prominently reduces the CXCL13 expression

Figure 3- MBD1 suppresses CXCL13 expresion in B-lymphoma cells

A. RARA binding sites on the human *CXCL13* promoter

B. CTCF binding sites on the human *CXCL13* promoter

C. p53 binding sites on the human *CXCL13* promoter

D. MYC binding sites on the human *CXCL13* promoter

E. MBD1 suppreses *CXCL13 expresion*

F. Knockdown efficiency of MBD1 and CTCF in Raji cells

G. MBD1 suppresses CXCL13 in HU depndent manner

H. HT29-cells treated with 5-Aza induces the *CXCL13* expression (GSE41586)

I. 5-Aza treatment of OCI-LY3 cells induce *CXCL13/CXCR5* expression

J. MYC promotes CXCL13 expresion in the Raji cells

Figure 4- Altered CTCF binding is associated with RNA-PolII enrichment at CXCL13 promoter

A. CTCF binding on *CXCL13* locus IN 114 cell lines

B. Quantification of CTCF binding on Site-1, Site-2, Site-3 of *CXCL13* locus in 114 cell types

C. CTCF binds to *CXCL13* promoter upon HU-treatment

D. Increased RNA-PolII binding at PP2 site upon HU-treatment

E. Treatment of Bobocat339 (TET inhibitor) reduced the *CXCL13* expresion in HCT116

Figure 5 Promoter conformational changes drive *CXCL13* expression in GCDBLs

A. *CCNG2* enhancer interaction with *CXCL13* locus

B. HiC interaction in Mantle cell lymphoma and normal B cells surrounding *CXCL13*

C. Virtual 4C data constructed from Hi-C data of Spleen

D. Schematic of superenhancer identification and 3C asay for the *CXCL13*-TSS1

E. Scheme of 3C asay based first and nested PCR

F. Distal super-enhancer of *CCNG2* locus interacts with *CXCL13* TSS1

G. SE1 and SE3 are evolutionary conserved among species

Figure- 6 Induced CXCL13 expression in GCDBLs deprives Light Zone migration

A. Migration of multiple B-lymphoma towards external CXCL13

B higher migration of Raij towards CXCL12 (DZ chemokine)

C. Higher migration index of Raji cells towards CXCL12 than CXCL13

D. CXCL13-Raji cells exhibits enhanced migration

E. CXCL13 IP in Raji

F. Intracellular CXCR5 in Raji cells

G. CXCR5 receptor is drastically internalized in GCDBLs

Figure 7. CXCR5-CXCL13 axis promotes cell cycle progression and counteracts the p53 stablity

A. Schematic of cell synchronization and proliferation assay

B. Higher cellular proliferation in *Raji-CXCR5^oe* cells

C. CXCR5-CXCL13 axis promotes *CCR7* expression

D. CXCR5-CXCL13 axis reduced *TP53*

E. CXCR5-CXCL13 axis counteracts p53 target genes

F. Upregulation of PTP4A1 upon DNA damage in Raji cells

G. Schematic of p53 degradation by ubiquitin ligases

H. Upregulation of *PIRH2*, *COP1* and downregulation of *HAUSP* mRNA in in *Raji-CXCR5^oe*

I. CXCR5-CXCL13 axis affects p53 target genes in HCT116

J. CXCR5-CXCL13 axis promotes *PIRH2* in the HepG2 cells

Supplementary Figure 1

A.

B. UVM survival curver

C. GBM survival curve

D. AML survival curve

**E. Full blots to detect CXCL13
(12% PAGE, 20 min transfer)**

Supplemetary Figure 2 Nutlin-3a treatment can activate mutant p53 in Raji cells

A. Bobocat-339 treatment reduces the CXCL13 expression in HCT116 cells

B. Nutlin-3a treatment of Raji cells

C. Nutlin-3a treatment can activate the mt-p53

Supplementary figure 3 - CTCF, RARA, and Myc binding sites are evolutionalry conserved on mouse CXCL13 pro

A. RARA-binding sites on the mouse *Cxcl13* locus

B. CTCF binding sites on the mouse *Cxcl13* locus

C. Myc binding sites on the mouse *Cxcl13* locus

D.

E.

Supplementary Figure 4

A. Genetic location of human *CXCL13* locus and exon-intron distribution

B. CTCF binding on *CXCL13* locus IN 114 cell lines

Site-1   Site-2   Site-3

Supplementary Figure 5- PBMC do not express the *CXCL13*

A. Human PBMC does not express *CXCL13*

Supplementary figure 6

A. CXCR5 receptor is not mutated in the Raji cells.

Query	1	MNYPLTLEMDLENLEDLFWELDRLDNYNDTSLVENHLCPATEGPLMASFKAVFVPVAYSL	60
		MNYPLTLEMDLENLEDLFWELDRLDNYNDTSLVENHLCPATEGPLMASFKAVFVPVAYSL	
P32302 CXCR5_HUMAN	1	MNYPLTLEMDLENLEDLFWELDRLDNYNDTSLVENHLCPATEGPLMASFKAVFVPVAYSL	60
Query	61	IFLLGVIGNVLVLVILERHRQTRSSTETFLFHLAVADLLLVFILPFAVAEGSVGWVLGTF	120
		IFLLGVIGNVLVLVILERHRQTRSSTETFLFHLAVADLLLVFILPFAVAEGSVGWVLGTF	
P32302 CXCR5_HUMAN	61	IFLLGVIGNVLVLVILERHRQTRSSTETFLFHLAVADLLLVFILPFAVAEGSVGWVLGTF	120
Query	121	LCKTVIALHKVNFYCSSLLLACIAVDRYLAIVHAVHAYRHRRLLSIHITCGTIWLVGFLL	180
		LCKTVIALHKVNFYCSSLLLACIAVDRYLAIVHAVHAYRHRRLLSIHITCGTIWLVGFLL	
P32302 CXCR5_HUMAN	121	LCKTVIALHKVNFYCSSLLLACIAVDRYLAIVHAVHAYRHRRLLSIHITCGTIWLVGFLL	180
Query	181	ALPEILFAKVSQGHHNNSLPRCTFSQENQAETHAWFTSRFLYHVAGFLLPMLVMGWCYVG	240
		ALPEILFAKVSQGHHNNSLPRCTFSQENQAETHAWFTSRFLYHVAGFLLPMLVMGWCYVG	
P32302 CXCR5_HUMAN	181	ALPEILFAKVSQGHHNNSLPRCTFSQENQAETHAWFTSRFLYHVAGFLLPMLVMGWCYVG	240
Query	241	VVHRLRQAQRRPQRQKAVRVAILVTSIFFLCWSPYHIVIFLDTLARLKAVDNTCKLNGSL	300
		VVHRLRQAQRRPQRQKAVRVAILVTSIFFLCWSPYHIVIFLDTLARLKAVDNTCKLNGSL	
P32302 CXCR5_HUMAN	241	VVHRLRQAQRRPQRQKAVRVAILVTSIFFLCWSPYHIVIFLDTLARLKAVDNTCKLNGSL	300
Query	301	PVAITMCEFLGLAHCCLNPMLYTFAGVKFRSDLSRLLTKLGCTGPASLCQLFPSWRRSSL	360
		PVAITMCEFLGLAHCCLNPMLYTFAGVKFRSDLSRLLTKLGCTGPASLCQLFPSWRRSSL	
P32302 CXCR5_HUMAN	301	PVAITMCEFLGLAHCCLNPMLYTFAGVKFRSDLSRLLTKLGCTGPASLCQLFPSWRRSSL	360
Query	361	SESENATSLTTF	372
		SESENATSLTTF	
P32302 CXCR5_HUMAN	361	SESENATSLTTF	372

Supplementary figure-7 CXCR5 overexpression induces the CXCL13 expression via a positive feedback

A. Higher CXCR5 levels in *Raji-CXCR5^oe* cells

B. Higher *CXCR5* mRNA levels in *Raji-CXCR5^oe* cells

C. Higher *CXCL13* mRNA levels in *Raji-CXCR5^oe* cells

D. CXCR5 overexpression does not induce the CXCL13 migration in Raji cells

E. *ARF1*, *ARFBP1*, and *MDM2* mRNA levels in *Raji-CXCR5^oe* cells