Type of Article: Genomic Resources

Title: Developing simple DNA extraction and PCR-RFLP for *MALE STERILITY 4* (*MS4*)

gene in *Cryptomeria japonica* D. Don: Toward an environmentally friendly protocol

Author names: Saneyoshi Ueno[1*], Yukiko Ito[2] , Yoichi Hasegawa[1] , Yoshinari Moriguchi[3]

Author affiliations:

[1]Department of Forest Molecular Genetics and Biotechnology, Forestry and Forest Products Research

Institute, 1 Matsunosato, Tsukuba, 305-8687, JAPAN

[2]Niigata Prefectural Forest Research Institute, 2249-5 Unotoro, Murakami, 958-0264, JAPAN

[3]Department of Agriculture, Faculty of Agriculture, Niigata University, 8050 Ikarashi 2 no-cho, Nishi-ku,

Niigata, 950-2181, JAPAN

* Corresponding author (email address): Saneyoshi Ueno (saueno@ffpri.affrc.go.jp)

ORCID:

S.U.: 0000-0001-5571-0622, Y.H.: 0000-0003-4729-2090, Y.M.: 0000-0002-8535-1517

Abstract

This study presents the development of a simple DNA extraction method and a novel

PCR-RFLP protocol for genotyping the *MALE STERILITY 4* (*MS4*) gene in *Cryptomeria japonica* as a

proof of concept. Traditional CTAB-based DNA extraction methods, while effective, involve hazardous

chemicals and require high-speed centrifugation, which are impractical in many field settings. Our

approach utilizes a household dish detergent-based buffer, sodium chloride, and polyvinylpyrrolidone

K-30 to extract DNA from *C. japonica* needle leaf tissues—without the need for liquid nitrogen. The

simplicity of this method makes it more accessible and environmentally friendly. The extracted DNA was

successfully used in PCR-RFLP analysis, targeting a single nucleotide polymorphism in the *MS4* gene,

demonstrating its efficacy for genotyping. The PCR-RFLP markers reliably discriminated between

individual genotypes, confirming the practical application of our simple extraction method, even for

conifers containing inhibitory substances. This technique is particularly advantageous for use in

arboretums and field stations, where the use of hazardous chemicals, specialized equipment, and a draft

chamber is limited. Our study contributes to genetic resource management by providing an easy, reliable,

and safer method for DNA extraction and genetic analysis.

Key words: *Cryptomeria japonica*, sugi, DNA extraction, dish detergent, PCR-RFLP, male sterility

Statements and Declarations

Competing interests: The authors declare no competing interests.

Introduction

Nucleic acid extraction is a starting process for genetic analysis and has been a topic of

research for many years (Yadav et al. 2018; Dairawan and Shetty 2020; Schenk et al. 2023). The DNA

extraction from plant tissue is difficult compared to that from animal tissue, because it contains

poly-saccharide, polyphenolics and other secondary metabolites that prevent nucleic acid extraction. This

challenge is particularly pronounced in trees, especially conifers like *Cryptomeria japonica*, which are

known for their high levels of polyphenolics and polysaccharides. These compounds interfere with DNA

extraction and PCR amplification, making the process more complex than with other plant species.

Developing a method that can effectively extract DNA from such challenging plant material is crucial for

advancing genetic studies in these species. In 1980, Murray and Thompson (Murray and Thompson

1980) published a DNA extraction method using CTAB (cetyl-trimethylammonium bromide) which

efficiently removes polysaccharides from plant tissue. Currently the CTAB-based method is used to

extract DNA/RNA from many plant species with wide downstream uses for restriction digestion and

PCR experiments without further purification (Doyle and Doyle 1987; Le Provost et al. 2007). One

drawback of the method is the use of hazardous chemicals such as chloroform and 2-mercaptoethanol,

which are recognized internationally as hazardous substances. Specific handling must comply with the

regulations and guidelines of each country. In addition to the use of hazardous chemicals, the CTAB

protocol uses centrifugal forces of up to 20,000 g. Unfortunately, arboretums and field stations are rarely

equipped with such centrifuges.

One of the simplest methods to extract DNA is the use of chelating resin, InstaGene (Bio-Rad

Laboratories, Inc.), which requires only heating the tissue with the resin matrix for DNA extraction.

Tsuruta et al. (2021) used InstaGene Matrix to extract DNA from somatic embryogenic cells of *C.*

japonica for marker assisted selection of *MS1* (*MALE STERILITY 1*) gene (Tsuruta et al. 2021). The

method was further applied for leaf, pollen, and inner bark tissue of *C. japonica* from which DNA was

extracted for *MS1* genotyping (Ueno et al. 2022). Although the InstaGene extraction is simple, the

amount of tissue (5–10 mg leaf) used with InstaGene Matrix is a key factor for the success of DNA

extraction (Ueno et al. 2022). Too much tissue results in a viscous solution, leading to failure in PCR.

Moreover, the process of trial and error with InstaGene (e.g., optimizing the amount of InstaGene matrix

used, the duration of heating the samples with the InstaGene matrix, and the PCR thermal profile) can be

a time-consuming task, especially for those inexperienced in DNA experiments.

In this study, we aim to develop a simple DNA extraction method and PCR-RFLP (polymerase

chain reaction-restriction fragment length polymorphism) genotyping technique for the recently identified

MS4 (*MALE STERILITY 4*) gene in *C. japonica* (Kakui et al. 2023). The mutation in *MS4* involves a

single nucleotide polymorphism (SNP) that results in an amino acid substitution. Although a KASP

(Kompetitive Allele Specific PCR) (Dipta et al. 2024) marker for the *MS4* gene was recently developed

(Watanabe et al. 2024), its application requires specialized equipment such as a real-time PCR machine,

which may not be readily available in arboretums or field stations. Another potential approach is

developing allele-specific PCR markers, but this method requires extensive optimization of PCR

conditions, even though Locked Nucleic Acids (LNA) are included in PCR primers (Latorra et al. 2003),

making it more challenging and time-consuming. Therefore, we focused on developing a PCR-RFLP

technique that serves as a practical example of applying our simple DNA extraction method. This

approach, which avoids the use of hazardous chemicals, aims to provide an accessible, environmentally

friendly tool for genetic analysis, offering a versatile solution for researchers in various settings.

Materials and Methods

Needle leaves from five *C. japonica* individuals (Table 1) were sampled at Niigata Prefectural

Forest Research Institute on July 22, 2022, and kept refrigerated at 4 °C until DNA extraction. One

needle leaf (approximately 5 mg) was ground in 1 mL of dish detergent buffer (10 % dish detergent

'Yashinomi Hi-power' (Saraya Co., Ltd.), 10 % NaCl, and 5% polyvinylpyrrolidone K-30 (PVP) (NACALAI

TESQUE) at room temperature using mortar and pestle (Fig. 1). The dish detergent contained 32 % surfactant

(sodium alkyl ether sulfates, alkyl betaines, and fatty acid alkanolamides), auxiliary cleaning agents, and

stabilizers. The resulting 'green juice' (approximately 500 µL) was transferred to a 1.5 mL tube, centrifuged for

three minutes with a CHIBITAN-R microcentrifuge (Hitachi Koki, Co. Ltd.) at the maximum speed of 6,200

rpm, and 300 µL of supernatant was then collected in a new 1.5 mL tube. Twice the volume (600 µL) of cold

100 % ethanol was added to the tube and mixed with inverted shaking. The solution was centrifuged for 10 min

at room temperature using the CHIBITAN-R, and the supernatant was discarded by decantation. 600 µL of

70 % ethanol was then added to the tube to wash the precipitate, spun briefly by CHIBITAN-R, and discarded

by decantation. The precipitate was air-dried at room temperature, and dissolved in 100 µL of TE buffer

(NACALAI TESQUE). The DNA concentration was measured by Qubit Fluorometer (Thermo Fisher

Scientific).

In order to demonstrate the practical application for the extracted DNA, a PCR-RFLP protocol was

developed to genotype the male-sterile causative mutation of *MS4* in *C. japonica* (Fig. 2) as a model system.

PCR primer pairs to amplify the target region were designed by Primer3Plus (Untergasser et al. 2012).   The

SEQUENCE_TEMPLATE contained the CJt034360 sequence (DDBJ accession number: ICQT01013259)

with the *PRIMER_MIN_SIZE* 23, *PRIMER_OPT_SIZE* 23, *PRIMER_MAX_POLY_X* 2,

PRIMER_PAIR_MAX_DIFF_TM 2, and SEQUENCE_TARGET 301,1 (target causative SNP coordinate

(301) along the *SEQUENCE_TEMPLATE*, and the length of 1). The enzymatic function of *MS4* is TKPR1

(tetraketide alpha-pyrone reductase 1), the dysfunction of which causes male sterility in not only *C. japonica*,

but also *Arabidopsis* (Grienenberger et al. 2010) and rice (Xu et al. 2019). Restriction enzymes to discriminate

the mutant and wild type alleles in *MS4* were searched using the REHUNT program (Cheng et al. 2018). Four

samples (Table 1), whose *MS4* genotypes were established elsewhere by artificial crossing (Miyajima et al.

2010; Hasegawa et al. 2018; Hirayama et al. 2021) and whose DNA was extracted using a modified CTAB

method (Hasegawa et al. 2020), were used in the development of PCR-RFLP protocol. The PCRs were carried

out in a 10 µL reaction containing 1 µL (about 10 ng) of template DNA, 1 µL of each of primer (2 µM stock

solution; forward: AGCCATGAATGGAACAACTGTGT, and reverse:

GGAGGAAGGCTCGGTCCTATTAT), resulting in a final concentration of 0.2 µM for each primer, and 5 µL

of QIAGEN Multiplex PCR Master Mix (Qiagen). The thermal profiles used were as follows: the initial

denaturation at 95 °C for 15 minutes, followed by 40 cycles of 95 °C for 30 sec, 60 °C for 90 sec, and 72 °C for

60 sec with the final extension of 72 °C for 10 min. The resulting products (5 µL) were digested with the

restriction enzyme *Hae*III (NIPPON GENE) in a 10 µL of reaction containing 2–5 µL of PCR products, 1 µL

of 10× Buffer M (resulting in a final concentration of 1×) , and 0.1 µL of *Hae*III (high concentration: 30 – 100

units/µL) and incubated for at least three hours or overnight at 37 °C. The PCR products and

restriction-digested products were run on 2 % agarose gels, stained with ethidium bromide, and visualized on a

UV transilluminator.

Reproducibility of this simple DNA extraction and PCR-RFLP protocol was confirmed with

current year needles sampled from a single individual, `Y938` (Table 1) growing at Forestry and Forest

Products Research Institute, Tsukuba, Japan on October 11, 2024.

Results and Discussion

The DNA extracted by the simple method yielded concentrations ranging from 1.01 to 2.11

ng/µL with an average of 1.66 ng/µL (N=8). The number of needles required to reach a sample weight of

5 mg varied, ranging from one to three needle leaves, depending on the individual sample. The

PCR-RFLP protocol developed for *MS4* successfully discriminated the individual genotypes (Fig. 3).

When DNA extracted by the dish detergent buffer was used for this protocol, expected results were also

obtained, demonstrating the successful application of the simple DNA extraction method in this study.

We confirmed the reproducibility of this simple DNA extraction and PCR-RFLP protocol by conducting

eight independent DNA extraction and PCR-RFLP experiment using the same single individual (Fig. 3).

In addition, one of the co-author (Ms. Ito), a beginner of DNA work, has already genotyped over 100

samples with a failure rate of less than 5%. This suggests that the method is not only reproducible but

also robust across different operators. If the sample is sufficiently ground, a white precipitate is visible at

the bottom of the tube after centrifugation during ethanol precipitation, indicating successful DNA

extraction. Even if the precipitate is not visible, the target fragment (1.5 kb) was still typically amplified

by PCR. These results indicate that the method is reproducible and robust across a wide variety of

samples and operators. Furthermore, using a homogenizer can improve stability and increase DNA yield.

Following the same protocol, the Multi-beads Shocker (MB3000; Yasui Kikai Corp., Osaka, Japan) at

2500 rpm for 20 sec achieved a concentration of 5 ng/µL, with a total yield of 500 ng. This demonstrates

the method's potential for high-throughput applications, making it suitable for handling large numbers of

samples efficiently without liquid nitrogen.

Because DNA extraction is a basic starting experiment of molecular biology, biology classes in

secondary education conduct DNA extraction experiments using commercially available detergent with

simple procedures (Ogata et al. 2010; de Andrade et al. 2024). Although these classes commonly use soft

tissue plant materials including broccoli flower buds and strawberries, no reports were found for DNA

extraction from coniferous plants using dishwashing detergent. In the current study, we developed a

simple DNA extraction method with commercially available dishwashing detergent for use in marker

assisted management of arboretum or field stations, where the use of hazardous chemicals is not

preferable and powerful centrifuges are not available. When DNA extracted with this simple detergent

buffer was run on agarose gels, DNA bands were rarely observed. This is probably due to the small

amount of DNA extracted that is often fragmented. However, PCR successfully amplified the 1.5 kb

fragment, though the amplification of primer dimers or non-specific amplification occurred (Fig. 3).

While polymerases, such as MightyAmp DNA polymerase (Lu et al. 2012), may be suitable for use with

crude DNA that contains PCR inhibitors, this study used HotStartTaq DNA polymerase in Multiplex Kit

(Qiagen) which is widely used for genetic analysis with microsatellite markers for clonal identification

and parentage analysis. Additionally, since the DNA extracted using this method could also be utilized

with restriction enzymes, which are generally more sensitive to inhibitors than polymerases in PCR, it

suggests that the quality of the extracted DNA is high enough for various molecular biology applications.

For next-generation sequencing applications, this DNA extraction method shows potential compatibility

with amplicon sequencing, such as MIG-seq (Suyama and Matsuki 2015). Although the yield and quality

may be limiting factors for RADseq (Baird et al. 2008; Peterson et al. 2012), exploring its application to

RADseq could be of interest and presents a challenge for future research.

The DNA extraction buffer used in this study (Fig. 1) comprised commercially available

dishwashing detergent, NaCl, and PVP, each contributing to the effectiveness of the extraction process.

The dishwashing detergent acts as a surfactant, breaking down cell membranes and denaturing proteins,

thus facilitating the release of DNA into the solution. There are several brands of dishwashing detergent

in the market. Although each product has different ingredients, comparative studies have indicated that

any dishwashing detergent could extract DNA (Ogata et al. 2010). Because the concentration of detergent

is different for each brand, preliminary experiments to determine the most effective concentration of

detergent is necessary. In the current study, we initially tried to use 1 µL of 'green juice' extracted by an

extraction buffer containing only 2 % dish washing detergent, directly as a template DNA for PCR.

Although 1.5 kbp PCR products were successfully obtained, we failed to digest the PCR products by

*Hae*III restriction enzyme, probably due to inhibiting contaminants in the 'green juice.' We therefore

included salting-out/salting-in for proteins, and ethanol precipitation for DNA in the protocol (Fig. 1).

NaCl is indispensable to precipitate DNA by ethanol, while it is expected to work as salt to prevent

proteins from dissolving in or precipitating from the buffer. Instead of using dishwashing detergent,

laundry detergent may be used (Nasiri et al. 2005). However, measuring a specific amount of detergent

powder and dissolving it into a buffer may be more tedious than using liquid type detergent, though any

commercially available products commonly found in the laboratory could be a key factor in terms of cost

and convenience. The PVP, the other ingredient in the extraction buffer (Fig. 1), was included, because it

efficiently removes phenolic compounds, which interact with DNA and prevent DNA extraction

(Heikrujam et al. 2020). Extraction buffer without PVP led to failure in *Hae*III restriction digestion in the

current protocol for *C. japonica*, though PCR was successful (data not shown). If samples contain less

phenolic compounds or only PCR application is needed, PVP could be removed from the extraction

buffer.

In conclusion, we successfully developed a simple and effective DNA extraction method that is

applicable to conifers, a plant group traditionally difficult to process due to inhibitory substances. The

successful application of this method in genotyping the *MS4* gene in *C. japonica* suggests its potential

utility in a wide range of plant species, making it a valuable tool for genetic analysis in settings where

access to specialized equipment is limited. This approach broadens the scope of molecular biology

applications and offers a practical solution for DNA extraction in diverse and challenging environments.

Acknowledgments

This research was supported by Bio-oriented Technology Research Advancement Institution

(BRAIN) Grant (JPJ007097; Project ID 28013B) and FFPRI Grant (#201421, #201906). Samples used in

the simplified DNA extraction method were collected by Mr. Iwai of the Niigata Prefectural Forest

Research Institute. The authors would like to thank Enago (www.enago.jp) for the English language

review.

Data Archiving Statement

The sequence data used in this study are archived in the DDBJ under the accession number

ICQT01013259.

References

Baird NA, Etter PD, Atwood TS, et al (2008) Rapid SNP discovery and genetic mapping using

sequenced RAD markers. PLoS One 3:e3376

Cheng Y-H, Liaw J-J, Kuo C-N (2018) REHUNT: a reliable and open source package for restriction

217   enzyme hunting. BMC Bioinformatics 19:178

218   Dairawan M, Shetty PJ (2020) The Evolution of DNA Extraction Methods. AJBSR 8:39

219   de Andrade BS, Dias BTC, da Silva ECS, et al (2024) DNA extraction from strawberry (*Fragaria*

220   *ananassa*) as a practical tool for teaching genetic content. Editora 719–732

221   Dipta B, Sood S, Mangal V, et al (2024) KASP: a high-throughput genotyping system and its applications

222   in major crop plants for biotic and abiotic stress tolerance. Mol Biol Rep 51:508

223   Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue.

224   PHYTOCHEMICAL BULLETIN 19:11–15

225   Grienenberger E, Kim SS, Lallemand B, et al (2010) Analysis of TETRAKETIDE α-PYRONE

226   REDUCTASE function in Arabidopsis thaliana reveals a previously unknown, but conserved,

227   biochemical pathway in sporopollenin monomer biosynthesis. Plant Cell 22:4067–4083

228   Hasegawa Y, Ueno S, Matsumoto A, et al (2018) Fine mapping of the male-sterile genes (*MS1*, *MS2*,

229   *MS3*, and *MS4*) and development of SNP markers for marker-assisted selection in Japanese

230   cedar (*Cryptomeria japonica* D. Don). PLoS One 13:e0206695

231   Hasegawa Y, Ueno S, Wei F-J, et al (2020) Development of diagnostic PCR and LAMP markers for

232   *MALE STERILITY 1* (*MS1*) in *Cryptomeria japonica* D. Don. BMC Res Notes 13:457

233   Heikrujam J, Kishor R, Behari Mazumder P (2020) The chemistry behind plant DNA isolation protocols.

234   In: Biochemical Analysis Tools - Methods for Bio-Molecules Studies. IntechOpen

235   Hirayama S, Iwai J, Higuchi Y, et al (2021) Selection of trees with male sterile genes except for *MALE*

236   *STERILITY 1* in *Cryptomeria japonica* D. don (In Japanese). J Jpn For Soc 103:161–167

Kakui H, Ujino-Ihara T, Hasegawa Y, et al (2023) A single-nucleotide substitution of *CjTKPR1*

determines pollen production in the gymnosperm plant *Cryptomeria japonica*. PNAS Nexus

2:gad236

Latorra D, Campbell K, Wolter A, Hurley JM (2003) Enhanced allele-specific PCR discrimination in

SNP genotyping using 3' locked nucleic acid (LNA) primers. Hum Mutat 22:79–85

Le Provost G, Herrera R, Paiva JAP, et al (2007) A micromethod for high throughput RNA extraction in

forest trees. Biol Res 40:291–297

Lu Q, Hu H, Mo J, Shu L (2012) Enhanced amplification of bacterial and fungal DNA using a new type

of DNA polymerase. Australas Plant Pathol 41:661–663

Miyajima D, Yoshii E, Hosoo Y, Taira H (2010) Cytological and genetic studies on male sterility in

Cryptomeria japonica D. Don (Shindai 8) (In Japanese). Journal of the Japanese Forest Society

92:106–109

Murray MG, Thompson WF (1980) Rapid isolation of high molecular weight plant DNA. Nucleic Acids

Res 8:4321–4325

Nasiri H, Forouzandeh M, Rasaee MJ, Rahbarizadeh F (2005) Modified salting-out method: high-yield,

high-quality genomic DNA extraction from whole blood using laundry detergent. J Clin Lab

Anal 19:229–232

Ogata F, Kawakami N, Naito S, Tsuzuki H (2010) An examination about the DNA extraction conditions

of Broccoli (*Brassica oleracea* var. *italica*) (In Japanese). Bulletin of Junshin Junior College

15–18

Peterson BK, Weber JN, Kay EH, et al (2012) Double digest RADseq: an inexpensive method for de

novo SNP discovery and genotyping in model and non-model species. PLoS One 7:e37135

Schenk JJ, Becklund LE, Carey SJ, Fabre PP (2023) What is the "modified" CTAB protocol?

Characterizing modifications to the CTAB DNA extraction protocol. Appl Plant Sci 11:e11517

Suyama Y, Matsuki Y (2015) MIG-seq: an effective PCR-based method for genome-wide

single-nucleotide polymorphism genotyping using the next-generation sequencing platform. Sci

Rep 5:16963

Tsuruta M, Maruyama TE, Ueno S, et al (2021) Marker-assisted selection for pollen-free somatic plants

of sugi (Japanese cedar, *Cryptomeria japonica*): A simple and effective methodology for

selecting male-sterile mutants with ms1-1 and ms1-2. Front Plant Sci 12:748110

Ueno S, Tsuruta M, Moriguchi Y, Hasegawa Y (2022) A manual to genotype male sterility gene (*MS1*)

mutations in *Cryptomeria japonica* D. Don (In Japanese)

Untergasser A, Cutcutache I, Koressaar T, et al (2012) Primer3--new capabilities and interfaces. Nucleic

Acids Res 40:e115

Watanabe M, Ueno S, Hasegawa Y, et al (2024) Development and application of a KASP marker for

marker-assisted selection against the male-sterile gene *MALE STERILITY 4* (*MS4*) in Japanese

cedar (*Cryptomeria japonica* D. Don). New Forests 1–14

Xu D, Qu S, Tucker MR, et al (2019) Ostkpr1 functions in anther cuticle development and pollen wall

formation in rice. BMC Plant Biol 19:104

Yadav S, Ram C, Singh S, Rana MK (2018) Plant genomic DNA isolation-the past and the present, a

review. Ind J Plant Genet Resour 31:315

Figure legends

Fig. 1 Workflow of the simple DNA extraction method

Fig. 2 Schematic representation of *MS4* (*TKPR1*) and PCR-RFLP in *Cryptomeria japonica*

Yellow rectangles indicate CDS (coding sequences), which are divided by intron sequences shown by

black lines. *MS4* consists of five exons and introns with a total of 3,323 bp from the start to the stop

codon. The CDS is translated into 333 amino acid residues, the 82^{nd} of which is cysteine (C) and arginine

(R) for the wild type and the mutant, respectively. This is the causative mutation of male sterility by *MS4*.

The restriction enzyme, *Hae*III, can recognize the GGCC, the mutant nucleic acid sequence, while the

wild type sequence cannot be cut by the enzyme. The PCR-RFLP primer pairs are shown by red

triangles; the PCR products from *ms4* allele contain two *Hae*III recognition sequences shown by black

triangles, the C82R site, and that in the third intron common with the products from the wild type allele.

The resulting electrophoretic profile, with gel wells in gray, shows thee, four, and two bans for *ms4*/*ms4*,

Ms4/*ms4*, and *Ms4*/*Ms4* genotype, respectively

Fig. 3 Photographic images for agarose gel electrophoresis

(a) Development of PCR-RFLP for *MS4* gene: PCR products (lanes 1–4) and *HaeIII*-digested products

(lanes 5–8) for `Shindai 8,` `S8HK5,` `Kamikiri Niigata-58,` `Higashikanbara-5` with genotypes of

ms4/*ms4*, *Ms4*/*ms4*, *Ms4*/*ms4*, and *Ms4*/*Ms4*, respectively. (b) DNA extracted by the simple extraction

using the dish detergent buffer was used for PCR-RFLP genotyping. *Hae*III-digested products for

`Shindai 8,' `Kamikiri Niigata-58,` `S8 × Gosenshi-1,` `Setsugai Niigata-1,` and `Kitakanbara-1` are in

lanes 9–13. The smallest bands (less than 100 bp) may result from primer dimers or unspecific

amplification products. (c) Reproducibility was tested by independently extracting DNA eight times from

the same sample individual (`Y938`) using the dish detergent buffer, followed by PCR-RFLP analysis:

PCR products (lanes 14–19 and 26–27) and *HaeIII*-digested products (lanes 20–25 and 28–29). In all (a),

(b) and (c), 100 bp ladder was run on the lane M

Tables and Figures

Table 1 Japanese cedar samples used in the current study

No[#]	ID	Male flower phenotype	*MS4* genotype[$]
1	Shindai 8	Sterile	*ms4/ms4*
2	S8HK5	Fertile	*Ms4/ms4*
3	Kamikiri Niigata-58	Fertile	*Ms4/ms4*
4	Higashikanbara-5	Fertile	*Ms4/Ms4*
5	S8 × Gosenshi-1	Fertile	*Ms4/ms4*
6	Setsugai Niigata-1	Fertile	*Ms4/Ms4*
7	Kitakanbara-1	Fertile	*Ms4/Ms4*
8	Y938	Fertile	*Ms4/Ms4*

[#] For development of the PCR-RFLP protocol, trees No 1–4 were used, while for development of the

simple DNA extraction method, trees No 1 and 4–7 were used. Tree No 8 was used to test the

reproducibility of the protocol.

[$] Genotypes were known previously for trees No 1–5 (Miyajima *et al*. 2010; Hasegawa *et al*. 2018;

Hirayama *et al*. 2021), while they were unknown for trees No 6–8 before the present study.

Fig. 1

Fig. 2

Fig. 3