Quantitative physiology and biomass composition of *Cyberlindnera jadinii* in

ethanol-grown cultures

Marcel A. Vieira-Lara[#], Marieke Warmerdam, Erik A. F. de Hulster, Marcel van den Broek, Jean-

Marc Daran, Jack T. Pronk*

Department of Biotechnology, Delft University of Technology, van der Maasweg 9, 2629, HZ

Delft, the Netherlands

Corresponding authors:

[#]M.A.VieiraLara@tudelft.nl

*J.T.Pronk@tudelft.nl, tel. +31 15 2782473

17 Running title: From ethanol to protein with *Cyberlindnera jadinii*

Abstract

Background

Elimination of greenhouse gas emissions in industrial biotechnology requires replacement of carbohydrates by alternative carbon substrates, produced from CO_2 and waste streams. Ethanol is already industrially produced from agricultural residues and waste gas and is miscible with water, self-sterilising and energy-dense. The yeast *C. jadinii* can grow on ethanol and has a history in the production of single-cell protein (SCP) for feed and food applications. To address a knowledge gap in quantitative physiology of *C. jadinii* during growth on ethanol, this study investigates growth kinetics, growth energetics, nutritional requirements, and biomass composition of *C. jadinii* strains in batch, chemostat and fed-batch cultures.

Results

In aerobic, ethanol-limited chemostat cultures, *C. jadinii* CBS 621 exhibited a maximum biomass yield on ethanol ($Y_{x/s}^{max}$) of 0.83 $g_{biomass} \cdot (g_{ethanol})^{-1}$ and an estimated maintenance requirement for ATP (m_{ATP}) of 2.7 $mmol \cdot (g_{biomass})^{-1} \cdot h^{-1}$. Even at specific growth rates below 0.05 h^{-1}, a stable protein content of approximately 0.54 $g_{protein} \cdot (g_{biomass})^{-1}$ was observed. At low specific growth rates, up to 17% of the proteome consisted of alcohol dehydrogenase proteins, followed by aldehyde dehydrogenases and acetyl-CoA synthetase. Of 13 *C. jadinii* strains evaluated, 11 displayed fast growth on ethanol ($\mu_{max} > 0.4$ h^{-1}) in mineral medium without vitamins, and CBS 621 was found to be a thiamine auxotroph. The prototrophic strain *C. jadinii* CBS 5947 was grown on an inorganic salts medium in fed-batch cultures (10-L scale) fed with pure ethanol. Biomass concentrations in these cultures increased up to 100 $g_{biomass} \cdot (kg_{broth})^{-1}$, with a biomass yield of 0.65 $g_{biomass} \cdot (g_{ethanol})^{-1}$. Model-based simulation, based on quantitative parameters determined in chemostat cultures, adequately predicted biomass production. A different protein content of chemostat- and fed-batch-grown biomass (54% and 42%, respectively) may reflect the more dynamic conditions in fed-batch cultures.

Conclusions

Analysis of ethanol-grown batch, chemostat and fed-batch cultures provided a quantitative physiology baseline for fundamental and applied research on *C. jadinii*. Its high maximum growth rate, high energetic efficiency of ethanol dissimilation, simple nutritional requirements and high protein content, make *C. jadinii* a highly interesting platform for production of SCP and other products from ethanol.

Keywords (3-10):

Ethanol, yeast, *Candida utilis*, protein, physiology, growth rate, chemostat

Introduction

Establishing a circular, net-zero emission industrial biotechnology industry requires replacing refined sugars, which are currently the predominant carbon sources for industrial fermentation processes, with feedstocks derived from CO_2 and renewable waste streams (1, 2). Especially for products that require ATP-intensive synthesis, achieving high titres, rates, and yields can be facilitated by first converting primary carbon sources into smaller organic molecules. These molecules can then serve as substrates in separate, intensive aerobic fermentation processes (1). In addition to microbial fermentation of lignocellulosic residues from agriculture and forestry (3), such small molecules can be produced by gas fermentation (4), electrochemical production from CO_2 (1, 5) and, potentially, from microbial electrosynthesis (6, 7). The resulting organic molecules include C_1 compounds such as formic acid and methanol as well as the C_2 compounds acetic acid and ethanol (1).

Ethanol is an energy-dense substrate that is completely miscible in water and, at high concentrations, self-sterilising. Especially at low pH, ethanol is less toxic than other C_1 and C_2 compounds generated from CO_2. Ethanol can be produced electrochemically from CO_2 at lab-scale with a Faraday efficiency of 68% (8). Production of ethanol from agricultural residues by yeast (0.3-0.5 $g_{ethanol} \cdot (g_{sugar})^{-1}$) (9) and from waste gases by *Clostridium autoethanogenum* (up to 200 $mmol_{ethanol} \cdot (g_{biomass})^{-1}$ and 30% carbon recovery in ethanol) (4, 10, 11) have both transitioned to industrial scale. Ethanol can be readily assimilated by a variety of microorganisms (12), including yeast species that are already applied in large-scale sugar-based fermentation processes. However, compared to the wealth of information on quantitative physiology of sugar-grown aerobic cultures of industrially relevant yeast species, quantitative information on the stoichiometry, kinetics, and energetics of ethanol-grown cultures of such yeasts is scarce (13).

After diffusion over the yeast plasma membrane (14), ethanol is first converted to acetyl-CoA.

This conversion involves the enzymes NAD^+-dependent alcohol dehydrogenase,

$NAD(P)^+$-dependent acetaldehyde dehydrogenase and acetyl-CoA synthetase. Further

dissimilation and assimilation of acetyl-CoA involves the tricarboxylic acid (TCA) cycle,

glyoxylate cycle and gluconeogenesis. Dissimilation of ethanol to CO_2 generates reduced

cofactors (5 NAD(P)H and 1 $FADH_2$) and requires input of one ATP-equivalent due to the

combination of the investment of 2 ATP equivalents in the acetyl-CoA synthetase reaction (15)

and the formation of 1 ATP equivalent by substrate-level phosphorylation in the TCA cycle.

Energy coupling of the subsequent reoxidation of NADH and $FADH_2$ by mitochondrial

respiration has a strong impact on the net ATP yield from ethanol dissimilation and, thereby, on

product yields on ethanol and on oxygen. The efficiency of this energy coupling is expressed by

the P/O ratio, which indicates how many moles of ATP oxidative phosphorylation generates

when a single electron pair (for example derived from NADH) is transferred to oxygen.

Involvement of an active multi-subunit proton-pumping NADH dehydrogenase ('Complex

I') in the mitochondrial respiratory chains of yeasts strongly influences their P/O ratio (16, 17).

Several industrially relevant yeasts, such as *Saccharomyces cerevisiae* and *Kluyveromyces*

lactis, lack the genetic information for synthesizing Complex I and, instead, only harbour single-

subunit, non-proton-translocating NADH dehydrogenases (18, 19). Other yeasts encode the

genetic information for synthesizing both Complex I and non-proton-translocating NADH

dehydrogenases (20). In such yeasts, energy coupling of respiration depends on how regulation

of protein synthesis and/or activity determine the distribution of NADH oxidation over Complex I

and less efficient systems.

The ascomycete budding yeast *Cyberlindnera jadinii* can grow on a variety of carbon

sources, including sugars, organic acids and alcohols, and on nitrogen sources including

ammonium, nitrate, urea and amino acids (21). Although no extensive quantitative physiology

studies have been published on ethanol-grown cultures of *C. jadinii*, this yeast has been reported to show high biomass yields on ethanol ($g_{biomass} \cdot (g_{ethanol})^{-1}$) (18). This observation is consistent with the presence, in the genome of *C. jadinii*, of the genetic information for encoding a Complex I NADH dehydrogenase (22). Based on dry weight, glucose-grown biomass of *C. jadinii* contains approximately 50% protein (18). *C. jadinii* strains have been reported to grow without vitamin supplementation (23), which enables simple processes and may offer options to select strains with high vitamin content.

C. jadinii is a polyploid teleomorph of *Candida utilis* (21, 24), which has a long history of safe use in biotechnology. *C. jadinii* biomass can be used as single-cell protein (SCP) as a food ingredient (25) and for animal nutrition (26). Nutritional studies have shown that *C. jadinii* SCP can functionally replace conventional protein sources in fish, chicken and swine feed (27-29). A historical example of SCP is Pruteen®, a product launched by Imperial Chemical Industries (ICI) in 1980, produced from hydrocarbon-derived methanol using *M. methylotrophus* (Matassa et al., 2016; Westlake, 1986). Although Pruteen® production was discontinued for economic reasons, interest in SCP as a viable protein source in animal feed continues to grow given the emergence of alternative feedstocks (30, 31).   SCP has the potential to alleviate negative environmental impacts of food systems (32-35) due to a lower carbon footprint, water use and land usage than those of other protein sources (36).

Large-scale industrial production of microbial biomass typically involves aerobic fed-batch cultures. Because of oxygen transfer limitations in industrial bioreactors, achieving high biomass concentrations in such processes requires a decreasing specific growth rate profile (37). For understanding and predicting microbial performance in such industrial contexts, knowledge on growth-rate-dependent physiology, with special attention for maintenance-energy requirements, biomass yields on ethanol and oxygen, and, in the case of SCP production, biomass composition is essential. This study therefore aims to investigate the quantitative

physiology of *C. jadinii* with ethanol as a substrate and its implications for SCP production. To

this end, we studied growth stoichiometry, growth energetics, vitamin requirements and biomass

composition in batch and chemostat cultures of *C. jadinii* strains. Based on quantitative

information derived from these experiments, a fed-batch cultivation protocol for SCP production

from pure ethanol as carbon source was designed and conducted at 10-L scale.

Materials & Methods

Yeast strains and media. *Cyberlindnera jadinii* strains CBS 621, CBS 567, CBS 841, CBS 842,

CBS 890, CBS 1516, CBS 1517, CBS 1600, CBS 2160, CBS 4511, CBS 5609, CBS 5947, and

CBS 7232 were obtained from the Westerdijk Institute (Utrecht, The Netherlands).

Saccharomyces cerevisiae CEN.PK113-7D was provided by Dr. Peter Kötter (Goethe

University, Frankfurt, Germany). Cultures were grown on a synthetic medium (SM) containing 5

$g{\cdot}L^{-1}$ $(NH_4)_2SO_4$, 3 $g{\cdot}L^{-1}$ KH_2PO_4 0.5 $g{\cdot}L^{-1}$ $MgSO_4{\cdot}7H_2O$, 15 $mg{\cdot}L^{-1}$ EDTA, 4.5 $mg{\cdot}L^{-1}$

$ZnSO_4{\cdot}7H_2O$, 0.3 $mg{\cdot}L^{-1}$ $CoCl_2{\cdot}6H_2O$, 1 $mg{\cdot}L^{-1}$ $MnCl_2{\cdot}H_2O$, 0.3 $mg{\cdot}L^{-1}$ $CuSO_4{\cdot}5H_2O$, 4.5 $mg{\cdot}L^{-1}$

$CaCl_2{\cdot}2H_2O$, 3 $mg{\cdot}L^{-1}$ $FeSO_4{\cdot}7H_2O$, 0.4 $mg{\cdot}L^{-1}$ $Na_2MoO_4{\cdot}2H_2O$, 1 $mg{\cdot}L^{-1}$ H_3BO_3, 0.1 $mg{\cdot}L^{-1}$ KI.

The pH of the medium was adjusted to 6.0 with 1 M KOH. After autoclaving at 120 °C for 20

min, 1 $mL{\cdot}L^{-1}$ of a filter-sterilized vitamin solution (0.05 $g{\cdot}L^{-1}$ D-(+)-biotin, 1.0 $g{\cdot}L^{-1}$ D-calcium

pantothenate, 1.0 $g{\cdot}L^{-1}$ nicotinic acid, 25 $g{\cdot}L^{-1}$ *myo*-inositol, 1.0 $g{\cdot}L^{-1}$ thiamine hydrochloride, 1.0

$g{\cdot}L^{-1}$ pyridoxal hydrochloride, 0.2 $g{\cdot}L^{-1}$ 4-aminobenzoic acid) (38) was aseptically added. In

some experiments, individual or all vitamins were omitted as indicated. Glucose solutions

autoclaved at 110 °C for 20 min or pure ethanol were aseptically added to the medium along

with the vitamin solution. For strain storage, 30% (v/v) glycerol (final concentration) was added

to shake-flask cultures grown overnight at 200 rpm and 30 °C in YPD medium (10 $g{\cdot}L^{-1}$ yeast

extract, 20 $g{\cdot}L^{-1}$ peptone and 20 $g{\cdot}L^{-1}$ glucose) and 1 mL samples were kept at -80 °C.

Shake flask and microtiter plate cultivation. Shake-flask cultures, grown in 500-mL flasks

containing 100 mL SM supplemented with either 20 $g{\cdot}L^{-1}$ D-glucose (SMD) or 7.5 $g{\cdot}L^{-1}$ ethanol

(SME), were incubated at 30 °C and 200 rpm in Innova incubators (Brunswick Scientific Edison,

NJ). Initial cultures were inoculated from frozen glycerol samples, incubated overnight, and used

for a second inoculation in the same medium. Upon reaching early exponential phase, these

pre-cultures were used to inoculate a third culture, either in shake flasks or microtiter plates, at

an initial OD_{660} of 0.2-1.0. Absorbance at 660 nm (OD_{660}) was measured with a Jenway 7200

spectrophotometer after accurate dilution of samples to yield OD_{660} values between 0.1 and 0.3.

Specific growth rates were calculated by fitting exponential phase data (OD 0.2 – 5.0) to the

equation $In\ OD(t) = In\ OD(0) + \mu_{max} \cdot t$, in which μ_{max} (h^{-1}) represents the maximum specific growth

rate. Cultivation in 96-well plates was done in a Growth Profiler 960 (Enzyscreen BV,

Heemstede, the Netherlands), operated at 30 °C and at 250 rpm. Images from the bottom of the

incubator plates were taken every 20 min. The number of green pixels per picture (Green Value)

was quantified with the GP960Viewer software. Green Values correlate with OD_{660} and were

used to calculate maximum specific growth rates from serial dilutions (39). Mathematical details

and assumptions of the method are described in Supp. Material S3.

Chemostat cultivation. Chemostat cultures of *C. jadinii* and *S. cerevisiae* were grown at 30 °C

in 1.5 L bioreactors connected to an ez-Control (Applikon Getinge, Delft, The Netherlands).

Glucose- and ethanol-limited cultures were grown on SM with 5 $g \cdot L^{-1}$ glucose and 3.75 $g \cdot L^{-1}$

ethanol, respectively, unless otherwise stated, with 0.2 $g \cdot L^{-1}$ Pluronic PE 6100 anti-foaming

agent (BASF, Ludwigshafen, Germany). The working volume of cultures was kept at 1.0 L with

an electrical level sensor connected to the effluent pump. Unless stated otherwise, culture pH

was maintained at 5.0 by automatic addition of 2 M KOH. Cultures were stirred at 800 rpm with

2 Rushton impellers and sparged with air at 0.5 L/min. CO_2 and O_2 contents in the in- and off-

-gas were measured with a multi-gas analyser (MultiExact 4100 Multi-gas Analyser, Servomex).

Dissolved oxygen concentration was monitored with an autoclavable oxygen electrode

(Applisens, Schiedam The Netherlands). After inoculation and a batch cultivation phase, the

continuous cultivation mode was initiated when a sharp decrease of the CO_2 output indicated

substrate depletion. At least two pre-steady-state samples were taken from the effluent line

when at least 5 volume changes had passed since the onset of continuous cultivation. When

values of dry weight and CO_2 differed by less than 5% for two consecutive volume changes,

steady-state samples were collected directly from the culture. Carbon recoveries, calculated

based on a yeast biomass carbon content of 45.5% (w/w) (40) were between 95 and 105%

unless stated otherwise. Maintenance coefficient and maximum biomass yield on substrate

were calculated by linear regression of substrate consumption rates (q_s) plotted against the

dilution rate (D) according to the Pirt equation (eq. 1; (41).

$$q_S = \frac{D}{Y_{x/s}^{max}} +\ m_S \ (1)$$

Fed-batch cultivation. Fed-batch cultivation was conducted at 30 °C in 20-L Bio Bench

reactors (Applikon Getinge, Delft, The Netherlands) with an initial working volume of 8.0 L.

Stirrer speed was kept at 800 rpm and pressurised air (20.95% O_2, 0.04% CO_2) was supplied at

a flow rate of 7 $L \cdot min^{-1}$. An overpressure of 1.0 bar (2.0 bar total pressure) was applied

throughout cultivation to enhance oxygen transfer. A polarographic oxygen electrode (model 32

275 6800; Ingold) was used to monitor the dissolved oxygen concentration to ensure it

remained above 15% of air saturation (42).

The medium used for fed-batch cultivation contained 2 $g \cdot L^{-1}$ $(NH_4)_2SO_4$, 20 $g \cdot L^{-1}$ KH_2PO_4 6 $g \cdot L^{-1}$

$MgSO_4 \cdot 7H_2O$, and trace elements (150 $mg \cdot L^{-1}$ EDTA, 45 $mg \cdot L^{-1}$ $ZnSO_4 \cdot 7H_2O$, 3 $mg \cdot L^{-1}$

$CoCl_2 \cdot 6H_2O$, 10 $mg \cdot L^{-1}$ $MnCl_2 \cdot H_2O$, 3 $mg \cdot L^{-1}$ $CuSO_4 \cdot 5H_2O$, 45 $mg \cdot L^{-1}$ $CaCl_2 \cdot 2H_2O$, 30 $mg \cdot L^{-1}$

$FeSO_4 \cdot 7H_2O$, 4 $mg \cdot L^{-1}$ $Na_2MoO_4 \cdot 2H_2O$, 10 $mg \cdot L^{-1}$ H_3BO_3, 1 $mg \cdot L^{-1}$ KI) and 0.2 $g \cdot L^{-1}$ Pluronic PE

6100 antifoam. Ammonium hydroxide (25% w/v) was used to control culture pH at 5.0 and to

simultaneously provide on-demand nitrogen-source supply. For the initial batch phase, which

was inoculated at an OD_{660} of 0.2, 12 $g \cdot L^{-1}$ ethanol was used as carbon source. When the end of

the exponential growth phase was indicated by a simultaneous decrease of CO_2 production and

an increase in the dissolved-oxygen concentration, the fed-batch phase was initiated. During

this phase, absolute ethanol was used as carbon source and pumped into the reactor with a

controllable Masterflex model 7518-00 pump (Cole-Parmer Instrument Company, Chicago, IL).

The feed rate was controlled by a MFCS SCADA Software (Sartorius Stedim Biotech GmbH,

Göttingen, Germany) according to equation 2. Over the duration of the fed batch phase, this

feed profile resulted in flow rates between 0.006 and 0.050 $kg_{ethanol} \cdot h^{-1}$.

$$F_{ethanol}(t) = \left(\frac{\mu}{Y_{x/s}^{max}} + m_s\right) \cdot c_{x,0} \cdot W_0 \cdot e^{\mu \cdot t} \quad (2)$$

In equation 2, $F_{ethanol}$ indicates the feed rate of ethanol in $kg_{ethanol} \cdot h^{-1}$, $c_{x,0}$ the initial biomass

concentration ($g_{biomass} \cdot (kg_{broth}^{-1})$), W_0 the initial culture weight (kg_{broth}) and t the time (h) after

starting the feed. $Y_{x/s}^{max}$ is the maximum biomass yield on substrate in the absence of

maintenance ($g_{biomass} \cdot (g_{ethanol}^{-1})$) and m_s is the maintenance coefficient ($g_{ethanol} \cdot (g_{biomass}^{-1}) \cdot h^{-1}$), both

calculated from chemostat experiments. The specific growth rate was kept at 0.10 h^{-1} until the

dissolved oxygen concentration decreased to 15% of air saturation. At that point, the feed was

changed from exponentially increasing to constant. The fed-batch process was simulated in

Python essentially as described by Hensing et al., 1995 (37). The volumetric mass transfer

coefficient ($k_L a$) (43) was experimentally determined to be 210 ± 11 h^{-1} with the dynamic

technique of absorption (44), and a value of 200 h^{-1} was used for the simulation. Scripts are

found in Appendix 1.

Analytical methods. Biomass dry weight in culture samples was measured as previously

described (45). Briefly, dried nitrocellulose filters (pore size 0.45 µm, Gelman Laboratory, Ann

Arbor, MI) stored at 80 °C were weighed and used to filter 10.0 mL culture samples. Fed-batch

samples were diluted in demineralized water to a concentration < 5 $g_{biomass} \cdot (kg_{broth})^{-1}$ prior to dry

weight measurements. To correct for increased broth density at the end of the process, density

was measured by weighing 10.0 mL of broth and used to convert values from $g_{biomass} \cdot (L_{broth})^{-1}$ to

$g_{biomass} \cdot (kg_{broth})^{-1}$ (for samples with > 70 $g_{biomass} \cdot (kg_{broth})^{-1}$). After washing with demineralized

water, filters were dried for 20 min at 350 W output in a microwave (Bosch, Stuttgart, Germany)

and weighed again. For determination of metabolite concentrations in culture supernatants,

samples were rapidly quenched with cold steel beads and filtered as described by Mashego et

al., 2003 (46). Concentrations of glucose and metabolites were determined by HPLC (47).

Biomass-specific rates of production and consumption of substrates and products, respectively,

as well as carbon recoveries, were calculated with a script available from Appendix 2.

Determination of biomass composition. Protein content of yeast biomass was determined

with the Biuret method according to Verduyn et al. (16), and lipid content according to Izard &

Limberger, and Johnson et al. (48, 49) with modifications. Briefly, cell suspension (20 mL) was

centrifuged at 10,000 g for 5 min. The pellet was washed and resuspended in demineralized

water to give a total volume of 10.0 mL H_2O and stored at -20 °C. For protein measurements,

thawed samples were mixed with 3 M NaOH (2:1), boiled at 100 °C for 10 min and cooled on

ice. After adding $CuSO_4 \cdot 5H_2O$ (2.5% w/v) to samples (1:3), they were incubated at room

temperature for 5 min and centrifuged at 13,000 rpm for 5 min. The absorbance of the clear

supernatant was then measured at 510 nm. A solution of 5 $g \cdot L^{-1}$ bovine serum albumin (Merck-

Sigma A9647) was used to construct a standard curve. For total lipid determination, 100 µL of

sample were mixed with 1.0 mL H_2SO_4 and vortexed. Samples were incubated at 100 °C for 10

min and cooled for 5 min. Subsequently, 2.5 mL of phosphoric acid-vanillin reagent (1.2 $g \cdot L^{-1}$

vanillin and 68% phosphoric acid) were added, samples were vortexed and incubated at 37 °C

for 15 min. Samples were cooled to room temperature and absorbance was read at 530 nm.

Standard solutions of sunflower oil (Merck-Sigma S5007) in chloroform (0 – 1.0 mg/mL) were

used to construct a calibration curve. Chloroform was fully evaporated prior to H_2SO_4 addition.

To measure RNA and carbohydrate contents of yeast biomass, culture samples were washed

once with demineralized water and pellets containing 3 to 5 mg biomass were stored at -20 °C.

A 2.0 g·L^{-1} solution of yeast RNA (Sigma 10109223001) was prepared in 10 mM Tris-HCl, pH

8.0 and also stored at -20 °C. Before analysis, the stock solution was diluted to obtain standard

solutions with RNA concentrations ranging from 5 to 60 mg·L^{-1}. RNA was extracted (50) and

quantified by measuring absorbance at 260 nm (U-3010 spectrophotometer, Hitachi). Total

carbohydrate contents in biomass were quantified according to (51). Briefly, 5 mL of

concentrated H_2SO_4 were added to 1 mL of sample, followed by the addition of 1 mL of phenol

solution (50 g·L^{-1}). Samples were incubated at 100 °C for 15 min and subsequently cooled to

room temperature. Absorbance was measured at 488 nm. A standard solution of glucose and

mannose (2:1) was used to construct a calibration range (0.0 - 0.2 g·L^{-1}). To correct for the

presence of nucleic acid pentoses, relative extinction coefficients of 0.36 and 0.26 (versus the

coefficient obtained from the standard curve) were used for RNA and DNA, respectively

(adapted from (40)). The following formula was used:

$$Abs_{sample} = \varepsilon_{standard} \cdot [carbohydrates] + \varepsilon_{RNA} \cdot [RNA] + \varepsilon_{DNA} \cdot [DNA] \tag{3}$$

Where Abs stands for absorbance, $\varepsilon_{standard}$ was obtained from freshly prepared standard curves,

RNA content was measured, and DNA content was assumed to be 0.5% of the biomass dry

weight for all samples.

Proteome analysis. Biomass samples (20 mL, containing approximately 3 g biomass·L^{-1}) were

centrifuged and washed once with demineralized water. Pellets were resuspended in cold

methanol (-20°C) and lysed with a Precellys Homogenizer (Bertin technologies SAS, Montigny-

le-Bretonneux, France) at 6,800 rpm for 15 s. Total protein concentrations of the resulting

samples were measured by Qubit protein assay, and proteins were isolated by Bligh and Dyer

extraction. Isolated proteins were processed further by reduction, alkylation, and trypsin

digestion (52). Samples were analysed in technical triplicates by liquid chromatography coupled

to tandem mass spectrometry (LC-MS/MS) using a Vanquish UHPLC coupled to a Qexactive

Plus (Thermo Fisher Scientific, Waltham, USA). Peptides were separated using reverse-phase

chromatography with a gradient of water with 0.1% formic acid (solvent A) and 20% water and

0.1% formic acid in acetonitrile (solvent B) from 5% B to 60% B in 52 min. Data-independent

acquisition (DIA) was performed with a resolution setting of 17,500 within the 400 to 1,200 m/z

range and a maximum injection time of 20 ms, followed by high-energy collision-induced

dissociation activated (HCD) MS/MS scans with a resolution setting of 17,500 and injection time

set to auto. Four DIA MS/MS fragmentation scans were performed after each full scan and the

mass range was divided into 100 m/z windows. Raw files were analysed with the Pulsar search

engine and Spectronaut, version 17.5, against the annotated proteins of *C. jadinii* (UniProtKB,

strain ATCC 18201/CBS 1600 [www.uniprot.org]). Label-free quantification was performed using

the top three peptides measured for each protein. Retention time realignment was based on

local (non-linear) regression.

The obtained signal per protein (*raw signal (i)*, proportional to protein copy number) was

multiplied by the respective molecular weight, yielding a *signal (i)*. Subsequently, the *signal (i)*

obtained per protein was divided by the sum of all *signals (i)*'s obtained in the sample and

multiplied by 100. This resulted in the approximate percentage (in weight/weight) by which each

protein contributed to the total proteome mass.

Amino acid analysis. Culture samples were washed once with demineralized water and pellets

containing approximately 50 mg biomass were stored at -20 °C. Total amino acid content of

yeast biomass was determined after hydrolysis of the biomass according to the Accq-Tag

method (Waters). Before hydrolysis, samples were weighed in a headspace vial, which was

subsequently flushed with nitrogen to remove oxygen. Hydrolysis with 6 M HCl was performed

to release free amino acids from proteins and peptides. Free amino acids were derivatized on

the NH_2 terminus with (AQC) 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate. For complete

conversion, samples were incubated for 10 min at 55 °C. Derivatized amino acids were

analysed using UPLC reversed-phase chromatography with diode array detection at 260 nm

(Acquity UPLC, Waters Corporation, Milford, USA). Derivatized amino acids were

chromatographically separated with an Acquity UPLC BEH C18 1.7 μm 2.1×100 mm (Waters)

analytical column and a gradient with Mobile phase A: AccQ·Tag Ultra Eluent A, and Mobile

phase B: AccQ·Tag Ultra Eluent B, starting at 0.1% B to 9.1% B in 5.74 min increasing to 21.2%

B at 7.74 min followed by a ramp to 59.6% B at 8.04 min with a flow rate of 0.7 mL/min.

Glutamine, asparagine, tryptophan and cysteine could not be determined with this method

because glutamine and asparagine are converted to glutamic acid and aspartic acid, and

tryptophan breaks down during hydrolysis.

Whole-genome sequencing. Genomic DNA was isolated from culture samples containing

approximately $7.0 \cdot 10^{9}$ cells with the Qiagen 100/G kit (Qiagen, Hilden, Germany), according to

manufacturer's instructions. Genomic DNA of *C. jadinii* CBS 621 was sequenced in-house on an

Illumina MiSeq sequencer (Illumina, San Diego, CA) to obtain a paired-end library with an

insert-size of 550 bp and 300 bp read length using TruSeq Nano DNA library preparation,

yielding 6.4 Gb of genomic DNA sequence. *C. jadinii* CBS 5947 was sequenced at Macrogen on

an Illumina NovaSeq sequencer to obtain a paired-end library with an insert-size of 550 bp and

151 bp read length using TruSeq DNA PCR-free library preparation, yielding 3.5 Gb of genomic

DNA sequence. Reads of strains CBS 621 and CBS 5947 were mapped against the *C. jadinii*

NBRC 0988 genome (BioProject: PRJDB11630; GCA_024346625.1) using BWA (version

0.7.15) (53). Alignments were processed using SAMtools (version 1.3.1.) (54) and variants were

called by applying Pilon (version 1.18) (55). Heterozygous single-nucleotide polymorphism

(SNP) ratios of each chromosome were extracted from the Pilon produced variant calling file

(VCF) and the distributions were plotted by using ggplot2 (56). *De novo* assembly was

performed on the Illumina reads with the SPAdes genome assembler (version 3.9.0) (57).

Sequencing data are available at NCBI (https://www.ncbi.nlm.nih.gov/) under the BioProject accession number PRJNA1153750.

Vitamin measurements. Analyses of selected vitamins (thiamine, pyridoxine, nicotinic acid and riboflavin) were conducted via Elisa by SGS Nederland B.V., division SGS Food Lab. Biomass samples containing circa 3 $g_{biomass}$ were used.

Software and Statistics. Scripts used for calculations were written in Python (v3.10), using Jupyter Notebook (v 6.4.11). Data visualisation and statistical analysis were conducted in GraphPad Prism (v9.0, GraphPad Software, San Diego, CA).

Results

1. **Biomass yield and protein content in glucose- and ethanol-grown chemostat cultures**

To verify earlier reports that *C. jadinii* shows higher biomass yields on ethanol than *S. cerevisiae* (16, 18), aerobic ethanol-limited chemostat cultures of the model strains *C. jadinii* CBS 621 (18, 58, 59) and *S. cerevisiae* CEN.PK113-7D (40, 60-62) were grown at a dilution rate of 0.10 h^{-1}. Consistent with earlier reports (18), steady-state cultures of *S. cerevisiae* CEN.PK113-7D showed a biomass yield of 0.59 ± 0.01 $g_{biomass} \cdot (g_{ethanol})^{-1}$ (Fig 1A). In four independent chemostat cultures of *C. jadinii* CBS 621, biomass yields were consistently 23% higher (0.74 ± 0.01 $g_{biomass} \cdot (g_{ethanol})^{-1}$) than in those of *S. cerevisiae*. The biomass yields on ethanol of *C. jadinii* CBS 621 were higher than previously reported for chemostat cultures of this strain (0.69 $g_{biomass} \cdot (g_{ethanol})^{-1}$) (16, 18). In contrast to the ethanol-limited cultures, glucose-limited chemostat cultures of *C. jadinii* CBS 621 and *S. cerevisiae* CEN.PK113-7D showed only 6% difference in biomass yield (0.51 ± 0.01 $g_{biomass} \cdot (g_{glucose})^{-1}$ and 0.48 ± 0.00 $g_{biomass} \cdot (g_{glucose})^{-1}$, respectively; Fig 1A). Both species showed lower biomass yields on oxygen in ethanol-limited cultures than in glucose-limited cultures (Fig 1B). However, biomass yields on oxygen in ethanol-limited cultures

of *C. jadinii* were approximately 50% higher in *C. jadinii* than *S. cerevisiae* (Table S1). The

different biomass yields of *C. jadinii* and *S. cerevisiae* on ethanol may reflect the presence and

absence, respectively, of a proton-pumping respiratory complex I in these two yeast species

(16, 22).

Consistent with earlier reports (18, 40), the highest protein content in yeast biomass (0.54

$g_{protein} \cdot (g_{biomass})^{-1}$) was observed in the ethanol-limited *C. jadinii* cultures and the lowest protein

content in the glucose-limited *S. cerevisiae* cultures (0.36 $g_{protein} \cdot (g_{biomass})^{-1}$) (Fig 1C). In the

ethanol-limited cultures of *C. jadinii*, the protein yield on ethanol reached 0.40 $g_{protein} \cdot (g_{ethanol})^{-1}$

(Fig 1D).

2. **Proteome and amino-acid composition of ethanol-grown *C. jadinii***

To further investigate the biomass composition of the ethanol- and glucose-limited chemostat

cultures of *C. jadinii*, proteome analyses were performed. Of 2596 detected proteins, 1656 were

identified with at least 3 different peptides (Supp. Material S1). In the ethanol-limited *C. jadinii*

cultures, the 10 most abundant proteins comprised approximately 25% of the proteome. Two

isoforms each of alcohol dehydrogenase (*Cj*Adh) and acetaldehyde dehydrogenase (*Cj*Ald), as

well as acetyl-CoA synthetase 1 (*Cj*Acs1), which together catalyse the first three steps in

ethanol catabolism (Fig 1E), accounted for 6 of the 10 most highly abundant proteins (Fig 1F).

Key proteins in ethanol assimilation (isocitrate lyase and malate synthase) were also very

highly expressed. While their levels were clearly higher in ethanol-grown cultures, the 6 proteins

involved in ethanol catabolism were also abundant in cells grown under glucose-limitation (Fig

1F). In contrast, levels of elongation factor 1-α and the α-subunit of the F_1F_o-ATP synthase were

similar in glucose-limited and ethanol-limited cultures (Fig 1F).

In view of the 75% higher respiration rate in ethanol-limited cultures than in glucose-limited

cultures of *C. jadinii* CBS 621 (4.4 vs 2.5 mmol $O_2 \cdot (g_{biomass})^{-1} \cdot h^{-1}$, Table S1), levels of proteins

involved in oxidative phosphorylation were also investigated. A set of 71 proteins involved in

oxidative phosphorylation that were represented by at least 2 independent peptides and

included subunits of all 4 respiratory complexes and F_1F_o-ATP synthase (Supp. Material S1)

showed, on average, a 60% higher level in ethanol-limited cultures than in glucose-limited

cultures. Complex I subunits (28 detected) were, on average, 75% higher in ethanol- compared

to glucose-limited samples (Fig. S1A).

The amino acid composition of *C. jadinii*, which is important for food and feed applications of

microbial biomass (31), was analysed and compared to that of *S. cerevisiae*. Biomass from

ethanol- and glucose-grown chemostat cultures of the two yeasts showed similar amino acid

composition, with high glutamate/glutamine levels (Table S2). A low content of methionine in

glucose- and ethanol-grown chemostat cultures of both yeasts (1.35 to 1.80% of the proteome,

Fig 1G) was previously reported for multiple yeasts species (27). The sum of the analysed

amino acid contents followed the same trend as total protein measurements (Fig S1B) but

resulted in lower absolute values (Fig S1C). There is a growing interest in using yeast biomass

in aquaculture to produce fish feed and thereby to curtail overfishing for fishmeal (27, 31, 32). A

comparison of amino acid levels in ethanol-grown *C. jadinii* biomass with reference values for

essential amino acids in fish feed identified methionine as limiting (Fig S1D), with all other

amino acids at or above required levels, assuming yeast biomass as the only diet component.

Figure 1: Effect of growth-limiting carbon source on the physiology of *C. jadinii* and proteome composition. *S. cerevisiae* CEN.PK113-7D and *C. jadinii* CBS 621 were grown in steady-state, aerobic ethanol-limited and glucose-limited chemostats at a dilution rate of 0.10 h^{-1}. (A, B) Biomass yields on substrate and oxygen, respectively. (C) Protein content as a percentage of biomass. (D) Protein yields on substrate. (E) Enzymes involved in conversion of ethanol to acetyl-CoA in yeasts. Adh: alcohol

dehydrogenase, Ald: aldehyde dehydrogenase, Acs: acetyl-CoA synthetase. (F) Top 10 most abundant

proteins (with at least 3 identified peptides) in the proteome of ethanol-grown *C. jadinii* versus the levels

of the same proteins in glucose-limited chemostats in the same species. *Cj* stands for *Cyberlindnera*

jadinii. (G) Protein amino acid residue composition (as a percentage of the total amino acid residue pool).

H stands for highest, and L for lowest amino acids in abundance for all analysed species and conditions.

#p<0.05, ##p<0.01, ###p<0.001, *S. cerevisiae* vs *C. jadinii*, same substrate. *p<0.05, **p<0.01,

***p<0.001, same yeast species, ethanol- (EtOH) vs glucose-grown (Gluc), n=2.

3. Maximum biomass yield and maintenance-energy requirement of *C. jadinii* in ethanol-

limited chemostat cultures

To analyse how specific growth rate affects biomass yield and to estimate the maintenance

energy requirements of *C. jadinii* CBS 621, ethanol-limited chemostat cultures were grown at

different dilution rates. To define a range of dilution rates, maximum specific growth rates (μ_{max})

of *C. jadinii* were determined in batch cultures grown on synthetic medium with ethanol or

glucose as sole carbon source (Fig 2A). Values of μ_{max} on ethanol and glucose were 0.43 ± 0.01

h^{-1} and 0.57 ± 0.01 h^{-1}, respectively (Fig 2B). Both values are higher than the reported μ_{max} of *S.*

cerevisiae CEN.PK113-7D on synthetic medium with glucose (0.39 h^{-1}) (63).

Ethanol-limited chemostat cultures of *C. jadinii* were grown at dilution rates ranging from 0.025

to 0.30 h^{-1}. Over this range of specific growth rates, biomass yields on ethanol ranged from 0.59

to 0.81 $g_{biomass}\cdot(g_{ethanol})^{-1}$ (Fig 2C, Table S3). Using the Pirt equation (eq. 1), we derived

maximum biomass yields on ethanol and oxygen based on the linear fits shown in Fig 2D-E.

From the same data, maximum biomass yields of 0.83 $g_{biomass}\cdot(g_{ethanol})^{-1}$ and 28 $g_{biomass}\cdot(mol\ O_2)^{-1}$

on oxygen were determined. The maximum yield on ethanol coincided with a theoretical value

of 0.83 - 0.84 $g_{biomass}\cdot(g_{ethanol})^{-1}$, estimated based on the assimilation equation for ethanol by *C.*

jadinii (13, 16). This observation suggests that, at high specific growth rates, growth of *C. jadinii*

CBS 621 is close to being carbon-limited rather than energy-limited. This means that the

amount of ATP generated by respiratory reoxidation of the reduced redox cofactors formed

during assimilation of ethanol almost exactly matches the amount of ATP needed for this

assimilation process.

From Fig 2D-E, we derived maintenance coefficients for ethanol (m_s) and oxygen (m_{O2}) of 0.28

$mmol_{ethanol}\cdot(g_{biomass})^{-1}\cdot h^{-1}$ and 0.76 mmol $O_2\cdot(g_{biomass})^{-1}\cdot h^{-1}$, respectively (Fig 2F). Maintenance

relates to use of energy substrate for processes that are not associated with growth, such as

homeostasis of membrane potential and turnover of cellular components (64) and explains the

decline in biomass yield as the dilution rate approaches zero (Fig 2C). Assuming an *in vivo* P/O

ratio of 1.8 *for C. jadinii* (16) and growth-rate independent maintenance-energy requirements

((41), Eq. 1), we estimated an ATP requirement for maintenance (m_{ATP}) of 2.7 mmol ATP$\cdot$(g

biomass)$^{-1}\cdot h^{-1}$ (Fig 2F). This value is over 4-fold higher than the estimated m_{ATP} for *S. cerevisiae*

grown aerobically in glucose-limited cultures (0.63 $mmol_{ATP}\cdot(g_{biomass}^{-1})\cdot h^{-1}$) (60).

A previous publication reported that *C. jadinii* shows higher biomass yields at pH 4.0 than at pH

5.0 (65). However, we did not observe a meaningful change in biomass yield when ethanol-

limited cultures were grown at pH 4.0 instead of 5.0, while protein contents of cultures grown at

pH 4.0 were lower (Fig S2A-C). Co-feeding with formic acid, whose metabolism in yeasts is

strictly dissimilatory, can supply additional NADH and thereby increase ATP availability for

assimilation of the carbon source (18, 66, 67). Formic acid co-feeding was conducted in cultures

grown at pH 5.0 (no expected toxicity (68)). This only marginally increased the biomass yield

and protein content in ethanol-limited cultures, resulting in a protein yield on ethanol of 0.43 g

$_{protein}\cdot(g_{ethanol})^{-1}$ (Fig S2D-F). The effect of formic acid co-feeding was more pronounced in

glucose-grown chemostat cultures, in which protein yields on glucose increased from 0.25 to

0.33 $g_{protein}\cdot(g_{glucose})^{-1}$ (Fig S2F-H).

4. Impact of specific growth rate on protein content and alcohol dehydrogenase levels in

ethanol-limited chemostat cultures

In aerobic, glucose-limited chemostat cultures of yeasts, the protein content of the biomass

typically increases with increasing specific growth rate (40, 65, 69, 70). In contrast, aerobic

ethanol-limited chemostat cultures of *C. jadinii* CBS 621 showed slightly lower protein contents

with increasing growth rates (Fig 2G). Similarly, while RNA content was reported to positively

correlate with specific growth rate in *S. cerevisiae* (40, 71), ethanol-limited chemostat cultures of

C. jadinii CBS 621 showed essentially stable RNA contents across a range of dilution rates (Fig

2G). In contrast to protein and RNA contents, contents of carbohydrates and lipids in biomass

increased with specific growth rate, representing 26 and 11%, respectively, of the biomass at D

= 0.30 h^{-1} (Fig. 2H). Yields of all four macromolecular biomass components on ethanol showed

a growth-rate dependent increase (Fig S2G-H), reflecting the trend of total biomass yields (Fig

2C).

Many studies report that the fraction of microbial proteomes allocated to the translation

machinery increases with increasing specific growth rate (72-74). To investigate whether

ethanol-grown *C. jadinii* exhibited a similar correlation, proteomics analyses were performed on

samples from cultures grown at 0.025, 0.10, 0.21 and 0.30 h^{-1}. The resulting proteome dataset

(Supp. Material S2) was manually curated into 7 categories based on the Gene Ontology

annotation obtained from UniProt (75). Levels of 357 proteins correlated well with specific

growth rate (R^2 > 0.90; 273 positive and 84 negative correlations; Supp. Material S2). The most

prominent effect was a negative correlation of the abundance of proteins involved in ethanol

catabolism (*Cj*Adh, *Cj*Ald and *Cj*Acs) with growth rate. These proteins occupied approximately

27% of the proteome at 0.025 h^{-1} and 12% at 0.3 h^{-1}. In contrast, the fraction of proteins

involved in oxidative phosphorylation (OXPHOS) increased from 4.8 to 7.4% (Fig. 2I), which

positively correlated with oxygen consumption rates (Fig S3A). These abundances are similar to

the proteome contribution of OXPHOS proteins of approximately 5% in *S. cerevisiae* and

Issatchenkia orientalis (*Pichia kudriavzevii*) (72, 76). The fractions of TCA-cycle enzymes and

proteins involved in amino acid synthesis both increased from approximately 4% at 0.025 h^{-1} to

9% at 0.3 h^{-1}, while the fraction of the proteome allocated to the translation machinery increased

from 22 to 29% (Fig 2I).

Of the proteins involved in ethanol catabolism, *Cj*Adh had the strongest impact on the

composition of the proteome. Total *Cj*Adh levels (3 most abundant isoforms, Fig S3B)

accounted for 17% and 5% of the proteome at 0.025 h^{-1} and 0.3 h^{-1}, respectively (Fig 2J),

followed by *Cj*Ald and *Cj*Acs (Fig S3C-D). Given the almost linear correlation between *Cj*Adh

levels and dilution rate, we estimated *Cj*Adh's specific maximum activity (v_{max}). When these

estimates were plotted against measured biomass-specific ethanol consumption rates (which

should equal the *in vivo* flux through *Cj*Adh), the curves intercepted at D = 0.39 h^{-1} (Fig 2K),

which is very close to the maximum specific growth rate of *C. jadinii* on ethanol (0.43 h^{-1}). This

comparison suggests that the maximum specific growth rate of *C. jadinii* CBS 621 on ethanol

may be constrained at the proteome by the level of alcohol dehydrogenase.

Figure 2: Growth rates and physiological characterization of *C. jadinii* in batch and ethanol-limited chemostat cultures at varying dilution rates. (A) Optical density at 660 nm (OD$_{660}$) as a function of time and (B) maximum growth rates (μ$_{max}$) of glucose- and ethanol-grown batch cultures. (C) Biomass yield on ethanol, (D) biomass-specific ethanol uptake rate (q$_{ethanol}$) and (E) biomass-specific oxygen

uptake rates (q_{O_2}) in aerobic, steady-state, ethanol-limited chemostat cultures of *C. jadinii* CBS 621. (F)

Maintenance requirements for ethanol ($m_{ethanol}$) and oxygen (m_{O2}) calculated as the y-intercept of the fitted

curves shown in D and E, respectively. Maintenance ATP requirements (m_{ATP}) were estimated from $m_{O2,}$

assuming an *in vivo* P/O ratio of 1.8 (16). (G) and (H) Quantification of the main components of the

biomass in % of dry weight: proteins (Prot), RNA, carbohydrates (Carb) and lipids. (I) Protein allocation in

ethanol-limited chemostat cultures grown at different dilution rates. Categories were manually curated

and represent (in ascending order) Ethanol catabolism (first three steps shown in Fig 1F), Carbohydrate

metabolism (Carb.), Oxidative Phosphorylation (OXPHOS), TCA cycle (including the glyoxylate shunt

reactions), Amino acid metabolism, Translation and Others. The last category encompasses proteins from

pathways with a lower proteome allocation or for which no pathway was assigned. (J) Total alcohol

dehydrogenase protein (*Cj*Adh, % of proteome) as a function of dilution rate. (K) Estimated v_{max} for

*Cj*Adh, assuming a k_{cat} of 150 min^{-1} for all isoforms (Brenda database (77), EC 1.1.1.1, average reported

value for yeasts) and a proteome comprising 50% of the dry weight (w/w), plotted together with the

ethanol consumption rates ($q_{ethanol}$) from bioreactor experiments. 0.39 h^{-1} represents the intersection

between both lines. Open symbols show carbon recovery of around 107%, while closed symbols

represent carbon recoveries of 100±5%. **p<0.001, n=2.

5. *C. jadinii* CBS 621 is a thiamine auxotroph

The chemically defined medium used in the experiments described above contained seven B-

type vitamins that are commonly included in defined media for yeast cultivation (78). However,

although vitamin-independent growth of other *C. jadinii* strains has been reported (23), the

vitamin dependence of strain CBS 621 has not been investigated. To assess vitamin

requirements of this strain, it was grown in shake-flask cultures on 'dropout' synthetic media

from which a single vitamin was omitted. In addition, growth was measured on a synthetic

medium from which all 7 vitamins were omitted (Δvitamins). In these experiments, only the

Δthiamine and Δvitamins media showed severely compromised growth (Fig 3A-B). When, after

24 h incubation in vitamin-free medium with ethanol, culture samples were used to inoculate

new flasks, the growth deficiency after 24 h further incubation in vitamin-free medium was even

more pronounced (Fig 3B). Addition of thiamine sufficed to restore growth (Fig 3C).

To investigate the apparent thiamine auxotrophy *C. jadinii* CBS 621, its genome was sequenced

and *de novo* assembled. The estimated genome size of 17.9 Mb (Table S4) is larger than that of

other *C. jadinii* strains (21). This may be due to its high degree of heterozygosity, which

generated alternative alleles for heterozygous regions during assembly. Based on the

distribution of SNPs in the heterozygous regions across the genome, CBS 621 is most probably

triploid (Fig S4A). A search for genes encoding proteins in thiamine biosynthesis (*CjTHI4*,

CjTHI5, *CjTHI6* and *CjTHI20*) (78) was performed in UniProt, using *C. jadinii* as a taxonomy

identifier. The annotated amino acid sequences for the type strain CBS 1600 were then used to

identify homologs in strain CBS 621 with the BLAST tool (Fig 3D). This homology search

yielded matches in strain CBS 621 with *CjTHI5*, *CjTHI6* and *CjTHI20*, but not with *CjTHI4*, which

encodes thiamine thiazole synthase, a single-turnover 'suicide enzyme' (78) involved in

thiamine biosynthesis. Based on these results, we concluded that loss of the *CjTHI4* gene

rendered *C. jadinii* CBS 621 a thiamine auxotroph.

Figure 3: Growth of *C. jadinii* CBS 621 on single-vitamin dropout and vitamin-free media. (A)

Growth curves of *C. jadinii* CBS 621 on single-vitamin dropout media in shake flasks grown on synthetic

medium with ethanol as a sole carbon source. Δ"vitamin name" indicates absence of only the indicated

vitamin from the standard vitamin mix. *Complete* indicates presence of all vitamins and Δ*vitamins*

indicates absence of all vitamins. (B) OD_{660} after 24h of cultivation in shake flasks. (C) OD_{660} after further

24h of cultivation in shake flasks for samples from indicated groups (x-axis) that were reinoculated at OD

0.5 in the presence or absence of thiamine, n=2. (D) Overview of tBLASTn search to identify genes

coding for enzymes in the biosynthesis of thiamine in CBS 621. The reference GenBank IDs used as

queries were retrieved from UniProt, based on searches containing the respective protein name and

species (*C. jadinii* CBS 1600). The tBLASTn algorithm was run locally using default settings.

6. Cultivation of prototrophic *C. jadinii* strains on ethanol on a vitamin-free medium

To evaluate the physiology of potentially prototrophic *C. jadinii* strains on ethanol, twelve

additional strains were selected, excluding those isolated from animal material unless a genome

sequence was available. A serial-dilution based method (Fig 4A-B) was used to calculate their

specific growth rates in microtiter plates (39). Of the tested *C. jadinii* strains, only CBS 621 and

CBS 7232 failed to grow on vitamin-free medium (Fig 4C). Growth of these strains was rescued

by adding of thiamine and biotin, respectively (Fig S5A). The maximum specific growth rates of

the strains ranged from 0.13 - 0.44 h^{-1} (Fig 4C). Most of the variation in the dataset could be

attributed to strain differences (p<0.001). Vitamin absence also significantly impacted the

specific growth rates, leading to an average 10-15% lower growth rate (p<0.001) (Fig 4C). The

final Green Value of the growth curves (Fig S5B) showed less variation among samples.

Three *C. jadinii* strains were selected for further investigation: CBS 5609 (highest $μ_{max}$), CBS

1600 (largest difference in $μ_{max}$ caused by vitamin absence) and CBS 5947 (among the top

three strains with the highest final Green Value). Maximum specific growth rates of these three

strains were measured in shake flasks to validate the serial dilution method. Although the effect

of vitamin absence on CBS 1600 was less pronounced (Fig 4D), values obtained with the two

methods corresponded well (Fig 4E, S5C). Publicly available genome sequences from strains

CBS 1600 and CBS 5609 (24, 79), indicated the presence of a complete *THI4* gene. Presence

of *THI4* in strain CBS 5947 was confirmed by whole-genome sequencing (Table S4). Like

strains CBS 5609 and CBS 621, *C. jadinii* CBS 5947 appeared to be triploid based on its SNP

distribution (Fig S4B).

To investigate biomass yields and protein contents of the three selected strains during growth

on ethanol in vitamin-free medium, they were grown in ethanol-limited chemostats at 0.10 h^{-1} in

the absence of supplemented vitamins (Table S5). *C. jadinii* CBS 5609, the strain with the

highest μ_{max} on ethanol (0.46 h^{-1}, Fig 4D) showed the lowest biomass yield on ethanol and

oxygen (Fig 4F-G). Strains CBS 1600 and CBS 5947 showed nearly the same biomass yields

on ethanol (0.72 and 0.74 $g_{biomass} \cdot (g_{ethanol})^{-1}$, respectively (Fig 4F-H). All three strains showed a

protein content of approximately 54% (Fig 4H). Vitamin-supplemented ethanol-limited cultures

of strains CBS 1600 and CBS 621 showed the same biomass yield on ethanol and protein

content (Fig 4F-H). For strain CBS 1600, these results indicate that omitting vitamins from

synthetic media does not affect biomass yield and protein content during ethanol-limited growth

at 0.10 h^{-1}.

Figure 4: Characterization of *C. jadinii* strains in microtiter plates, shake flasks and bioreactors concerning vitamin dependency. (A) Representative growth curves of serially diluted CBS 621 in microtiter plates, ranging from 1 to 32× dilution, initial OD_{660} 0.2 to 0.00625, respectively. (B) Calculation of specific growth rates via serial dilution method. The time (h) required for a specific dilution to reach a green value of 20 was plotted against the natural logarithm of the respective dilution (1-32). (C) Specific growth rates for 13 *C. jadinii* strains grown on ethanol in the presence or absence of vitamins, calculated with the serial dilution method, n.d: non-detectable. (D) Specific growth rates of 4 *C. jadinii* strains grown on ethanol in shake flasks in the presence or absence of vitamins. (E) Correlation of specific growth rates obtained in microtiter plates (x-axis) with those measured in shake flasks (y-axis) for strains grown in the presence of vitamins. (F-H) Biomass yields on ethanol, oxygen and protein content for selected strains grown in ethanol-limited chemostats (dilution rate 0.10 h^{-1}). Where indicated by Δvit, no vitamins were added to the cultivation, n=2-3.

7. High-cell-density fed-batch cultivation of *C. jadinii* on ethanol in a mineral, vitamin-free medium.

Of the vitamin-prototrophic *C. jadinii* strains, CBS 5947 showed the highest biomass yield in ethanol-limited chemostat cultures (0.74 $g_{biomass} \cdot (g_{ethanol})^{-1}$, Fig 4F). To evaluate whether this strain could be grown to high biomass densities on ethanol in a vitamin-free medium, a fed-batch protocol was developed. After an initial batch phase, a biomass concentration of 5.5 $g_{biomass} \cdot (kg_{broth})^{-1}$ was reached in two independent reactors. For the subsequent fed-batch regime, pure ethanol was continuously fed and 25% NH_4OH was used as pH titrant and nitrogen source. To increase oxygen solubility, 1 bar overpressure was applied since the beginning of the batch phase (Fig 5A). The initial specific growth rate was set at 0.10 h^{-1} by applying a preprogrammed exponentially increasing ethanol feed rate (Fig 5B). When dissolved oxygen concentration in the reactor decreased to a concentration corresponding to 15% of air saturation at atmospheric pressure (Fig S6A), the ethanol feed rate was first fixed at 0.047 $kg_{ethanol} \cdot h^{-1}$ and, after 5 additional hours, at 0.035 $kg_{ethanol} \cdot h^{-1}$ (Fig 5B) to prevent oxygen limitation. In total, approximately 1.4 kg of ethanol was consumed in each reactor within 48 h (Fig 5C), leading to a biomass concentration of 98 ± 4 $g_{biomass} \cdot (kg_{broth})^{-1}$ (Fig 5D). Consistent with the cultures being ethanol-limited, the ethanol concentration in culture samples remained below detection limit throughout the fed-batch phase. From the time-dependent increase of the amount of biomass in the reactor, we calculated an approximate specific growth rate of 0.02 h^{-1} at the end of the fed-batch phase (Fig 5E). A decline in growth rate was also evident from the slope of the CO_2 profile during fermentation (Fig S6B). The fed-batch process was simulated (37) based on the parameters calculated from Fig 2D-F and the feed profile shown in Fig 5B. Experimentally determined specific growth rates and ethanol consumption were consistent with model predictions (Fig 5C-E).

The protein content remained at 50% during the exponential feed phase of the fed-batch culture

(Fig 5F). However, in contrast to observations in steady-state chemostat cultures (Fig 2) in

which protein concentrations did not decline at low specific growth rates, a decrease of the

protein content to 42% was observed towards the end of the fed-batch cultures (Fig 5F). This

indicates that the protein content decreased when the growth rate was dynamically changed

from 0.10 to 0.022 h^{-1}. In the two fed-batch experiments, an average of 405 g protein was

produced (Fig 5G). Based on the total ethanol consumed and total biomass and protein

produced during the fed-batch phase, biomass, and protein yields of 0.65 $g_{biomass} \cdot (g_{ethanol})^{-1}$ and

0.28 $g_{protein} \cdot (g_{ethanol})^{-1}$, respectively, were calculated (Fig 5H). Production of low concentrations of

acetate and succinate in the fed-batch processes (~10 mM final concentrations) (Fig S6C)

played a negligible role in the total carbon recovery (Fig S6D-E).

In the fed-batch cultures, strain CBS 5947 needed to synthesize vitamins to sustain growth.

Levels of thiamine, riboflavin, niacin, and pyridoxine in fed-batch grown *C. jadinii* CBS 5947

ranged from 1.7 – 260 mg $\cdot (kg_{biomass})^{-1}$ (Table S6) . When compared with the vitamin contents in

aquaculture diets (80), ethanol-grown *C. jadinii* biomass showed higher levels of niacin,

riboflavin and pyridoxine, while thiamine content of the yeast biomass was approximately 3-fold

lower (Table S6).

Figure 5: Growth of *C. jadinii* CBS 5947 on vitamin-free medium in aerobic, ethanol-limited fed-batch cultures. Aerobic fed-batch fermentation was conducted at 30 °C, at pH 5.0 and at 1 bar overpressure, with a feed of pure ethanol. A 25% (w/v) NH_4OH solution was used for simultaneous pH control and nitrogen supplementation. (A) Scheme of reactor and critical parameters related to the process (fed-batch phase). W_0: initial weight of the culture broth (kg), $c_{x,0}$: initial biomass concentration ($kg_{biomass} \cdot (kg_{broth})^{-1}$), μ_0: initial specific growth rate in fed-batch phase (h^{-1}), DW = biomass dry weight concentration (g biomass$\cdot(kg_{broth})^{-1}$). Figure created in BioRender. Vieira-Lara, M. (2024) BioRender.com/z76n618. (B) Profile of feed rate during the fed-batch phase. (C-E) Consumed ethanol, biomass dry weight, and growth rate (μ) during the fed-batch phase. Symbols represent data and grey lines refer to simulations. (F-G) protein percentage in the biomass and total protein content during the

fed-batch phase. (H) Total biomass and protein yield on ethanol based on the whole fed-batch process.

Data are shown for 2 independent reactors (mean ± SD).

Discussion

Fast growth of industrial microorganisms is especially beneficial during batch cultivation and

during the initial exponential growth phase of aerobic fed-batch fermentation processes, when

oxygen transfer is not yet limiting. In a recent screening of 52 ascomycete budding yeasts, we

noted that *C. jadinii* CBS 621 was the fastest-growing strains on a synthetic medium containing

ethanol as sole carbon source (13). In the present study, multiple *C. jadinii* strains showed

specific growth rates on ethanol close to 0.4 h^{-1} (doubling time 1.7 h) in batch cultures grown on

a synthetic medium supplemented with a standard vitamin mixture (Fig 2B, 4C-D). With two

exceptions, the investigated *C. jadinii* strains were prototrophic and showed maximum specific

growth rates on ethanol in vitamin-free media that were only 10-15% lower than in vitamin-

supplemented medium. Use of vitamin-free media offers advantages in terms of cost and

simplicity of medium preparation and sterilization and, potentially, reduction of contamination

risks (78). Moreover, use of a naturally prototrophic yeast strain precludes the need for

extensive metabolic engineering strategies, such as recently applied to *S. cerevisiae* (62), to

eliminate full vitamin auxotrophies. The thiamine auxotrophy of our initial model strain *C. jadinii*

CBS 621 (Fig 3A-C) could be attributed to absence of the *CjTHI4* gene. Thiamine auxotrophy of

this strain may enable use of *CjTHI4* as a selectable marker in ongoing research to make *C.*

jadinii fully accessible to Cas9-based genome editing (81).

Biomass yields of *C. jadinii* CBS 621 in ethanol-limited chemostat cultures with vitamin

supplementation, and of two prototrophic *C. jadinii* strains in vitamin-free media, were among

the highest reported for yeasts (13) (Fig 4F). The maximum biomass yield ($Y_{x/s}^{max}$) of *C. jadinii*

CBS 621 on ethanol of 0.83 $g_{biomass} \cdot (g_{ethanol})^{-1}$ in vitamin-supplemented, ethanol-limited

chemostat cultures even matched the theoretical maximum yield on ethanol of 0.83 - 0.84

$g_{biomass} \cdot (g_{ethanol})^{-1}$ (13, 16) for ethanol-limited growth with ammonium as the nitrogen source. A

previous study on strain CBS 621 that was based on measurements at a single dilution rate

(0.10 h^{-1}) reported lower biomass yields than observed in our study (18). A similarly high

biomass yield on ethanol (0.82 $g \cdot g^{-1}$) was reported for batch cultures of the related species

Cyberlindnera saturnus (12). We consistently measured high biomass yields in replicate

cultures of different *C. jadinii* strains, both on vitamin-supplemented and vitamin-free media,

checked carbon recoveries of the chemostat cultures and confirmed that no biomass retention

occurred (82) and therefore trust our reported data.

Ethanol-limited chemostat cultures of *C. jadinii* showed a 25% higher biomass yield on ethanol

than cultures of *S. cerevisiae* grown under the same set of conditions (Fig. 1). This difference

can be attributed to the presence and absence, respectively, of genes encoding a proton-

pumping Complex I NADH dehydrogenase in these two yeasts (13). In contrast to the difference

observed in ethanol-limited cultures (16, 18, 58), biomass yields of *C. jadinii* and *S. cerevisiae* in

aerobic glucose-limited cultures differed by only 6% (Fig 1). This observation suggests that the

contribution of Complex I to respiratory reoxidation of NADH by *C. jadinii* may be carbon-source

dependent. A deeper understanding of the role and regulation of Complex I in *C. jadinii* may

contribute to improving yields of biomass and other ATP-intensive products, also on carbon-

and energy sources other than ethanol. Our current lack of understanding on this subject is

illustrated by the observation that, counterintuitively, levels of the non-proton-pumping single-

subunit NADH dehydrogenase *Cj*Ndi1 were approximately 150-fold higher in the proteome of

ethanol-limited cultures of *C. jadinii* than in glucose-limited cultures (Fig S1A, Supp Material S1).

Further research should also address the question of how subcellular localization of key

enzymes and/or redox shuttle mechanisms enable reoxidation of NADH generated by ethanol

and acetaldehyde dehydrogenases, enzymes that can occur in the cytosol as well as in the

mitochondrial matrix of yeasts (61), by Complex I, whose catalytic site is in the mitochondrial

matrix.

The estimated growth-rate independent maintenance energy requirement for ATP (m_{ATP}) of

ethanol-limited cultures of *C. jadinii* (2.7 mmol ATP$\cdot$($g_{biomass}^{-1}$)$\cdot h^{-1}$, Fig 2F) was higher than the

m_{ATP} of 1.2 mmol ATP$\cdot$($g_{biomass}$)$^{-1}\cdot h^{-1}$ estimated from a study on aerobic, glucose-limited

chemostat cultures of the Complex-I containing yeast *Komagataella phaffii*, grown at dilution

rates ranging from 0.025 to 0.10 h^{-1} (83). In energy-substrate-limited cultures, a high

maintenance energy requirement most strongly affects biomass yields at low specific growth

rates, where it needs a large fraction of the substrate to be respired instead of converted to

biomass. Consequently, and consistent with model-based simulations that included the high

maintenance requirement of *C. jadinii*, high-cell-density fed-batch cultures did not attain the

biomass yields reached in ethanol-limited chemostat cultures grown at intermediate rates (Fig

3C, 4F, 5H). Protein turnover has been reported to account for a large share of the ATP

requirement for cellular maintenance in *S. cerevisiae* (84). Interestingly, two heterologous-

protein producing *K. phaffii* strains were reported to exhibit 2-fold higher maintenance-energy

requirements than a reference strain (83, 85). This raises the question of whether the high m_{ATP}

of *C. jadinii* may be related to its high protein content, and particularly the remarkably high levels

of specific proteins involved in ethanol metabolism.

Under substrate-limited conditions, competitiveness of microorganisms is determined by their

affinity for the growth-limiting substrate, i.e. the ability to sustain a high biomass-specific

consumption rate (q_s) at a low concentration (c_s) of that substrate. Based on the Monod

equation ($q_s = q_{s,max} (c_s/(c_s + K_s))$), affinity can be defined as $q_{s,max}/K_S$ (86), in which $q_{s,max}$ is the

maximum biomass-specific substrate uptake rate, and K_s is the value of c_s at which $q_s =$

$\frac{1}{2}\cdot q_{s,max}$. At low specific growth rates, a large fraction of the proteome of ethanol-limited cultures

of *C. jadinii* was allocated to enzymes that catalyse the first three reactions of ethanol

metabolism (Fig 2I) and, in particular, to alcohol dehydrogenase (Fig 2J). This observation

indicates that, at low specific growth rates, *C. jadinii* increases its affinity for ethanol by

increasing $q_{s,max}$. This adaptation strategy resembles regulation of methanol oxidase, the first

enzyme in methanol metabolism, in *Ogataea polymorpha*. In slow-growing (0.03 h^{-1}) methanol-

limited cultures of this methylotrophic yeast, 20% of its proteome consisted of methanol oxidase

(87), which closely resembles the 17% contribution of alcohol-dehydrogenases in ethanol-

limited *C. jadinii* cultures grown at 0.025 h^{-1}. We hypothesize that this '$q_{s,max}$ adaptation' to

ethanol-limited growth contributes to the observation that, despite lower rates of assimilation

and dissimilation at low specific growth rates, *C. jadinii* exhibited an almost growth-rate

independent protein content of ca. 54% in ethanol-limited chemostat cultures.

Based on the quantitative data obtained from chemostat experiments, we designed a feed

profile for high-cell density ethanol-limited fed-batch cultures. A previous study also described

based fed-batch cultivation of *C. jadinii* on ethanol, but applied discontinuous ethanol feeding

and reached a maximum biomass concentration of 15 $g_{biomass} \cdot (kg_{broth})^{-1}$ (88). By controlled

feeding of pure ethanol, which eliminated the need for sterilisation of the feed line, the fed-batch

cultures reached 100 $g_{biomass} \cdot (kg_{broth})^{-1}$ in 48 h of cultivation (Fig 5D). Experimental data on

biomass formation were in excellent agreement with model predictions. However, the protein

content of biomass harvested at the end of high-cell density fed-batch cultures, when the

specific growth rate had decreased to 0.02 h^{-1}, had decreased to only 42%. Further research is

needed to assess whether this marked difference with the protein content in slow-growing

chemostat cultures is due to the dynamic growth-rate profile in the fed-batch cultures, imperfect

mixing of the pure ethanol feed, or other factors.

Conclusions

Based on different analyses on ethanol-grown batch, chemostat and fed-batch cultures, the

present study provides a quantitative physiology baseline for fundamental and applied research

on *C. jadinii*. Its high maximum growth rate, high energetic efficiency of ethanol dissimilation,

simple nutritional requirements and high protein content, make this yeast a highly interesting

platform for production of SCP and other products from ethanol produced by low-emission

technologies. Especially for SCP applications, an observed difference in protein content

between ethanol chemostat and fed-batch cultures that were grown over the same range of

specific growth rates, merits further investigation. From a fundamental science perspective, our

results identify questions about regulation, energy coupling and compartmentation of NADH

reoxidation during growth of yeasts and fungi during growth on non-conventional substrates. In

addition, the results underline the key role of maintenance energy requirements in high-cell-

density fed-batch processes.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All data generated or analysed during this study are included in this published article and its

supplementary information files.

Competing interests

The company dsm-firmenich owns intellectual property rights of technology discussed in this

paper. MAVL and JTP are co-inventors on patent applications related to the present work.

Funding

Co-funding was obtained from dsm-firmenich and from a supplementary grant 'TKI- Toeslag' for

Topconsortia for Knowledge and Innovation (TKI's) of the Netherlands Ministry of Economic

Affairs and Climate Policy (CHEMIE.PGT.2021.003). The PhD project of MW is funded from a

Stevin Prize awarded to JTP by the Dutch Research Council (NWO).

Authors' contributions

MAVL: investigation, methodology, analysis, data visualization, writing (original draft, reviewing

and editing). MW: investigation, methodology, analysis, writing (reviewing and editing). EdH and

MvdB: investigation, methodology. JMD: conceptualization, writing (reviewing and editing). JTP:

conceptualization, supervision, writing (reviewing and editing).

Acknowledgements

The authors thank Clara Carqueija Cardoso for the help with DNA sequencing, Christiaan

Mooiman for assistance with the fed-batch experiments, Marijke Luttik for the help with growth

experiments, Chris Klomp and Niels Tiemersma for evaluating the methods for biomass

composition and help with bioreactor cultivation, and Jordy Elfering for the testing of the serial

dilution method and screening and for help with bioreactor cultivation. We thank André Vente for

the amino acid and proteome analyses, and Mickel Jansen and Tjeerd van Rij for the input in

the project.

References

1.   Claassens NJ, Cotton CAR, Kopljar D, Bar-Even A. Making quantitative sense of

electromicrobial production. Nature Catalysis. 2019;2(5):437-47.

2.   Leger D, Matassa S, Noor E, Shepon A, Milo R, Bar-Even A. Photovoltaic-driven

microbial protein production can use land and sunlight more efficiently than conventional crops.

Proc Natl Acad Sci U S A. 2021;118(26).

3.   Lynd LR, Beckham GT, Guss AM, Jayakody LN, Karp EM, Maranas C, et al. Toward

low-cost biological and hybrid biological/catalytic conversion of cellulosic biomass to fuels.

Energy & Environmental Science. 2022;15(3):938-90.

4.   Martin ME, Richter H, Saha S, Angenent LT. Traits of selected Clostridium strains for

syngas fermentation to ethanol. Biotechnol Bioeng. 2016;113(3):531-9.

5.   Smith WA, Burdyny T, Vermaas DA, Geerlings H. Pathways to Industrial-Scale Fuel Out

of Thin Air from CO2 Electrolysis. Joule. 2019;3(8):1822-34.

6.   Christodoulou X, Okoroafor T, Parry S, Velasquez-Orta SB. The use of carbon dioxide in

microbial electrosynthesis: Advancements, sustainability and economic feasibility. Journal of

CO2 Utilization. 2017;18:390-9.

7.   Cabau-Peinado O, Winkelhorst M, Stroek R, de Kat Angelino R, Straathof AJJ, Masania

K, et al. Microbial electrosynthesis from CO(2) reaches productivity of syngas and chain

elongation fermentations. Trends Biotechnol. 2024.

8.   Abdinejad M, Farzi A, Möller-Gulland R, Mulder F, Liu C, Shao J, et al. Eliminating

redox-mediated electron transfer mechanisms on a supported molecular catalyst enables CO2

conversion to ethanol. Nature Catalysis. 2024.

9.   Jansen MLA, Bracher JM, Papapetridis I, Verhoeven MD, de Bruijn H, de Waal PP, et al.

Saccharomyces cerevisiae strains for second-generation ethanol production: from academic

exploration to industrial implementation. FEMS Yeast Res. 2017;17(5).

10.   Takors R, Kopf M, Mampel J, Bluemke W, Blombach B, Eikmanns B, et al. Using gas

mixtures of CO, CO(2) and H(2) as microbial substrates: the do's and don'ts of successful

technology transfer from laboratory to production scale. Microb Biotechnol. 2018;11(4):606-25.

11.   Ingelman H, Heffernan JK, Harris A, Brown SD, Shaikh KM, Saqib AY, et al. Autotrophic

adaptive laboratory evolution of the acetogen Clostridium autoethanogenum delivers the gas-

fermenting strain LAbrini with superior growth, products, and robustness. N Biotechnol.

2024;83:1-15.

12.   Van Peteghem L, Sakarika M, Matassa S, Rabaey K. The Role of Microorganisms and

Carbon-to-Nitrogen Ratios for Microbial Protein Production from Bioethanol. Appl Environ

Microbiol. 2022;88(22):e0118822.

13.   Warmerdam M, Vieira-Lara MA, Mans R, Daran JM, Pronk JT. Specific growth rates and

growth stoichiometries of Saccharomycotina yeasts on ethanol as sole carbon and energy

substrate. bioRxiv (preprint). 2024.

14.   Kotyk A, Alonso A. Transport of ethanol in baker's yeast. Folia Microbiol (Praha).

1985;30(1):90-1.

15.   Kozak BU, van Rossum HM, Niemeijer MS, van Dijk M, Benjamin K, Wu L, et al.

Replacement of the initial steps of ethanol metabolism in Saccharomyces cerevisiae by ATP-

independent acetylating acetaldehyde dehydrogenase. FEMS Yeast Res. 2016;16(2):fow006.

16.   Verduyn C, Stouthamer AH, Scheffers WA, van Dijken JP. A theoretical evaluation of

growth yields of yeasts. Antonie Van Leeuwenhoek. 1991;59(1):49-63.

17.   Juergens H, Hakkaart XDV, Bras JE, Vente A, Wu L, Benjamin KR, et al. Contribution of

Complex I NADH Dehydrogenase to Respiratory Energy Coupling in Glucose-Grown Cultures of

Ogataea parapolymorpha. Appl Environ Microbiol. 2020;86(15).

18.   Verduyn C. Physiology of yeasts in relation to biomass yields. Antonie Van

Leeuwenhoek. 1991;60(3-4):325-53.

19.   Joseph-Horne T, Hollomon DW, Wood PM. Fungal respiration: a fusion of standard and

alternative components. Biochim Biophys Acta. 2001;1504(2-3):179-95.

20.   Nosek J, Fukuhara H. NADH dehydrogenase subunit genes in the mitochondrial DNA of

yeasts. J Bacteriol. 1994;176(18):5622-30.

21.   Sousa-Silva M, Vieira D, Soares P, Casal M, Soares-Silva I. Expanding the Knowledge

on the Skillful Yeast Cyberlindnera jadinii. J Fungi (Basel). 2021;7(1).

22.   Averet N, Jobin ML, Devin A, Rigoulet M. Proton pumping complex I increases growth

yield in Candida utilis. Biochim Biophys Acta. 2015;1847(10):1320-6.

23.   Burkholder PR, McVeigh I, Moyer D. Studies on some growth factors of yeasts. Journal

of Bacteriology. 1944;48(4):385-91.

24.   Rupp O, Brinkrolf K, Buerth C, Kunigo M, Schneider J, Jaenicke S, et al. The structure of

the Cyberlindnera jadinii genome and its relation to Candida utilis analyzed by the occurrence of

single nucleotide polymorphisms. J Biotechnol. 2015;211:20-30.

25.   Lallemand. Exploring Lallemand's Torula Yeast: A Remarkable Journey [Available from:

https://bio-lallemand.com/savory-ingredients/exploring-lallemands-torula-yeast-specialties/.

26.   Solomons G, Litchfield JH. Single cell protein. Critical Reviews in Biotechnology.

1983;1(1):21-58.

27.   Agboola JO, Lapeña D, Øverland M, Arntzen MØ, Mydland LT, Hansen JØ. Yeast as a

novel protein source - Effect of species and autolysis on protein and amino acid digestibility in

Atlantic salmon (Salmo salar). Aquaculture. 2022;546.

28.   Cruz A, Sterten H, Steinhoff FS, Mydland LT, Overland M. Cyberlindnera jadinii yeast as

a protein source for broiler chickens: effects on growth performance and digestive function from

hatching to 30 days of age. Poult Sci. 2020;99(6):3168-78.

29.   Cruz A, Tauson A-H, Matthiesen CF, Mydland LT, Øverland M. Cyberlindnera jadinii

yeast as a protein source for growing pigs: Effects on protein and energy metabolism. Livestock

Science. 2020b;231.

30.   Matassa S, Boon N, Pikaar I, Verstraete W. Microbial protein: future sustainable food

supply route with low environmental footprint. Microb Biotechnol. 2016;9(5):568-75.

31.   Agboola JO, Øverland M, Skrede A, Hansen JØ. Yeast as major protein-rich ingredient

in aquafeeds: a review of the implications for aquaculture production. Reviews in Aquaculture.

2020;13(2):949-70.

32.   Jones SW, Karpol A, Friedman S, Maru BT, Tracy BP. Recent advances in single cell

protein use as a feed ingredient in aquaculture. Curr Opin Biotechnol. 2020;61:189-97.

33. Nyyssola A, Suhonen A, Ritala A, Oksman-Caldentey KM. The role of single cell protein

in cellular agriculture. Curr Opin Biotechnol. 2022;75:102686.

34. Ritala A, Hakkinen ST, Toivari M, Wiebe MG. Single Cell Protein-State-of-the-Art,

Industrial Landscape and Patents 2001-2016. Front Microbiol. 2017;8:2009.

35. Spalvins. Improving single cell protein yields and amino acid profile via mutagenesis:

review of applicable amino acid inhibitors for mutant selection. 2021.

36. Choi KR, Yu HE, Lee SY. Microbial food: microorganisms repurposed for our food.

Microb Biotechnol. 2022;15(1):18-25.

37. Hensing MCM, Vrouwenvelder JS, Hellinga C, Van Dijken JP, Pronk JT. Use of

chemostat data for modelling extracellular-inulinase production by Kluyveromyces marxianus in

a high-cell-density fed-batch process. Journal of Fermentation and Bioengineering.

1995;79(1):54-8.

38. Verduyn C, Postma E, Scheffers WA, Van Dijken JP. Effect of benzoic acid on metabolic

fluxes in yeasts: a continuous-culture study on the regulation of respiration and alcoholic

fermentation. Yeast. 1992;8(7):501-17.

39. Hazan R, Que YA, Maura D, Rahme LG. A method for high throughput determination of

viable bacteria cell counts in 96-well plates. BMC Microbiol. 2012;12:259.

40. Lange HC, Heijnen JJ. Statistical reconciliation of the elemental and molecular biomass

composition of Saccharomyces cerevisiae. Biotechnol Bioeng. 2001;75(3):334-44.

41. Pirt SJ. The maintenance energy of bacteria in growing cultures. Proc R Soc Lond B Biol

Sci. 1965;163(991):224-31.

42. van Hoek P, de Hulster E, van Dijken JP, Pronk JT. Fermentative capacity in high-cell-

density fed-batch cultures of baker's yeast. Biotechnol Bioeng. 2000;68(5):517-23.

43. Puiman L, Elisiário MP, Crasborn LML, Wagenaar LECH, Straathof AJJ, Haringa C. Gas

mass transfer in syngas fermentation broths is enhanced by ethanol. Biochemical Engineering

Journal. 2022;185.

44.   Garcia-Ochoa F, Gomez E. Bioreactor scale-up and oxygen transfer rate in microbial

processes: an overview. Biotechnol Adv. 2009;27(2):153-76.

45.   Postma E, Verduyn C, Scheffers WA, Van Dijken JP. Enzymic analysis of the crabtree

effect in glucose-limited chemostat cultures of Saccharomyces cerevisiae. Appl Environ

Microbiol. 1989;55(2):468-77.

46.   Mashego MR, van Gulik WM, Vinke JL, Heijnen JJ. Critical evaluation of sampling

techniques for residual glucose determination in carbon-limited chemostat culture of

Saccharomyces cerevisiae. Biotechnol Bioeng. 2003;83(4):395-9.

47.   Guadalupe Medina V, Almering MJ, van Maris AJ, Pronk JT. Elimination of glycerol

production in anaerobic cultures of a Saccharomyces cerevisiae strain engineered to use acetic

acid as an electron acceptor. Appl Environ Microbiol. 2010;76(1):190-5.

48.   Johnson KR, Ellis G, Toothill C. The sulfophosphovanillin reaction for serum lipids: a

reappraisal. Clin Chem. 1977;23(9):1669-78.

49.   Izard J, Limberger RJ. Rapid screening method for quantitation of bacterial cell lipids

from whole cells. J Microbiol Methods. 2003;55(2):411-8.

50.   Benthin S, Nielsen J, Villadsen J. A simple and reliable method for the determination of

cellular RNA content. Biotechnology Techniques. 1991;5(1):39-42.

51.   Herbert D, Phipps PJ, Strange RE. Chapter III Chemical Analysis of Microbial Cells.

Methods in Microbiology1971. p. 209-344.

52.   Hebert AS, Richards AL, Bailey DJ, Ulbrich A, Coughlin EE, Westphall MS, et al. The

one hour yeast proteome. Mol Cell Proteomics. 2014;13(1):339-47.

53.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009;25(14):1754-60.

54.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.

55.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an

integrated tool for comprehensive microbial variant detection and genome assembly

improvement. PLoS One. 2014;9(11):e112963.

56.   Wickham H, Wickham H. ggplot2: Springer; 2016.

57.   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes De

Novo Assembler. Curr Protoc Bioinformatics. 2020;70(1):e102.

58.   van Urk H, Postma E, Scheffers WA, van Dijken JP. Glucose transport in crabtree-

positive and crabtree-negative yeasts. J Gen Microbiol. 1989;135(9):2399-406.

59.   Bruinenberg PM, van Dijken JP, Scheffers WA. An enzymic analysis of NADPH

production and consumption in Candida utilis. J Gen Microbiol. 1983;129(4):965-71.

60.   Vos T, Hakkaart XD, de Hulster EA, van Maris AJ, Pronk JT, Daran-Lapujade P.

Maintenance-energy requirements and robustness of Saccharomyces cerevisiae at aerobic

near-zero specific growth rates. Microb Cell Fact. 2016;15(1):111.

61.   Bakker BM, Bro C, Kotter P, Luttik MA, van Dijken JP, Pronk JT. The mitochondrial

alcohol dehydrogenase Adh3p is involved in a redox shuttle in Saccharomyces cerevisiae. J

Bacteriol. 2000;182(17):4730-7.

62.   Ehrmann AK, Wronska AK, Perli T, de Hulster EAF, Luttik MAH, van den Broek M, et al.

Engineering Saccharomyces cerevisiae for fast vitamin-independent aerobic growth. Metab

Eng. 2024;82:201-15.

63.   Wronska AK, Haak MP, Geraats E, Bruins Slot E, van den Broek M, Pronk JT, et al.

Exploiting the Diversity of Saccharomycotina Yeasts To Engineer Biotin-Independent Growth of

Saccharomyces cerevisiae. Appl Environ Microbiol. 2020;86(12):e00270-20.

64.   van Bodegom P. Microbial maintenance: a critical review on its quantification. Microb

Ecol. 2007;53(4):513-23.

65.   Paredes-Lopez O, Camargo-Rubio E, Ornelas-Vale A. Influence of specific growth rate

on biomass yield, productivity, and compostion of Candida utilis in batch and continuous culture.

Appl Environ Microbiol. 1976;31(4):487-91.

66.   van Winden WA, Mans R, Breestraat S, Verlinden RAJ, Mielgo-Gomez A, de Hulster

EAF, et al. Towards closed carbon loop fermentations: Cofeeding of Yarrowia lipolytica with

glucose and formic acid. Biotechnol Bioeng. 2022;119(8):2142-51.

67.   Bruinenberg PM, Jonker R, Vandijken JP, Scheffers WA. Utilization of Formate as an

Additional Energy-Source by Glucose-Limited Chemostat Cultures of *Candida utilis* CBS 621

and *Saccharomyces cerevisiae* CBS 8066 - Evidence for the Absence of Transhydrogenase

Activity in Yeasts. Arch Microbiol. 1985;142(3):302-6.

68.   Overkamp KM, Kotter P, van der Hoek R, Schoondermark-Stolk S, Luttik MA, van Dijken

JP, et al. Functional analysis of structural genes for NAD(+)-dependent formate dehydrogenase

in Saccharomyces cerevisiae. Yeast. 2002;19(6):509-20.

69.   Mor JR, Fiechter A. Continuous cultivation of Saccharomyces cerevisiae. I. Growth on

ethanol under steady-state conditions. Biotechnology and Bioengineering. 1968;10(2):159-76.

70.   Fonseca GG, Gombert AK, Heinzle E, Wittmann C. Physiology of the yeast

Kluyveromyces marxianus during batch and chemostat cultures with glucose as the sole carbon

source. FEMS Yeast Res. 2007;7(3):422-35.

71.   Waldron C, Lacroute F. Effect of growth rate on the amounts of ribosomal and transfer

ribonucleic acids in yeast. J Bacteriol. 1975;122(3):855-65.

72.   Elsemman IE, Rodriguez Prado A, Grigaitis P, Garcia Albornoz M, Harman V, Holman

SW, et al. Whole-cell modeling in yeast predicts compartment-specific proteome constraints that

drive metabolic strategies. Nat Commun. 2022;13(1):801.

73.   Bruggeman FJ, Planque R, Molenaar D, Teusink B. Searching for principles of microbial

physiology. FEMS Microbiol Rev. 2020;44(6):821-44.

74.   Klumpp S, Scott M, Pedersen S, Hwa T. Molecular crowding limits translation and cell
growth. Proc Natl Acad Sci U S A. 2013;110(42):16754-9.

75.   Consortium U. UniProt: a worldwide hub of protein knowledge. Nucleic acids research.
2019;47(D1):D506-D15.

76.   Shen Y, Dinh HV, Cruz E, Call CM, Baron H, Ryseck R-P, et al. Proteome capacity
constraints favor respiratory ATP generation. 2022.

77.   Schomburg I, Jeske L, Ulbrich M, Placzek S, Chang A, Schomburg D. The BRENDA
enzyme information system-From a database to an expert system. J Biotechnol. 2017;261:194-
206.

78.   Perli T, Wronska AK, Ortiz-Merino RA, Pronk JT, Daran JM. Vitamin requirements and
biosynthesis in Saccharomyces cerevisiae. Yeast. 2020;37(4):283-304.

79.   Yasukawa T, Oda AH, Nakamura T, Masuo N, Tamura M, Yamasaki Y, et al. TAQing2.0
for genome reorganization of asexual industrial yeasts by direct protein transfection. Commun
Biol. 2022;5(1):144.

80.   Council NR. Nutrient Requirements of Fish and Shrimp. Washington, DC: The National
Academies Press; 2011. 392 p.

81.   Gu L, Zhang R, Fan X, Wang Y, Ma K, Jiang J, et al. Development of CRISPR/Cas9-
Based Genome Editing Tools for Polyploid Yeast Cyberlindnera jadinii and Its Application in
Engineering Heterologous Steroid-Producing Strains. ACS Synth Biol. 2023;12(10):2947-60.

82.   Noorman H, Baksteen J, Heijnen J, Luyben CA. The bioreactor overflow device: an
undesired selective separator in continuous cultures? Microbiology. 1991;137(9):2171-7.

83.   Rebnegger C, Vos T, Graf AB, Valli M, Pronk JT, Daran-Lapujade P, et al. Pichia
pastoris Exhibits High Viability and a Low Maintenance Energy Requirement at Near-Zero
Specific Growth Rates. Appl Environ Microbiol. 2016;82(15):4570-83.

84.   Hong KK, Hou J, Shoaie S, Nielsen J, Bordel S. Dynamic 13C-labeling experiments

prove important differences in protein turnover rate between two Saccharomyces cerevisiae

strains. FEMS Yeast Res. 2012;12(7):741-7.

85.   Maurer M, Kuhleitner M, Gasser B, Mattanovich D. Versatile modeling and optimization

of fed batch processes for the production of secreted heterologous proteins with Pichia pastoris.

Microb Cell Fact. 2006;5(1):37.

86.   Button DK. Kinetics of nutrient-limited transport and microbial growth. Microbiol Rev.

1985;49(3):270-97.

87.   van Dijken JP, Otto R, Harder W. Growth of Hansenula polymorpha in a methanol-

limited chemostat. Physiological responses due to the involvement of methanol oxidase as a

key enzyme in methanol metabolism. Arch Microbiol. 1976;111(1-2):137-44.

88.   Watteeuw CM, Armiger WB, Ristroph DL, Humphrey AE. Production of single cell

protein from ethanol by fed-batch process. Biotechnology and Bioengineering. 1979;21(7):1221-

37.