Effects of lipid membranes on RNA catalytic activity and stability

Authors: Tomasz Czerniak[a], James P. Saenz[a, b*]

[a] Technische Universität Dresden, B CUBE Center for Molecular Bioengineering, 01307

Dresden, Germany

[b] Technische Universität Dresden, Faculty of Medicine, Dresden 01307, Germany

*corresponding author and lead contact: james.saenz@tu-dresden.de

Abstract

RNA plays crucial roles in cellular organization and metabolism, and modulating its

activity is essential for maintaining cellular functions. RNA activity, involving both catalytic

(ribozymes) and translation processes, is controlled via myriad of different mechanisms

involving different binding partners such as proteins and smaller polar solutes. We previously

reported that lipid membranes can directly interact with the artificial R3C ribozyme changing

its activity, however the effect of lipids on naturally occurring ribozymes remains unknown.

Here, we report that both catalytic activity as well as RNA integrity can be controlled by the

presence of different lipid membranes. Lipid gel membranes decreased the activity of

hepatitis delta virus (HDV) and increased the hammerhead (HH) ribozyme reaction yield. The

presence of lipid liquid membrane lattices triggered RNA degradation with greater

degradation occurring in the single-stranded regions of RNA. The interplay between RNA

activity and stability in the presence of different lipid membranes introduces multiple

possibilities, where different combinations of ribozyme and lipid membrane composition

could produce different effects on activity. Taken together, these observations support the

hypothesis that the activity of both natural and artificial RNAs can be modulated by lipid

membranes which, in turn, contribute to the development of novel riboswitch-like molecules,

and lipid membrane-based RNA-biosensors.

Introduction

Ribozymes are catalytically active RNA molecules which play crucial role in the cellular

as well as viral metabolism[1–4]. The activity of ribozymes is strictly dependent on correct RNA

folding, and on the presence of interaction partners such as divalent cations. For example,

hepatitis delta virus (HDV), a pathogen involved in liver cirrhosis development, contains the

crucial ribozyme sequence involved in HDV replication in which activity is strictly dependent

on a characteristic pseudoknot folding[5,6] and the presence of divalent cations. Similarly,

another class of ribozyme, hammerhead (HH), involved, for example, in viroid and retrozyme

replication, also requires specific structures in order to be catalytically active, however it does

not necessarily require divalent cations[7,8]. Ribosome and RNAseP activity are modified by

interaction with proteins[9–11]. These phenotypes offer a range of possibilities in which different

natural ribozymes can be controlled in different ways using relatively simple changes in the

ribozyme microenvironment.

Most small ribozymes are capable of maintaining their full catalytic activity without
other macromolecules[2,12,13]. It was discovered, however, that RNA activity can be changed
through the interaction with small solutes such as ATP, theophylline, and cyclic di-guanosine
monophosphate in the special RNA classes such as aptazymes and riboswitches[14–17].
Moreover, recent research has shown that ribozyme activity can be also controlled through
interactions with much larger binding partners such as single-component lipid membranes[18].
It is, however, unclear how different lipid membranes (various lipid composition, and
membrane fluidity) can further tune the activity of ribozymes.
50
RNA-lipid interactions have been reported to involve electrostatic interactions[19–21],
but also, hydrophobic interactions have been proposed (i.e. RNA penetrating lipid membrane
lattice)[22–26]. RNA-lipid interplay can result in functional outcomes and regulatory mechanisms
that influence RNA phenotype and activity. For instance, the interaction, which can be tuned
both by the RNA structure and sequence, could change RNA conformation, which can
potentially lead to changes in ribozyme activity[18,24,25,27]. Thus, it is potentially possible to use
lipids to control the activity of ribozymes with different activity mechanisms, sequences, and
structures.
59
The effect of lipid membranes on artificial R3C ligase[28] was previously demonstrated,
showing that changes in the sequence of R3C ligase substrate significantly changed
characteristics of the ribozyme activity. Namely, the presence of lipid gel membranes
improved both the interaction between R3C ligase and its short RNA substrate via sequence-
dependent RNA-lipid binding[18]. It is, however, unclear how the presence of lipid membranes
can influence the non-artificial ribozyme activity directly by the binding of the ribozyme
molecules.
67
Here, we show how different ribozymes can be affected by interactions with lipid gel
membranes. The activities of hammerhead and HDV ribozymes were influenced differently.
The HDV ribozyme showed inhibition, while the hammerhead ribozyme exhibited an increase
in reaction rate. We further report lipid-membrane dependent RNA-degradation which
depends on membrane fluidity. Taken together, both activity changes (R3C, HH, HDV) as well
as lipid-triggered degradation create new level of control which might be employed in
synthetic as well as molecular biology, and help to understand riboregulatory processes.
75
Methods
Materials
HEPES, magnesium chloride, calcium chloride, ammonium chloride were purchased in
Carl Roth. Lead acetate, choline chloride, glycerophosphocholine, cholesterol were purchased
in Sigma Aldrich. Dioleoylphosphatidylcholine (DOPC), dipalmitoylphosphatidylcholine
(DPPC), diarachidoylphosphatidylcholine (20:0 PC) were purchased in Avanti Polar Lipids.
SybrGold was purchased in Thermo Fisher. All buffers were prepared in MilliQ water (Merck).
All chemicals were used without further purification.

RNA synthesis and purification

HDV and HH constructs were obtained via standard in vitro T7 transcription (**Suppl.**
Table 1), overnight 37 °C. RNA was separated on denaturing polyacrylamide gel, post-stained
with SybrGold, and excised from the gel using the razor blade. Gel slices were then crush-and-
soaked in the Tris-borate-EDTA buffer, 90 °C, for 1 hour. Obtained RNA was ethanol
precipitated, washed with 70% ethanol and resuspended in MilliQ water. Concentration and
integrity of RNA were estimated using absorbance readout and denaturing PAGE, respectively.
6-FAM labelled hammerhead substrate was purchased at IDT and used without further
purification.

Lipid vesicle preparation

Lipid stocks dissolved in chloroform were pipetted into a glass vial and briefly
evaporated under a steady flow of nitrogen gas. To remove organic solvent residues, a lipid
film was dried under vacuum overnight. To obtain multi-layer vesicles (MLVs), lipid films were
hydrated in the reaction buffer (10 mM HEPES pH 7, 5 mM $CaCl_2$ and 5 mM $MgCl_2$). Liposomes
were shaken above the melting temperature of the lipids for one hour with brief vortexing
every 15 minutes. A cloudy liposome suspension was transferred to the Eppendorf SafeLock
tube and freeze-and-thawed 10x in liquid nitrogen/60 °C thermoshaker (1'/5', respectively),
to reduce multi-lamellarity of the vesicles. Liposome suspension was then extruded 17x
through 100 nm polycarbonate filters to achieve a consistent size distribution of vesicles.
Liposome stocks were kept at 4 °C.

Activity assays

All of the activity assays were performed in the 10 mM HEPES pH 7, 5 mM $CaCl_2$ and 5
mM $MgCl_2$. Hammerhead (67 nM ribozyme, 133 nM substrate, 30 uL reaction volume) and
HDV (14 nM, 30 uL reaction) incubations were conducted in the temperature cycling
conditions (15 cycles 24 °C -> 60 °C, 3' each) to enhance RNA refolding[5]. After the incubation,
RNA was ethanol precipitated and resolved on denaturing PAGE. Hammerhead activity was
measured using the normalized product intensity to sum of product and substrate intensities.
HDV activity readouts were based on the SybrGold-post stained gels using the ratio of un-
reacted HDV-tRNA band to the sum of tRNA and HDV-tRNA bands. Hammerhead kinetic
analysis in diluted conditions (5 nM ribozyme, 10 nM substrate, 400 uL total reaction volume)
was performed with 100 μM lipids (either DPPC or 20:0 PC). The reactions were stopped after
3 temperature cycles.

Degradation readouts

RNA was incubated with different additives (lipid membranes, parts of lipid
headgroup, lead acetate); after the incubation RNA was ethanol precipitated, separated on
denaturing PAGE, and post-stained with SybrGold.

Results

The catalytic activity of RNA is modified by the presence of lipid membranes

We previously demonstrated an RNA-sequence dependent modification of ribozyme catalytic activity based on the presence of lipid gel membranes[18]. The model system was based on a trans-acting single-turnover R3C ligase and the activity changes were based rather on the substrate-lipid membrane interaction and not direct ribozyme-membrane interactions. Here, to diversify the potential modes of action, we chose different RNA with different mechanisms of activity: trans and cis acting ribozymes which trigger trans- and cis-cleavage of RNA substrates, hammerhead and HDV ribozyme, respectively. Both ribozymes bind to the lipid gel membranes with relatively high partition coefficient values ($>1\times10^6$)[18] and with lipid:nt ratios around 1:1 (**Fig. 1a, b**).

Activity assays of HDV and HH remarkably show different behaviors in the presence of lipids. The hammerhead final reaction yield was not affected, whereas the yield for the HDV ribozyme reaction decreased in the presence of the lipid gel membranes in a concentration-dependent way. Interestingly, the overnight incubation of HDV construct did not give a 100% reaction yield (**Suppl. Fig. 1**) which suggests that a fraction of HDV molecules bound to the lipid membrane are unable to perform the cleavage reaction, most likely due to impaired refolding. Indeed, the HDV ribozyme can fold into multiple different structures of which only a subset are catalytically active[5]. Hammerhead ribozyme, on the other hand, as the shorter molecule folds into one dominant structure. The presence of lipid gel membranes most likely does not bias its folding, thus no differences in the final reaction yield were observed (**Fig. 1b**). These differential lipid-dependent responses open a potential opportunity to create a system in which RNA activity can be selectively influenced through interactions with lipid membranes. As a proof of principle, we used hammerhead-HDV construct[29] in which both ribozymes act together creating characteristic digestion pattern. As expected, the presence of lipid gel membranes selectively inhibited the activity of HDV ribozyme whereas left hammerhead functionality unbiased (**Fig. 1c**).

To estimate if gel membranes have any beneficial effect on hammerhead ribozyme activity (e.g. through up-concentration of RNAs on the lipid surfaces, increased RNA-RNA contacts) we performed a time-lapse experiment. We observed that the overall reaction rate (yield over time, **Suppl. Fig. 2**) increased in the lipid gel membrane system (**Fig. 2**). A phase-transition condition (lipid membrane coming through liquid-gel phase transition during the reaction temperature cycling), which are closer to the biological systems than the solid gel membranes, also showed significant increase of the hammerhead reaction yield proving that the membrane does not necessarily have to be in the gel state over the course of the reaction; moreover, when the membrane is in a liquid phase, RNA can possibly diffuse off from the liquid membranes due to low binding affinity[18,21,30] allowing the RNA to freely interact with other interaction partners.

The presence of lipid liquid membranes triggers RNA degradation

It was previously shown that most RNA sequences do not bind well to biologically relevant lipid liquid membranes. However, Suga et al. reported that RNA binds to lipid liquid membranes which might affect its structure and catalytic activity[24,25,27]. Motivated by the unclear role of liquid membranes in the RNA-lipid systems we performed an activity assay in parallel with a binding assay of HDV ribozyme. We hypothesized that liquid membranes should not affect RNA activity due to the lack of a stiff, RNA-structure-locking gel membrane. High binding and inhibition of activity was observed for gel membranes which is in line with previous observations. Additionally, no significant binding nor effect on HDV activity due to the presence of liquid disordered membranes took place, and only slight HDV binding and inhibition was observed for stiffer liquid ordered membranes (**Fig. 3a**), however, we noticed that the HDV sequence undergoes degradation generating different RNA pattern than expected (this process is dependent on the lipid membrane concentration, **Suppl. Fig. 3**). To investigate which part of the lipid membrane is responsible for RNA degradation, RNA was incubated with different chemical components of the phospholipid headgroup (**Fig. 3b**). None of the components of the lipid headgroup can create stable lipid bilayers, and all are soluble in the buffer solution. No enhanced RNA degradation was observed in the presence of choline, ammonium ions, phosphate ions, and glycerophosphocholine, which proves that the presence of the liquid membrane lattice is crucial for triggering enhanced RNA degradation. The RNA degradation is dependent on lipid membrane fluidity with highest degradation observed for lipid liquid membranes, followed by liquid ordered and gel lattices (**Fig. 3c**, left panel). We noted that RNA degrades in a specific pattern rather than generating a smear of degradation products which most likely is due to structure-dependent RNA cleavage: typically single stranded parts of RNA are more prone to degrade[31,32].

To investigate if the observed RNA degradation pattern represents selective cleavage of the single stranded segments we incubated the RNA in the presence of lead which preferentially induces the cleavage at the single stranded regions. We observed that lead generated a pattern similar to lipid-dependent cleavage, however it also generated more degradation products. It is most likely that lead, as a metal ion, is smaller than lipid membranes and has better access to all parts of RNA compared with bulky lipid membrane lattice. We hypothesize that the presence of lipid liquid membranes allows single stranded parts of RNA to penetrate the hydrophobic core of the membrane[22–26] which changes chemical environment triggering specific RNA degradation.

Discussion

RNA activity is involved in a diverse range of biologically relevant processes such as viral replication[3], retrotransposon migration[4], RNA maturation[2], and gene expression control[14]. RNA activity has been shown to be regulated through the interaction with small solutes, as well as by proteins[9,10,14,15]. Here, we demonstrate the potential to control the activity of naturally occurring RNAs with lipid membranes.

Here, we characterized how the activity of catalytic RNAs was affected by lipid membranes in a gel and liquid state, which significantly differ in their fluidity. Our results extend previous findings that RNA-lipid interactions are sensitive to several factors such as membrane fluidity, lipid composition, and RNA sequence and structure[18,21,30,33–35]. For example, the activity of R3C ligase ribozyme can be controlled through interactions with lipid gel membranes in a sequence and structure-dependent manner[18]. Moreover, the interaction with lipid membranes influenced RNA structure which could, in turn, also affect RNA activity. Here, we observed lipid-based RNA-activity changes for two different naturally occurring ribozymes, HDV and HH. The HDV ribozyme was partially inhibited, likely due to the refolding issues in the presence of stiff crystal-like lipid gel membrane[36], whereas HH remained active and even accelerated under similar conditions (**Fig. 1, 2**). Alternatively, steric hindrance from the bulky lipid membranes as well as direct nucleotide-lipid interactions, may have played a role[26,37]. Remarkably, the presence of biologically relevant lipid liquid membranes affected RNA integrity, resulting in partial cleavage with characteristic patterns that depended on the presence of the lipid liquid membrane lattice (**Fig. 3**).

Our results suggest that the viroid, HDV, and retrozyme replication could also potentially be affected by the presence of the lipid membranes. Our observations of simple RNA-lipid based systems create a framework to understand if and how lipid membranes can be involved in RNA metabolism. For instance, mRNA degradation might occur at specific membrane fluidities, either activating or inhibiting its translation activity. Additionally, lipid-sensitive riboswitches, might represent another level of gene expression in which RNA is activated by lipid membranes with specific composition or property. Lipid-membrane co-localized RNAs have been found *in vivo* in the form of mRNAs[38–40], lncRNAs[41,42], and glycoRNAs[43]. Recently it was also shown that cell-extracted RNA species can bind to lipid membranes[44]. In some cases, the mechanism of binding is not entirely clear, however it was proposed that some of the RNA sequences can have affinity towards the lipid membranes[33–35,45]. The lipid-driven RNA-activity changes might have several implications in cellular biology. The close proximity of catalytic RNAs, such as ribosomes, to different lipid membranes might have the influence on the RNA activity and integrity which further could influence cellular homeostasis. It was demonstrated that lipids forming the nuclear envelope increase RNA stability[46] which, for example, might improve the activity of RNAseP, a ribozyme involved in the small RNA processing.

251   Development of lipid-binding RNAs could potentially have applications for the artificial
252 introduction of lipid-binding RNA species *in vivo*. For example, the mRNA translation process
253 could be co-localized with the lipid membranes which would provide a means to control the
254 localization of membrane protein translation[39,47]. Creating new aptazymes and bi-functional
255 aptamers could also create diverse and controllable functional outcomes; it was reported that
256 merging the lipid-binding sequences with tryptophan aptamers increased tryptophan lipid
257 membrane permeability[48]. Modifying existing RNAs with lipid-binding motifs might play a role
258 in the exosome packing processes[45,49–51]. Novel lipid-based aptazymes and riboswitches could
259 generate a pool of lipid-dependent feedback loops controlling further the stability and activity
260 of modified ribozymes and mRNAs. Lastly, a bioinformatic analysis of lipid-binding motifs in
261 currently developed lipid-binding RNAs would help to find naturally existing lipid-binding
262 RNAs.
263
264   Merging our observations of modified RNA activity (R3C ligase, hammerhead and HDV
265 ribozymes) and altered RNA stability ((including RNA degradation and lipid-based cleavage of
266 other RNA species) in the presence of lipid membranes provides a frame fork for developing
267 functional systems based on the interactions between RNA and lipids. This framework lays the
268 groundwork for understanding of RNA-lipid interplay in living cells, as well as in the synthetic
269 and prebiotic systems.
270

271 **Acknowledgements**
272   The authors would like to thank Mario Mörl lab for helpful discussions, as well as for
273 the aliquot of the HH-tRNA-HDV construct. This work was supported by the B CUBE of the TU
274 Dresden, a German Federal Ministry of Education and Research BMBF grant (to J.S., project
275 03Z22EN12), and a VW Foundation ''Life'' grant (to J.S., project 93090).
276

277 **Author contributions**
278   T.C. performed and analyzed experiments, conceptualized and wrote the manuscript.
279 J.P.S acquired the funding, conceptualized, and edited the manuscript.
280

281 **Conflict of interest**
282   The authors declare no conflict of the interest.

References

1. Cech, T. R. The Ribosome Is a Ribozyme. *Science* **289,** 878–879 (2000).

2. Kruger, K., Grabowski, P. J., Zaug, A. J., Sands, J., Gottschling, D. E. & Cech, T. R. Self-splicing
RNA: Autoexcision and autocyclization of the ribosomal RNA intervening sequence of
tetrahymena. *Cell* **31,** 147–157 (1982).

3. Jeng, K. S., Daniel, A. & Lai, M. M. A pseudoknot ribozyme structure is active in vivo and
required for hepatitis delta virus RNA replication. *J. Virol.* **70,** 2403–2410 (1996).

4. Cervera, A., Urbina, D. & Peña, M. de la. Retrozymes are a unique family of non-autonomous
retrotransposons with hammerhead ribozymes that propagate in plants through circular
RNAs. *Genome Biol.* **17,** 135 (2016).

5. Wu, H. N. & Lai, M. M. RNA conformational requirements of self-cleavage of hepatitis delta
virus RNA. *Mol Cell Biol* **10,** 5575–5579 (1990).

6. Ferré-D'Amaré, A. R., Zhou, K. & Doudna, J. A. Crystal structure of a hepatitis delta virus
ribozyme. *Nature* **395,** 567–574 (1998).

7. Murray, J. B., Seyhan, A. A., Walter, N. G., Burke, J. M. & Scott, W. G. The hammerhead,
hairpin and VS ribozymes are catalytically proficient in monovalent cations alone. *Chem. Biol.*
5, 587–595 (1998).

8. Curtis, E. A. & Bartel, D. P. The hammerhead cleavage reaction in monovalent cations. *Rna*
7, 546–552 (2001).

9. Marquez, S. M., Chen, J. L., Evans, D. & Pace, N. R. Structure and Function of Eukaryotic
Ribonuclease P RNA. *Mol. Cell* **24,** 445–456 (2006).

10. Nissen, P., Hansen, J., Ban, N., Moore, P. B. & Steitz, T. A. The Structural Basis of Ribosome
Activity in Peptide Bond Synthesis. *Science* **289,** 920–930 (2000).

11. Pool, M. R., Stumm, J., Fulga, T. A., Sinning, I. & Dobberstein, B. Distinct Modes of Signal
Recognition Particle Interaction with the Ribosome. *Science* **297,** 1345–1348 (2002).

12. Horning, D. P. & Joyce, G. F. Amplification of RNA by an RNA polymerase ribozyme. *Proc*
National Acad Sci **113,** 9786–9791 (2016).

13. Paul, N. & Joyce, G. F. A self-replicating ligase ribozyme. *Proc National Acad Sci* **99,** 12733–
12740 (2002).

14. Sudarsan, N., Lee, E. R., Weinberg, Z., Moy, R. H., Kim, J. N., Link, K. H. & Breaker, R. R.
Riboswitches in Eubacteria Sense the Second Messenger Cyclic Di-GMP. *Science* **321,** 411–413
(2008).

15. Robertson, M. P. & Ellington, A. D. Design and optimization of effector-activated ribozyme
ligases. *Nucleic Acids Res* **28,** 1751–1759 (2000).

16. Tang, J. & Breaker, R. R. Rational design of allosteric ribozymes. *Chem Biol* **4,** 453–459
(1997).

17. Sekella, P. T., Rueda, D. & Walter, N. G. A biosensor for theophylline based on fluorescence
detection of ligand-induced hammerhead ribozyme cleavage. *RNA* **8,** 1242–1252 (2002).

18. Czerniak, T. & Saenz, J. P. Lipid membranes modulate the activity of RNA through
sequence-dependent interactions. *Proc National Acad Sci* **119,** e2119235119 (2022).

19. Budker, V. G., Godovikov, A. A., Naumova, L. P. & Slepneva, I. A. Interaction of
polynucleotides with natural and model membranes. *Nucleic acid research* (1980).

20. Budker, V. G., Kazatchkov, Yu. A. & Naumova, L. P. Polynucleotides adsorb on
mitochondrial and model lipid membranes in the presence of bivalent cations. *Febs Lett* **95,**
143–146 (1978).

21. Pannwitt, S., Slama, K., Depoix, F., Helm, M. & Schneider, D. Against Expectations:
Unassisted RNA Adsorption onto Negatively Charged Lipid Bilayers. *Langmuir* **35,** 14704–
14711 (2019).

22. Michanek, A., Yanez, M., Wacklin, H., Hughes, A., Nylander, T. & Sparr, E. RNA and DNA
Association to Zwitterionic and Charged Monolayers at the Air–Liquid Interface. *Langmuir* **28,**
9621–9633 (2012).

23. Michanek, A., Kristen, N., Höök, F., Nylander, T. & Sparr, E. RNA and DNA interactions with
zwitterionic and charged lipid membranes — A DSC and QCM-D study. *Biochimica Et
Biophysica Acta Bba - Biomembr* **1798,** 829–838 (2010).

24. Suga, K., Umakoshi, H., Tomita, H., Tanabe, T., Shimanouchi, T. & Kuboi, R. Liposomes
destabilize tRNA during heat stress. *Biotechnol J* **5,** 526–529 (2010).

25. Suga, K., Tanabe, T., Tomita, H., Shimanouchi, T. & Umakoshi, H. Conformational change
of single-stranded RNAs induced by liposome binding. *Nucleic Acids Res* **39,** 8891–8900 (2011).

26. Marty, R., N'soukpoé-Kossi, C. N., Charbonneau, D. M., Kreplak, L. & Tajmir-Riahi, H.-A.
Structural characterization of cationic lipid–tRNA complexes. *Nucleic Acids Res* **37,** 5197–5207
(2009).

27. Suga, K., Tanaka, S. & Umakoshi, H. Liposome membrane can induce self-cleavage of RNA
that models the core fragments of hammerhead ribozyme. *Eur Biophys J* **45,** 55–62 (2016).

28. Rogers, J. & Joyce, G. F. The effect of cytidine on the structure and function of an RNA
ligase ribozyme. *Rna* **7,** 395–404 (2001).

29. Schürer, H., Lang, K., Schuster, J. & Mörl, M. A universal method to produce in vitro
transcripts with homogeneous 3' ends. *Nucleic Acids Res.* **30,** e56–e56 (2002).

30. Janas, T., Janas, T. & Yarus, M. Specific RNA binding to ordered phospholipid bilayers.
Nucleic Acids Res **34,** 2128–2136 (2006).

31. Soukup, G. A. & Breaker, R. R. Relationship between internucleotide linkage geometry and
the stability of RNA. *Rna* **5,** 1308–1325 (1999).

32. Ciesiolka, J., Lorenz, S. & Erdmann, V. A. Different conformational forms of Escherichia coli
and rat liver 5S rRNA revealed by Pb(II)-induced hydrolysis. *Eur J Biochem* **204,** 583–589
(1992).

33. Vlassov, A., Khvorova, A. & Yarus, M. Binding and disruption of phospholipid bilayers by
supramolecular RNA complexes. *Proc National Acad Sci* **98,** 7706–7711 (2001).

34. Vlassov, A. & Yarus, M. Interaction of RNA with Phospholipid Membranes. *Mol Biol+* **36,**
389–393 (2002).

35. Khvorova, A., Kwak, Y.-G., Tamkun, M., Majerfeld, I. & Yarus, M. RNAs that bind and change
the permeability of phospholipid membranes. *Proc National Acad Sci* **96,** 10649–10654 (1999).

36. Swadling, J. B., Suter, J. L., Greenwell, H. C. & Coveney, P. V. Influence of Surface Chemistry
and Charge on Mineral–RNA Interactions. *Langmuir* **29,** 1573–1583 (2013).

37. Xu, Y., Huang, S.-W., Ding, H.-M. & Ma, Y.-Q. Molecular dynamics simulations on the
interactions between nucleic acids and a phospholipid bilayer. *Chin. Phys. B* **33,** 028701 (2024).

38. Jambor, H., Surendranath, V., Kalinka, A. T., Mejstrik, P., Saalfeld, S. & Tomancak, P.
Systematic imaging reveals features and changing localization of mRNAs in Drosophila
development. *Elife* **4,** e05003 (2015).

39. Kannaiah, S., Livny, J. & Amster-Choder, O. Spatiotemporal Organization of the E. coli
Transcriptome: Translation Independence and Engagement in Regulation. *Mol. Cell* **76,** 574-
589.e7 (2019).

40. Janas, T., Sapoń, K., Janas, T. & Yarus, M. Specific binding of VegT mRNA localization signal
to membranes in Xenopus oocytes. *Biochimica Et Biophysica Acta Bba - Mol Cell Res* **1868,**
118952 (2021).

41. Li, R.-H., Tian, T., Ge, Q.-W., He, X.-Y., Shi, C.-Y., Li, J.-H., Zhang, Z., Liu, F.-Z., Sang, L.-J.,
Yang, Z.-Z., Liu, Y.-Z., Xiong, Y., Yan, Q., Li, X., Ju, H.-Q., Liu, J., Wang, L.-J., Shao, J.-Z., Wang,
W., Zhou, T. & Lin, A. A phosphatidic acid-binding lncRNA SNHG9 facilitates LATS1 liquid–liquid
phase separation to promote oncogenic YAP signaling. *Cell Res* **31,** 1088–1105 (2021).

42. Lin, A., Hu, Q., Li, C., Xing, Z., Ma, G., Wang, C., Li, J., Ye, Y., Yao, J., Liang, K., Wang, S., Park,
P. K., Marks, J. R., Zhou, Y., Zhou, J., Hung, M.-C., Liang, H., Hu, Z., Shen, H., Hawke, D. H., Han,
L., Zhou, Y., Lin, C. & Yang, L. The LINK-A lncRNA interacts with PtdIns(3,4,5)P3 to hyperactivate
AKT and confer resistance to AKT inhibitors. *Nat Cell Biol* **19,** 238–251 (2017).

43. Flynn, R. A., Pedram, K., Malaker, S. A., Batista, P. J., Smith, B. A. H., Johnson, A. G., George,
B. M., Majzoub, K., Villalta, P. W., Carette, J. E. & Bertozzi, C. R. Small RNAs are modified with
N-glycans and displayed on the surface of living cells. *Cell* **184,** 3109-3124.e22 (2021).

44. Bayona-Hernandez, A., Guerra, S., Jiménez-Ramirez, I. A., Sztacho, M., Hozak, P.,
Rodriguez-Zapata, L. C., Pereira-Santana, A. & Castaño, E. LIPRNAseq: a method to discover
lipid interacting RNAs by sequencing. *Mol. Biol. Rep.* **50,** 6619–6626 (2023).

45. Mańka, R., Janas, P., Sapoń, K., Janas, T. & Janas, T. Role of RNA Motifs in RNA Interaction
with Membrane Lipid Rafts: Implications for Therapeutic Applications of Exosomal RNAs. *Int J*
Mol Sci **22,** 9416 (2021).

46. Micheli, M., Albi, E., Leray, C. & Magni, M. V. Nuclear sphingomyelin protects RNA from
RNase action. *FEBS Lett.* **431,** 443–447 (1998).

47. Nevo-Dinur, K., Nussbaum-Shochat, A., Ben-Yehuda, S. & Amster-Choder, O. Translation-
Independent Localization of mRNA in E. coli. *Science* **331,** 1081–1084 (2011).

48. Janas, T., Janas, T. & Yarus, M. A membrane transporter for tryptophan composed of RNA.
Rna **10,** 1541–1549 (2004).

49. Janas, T., Janas, M. M., Sapoń, K. & Janas, T. Mechanisms of RNA loading into exosomes.
Febs Lett **589,** 1391–1398 (2015).

50. Batagov, A. O., Kuznetsov, V. A. & Kurochkin, I. V. Identification of nucleotide patterns
enriched in secreted RNAs as putative cis-acting elements targeting them to exosome nano-
vesicles. *Bmc Genomics* **12,** S18–S18 (2011).

51. Janas, T., Sapoń, K. & Janas, T. Selection of bifunctional RNAs with specificity for arginine
and lipid membranes. *FEBS Lett.* **598,** 1061–1079 (2024).

Fig. 1 Lipid gel membranes act differently on different ribozymes. a) HDV ribozyme self cleaves under temperature cycling conditions. The reaction yield (green curve) decreases with increasing lipid binding (orange curve). **b)** HH ribozyme cleaves its substrate which generates shorter RNA products. Despite significant lipid binding (orange curve) there is no change in the total reaction yield (green datapoints). **c)** HH-tRNA-HDV fusion construct (top band) self-cleaves in two separate spots generating three products: HH, HDV, and tRNA. The presence of lipid gel membranes selectively partially inhibits HDV activity generating half-cleaved intermediate. The obtained binding and activity fits are described by Hill's function.

Fig. 2 Increased reaction rate of hammerhead ribozyme in the presence of lipid gel membranes. The hammerhead ribozyme reaction was stopped after 3 temperature cycles in which the reaction did not reach completion for lipid—free system (**Suppl. Fig. 2**). The reaction yield is significantly higher for the lipid-based systems.

a) Lipid membranes trigger HDV degradation

b) Lipid membrane lattice is necessary to degrade RNA

c) Sequence-dependent degradation pattern

Fig. 3 Lipid membranes induce RNA degradation.
a) HDV-tRNA construct was incubated in the presence of 1 mM lipids. RNA and lipids were then and analyzed on the denaturing PAGE. W – reaction before separation, SN – supernatant (lipid free fraction), P – pellet (lipid bound fraction). **b)** HDV RNA was incubated in the presence of 1 mM additives at 20 °C for 24 hours. The presence of water-soluble additives did not influence RNA integrity whereas liquid disordered lipid membranes (DOPC) induced RNA degradation. **c)** 50 ng of HH RNA was incubated for 3 hours at room temperature with 1 mM lipids with different membrane fluidities. Ld – liquid disordered (DOPC), Lo – liquid ordered (DPPC:Chol30%), s – gel phase (DPPC). Additionally, the RNA was incubated for 10' in the presence of 10 mM lead acetate (Pb) which preferentially triggers RNA cleavage at the single-stranded parts of RNA. Dashed line on the right gel symbolizes that a middle lane was removed from the original gel image.