Species limits and hybridization in Andean leaf-eared mice (*Phyllotis*)

Marcial Quiroga-Carmona[1,2,3], Schuyler Liphardt[4], Naim M. Bautista[1], Pablo Jayat[5,6], Pablo Teta[7],

Jason L. Malaney[8], Tabitha McFarland[9,10], Joseph A. Cook[9,10], L. Moritz Blumer[11], Nathanael D.

Herrera[4], Zachary A. Cheviron[4], Jeffrey M. Good[4], Guillermo D'Elía[2,3], Jay F. Storz[1]

[1]School of Biological Sciences, University of Nebraska, Lincoln, NE, United States

[2]Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias, Universidad Austral de Chile,

Valdivia, Chile

[3]Colección de Mamíferos, Facultad de Ciencias, Universidad Austral de Chile, Campus Isla Teja,

Valdivia, Chile

[4]Division of Biological Sciences, University of Montana, Missoula, MT, United States

[5]Unidad Ejecutora Lillo (CONICET-Fundación Miguel Lillo), San Miguel de Tucumán, Argentina

[6]Departamento de Ciencias Básicas y Tecnológicas, Universidad Nacional de Chilecito (UNdeC),

Argentina

[7]División Mastozoología, Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Ciudad

Autónoma de Buenos Aires, Argentina

[8]New Mexico Museum of Natural History and Science, Albuquerque, NM, United States

[9]Museum of Southwestern Biology, University of New Mexico, Albuquerque, NM, United States

[10]Department of Biology, University of New Mexico, Albuquerque, NM, United States

[11]Department of Genetics, University of Cambridge, Cambridge, United Kingdom

Correspondence:

Jay F. Storz

School of Biological Sciences

University of Nebraska

Lincoln, Nebraska, USA

E-mail: jstorz2@unl.edu

ABSTRACT

Leaf-eared mice (genus *Phyllotis*) are among the most widespread and abundant small mammals in the Andean Altiplano, but species boundaries and distributional limits are often poorly delineated due to sparse survey data from remote mountains and high-elevation deserts. Here we report a combined analysis of mitochondrial DNA variation and whole-genome sequence (WGS) variation in *Phyllotis* mice to delimit species boundaries, to assess the timescale of diversification of the group, and to examine evidence for interspecific hybridization. Estimates of divergence dates suggest that most diversification of *Phyllotis* occurred during the past 3 million years. Consistent with the Pleistocene Aridification hypothesis, our results suggest that diversification of *Phyllotis* largely coincided with climatically induced environmental changes in the mid- to late Pleistocene. Contrary to the Montane Uplift hypothesis, most diversification in the group occurred well after the major phase of uplift of the Central Andean Plateau. Species delimitation analyses revealed surprising patterns of cryptic diversity within several nominal forms, suggesting the presence of much undescribed alpha diversity in the genus. Results of genomic analyses revealed evidence of ongoing hybridization between the sister species *Phyllotis limatus* and *P. vaccarum* and suggest that the contemporary zone of range overlap between the two species may represent an active hybrid zone.

Running title: Species limits of Andean mice

Keywords: Altiplano, Andes, geographic range limits, introgression, Puna de Atacama, species delimitation

1. INTRODUCTION

Leaf-eared mice in the genus *Phyllotis*, Waterhouse 1873, are emblematic mammals of the Andean Altiplano and have an exceptionally broad latitudinal distribution in South America, from Ecuador to the northern coast of the Strait of Magellan (Steppan & Ramírez, 2015). The genus has an even more impressive elevational distribution: Whereas *P. darwini* is found at sea level along the desert coastline of northern Chile, and species like *P. anitae*, *P. nogalaris*, and *P. osilae* are found in humid, lowland Yungas forests on the eastern sub-Andean slopes (Jayat et al., 2016), other taxa such as *P. vaccarum* have been documented at extreme elevations (>6000 m above sea level) on the upper reaches and summits of some of the highest peaks in the Andean Cordillera (Storz et al., 2020; Steppan et al., 2022; Storz et al., 2023, 2024). Although *Phyllotis* mice are among the most widespread and abundant small mammals in the Andean Altiplano and adjacent lowlands, the taxonomic status and range limits of many species are not well-resolved due to sparse survey data from remote mountains and high-elevation deserts (puna). The resultant gaps in sampling coverage have hindered a complete assessment of species richness and geographic distributions of *Phyllotis* mice.

Over the last two decades, *Phyllotis* has been subject to several taxonomic assessments that have helped resolve species limits and phylogenetic relationships (Jayat et al., 2007, 2016, 2021; Ojeda et al., 2021; Steppan et al., 2007; Rengifo & Pacheco, 2015, 2017; Teta et al., 2018, 2022). There are currently 26 recognized species of *Phyllotis*, and the genus comprises three main clades, commonly referred to as the *andium-amicus*, *osilae*, and *darwini* species groups (Rengifo & Pacheco, 2017; Steppan, 1993, 1995; Steppan et al., 2007; Steppan & Ramírez, 2015; Teta et al., 2022). The *darwini* group is the most speciose and includes several species that are broadly co-distributed in the Atacama Desert and Andean dry puna: *P. caprinus*, '*P. chilensis*' (*sensu* Pearson, 1958; referred to as '*P. posticalis-P. rupestris*' by Ojeda et al., 2021), *P. darwini*, *P. limatus*, *P. magister*, and *P. vaccarum* (Jayat et al., 2021; Ojeda et al., 2021; Steppan & Ramírez, 2015; Storz et al., 2024; Teta et al., 2022). This set of closely related species form part of the so-called *P. xanthopygus* complex (Steppan et al., 2007; Walker et al., 1984). In northeastern Chile and bordering regions of Argentina and Bolivia, the ranges of several of these species potentially overlap (Figure 1a), but in most cases the distribution limits are not clearly defined. We often do not know the extent to which species ranges overlap across Andean elevational gradients, which is important for understanding the relative roles of competitive exclusion and physiological tolerances in shaping elevational patterns of species turnover and for detecting distributional shifts in response to climate change.

In this same region, genomic delimitation of species boundaries between *P. limatus* and *P. vaccarum* in northern Chile led to a dramatically revised understanding of the latitudinal and elevational range limits of the former species (Storz et al., 2024). Previously inferred range limits of *P. limatus* were found to be in error because specimens from the highest elevations and most southern

latitudes had been mis-identified as *P. limatus* on the basis of mitochondrial (mt) DNA and were later

identified as *P. vaccarum* on the basis of whole-genome sequence data (Storz et al., 2024). The fact

that some *P. vaccarum* carry mtDNA haplotypes more closely related to those of *P. limatus* suggests a

history of introgressive hybridization and/or incomplete lineage sorting. In addition to highlighting the

importance of using multilocus data to define species limits, the observed mitonuclear discordance in

P. limatus and *P. vaccarum* suggests the possibility of hybridization between other pairs of *Phyllotis*

species in regions of historical or contemporary range overlap.

Here we report a combined analysis of mtDNA variation and whole-genome sequence (WGS)

variation in *Phyllotis* mice to delimit species boundaries, to assess the timescale of diversification of

the group, and to examine evidence for interspecific hybridization. The analysis is principally focused

on a large set of vouchered specimens that we collected over the course of five high-elevation survey

expeditions in the Puna de Atacama, Central Andes (2020-2023), in conjunction with additional

collecting trips in the surrounding Altiplano and adjoining lowlands in Argentina, Bolivia, and Chile. The

genomic analysis is primarily focused on members of the *P. darwini* species group that have

overlapping or potentially overlapping ranges.

2. MATERIAL AND METHODS

2.1 SPECIMEN COLLECTION

We collected representatives of multiple species of *Phyllotis* during the course of small-mammal

surveys in the Altiplano and adjoining lowlands on both sides of the Andean Cordillera in Chile, Bolivia,

and Argentina. We captured all mice using Sherman live traps, in combination with Museum Special

snap traps at some localities. We sacrificed animals in the field, prepared them as museum

specimens, and preserved liver tissue in ethanol as a source of genomic DNA. All specimens are

housed in the mammal collections of the Universidad Austral de Chile, Valdivia, Chile (UACH),

Colección Boliviana de Fauna, La Paz, Bolivia (CBF), Centro Regional de Investigaciones Científicas

y Transferencia Tecnológica de La Rioja, La Rioja, Argentina (CRILAR), Centro Nacional Patagónico,

Chubut, Argentina (CNP), Fundación-Instituto Miguel Lillo, Tucumán, Argentina (CML), Museo

Argentino de Ciencias Naturales "Bernardino Rivadavia", Ciudad Autónoma de Buenos Aires,

Argentina (MACN-Ma), or the Museum of Southwestern Biology, New Mexico, USA (MSB). We

identified all specimens to the species level based on external characters (Jayat et al., 2021; Steppan

& Ramírez, 2015; Teta et al., 2022) and, as described below, we later confirmed field-identifications

with DNA sequence data.

In Chile, all animals were collected in accordance with permissions to JFS, MQC, and GD from

the following Chilean government agencies: Servicio Agrícola y Ganadero (6633/2020, 2373/2021,

5799/2021, 3204/2022, 3565/2022, 911/2023 and 7736/2023), Corporación Nacional Forestal

(171219, 1501221, and 31362839), and Dirección Nacional de Fronteras y Límites del Estado

(DIFROL, Autorización de Expedición Científica #68 and 02/22). In Bolivia, all animals were collected

in accordance with permissions to JFS (Resolución Administrativa 026/09) and JAC (DVS-CRT-02/91)

from the Ministerio de Medio Ambiente y Agua, Estado Plurinacional de Bolivia. In Argentina, all

animals were collected in accordance with the following permissions to JPJ from the Ministerio de

Ambiente y Cambio Climático de Jujuy: Expte. N° P4-00402-21 Disp. S.A. N° 001/22, Expte. N° P4 -

00158 -22 Disp. S.A. N° 007/22 and Expte. N° 677-330-2021. All live-trapped animals were handled in

accordance with protocols approved by the Institutional Animal Care and Use Committee (IACUC) of

the University of Nebraska (project ID's: 1919, 2100), IACUC of the University of New Mexico (project

ID's: 16787 and 20405), and the bioethics committee of the Universidad Austral de Chile (certificate

456/2022).

2.2 SEQUENCE DATA

To maximize geographic coverage in our survey of mtDNA variation, we generated sequence data for

a subset of our own voucher specimens (*n*=269) and supplemented this dataset with publicly available

Phyllotis sequences from GenBank (*n*=180). This sequence dataset, based on a total of 449

specimens, includes 20 of the 26 nominal species that are currently recognized within the genus

Phyllotis. We used a subset of our newly collected voucher specimens (*n*=137) for the analysis of

WGS variation.

2.3 MITOCHONDRIAL DNA VARIATION

For the analysis of mtDNA variation, we extracted DNA from liver samples and PCR-amplified the first

801 base pairs of the *cytochrome b* (*cytb*) gene using the primers MVZ 05 and MVZ 16 (Smith and

Patton 1993), following protocols of Cadenillas and D'Elía (2021). Of the 269 *cytb* sequences that we

generated from our own set of voucher specimens, 89 were published previously (Storz et al., 2020,

2024; GenBank accession numbers: OR784643-OR784661, OR799565-OR799614, and OR810731-

OR810743). We deposited all newly generated sequences in GenBank (accession numbers: XXX-

XXX [pending]). The newly generated sequences derive from voucher specimens housed in the

Argentine, Bolivian, Chilean, and US collections mentioned above (section 2.1).

2.4 PHYLOGENY ESTIMATION

As outgroups for the phylogenetic analysis, we used *cytb* sequences from five other phyllotine rodents

(*Auliscomys boliviensis*, JQ434420; *A. pictus*, U03545; *A. sublimis*, U03545; *Calomys musculinus*,

HM167822; and *Loxondontomys micropus*, GU553838). The final set of 454 sequences was aligned

with MAFFT v7 (Katoh et al., 2017) using the E-INS-i strategy to establish character primary

homology. The aligned matrix was visually inspected with AliView v1.26 (Larsson, 2014) to check for

the presence of internal stop codons and shifts in the reading frame. Pairwise genetic distances and

their standard errors (p-dist./SE) were calculated using MEGA X 10.1.8 (Kumar et al., 2018).

Redundant *cytb* sequences were identified and discarded using the functions *FindHaplo* and

haplotype in the *sidier* (Pajares, 2013) and *haplotypes* (Aktas, 2023) R packages, respectively. The

final matrix of nonredundant sequences included a total of 287 haplotypes.

The nucleotide substitution model (HKY + I + G) that provided the best fit to the nonredundant

cytb data matrix was selected based on the Bayesian Information Criterion (BIC) using ModelFinder

(Kalyaanamoorthy et al., 2017). Genealogical relationships among haplotypes of *Phyllotis* species

were estimated via Maximum Likelihood (ML) and Bayesian Inference (BI). The ML analysis was

performed using IQ-TREE (Trifinopoulos et al., 2016), with perturbation strength set to 0.5 and the

number of unsuccessful iterations set to 100. Nodal support was assessed through 1000 ultrafast

bootstrap replicates (UF; Minh et al., 2013). BI was implemented with BEAST 2 v2.6.7 (Bouckaert et

al., 2014), which was also used to estimate divergence dates among *Phyllotis* species. A gamma site

model was selected with the substitution model set to HKY. The gamma shape parameter (exponential

prior, mean 1.0) and proportion of invariant sites (uniform distribution, 0.001–0.999, lower and upper

bounds) were estimated. To prevent the sampling of excessively small values for the HKY

exchangeability rates, the prior sampling distribution was set to gamma with a shape parameter

(alpha) of 2.0 and a scale parameter (beta) of 0.5. The clock model was set to Relaxed Log Normal

with an estimated clock rate. The calibrated Yule model was selected to parameterize fossil

calibrations. For the mean branch rate (ucldMean), an exponential sampling distribution was applied

with a mean of 10.0 and no offset. Given that variation in substitution rates among branches is low and

evidence suggests that molecular evolution is largely clock-like across Phyllotini (Parada et al., 2013),

standard deviation in rates across branches (ucldStdev) was converted to an exponential prior

distribution with a mean of 0.3337 and no offset. Since the fossil record for *Phyllotis* is not sufficient to

establish primary calibration points (Pardiñas et al., 2002), we used secondary calibration points

based on an estimated phylogeny of the subfamily Sigmodontinae (Parada et al., 2015). We used

95% credibility intervals for estimated crown ages of the genus *Phyllotis* (3.35-6.66 Mya), the *darwini*

species group (4.51-1.77 Mya), and the *xanthopygus* species complex (1.33-2.46 Mya). We performed

two runs of 600 x 10^6 MCMC generations with trees sampled every 4 x 10^3 steps, yielding 15,001

samples for parameter estimates. Effective sample sizes greater than 200 for all parameters (i.e.,

stable values of convergence) were verified using Tracer v1.7.1 (Rambaut et al., 2018). Runs were

combined with LogCombiner v2.6.7 (Bouckaert et al., 2014), using a 10% burn-in that was determined

by examining individual traces. The first 10% of estimated trees were discarded and the remainder

were used to construct a maximum clade credibility tree with posteriori probability values (PP) and age

estimates employing TreeAnnotator v2.6.2 (Rambaut & Drummond, 2019).

2.5 ASSESSMENT OF SPECIES LIMITS WITHIN THE *P. XANTHOPYGUS* SPECIES COMPLEX

To delimit species within the *P. darwini* group, we employed the Bayesian time calibrated-ultrametric

tree estimated with BEAST 2 and two single-locus coalescent methods: The General Mixed Yule

Coalescent model (GMYC; Pons et al., 2006; Fujisawa & Barraclough, 2013) and the Poisson Tree

Processes (PTP; Zhang et al., 2013). Both methods are based on the fit of different mixed models (the

General Mixed Yule Coalescent model in the case of the GMYC, and the Poisson Tree Processes in

the case of the PTP) to processes of interspecific diversification and/or genealogical branching within

species (Fujisawa & Barraclough, 2013; Zhang et al., 2013). These methods were implemented via

their online web servers: https://species.h-its.org/gmyc/ and http://species.h-its.org/ptp/, respectively.

The Bayesian implementations of these methods (b-GMYC: Reid & Carstens, 2012; b-PTP: Zhang et

al., 2013) were also employed to account for uncertainty in gene tree estimation. The b-GMCY

analysis was implemented in R via the *b-GMCY* R package (Reid and Carstens 2012), which offers

estimates of the posterior marginal probabilities for candidate species, setting a post-burn-in sample of

1000 trees sampled from the posterior distribution of trees. For all parameters, priors were set as

default (i.e., t1 and t2 were set at 2 and 100, respectively), and the analysis was completed with 50 x

10^3 generations, burning 10% of these and with a thinning interval of 1000 samples. The b-PTP

analysis was implemented in the associated online web server (http://species.h-its.org/b-ptp/) with

default values (i.e., 100 x 10^3 MCMC, thinning of 100 and burning of 0.1). Branch lengths are

proportional to coalescence times in the GMYC model, whereas they are proportional to the number of

nucleotide substitutions in the PTP model (Dellicour & Flot, 2018).

2.6 WHOLE-GENOME SEQUENCE DATA

We generated low-coverage whole-genome sequence (WGS) data for a subset of 137 *Phyllotis*

specimens that were included in the *cytb* data matrix, which we analyzed in conjunction with a

chromosome-level reference genome for *Phyllotis vaccarum* (Storz et al., 2023). Depth of coverage

ranged from 1.04x to 24.06X (median = 2.58X). According to field identifications and *cytb* haplotypes,

this set of specimens represented a total of 11 species (*P. anitae*, *P. camiari*, *P. caprinus*, *P. chilensis*,

P. darwini, *P. limatus*, *P. magister*, *P. nogalaris*, *P. pehuenche*, *P. vaccarum*, and *P. xanthopygus*),

several of which have potentially overlapping ranges (Figure 1a). All species other than *P. anitae* and

P. nogalaris are members of the *darwini* species group. Of the 137 vouchered specimens included in

the genomic analysis, data for 61 specimens representing *P. chilensis*, *P. limatus*, *P. magister*, and *P.*

vaccarum were published previously (Storz et al., 2024).

2.6.1 *Genomic library preparation and whole-genome sequencing*

All library preparations for whole genome resequencing experiments were conducted in the University

of Montana Genomics Core facility. We extracted genomic DNA from ethanol-preserved liver tissue

using the DNeasy Blood and Tissue kit (Qiagen). We used a Covaris E220 sonicator to shear DNA

and we then prepared genomic libraries using the KAPA HyperPlus kit (Roche). Individual libraries

were indexed using KAPA UDI's and pooled libraries were sent to Novogene for Illumina paired-end

150 bp sequencing on a Novaseq X.

2.6.2 *Read quality processing and mapping to the reference genome*

We used fastp 0.23.2 (Chen et al., 2018) to remove adapter sequences, and to trim and filter low-

quality reads from sequences generated from library preparations. We used a 5 bp sliding window to

remove bases with a mean quality less than 20 and we discarded all reads <25 bp. We merged all

overlapping reads that passed filters and retained all reads that could not be merged or whose paired

reads failed filtering. We separately mapped merged reads, unmerged but paired reads, and unpaired

reads to the *P. vaccarum* reference genome with BWA 0.7.17 (Li & Durbin, 2009) using the mem

algorithm with the -M option which flags split reads as secondary for downstream compatibility. We

sorted, merged, and indexed all resulting binary alignment maps with SAMtools 1.15.1 (Li et al., 2009)

and used picard 2.27.4 to detect and remove PCR duplicates. We used GATK 3.8 (McKenna et al.,

2010) to perform local realignment around targeted indels to generate the final BAM files.

2.6.3 *Mitochondrial genome assembly*

a *de novo* assembly of the mitochondrial genome of *Phyllotis vaccarum* (specimen UACH8291) as a

seed sequence, we used NOVOplasty 4.3.3 (Dierckxsens et al., 2017) to generate *de novo*

mitochondrial genome assemblies for all other *Phyllotis* specimens. We annotated assembled

mitochondrial genomes with MitoZ to identify coding sequences and we generated a multiple

alignment of coding sequence with MAFFT 7.508 (Katoh & Standley, 2013), using the --auto flag to

determine the best algorithm given the data.

2.7 ANALYSIS OF WHOLE-GENOME SEQUENCE VARIATION IN *PHYLLOTIS*

First, we randomly downsampled all higher coverage samples to the median coverage (2.58X) using

SAMtools 1.17 to avoid artifacts associated with variation in coverage across samples that can impact

inferences of population structure. We calculated genotype likelihoods for scaffolds 1-19 (covering

>90% of the *Phyllotis* genome) for all samples in ANGSD 0.939 (Korneliussen et al., 2014). We used -

GL 2 to specify the GATK model for genotype likelihoods, retained only sites with a probability of being

variable >1e-6 with -SNP_pval 1e-6. We filtered out bad and non-uniquely mapped reads with -

remove_bads 1 and -uniqueOnly 1, respectively, and only retained reads and bases with a mapping

quality higher than 20. We adjusted mapping quality for excessive mismatches with -C 50. We used

PCAngsd v.0.99.0 (Meisner & Albrechtsen, 2018) to calculate the covariance matrix from genotype

likelihoods and used a minor allele frequency filter of 0.05. Finally, we calculated eigenvectors and

plotted the first, second, and third principal components using the R package *ggplot2* (Wickham,

2016).

Based on results of our genus-wide genomic PCA, we recalculated genotype likelihoods and

performed additional genomic analyses on a subset of *P. vaccarum* and *P. limatus* specimens (*n*=51

and 20, respectively). To test for admixture between *P. vaccarum* and *P. limatus*, we calculated

ancestry proportions with NGSadmix (Skotte et al., 2013). To alleviate computational costs associated

with NGSadmix we generated a reduced SNP set by sampling every hundredth SNP calculated by

ANGSD. We ran NGSadmix with K=1-10 with ten iterations for each K value with a random starting

seed and a minor allele frequency filter of 0.05. We evaluated the optimal K value using EvalAdmix

0.95 which calculates the pairwise covariance matrix of residuals of model fit. The results of

EvalAdmix determined K=2 as the optimal value of K. We combined individual runs for each K value

with the R package PopHelper 2.3.1 to average estimates of ancestry across runs.

2.8 GENOMIC PATTERNING OF ADMIXTURE

To examine the genomic patterning of mixed *P. vaccarum*/*P. limatus* ancestry, we conducted a

windowed PCA of nucleotide variation. We used the script windowed_pcangsd.py

(10.5281/zenodo.8127993) to compute the first principal component in 90% overlapping 1 Mbp

windows along chromosomes 1 to 19, using the subset of 51 *P. vaccarum* and 20 *P. limatus* samples

and employing minor allele frequency threshold of 0.01. For visualization we excluded outlier windows

(those with less than 0.3 % informative sites and those featuring the largest 0.005 % absolute PC1

values across the genome). For consistency we polarized PC1 orientation by its sign for chromosome

1 since polarity is arbitrary in principal component analyses.

3. RESULTS

The analysis of mtDNA data was based on a total of 449 *Phyllotis* specimens from 169 localities that

span most of the distributional range of the genus (Figure 1b). For the analysis of WGS variation, we

used a subset of 137 vouchered specimens representing 11 nominal species of *Phyllotis* that have

overlapping or potentially overlapping ranges in Argentina, Bolivia, and Chile. *Phyllotis vaccarum* is

one of the most broadly distributed species in this region and different parts of its range potentially

overlap with those of *P. caprinus*, *P. chilensis*, *P. darwini*, *P. limatus*, *P. magister*, and *P. pehuenche*

(Figure 1a). We therefore concentrated much of our sampling efforts on these zones of range overlap

to examine evidence of introgressive hybridization.

3.1 PHYLOGENETIC RELATIONSHIPS AND DIVERGENCE TIMES

At the level of the genus *Phyllotis*, phylogeny estimates based on BI and ML both recovered three

main clades corresponding to the *andium-amicus*, *osilae*, and *darwini* species groups (Figure 2 and

Figure S1). In the BI analysis, the *andium-amicus* and *osilae* clades were recovered as sister groups

(Bayesian Posterior Probability [PP] = 1) (Figure 2), whereas the ML analysis placed the *osilae* clade

as sister to the clade formed by *andium-amicus* and *darwini* (Bootstrap Percentage [BP] = 53) (Figure

S1). Within the *darwini* group, BI and ML analyses generally recovered the same set of relationships

within the *P. xanthopygus* complex, with the exception that the BI phylogeny placed *P. pehuenche* and

P. xanthopygus as sister (PP = 1; Figure 2), whereas the ML phylogeny placed *P. xanthopygus* as

sister to the clade containing *P. caprinus*, *P. limatus*, *P. vaccarum*, and *P. pehuenche* (BP = 70; Figure

S1).

The median estimated crown age for the genus *Phyllotis* was 4.28 Mya with a 95% Highest

Posterior Distribution (HPD) of 3.07-5.63 Mya, a range that almost spans the entire Pliocene. Crown

ages and associated HPD's for the clades corresponding to the species groups *andium-amicus*,

osilae, and *darwini*, were 1.36 (0.66-2.28), 1.26 (0.68-2.23), and 2.83 Mya (1.98-3.72), respectively.

Within each of these three groups, most species diverged during the last ~2 Mya and there appears to

have been a pulse of speciation during the mid to late Pleistocene.

The species delimitation analyses were consistent in recognizing each of the 20 nominal forms

of *Phyllotis* represented in the full *cytb* dataset. Different delimitation approaches identified 36-37

distinct units (Figure 3). Results of the delimitation analyses suggest that *P. caprinus*, *P. chilensis*, *P.*

darwini, *P. magister*, and *P. vaccarum* may each represent complexes of multiple species. The

internal subdivisions identified within *P. caprinus* and *P. darwini*, and some of those identified within *P.*

chilensis, have allopatric distributions (Figure S2). Results of the GMYC and PTP delimitation

analyses differed in the number of units identified within *P. vaccarum* and *P. pehuenche*. The GMYC

and b-GMYC analyses identified six distinct units within *P. vaccarum* and recognized *P. pehuenche* as

a single unit. By contrast, the PTP and b-PTP implementations recognized three distinct units within

both *P. vaccarum* and *P. pehuenche*.

Levels of mitochondrial differentiation between pairs of *Phyllotis* species are highly variable, with

estimated *p*-distances ranging from 2.73% (SE = 0.004) between the sister species *P. limatus* and *P.*

vaccarum, to 17.28% (SE = 0.013) between *P. gerbilus* and *P. nogalaris* (Table 1). The mean *p*-

distance between nominal species within the genus *Phyllotis* is 7.55% (SE = 0.005). Within the

Phyllotis xanthopygus species complex, the maximum *p*-distance is 10.82% between *P. pehuenche*

and *P. chilensis* (Table 1). We also estimated *p*-distances between internal subdivisions (candidate

species) within several nominal forms that were identified as significant in the species delimitation

analyses. In these cases, pairwise *p*-distances ranged from 1.81% (SE = 0.003) between subdivisions

within *P. magister* to 9.32% (SE = 0.011) between the most divergent subdivisions within *P. chilensis*

(Table S1).

3.2 GENOMIC ASSESSMENT OF SPECIES LIMITS

To further examine species limits suggested by the analysis of *cytb* sequence variation, we generated

low-coverage WGS data for representative subsets of specimens from 11 nominal species, several of

which have overlapping ranges in the Altiplano and/or adjoining lowlands. We also derived an

alignment of whole mitochondrial genomes from the WGS data. Whereas the BI and ML analyses of

cytb variation yielded some conflicting estimates of species relationships within the *P. xanthopygus*

complex (Figures 2 and S1), the ML phylogeny estimate based on complete mitochondrial genomes

confirmed the close relationship between *P. pehuenche* and *P. xanthopygus* and placed them sister to

the clade comprising *P. caprinus*, *P. limatus*, and *P. vaccarum* (BP = 100) (Figure 4).

In a PCA of genome-wide variation, PC1, PC2, and PC3 captured 36.8%. 23.2%, and 7.15% of

the total variation, respectively (Figure 5a,b). Samples of *P. darwini* from the northern and southern

portions of the species range separated into two highly distinct clusters (Figure 5a,b). The distinct

clusters of *P. darwini* specimens identified in the genomic PCA are perfectly congruent with two

divergent mtDNA subclades that were identified as significant internal subdivisions in the species

delimitation analysis (Figure 3). Using coding sequence of the complete mitochondrial genome, the

estimated *p*-distance between the northern and southern subdivisions of *P. darwini* was 7.25% (SE =

0.002) (Table S2).

The sister species *P. limatus* and *P. vaccarum* were not readily distinguishable along first two

PC axes (Figure 5a), but they were cleanly separated along PC3 (Figure 5b). One specimen, GD2350,

which was identified as *P. limatus* on the basis of mtDNA, fell in between the two distinct clusters of *P.*

limatus and *P. vaccarum* samples in PC3 space (Figure 5b). The GD2350 specimen was collected in

the narrow zone of range overlap between *P. limatus* and *P. vaccarum* in northern Chile, within 200-

250 km of localities where *P. vaccarum* specimens were found to carry *limatus*-like mtDNA haplotypes

(Figure 5c). Individual admixture proportions estimated with NGSadmix also distinguished *P. limatus*

and *P. vaccarum* samples as genetically distinct clusters, and GD2350 was assigned approximately

equal admixture proportions of the two species (Fig. 5d).

A sliding window analysis of PC1 comprising the full sample of *P. limatus* and *P. vaccarum*

specimens revealed a mosaic patterning of variation along the genome of GD2350, as autosomal

segments alternated between three main patterns: (*i*) homozygous for *P. limatus* ancestry, (*ii*)

homozygous for *P. vaccarum* ancestry, or (*iii*) heterozygous, falling approximately halfway in between the two species (Figure 6).

3.3 REVISED GEOGRAPHIC RANGE LIMITS OF *PHYLLOTIS* SPECIES

The integrated analysis of mtDNA and WGS data enabled us to delineate the geographic range limits of several species in the Puna de Atacama and surrounding regions. The mice identified as *P. caprinus* that we collected in southern Bolivia significantly extend the species' known range to the north (Figure 7). Another possibility suggested by results of the species delimitation analysis (Figure 3) is that the northernmost Bolivian specimens do not represent extralimital records of *P. caprinus*, but may instead represent a new, undescribed species that is closely related to the form currently recognized as *P. caprinus*. In the case of *P. chilensis*, our specimens from the Chilean regions of Arica y Parinacota, Tarapacá, and Antofagasta extend the species' known range to the west (Figure 7).

Our records for *P. vaccarum* indicate that this primarily highland species is replaced by *P. darwini* at elevations <2500 m on the western slope of the Andes, but – beyond the northernmost limits of *P. darwini* – the range of *P. vaccarum* extends all the way to sea level along a narrow stretch of coastline in northern Chile (Figure 7). On the eastern slope of the Andes, our records from northwestern Argentina indicate that the species does not occur <1200 m, as it is replaced by *P. anitae* and *P. nogalaris* in lowland Yungas forests. Further south along the eastern slope of the Cordillera where humid lowland forests give way to arid steppe and Monte habitats, our lowest elevation records of *P. vaccarum* were from 765-1158 m in the Argentine provinces of Catamarca, Neuquén, and Mendoza, but the majority of records are from elevations >1200 m.

4. DISCUSSION

4.1 MOST DIVERSIFICATION OF *PHYLLOTIS* OCCURRED IN THE PLEISTOCENE

Estimating divergence times of Sigmodontine rodents has been difficult due to a lack of suitable fossils that could be used to calibrate molecular data (Salazar-Bravo et al., 2013). Previous studies placed the basal split of *Phyllotis* in the Pliocene (3.0–5.1 Mya) and the basal split of the *P. xanthopygus* species complex in the Pliocene-Pleistocene transition (1.6–2.3 Mya) using a maximum likelihood clock estimate of 7.3% divergence per Mya (Steppan et al., 2004, 2007). Riverón (2011) estimated a similar Pliocene basal split for *Phyllotis* (2.83-4.05 Mya) using an analogous strict-clock estimate. Our secondary calibration estimations suggest a similar timing of diversification of *Phyllotis*, with an estimated initial divergence 4.28 Mya (95% HPD = 3.07-5.63 Mya) and subsequent diversification of the *P. xanthopygus* complex 2.83 Mya (95% HPD = 1.99-3.72 Mya). However, the divergence time estimates should be always interpreted with caution due to uncertainty about the applied calibrations

(Steppan et al., 2007; Parham et al., 2012).

In principle, the diversification of *Phyllotis* could have been spurred by mountain uplift and/or

climate-related environmental changes at the end of the Pliocene and the beginning of the

Pleistocene. The Central Andean Plateau experienced the most significant phase of uplift in the late

Miocene-Pliocene (Gregory-Wodzicki 2000). The montane uplift hypothesis therefore predicts that

diversification of *Phyllotis* would have started well before the end of the Pliocene (2.6 Mya). It is also

possible that diversification occurred more recently, and independently of Andean uplift, during periods

of climate-induced environmental change in the Pleistocene. For example, the mid-Pleistocene

Transition (MPT; 1.25–0.70 Mya) was associated with a major shift in global climate periodicity that

produced a persistent global aridification trend (Herbert, 2023). Thus, the Pleistocene Aridification

hypothesis predicts that diversification of *Phyllotis* would have occurred more recently than the

Andean uplift, coinciding with periods of climate change that were not directly related to orogenic

events.

Using secondary calibrations for divergence time estimates, our results suggest a progressive

diversification of *Phyllotis* during the past 3 million years with divergence times for most species

coinciding with glacial cycles in the mid- to late Pleistocene (Figure 2). Basal splits in two of the three

main *Phyllotis* clades (the *andium-amicus* and *osilae* species groups) occurred prior to the MPT (0.7-

1.25 Mya), whereas the basal split within the *darwini* group is estimated to have occurred 2.1 Mya

(95% HPD = 1.56-1.86 Mya) at the Pliocene-Pleistocene boundary. Within each of the three main

clades, most diversification occurred within the past ~1.5-1.8 Mya. Thus, our results suggest that most

diversification of *Phyllotis* occurred well after the late Miocene-Pliocene phase of Andean uplift.

4.2 ALPHA DIVERSITY WITHIN THE *PHYLLOTIS DARWINI* SPECIES GROUP

Based on results of our phylogenetic reconstructions and species delimitation analyses, we can

identify at least 10 clades that are referable to traditionally recognized species within the *Phyllotis*

darwini species group (Figures 2, 3, and 4). However, results of the species delimitation analysis

clearly show that some of these nominal forms may be polytypic or could be split into more species

after further taxonomic work. There appears to be potential for the existence of cryptic species within

nominal forms that are currently recognized as *P. caprinus*, *P. chilensis*, *P. darwini*, *P. magister*, and

P. vaccarum (Figure 3).

The Bolivian specimens of *P. caprinus* from Chuquisaca (MSB237236) and Cochabamba

(MSB238568) constitute a clade with a high degree of mitochondrial differentiation relative to the

remaining Argentine specimens that are referable to *P. caprinus* as currently recognized (*p*-

distance=5.6%, SE=0.008) (Figure 3). *Phyllotis darwini* and *P. chilensis* also exhibit north-south

patterns of internal substructure (Figure S2a,b), with highly distinct units identified by the species

delimitation analyses (Figure 3). In the case of *P. darwini*, divergence between northern and southern

mtDNA clades is also apparent at the whole-genome level (Figure 5a,b). Consistent with results of

Ojeda et al. (2021), the clade that includes specimens that we refer to as *P. chilensis* appears likely to

contain multiple cryptic species with apparently allopatric distributions in Peru (Figure S2b). Although

Ojeda et al. (2021) referred to this group as the "*P. posticalis-P. rupestris*" clade, geographic

considerations of type localities suggest that "*P. chilensis*" is a more appropriate name for the

subclade with the southern-most distribution in northeastern Chile, southwestern Bolivia, and

northwestern Argentina (Hershkovitz, 1962; Mann, 1945; Thomas, 1912). Here and elsewhere (Storz

et al., 2014), we followed Mann (1945) and Pearson (1958) in using the name "*P. chilensis*" for the

mice in this subclade that we collected in Altiplano of northern Chile, southwestern Boliva, and

northwestern Argentina. For the subclade with the northern-most distribution in this group, it seems

reasonable to use the name *posticalis* because it includes a specimen from the vicinity of the

associated type locality in the Department of Junín, Peru (Thomas, 1912).

In *P. vaccarum*, one *cytb* haplogroup that was identified as a distinct unit in the species

delimitation analysis is sister to a clade formed by haplotypes of *P. limatus*. The *P. vaccarum* mice that

harbor *limatus*-like mtDNA haplotypes are not distinguishable from other *P. vaccarum* at the whole-

genome level (Storz et al. 2024). In this particular case of mitonuclear discordance, identified mtDNA

subdivisions are clearly not reflective of cryptic species within *P. vaccarum*.

4.3 EVIDENCE FOR INTERSPECIFIC HYBRIDIZATION

The genomic data revealed clear-cut evidence of ongoing hybridization between *P. limatus* and *P.*

vaccarum (Figure 5d and Figure 6), suggesting that introgression is a plausible explanation for the

sharing of mtDNA haplotypes between the two species (Figure 5c; see Storz et al., 2024). The

GD2350 specimen carries *P. limatus* mtDNA but harbors approximately equal genome-wide admixture

proportions from *P. limatus and P. vaccarum* (Figure 5d). At face value, the approximately equal

admixture proportions suggest that GD2350 could be a first generation (F1) interspecific hybrid that

has received one haploid complement of chromosomes from each parent. However, in the windowed

PCA, an F1 hybrid would be expected to continuously localize halfway between the two divergent

parental stocks. Contrary to that expectation, tracts across the genome of GD2350 were either

homozygous for *P. vaccarum* ancestry, homozygous for *P. limatus* ancestry, or heterozygous (i.e.,

combining both species' genomes)(Figure 6). The mosaic patterning of nucleotide variation appears to

reflect one or more rounds of recombination subsequent to an initial *P. limatus* x *P. vaccarum*

hybridization event and suggests that GD2350 is the product of an F2 or more advanced-stage

intercross. Given that GD2350 was assigned roughly equal admixture proportions for both species

(Figure 5c), it is likely that the zone of range overlap between *P. limatus* and *P. vaccarum* in northern

Chile represents a zone of ongoing hybridization. Although the observed pattern of genomic
mosaicism in GD2350 could have been produced by a balanced number of backcrossing events with
both parental species, we regard ongoing matings between hybrids as a more likely scenario. More
intensive collecting from the zone of range overlap between *P. limatus* and *P. vaccarum* will be
required to assess the pervasiveness of hybridization between the two species.

Aside from the evidence of hybridization and mitonuclear discordance between *P. limatus* and
P. vaccarum, which also happen to be the only pair of sister species with overlapping ranges within the
P. darwini group, all remaining *Phyllotis* specimens that grouped together in the *cytb* phylogeny were
also identified as distinct groupings in the analysis of WGS data (Figure 5a,b).

4.4 A REVISED UNDERSTANDING OF GEOGRAPHICAL RANGE LIMITS OF *PHYLLOTIS* MICE
The use of sequence data to confirm the identities of all collected specimens provided new information
about geographic range limits and revealed notable range extensions for several *Phyllotis* species
(Figure 7). The westward range extension of *P. chilensis* in northern Chile is noteworthy because only
P. limatus and *P. magister* had been previously recorded in this zone (Steppan & Ramírez, 2015:
Ojeda et al., 2021). We collected *P. chilensis* from a number of extremely high-elevation localities in
northern Chile and western Bolivia, including multiple specimens from 5221 m on the flanks of Volcán
Parinacota and 5027 m on the flanks of Volcán Acotango in western Bolivia. Such records highlight the
importance of surveying environmental extremes to accurately characterize geographic range limits,
especially for taxa like *Phyllotis* that are known to inhabit extreme southern latitudes in Patagonia,
extreme elevations in the Central Andes, and extreme arid zones in the Atacama Desert. *P. vaccarum*
was previously documented to have the broadest elevational range of any mammal, from the coastal
desert of northern Chile to the summits of >6700 m volcanoes (Storz et al., 2020, 2024). The species
has a similarly broad elevational range on the eastern slope of the Andes, but the lower range limit
depends on the nature of the low elevation biome (Jayat et al., 2021; Riverón, 2011). In northwest
Argentina, the species appears to have a lower range limit >1200 m, as it is replaced by species in the
osilae group in humid Yungas forests. In central western Argentina, *P. vaccarum* reaches elevations
<1000 m in arid Patagonian steppe and Monte habitats.

5. CONCLUSIONS
Many previous phylogenetic assessments of *Phyllotis* have been limited to existing material in
zoological collections, so geographic coverage is often quite sparse and uneven. Our intensive
collecting in the Andean Altiplano and surrounding lowlands enabled us to fill key gaps in geographic
coverage. By integrating vouchered specimen records with species identifications based on mtDNA
and WGS data, we now have a better understanding of geographic range limits for species in the *P.*

darwini group. The delimitation of genetically distinct units within several named forms indicates the presence of much undescribed alpha diversity in *Phyllotis*, as pointed out by previous authors (e.g., Ojeda et al., 2021; Jayat et al., 2021). Although much of the diversification of *Phyllotis* may have occurred in the Andean highlands, our divergence date estimates suggest that diversification of these mice was not associated with the major phase of uplift of the Central Andean Plateau in the Miocene-late Pliocene. Instead, most lineage splitting was associated with climatically induced environmental changes in the mid- to late Pleistocene.

Within the *P. xanthopygus* complex, *P. limatus* and *P. vaccarum* represent the only species for which we observed mitonuclear discordance and documented ongoing hybridization. This example demonstrates that interspecific hybridization occurs in *Phyllotis*, but more intensive collecting in zones of range overlap between species will be required to assess the pervasiveness of introgressive hybridization in the group.

AUTHOR CONTRIBUTIONS

MQ-C, GD, and JFS designed the study, MQ-C, NMB, GD, PJ, PT, and JFS performed the fieldwork, MQ-C, SL, JLM, and TM performed the laboratory work, MQ-C, SL, JLM, TM, JAC, LMB, NDH, ZAC, JMG, GD, and JFS performed data analysis and/or helped with interpretation, MQ-C and JFS wrote the initial draft of the manuscript, and all authors read and approved it.

ACKNOWLEDGMENTS

We thank Mario Pérez-Mamani and Juan Carlos Briceño for assistance and companionship in the field, and we thank José Urquizo for helpful comments and discussion.

FUNDING INFORMATION

This work was funded by grants to JFS from the National Institutes of Health (R01 HL159061), National Science Foundation (OIA-1736249 and IOS-2114465), and National Geographic Society (NGS-68495R-20) and a grant to GD from the Fondo Nacional de Desarrollo Científico y Tecnológico (Fondecyt 1221115).

ORCID

Marcial Quiroga-Carmona: 0000-0002-2321-7777

Schuyler Liphardt: 0000-0001-8370-8722

Naim M. Bautista: 0000-0003-0634-0842

Pablo Jayat: 0000-0002-6838-2987

Pablo Teta: 0000-0001-8694-0498

Jason L. Malaney: 0000-0002-3187-7652

Tabitha McFarland: 0009-0002-9211-682X

Joseph A. Cook: 0000-0003-3985-0670

Moritz Blumer: 0000-0002-5775-1767

Nathanael D. Herrera: 0000-0002-5039-8876

Zachary A. Cheviron: 0009-0006-2089-5579

Jeffrey M. Good: 0000-0003-0707-5374

Guillermo D'Elía: 0000-0001-7173-2709

Jay F. Storz: 0000-0001-5448-7924

556 REFERENCES

Aktas C (2023) haplotypes: manipulating DNA sequences and estimating unambiguous haplotype

network with statistical parsimony. R package version 1.1.3.1. https://CRAN.R-

project.org/package=haplotypes

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, Suchard MA, Rambaut A, Drummond

AJ. (2014) BEAST 2: a software platform for Bayesian evolutionary analysis. PLoS

Computational Biology, 10(4), e1003537. https://doi.org/10.1371/journal.pcbi.1003537

Cadenillas R, D'Elía G (2021) Taxonomic revision of the populations assigned to *Octodon degus*

(Hystricomorpha: Octodontidae): With the designation of a neotype for *Sciurus degus* G. I.

Molina, 1782 and the description of a new subspecies. Zool Anz 292:14–28.

https://doi.org/10.1016/j.jcz.2021.02.008

Chen S, Zhou Y, Chen Y, & Gu J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics, *34*, i884-i890.

Dellicour, S., & Flot, J. F. (2018). The hitchhiker's guide to single-locus species delimitation. Molecular

Ecology Resources, 18(6), 1234-1246.

Fujisawa T, Barraclough TG .2013. Delimiting Species Using Single-Locus Data and the Generalized

Mixed Yule Coalescent Approach: A Revised Method and Evaluation on Simulated Data Sets.

Syst Biol 62(5):707-724. https://doi.org/10.1093/sysbio/syt033

Dierckxsens, N., P. Mardulyn, and G. Smits. 2017. NOVOPlasty: de novo assembly of organelle

genomes from whole genome data. Nucleic Acids Research 45:e18.

Gregory-Wodzicki, K. M. (2000). Uplift history of the Central and Northern Andes: a review. Geological

society of America bulletin, 112(7), 1091-1105.

Herbert, T. D. (2023). The mid-Pleistocene climate transition. Annual review of earth and planetary

sciences, 51(1), 389-418.

Jayat, J. P., D'Elia, G., Pardiñas, U. F. J., & Namen, J. G. (2007). A new species of *Phyllotis* (Rodentia,

Cricetidae, Sigmodontine) from the upper montane forest of the Yungas of north-western

Argentina. In D. A. Kelt, E. P. Lessa, J. Salazar-Bravo, & J. L. Patton (eds.). The quintessential

naturalist: honoring the life and legacy of Oliver P. Pearson, (pp. 775–798). University of

California. Publications in Zoology, 134.

https://doi.org/10.1525/california/9780520098596.003.0022

Jayat, J. P., Ortiz, P. E., González, F. R., D'Elía, G. (2016). Taxonomy of the *Phyllotis osilae* species

group in Argentina; the status of the "Rata de los nogales" (*Phyllotis nogalaris* Thomas, 1921;

Rodentia: Cricetidae). Zootaxa, 4083, 397–417. https://doi.org/10.11646/zootaxa.4083.3.5

Jayat, J. P., P. Teta, A. A. Ojeda, S. J. Steppan, J. M. Osland, P. E. Ortiz, A. Novillo, C. Lanzone, and

R. A. Ojeda (2021) The *Phyllotis xanthopygus* complex (Rodentia, Cricetidae) in Central Andes,

systematics and description of a new species. Zoologica Scripta 50:689–706.

Katoh, K., and D. M. Standley (2013) MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Molecular Biology and Evolution 30:772–780.

Katoh K, Rozewicki J, Yamada KD (2017) MAFFT online service: multiple sequence alignment,

interactive sequence choice and visualization. Brief Bioinform 20(4):1160-1166.

https://doi.org/10.1093/bib/bbx108

Korneliussen, T. S., A. Albrechtsen, and R. Nielsen (2014) ANGSD: analysis of next generation

sequencing data. BMC Bioinformatics 15:356.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: molecular evolutionary genetics

analysis across computing platforms. Mol Biol Evol 35(6):1547-1549.

https://doi.org/10.1093/molbev/msy096

Kalyaanamoorthy S, Minh BQ, Wong TK, von Haeseler A, Jermiin LS (2017) ModelFinder: fast model

selection for accurate phylogenetic estimates. Nature methods 14: 587-589.

https://doi.org/10.1038/nmeth.4285

Larsson A (2014) AliView: a fast and lightweight alignment viewer and editor for large data sets.

Bioinformatics 30(22):3276-3278. https://doi.org/10.1093/bioinformatics/btu531

Li, H., and R. Durbin (2009) Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics, *25*,1754-1760.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth et al. (2009) The sequence

alignment/map format and SAMtools. *Bioinformatics*, *25*, 2078-2079.

McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernytsky, K. Garimella, et al.,

(2010) The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation

DNA sequencing data. *Genome Research*, *20*,1297-1303.

Meisner, J., and A. Albrechtsen. 2018. Inferring population structure and admixture proportions in low-

depth NGS data. Genetics 210:719–731.

Minh, B. Q., Thi Nguyen, M.A., von Haeseler, A.(2013). Ultrafast Approximation for phylogenetic
bootstrap. Molecular Biology and Evolution, 30(5), 1188–1195.

Ojeda, A. A., P. Teta, J. P. Jayat, C. Lanzone, P. Cornejo, A. Novillo, and R. A. Ojeda. 2021.
Phylogenetic relationships among cryptic species of the *Phyllotis xanthopygus* complex
(Rodentia, Cricetidae). Zoologica Scripta 50:269–281.

Pajares AJM (2013). "SIDIER: substitution and indel distances to infer evolutionary relationships."
Methods in Ecology and Evolution, 4, 1195-1200.

Parada A, Pardiñas UF, Salazar-Bravo J, D'Elía G, Palma RE. 2013. Dating an impressive Neotropical
radiation: molecular time estimates for the Sigmodontinae (Rodentia) provide insights into its
historical biogeography. Molecular Phylogenetics and Evolution 66:960–968.
https://doi.org/10.1016/j.ympev.2012.12.001

Parada A, D'Elía, G., Palma, R.E. (2015). The influence of ecological and geographical context in the
radiation of Neotropical sigmodontine rodents. BMC Evolutionary Biology 15:172.

Pardiñas, U. F. J., D'Elía, G., and Ortiz, P. E. (2002). Sigmodontinos fósiles (Rodentia, Muroidea,
Sigmodontinae) de América del Sur: estado actual de su conocimiento y prospectiva.
Mastozoología Neotropical 9(2): 209–252.

Parham JF, Donoghue PC, Bell CJ et al. (2012) Best practices for justifying fossil calibrations.
Systematic Biology, 61, 346–359

Pearson, O. P. (1958). A taxonomic revision of the rodent genus *Phyllotis*. University of California,
Publications in Zoology, 56, 391–496.

Pons J, Barraclough TG, Gómez-Zurita J, Cardoso A, Duran DP, Hazell S, Kamoun S, Sumlin WD,
Vogler AP (2006) Sequence-based species delimitation for the DNA taxonomy of undescribed
insects. Syst Biol 55(4):595-609. https://doi.org/10.1080/10635150600852011

Rambaut A, Drummond AJ .2019. TreeAnnotator v2 6.0–MCMC output analysis. Software
development. Part of Beast, 2.

Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA .2018. Posterior summarization in Bayesian
phylogenetics using Tracer 1.7. Systematic Biology 67(5): 901-904.
https://doi.org/10.1093/sysbio/syy032

R Core Team (2020) R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria

Reid, N.M., Carstens, B.C. Phylogenetic estimation error can decrease the accuracy of species
delimitation: a Bayesian implementation of the general mixed Yule-coalescent model. BMC
Evol Biol 12, 196 (2012). https://doi.org/10.1186/1471-2148-12-196

Rengifo, E. M., & Pacheco, V. (2015). Taxonomic revision of the Andean leaf–eared mouse, *Phyllotis*
andium Thomas 1912 (Rodentia: Cricetidae), with the description of a new species. Zootaxa,

651   4018, 349–380. https://doi.org/10.11646/zootaxa.4018.3.2

652 Rengifo, E. M., & Pacheco, V. (2017). Phylogenetic position of the Ancash leaf-eared mouse *Phyllotis*

653   *definitus* Osgood 1915 (Rodentia: Cricetidae). Mammalia, 82, 153–166.

654   https://doi.org/10.1515/mammalia-2016-0138

655 Rengifo, E. M., J. Brito, J. P. Jayat, R. Cairampoma, A. Novillo, N. Hurtado, I. Ferro, et al. (2022)

656   Andean non-volant small mammals: a dataset of community assemblages of non-volant small

657   mammals from the high Andes. Ecology 103:e3767.

658 Riverón, S. (2011). Estructura poblacional e historia demográfica del "pericote patagónico" *Phyllotis*

659   *xanthopygus* (Rodentia: Sigmodontinae) en Patagonia Argentina. Ph. D. Dissertation,

660   Universidad de la República, Uruguay, pp. 100.

661 Salazar-Bravo, J., Pardiñas, U. F., D'Elía, G. (2013). A phylogenetic appraisal of Sigmodontinae

662   (Rodentia, C ricetidae) with emphasis on phyllotine genera: systematics and biogeography.

663   *Zoologica Scripta*, *42*(3), 250-261.

664 Sikes RS, Gannon WL (2011) Guidelines of the American Society of Mammalogists for the use of wild

665   mammals in research. J Mammal 92(1):235-253. https://doi.org/10.1093/jmammal/gyw078

666 Skotte, L., Korneliussen, T.S., & Albrechtsen, A. (2013) Estimating individual admixture proportions

667   from next generation sequencing data. Genetics, 195(3), 693-702.

668 Smith MF, Patton JL (1993) The diversification of South American murid rodents: evidence from

669   mitochondrial DNA sequence data for the akodontine tribe. Biological Journal of the Linnean

670   Society, 50, 149–177.

671 Steppan, S. J. (1993). Phylogenetic relationships among the Phyllotini (Rodentia: Sigmodontinae)

672   using morphological characters. Journal of Mammalian Evolution, 1, 187–213.

673   https://doi.org/10.1007/BF01024707

674 Steppan, S. J. (1995). Revision of the tribe Phyllotini (Rodentia: Sigmodontinae), with a phylogenetic

675   hypothesis for the Sigmodontinae. Fieldiana Zoology, New Series, 80, 1–112.

676   https://doi.org/10.5962/bhl.title.3336

677 Steppan, S. J., Adkins, R. M., & Anderson, J. (2004) Phylogeny and divergence-date estimates of

678   rapid radiations in muroid rodents based on multiple nuclear genes. Systematic Biology,

679   53(4), 533-553. https://doi.org/10.1080/10635150490468701

680 Steppan, S. J., Ramírez, O., Banbury, J., Huchon, D., Pacheco, V., Walker, L. I., & Spotorno, A. E.

681   (2007). A molecular reappraisal of the systematics of the leaf–eared mice Phyllotis and their

682   relatives. In D. A. Kelt, E. P. Lessa, J. Salazar-Bravo, & J. L. Patton (Eds.), The Quintessential

683   Naturalist: Honoring the Life and Legacy of Oliver Pearson (pp. 799–826). University of

684   California, Publications of Zoology, California.

685 Steppan, S. J., Zawadzki, C., & Heaney, L. R. (2003). Molecular phylogeny of the endemic Philippine

rodent Apomys (Muridae) and the dynamics of diversification in an oceanic archipelago.

Biological Journal of the Linnean Society, 80(4), 699-715.

Steppan, S. J., Ramírez, O. (2015). Genus *Phyllotis* Waterhouse, 1837. In: J. L. Patton, U. F. J.

Pardiñas, & G. D'Elía (Eds.), Mammals of South America, Volume 2, Rodents (pp 535–555).

The University of Chicago Press. Chicago and London.

https://doi.org/10.7208/chicago/9780226169606.001.0001.

Steppan, S. J., T. Bowen, M. R. Bangs, M. Farson, J. F. Storz, M. Quiroga-Carmona, G. D'Elía, et al.

(2022). Evidence of a population of leaf-eared mice (*Phyllotis vaccarum*) above 6000 m in the

Andes and a survey of high-elevation mammals. Journal of Mammalogy 103:776–785.

Storz, J. F., Quiroga-Carmona, M., Opazo, J. C., Bowen, T., Farson, M., Steppan, S. J., & D'Elía, G.

(2020). Discovery of the world's highest-dwelling mammal. Proceedings of the National

Academy of Sciences of the United States of America, 117, 18169–18171.

https://doi.org/10.1073/pnas.2005265117

Storz, J. F., S. Liphardt, M. Quiroga-Carmona, N. M. Bautista, J. C. Opazo, T. B. Wheeler, G. D'Elía,

and J. M. Good. 2023. Genomic insights into the mystery of mouse mummies on the summits

of Atacama volcanoes. Current Biology 33:R1040–R1042.

Storz, J. F., Quiroga-Carmona, M., Liphardt, S., Bautista, N. M., Opazo, J. C., Rico Cernohorska, A.,

Salazar-Bravo, J., Good, J. M., & D'Elía, G. (2024). Extreme high-elevation mammal surveys

reveal unexpectedly high upper range limits of Andean mice. The American Naturalist, 203(6),

726-735. https://doi.org/10.1086/729513

Thomas, O. (1912) 1912. New bats and rodents from South America. Ann. Mag. Nat. Hist., (8), 10:

403-411.

Teta, P., Jayat, J. P., Lanzone, C., Novillo, A., Ojeda, A., & Ojeda, R. A. (2018). Geographic variation in

quantitative skull traits and systematics of southern populations of the leaf-eared mice of the

Phyllotis xanthopygus complex (Cricetidae, Phyllotini) in southern South America. Zootaxa,

4446, 68–80. https://doi.org/10.11646/zootaxa.4446.1.5

Teta, P., Jayat, J. P., Steppan, S. J., Ojeda, A. A., Ortiz, P. E., Novillo, A., ... & Ojeda, R. A. (2022).

Uncovering cryptic diversity does not end: a new species of leaf-eared mouse, genus

Phyllotis (Rodentia, Cricetidae), from Central Sierras of Argentina. Mammalia, 86(4), 393-405.

Trifinopoulos J, Nguyen LT, von Haeseler A, Minh BQ (2016) W-IQ-TREE: a fast online phylogenetic

tool for maximum likelihood analysis. Nucleic Acids Res 44: W232–W235.

https://doi.org/10.1093/nar/gkw256

Walker, L. I., Spotorno, A. E., & Arrau, J. (1984). Cytogenetic and reproductive studies of two nominal

subspecies of *Phyllotis darwini* and their experimental hybrids. Journal of Mammalogy, 65,

220–230. https://doi.org/10.2307/1381161

Wickham, H. 2016. ggplot2: elegant graphics for data analysis. 2nd ed. Springer, Cham.

Zhang J, Kapli P, Pavlidis P, Stamatakis A (2013) A general species delimitation method with

723   applications to phylogenetic placements. Bioinformatics 29(22):2869-2876.

724   https://doi.org/10.1093/bioinformatics/btt499

TABLES

Table 1. Mean *cytb p*-distances between pair of species of *Phyllotis* (below diagonal). Mean values for intraspecific *p*-distances are shown in bold on the diagonal. Standard errors (SE) for each estimate of pairwise distance is shown above the diagonal.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1. *P. amicus*	--	0.998	1.221	1.205	1.204	1.178	1.241	1.219	1.011	1.114	1.255	1.289	1.223	1.154	1.165	1.050	1.121	1.215	1.159	1.303
2. *P. andium*	11.498	**5.381**	1.193	1.193	1.195	1.082	1.075	1.061	1.111	1.125	1.049	1.260	1.138	0.920	1.234	1.112	0.559	1.286	1.120	1.134
3. *P. anitae*	14.414	12.032	**1.253**	1.335	1.237	1.355	1.166	1.347	1.398	1.387	1.259	1.083	1.078	1.156	1.482	1.322	1.224	1.071	1.366	1.343
4. *P. bonariensis*	15.855	13.615	14.052	**0.749**	0.877	0.864	1.044	1.238	1.180	0.978	0.964	1.276	1.252	1.132	0.879	1.022	1.140	1.315	0.886	0.880
5. *P. camiari*	14.657	13.048	12.119	8.514	**1.049**	0.886	1.000	1.244	1.256	1.036	0.986	1.214	1.225	1.160	1.001	1.019	1.306	1.290	0.988	0.938
6. *P. caprinus*	16.062	13.625	13.616	8.704	9.218	**4.061**	1.072	1.207	1.260	0.822	1.034	1.288	1.178	1.146	0.911	0.877	1.215	1.148	0.787	0.914
7. *P. darwini*	15.844	13.466	13.597	12.196	12.774	12.469	**3.305**	1.334	1.318	1.020	1.062	1.254	1.198	1.170	1.116	1.133	1.235	1.178	1.016	1.079
8. *P. definitus*	12.453	10.873	14.994	15.105	15.341	15.807	15.459	**0.001**	1.280	1.269	1.460	1.347	1.391	0.995	1.354	1.385	1.125	1.327	1.232	1.234
9. *P. gerbilus*	6.173	12.031	15.722	15.432	15.240	15.931	16.540	12.638	**0.274**	1.211	1.302	1.486	1.385	1.236	1.135	1.174	1.269	1.270	1.219	1.337
10. *P. limatus*	14.237	12.813	13.227	8.636	9.014	7.250	11.946	14.827	14.435	**0.512**	1.021	1.350	1.324	1.052	0.976	0.999	1.284	1.283	0.396	1.046
11. *P. magister*	14.915	12.630	13.351	10.609	10.575	11.008	10.852	14.961	15.877	9.711	**1.568**	1.139	1.082	1.176	1.030	1.002	1.125	1.135	0.976	0.956
12. *P. nogalaris*	16.105	14.206	11.259	14.232	14.157	16.030	15.114	15.698	17.284	14.566	14.328	--	1.035	1.146	1.373	1.352	1.302	1.067	1.296	1.320
13. *P. osilae*	14.723	12.551	10.068	14.082	13.900	14.708	15.349	15.721	16.283	14.043	13.301	10.205	**3.125**	1.158	1.331	1.252	1.185	0.910	1.309	1.197
14. *P. pearsoni*	12.406	9.862	13.181	14.286	14.361	15.353	14.387	7.103	12.948	13.878	13.234	15.664	14.210	--	1.214	1.118	0.952	1.202	0.986	1.158
15. *P. pehuenche*	15.874	14.086	14.730	9.277	10.602	9.128	13.325	16.252	16.280	8.894	11.256	15.689	15.516	15.499	**1.449**	0.950	1.270	1.279	0.961	1.009
16. *P. chilensis/posticalis*	15.236	14.469	14.023	9.507	9.632	9.674	12.576	16.412	15.647	9.095	10.992	14.492	15.134	14.687	10.820	**1.578**	1.180	1.172	0.980	0.997
17. *P. stenops*	11.857	4.801	11.710	13.111	12.797	13.393	13.883	11.093	12.250	12.668	11.744	14.680	11.732	10.056	14.407	14.184	**0.252**	1.289	1.266	1.246
18. *P. tucumanus*	14.591	12.660	10.032	13.836	13.182	14.074	14.370	14.571	15.768	14.041	12.413	10.189	6.811	13.962	15.467	14.241	11.909	--	1.328	1.338
19. *P. vaccarum*	15.351	13.194	14.289	8.512	9.186	7.304	12.260	15.270	15.453	2.733	10.171	14.894	14.756	13.973	9.170	9.513	13.164	14.946	**2.224**	0.981
20. *P. xanthopygus*	15.205	12.941	14.237	8.010	9.383	8.304	11.199	15.837	15.032	8.668	10.461	14.008	13.467	14.902	9.540	10.279	12.910	13.890	8.737	**0.829**

FIGURE LEGENDS

Figure 1. Distribution limits of *Phyllotis* species and geographic sampling coverage in the Central Andes and adjoining lowlands. A) Ranges of *Phyllotis* mice in the *P. darwini* species group, based on patterns of morphological and DNA marker variation (Jayat et al., 2021; Ojeda et al., 2021; Steppan & Ramírez, 2015; Storz et al., 2024). B) Distribution of 169 sampling localities, representing sites of origin for 449 *Phyllotis* specimens used in the survey of *cytb* and WGS variation.

Figure 2. Calibrated maximum clade credibility tree showing Bayesian estimates of phylogenetic relationships and divergence times within the genus *Phyllotis*. Estimates of the 95% Highest Posterior Distributions interval for the divergence times are shown for main clades. Node support is shown only for those cases in which Bayesian posterior probability values were <1. Specimens in the clade labeled '*P. vaccarum**' carry *cytb* haplotypes that group with haplotypes of *P. limatus*, even though whole-genome sequence data confirmed their identity as *P. vaccarum* (Storz et al., 2024).

Figure 3. Maximum clade credibility depicting the delimitation schemes inferred from GMYC (red bars) and PTP (blue bars). Gaps in the vertical bars denote units delimited by each method, and asterisks denote splits with support values >0.75. Continuous gray bars denote current taxonomic designations for nominal species. Terminal labels depict the haplotype classes of sequences that were retained to construct the non-redundant matrix of *cytb* haplotypes. Specimens in the clade labeled '*P. vaccarum**' carry *cytb* haplotypes that group with haplotypes of *P. limatus*, even though whole-genome sequence data confirmed their identity as *P. vaccarum* (Storz et al., 2024).

Figure 4. Maximum likelihood tree estimated from coding sequence of complete mitochondrial genomes for a set of 11 nominal *Phyllotis* species. Numbers adjacent to internal nodes denote ultrafast bootstrap support values for each clade. Within the taxon currently recognized as *P. darwini*, the species delimitation analysis identified two highly distinct subdivisions (see Figure 3). Representatives of both internal subdivisions form distinct clades in the mitogenome tree, which we labeled '*P. darwini* south' and '*P. darwini*' north.

Figure 5. Genomic variation among species of *Phyllotis* based on 137 samples representing 11 nominal species. A) Genomic principal component analysis (PCA) of genome-wide variation (PC1 vs PC2). Two distinct clusters of nominal *P. darwini* specimens, '*darwini* South' and '*darwini* North', are distinguished along the PC1 axis. B) Plot of PC1 vs PC3 separates *P. limatus* and *P. vaccarum* along the PC3 axis, and reveals a single specimen, GD2350 (designated *P. limatus* on the basis of mtDNA haplotype), that has a PC3 score intermediate between the two species. C) Map of collecting localities and distribution limits of *P. limatus* and *P. vaccarum*. GD2350 comes from a site located in a narrow zone of range overlap between the two species in northern Chile. The map also shows the distribution of mice that are identified as *P. vaccarum* on the basis of whole-genome sequence data, but which carry mtDNA haplotypes that are more closely related to those of *P. limatus* (denoted as '*P. vaccarum**' in the inset tree diagram). D) Structure plot showing clear distinction between *P. limatus* and *P. vaccarum* (*n*=20 and 51, respectively). The putative hybrid specimen, GD2350, was assigned almost exactly equal ancestry proportions from the two species.

Figure 6. Windowed PCA of a *P. vaccarum* x *P. limatus* hybrid. PC1 was computed in overlapping 1 Mbp windows along the genome for a subset of 50 *P. vaccarum* (green), 20 *P. limatus* (blue), and the putative hybrid, GD2350 (red). Mean PC1 values for each species are shown as white lines and the mean value between both species' averages is shown as a grey line. GD2350 features a mosaic genome, with its local ancestry alternating between *P. vaccarum, P. limatus*, or a point intermediate between the two species. (A) Windowed PCA of chromosomes 1-19. (B) High resolution visualization of PC 1 along chromosome 1.

Figure 7. Revised distribution limits of species in the *Phyllotis darwini* species group based on mtDNA

and WGS data. Filled circles denote collection localities that helped define geographic range limits.

'xanthopygus' complex

'darwini' group

'osilae' group

P. limatus

P. vaccarum*

P. vaccarum

P. caprinus

P. pehuenche

P. xanthopygus

P. camiari

'P. chilensis'

P. magister

P. darwini north

P. darwini south

P. nogalaris

P. anitae

0.03