Ethanol reduces grapevine water consumption by limiting transpiration

Neila Ait Kaci[1], Alice Diot[1], Beatrice Quinquiry[1], Olivier Yobregat[2], Anne Pellegrino[3], Pierre Maury[1]*, Christian Chervin[1]*

Addresses:

[1] AgroToulouse, Toulouse INP, Université de Toulouse, UMR AGIR et LRSV, BP 32607, 31326 Castanet-Tolosan, France

[2] Institut Français de la Vigne et du Vin, V'Innopôle Sud-Ouest – 1920 route de Lisle sur Tarn, 81310 Peyrole, France

[3] UMR LEPSE, University of Montpellier, INRAE, Institut Agro Montpellier, 34060 Montpellier, France

*corresponding authors: pierre.maury@toulouse-inp.fr; christian.chervin@toulouse-inp.fr

Keywords: drought acclimation, ethanol, Gamay cuttings, hormesis, *Vitis vinifera*, transpiration.

Abbreviations

DWS – Days of Water Stress

EtOH – Ethanol

FTSW – Fraction of Transpirable Soil Water

FTSWt – Fraction of Transpirable Soil Water threshold

HSPs – Heat Shock Proteins

NTR – Normalised transpiration rate

sHSPs – small Heat Shock Proteins

WS – Water Stress

WW – Well Watered

Abstract

Studies suggest that ethanol (EtOH), triggers plant adaptation to various stresses at low concentrations (10 μM to 10 mM). This study investigates whether EtOH induces drought acclimation in grapevine, as demonstrated previously in *Arabidopsis*, rice, and wheat. Preliminary results with bare root Gamay cuttings showed that those pre-treated with 10 μM EtOH aqueous solutions lost fewer leaves when deprived of water compared to controls. Subsequently, we ran a potted-cutting experiment with progressive soil water deficit. Plants pre-treated with EtOH solutions (0.4 and 250 mM) exhibited slower depletion of the fraction of transpirable soil water (FTSW), compared to controls. While 0.4 and 250 mM EtOH tended to decrease transpiration in early days, these EtOH pre-treated plants maintained higher leaf transpiration than controls after 10 days of soil water depletion. The transpiration response to FTSW was affected by EtOH application. EtOH pre-treatments limited plant leaf expansion without increasing leaf senescence, and increased root dry mass. The grapevine responses to EtOH priming followed typical hormesis curves. RNA-seq data revealed transcripts related to this EtOH priming effect. These results suggest that EtOH improves grapevine acclimation to drought, leading to potential water-savings in wine growing regions prone to high water shortages, linked to climate change.

Introduction

Many recent studies show that ethanol at physiological concentrations can have positive roles in plant development (Diot *et al*., 2024a). In particular, the work of Bashir *et al*. (2022) demonstrated that ethanol could induce drought acclimation in several plant species, including Arabidopsis, rice, and wheat. This induced acclimation is particularly interesting in the context of climate change, which is expected to increase water stress in certain geographical areas, such as the Mediterranean region (Cramer *et al*., 2018; IPCC, 2021). Given that viticulture is particularly developed in this region (OIV, 2022), we sought to test the effect of ethanol on grapevine under controlled conditions of water stress, to verify if the observations of Bashir *et al.* (2022) are applicable to this species.

Initially, experiments were conducted on grapevine cuttings with bare roots in water, with the main observation being the percentage of dried leaves one week after rehydration. Then, trials were carried out on potted grapevine cuttings, monitoring the weight of the pots

to estimate plant transpiration, as this is a commonly monitored parameter in water stress experiments (Rodrigues *et al.*, 2008; Huang *et al.*, 2020). This gravimetric method enabled the estimation of the fraction of transpirable soil water (FTSW), a variable widely used as an indicator of the soil water deficit imposed on the plant (Sinclair, 2005). Thus, the response of plant transpiration to soil drying can be characterised by the FTSW threshold, i.e. the soil water status above which the plant ecophysiological processes are sensitive to soil drying. According to the FTSWt, the eco-physiological processes showed differences in sensitivity to water deficit: leaf expansion exhibits the highest sensitivity, followed by plant transpiration, and leaf senescence (Kang *et al.*, 2024). Our innovative results about ethanol altering grapevine transpiration lead us to propose a companion paper to our RNA-seq data and results obtained regarding grapevine resistance to heat stress (Diot *et al.*, 2024b). We hope to quickly stimulate further research to find the appropriate conditions of EtOH application for the various varieties and geographical situations encountered in vineyards affected by water stress episodes.

Material and Methods

Plant material

Grapevine cuttings were made from one-year-old dormant Gamay shoots, clone 787, from a collection of genetic material whose plant material is tested to be safe for the main vine viruses (IFV Sud-ouest, Peyrole, France). The cuttings were prepared as described by Mullins (1966), with the following modifications. Single-node cuttings, 5-cm long with two cutter scars at the bottom to facilitate rooting, were disinfected with active chlorine (1.5% v/v) for one minute and rinsed three times with running water. Then, 12 cuttings were placed in 250 ml plastic crates filled with 150 ml water and covered with aluminium foil to hold the cuttings up, the lower end being immersed in water, the bud ending up upwards (Fig. 1A). The cuttings were left for one month in a growing chamber at 25°C/20°C 16h/8h (day/night) with light set to 250 $\mu mol.m^{-2}.s^{-1}$. The water level was maintained to the initial filling mark.

Two experiments were conducted, one with bare root cuttings (Exp. 1), one with potted cuttings (Exp. 2).

Ethanol treatments applied to bare-root cuttings

In Exp. 1, the water was removed and 150 ml of freshly prepared ethanol solutions at various concentrations were poured into the crates and left for 24 h to allow ethanol priming. Then, all solutions were removed and the cuttings were left without water for 38h (duration determined by preliminary trials). After this drying period, 150 ml of water were poured into each crate, and the cuttings were left to recover in the growing chamber. The percentage of dry leaves was assessed one week later.

Ethanol treatments applied to potted cuttings

In Exp. 2, the one-month cuttings (Fig. 1B) were transferred into 105 ml nursery plastic pots, filled with P.A.M.2 Proveen→ substrate (Proveen, Netherlands), and water saturated. Plants were left to settle down and develop roots for one week in a growing chamber 25°C/20°C (day/night) with a 16 h photoperiod providing 250 $\mu mol.m^{-2}.s^{-1}$ photosynthetically active radiation at the top of the plants. The plants were maintained at maximum Fraction of Transpirable Soil Water (FTSW = 1) over the week. The pots were covered with aluminium foil on the top, and an additional 3 mm polystyrene sheet (Depron→, Germany) was added onto the top to limit soil evaporation (Fig. 1D, E, F, G, J). Evaporation did not exceed 10% of the FTSW after 30 days (Supplementary Fig. S1). Then, the day prior to the beginning of the water deficit, 52.5 ml of ethanol solutions at various concentrations were poured onto the pots, after removing temporarily the bottom aluminium foil to allow run-off of the solution into a saucer. The ethanol concentrations tested in this study were 0, 0.4, 2, 10, 50 and 250 mM. These concentrations correspond to ethanol/water ratios of 0, 0.002, 0.012, 0.06, 0.29, and 1.46% (v/v), respectively. The range was chosen with 5-fold increments to minimise errors from soil variability. Pairs of plants one well-watered (WW) and one water-stressed (WS) were established for all EtOH doses. A total of eight blocks were set-up, in each block we added one pot without any plant, to measure the evaporation of the pot.

Soil water status measurements

The day following the ethanol treatment (Exp. 2), the watering of Water Stress "WS" pots was stopped, while the Well Watered "WW" pots were maintained at FTSW = 1. FTSW was calculated from pot weights, as described previously (Lebon *et al*., 2006). The daily FTSW was calculated from pot weight on d day (pot weight d), the pot weight at full soil water capacity (pot weight at fc), and the pot weight when transpiration of the non-irrigated pot was

less than 10% of WW controls of the same EtOH treatment (pot weight 10%). Prior to water deficit treatment, pots were irrigated to saturation, and after the excess water had drained, the pots were weighed to estimate the pot mass at full soil water capacity. Each pot was weighed daily, twice for the WW, before and after re-watering. $FTSW_d$ was then determined as follow (eq.1):

$$FTSW_d = (\text{pot weight d} - \text{pot weight 10\%}) / (\text{pot weight fc} - \text{pot weight}) \qquad (eq.\ 1)$$

Grapevine transpiration and leaf area measurements

Water loss for individual plants (T) over 24h (Exp. 2) was calculated from the pot weights difference, considering pot weight after irrigation (d-1 day) and before irrigation (d day) for WW. Then, transpiration rate per leaf area (Tr) was determined by dividing the daily pot weights difference by the total leaf area values measured as close as possible to the transpiration date (eq.2).

$$Tr\ (g.cm^{-2}) = (\text{pot weight d} - \text{pot weight d-1}) / \text{total leaf area} \qquad (eq.\ 2)$$

The leaf area of all plants was estimated using a regression between the longest leaf vein and the leaf area of 230 leaves (Supplementary Fig. S2). The leaf area was determined with the "easy leaf area" application (https://www.quantitative-plant.org/software/easy-leaf-area). For this purpose, leaf main vein lengths were measured twice, once around day 3 and once around day 8. The calculation of plant transpiration of days 2, 3, 4 and 5 was performed using the first set of leaf area estimates, and the transpiration of days 6, 7, 8, 9 and 10 was calculated using the second set of leaf estimates.

Normalised transpiration rate (NTR)

Daily plant transpiration values (T in $g.plant^{-1}.d^{-1}$) were normalised as described by King and Purcell (2017), according to equation 3. Briefly, a correction factor (CF) was calculated to adjust for differences in transpiration among plants within an EtOH treatment under WW conditions. The correction factor was calculated while all plants were WW (before Day 0) by dividing the water loss of individual plants by the average water loss of the WW plants at the same EtOH dose. A second correction was made to account for differences in transpiration that occurred among days after initiation of water deficit treatment. Water loss for individual plants was divided by the average water loss for WW plants of the same EtOH treatment (T_WW average). Normalised transpiration for each plant for a given weighing interval was:

163    NTR = [T /(T_WW average)]/CF   (eq. 3)

Modelling the response of normalised transpiration rate to progressive soil water deficit

166    The response of NTR at the whole plant level to FTSW was fitted according to equation

167 2 (Muchow and Sinclair, 1991), in which "a" is the model parameter describing the response

168 of NTR to FTSW, "a" characterise the curvature of the fit. As the "a" parameter increases, the

169 sensitivity of the NTR strengthens at a given same FTSW value. The FTSW threshold (FTSWt)

170 represents the FTSW at which NTR (modelled by equation 2) starts to be influenced by soil

171 water deficit. FTSWt was reached when the NTR maximum was reduced by 5%, as proposed

172 for nonlinear models by Sadras and Milroy (1996). A higher FTSWt indicates a higher sensitivity

173 to water deficit (Andrianasolo *et al.*, 2016).

174    NTR = 1/[1 + 9 x exp(a x FTSW)]   (eq. 4)

Re-watering of WS plants and experiment ending (Exp.2)

177    After preliminary tests, to find a parameter to decide the time at which the WS plants

178 will be re-watered, it was decided to choose the wilting of at least two top leaves in the canopy

179 for Exp. 2. We thought initially to choose the time when plants reached FTSW = 0, but some

180 plants showed severe wilting before reaching FTSW = 0. The plants were then re-watered with

181 30 ml the first day, then the amount of water was calculated daily to reach FTSW = 0.95. The

182 plants were harvested one week after re-watering. The WW plants and their corresponding

183 WS plants (being part of the same pair, as defined above) were harvested the same day. The

184 harvest period was spread over a week from day 28 to day 34 after the beginning of Exp. 2.

185 On the harvest day, all leaves and roots were collected and the leaf area was estimated directly

186 with Easy Leaf Area application. The percentage of dried leaves was calculated over the seven

187 oldest leaves, as at day 0 there was a minimum of seven leaves per plant (example of dry

188 leaves in Fig. 1I). Finally, the roots were dried at 60°C for at least 72h, and then weighed.

RNA-seq analyses on grapevine cell cultures

191 RNA-seq analyses have been performed on *Vitis vinifera* cv. Gamay Fréaux cell cultures. The

192 complete material and method is described in Diot *et al*. (2024b). However, the following

193 modifications in the formal analysis have been done: the relative estimated transcript counts

194 have been performed only with the two conditions used in this study (comparison between

6h_0mM and 6h_1mM) and for this analysis, the thresholds for Differentially Expressed Genes

were set at |Log2 Fold Change| > 0.5, and padj < 0.1. The raw data discussed in this publication

have been deposited in NCBI's Gene Expression Omnibus (Edgar *et al.*, 2002) and are

accessible through GEO Series accession number GSE275842

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE275842).

Statistical treatments

Data normality and homoscedasticity hypotheses were first assessed using

respectively Shapiro-Wilk tests and Levene tests (SWL tests). When SWL tests were successful,

ANOVA were used to assess the effects of water treatment, ethanol treatments, their

respective interactions on transpiration, green leaf area at harvest, leaf senescence (dried

leaves), root dry weight, FTSW threshold, and "a" parameter describing the response of NTR

to FTSW; and multi-comparisons tests were assessed with Tukey HSD tests. When SWL tests

were not successful, we used rank-based nonparametric Kruskal-Wallis tests followed by

pairwise Wilcoxon tests. We also completed these analyses with pairwise tests: Student tests

when normality and homoscedasticity were respected, otherwise, we used pairwise Wilcoxon

tests. Differences were considered significant when p < 0.05. Fits of NTR to FTSW were

performed using the non-linear least squares (NLS) regression model. The goodness of fit was

assessed by calculating the Rsquare (R^2), root mean square error (RMSE), and mean squared

error (MSE). All fitting and statistical analyses were performed with Rstudio version 4.2.1,

using the packages 'car' and 'rstatix'.

Results

Ethanol priming reduced the percentage of dried leaves in bare root cuttings subjected to a

temporary water depletion

Figure 1 shows the various stages of the culture of Gamay single-node cuttings, from

the bare root initial one-month growing stages in water and plastic crates (Fig. 1A) then

potting stages (Fig. 1B, C) and the growing chamber in environmentally controlled conditions

(Fig. 1J). A preliminary experiment (Exp.1) was conducted with the bare root cuttings, by

priming them with EtOH at various concentrations, before removal of the aqueous solutions

for 38h, then re-filling the crates with water. After an additional week, we observed the amount of dried leaves (Fig. 2). The priming with EtOH reduced the percentage of dried leaves from 35.5% (controls at 0 mM) to 17.7% (EtOH at 0.01 mM), without being significant. But, as observed with increasing EtOH doses, there was a clear trend of reduction of the dried leaves compared to controls. Instead of running more bare root trials to reach significance levels, with conditions which are not natural, we decided to run a larger trial, with cuttings in light potting soil (Exp. 2), and to follow a larger set of parameters. At least, these conditions can be met in some nurseries.

Ethanol maintained higher Fractions of Transpirable Soil Water than controls both for well watered and dried pots

The changes in FTSW over a three-week period in potted single-node cuttings (Exp. 2) were plotted for WW pots (Fig. 3A) and WS pots (Fig. 3B). FTSW of WW pots were brought up daily to FTSW of 0.95. The results show the FTSW just before re-watering the pots. Most pots remained above 0.7, regardless of the EtOH concentration used for priming, which was a one-time-24 h-treatment preceding day 0. The FTSW of WS pots gradually decreased after their last irrigation on day 0, and we observed that, in plants primed with 250 mM EtOH, the FTSW decreased more slowly than in other conditions. Interestingly, plants primed with 250 mM also kept a higher FTSW than controls for WW plants (Fig. 3A), even if it was less obvious than in WS plants. We also noticed slight differences around day 5, between WS controls and WS plants primed with 0.4 and 10 mM EtOH, with EtOH-primed cultures maintaining a higher FTSW than controls. Finally, given that there was a negligible decrease in FTSW in pots without plants due to soil evaporation (Supplementary Fig. S1), we assumed that changes in FTSW were primarily due to plant transpiration and tested this hypothesis in a series of subsequent analyses.

Ethanol slowed down the grapevine transpiration

Figure 4 shows the changes in transpiration per unit of leaf area from day 2 to day 10, a period with the biggest FTSW differences between the various treatments (Fig. 3). The full statistics regarding the Figure 4 graphs are given in Supplementary Table S1, to avoid overloading the transpiration graphs. The transpiration of WS plants started to be significantly lower than WW plants from day 6 (Fig. 4; Supplementary Table S1). It is noticeable that 250 mM EtOH reduced significantly plant transpiration, in comparison to controls, from day 4 to

day 5 in WS plants, and from day 3 to day 8 in WW plants. We also observed that, compared to the controls, the transpiration of WS plants treated with EtOH showed opposite trends at the beginning and at the end of the experiment. For example, cuttings treated with 250 mM of EtOH had a lower rate of transpiration than controls at days 4 and 5, but displayed higher transpiration than controls at days 9 and 10. Figure 4 also shows that WS plants which transpired less than controls at the beginning of the experiment, were transpiring more than controls at the end, this was particularly noticeable in 0.4 and 250 mM treated plants, thus confirming trends of smaller FTSW drop observed in Figure 3. Indeed, when pooling the transpiration results from day 2 to day 6, there was a significant transpiration decrease in the plants treated with 0.4 mM EtOH compared to controls (data not shown). We also noticed that there was no significant difference between WS plant transpiration at days 8 to 10, when testing with multiple comparisons obtained after ANOVA (Supplementary Table S1). But when testing the results with pairwise tests, Student or Wilcoxon (Supplementary Table S2), significant differences were noticed at days 8 to 10, as pre-treating with 250 mM led to higher transpiration than in plants pre-treated with 0, 2 and 50 mM EtOH.

Finally, we observed a bell-shaped distribution in the transpiration rates according to increasing EtOH doses (not taking the controls into account), particularly noticeable in WW and WS at day 10 (Supplementary Fig. S3). This is classical of a hormesis effect, which will be discussed later.

Ethanol altered the plant transpiration response to FTSW

The normalised transpiration rate (NTR) response to FTSW is shown in Figure 5. According to the values of R^2, RMSE, MSE (Fig. 5), the performance of non-linear fitting was satisfactory. NTR started at 1 when the plants transpired at the same rate as the WW plants within the same EtOH treatment, then dropped to lower values when FTSW became a limiting factor for transpiration. The FTSW threshold (FTSWt) from which NTR dropped below 0.95 was determined. We observed that priming plants with 250 mM led the plants to switch the FTSWt to higher values (FTSWt = 0.59) than plants primed with lower doses (FTSWt < 0.46) (detailed values and stats are given in Supplementary Table S3). Regarding the "a" parameter, the curvature coefficient, pairwise comparisons showed significant differences between 0.4 mM and 50 mM, compared to 250 mM. Thus, priming plants with 250 mM EtOH showed earlier regulation of transpiration in response to a progressive soil drying.

Soil water deficit and ethanol differentially impacted the leaf development and senescence

As shown in Figure 6 the soil water deficit limited the leaf expansion over the 4-week trial. At the end of the experiment, the average leaf area of the WW control plants was just above 200 cm² (Fig. 6A), when the one of the WS control plants was just above 100 cm² (Fig. 6B). The EOH priming also reduced the leaf expansion, this was particularly noticeable on the plants primed with 250 mM EtOH which were harbouring an average leaf area twice smaller than the controls (0 mM EtOH) in WW plants (Fig. 6A) and in WS plants (Fig. 6B).
The results of Figure 7A showed that, at harvest, WW plants harboured around 10% of dried leaves, while WS plants harboured around 80% of dried leaves (Fig. 7B). In none of these two series, WW and WS, the ethanol changed significantly the % of dried leaves.

Soil water deficit reduced root development

Regardless of the EtOH priming, all the root systems of WW plants reached dry weights around 500 mg (Fig. 8A), while the ones of WS plants hardly exceeded 200 mg.
Similarly to the transpiration rates (Fig. 4), the WW plants displayed a trend of hormetic response to EtOH, with a boost of root dry weight for EtOH doses around 2mM. The root system at this latter dose presented an increase in dry weight by 20% compared to the WW controls at 0 mM. This trend was also visible in WS plants (Fig. 8B), but its intensity was lowered by the water deficit. However, none of these responses were significantly different from controls.

RNA-seq analysis highlights some metabolisms potentially involved

As we ran RNA-seq experiments in parallel to this study, using a similar Gamay cultivar (Diot *et al.*, 2024b), we analysed this RNA-seq data focussing on metabolisms described by Bashir *et al*. (2022) as major causes of ethanol-induced resistance to drought in other plant species they studied. The Figure 9 shows that some genes involved in metabolisms of abscisic acid (ABA), aldehyde dehydrogenase (Ald DH), gamma-aminobutyric acid (GABA), neoglucogenesis, starch and sucrose were differentially regulated 6 hours after priming Gamay cell cultures with 1 mM ethanol. We also checked the responses of genes involved in calcium related metabolisms and found many whose expression was either up- or down-regulated by ethanol priming. But no strong trend was observed, either upward or downward in any particular metabolism. The number of calcium related genes was standing out, but this is also due to the fact that calcium related genes are more represented in Differentially

Expressed Genes (DEGs) than genes related to the metabolisms cited above. Finally, it is worth noticing that the only three sucrose related DEGs were all down regulated.

Discussion

Ethanol priming and grapevine bare root cuttings

Water is becoming a scarce resource and the risk of excessive drought linked to climate change is a serious concern for vineyard management (van Leeuwen *et al*., 2024). Our observations may provide solutions to limit water use by the grapevine, possibly by limiting transpiration, following an ethanol priming at very low doses, close to physiological values obtained in Gamay cells, 100 µM to 2 mM (Diot *et al.*, 2024b); however we did not measure endogenous EtOH in Gamay cuttings. In a preliminary experiment with bare root cuttings, we observed a better acclimation to drought induced by ethanol priming (Fig. 2), but using a non-physiological drought simulation, i.e. removing suddenly the water from bare-root cuttings and leaving the plants without water for more than 30 hours. Nevertheless, the obvious reduction in dry leaves observed in plants pre-treated with 10 µM EtOH, one week after re-watering, was noteworthy; this latter concentration corresponds to 0.0006% of ethanol in water, which is extremely low, but has been hypothesised to be in the range of detectable level of EtOH for organisms to use as a signal to adapt to stressful environments, by eliciting physiological changes (Diot *et al*. 2024a).

Ethanol reduces water use by potted cuttings

These preliminary results on bare root cuttings prompted us to carry out more physiological tests with plants in soil, in order to apply water deficit kinetics on root systems closer to vineyard conditions. The FTSW dropped below 0.1 between 10 and 15 days in most water-deprived plants (Fig. 3B), which is consistent with the dynamics reported in other studies on grapevine with different pot sizes and various climatic conditions (Lebon *et al.*, 2006). The most obvious effect of ethanol priming was the slower drop of FTSW after priming with 250 mM (Fig. 3B), a limitation of soil water use that was also observable in well-watered plants pre-treated with a similar EtOH dose (Fig. 3A).

Ethanol reduces cuttings transpiration

As pots without plants showed marginal FTSW decrease over the period of experiment (Supplementary Fig. S1), we assumed that all water loss was due to transpiration. And as it was demonstrated by Bashir *et al*. (2022) in three other plant species, the EtOH priming reduced plant transpiration in grapevine too (Fig. 4). The strongest effect was obtained with 250 mM EtOH, which corresponds to a dilution of 1.4% EtOH/water (v/v). This EtOH concentration reduced transpiration in both well-watered and water-deprived plants, as soon as two days after the priming, even if the transpiration drop became significant only after three days. Interestingly, the WS plants treated with this 250 mM concentration maintained a higher transpiration rate than WS controls (pre-treated with water alone) after 8 days of experiment This effect could be beneficial in cases of combined water deficit and heat stress, as transpiration is critical to cool down leaf temperature (Costa *et al.*, 2012). Even more interesting, we want to attract reader attention to a similar set of responses obtained with the EtOH smallest dose, 0.4 mM which corresponds to 0.002% EtOH/water (v/v). Indeed, the reduction of transpiration was also observed in the early days of experiment, but to a lesser extent than 250 mM, and where the maintained transpiration was also observed at the end of the 10 day period. Even if some of these effects are not significant in our experiment conditions, the drop of transpiration induced by 0.4 mM, compared to water stress controls (pre-treated with water alone), was noticeable from day 2 to day 4, and it was lower by up to 20%. This could generate big water savings at a vineyard scale, and definitely prompt us to test these effects at a larger scale. As in the 250 mM conditions, the 0.4 mM EtOH also maintained a higher transpiration rate than water stress controls after 8 to 10 days of experiment.

Transpiration variation follows a hormesis response

These effects obtained at low and high EtOH are logical, as EtOH is known to generate hormesis responses (Calabrese and Baldwin, 2003; Diot *et al.*, 2024a). These responses harbour a bell shaped curve as a function of increasing concentration of the priming chemical. This is quite obvious in all curves of the well-watered plants (Fig. 4). To illustrate this point, we reproduced the plots of day 10 with bell-shaped curves in Supplementary Fig. S5. The green curve highlights an increased transpiration at low ethanol doses in WW plants. Remarkedly, the water stress led to inversing the hormesis response, highlighted by the red curve in WS plants. Hormesis responses are known to be the initial parts of sigmoid curves (Kendig *et al.*,

2010), and it is known that environmental conditions can modulate hormesis responses in plants (Belz and Cedergreen, 2010). These latter showed that environmental factors, such as temperature and light, modulated the stimulation of lettuce growth by low doses of a phytotoxin, parthenin. However, further experiments would be necessary to bring more explanations about modulation of grapevine transpiration in response to a combination of EtOH priming and water stress.

The progressive soil drying affects plant transpiration; the FTSWt on plants not subjected to ethanol priming was equal to 0.42. This value is very close to 0.4, the value reported for grapevines in previous studies using the two-segment plateau regression procedure (Lebon *et al.*, 2003, Hofmann *et al.*, 2014). FTSWt values calculated using the two-segment plateau regression procedure resulting in lower values than those obtained in our study with the inverse exponential model (Kang *et al.*, 2024). The response of transpiration to FTSW is employed in various dynamic crop models to monitor vineyards water status (Lebon *et al.*, 2003; Valdés-Gomez *et al.,* 2009 ; Ramos and Casasnovas, 2014; Er-Raki *et al.,* 2021), and it is worth noting that ethanol is modulating this response. According to the "a" parameter describing the response of NTR to FTSW, priming with 250 mM EtOH showed a higher "a" than smaller doses, which indicates an earlier regulation of transpiration in response to a progressive soil drying (FTSWt = 0.42 vs 0.59 for 0 mM EtOH and 250 mM, respectively). This response is associated with a reduction of transpiration and leaf growth even under non-limiting soil water conditions. It should be concluded that high doses of ethanol reduce plant transpiration regardless of the water status of the soil, with a greater response when the soil dries out.

Ethanol effects on leaf and root developments

The hormesis effect was also observed when looking at the green leaf area at harvest, around four weeks after trial beginning (Fig. 6) with a slight stimulation of leaf area at 2 mM, regardless of the water-deficit conditions. These results also showed that high doses of ethanol (250 mM) limited the leaf development whatever the water deficit (in WW and in WS), probably due the beginning of a toxic effect of ethanol at this high concentration, already visible at 50 mM. This toxic response to exogenous ethanol has already been observed in oilseed rape by Wu *et al.* (2019). However, the water deficit induced a stronger effect than ethanol, clearly limiting leaf development in all WS cases compared to WW cases.

The observation of high percentages of dry leaves, mainly in WS plants (Fig. 7), was as

expected a classical plant response to water deficit (Fischer and Turner, 1978). The ethanol

was not significantly modulating this drought effect in WS plants, even if there was a slight

trend to increasing it at the highest dose. Because the choice of the harvest date was based

on the wilting of top leaves, and because the 250 mM treated plants showed nearly no wilting,

they were probably harvested too late. Nevertheless, the absence of ethanol effect on the

percentage of dry leaves in WW plants (Fig. 7), always below 10%, proved that we did not use

highly toxic ethanol concentrations.

The last phenotype we described in this experiment was the dry weight of the root

systems (Fig. 8). Once again, we observed a hormesis effect in the WW plants with an

increased root weight from 500 mg in controls to 600 mg in 2 mM EtOH treated plants, thus a

+20% increase in dry weight. This trend was not significant, with a t-test p-value = 0.15,

nevertheless this boosting effect of ethanol on root development has already been observed

in various plant species, such as mung bean (Bhattacharya *et al*., 1985) and oilseed rape (Wu

et al., 2019). At higher EtOH doses, above 2 mM, the dry root weight decreased to control

levels.

Regulations at the transcription level, potential roles for small heat shock proteins

In an attempt to find explanations for the differences in transpiration, we used the

RNA-seq data from a companion paper (Diot *et al*. 2024b), performed on Gamay cells, to

screen transcript accumulation in response to an exogenous ethanol application. First, we

must highlight that Gamay Freaux is genetically very close to Gamay, as it is Gamay mutant

(see cultivars numbers 4377 and 4382 in https://vivc.de/). We searched particularly for

markers of the metabolisms revealed by Bashir *et al.* (2022) as major causes of ethanol-

induced resistance to drought: ABA, Ald DH, GABA, neoglucogenesis, starch and sucrose.

However, no marker stood out as a critical consequence of the priming by ethanol, even if a

slight trend for down regulation of sucrose related genes was observed. Nevertheless, the

RNA-seq data are available through the companion paper (Diot *et al.*, 2024b) and it may be a

resource for teams looking for a particular gene or sets of genes. We also checked for

regulation of calcium related gene transcripts, as calcium involvement has been described in

stomatal closure (Huang *et al.*, 2019), but there was no tendency for up or down regulation of

calcium related genes. Diot *et al.* (2024b) observed a strong induction of small heat shock proteins (sHSPs) after ethanol priming of Gamay cells. The link between sHSPs and drought stress is known (Rizhsky *et al.*, 2002; Sarkar *et al.*, 2009) and Reddy *et al.* (2014) have observed the induction of class I and II sHSPs following drought stress in barley. Finally, it is worth noting that heat shock protein families, HSC70 and HSP90, are known to be involved in stomatal closure (Clément *et al.*, 2011). Whether small HSPs are also involved in stomatal closure would deserve further research, but our observations: EtOH priming reduced transpiration, as shown in this article, and EtOH priming induced a strong sHSPs up-regulation, as shown in the RNA-seq data (Diot *et al.*, 2024b), highlight a potential positive correlation.

Conclusion

We showed that ethanol priming induces a slowdown of the grapevine transpiration, which was obvious with the 250 mM dose, but could also be effective at much lower dose around 0.4 mM. This latter concentration corresponds to an ethanol dilution of 0.002% in water, which would be easy and economical to implement; even though the ethanol dilution of 1.4% in water, equivalent to 250 mM, is still relatively low. Confirming the promising effects of such low doses would deserve further studies with older vines, either at greenhouse or vineyard scales. Additionally, we also observed that grapevine responds to the range of ethanol concentrations in a typical hormesis way. This clearly requires more studies to better understand this type of biochemical responses, linked to chemical priming or hormesis, which are now attracting more attention from the scientific community (Sako *et al.*, 2020; Salinitro *et al.* 2021). Finally, the links between sHSPs, ethanol priming and plant resistance to drought also deserves more attention.

Supplementary data

The following supplementary data are available at JXB online.

Fig. S1. Variations of FTWS with time in pots without plants

Fig. S2. Regression between the longest leaf vein and the leaf area

Fig. S3. Hormesis effects on transpiration, leading to bell-shape curves

Table S1. Multiple comparisons on Fig. 4 data

Acknowledgments

We thank EUR TULIP and the Occitanie Region for the PhD grant to AD. We also thank M. Bouzayen and J. Pirrello for GBF team co-direction and side-funding, L. Lemonnier and D. Saint-Martin for help in cell and plant cultures.

Author contributions

NAK, AP, PM and CC conceptualised the project and designed the research. OY provided the Gamay cuttings. NAK, BQ, PM and CC performed the cuttings experimentations. AD performed RNA-seq experiments on Gamay cells. NAK, AD, PM and CC ran statistical tests. All co-authors contributed to writing and editing the manuscript.

Conflict of interest

The authors declare they have no conflict of interest.

Funding

This work was supported by Toulouse-INP which provided an ATER fellowship to NAK. The École Universitaire de Recherche TULIP-GS (ANR-18-EURE-0019), provided half of a PhD grant to AD and the other half was provided by the Occitanie Region. We also acknowledge the financial support for operating our laboratories, from INRAE, CNRS and Toulouse INP.

Data availability

The data underlying this article will be shared upon request to the corresponding authors.

Special Issue

This article is part of the Special Issue 'Grapevine'. See the companion article Diot *et al*., 2024b.

References

Andrianasolo FN, Casadebaig P, Langlade N, Debaeke P, Maury P. 2016. Effects of plant growth stage and leaf aging on the response of transpiration and photosynthesis to water deficit in sunflower. Functional Plant Biology **43**, 797-805.

Bashir K, Todaka D, Rasheed S, Matsui A, Ahmad Z, Sako K, Utsumi Y, Vu AT, Tanaka M, Takahashi S, Ishida J. 2022. Ethanol-mediated novel survival strategy against drought stress in plants. Plant and Cell Physiology **63**(9), 1181-1192.

Belz RG, Cedergreen N. 2010. Parthenin hormesis in plants depends on growth conditions. Environmental and Experimental Botany **69**, 293-301**.**

Bhattacharya S, Bhattacharya NC, Bhatnagar VB. 1985. Effect of ethanol, methanol and acetone on rooting etiolated cuttings of Vigna radiata in presence of sucrose and auxin. Annals of Botany **55**: 143-145.

Calabrese EJ, Baldwin LA. 2003. Ethanol and hormesis. Critical Reviews in Toxicology **33**, 407–424.

Clément M, Leonhardt N, Droillard MJ, Reiter I, Montillet JL, Genty B, Laurière C, Nussaume L, Noël LD. 2011. The cytosolic/nuclear HSC70 and HSP90 molecular chaperones are important for stomatal closure and modulate abscisic acid-dependent physiological responses in Arabidopsis. Plant Physiology **156,** 1481-1492.

Costa JM, Ortuño MF, Lopes CM, Chaves MM. 2012. Grapevine varieties exhibiting differences in stomatal response to water deficit. Functional Plant Biology, **39**, 179-189

Cramer W, Guiot J, Marini K, Verger A. 2018. Climate change and interconnected risks to sustainable development in the Mediterranean. Nature Climate Change **8**, 972-980.

Diot A, Groth G, Blanchet S, Chervin C. 2024a. Responses of animals and plants to physiological doses of ethanol: a molecular messenger of hypoxia? FEBS Journal **291**, 1102-1110.

Diot A, Madignier G, Maza E, Djari A, Di Valentin O, Chen Y, Blanchet S, Chervin C. 2024b. Responses of grapevine cells to physiological doses of ethanol, among which induced resistance to heat stress. Submitted to Journal of Experimental Botany (companion paper).

Edgar R, Domrachev M, Lash AE. 2002. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. Nucleic acids research **30**(1), 207-210.

Er-Raki S, Bouras E, Rodriguez JC, Watts CJ, Lizarraga-Celaya C, Chehbouni A. 2021. Parameterization of the AquaCrop model for simulating table grapes growth and water productivity in an arid region of Mexico. Agricultural Water Management **245**, 106585.

Fischer RA, Turner NC. 1978. Plant Productivity in the Arid and Semiarid Zones. Annual Reviews of Plant Biology **29**, 277-317.

Hofmann M, Lux R, Schultz HR. 2014. Constructing a framework for risk analyses of climate change effects on the water budget of differently sloped vineyards with a numeric simulation

using the Monte Carlo method coupled to a water balance model. Frontiers in Plant Science **5**, 645.

Huang S, Waadt R, Nuhkat M, Kollist H, Hedrich R, Roelfsema MRG. 2019. Calcium signals in guard cells enhance the efficiency by which abscisic acid triggers stomatal closure. New Phytologist **224**, 177-187.

Huang Z, Liu F, Bao Y, Zou Z, Jiang Z. 2020. Water stress timing effects on gas exchange and water use efficiency of grapevine leaves. Scientia Horticulturae **267**, 109346.

IPCC. 2021. Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J.B.R., Maycock, T.K. , Waterfield, T., Yelekçi, O., Yu, R. & Zhou, B. (eds.)]. Cambridge University Press.

Kang L, Debaeke P, Schoving C, Maury P. 2024. Responses of Leaf Expansion, Plant Transpiration and Leaf Senescence of Different Soybean (*Glycine max*.(L.) Merr.) Genotypes to Soil Water Deficit. Journal of Agronomy and Crop Science **210**, e12746.

Kendig EL, Le HH, Belcher SM. 2010. Defining Hormesis: Evaluation of a Complex Concentration Response Phenomenon. International Journal of Toxicology **29**, 235-246.

King CA, Purcell LC. 2017. Evaluation of methods for estimating transpiration response to soil drying for container-grown plants. Crop Science **57**, 2143–2148.

Lebon E, Dumas V, Pieri P, Schultz HR. 2003. Modelling the seasonal dynamics of the soil water balance of vineyards. Functional Plant Biology **30**, 699–710.

Lebon E, Pellegrino A, Louarn G, Lecoeur J. 2006. Branch Development Controls Leaf Area Dynamics in Grapevine (*Vitis vinifera*) Growing in Drying Soil. Annals of Botany **98**, 175–185.

Muchow RC, Sinclair TR. 1991. Water Deficit Effects on Maize Yields Modeled under Current and "Greenhouse" Climates. Agronomy Journal **83**, 1052–1059.

Mullins MG. 1966. Test-plants for investigations of the physiology of fruiting in *Vitis vinifera* L. Nature **209**, 419-420.

Naulleau A, Gary C, Prévot L, Hossard L. 2021. Evaluating strategies for adaptation to climate change in grapevine production — a systematic review. Frontiers in Plant Science **11**, 607859.

OIV. 2022. State of the World Vitivinicultural Sector. OIV Report. https://www.oiv.int/sites/default/files/documents/OIV_State_of_the_world_Vine_and_Wine_sector_in_2022_2.pdf

Ramos M, Casasnovas JM. 2014**.** Soil water variability and its influence on transpirable soil water fraction with two grape varieties under different rainfall regimes. Agriculture Ecosystems & Environment **185**, 253–262.

Reddy PS, Kavi Kishor PB, Seiler C, Kuhlmann M, Eschen-Lippold L, Lee J, Reddy MK, Sreenivasulu N. 2014. Unraveling regulation of the small heat shock proteins by the heat shock factor HvHsfB2c in barley: its implications in drought stress response and seed development. PLoS One. **9**, e89125.

Rizhsky L, Liang H, Mittler R. 2002. The combined effect of drought stress and heat shock on gene expression in tobacco. Plant Physiol. **130**, 1143-51.

Rodrigues ML, Santos TP, Rodrigues AP, de Souza CR, Lopes CM, Maroco JP, Chaves MM. 2008. Hydraulic and chemical signaling in the regulation of stomatal conductance and plant water use in field grapevines growing under deficit irrigation. Functional Plant Biology **35**, 565-579.

Sadras VO, Milroy SP. 1996. Field crops research soil-water thresholds for the responses of leaf expansion and gas exchange: a review. Field Crops Research **47**, 253-266,

Sako K, Nguyen HM, Seki M. 2020. Advances in chemical priming to enhance abiotic stress tolerance in plants. Plant and Cell Physiology **61**, 1995-2003.

Salinitro M, Mattarello G, Guardigli G, Odajiu M, Tassoni A. 2021. Induction of hormesis in plants by urban trace metal pollution. Scientific Reports **11**, 20329.

Sarkar NK, Kim YK, Grover A. 2009. Rice sHsp genes: genomic organization and expression profiling under stress and development. BMC Genomics. **10**, 393.

Sinclair TR. 2005. Theoretical analysis of soil and plant traits influencing daily plant water flux on drying soils. Agronomy Journal **97**, 1148–1152.

Valdés-Gómez H, Celette F, García de Cortázar-Atauri I, Jara-Rojas F, Ortega-Farías S, Gary C. 2009. Modelling soil water content and grapevine growth and development with the stics crop-soil model under two different water management strategies. OENO One **43**, 13–28.

Van Leeuwen C, Sgubin G, Bois B, Ollat N, Swingedouw D, Zito S, Gambetta GA. 2024. Climate change impacts and adaptations of wine production. Nature Reviews Earth & Environment **5**, 258–275.

Wu Z, Yang L, Jiang L, Zhang Z, Song H, Rong X, Han Y. 2019. Low concentration of exogenous ethanol promoted biomass and nutrient accumulation in oilseed rape (*Brassica napus L*.). Plant Signaling & Behavior **14**, 1681114.

Fig. 1. Various stages of the culture of Gamay cuttings: **A)** bare root cuttings in water; **B-C)** repotting after one month; **D-G)** variability of cuttings; **H)** measurement of leaf surface with Easy Leaf Area; **I)** example of dry leaves on the bottom at the end of the experiment; **J)** set-up in the growth chamber.

Fig. 2: Effects of ethanol (EtOH) priming on Gamay one-node cuttings with bare roots (preliminary study); One-month-old cuttings were primed with EtOH at various concentrations for 24h, then left without water for 38h, then assessed for dry leaves one week after rewatering; 4 replicates of 10 cuttings per concentration. Multiple comparisons with t-tests against controls (0).

Fig. 3. Effects of ethanol (EtOH) priming on FTSW as a function of time after « soil water deficit » initiation in Gamay one-node cuttings growing on light potting soil. **A)** well watered plants, WW, **B)** water stress plants, WS, irrigation stopped at day 1. n = 8, error bars show S.E.

Fig. 4. Effects of ethanol (EtOH) priming on transpiration (Tr) of Gamay one-node cuttings in pots, from day 2 to day 10 after the initiation of soil water deficit n = 8, error bars show S.E.; all statistical details in Supp. Table S1; multiple comparisons: capital letters to compare within WW; small caps to compare within WS; * stands for significant differences at 0.05 between WW and WS within the same EtOH dose.

Fig. 5. Response of normalised transpiration rate (NTR) at the whole plant level to the fraction of transpirable soil water (FTSW) for six ethanol priming doses 0, 0.4, 2, 10, 50 and 250 mM. The curve fitting and calculations have been performed with the R software as described in material and methods. "a" represents the curvature of the fit; RMSE stands for root mean square error; MSE stands for mean square error; Rsquared is the goodness of fit; FTSWt stands for FTSW threshold.

Fig. 6. Impact of ethanol (EtOH) priming on leaf area at harvest, n = 7 plants, error bars show SE. Multiple comparisons by Tukey HSD. Capital letters to compare within WW, small caps to compare within WS, and * to show p < 0.05 when comparing water deficit effect within a similar EtOH dose.

Fig. 7: Impact of ethanol (EtOH) priming on the % of dry leaves at harvest, n = 7 plants, error bars show SE. No significant difference due to EtOH within WW and WS by Tukey HSD, but * show a p < 0.05 in all comparisons of water deficit effect within each EtOH dose.

Fig. 8. Impact of ethanol (EtOH) priming on the root development at harvest, n = 7 plants, error bars show SE. No significant difference due to EtOH within WW and WS by Tukey HSD, but * show a p < 0.05 in all comparisons of water deficit effect within each EtOH dose.

Fig. 9. Differentially Expressed Genes linked to metabolisms related to stomatal closure, for the comparison 0 vs. 1 mM EtOH priming of Gamay cell cultures, 6 hours after treatment (see companion paper Diot *et al.*, 2024b).

Supplementary Data

Supplementary Fig. S1. Variation of FTSW over time in pots without plants, n = 8 pots, error bars show SE.

Supplementary Fig. S2. Regression between the longest vein and the leaf area, n = 233 leaves measured in 30 different Gamay one-node cuttings, dedicated to this purpose, in the same experiment and similar growing conditions.

Supplementary Fig. S3. Hormesis effects on transpiration, leading to bell-shape curves (green for WW, red for WS) as a function of increasing ethanol doses. The water stress is changing the bell-shape curve, and such an environmental effect on the hormesis response has been described by Belz and Cedergreen (2010).

Supplementary Table S1. Post-hoc multiple comparisons issued from ANOVA on Tr values (data in Fig. 4) performed with R software.

Tab 1 - Water treatment effect on NTR with EtOH treatments pooled, and EtOH treatments effect on NTR with water treatments pooled, as a function of time (02 to 10 days)

		02 days		03 days		04 days		05 days		06 days		07 days		08 days		09 days		10 days	
		Mean NTR	Wilcoxon	Mean NTR	Wilcoxon	Mean NTR	Wilcoxon	Mean NTR	Wilcoxon	Mean NTR	Tukey HSD	Mean NTR	Tukey HSD	Mean NTR	Wilcoxon	Mean NTR	Tukey HSD	Mean NTR	Wilcoxon
Water	WW	0.0748	a	0.0775	a	0.0892	a	0.0904	a	0.0570	a	0.0762	a	0.0590	a	0.0715	a	0.0702	a
	WS	0.0753	a	0.0773	a	0.0853	a	0.0817	a	0.0454	b	0.0480	b	0.0297	b	0.0296	b	0.0248	b
EtOH	0 mM	0.0751	AB	0.0816	A	0.0951	A	0.0950	A	0.0567	A	0.0645	A	0.0455	A	0.0509	A	0.0475	A
	0,4 mM	0.0693	AB	0.0732	A	0.0834	A	0.0838	A	0.0541	A	0.0683	A	0.0499	A	0.0556	A	0.0512	A
	2 mM	0.0872	A	0.0942	A	0.1058	A	0.0975	A	0.0526	A	0.0640	A	0.0465	A	0.0527	A	0.0507	A
	10 mM	0.0758	AB	0.0808	A	0.0969	A	0.0984	A	0.0528	A	0.0645	A	0.0452	A	0.0518	A	0.0509	A
	50 mM	0.0818	A	0.0786	A	0.0839	A	0.0818	AB	0.0507	A	0.0606	A	0.0417	A	0.0490	A	0.0433	A
	250 mM	0.0604	B	0.0525	B	0.0532	B	0.0551	B	0.0399	A	0.0525	A	0.0399	A	0.0481	A	0.0467	A

Tab 2 - Interaction of water and EtOH treatment effects on NTR, as a function of time (02 to 10 days)

	02 days		03 days		04 days		05 days		06 days		07 days		08 days		09 days		10 days	
Water x Ethanol	p = 0,4		p = 0,0227		p < 0,01		p = 0,016		p = 0,1		p = 0,02		p < 0,01		p < 0,01		p < 0,01	
WW		Wilcoxon		Wilcoxon		Wilcoxon		Wilcoxon		Tukey HSD		Tukey HSD		Tukey HSD		Tukey HSD		Tukey HSD
0 mM	0.0726	A	0.0786	AB	0.0934	AB	0.0949	AB	0.0609	AB	0.0813	AB	0.0638	A	0.0775	AB	0.0748	A
0,4 mM	0.0695	A	0.0748	AB	0.0860	AB	0.0884	AB	0.0574	AB	0.0784	AB	0.0619	A	0.0722	AB	0.0727	A
2 mM	0.0890	A	0.0967	A	0.1116	A	0.1111	A	0.0656	A	0.0877	A	0.0689	A	0.0831	A	0.0818	A
10 mM	0.0782	A	0.0822	AB	0.1009	A	0.1044	AB	0.0601	AB	0.0803	AB	0.0610	A	0.0734	AB	0.0743	A
50 mM	0.0821	A	0.0803	AB	0.0898	AB	0.0898	AB	0.0608	AB	0.0802	AB	0.0614	A	0.0775	AB	0.0724	A
250 mM	0.0588	A	0.0527	B	0.0531	B	0.0531	B	0.0382	B	0.0505	B	0.0380	B	0.0466	B	0.0460	A
WS																		
0 mM	0.0777	a	0.0847	a	0.0968	a	0.0951	a	0.0525	a	0.0478	a	0.0273	a	0.0243	a	0.0202	a
0,4 mM	0.0692	a	0.0719	a	0.0811	a	0.0798	a	0.0513	a	0.0596	a	0.0393	a	0.0411	a	0.0323	a
2 mM	0.0853	a	0.0913	a	0.0992	a	0.0819	a	0.0377	a	0.0370	a	0.0210	a	0.0180	a	0.0152	a
10 mM	0.0730	a	0.0792	a	0.0924	a	0.0915	a	0.0444	a	0.0465	a	0.0271	a	0.0272	a	0.0240	a
50 mM	0.0814	a	0.0770	a	0.0780	a	0.0739	a	0.0405	a	0.0411	a	0.0219	a	0.0206	a	0.0141	a
250 mM	0.0630	a	0.0521	a	0.0533	b	0.0583	a	0.0427	a	0.0556	a	0.0429	a	0.0504	a	0.0476	a

Tab 3 - Interaction of WW and WS treatment effects on NTR, as a function of time (02 to 10 days)

	02 days		03 days		04 days		05 days		06 days		07 days		08 days		09 days		10 days	
WW x WS																		
Ethanol	diff of means	p	diff of means	p	diff of means	p	diff of means	p	diff of means	p	diff of means	p	diff of means	p	diff of means	p	diff of means	p
0 mM	did not test*		0.0061	0.624	0.0034	0.818	0.0002	0.992	0.0084	0.282	0.0335	0.002	0.0365	<0.001	0.0533	<0.001	0.0546	<0.001
0,4 mM			0.0029	0.824	0.0049	0.745	0.0087	0.627	0.0061	0.449	0.0188	0.089	0.0226	0.004	0.0311	0.005	0.0404	<0.001
2 mM	*no factor showed		0.0055	0.674	0.0125	0.411	0.0292	0.104	0.0279	<0.001	0.0507	<0.001	0.0479	<0.001	0.0651	<0.001	0.0667	<0.001
10 mM	sig. effect by 2W ANOVA		0.0031	0.814	0.0085	0.574	0.0129	0.469	0.0157	0.053	0.0338	0..003	0.0353	<0.001	0.0463	<0.001	0.0503	<0.001
50 mM			0.0033	0.819	0.0119	0.483	0.0159	0.425	0.0202	0.026	0.0391	0.002	0.0394	<0.001	0.0569	<0.001	0.0583	<0.001
250 mM			0.0011	0.939	0.0001	0.996	0.0048	0.812	0.0040	0.66	0.0043	0.73	0.0048	0.581	0.0026	0.829	0.0002	0

Supplementary Table S2. Pairwise comparisons on Tr values (data in Fig. 4) performed with R software.

Day 2

WW

Ethanol Doses	0	0.4	2	10	50	250
0	/	0.786	0.257	*0.959*	0.398	0.206
0.4		/	0.216	*0.867*	0.310	0.379
2			/	*0.574*	0.666	0.056
10				/	*0.414*	*0.281*
50					/	0.040
250						/

WS

Ethanol Doses	0	0.4	2	10	50	250
0	/	0.395	0.576	0.691	0.745	0.241
0.4		/	0.157	0.660	0.127	0.421
2			/	0.351	0.763	0.122
10				/	0.411	0.347
50					/	0.059
250						/

Day 3

WW

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.759	0.286	0.780	0.897	0.024
0.4		/	0.226	0.582	0.693	0.054
2			/	0.414	0.400	0.017
10				/	0.898	0.025
50					/	0.032
250						/

WS

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.234	0.677	0.669	0.511	0.022
0.4		/	0.163	0.468	0.492	0.023
2			/	0.452	0.354	0.035
10				/	0.841	0.038
50					/	0.008
250						/

Day 4

WW

Ethanol Doses	0	0.4	2	10	50	250
0	/	0.622	0.342	0.671	0.827	0.006
0.4		/	0.200	0.411	0.811	0.014
2			/	0.610	0.312	*0.001*
10				/	0.573	0.009
50					/	0.013
250						/

WS

Ethanol Doses	0	0.4	2	10	50	250
0	/	0.013	*0.779*	0.755	0.133	0.003
0.4		/	*0.397*	0.376	0.750	0.009
2			/	*0.710*	*0.366*	*0.010*
10				/	0.334	0.022
50					/	0.037
250						/

Day 5

WW

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.698	0.440	0.642	0.769	0.011
0.4		/	0.287	0.439	0.934	0.016
2			/	0.775	0.343	0.008
10				/	0.502	0.014
50					/	0.014
250						/

WS

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.193	*0.121*	0.845	0.149	0.008
0.4		/	*0.613*	0.513	0.662	0.064
2			/	*0.383*	*0.836*	*0.432*
10				/	0.418	0.136
50					/	0.273
250						/

Day 6

WW

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.699	0.589	0.918	0.990	0.014
0.4		/	0.331	0.702	0.683	0.022
2			/	0.433	0.552	0.002
10				/	0.918	0.002
50					/	0.006
250						/

WS

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.880	0.114	0.351	0.213	0.224
0.4		/	0.159	0.444	0.279	0.316
2			/	0.501	0.789	0.597
10				/	0.711	0.849
50					/	*1.000*
250						/

Day 7

WW

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.829	0.603	0.929	0.931	0.017
0.4		/	0.440	0.844	0.884	0.021
2			/	0.418	0.498	0.002
10				/	0.984	0.001
50					/	0.005
250						/

WS

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	*0.328*	*0.336*	*0.955*	*0.950*	*0.284*
0.4		/	0.106	0.346	0.232	0.783
2			/	0.421	0.749	0.118
10				/	0.689	0.450
50					/	0.289
250						/

Day 8

WW

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.841	0.543	*0.798*	0.776	0.003
0.4		/	0.440	*0.779*	0.949	0.007
2			/	*0.279*	0.362	*0.000*
10				/	*0.950*	*0.001*
50					/	0.002
250						/

WS

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.192	0.352	0.987	0.471	0.041
0.4		/	0.069	0.238	0.111	0.740
2			/	0.433	0.896	0.009
10				/	0.546	0.080
50					/	0.021
250						/

Day 9

WW

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.696	0.686	0.729	0.997	*0.072*
0.4		/	0.387	0.904	0.679	0.032
2			/	0.373	0.674	0.006
10				/	0.703	0.008
50					/	0.018
250						/

WS

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.168	0.282	0.736	0.564	0.003
0.4		/	0.083	0.331	0.152	0.542
2			/	0.320	0.696	0.001
10				/	0.512	0.055
50					/	0.004
250						/

Day 10

WW

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	0.872	0.579	0.966	0.856	0.021
0.4		/	0.480	0.876	0.979	0.027
2			/	0.481	0.479	*0.014*
10				/	0.851	0.003
50					/	*0.073*
250						/

WS

Ethanol Doses (mM)	0	0.4	2	10	50	250
0	/	*0.382*	*0.232*	*0.955*	*0.282*	*0.011*
0.4		/	*0.232*	*0.613*	*0.181*	*0.093*
2			/	*0.456*	*0.945*	*0.010*
10				/	0.291	0.060
50					/	0.002
250						/

numbers in standard font obtained by Student test
numbers in italic font obtained by Wilcoxon test

Supplementary Table S3. Parameters describing the NTR response to FTSW variations, modulated by six ethanol doses. All fitting and calculations details are given in Material and Methods

EtOH	a mean	a group	FTSWt mean	FTSWt group	R^2
0 mM	-12.25	A	0.42	*a*	0.95
0.4 mM	-12.21	A	0.39	*a*	0.92
2 mM	-10.63	A	0.42	*a*	0.96
10 mM	-11.17	A	0.46	*a*	0.87
50 mM	-13.98	A	0.37	*a*	0.89
250 mM	-8.35	A	0.59	*a*	0.86

Groups in standard font obtained by Tukey HSD test
Groups in italic font obtained by Wilcoxon test

After anova on the "a", the effect of the treatment was not significant (p = 0.1)
After anova on the "FTSWt", the effect of the treatment was not significant (p = 0.06)
Then, we ran pairwise comparisons with other tests (see below)

coefficient "a"	0 mM	0.4 mM	2 mM	10 mM	50 mM	250 mM
0 mM	/	0.51	0.28	1.00	0.41	0.15
0.4 mM		/	0.26	0.48	0.41	0.04
2 mM			/	0.80	0.089	0.23
10 mM				/	0.21	0.23
50 mM					/	0.02
250 mM						/

numbers in standard font obtained by student test
numbers in in italic font obtained by wilcoxon test

FTSWt	0 mM	0.4 mM	2 mM	10 mM	50 mM	250 mM
0 mM	/	0.60	0.27	1.00	0.56	*0.08*
0.4 mM		/	0.26	*0.82*	0.70	*0.06*
2 mM			/	0.80	0.11	0.17
10 mM				/	0.35	0.22
50 mM					/	*0.09*
250 mM						/

numbers in standard font obtained by student test
numbers in in italic font obtained by wilcoxon test