TITLE
Host and antibiotic jointly select for greater virulence in *Staphylococcus aureus*

AUTHORS
Michelle Su*[1], Kim L. Hoang*[1], McKenna Penley[2], Michelle H. Davis[1], Jennifer D. Gresham[2,3], Levi T. Morran[2], Timothy D. Read[1]

*these authors contributed equally

AFFILIATIONS
1. Emory University School of Medicine, Division of Infectious Diseases, Atlanta, Georgia, USA
2. Department of Biology, Emory University, Atlanta, Georgia, USA
3. Science Department, Penn State Scranton, Dunmore, Pennsylvania, USA

CORRESPONDING AUTHOR
TDR: tread@emory.edu

ACKNOWLEDGEMENTS
The authors are grateful for insightful discussion with members of the Read and Morran labs, and Shaun Brinsmade for reading the manuscript. We thank Carmen Alvarez, Jason Chen, and Emily Stevens for help with experimental protocols.

FUNDING
MS was supported by the Antimicrobial Resistance and Therapeutic Discovery Training Program funded by NIAID T32 award AI106699. TDR was supported by National Institutes of Health (NIH) AI139188 and AI158452. KLH and TDR were supported by the Office of Advanced Molecular Detection, Centers for Disease Control and Prevention Cooperative Agreement Number CK22-2204 through contract 40500-050- 23234506 from the Georgia Department of Public Health. LTM was supported by a National Science Foundation Division of Environmental Biology (NSF-DEB) grant 1750553.

AUTHOR CONTRIBUTIONS
MS, LTM, and TDR conceptualized the project. MS, KLH, MP, MHD, and JDG conducted the experiments. KLH and MS analyzed the data and drafted the manuscript. All authors contributed to manuscript revision.

DATA AVAILABILITY
Raw sequences have been deposited in the NCBI Sequence Read Archive under the BioProject accession number PRJNA1162848, and phenotypic data deposited in the Figshare Repository (10.6084/m9.figshare.28745558).

CONFLICTS OF INTERESTS
The authors declare no conflicts of interests.

ABSTRACT

Widespread antibiotic usage has resulted in the rapid evolution of drug-resistant bacterial pathogens and poses significant threats to public health. Resolving how pathogens respond to antibiotics under different contexts is critical for understanding disease emergence and evolution going forward. The impact of antibiotics has been demonstrated most directly through *in vitro* pathogen passaging experiments. Independent from antibiotic selection, interactions with hosts have also altered the evolutionary trajectories and fitness landscapes of pathogens, shaping infectious disease outcomes. However, it is unclear how interactions between hosts and antibiotics impact the evolution of pathogen virulence. Here, we evolved and re-sequenced *Staphylococcus aureus,* a major bacterial pathogen, varying exposure to host and antibiotics to tease apart the contributions of these selective pressures on pathogen adaptation. After 12 passages, *S. aureus* evolving in *Caenorhabditis elegans* nematodes exposed to a sub-minimum inhibitory concentration of antibiotic (oxacillin) became highly virulent, regardless of whether the ancestral pathogen was methicillin-resistant (MRSA) or methicillin-sensitive (MSSA). Host and antibiotic exposure selected for reduced drug susceptibility in MSSA lineages while increasing MRSA total growth outside hosts. We identified mutations in genes involved in complex regulatory networks linking virulence and metabolism, including *codY*, *agr*, and *gdpP*, suggesting that rapid adaptation to infect hosts may have pleiotropic effects. In particular, MSSA populations under selection from host and antibiotic accumulated mutations in the global regulator gene *codY*, which controls biofilm formation in *S. aureus.* These populations had indeed evolved more robust biofilms—a trait linked to both virulence and antibiotic resistance—suggesting evolution of one trait can confer multiple adaptive benefits. Mutations that arose in these genes were also enriched in clinical isolates associated with systemic infections in humans. Despite evolving in similar environments, MRSA and MSSA populations—differing only in the presence of an intact accessory gene (*mecA*)—proceeded on divergent evolutionary paths, with MSSA populations exhibiting more similarities across replicate populations. Our results underscore the importance of considering the host context as a critical driver of pathogen traits like virulence and antibiotic resistance.

KEYWORDS

Antibiotic resistance, virulence evolution, experimental evolution, *Staphylococcus aureus*, pathogen adaptation

INTRODUCTION

Antibiotic resistance is a global crisis (Larsson and Flach, 2022). Pathogens that evolve resistance and are able to survive antibiotic therapy can go on to cause disease and transmit to other hosts; hence antibiotic resistance and virulence are intertwined (Geisinger and Isberg, 2017). Selective pressures like antibiotic usage and host defenses have independently been shown to alter the evolution of pathogen traits (Herren and Baym, 2022; Kim L Hoang et al., 2024; Kubinak et al., 2012; Larsson and Flach, 2022; Toprak et al., 2012). However, pathogens face multiple selective pressures in their environment, and selection by these forces can interact synergistically to alter evolutionary rates and trajectories relative to individual pressures alone (Merlo et al., 2020; Sharma et al., 2020). It is unclear how virulence evolves in the face of more than one selective pressure and whether this trait is constrained or facilitated by antibiotic resistance. These dynamics are even less understood in the early stages of disease emergence, where pathogens face a different suite of selection dynamics than pathogens having reached equilibrium (Day et al., 2022; Visher et al., 2021). Identifying conditions under which antibiotic resistance and virulence evolve in novel host populations will shed light on how diseases emerge and potential mitigation measures.

The strength of selection impacts pathogen evolution (Cisneros-Mayoral et al., 2022; Mahrt et al., 2021; Oz et al., 2014; Wistrand-Yuen et al., 2018). While antibiotic therapy administers concentrations that should kill all pathogen cells, low concentrations of antibiotics (sub-minimum inhibitory concentration, or sub-MIC) are pervasive in a variety of settings, including in natural environments (e.g., water bodies and soil) due to pollution and biological waste (Larsson and Flach, 2022). Sub-MICs allow susceptible populations to continue dividing, affording opportunities for mutations conferring greater resistance to emerge (Andersson and Hughes, 2014). Ultimately, selection by sub-MICs can lead to highly resistant pathogens (Gullberg et al., 2011; Wistrand-Yuen et al., 2018), which tend to incur less fitness costs compared to those under selection by high antibiotic concentrations (Westhoff et al., 2017). Importantly, mutations not related to antibiotic resistance can arise, such as those involved in adaptation to the growth environment (Pereira et al., 2023). Low concentrations of antibiotics can also alter expression of virulence factors *in vitro* across a broad range of pathogens (Braga et al., 2000; El-Houssaini et al., 2019; Haddadin et al., 2010; Khan et al., 2020), suggesting that they can affect virulence during infection. Sub-MIC antibiotics therefore have the potential to alter the early stages of disease emergence in host populations, particularly for those pathogens that can spend parts of their life history outside the host. Taken together, exposure to low antibiotic concentrations likely alters how pathogens interact with their hosts, shaping evolution of both virulence and antibiotic resistance. However, there is a knowledge gap in the interaction between antibiotics and pathogen evolution *in vivo* (Windels et al., 2020).

The host can exert strong selection on pathogens, especially during the period when a new organism is infected (e.g., a zoonotic transition). Infection of new host individuals tends to bottleneck pathogen populations (Bacigalupe et al., 2019; Klemm et al., 2016). These bottlenecks can confer a fitness advantage to resistant bacteria in the face of low antibiotic concentrations (McVicker et al., 2014). In addition to defenses like the immune system, the host environment differs drastically in nutrient availability compared to rich laboratory media (Windels et al., 2020). For example, *Staphylococcus aureus* passaged through macrophage cell lines had increased pathogen survival as well as resistance to antibiotics; these traits were lost when the pathogen was exposed to nutrient-rich media (Alves et al., 2024). Similarly, the human defensive peptide b-defensin 3 can maintain reduced susceptibility of *S. aureus* to the antibiotic vancomycin (Fait et al., 2023). Adapting to different niches within a host can also bring about drastic genomic changes. *Salmonella enterica* transitioning from an intestinal to a systemic lifestyle exhibited genome degradation in genes no longer necessary for inhabiting the gastrointestinal tract (Klemm et al.,

128   2016). While sub-MIC antibiotic exposure and host factors have been shown to independently
129   shape pathogen evolution, it is unclear how the collective actions of both selective pressures
130   affect the evolution of virulence and antibiotic resistance.
131
Staphylococcus aureus is an opportunistic pathogen that commonly colonizes nares and skin in
humans but can invade internal organs and blood, causing systemic infections and bacteremia
(Tsouklidis et al., 2020). In 2017, more than 119,000 bacteremia infections caused by *S. aureus*
occurred in the United States with a mortality rate of 18% (Kourtis et al., 2019). *S. aureus*
colonizing host surfaces exhibit distinct genomic signatures against those isolated during
systemic infection, indicating that the pathogen employs different strategies to adapt to different
host sites (Giulieri et al., 2022). For example, hemolytic strains tend to be more abundant than
non-hemolytic ones in murine systemic infection models, whereas the opposite occurs for wound
models (Schwan et al., 2003). Once inside the host, *S. aureus* resists host immune function by
hindering or lysing immune cells (Kwiecinski and Horswill, 2020; Thomer et al., 2016). Sub-MIC
concentrations of beta-lactam antibiotics—which disrupts bacterial cell wall synthesis—modify
expression of virulence factors in *S. aureus*, increasing the expression of alpha-hemolysin (Hodille
et al., 2017). Treatment with beta-lactam antibiotics systemically exposes *S. aureus* to low
antibiotic levels at non-target host tissues (Nix et al., 1991). As an opportunistic pathogen, *S.
aureus* can also survive for extended periods of time outside of hosts and have been isolated
from environments such as veterinary clinics and schools, as well as natural and constructed
settings such as wastewater and beaches (Loeffler et al., 2005; Roberts et al., 2013; Steadmon
et al., 2023; Thapaliya et al., 2017). Combined with the increasing prevalence and concentrations
of antibiotics in the environment, these factors likely increase opportunities for exposure of *S.
aureus* to sub-MIC antibiotics, subsequently affecting its interaction with the host during infection.
152
Experimental evolution has been used to elucidate how different selective pressures impact
pathogen evolution. *In vitro* studies have yielded insights into the phenotypes and genetic loci
generated by longer-term antibiotic selection through passaging experiments lasting hundreds of
generations (Fait et al., 2023; Long et al., 2023; Westhoff et al., 2017; Wistrand-Yuen et al., 2018).
Conversely, *in vivo* studies have focused on transmitting pathogens through individual hosts for
a single to a handful of passages (Erler et al., 2024; Higazy et al., 2024; McVicker et al., 2014).
However, few systems have been suitable to examine how antibiotics affect pathogen adaptation
within a host context. In this study, we take advantage of the suitability of a multicellular host,
Caenorhabditis elegans nematodes, to study designs of high replication across an appreciable
temporal scale. The *S. aureus* used here was originally isolated from a skin and soft tissue
infection of a human inmate (Diep et al., 2006) and thus is a novel pathogen to *C. elegans (Ekroth
et al., 2021)*. Nonetheless, *S. aureus* can kill nematodes by colonizing the host intestine and lysing
cells, inducing expression of defense genes in nematodes with roles conserved in humans
(Irazoqui et al., 2010; Sifri et al., 2003). Virulence screens in *C. elegans* using *S. aureus*
transposon libraries also showed overlapping results to other infection models (Bae et al., 2004;
Begun et al., 2005), with the degree of *S. aureus* virulence in *C. elegans* reflecting disease
severity of human infections (Wu et al., 2013, 2010). Experimentally evolving *S. aureus* in *C.
elegans* thus allows us to track the early stages of virulence and antibiotic resistance evolution in
novel host populations with the potential to identify conserved genomic regions underlying
evolved traits.
173
Here, we directly test the impact of host and sub-MIC antibiotic exposure on pathogen evolution.
Selection exerted by two forces may impede the pathogen's response to one or both forces (Merlo
et al., 2020). Adaptation may require resources to be expended toward either virulence or
antibiotic resistance, leading to a trade-off between these traits (Ferenci, 2016). Alternatively,
weaker selection from sub-MIC antibiotics may interact synergistically with hosts and facilitate the

evolution or maintenance of high virulence and antibiotic resistance. Sub-MIC antibiotics can favor
no-cost mutations (Westhoff et al., 2017), wherein pathogens can rapidly adapt without affecting
other traits (Visher et al., 2021). Because emerging pathogens tend to be far from the optimum of
the fitness landscape, we expect pleiotropic or large-effect mutations to play a role (Bomblies and
Peichel, 2022). We took advantage of the tractability of the system to determine how virulence,
fitness, and antibiotic resistance are connected and how multiple selective pressures shape the
evolutionary trajectory of two *S. aureus* isolates differing only in their antibiotic susceptibility.
186
Carriage of Staphylococcal cassette chromosome *mec* (SCC*mec*), which encodes *mecA*, an
accessory gene that provides resistance against beta-lactam antibiotics, is a major mechanism
for existing antibiotic resistance in *S. aureus* (Shore and Coleman, 2013). We passaged two *S.*
aureus isogenic strains, one with existing resistance (*mecA+*, MRSA) and one with a transposon
insertion in *mecA* (*mecA-*, MSSA) under selection by *C. elegans* and a sub-MIC level of oxacillin,
a beta-lactam antibiotic that has replaced methicillin as a therapy for Staphylococcal infections.
We selected for virulence in both strains to determine whether antibiotic resistance can hinder the
evolution of greater virulence. We then quantified the ability of evolved pathogens to kill hosts as
the metric for virulence. We also assessed whether populations maintained their ability to
hemolyze red blood cells—an indicator of virulence expression in both humans and nematodes
(Sifri et al., 2003). Because growth outside the host is important for transmission of opportunistic
pathogens, we quantified the *in vitro* growth and MIC of evolved pathogens. Finally, we identified
mutations potentially underlying evolved traits and compared them against a database of over
80,000 *S. aureus* genomes to ascertain whether these mutations may be associated with
adaptation to different human host sites.
202
RESULTS
Host and sub-MIC antibiotic exposure selected for greater virulence in both MRSA and
MSSA.
We experimentally evolved two *S. aureus* isogenic variants of USA300 JE2 (Diep et al., 2006),
MRSA and MSSA, with or without a host and sub-MIC antibiotic exposure. The ancestral MRSA
and MSSA isolates did not differ in terms of total growth without oxacillin, but exhibited a slight
decline in sub-MIC oxacillin (Figures 1A and 1B). For each ancestor, we passaged six
independently evolving populations under each condition 12 times, for a total of 48 evolved *S.*
aureus populations (Figure 1C).
212

214
Figure 1. Experimental evolution design. A) Growth curves and B) total growth of ancestral
MRSA and MSSA populations *in vitro* with and without sub-MIC oxacillin. Different letters
indicate significant differences (χ^2_1 = 6.39, P = 0.01). C) MRSA and MSSA were passaged 12
times with or without hosts, in the presence or absence of a sub-MIC of the antibiotic oxacillin.
Each treatment consisted of six independently evolving replicate populations. Experimental
evolution treatment abbreviations are indicated in the purple and yellow boxes. Error bars
indicate standard errors.
222

To assess the changes in virulence of evolved *S. aureus*, we measured the mortality of *C. elegans*
infected with evolved pathogens. While the MRSA and MSSA ancestors caused similar levels of
host mortality (dotted and dashed lines in Figure 2A), there was a significant treatment effect for
evolved pathogens (Figure 2A). For the MRSA genotype, host and oxacillin exposure selected for
the greatest virulence. Conversely, pathogens evolved in the absence of either pressure exhibited
attenuated virulence. These populations also had significantly greater variance compared to those
under selection from host and oxacillin, and those under solely host selection. For MSSA, host
and oxacillin exposure similarly favored greater virulence over the other three conditions. There
was no difference between variances.
232
We also characterized the hemolytic activity of evolved populations, which correlates with the
ability to secrete extracellular toxins and virulence (Cheung et al., 2012). At least one population
from most treatments had lost the ancestral ability to hemolyze sheep's blood (Figure 2B). Sub-
MIC oxacillin maintained hemolytic activity, suggesting that constant exposure to low
concentrations of oxacillin favored retention of extracellular toxicity. While hemolysis was not
necessary for increased host-killing, this ability was more often found in populations causing
greater nematode mortality (Figure 2C). As virulence was still heightened in pathogens unable to
destroy red blood cells (e.g., MSSA pathogens evolving without host or oxacillin caused greater
than 90% mean mortality despite none being hemolysis-positive), other virulence factors may be
compensating for the absence of hemolysis.
243
Natural populations of *S. aureus* exhibit variation in hemolytic ability (King et al., 2016), even
within an individual host (McAdam et al., 2011). We thus hypothesized that a significant selective
pressure like oxacillin would favor little variation of this trait within a population. We determined
the hemolysis status of isolates from each evolved population (10 isolates x 48 populations)—
most populations were in consensus (Figure S1). Furthermore, oxacillin still has a significant
effect, where hemolysis was maintained when under sub-MIC exposure selection (Figure 2D).
Oxacillin favored less hemolytic diversity in MSSA populations, further demonstrating the
interaction between antibiotics and virulence, and that antibiotics can shape variation in virulence
traits.

254
Figure 2. Evolution of virulence is facilitated by exposure to both host and sub-MIC
antibiotic. A) Virulence in terms of *C. elegans* mortality. Dashed and dotted lines indicate
respective ancestral virulence. Shaded areas indicate standard errors of technical replicates of
ancestral virulence. Different letters indicate significant differences (χ^2_7 = 460.43, P < 0.001.
MRSA Levene's test for homogeneity of variance: $F_{3,20}$ = 4.06, P = 0.021. MSSA: $F_{3,20}$ = 0.99, P
= 0.417). B) Virulence in terms of the ability to hemolyze sheep's blood, assayed at the
population level (oxacillin: χ^2_1 = 5.82, P = 0.016). The y-axis indicates the number of evolved
populations for each category. C). Host mortality from A) grouped by hemolysis status in B)
(Kruskal-Wallis χ^2_1 = 3.97, P = 0.046). D) The proportion of colonies sampled from each
evolved population that are able to hemolyze sheep's blood (oxacillin: χ^2_1 = 23.30, P < 0.001.

Levene's test for homogeneity of variance: $F_{1,22}$ = 26.06, P < 0.001). Error bars indicate standard
errors. *P < 0.05
267
268
Host and sub-MIC antibiotic synergistically promoted growth of MRSA outside hosts and
reduced drug susceptibility in MSSA.
We measured the *in vitro* growth of evolved populations to evaluate how pathogen fitness outside
the host had been impacted. We used rich media to replicate the conditions under which *S. aureus*
evolved during the experiment. Importantly, rich media reduced the risk of introducing additional
selective pressures than those being tested. In media without oxacillin, there were no significant
differences in total growth between treatments (Figure 3A). However, in sub-MIC oxacillin, MRSA
populations under selection from both pressures exhibited the greatest growth (Figure 3B), with
some achieving more growth than those in the absence of oxacillin. By contrast, exposure to sub-
MIC oxacillin alone yielded the lowest growth, suggesting a fitness cost. Similarly, MSSA
populations exposed to host and sub-MIC oxacillin exhibited a moderate increase in growth
compared to other combinations of selective pressures.
281
We then determined the MICs of evolved populations to ascertain how host and sub-MIC
exposure affected the evolution and maintenance of antibiotic resistance. For the MRSA
genotype, all but two populations retained the ancestral level of oxacillin resistance (16ug/mL
oxacillin; Figure 3C). During evolution in hosts without antibiotic selection, two populations lost
their resistance and exhibited similar susceptibility as the MSSA ancestor (0.25ug/mL). For the
MSSA genotype, host and oxacillin exposure selected for decreased antibiotic sensitivity, up to
8-fold the ancestral MIC. These results suggested sub-MIC exposure combined with host factors
potentiated the increase in antibiotic resistance in MSSA. One population under selection from
solely the host also had an increased MIC, supporting previous evidence showing non-antibiotic
selective pressures, such as a host in our study, can select for reduced antibiotic susceptibility
(Koch et al., 2014). There was no variation in antibiotic resistance within MRSA and little variation
within MSSA across the most genetically diverse populations (Figure S2).
294

295
Figure 3. **Host and sub-MIC antibiotic selection facilitated pathogen growth in antibiotics**.
Pathogen *in vitro* growth A) without oxacillin (χ^2_7 = 9.24, P = 0.24) and B) in sub-MIC oxacillin
(χ^2_7 = 106.16, P < 0.001). Different letters indicate significant differences. C). Oxacillin MIC of
evolved populations (Fisher's exact test P < 0.001). The y-axis indicates the number of evolved
populations for each category. Error bars indicate standard errors. **P < 0.01

Parallel evolution of regulators of virulence and antibiotic resistance across evolved
populations.
We conducted whole-genome sequencing of populations to identify mutations arisen from host
and antibiotic selection. Below we focus on mutations that had swept to fixation (Figure 4A, Table
S1); Figures S3 and S4 summarize mutations that were below 100% frequency in each
population. Populations evolved from the MSSA ancestor had significantly fewer mutations
(excluding those intergenic or synonymous) than MRSA populations (Figure S5; χ^2_1 = 5.28, P =
0.022), potentially due to the slightly reduced growth of MSSA in the presence of oxacillin (Figure
1B). Sub-MIC oxacillin selection also resulted in more mutations than in its absence (χ^2_1 = 5.92,
P = 0.015), although this is likely driven by MRSA populations. While this result is consistent with
the role of antibiotics in increasing mutation rates in bacteria (Revitt-Mills and Robinson, 2020),
there were only two mutations in DNA and mismatch repair genes (*mutL* and *recA*), suggesting
repair genes were not the sole mechanism involved. Of the 32 indels detected, 19 were in host-
associated populations. Across all populations, many mutations occurred in the same gene (e.g.,
agr, *gdpP*, *graSR*, *pbpA*, and *saeRS*) but not the same amino acid (Table S2), indicating parallel
evolution at the gene level.

Between pathogens with the highest virulence—those under selection from both selective
pressures—populations arisen from MRSA had mutations in 25 genes and intergenic regions,
while those from MSSA had mutations in 12 genes and intergenic regions. Both treatments had

mutations in virulence regulators *codY* (Brinsmade, 2017) and *saeRS* (Montgomery et al., 2010)
suggesting some shared pathways to heightened virulence among the two treatments, but
mutations in these genes did not occur across all replicate populations. These treatments also
had mutations in *gdpP* and *pbpA*, which have been implicated in resistance to beta-lactam
antibiotics (Bilyk et al., 2022; Poon et al., 2022), with *gdpP* playing a role in MSSA resistance to
oxacillin in particular (Giulieri et al., 2020). Five out of six populations evolved from the MSSA
ancestor had a mutation in either gene, which may have contributed to the reduced oxacillin
sensitivity in these populations.
330
Regulatory genes were enriched with mutations (Figure 4A; one-sample Poisson test P < 0.001
for MRSA and MSSA genotypes). These included *agr*, *saeRS,* and *codY*. *agr* is a quorum sensor
that regulates the expression of virulence factors (Novick, 2003). All 22 mutations in *agr* were
found in populations that have lost the ability to hemolyze red blood cells. In contrast to the other
two regulators, *codY* is a transcriptional repressor found in gram-positive bacteria that controls
virulence by monitoring nutrient levels in the environment (Brinsmade, 2017).
337
Nine populations experienced large-scale deletions that included the entire SCC*mec* cassette.
Seven of the nine populations were host-associated. Deletion of ACME has been shown to
enhance the competitive fitness of *S. aureus* USA300 *in vivo* (Diep et al., 2008). The two MRSA
populations that lost their resistance (Figure 3C) had SCC*mec* deletions, suggesting that
resistance could be more costly when evolving in a host. The loss of SCC*mec* and ACME was
more often identified in populations exhibiting an increase in total growth from the ancestor
outside the host (Figure 4B).

A

REGULATORY GENE
IMPLICATED IN VIRULENCE
IMPLICATED IN ANTIBIOTIC RESISTANCE

NUMBER OF POPULATIONS

Tw

Extra

BC tra

murP
Na+/proline symporter
peptide ABC transporter substrate–binding protein
teichoic acid ABC transporter ATP–binding protein

Cellular maintenance   N–acetylmuramoyl–L–alanine amidase
nusA
recA
rpoB
Mobile genetic element   ACME and SCCmec
SCCmec

+host +ox
+host -ox
-host +ox
-host -ox
+host +ox
+host -ox
-host +ox
-host -ox

MRSA   MSSA

348
Figure 4. Regulatory genes likely played an important role in pathogen adaptation. A)
Mutations swept to fixation, excluding intergenic and synonymous mutations, grouped by
general function. The size of each point indicates how many populations had acquired at least
one mutation in the gene. Colored shapes next to genes indicate whether these genes are
regulatory or have been implicated in virulence or antibiotic resistance in the literature (see
Table S1). B) *In vitro* growth of populations with SCC*mec* and ACME deletions with or without
sub-MIC oxacillin (one-sample t-test t = 2.73, df = 8, P = 0.026). C) Change from ancestral

pathogen-induced mortality *vs*. mutations (*codY* one-sample t-test = 19.56, df = 7, P < 0.001;
gdpP: 17.02, df = 3, P = 0.002; *pbpA*: 7.54, df = 3, P = 0.012). D) Hemolysis status *vs*.
mutations (Fisher's exact test P < 0.001; *agr vs. alr*: P = 0.036; *argR*: P = 0.006; *codY*: P =
0.005; *gdpP*: P = 0.006; *purR*: P = 0.006). E) Oxacillin MIC *vs*. mutations (Fisher's exact test P
= 0.0015; *codY vs. brnQ1*: P = 0.043). F) Biofilm production of evolved populations. Different
letters indicate significant differences (MRSA: χ^2_3 = 7.24, P = 0.06. MSSA: χ^2_3 = 17.91, P <
0.001). The column widths in D) and E) corresponds to the number of mutations. Error bars
indicate standard errors. All evolved populations were sequenced except for one population
364   from the -host-ox treatment. *P < 0.05, **P < 0.01

Mutations in antibiotic resistance genes were regularly found in pathogen populations with
heightened virulence.
We identified mutations arising in specific genes that appeared more often than by chance in
measured traits to pinpoint loci potentially underlying evolved phenotypes. We focused on genes
where two or more mutations (excluding intergenic or synonymous mutations) had fixed during
the experiment regardless of treatment (Figure 4C). Mutations in three genes were regularly
identified in populations exhibiting significant increases in virulence from the ancestor: *codY*,
gdpP, and *pbpA*. Mutations in *agr* in general were not associated with changes in overall
virulence, but MSSA populations harboring mutations in this gene were more likely to exhibit
greater virulence compared to MRSA populations (Wilcoxon rank sum exact test P = 0.045).

Mutations in specific genes were often found in populations able to hemolyze red blood cells
(Figure 4D). There was a greater proportion of populations with *agr* mutations unable to hemolyze
red blood cells compared to *alr, argR, codY, gdpP,* and *purR*. There were also significant
differences between the mutations regularly identified in oxacillin-resistant populations evolved
from the MSSA ancestor (Figure 4E), where a greater proportion of populations with mutations in
codY had reduced oxacillin susceptibility compared to *brnQ1*. Exposure to oxacillin maintained
hemolysis in evolved populations (Figure 2B), suggesting low concentrations of oxacillin exerted
negative selection on the *agr* locus. By contrast, mutations in *agr* were often in populations
exhibiting loss of hemolytic activity, consistent with previous findings (Schwan et al., 2003).

Because mutations in *codY* appeared to be important for both virulence (Figure 4C) and antibiotic
resistance (Figure 4E), we hypothesized that traits controlled by *codY* were responsible for the
traits observed in MSSA populations under selection from both pressures. A potential mechanism
underlying changes in virulence and antibiotic sensitivity in MSSA may involve biofilm formation.
Biofilm is implicated in *S. aureus* pathogenesis as well as in dampening the efficacy of antibiotic
treatment (Long et al., 2023). In *C. elegans*, *S. aureus* biofilm enhances virulence and protects
the pathogen from host innate immune defenses (Begun et al., 2007). Both sub-MIC level of beta-
lactam antibiotics and null-mutants of *codY* induce robust biofilm formation and hemolysis activity
(Chen et al., 2021; Kuroda et al., 2007; Majerczyk et al., 2008). Taken together, exposure to sub-
MIC level of oxacillin and acquisition of potentially deleterious mutations in *codY* may underlie
both increased virulence and reduced antibiotic susceptibility. Our findings partially supported this
hypothesis: biofilm formation did not significantly differ for the MRSA genotype. By contrast,
biofilm production differed between treatments for the MSSA genotype (Figure 4F), with
populations exposed to both selective pressures forming more biofilm than those evolved without
host or either pressure, but not those evolved in solely sub-MIC oxacillin. While host and sub-
MIC oxacillin selection favored robust biofilms in MSSA, oxacillin by itself also increased biofilm
formation, suggesting low antibiotic concentrations contributed significantly to the evolution of
drug-sensitive populations. The loci underlying biofilm formation may be different in these two
treatments, since mutations in *codY* did not appear in populations under solely oxacillin selection
(biofilm is a polygenic trait in *S. aureus* (Zapotoczna et al., 2016)).

Mutations that arose during experimental evolution are regularly found in strains associated with human systemic infections.

We then determined whether mutations that arose in our experiment exist in natural *S. aureus* isolates. We conducted BLAST searches against a dataset of 83,383 high quality public *S. aureus* whole genome assemblies (Raghuram et al., 2024) (Figure 5A) and found matches against the majority of our mutations. We hypothesized that these mutations may be important for *S. aureus* adaptation to different environments. Indeed, for many genes implemented in virulence and antibiotic resistance, a greater proportion of natural isolates containing our mutations were found in blood and systemic infections compared to skin/nose/throat colonization than expected (Figure 5, statistics shown in Table S3). Mutations arisen in MRSA populations under selection from both host and sub-MIC oxacillin had the most matches to natural isolates (Figure 5B), further highlighting the importance of the interaction between these two selective forces. Overall, our results demonstrate that evolution experiments in an invertebrate model can give rise to clinically-relevant mutations in a major bacterial pathogen.

Figure 5. **Mutations that arose during experiment are enriched in isolates associated with systemic infection in humans**. A) Number of genomes in our public *S. aureus* genome dataset (see Methods) grouped by isolation source. General host-association refers to descriptions not specific enough to assign to other categories. We excluded samples that were

430   ambiguously specified or missing information. B) to H) Number of genomes in the database
431 containing mutations in the respective gene. I) Number of genomes in the dataset containing the
432   mutations arisen in *agr*. This gene is separate from the others to facilitate ease of visualization
433   due to the magnitude of the y-axis. Asterisks indicate significant difference in the proportion of
434 blood/systemic-associated genomes compared to the expected distribution in the dataset. *P <
435   0.05, **P < 0.01, ***P < 0.001
436

MRSA and MSSA exhibited divergent phenotypic and genomic evolution during adaptation

We conducted principal component analysis on the evolved traits (virulence, growth in media with
or without sub-MIC oxacillin, and biofilm production) in MRSA and MSSA populations to determine
the impact of selective pressures on overall pathogen phenotypic evolution (Figures 6A and 6B).
We found a significant effect of treatment for the MRSA genotype, where populations exposed to
host and sub-MIC oxacillin clustered together, largely separating from all other treatments (P <
0.05 for all pairwise comparisons). The MSSA genotype had a marginally non-significant effect of
treatment. Populations evolved from the MRSA ancestor had similar trajectories for growth in sub-
MIC oxacillin and virulence, whereas biofilm production and virulence were more correlated for
populations evolved from the MSSA ancestor. For MRSA populations, biofilm production and
growth without oxacillin also appeared to be positively correlated.

To determine how host and sub-MIC antibiotic shaped pathogen genomic evolution, we built
phylogenies using mutations that arose from the MRSA and MSSA ancestors (Figures 6C and
D). We compared Euclidean distances calculated from the frequency of each mutation to
determine the genetic distance from the ancestor—how fast each population was evolving—and
the genetic distance between populations within each treatment—how much each population had
diverged from one another (Betts et al., 2018; Ford et al., 2017). There was no significant
difference between treatments for MRSA populations in terms of genetic distance to the ancestor
(Figure S6A), suggesting that populations generally exhibited similar evolutionary rates. By
contrast, MSSA populations had a significant effect of treatment (Figure S6B), where populations
under selection from solely the host had greater genetic distance from the ancestor than
populations under selection from solely oxacillin (Tukey's post-hoc test P < 0.001) or under neither
selective pressure (P = 0.004).

For the MSSA group, populations within each treatment tended to exhibit more genetic similarity
(Figures 6C and 6D). For MRSA, treatments varied in how divergent populations were (Figure
S6C), where populations exposed to both selective pressures diverged more from each other
than populations in the other three treatments (Dunn's post-hoc test P < 0.05 for all pairwise
comparisons). Likewise, MSSA populations also differed in how much they varied (Figure S6D).
Populations under solely host selection had greater divergence from each other than all other
treatments (Tukey's post-hoc test P < 0.001 for all pairwise comparisons). By contrast,
populations under selection solely from sub-MIC oxacillin diverged the least (P < 0.05 for all
pairwise comparisons). These findings altogether suggest that even concentrations well below
the MIC differentially impacts sensitive and resistant pathogens. As MRSA and MSSA only
differed in the presence of an intact *mecA* gene at the start of the experiment, accessory genes
may play important roles in shaping bacterial evolution (Jackson et al., 2011).

Figure 6. **MRSA and MSSA underwent distinct evolutionary trajectories.** Principal component analysis of traits evolved from A) MRSA ancestors (PERMANOVA $F_{(3,23)}$ = 8.38, r^2 = 0.557, P = 0.001) and B) MSSA ancestors (PERMANOVA $F_{(3,23)}$ = 2.20, r^2 = 0.248, P = 0.064). Phylogenetic tree constructed from frequencies of mutations in populations evolving from C) MRSA ancestors (distance to ancestor: $F_{(3,20)}$ = 2.39, P: 0.099; distance between populations: Kruskal-Wallis χ^2_3 = 21.09, P < 0.001) and D) MSSA ancestors (distance to ancestor: $F_{(3,19)}$ = 10.44, P < 0.001; distance between populations: $F_{(3,51)}$ = 54.20, P < 0.001). Scale bar indicates Euclidean distance.

DISCUSSION

While antibiotics have existed for millions of years, the mass production and overuse of antibiotics by humans in the last century have exerted an unprecedented level of selection on pathogens (Spagnolo et al., 2021; Wright, 2007). In addition to antibiotics, pathogens face numerous selective pressures such as host defenses and competitors (Bashey, 2015; Cobey, 2014; Kim L Hoang et al., 2024). Opportunistic pathogens are especially likely to encounter diverse selective pressures due to their ability to proliferate outside a host. In this study, we evolved *S. aureus* with or without pre-existing antibiotic resistance under different combinations of host and sub-MIC antibiotics to explore the evolution of virulence and antibiotic resistance. We determined whether increased virulence trades-off with antibiotic resistance by directly selecting for virulence across all host-associated treatments. Nonetheless, we observed differing evolutionary trajectories, where exposure to oxacillin in host-associated treatments resulted in pathogens causing the greatest host mortality. Our experimental design allowed us to tease apart the contributions of the host, sub-MIC antibiotics, and their interaction to demonstrate that adaptation to novel hosts did

not require trade-offs in pathogen traits. Indeed, selection from both host and antibiotics actually
resulted in rapid pathogen adaptation, potentially creating the conditions under which a pathogen
may emerge in new hosts.

Selection from the host and sub-MIC antibiotic favored the greatest virulence and growth in
antibiotics, indicating that interactions between these pressures can shape both traits. Sub-MIC
antibiotics can facilitate adaptation to high drug concentrations and to the growth environment
(Andersson and Hughes, 2014; Pereira et al., 2023). Frequent exposure to low concentrations of
antibiotics in *S. aureus* may occur due to its ability to grow and survive in non-host environments
(Loeffler et al., 2005; Roberts et al., 2013; Steadmon et al., 2023; Thapaliya et al., 2017), At the
cellular level, sub-MIC antibiotics can affect *S. aureus* by directly binding to virulence factors,
modulating virulence regulators, and inducing the expression of toxins like hemolysin (Chen et
al., 2021; Hodille et al., 2017). Prior exposure to sub-MIC inside and outside hosts can thus
mediate traits involved in virulence upon host encounter. Our results further provided evidence
for potential links between responses to antibiotic selection and virulence. Mutations in genes
involved in resistance to antibiotics were found more often in populations with increased virulence,
suggesting that antibiotic adaptation may also favor evolution of virulence. Furthermore, in the
absence of oxacillin selection, populations lost their ability to hemolyze red blood cells, supporting
previous findings that antibiotics can influence expression of virulence factors (Andersson and
Hughes, 2014; Chen et al., 2021).

Likewise, host factors contributed to antibiotic resistance. All populations initially sensitive to
oxacillin exhibited increased MICs when under selection from both host and sub-MIC oxacillin.
One population evolving under selection from either solely sub-MIC antibiotics or solely the host
exhibited decreased oxacillin sensitivity, which is consistent with previous studies examining
hosts (McVicker et al., 2014) and sub-MIC (Pereira et al., 2023; Wistrand-Yuen et al., 2018)
exposure separately. The pathogens in these studies evolved for much longer than our study,
suggesting that having a host accelerated the evolution of antibiotic resistance, potentially due to
bottlenecks that favor resistant isolates (McVicker et al., 2014). Evolution in hosts also contributed
to pathogen growth outside hosts, an important trait for opportunistic pathogens as persistence in
the environment increases the likelihood of contact with new hosts. Host association also resulted
in more indels, including large-scale deletions of antibiotic resistance-conferring SCC*mec* and
ACME. Genome degradation is a common pattern found in the transition to long-term within-host
adaptation across diverse pathogens (Bentley and Parkhill, 2015; Klemm et al., 2016; Lawrence,
2005). Ultimately, exposure to hosts and antibiotics together could create a feedback loop where
both evolutionary pressures select for better adaptation in the other, resulting in highly virulent
and resistant pathogens that can persist outside hosts.

Several mutations arose in genes with primary roles in metabolism, such as *codY*. It is becoming
clear that pathogen metabolism plays a central role in its virulence (Feinbaum et al., 2012; Lindsay
et al., 2023) and antibiotic resistance (Zampieri et al., 2017). For example, mutations in
chromosomal- and plasmid-encoded metabolic genes are widely prevalent among *E. coli* strains
and protect them against antibiotic challenges (Lopatkin et al., 2021; Palomino et al., 2023).
Furthermore, antibiotics can directly select for reduced metabolic rates to dampen the deleterious
effects of drugs (Lopatkin et al., 2021), and increased resistance is more likely to evolve at high
nutrient concentrations (Windels et al., 2024). Similarly, metabolic efficiency can alter virulence
(Lindsay et al., 2023), while resource limitations can hinder the evolution of virulence (Pak et al.,
2024). In *S. aureus*, CodY de-represses expression of virulence factors under resource-depleted
conditions in order to acquire nutrients through the lysing of host cells (Brinsmade, 2017). Our
genomic data were consistent with the observation that *codY* may be mediating virulence through
control of *agr*: MSSA populations under selection from both host and sub-MIC oxacillin exhibited

the greatest virulence and was one of the two treatments without mutations in *agr*. These findings
suggested that the *codY* mutations in these populations may be deleterious, rendering CodY less
efficient in repressing *agr*-mediated virulence as a result. Future experiments may include
introducing these mutations into the ancestral background to directly link the mutations in these
genes to evolved virulence. As resources likely differ in abundance and type between hosts and
media, our findings illustrated the importance of taking into consideration the environment in which
pathogens evolve when evaluating traits of interest.
558
Regulatory genes in general acquired more mutations than expected by chance alone. This
finding supported previous work showing that regulatory elements were more likely to acquire
phenotype-altering mutations, especially in negative regulators (e.g., *codY*) (Lind et al., 2015).
Enrichment of mutations in regulatory genes has also been found in *E. coli* under antibiotic
selection and in *Pseudomonas aeruginosa* under host selection (Card et al., 2021; Jansen et al.,
2015). Pleiotropic genes like regulators may increase the capacity to adapt to multiple
environments. Because *C. elegans* is a novel host to *S. aureus*, selection is more likely to act on
genes that have large initial gains in the fitness landscape, such as those that have pleiotropic
effects (Bomblies and Peichel, 2022). Indeed, mutations in several metabolic regulators (*codY,*
purR, gpmA) and virulence regulators (*saeRS*) were mainly identified in populations involving
hosts. The resulting phenotypes of these mutations, such as increased biofilm production, can
then confer multiple benefits like increased virulence and antibiotic resistance. Our experimental
results revealed similar patterns to a comparative study analyzing over 2000 *S. aureus* genomes,
where mutations in genes with pleiotropic effects, such as *purR*, may have underly pathogen
adaptation during systemic infection (Giulieri et al., 2022). Thus, changes in genes of large effects
like global regulators appear to be critical mediators between different traits that altogether allow
bacteria to rapidly adapt to new and stressful environments.
576
Antibiotic resistance did not impede the evolution of virulence. Antibiotic resistance tends to be
costly in the absence of antibiotics; this cost can subsequently constrain evolution of traits
requiring cellular resources, such as virulence (Ferenci, 2016). However, mutations conferring
resistance can have little to no costs, depending on the pathogen species and drugs, and costs
can be offset by compensatory mutations (Melnyk et al., 2015). Sub-MIC levels favor low-cost
mutations (Westhoff et al., 2017), therefore virulence and antibiotic resistance may not trade-off
during disease emergence (Visher et al., 2021). Virulence may alternatively be costly in the
presence of antibiotic resistance—populations evolved from the MRSA ancestor without either
selective pressure exhibited the lowest virulence. The corresponding MSSA populations did not
have a similar decline in virulence, suggesting that relaxed selection attenuated virulence for
populations with existing resistance. Genomic differences between evolved MRSA and MSSA
further demonstrate diverging evolutionary pathways in antibiotic-resistant *versus* antibiotic-
sensitive populations. Oxacillin-exposed MSSA had no mutations in *agr* and retained their
hemolytic ability, while some MRSA populations gained *agr* mutations and lost the trait,
suggesting that other pathways are utilized to express virulence. Lastly, mutations in MRSA under
selection from both host and antibiotics were significantly enriched in blood and systemic *S.*
aureus isolated from humans, suggesting that antibiotics may have significant roles in MRSA
invasive infection. Acquiring mutations in genes that impact antibiotic resistance, such as *pbpA*
and *pbpB*, may allow *S. aureus* to persist longer in hosts. Additional direct tests are needed to
evaluate the role of these mutations in adaptation of *S. aureus* to different infection sites.
597
598 Recent work has shown that diverse genotypes of one pathogen species converge on similar
599 phenotypes when facing similar selective pressures (Filipow et al., 2023). Here, we showed that
600 antibiotic resistance status can lead otherwise isogenic pathogens to use different adaptive
601 strategies. Initially susceptible pathogens, facing novel selection from host and antibiotics,

exhibited reduced phenotypic and genomic variation over evolutionary time and thus may have had a limited number of pathways to adapt to new environments. By contrast, resistant populations had greater variance and thus may have had more routes to adapt. Overall, the potential for low-cost (Visher et al., 2021; Westhoff et al., 2017) and pleiotropic mutations (Bomblies and Peichel, 2022) may have created ideal conditions for a novel pathogen to establish in host populations regardless of its susceptibility to antibiotics. Our work demonstrated that responding to multiple pressures can result in rapid adaptation to new hosts and even evolution of traits not under selection, emphasizing the importance of non-canonical pathways (e.g., metabolism) in shaping these traits. Pathogen evolution in a tractable invertebrate animal model yielded phenotypes and genotypes similar to those identified in mammalian hosts, highlighting the utility of evolution experiments to identify potential ecological and genetic mechanisms that may give rise to pathogen traits conserved across systems. Our findings ultimately emphasize the importance of considering the host context in the evolution of antibiotic resistance. Integrating multiple traits, such as virulence, antibiotic resistance, and fitness may be critical in identifying the factors that facilitate host shifts and persistence of drug-resistant pathogens.

REFERENCES

Alves J, Vrieling M, Ring N, Yebra G, Pickering A, Prajsnar TK, Renshaw SA, Fitzgerald JR.
2024. Experimental evolution of Staphylococcus aureus in macrophages: dissection of a
conditional adaptive trait promoting intracellular survival. mBio. DOI:
https://doi.org/10.1128/mbio.00346-24

Andersson DI, Hughes D. 2014. Microbiological effects of sublethal levels of antibiotics. *Nature Reviews Microbiology*. DOI: https://doi.org/10.1038/nrmicro3270, PMID: 24861036

Azimi S, Roberts AEL, Peng S, Weitz JS, McNally A, Brown SP, Diggle SP. 2020. Allelic
polymorphism shapes community function in evolving Pseudomonas aeruginosa
populations. ISME Journal **14**:1929–1942. DOI: https://doi.org/10.1038/s41396-020-0652-0, PMID: 32341475

Bacigalupe R, Tormo-Mas MÁ, Penadés JR, Ross Fitzgerald J. 2019. A multihost bacterial
pathogen overcomes continuous population bottlenecks to adapt to new host species.
Science Advances **5**:1–15. DOI: https://doi.org/10.1126/sciadv.aax0063, PMID: 31807698

Bae T, Banger AK, Wallace A, Glass EM, Åslund F, Schneewind O, Missiakas DM. 2004.
Staphylococcus aureus virulence genes identified by bursa aurealis mutagenesis and
nematode killing.

Bashey F. 2015. Within-host competitive interactions as a mechanism for the maintenance of
parasite diversity. Philosophical Transactions of the Royal Society B: Biological Sciences
370. DOI: https://doi.org/10.1098/rstb.2014.0301, PMID: 26150667

Begun J, Gaiani JM, Rohde H, Mack D, Calderwood SB, Ausubel FM, Sifri CD. 2007.
Staphylococcal biofilm exopolysaccharide protects against Caenorhabditis elegans
immune defenses. PLoS Pathogens **3**:526–540. DOI:
https://doi.org/10.1371/journal.ppat.0030057, PMID: 17447841

Begun J, Sifri CD, Goldman S, Calderwood SB, Ausubel FM. 2005. Staphylococcus aureus
virulence factors identified by using a high-throughput Caenorhabditis elegans-killing
model. Infection and Immunity **73**:872–877. DOI: https://doi.org/10.1128/IAI.73.2.872-877.2005, PMID: 15664928

Bentley SD, Parkhill J. 2015. Genomic perspectives on the evolution and spread of bacterial
pathogens. Proceedings of the Royal Society B: Biological Sciences **282**. DOI:
https://doi.org/10.1098/rspb.2015.0488, PMID: 26702036

Betts A, Gray C, Zelek M, Maclean RC, King KC. 2018. High parasite diversity accelerates host
adaptation and diversification. Science **360**:907–911.

Bilyk BL, Panchal V V., Tinajero-Trejo M, Hobbs JK, Foster SJ. 2022. An interplay of multiple
positive and negative factors governs methicillin resistance in *Staphylococcus aureus*.
Microbiology and Molecular Biology Reviews **86**. DOI: https://doi.org/10.1128/mmbr.00159-21, PMID: 35420454

Blazanin M. 2023. gcplyr: an R package for microbial growth curve data analysis. bioRxiv
2023.04.30.538883.

Bomblies K, Peichel CL. 2022. Genetics of adaptation. Proceedings of the National Academy of
Sciences of the United States of America **119**. DOI:
https://doi.org/10.1073/pnas.2122152119, PMID: 35858399

Braga PC, Dal Sasso M, Sala MT. 2000. Sub-MIC concentrations of cefodizime interfere with
various factors affecting bacterial virulence. Journal of Antimicrobial Chemotherapy **45**:15–
25. DOI: https://doi.org/10.1093/jac/45.1.15, PMID: 10629008
Brinsmade SR. 2017. CodY, a master integrator of metabolism and virulence in Gram-positive
bacteria. *Current Genetics*. DOI: https://doi.org/10.1007/s00294-016-0656-5, PMID:
27744611
Card KJ, Thomas MD, Graves JL, Barrick JE, Lenski RE. 2021. Genomic evolution of antibiotic
resistance is contingent on genetic background following a long-term experiment with
Escherichia coli. Proceedings of the National Academy of Sciences of the United States of
America **118**:1–9. DOI: https://doi.org/10.1073/pnas.2016886118, PMID: 33441451
Chen J, Zhou H, Huang J, Zhang R, Rao X. 2021. Virulence alterations in staphylococcus
aureus upon treatment with the sub-inhibitory concentrations of antibiotics. *Journal of*
Advanced Research. DOI: https://doi.org/10.1016/j.jare.2021.01.008
Cheung GYC, Duong AC, Otto M. 2012. Direct and synergistic hemolysis caused by
Staphylococcus phenol-soluble modulins: Implications for diagnosis and pathogenesis.
Microbes and Infection **14**:380–386. DOI: https://doi.org/10.1016/j.micinf.2011.11.013,
PMID: 22178792
Cisneros-Mayoral S, Graña-Miraglia L, Perez-Morales D, Peña-Miller R, Fuentes-Hernández A.
2022. Evolutionary History and Strength of Selection Determine the Rate of Antibiotic
Resistance Adaptation. Molecular Biology and Evolution **39**. DOI:
https://doi.org/10.1093/molbev/msac185, PMID: 36062982
CLSI. n.d. Performance standards for antimicrobial susceptibility testing: CLSI approved
standard M100-S23 2013.
Cobey S. 2014. Pathogen evolution and the immunological niche. Annals of the New York
Academy of Sciences **1320**:1–15. DOI: https://doi.org/10.1111/nyas.12493, PMID:
25040161
Cordero OX, Wildschutte H, Kirkup B, Proehl S, Ngo L, Hussain F, Le Roux F, Mincer T, Polz
MF. 2012. Ecological populations of bacteria act as socially cohesive units of antibiotic
production and resistance. Science **337**:1228–1231. DOI:
https://doi.org/10.1126/science.1219385, PMID: 22955834
Day T, Kennedy DA, Read AF, Gandon S. 2022. Pathogen evolution during vaccination
campaigns. PLoS biology **20**:e3001804. DOI: https://doi.org/10.1371/journal.pbio.3001804,
PMID: 36149891
Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved microbes
from next-generation sequencing data using breseq. *Engineering and Analyzing*
Multicellular Systems: Methods and Protocols. p. 165–188. DOI:
https://doi.org/10.1007/978-1-4939-0554-6
Diep BA, Gill SR, Chang RF, Phan TH Van, Chen JH, Davidson MG, Lin F, Lin J, Carleton HA,
Mongodin EF, Sensabaugh GF, Perdreau-Remington F. 2006. Complete genome
sequence of USA300, an epidemic clone of community-acquired meticillin-resistant
Staphylococcus aureus. Lancet **367**:731–739. DOI: https://doi.org/10.1016/S0140-
6736(06)68231-7, PMID: 16517273
Diep BA, Stone GG, Basuino L, Graber CJ, Miller A, Des Etages SA, Jones A, Palazzolo-
Ballance AM, Perdreau-Remington F, Sensabaugh GF, DeLeo FR, Chambers HF. 2008.

The arginine catabolic mobile element and staphylococcal chromosomal cassette mec linkage: Convergence of virulence and resistance in the USA300 clone of methicillin-resistant Staphylococcus aureus. Journal of Infectious Diseases **197**:1523–1530. DOI: https://doi.org/10.1086/587907, PMID: 18700257

Ekroth AKE, Gerth M, Stevens EJ, Ford SA, King KC. 2021. Host genotype and genetic diversity shape the evolution of a novel bacterial infection. ISME Journal **15**:2146–2157. DOI: https://doi.org/10.1038/s41396-021-00911-3, PMID: 33603148

El-Houssaini HH, Elnabawy OM, Nasser HA, Elkhatib WF. 2019. Influence of subinhibitory antifungal concentrations on extracellular hydrolases and biofilm production by Candida albicans recovered from Egyptian patients. BMC Infectious Diseases **19**:1–9. DOI: https://doi.org/10.1186/s12879-019-3685-0, PMID: 30651066

Erler S, Cotter SC, Freitak D, Koch H, Palmer-Young EC, de Roode JC, Smilanich AM, Lattorff HMG. 2024. Insects' essential role in understanding and broadening animal medication. Trends in Parasitology **40**:338–349. DOI: https://doi.org/10.1016/j.pt.2024.02.003

Fait A, Andersson DI, Ingmer H. 2023. Evolutionary history of Staphylococcus aureus influences antibiotic resistance evolution. Current Biology **33**:3389-3397.e5. DOI: https://doi.org/10.1016/j.cub.2023.06.082, PMID: 37494936

Feinbaum RL, Urbach JM, Liberati NT, Djonovic S, Adonizio A, Carvunis AR, Ausubel FM. 2012. Genome-wide identification of Pseudomonas aeruginosa virulence-related genes using a Caenorhabditis elegans infection model. PLoS Pathogens **8**:11. DOI: https://doi.org/10.1371/journal.ppat.1002813, PMID: 22911607

Ferenci T. 2016. Trade-off mechanisms shaping the diversity of bacteria. Trends in Microbiology **24**:209–223. DOI: https://doi.org/10.1016/j.tim.2015.11.009, PMID: 26705697

Filipow N, Mallon S, Shewaramani S, Kassen R, Wong A. 2023. The impact of genetic background during laboratory evolution of Pseudomonas aeruginosa in a cystic fibrosis-like environment . Evolution **78**:566–578. DOI: https://doi.org/10.1093/evolut/qpad189

Ford SA, Kao D, Williams D, King KC. 2016. Microbe-mediated host defence drives the evolution of reduced pathogen virulence. Nature Communications **7**:1–9. DOI: https://doi.org/10.1038/ncomms13430

Ford SA, Williams D, Paterson S, King KC. 2017. Co-evolutionary dynamics between a defensive microbe and a pathogen driven by fluctuating selection. Molecular Ecology **26**:1778–1789. DOI: https://doi.org/10.1111/mec.13906

Geisinger E, Isberg RR. 2017. Interplay between antibiotic resistance and virulence during Disease promoted by multidrug-resistant bacteria. Journal of Infectious Diseases **215**:S9–S17. DOI: https://doi.org/10.1093/infdis/jiw402, PMID: 28375515

Giulieri SG, Guérillot R, Duchene S, Hachani A, Daniel D, Seemann T, Davis JS, Tong SYC, Young BC, Wilson DJ, Stinear TP, Howden BP. 2022. Niche-specific genome degradation and convergent evolution shaping Staphylococcus aureus adaptation during severe infections. eLife **11**:1–33. DOI: https://doi.org/10.7554/eLife.77809, PMID: 35708309

Giulieri SG, Guérillot R, Kwong JC, Monk IR, Hayes AS, Daniel D, Baines S, Sherry NL, Holmes NE, Ward P, Gao W, Seemann T, Stinear TP, Howden BP. 2020. Comprehensive genomic investigation of adaptive mutations driving the low-level oxacillin resistance phenotype in staphylococcus aureus. mBio **11**:1–18. DOI: https://doi.org/10.1128/mBio.02882-20, PMID: 33293382

Gullberg E, Cao S, Berg OG, Ilbäck C, Sandegren L, Hughes D, Andersson DI. 2011. Selection
of resistant bacteria at very low antibiotic concentrations. PLoS Pathogens **7**:1–9. DOI:
https://doi.org/10.1371/journal.ppat.1002158, PMID: 21811410
Haddadin RNS, Saleh S, Al-Adham ISI, Buultjens TEJ, Collier PJ. 2010. The effect of
subminimal inhibitory concentrations of antibiotics on virulence factors expressed by
Staphylococcus aureus biofilms. Journal of Applied Microbiology **108**:1281–1291. DOI:
https://doi.org/10.1111/j.1365-2672.2009.04529.x, PMID: 19778348
Herren CM, Baym M. 2022. Decreased thermal niche breadth as a trade-off of antibiotic
resistance. ISME Journal **16**:1843–1852. DOI: https://doi.org/10.1038/s41396-022-01235-
6, PMID: 35422477
Higazy D, Pham AD, van Hasselt C, Høiby N, Jelsbak L, Moser C, Ciofu O. 2024. In vivo
evolution of antimicrobial resistance in a biofilm model of Pseudomonas aeruginosa lung
infection . The ISME Journal **18**:1–12. DOI: https://doi.org/10.1093/ismejo/wrae036
Hoang Kim L, Read TD, King KC. 2024. Incomplete immunity in a natural animal-microbiota
interaction selects for higher pathogen virulence. Current Biology **34**:1–7.
Hoang Kim L., Read TD, King KC. 2024. Defense Heterogeneity in Host Populations Gives Rise
to Pathogen Diversity. The American Naturalist **204**:000–000. DOI:
https://doi.org/10.1086/731996
Hodille E, Rose W, Diep BA, Goutelle S, Lina G, Dumitrescu O. 2017. The Role of Antibiotics in
Modulating Virulence in Staphylococcus aureus. DOI: https://doi.org/10.1128/CMR
Hudzicki J. 2009. Kirb-Bauer disk diffusion susceptibility test protocol. American Society for
Microbiology 1–23.
Ibarra JA, Pérez-Rueda E, Carroll RK, Shaw LN. 2013. Global analysis of transcriptional
regulators in Staphylococcus aureus. BMC Genomics **14**. DOI:
https://doi.org/10.1186/1471-2164-14-126, PMID: 23442205
Irazoqui JE, Troemel ER, Feinbaum RL, Luhachack LG, Cezairliyan BO, Ausubel FM. 2010.
Distinct pathogenesis and host responses during infection of C. elegans by P. aeruginosa
and S. aureus. PLoS Pathogens **6**:1–24. DOI:
https://doi.org/10.1371/journal.ppat.1000982, PMID: 20617181
Jackson RW, Vinatzer B, Arnold DL, Dorus S, Murillo J. 2011. The influence of the accessory
genome on bacterial pathogen evolution. Mobile Genetic Elements **1**:55–65. DOI:
https://doi.org/10.4161/mge.1.1.16432
Jansen G, Crummenerl LL, Gilbert F, Mohr T, Pfefferkorn R, Thänert R, Rosenstiel P,
Schulenburg H. 2015. Evolutionary transition from pathogenicity to commensalism: Global
regulator mutations mediate fitness gains through virulence attenuation. Molecular Biology
and Evolution **32**:2883–2896. DOI: https://doi.org/10.1093/molbev/msv160, PMID:
26199376
Khan F, Lee JW, Javaid A, Park SK, Kim YM. 2020. Inhibition of biofilm and virulence properties
of Pseudomonas aeruginosa by sub-inhibitory concentrations of aminoglycosides.
Microbial Pathogenesis **146**. DOI: https://doi.org/10.1016/j.micpath.2020.104249, PMID:
32418905
King JM, Kulhankova K, Stach CS, Vu BG, Salgado-pabón W. 2016. Phenotypes and virulence
among Staphylococcus aureus USA100, USA200, USA300, USA400, and USA600 clonal
lineages. mSphere **1**:1–15. DOI: https://doi.org/10.1128/mSphere.00071-16.Editor

Klemm EJ, Gkrania-Klotsas E, Hadfield J, Forbester JL, Harris SR, Hale C, Heath JN, Wileman
T, Clare S, Kane L, Goulding D, Otto TD, Kay S, Doffinger R, Cooke FJ, Carmichael A,
Lever AML, Parkhill J, MacLennan CA, Kumararatne D, Dougan G, Kingsley RA. 2016.
Emergence of host-adapted Salmonella Enteritidis through rapid evolution in an
immunocompromised host. Nature Microbiology **1**:1–6. DOI:
https://doi.org/10.1038/nmicrobiol.2015.23, PMID: 27572160
Koch G, Yepes A, Förstner KU, Wermser C, Stengel ST, Modamio J, Ohlsen K, Foster KR,
Lopez D. 2014. Evolution of resistance to a last-resort antibiotic in staphylococcus aureus
via bacterial competition. Cell **158**:1060–1071. DOI:
https://doi.org/10.1016/j.cell.2014.06.046, PMID: 25171407
Kourtis AP, Hatfield ; Kelly, Baggs J, Mu ; Yi, See I, Epson E, Nadle ; Joelle, Kainer MA,
Dumyati G, Petit S, Ray SM, Ham D, Capers C, Ewing ; Heather, Coffin N, Mcdonald ; L
Clifford, Jernigan J, Cardo D. 2019. Vital signs: epidemiology and recent trends in
methicillin-resistant and in methicillin-susceptible Staphylococcus aureus bloodstream
infections-United States. Morbidity and Mortality Weekly Report **68**:214–219.
Kubinak JL, Ruff JS, Hyzer CW, Slev PR, Potts WK. 2012. Experimental viral evolution to
specific host MHC genotypes reveals fitness and virulence trade-offs in alternative MHC
types. Proceedings of the National Academy of Sciences of the United States of America
109:3422–3427. DOI: https://doi.org/10.1073/pnas.1112633109, PMID: 22323587
Kuroda H, Kuroda M, Cui L, Hiramatsu K. 2007. Subinhibitory concentrations of β-lactam induce
haemolytic activity in Staphylococcus aureus through the SaeRS two-component system.
FEMS Microbiology Letters **268**:98–105. DOI: https://doi.org/10.1111/j.1574-
6968.2006.00568.x, PMID: 17263851
Kwiecinski JM, Horswill AR. 2020. Staphylococcus aureus bloodstream infections: pathogenesis
and regulatory mechanisms. Current Opinion in Microbiology **53**:51–60. DOI:
https://doi.org/10.1016/j.mib.2020.02.005, PMID: 32172183
Larsson DGJ, Flach CF. 2022. Antibiotic resistance in the environment. *Nature Reviews*
Microbiology. DOI: https://doi.org/10.1038/s41579-021-00649-x, PMID: 34737424
Launay A, Wu CJ, Dulanto Chiang A, Youn JH, Khil PP, Dekker JP. 2021. In vivo evolution of
an emerging zoonotic bacterial pathogen in an immunocompromised human host. Nature
Communications **12**:1–12. DOI: https://doi.org/10.1038/s41467-021-24668-7, PMID:
34301946
Lawrence JG. 2005. Common themes in the genome strategies of pathogens. Current Opinion
in Genetics and Development **15**:584–588. DOI: https://doi.org/10.1016/j.gde.2005.09.007,
PMID: 16188434
Lees JA, Kremer PHC, Manso AS, Croucher NJ, Ferwerda B, Serón MV, Oggioni MR, Parkhill
J, Brouwer MC, van der Ende A, van de Beek D, Bentley SD. 2017. Large scale genomic
analysis shows no evidence for pathogen adaptation between the blood and cerebrospinal
fluid niches during bacterial meningitis. Microbial Genomics **3**. DOI:
https://doi.org/10.1099/mgen.0.000103, PMID: 28348877
Lind PA, Farr AD, Rainey PB. 2015. Experimental evolution reveals hidden diversity in
evolutionary pathways. eLife **2015**:1–17. DOI: https://doi.org/10.7554/eLife.07074, PMID:
25806684

Lindsay RJ, Holder PJ, Talbot NJ, Gudelj I. 2023. Metabolic efficiency reshapes the seminal
relationship between pathogen growth rate and virulence. Ecology Letters **26**:896–907.
DOI: https://doi.org/10.1111/ele.14218, PMID: 37056166
Loeffler A, Boag AK, Sung J, Lindsay JA, Guardabassi L, Dalsgaard A, Smith H, Stevens KB,
Lloyd DH. 2005. Prevalence of methicillin-resistant Staphylococcus aureus among staff
and pets in a small animal referral hospital in the UK. Journal of Antimicrobial
Chemotherapy **56**:692–697. DOI: https://doi.org/10.1093/jac/dki312, PMID: 16141276
Long DR, Penewit K, Lo HY, Almazan J, Holmes EA, Bryan AB, Wolter DJ, Lewis JD, Waalkes
A, Salipante SJ. 2023. In Vitro Selection Identifies Staphylococcus aureus Genes
Influencing Biofilm Formation. Infection and Immunity **91**. DOI:
https://doi.org/10.1128/iai.00538-22, PMID: 36847490
Lopatkin AJ, Bening SC, Manson AL, Stokes JM, Kohanski MA, Badran AH, Earl AM, Cheney
NJ, Yang JH, Collins JJ. 2021. Clinically relevant mutations in core metabolic genes confer
antibiotic resistance. Science **371**. DOI: https://doi.org/10.1126/science.aba0862, PMID:
33602825
Mahrt N, Tietze A, Künzel S, Franzenburg S, Barbosa C, Jansen G, Schulenburg H. 2021.
Bottleneck size and selection level reproducibly impact evolution of antibiotic resistance.
Nature Ecology and Evolution **53**:1275. DOI: https://doi.org/10.1038/s41588-021-00939-3,
PMID: 34493861
Majerczyk CD, Sadykov MR, Luong TT, Lee C, Somerville GA, Sonenshein AL. 2008.
Staphylococcus aureus CodY negatively regulates virulence gene expression. Journal of
Bacteriology **190**:2257–2265. DOI: https://doi.org/10.1128/JB.01545-07, PMID: 18156263
Martinez Arbizu P. 2020. pairwiseAdonis: Pairwise multilevel comparison using adonis.
McAdam PR, Holmes A, Templeton KE, Fitzgerald JR. 2011. Adaptive evolution of
Staphylococcus aureus during chronic endobronchial infection of a cystic fibrosis patient.
PLoS ONE **6**:1–5. DOI: https://doi.org/10.1371/journal.pone.0024301, PMID: 21912685
McVicker G, Prajsnar TK, Williams A, Wagner NL, Boots M, Renshaw SA, Foster SJ. 2014.
Clonal Expansion during Staphylococcus aureus Infection Dynamics Reveals the Effect of
Antibiotic Intervention. PLoS Pathogens **10**. DOI:
https://doi.org/10.1371/journal.ppat.1003959, PMID: 24586163
Mei M, Thomas J, Diggle SP. 2021. Heterogenous susceptibility to r-pyocins in populations of
pseudomonas aeruginosa sourced from cystic fibrosis lungs. mBio **12**. DOI:
https://doi.org/10.1128/mBio.00458-21, PMID: 33947755
Melnyk AH, Wong A, Kassen R. 2015. The fitness costs of antibiotic resistance mutations.
Evolutionary Applications **8**:273–283. DOI: https://doi.org/10.1111/eva.12196
Merlo LMF, Sprouffske K, Howard TC, Gardiner KL, Caulin AF, Blum SM, Evans P, Bedalov A,
Sniegowski PD, Maley CC. 2020. Application of simultaneous selective pressures slows
adaptation. Evolutionary Applications **13**:1615–1625. DOI:
https://doi.org/10.1111/eva.13062
Montgomery CP, Boyle-Vavra S, Daum RS. 2010. Importance of the global regulators agr and
SaeRS in the pathogenesis of CA-MRSA USA300 infection. PLoS ONE **5**:1–9. DOI:
https://doi.org/10.1371/journal.pone.0015177, PMID: 21151999
Nix DE, Goodwin SD, Peloquin CA, Rotella DL, Schentag JJ. 1991. Antibiotic tissue penetration
and its relevance: Impact of tissue penetration on infection response. Antimicrobial Agents

and Chemotherapy **35**:1953–1959. DOI: https://doi.org/10.1128/AAC.35.10.1953, PMID:
1759813

Novick RP. 2003. Autoinduction and signal transduction in the regulation of staphylococcal
virulence. Molecular Microbiology **48**:1429–1449. DOI: https://doi.org/10.1046/j.1365-
2958.2003.03526.x, PMID: 12791129

Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P,
Stevens M, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Carvalho
G, Chirico M, Caceres M De, Durand S, Evangelista H, FitzJohn R, Friendly M, Furneaux
B, Hannigan G, Hill M, Lahti L, McGlinn D, Ouellette M, Cunha ER, Smith T, Stier A, Braak
C Ter, Weedon J. 2024. vegan: Community Ecology Package.

Oz T, Guvenek A, Yildiz S, Karaboga E, Tamer YT, Mumcuyan N, Ozan VB, Senturk GH, Cokol
M, Yeh P, Toprak E. 2014. Strength of selection pressure is an important parameter
contributing to the complexity of antibiotic resistance evolution. Molecular Biology and
Evolution **31**:2387–2401. DOI: https://doi.org/10.1093/molbev/msu191, PMID: 24962091

Pak D, Kamiya T, Greischar MA. 2024. Proliferation in malaria parasites: How resource
limitation can prevent evolution of greater virulence. Evolution **78**:1287–1301. DOI:
https://doi.org/10.1093/evolut/qpae057, PMID: 38581661

Palomino A, Gewurz D, DeVine L, Zajmi U, Moralez J, Abu-Rumman F, Smith RP, Lopatkin AJ.
2023. Metabolic genes on conjugative plasmids are highly prevalent in Escherichia coli and
can protect against antibiotic treatment. ISME Journal **17**:151–162. DOI:
https://doi.org/10.1038/s41396-022-01329-1, PMID: 36261510

Paterson GK, Harrison EM, Murray GGR, Welch JJ, Warland JH, Holden MTG, Morgan FJE, Ba
X, Koop G, Harris SR, Maskell DJ, Peacock SJ, Herrtage ME, Parkhill J, Holmes MA.
2015. Capturing the cloud of diversity reveals complexity and heterogeneity of MRSA
carriage, infection and transmission. Nature Communications **6**. DOI:
https://doi.org/10.1038/ncomms7560, PMID: 25814293

Penley M, Morran L. 2018. Assessment of Caenorhabditis elegans Competitive Fitness in the
Presence of a Bacterial Parasite. Bio-Protocol **8**:1–14. DOI:
https://doi.org/10.21769/bioprotoc.2971

Pereira C, Warsi OM, Andersson DI. 2023. Pervasive Selection for Clinically Relevant
Resistance and Media Adaptive Mutations at Very Low Antibiotic Concentrations.
Molecular Biology and Evolution **40**:1–13.

Petit RA, Read TD. 2020. Bactopia: a Flexible Pipeline for Complete Analysis of Bacterial
Genomes. mSystems **5**. DOI: https://doi.org/10.1128/msystems.00190-20, PMID:
32753501

Poon R, Basuino L, Satishkumar N, Chatterjee A, Mukkayyan N, Buggeln E, Huang L, Nair V,
Argudín MA, Datta SK, Chambers HF, Chatterjee SS. 2022. Loss of GdpP function in
Staphylococcus aureus leads to β-lactam tolerance and enhanced evolution of β-lactam
resistance. Antimicrobial Agents and Chemotherapy **66**. DOI:
https://doi.org/10.1128/AAC.01431-21, PMID: 34843389

Raghuram V, Petit RA, Karol Z, Mehta R, Weissman DB, Read TD. 2024. Average nucleotide
identity-based Staphylococcus aureus strain grouping allows identification of strain-specific
genes in the pangenome. mSystems **9**. DOI: https://doi.org/10.1128/msystems.00143-24,
PMID: 38934646

Revitt-Mills SA, Robinson A. 2020. Antibiotic-Induced Mutagenesis: Under the Microscope.
Frontiers in Microbiology **11**. DOI: https://doi.org/10.3389/fmicb.2020.585175

Roberts MC, Soge OO, No D. 2013. Comparison of multi-drug resistant environmental
methicillin-resistant Staphylococcus aureus isolated from recreational beaches and high
touch surfaces in built environments. Frontiers in Microbiology **4**:1–8. DOI:
https://doi.org/10.3389/fmicb.2013.00074

Ruiz-Bedoya T, Wang PW, Desveaux D, Guttman DS. 2023. Cooperative virulence via the
collective action of secreted pathogen effectors. Nature Microbiology **8**:640–650. DOI:
https://doi.org/10.1038/s41564-023-01328-8, PMID: 36782026

Schwan WR, Langhorne MH, Ritchie HD, Stover CK. 2003. Loss of hemolysin expression in
Staphylococcus aureus agr mutants correlates with selective survival during mixed
infections in murine abscesses and wounds. FEMS Immunology and Medical Microbiology
38:23–28. DOI: https://doi.org/10.1016/S0928-8244(03)00098-1, PMID: 12900051

Sharma M, Quader S, Guttal V, Isvaran K. 2020. The enemy of my enemy: multiple interacting
selection pressures lead to unexpected anti-predator responses. Oecologia **192**:1–12. DOI:
https://doi.org/10.1007/s00442-019-04552-4, PMID: 31773313

Shore AC, Coleman DC. 2013. Staphylococcal cassette chromosome mec: Recent advances
and new insights. International Journal of Medical Microbiology **303**:350–359. DOI:
https://doi.org/10.1016/j.ijmm.2013.02.002, PMID: 23499303

Sifri CD, Begun J, Ausubel FM, Calderwood SB. 2003a. Caenorhabditis elegans as a model
host for Staphylococcus aureus pathogenesis. Infection and Immunity **71**:2208–2217. DOI:
https://doi.org/10.1128/IAI.71.4.2208-2217.2003, PMID: 12654843

Sifri CD, Begun J, Ausubel FM, Calderwood SB. 2003b. Caenorhabditis elegans as a model
host for Staphylococcus aureus pathogenesis. Infection and Immunity **71**:2208–2217. DOI:
https://doi.org/10.1128/IAI.71.4.2208-2217.2003, PMID: 12654843

Souvorov A, Agarwala R, Lipman DJ. 2018. SKESA: Strategic k-mer extension for scrupulous
assemblies. Genome Biology **19**:1–13. DOI: https://doi.org/10.1186/s13059-018-1540-z,
PMID: 30286803

Spagnolo F, Trujillo M, Dennehy JJ. 2021. Why do antibiotics exist? mBio **12**. DOI:
https://doi.org/10.1128/mBio.01966-21, PMID: 34872345

Steadmon M, Ngiraklang K, Nagata M, Masga K, Kiana L. Frank. 2023. Effects of water turbidity
on the survival of Staphylococcus aureus in environmental fresh and brackish waters.
Water Environment Research.

Stiernagle T. 2006. Maintenance of *C. elegans*. WormBook 2006. DOI:
https://doi.org/10.1895/wormbook.1.101.1, PMID: 18050451

Thapaliya D, Taha M, Dalman MR, Kadariya J, Smith TC. 2017. Environmental contamination
with Staphylococcus aureus at a large, Midwestern university campus. Science of the Total
Environment **599–600**:1363–1368. DOI: https://doi.org/10.1016/j.scitotenv.2017.05.080,
PMID: 28525941

Thomer L, Schneewind O, Missiakas D. 2016. Pathogenesis of Staphylococcus aureus
Bloodstream Infections. Annual Review of Pathology: Mechanisms of Disease **11**:343–364.
DOI: https://doi.org/10.1146/annurev-pathol-012615-044351, PMID: 26925499

Toprak E, Veres A, Michel JB, Chait R, Hartl DL, Kishony R. 2012. Evolutionary paths to
antibiotic resistance under dynamically sustained drug selection. Nature Genetics **44**:101–
105. DOI: https://doi.org/10.1038/ng.1034, PMID: 22179135
Tsouklidis N, Kumar R, Heindl SE, Soni R, Khan S. 2020. Understanding the Fight Against
Resistance: Hospital-Acquired Methicillin-Resistant Staphylococcus Aureus vs.
Community-Acquired Methicillin-Resistant Staphylococcus Aureus. Cureus. DOI:
https://doi.org/10.7759/cureus.8867
Vega NM, Gore J. 2017. Stochastic assembly produces heterogeneous communities in the
Caenorhabditis elegans intestine. PLoS Biology **15**:1–20. DOI:
https://doi.org/10.1371/journal.pbio.2000633, PMID: 28257456
Visher E, Evensen C, Guth S, Lai E, Norfolk M, Rozins C, Sokolov NA, Sui M, Boots M. 2021.
The three Ts of virulence evolution during zoonotic emergence. Proceedings of the Royal
Society B: Biological Sciences **288**. DOI: https://doi.org/10.1098/rspb.2021.0900, PMID:
34375554
Wahl LM, Gerrish PJ, Saika-Voivod I. 2002. Evaluating the impact of population bottlenecks in
experimental evolution. Genetics **162**:961–971. DOI:
https://doi.org/10.1093/genetics/162.2.961, PMID: 12399403
Westhoff S, Van Leeuwe TM, Qachach O, Zhang Z, Van Wezel GP, Rozen DE. 2017. The
evolution of no-cost resistance at sub-MIC concentrations of streptomycin in Streptomyces
coelicolor. ISME Journal **11**:1168–1178. DOI: https://doi.org/10.1038/ismej.2016.194,
PMID: 28094796
Windels EM, Cool L, Persy E, Swinnen J, Matthay P, Van Den Bergh B, Wenseleers T, Michiels
J. 2024. Antibiotic dose and nutrient availability differentially drive the evolution of antibiotic
resistance and persistence. ISME Journal **18**. DOI: https://doi.org/10.1093/ismejo/wrae070,
PMID: 38691440
Windels EM, Van Den Bergh B, Michiels J. 2020. Bacteria under antibiotic attack: Different
strategies for evolutionary adaptation. PLoS Pathogens **16**:1–8. DOI:
https://doi.org/10.1371/journal.ppat.1008431, PMID: 32379814
Wistrand-Yuen E, Knopp M, Hjort K, Koskiniemi S, Berg OG, Andersson DI. 2018. Evolution of
high-level resistance during low-level antibiotic exposure. Nature Communications **9**. DOI:
https://doi.org/10.1038/s41467-018-04059-1, PMID: 29686259
Wright GD. 2007. The antibiotic resistome: The nexus of chemical and genetic diversity. Nature
Reviews Microbiology **5**:175–186. DOI: https://doi.org/10.1038/nrmicro1614, PMID:
17277795
Wu K, Conly J, McClure JA, Elsayed S, Louie T, Zhang K. 2010. Caenorhabditis elegans as a
host model for community-associated methicillin-resistant Staphylococcus aureus. Clinical
Microbiology and Infection **16**:245–254. DOI: https://doi.org/10.1111/j.1469-
0691.2009.02765.x
Wu K, Zhang K, McClure J, Zhang J, Schrenzel J, Francois P, Harbarth S, Conly J. 2013. A
correlative analysis of epidemiologic and molecular characteristics of methicillin-resistant
Staphylococcus aureus clones from diverse geographic locations with virulence measured
by a Caenorhabditis elegans host model. European Journal of Clinical Microbiology and
Infectious Diseases **32**:33–42. DOI: https://doi.org/10.1007/s10096-012-1711-x, PMID:
22898726

Zampieri M, Enke T, Chubukov V, Ricci V, Piddock L, Sauer U. 2017. Metabolic constraints on
the evolution of antibiotic resistance. Molecular Systems Biology **13**:1–14. DOI:
https://doi.org/10.15252/msb.20167028, PMID: 28265005
Zapotoczna M, O'Neill E, O'Gara JP. 2016. Untangling the diverse and redundant mechanisms
of *Staphylococcus aureus* biofilm formation. PLoS Pathogens **12**:1–6. DOI:
https://doi.org/10.1371/journal.ppat.1005671, PMID: 27442433
1014
1015

MATERIALS AND METHODS
Bacteria and nematode strains
Staphylococcus aureus USA300 JE2 (NR-46543) and USA300 JE2 *mecA*-tn (Transposon Mutant NE1868) from the Nebraska Transposon Mutant Library were acquired from BEI resources (https://www.beiresources.org/). To ensure true isogenic ancestors, the *mecA*-tn was transduced into the USA300 JE2 background, selecting for the erythromycin resistance marker on the Mariner transposon, and the insertion site was confirmed by polymerase chain reaction. Transposon maintenance was confirmed by testing for resistance to erythromycin (5µg/mL) or by genome sequencing when available. In cases where transposon loss was verified by genome sequencing, strains were tested and shown to maintain sensitivity to oxacillin. As the primary function of the transposon was to knock out *mecA* function, loss of the transposon should not confound subsequent analyses as antibiotic sensitivity was maintained.

Caenorhabditis elegans strain N2 Bristol and *Escherichia coli* strain OP50 were provided by the Caenorhabditis Genetics Center, which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440). Nematodes were maintained on Nematode Growth Medium Lite (US Biological, Swampscott, MA) according to WormBook protocols (Stiernagle, 2006).

Experimental evolution
Selection for increased virulence of *S. aureus* (USA300 JE2 or USA300 JE2 *mecA-tn*) in *C. elegans* was performed by passaging *S. aureus* from dead hosts 24 hours post-infection. Half of the *S. aureus* populations were additionally subjected to antibiotic pressure during the passages before and after *C. elegans* selection with sub-MIC oxacillin exposure (0.03125µg/mL). This concentration was tested against both ancestral strains and did not substantially inhibit growth of the methicillin-sensitive USA300 JE2 *mecA*-tn strain.

For each passage of experimental evolution, *S. aureus* was grown in Brain Heart Infusion (BHI) broth with 4µg/mL colistin (to prevent *E. coli* contamination from the nematode intestine) ± 0.03125µg/mL oxacillin and incubated at 37°C overnight, shaking at 250rpm. The next day, 200µL of the overnight *S. aureus* culture was plated onto BHI agar and incubated at 28°C overnight. Approximately 1500 *C. elegans* were plated and allowed to consume *S. aureus* for 24 hours before 30 dead nematodes were picked from each plate. Nematodes were identified as dead by a lack of response to tapping by a platinum wire (Ford et al., 2017; Kim L Hoang et al., 2024). Bacterial extraction methods were adapted from (Vega and Gore, 2017). In brief, picked dead nematodes were washed once with M9 buffer supplemented with 0.1% Triton X-100, once with M9 buffer, then subjected to a 1:1000 bleach-M9 buffer solution for 15 minutes. Nematodes were then washed twice with M9 buffer and ground with a motorized pestle to extract *S. aureus* from the host intestine. *S aureus* populations evolving without hosts were obtained from nematode-free bacterial lawns with a loop. Populations of *S. aureus* were grown as previously before freezing and storage at -80°C. Twenty-five percent of each frozen culture was used as inoculum for the next round of experimental evolution. Preliminary experiments determined colony-forming units per milliliter dropped by approximately 50% after freezing. To maintain population diversity, the inoculum amount (25%), adjusted for freezing, was chosen to be within the dilution ratio presented in (Wahl et al., 2002) that minimizes the chance that rare beneficial mutations are lost. Twelve rounds of passage were completed for 2 *S. aureus* genotypes x 2 host treatments x 2 antibiotic treatments x 6 replicates = 48 populations. Whole populations were frozen for subsequent analyses. We conducted all assays to evaluate changes in evolved *S. aureus* at the population level instead of single isolates to represent the context under which *S. aureus* evolved in our experiment and generally how pathogens exist in nature and in hosts (Cordero et al., 2012; Launay et al., 2021; McAdam et al., 2011; Mei et al., 2021; Paterson et al., 2015). Population level assays also allow for potential interactions between genotypes that give rise to phenotypes

1067   like virulence (Ruiz-Bedoya et al., 2023) and antibiotic resistance (Azimi et al., 2020; Cordero et
1068   al., 2012) that would otherwise not be captured in single-isolate assays.
1069
Mortality assay
Nematodes were age-synchronized as in (Penley and Morran, 2018) and grown on *E.coli* OP50
until L4 larval stage (48 hours at 20˚C). Nematodes were subsequently washed off, counted, and
approximately 200 individuals were plated on *S. aureus* lawns on BHI plates. Another set of
nematodes were plated on *E. coli* OP50 to estimate the total number of nematodes. Plates of *S.
aureus* were prepared 24 hours prior by plating 200μL of an overnight population culture on BHI
agar and grown at 28˚C. After two days at 20˚C, plates were scored by counting live nematodes.
Host mortality data were analyzed using a generalized linear model with a binomial distribution
and followed by Tukey multiple-comparison tests to determine pairwise differences. To compare
variances across treatments, we conducted Levene's test for homogeneity of variance. All
statistical analyses, including those below, were conducted in R (version 4.2.0).
1081
Hemolysis phenotype assay
Populations of *S. aureus* were streaked onto Trypticase Soy Agar II with 5% sheep's blood plates
and incubated overnight at 37˚C. Bacteria were assessed as either hemolysis positive or negative
depending on whether there was a complete clearing around the colonies. We did not cross-
streak with RN4220, which produces beta-hemolysin, and allows for detection of delta-hemolysin.
Therefore, our assay mainly detects alpha-toxin and phenol-soluble modulins. Hemolysis data
were analyzed using a generalized linear model with a binomial distribution. To determine whether
there was an association between hemolysis status and host mortality, we compared the host
mortality means of hemolysis-positive vs. hemolysis-negative pathogens using a Kruskal–Wallis
test.
1092
1093   To determine whether there was variation in hemolysis ability within populations, we streaked 10
1094   colonies from each of the 48 evolved populations onto Trypticase Soy Agar II with 5% sheep's
1095   blood plates and incubated overnight at 37˚C. To compare the number of isolates able to
1096   hemolyze, we conducted a generalized linear model with a binomial distribution. To compare
1097   variances, we conducted Levene's test for homogeneity of variance.
1098
***In vitro* growth assay**
We measured growth in BHI to simulate conditions under which pathogens were grown in our
experiment. Rich media allowed us to determine growth when conditions are optimal for evolved
pathogens. Measuring growth in minimal media, while being more similar to conditions that *S.
aureus* may face in its environment, would introduce another selective pressure. Overnight
cultures of each evolved population was diluted 1:1000 in BHI broth, then 200μl of each dilution
was added to 96-well plates, covered with a Breathe-Easy sealing membrane (Sigma) to minimize
evaporation. The optical density at 600nm at 37˚C with shaking at 282 cycles per minute was
recorded every 10 minutes for 16 hours (1000 minutes) using a BioTek Synergy HTX Multimode
Reader. We randomly assigned each of the 48 evolved populations to the inner wells of the 96-
well plates, with at least three technical replicates per population. Each plate also included
technical replicates of each ancestor. Each ancestral strain was assayed with four technical
replicates per plate across four plates. These steps were repeated for media with 0.03125μg/mL
oxacillin. We used the R package *gcplyr* (Blazanin, 2023) to calculate the area under the curve of
density, a metric that quantifies the overall bacterial growth. We compared the growth of
populations across treatments using the mean area under the curve of the technical replicates for
each population. We used a linear model followed by Tukey multiple-comparison tests to
determine pairwise differences between the ancestors, and linear mixed models for evolved
populations grown with or without oxacillin.

1118
Antimicrobial susceptibility assay
Antimicrobial susceptibility testing for oxacillin was done according to Clinical & Laboratory
Standards Institute (CLSI) standard protocols for broth microdilution (CLSI, n.d.). In brief,
populations were streaked from frozen stock and re-streaked the next day. From the second day
culture, populations were resuspended in normal saline to a 0.5 McFarland standard. Then, these
cultures were diluted 1:20 in normal saline before 10µL was added to 90µL of CAMHB + 2% NaCl
with the appropriate concentration of oxacillin. Plates were incubated at 35˚C without shaking and
read at 24 hours. For statistical analysis of populations evolved from the MSSA ancestor, we set
a threshold such that anything above the ancestral susceptibility level (0.25ug/ml) was considered
a decrease in oxacillin sensitivity. We then compared the proportion of populations exhibiting a
decrease in oxacillin sensitivity with those that had not using Fisher's exact test with false
discovery rate-adjusted (FDR) p-values.
1131
To determine whether there was variation in antibiotic susceptibility within evolved populations,
we sampled 10 colonies from each of the four populations with the most number of mutations
(two from MRSA and two from MSSA), and two additional randomly selected populations. We
argued that variation would be the most observable in populations with the most genetic diversity.
We then followed the standard protocol for a Kirby-Bauer disk diffusion susceptibility test
(Hudzicki, 2009) using oxacillin. Briefly, we inoculated Mueller-Hinton agar plates (Hardy
Diagnostics) of overnight cultures with standardized OD_{600} and placed an oxacillin-impregnated
disk (HardyDisk AST Oxacillin, OX1) onto each plate. We measured the diameter of the zone of
inhibition after incubation at 37˚C overnight.
1141
Biofilm assay
Overnight cultures of each population were diluted 1:40 in BHI broth containing 0.5% glucose,
then 100µl of each dilution was added to the inner wells of 96-well plates (NEST Biotechnology,
flat-bottom, tissue culture-treated). Plates were incubated for 24 hours at 20˚C, the temperature
that nematodes experience during infection. Each well was washed three times with 200µl of
phosphate-buffered saline (PBS) to remove non-adherent cells. We then stained cells with 125µl
of 0.1% crystal violet for 20 minutes, followed by washing each well twice with 200µl PBS. We
then resuspended the crystal violet in 125µl of 30% acetic acid. After incubation for 10 minutes,
we measured the optical density at 595nm using a BioTek Synergy HTX Multimode Reader. We
randomly assigned each of the 48 evolved populations to the inner wells of the 96-well plates,
with at least three technical replicates per population. Each plate also included technical replicates
of each ancestor. We standardized the mean of technical replicates for each population against
the respective ancestor within each plate to control for variation across plates. We used mixed
linear models followed by Tukey multiple-comparison tests to assess pairwise differences
between treatments.
1157
Whole-genome shotgun sequencing and variant annotation
Bacterial genome DNA was extracted from whole populations using Wizard Genomic DNA
Purification Kit (Promega). Population sequencing allows us to capture the genetic variation within
populations, which is critical to understanding how pathogens evolve (Azimi et al., 2020; Ford et
al., 2017, 2016; Kim L Hoang et al., 2024; Kim L. Hoang et al., 2024; Mahrt et al., 2021). Genome
sequencing of these DNA samples was performed at the M-Gen center using a paired-end library
preparation method based on Illumina Nextera and sequenced on the NextSeq 550.
1165
1166 The raw data was quality trimmed and adapters removed using illumina-cleanup
1167 (https://github.com/rpetit3/illumina-cleanup) or with bactopia (Petit and Read, 2020). Mutations
1168 were called based on comparison to the reference USA300 JE2 genome (BioProject:

PRJNA224116, BioSample: SAMN06677988, Assembly: GCF_002085525.1) using breseq (Deatherage and Barrick, 2014). The presence of large-scale deletions and other genetic changes were confirmed by analysis of de novo assembled genomes performed by the SKESA assembler software (Souvorov et al., 2018). The ancestral populations were almost identical to the reference genome used except for three mutations that were not found in any evolved populations. Another three mutations in *bioD1, cls* and *ccdC* were present in all populations derived from the MSSA ancestor and likely occurred in the first passage from the frozen culture before experimental evolution was performed. We therefore removed these six mutations from the analysis. Raw sequences have been deposited in the NCBI Sequence Read Archive under the BioProject accession number PRJNA1162848.

The number of mutations (excluding synonymous or intergenic regions) across treatments were analyzed using a generalized linear model with a Poisson distribution. For mutation enrichment analysis, we followed the steps as described previously (Lees et al., 2017). Briefly, we calculated the "mutation rate" for each ancestor by dividing the total number of mutations by the reference genome size. We multiplied the mutation rate by the total length of regulatory genes (Ibarra et al., 2013) with mutations in our study to generate the expected number of mutations for these genes. We then conducted single-tailed Poisson tests to determine whether the actual number of mutations in regulatory genes was greater than expected.

For phenotypes *vs.* mutations, we considered only mutations occurring at least twice throughout the experiment. For host mortality *vs.* mutations and growth rate *vs.* SCC*mec* and ACME, we conducted a one sample t-test for each evolved pathogen against its respective ancestor. P-values were adjusted using the FDR. For hemolysis status *vs.* mutations and oxacillin MIC *vs.* mutations, we conducted Fisher's exact tests with Bonferroni corrections.

Comparing mutations in experiment against publicly available genomes
For each non-synonymous mutation in each gene, we created the FASTA file of the USA300 JE2 protein product and a cognate file of the mutant protein with the amino acid substitution edited manually. We ran tBLASTn (version 2.12.0) searches of both proteins against 83,383 high quality *S. aureus* whole genome assemblies (Raghuram et al., 2024). We filtered out matches to genomes with less than 95% identity. We screened for cases where mutants had a better match to a genome than the ancestral protein (higher bitscore).

We collected meta-data for BioSamples accession linked to each genome from NCBI Entrez using a Python script from Jason Stajich (https://github.com/stajichlab/biosample_metadata), then manually curated the "isolation_source" column to either blood/systemic infection, skin/nose/throat, general host-association, animal, environment, others, and missing (see Table S4 for specific terms). For BioSamples matching mutants within each treatment, we compared the proportion of blood/systemic infection *vs.* skin/nose/throat against the expected proportion (all BioSamples in the dataset) using a chi-square goodness-of-fit test.

Principal component analysis
We conducted a principal component analysis on the phenotypes quantified in evolved populations using the prcomp function in R for each *S. aureus* genotype. We performed permutational analysis of variances (PERMANOVA, 999 permutations) to test for differences between treatments using adonis2 of the *vegan* package (Oksanen et al., 2024) and pairwiseAdonis (Martinez Arbizu, 2020).

Genetic distance

We generated a matrix of Euclidian genetic distances (the square root of pairwise differences)
using the frequencies of mutations that arose for each *S. aureus* genotype, from which we
constructed phylogenies (Betts et al., 2018; Ford et al., 2017). We used pairwise differences
between the ancestor and each evolved populations to compare rates of evolution across
treatments, and pairwise differences between replicate populations within each treatment to
compare the degree of divergence across treatments. We compared rates of evolution for MRSA
and MSSA treatments, and degree of divergence for MSSA treatments, using linear models
followed by Tukey multiple-comparison tests. We compared the degree of divergence for MRSA
treatments using a Kruskal–Wallis test followed by Dunn's post-hoc test.

SUPPLEMENTAL FIGURES
2

3
4   **Figure S1**. Most colonies sampled matched their respective population in terms of the
5   ability to hemolyze sheep's blood (*i.e.,* over 50% of colonies having the same hemolysis
6   status as the population they were sampled from).

9 **Figure S2.** Zone of inhibition from Kirby-Bauer disk diffusion susceptibility test with oxacillin
10 for colonies sampled from each of the four populations with the most number of mutations
11 (two from MRSA and two from MSSA), and two additional randomly selected populations.

13
14 **Figure S3**. Count of all mutations (from 10-100% frequency) arisen in evolved populations.

16
17
18 **Figure S4.** Mutations at frequencies between 0.1 and < 1, excluding synonymous or
19 intergenic mutations, in each evolved population. Points with darker shades indicate more
20 than one mutation present. All mutations in MRSA +host +ox populations between 0.1 and <
21 1 are all either synonymous or intergenic. A table of all mutations and frequencies is
22 available on figshare (10.6084/m9.figshare.28745558)

23
24   **Figure S5**. A) Count and B) Mean of mutations swept to fixation in evolved populations.

Figure S6. Genetic distance between the ancestor and evolved populations for A) MRSA
and B) MSSA genotypes. Genetic distance between replicate populations within each
treatment for C) MRSA and D) MSSA genotypes. Different letters indicate significant
differences. Error bars indicate standard errors.

SUPPLEMENTAL TABLES

32 **Table S1**. Nonsynonymous mutations occurred in genes with known roles in virulence and
33 antibiotic resistance.

Gene	Role identified in previous studies	Number of mutations in our study	Notes
agr	regulator, virulence [1]	22	14 in MRSA 18 in -OX 10 in +HOST
saeRS	regulator, virulence [2]	4	3 in MRSA
codY	regulator, metabolism and virulence [3]	8	6 in MSSA 8 in +HOST 7 in +OX
graSR	regulator, virulence [4] and resistance to beta-lactam antibiotics [5]	4	4 in MRSA 4 in +OX
pbpA	resistance to beta-lactam antibiotics [6]	4	2 in MRSA 2 MSSA 4 in +OX
pbpB	resistance to beta-lactam antibiotics [7]	4	4 MRSA 4 +OX
gdpP	resistance to beta-lactam antibiotics [8]	4	1 in MRSA 3 in MSSA 4 in +OX
SCC*mec and* Arginine Catabolic Mobile Element (ACME)	resistance to beta-lactam antibiotics [9]	9	5 SCC*mec* (23.7 kb deletion) 4 also included ACME (54.7 kb deletion) 6 in MSSA 7 in +HOST 1 *arcA* (99 bp deletion, encoded by ACME) in MRSA
brnQ1	metabolism [10]	20	16 in -HOST
purr	virulence [11]	4	4 in +HOST 3 in MRSA 3 in +OX
gpmA	virulence [12]	2	2 in +HOST 2 in MSSA 2 in -OX

35 **Table S2**. Fixed mutations in genes and intergenic regions appearing in more than two
36 populations. SYN = synonymous, NONSYN = nonsynonymous
37

Gene	Evolution treatment	Population	Position	Ancestral codon	Evolved codon	Mutation type	Mutation category	Amino acid change
ACME and *SCCmec*	MRSA -host -ox	three	33633	54659-bp		DEL	indel	
	MSSA +host +ox	three	33633	54659-bp		DEL	indel	
	MSSA +host -ox	five	33633	54659-bp		DEL	indel	
		six	33633	54659-bp		DEL	indel	
B7H15_10520	MRSA +host +ox	six	2031543	T	C	SNP	SYN	I19I
	MRSA -host +ox	three	2031543	T	C	SNP	SYN	I19I
SCCmec	MRSA +host -ox	three	33636	23687-bp		DEL	indel	
		six	33636	23687-bp		DEL	indel	
	MSSA +host -ox	four	33636	23687-bp		DEL	indel	
		five	33636	23687-bp		DEL	indel	
	MSSA -host +ox	two	33636	23687-bp		DEL	indel	
acetyl-CoA synthetase/ antibiotic biosynthesis monooxy- genase	MRSA +host -ox	two	2749064	C	A	SNP	intergenic	
	MSSA +host -ox	one	2749081	T	A	SNP	intergenic	
agr	MRSA +host +ox	six	2151051	C	T	SNP	NONSYN	S164L
		six	2151048	A	G	SNP	NONSYN	E163G
	MRSA +host -ox	one	2151213	G	T	SNP	NONSYN	R218L
		two	2150580	A	C	SNP	NONSYN	E7A
		four	2150063	A	T	SNP	other	K256*
	MRSA +host -ox	five	2151272	A		DEL	indel	
		six	2150931	T		DEL	indel	
	MRSA -host +ox	three	2151051	C	T	SNP	NONSYN	S164L
		three	2151048	A	G	SNP	NONSYN	E163G
	MRSA -host -ox	two	2149664	A		DEL	indel	
		three	2151047	G	T	SNP	other	E163*
		four	2151272	A		DEL	indel	
		five	2149476	G	A	SNP	other	W60*
		six	2151212	C	T	SNP	other	R218*
	MSSA +host -ox	four	2150250	G	A	SNP	NONSYN	G318D
		five	2151233	A	T	SNP	other	K225*
		six	2150298	C	A	SNP	NONSYN	P334H
	MSSA -host -ox	one	2149635	C	A	SNP	other	S113*
		three	2150803	T	CAGTT	INS	indel	
		four	2149989	C	A	SNP	NONSYN	T231K
		six	2150700	C	A	SNP	NONSYN	A47D
		six	2150691	TT	AA	SUB	indel	

agrD/C	MRSA -host +ox	one	2149252	G	A	SNP	intergenic	
	MSSA -host -ox	two	2149281	T	G	SNP	intergenic	
alr	MRSA -host +ox	two	2190276	A	G	SNP	NONSYN	L280S
		six	2190276	A	G	SNP	NONSYN	L280S
	MRSA -host -ox	one	2190276	A	G	SNP	NONSYN	L280S
argR	MRSA +host +ox	four	1639665	G	C	SNP	NONSYN	T99R
	MRSA -host +ox	two	1639510	A	T	SNP	NONSYN	*151K
		six	1639510	A	T	SNP	NONSYN	*151K
	MRSA -host -ox	one	1639510	A	T	SNP	NONSYN	*151K
arginine repressor/- geranyl transferase	MRSA +host -ox	three	1639991	C	T	SNP	intergenic	
	MSSA +host -ox	four	1639991	C	T	SNP	intergenic	
		six	1639991	C	T	SNP	intergenic	
brnQ1	MRSA -host -ox	two	218284	A	G	SNP	NONSYN	L154S
		three	218456	A	T	SNP	NONSYN	F97I
		three	218451	G		DEL	indel	
		four	218425	108-bp		DEL	indel	
		five	217434	GCCGATAATGAA		DEL	indel	
		six	217538	97-bp		DEL	indel	
	MSSA +host +ox	three	218420	CA	AT	SUB	indel	
	MSSA +host -ox	two	218581	A	T	SNP	other	L55*
		three	217879	G	T	SNP	other	S289*
		five	218644	C	A	SNP	NONSYN	G34V
	MSSA -host +ox	one	218035	C	T	SNP	NONSYN	G237D
		two	218420	CA	AT	SUB	indel	
		three	218420	CA	AT	SUB	indel	
		four	218035	C	T	SNP	NONSYN	G237D
		five	218420	CA	AT	SUB	indel	
		six	218035	C	T	SNP	NONSYN	G237D
	MSSA -host -ox	one	218128	GCTAGCGCAT		DEL	indel	
		two	218644	C	A	SNP	NONSYN	G34V
		three	218644	C	A	SNP	NONSYN	G34V
		six	218644	C	A	SNP	NONSYN	G34V
codY	MRSA +host +ox	two	1275011	C	T	SNP	NONSYN	H159Y
		six	1274675	A	G	SNP	NONSYN	K47E
		one	1275155	G	T	SNP	NONSYN	A207S
		two	1275155	G	T	SNP	NONSYN	A207S
	MSSA +host +ox	four	1275155	G	T	SNP	NONSYN	A207S
		five	1275155	G	T	SNP	NONSYN	A207S
		six	1275155	G	T	SNP	NONSYN	A207S
	MSSA +host -ox	two	1274976	C	T	SNP	NONSYN	A147V
	MRSA +host +ox	six	87329	T	A	SNP	SYN	L624L

copper-translocating P-type ATPase	MRSA -host +ox	three	87329	T	A	SNP	SYN	L624L
ebh	MSSA +host -ox	five	1477894	C	T	SNP	NONSYN	E8357K
	MSSA -host -ox	two	1480506	G	A	SNP	NONSYN	P7486L
fmtB	MRSA +host +ox	six	2284788	G	T	SNP	other	
	MRSA -host +ox	three	2284788	G	T	SNP	other	
gdpP	MRSA +host +ox	three	18721	G	T	SNP	NONSYN	G307V
	MSSA +host +ox	one	18720	G	A	SNP	NONSYN	G307S
		four	18745	A	C	SNP	NONSYN	K315T
		five	19143	A	T	SNP	NONSYN	S448C
gpmA	MSSA +host -ox	one	2539086	G	A	SNP	NONSYN	R114C
	MSSA +host -ox	three	2538887	G	A	SNP	NONSYN	S180F
graSR	MRSA +host +ox	three	734510	A		DEL	indel	
		four	735131	C	G	SNP	other	S187*
		six	734085	T		DEL	indel	
	MRSA -host +ox	three	734085	T		DEL	indel	
Hydroxy-methyl-glutaryl-CoA synthase/-cysteine methyl-transferase	MRSA +host +ox	six	2685330	T	C	SNP	intergenic	
	MRSA -host +ox	three	2685330	T	C	SNP	intergenic	
pbpA	MRSA +host +ox	one	1189756	C	A	SNP	NONSYN	D350E
		four	1190201	C	G	SNP	NONSYN	H499D
	MSSA +host +ox	three	1189756	C	A	SNP	NONSYN	D350E
		six	1189758	G	T	SNP	NONSYN	W351L
pbpB	MRSA +host +ox	two	1520209	G	A	SNP	NONSYN	A132T
		five	1520434	G	A	SNP	NONSYN	A207T
	MRSA -host +ox	one	1520247	A	GGTTCTGAA	INS	indel	
		four	1520414	G	T	SNP	NONSYN	G200V
purR	MRSA +host +ox	one	532212	C	T	SNP	NONSYN	R96C
		three	532013	T	T	INS	indel	
		five	532013	T	T	INS	indel	
	MRSA -host +ox	four	532240	T	T	INS	indel	
saeRS	MRSA +host +ox	two	782595	G	T	SNP	NONSYN	P74T
		five	782577	C	T	SNP	NONSYN	A80T
	MRSA +host -ox	two	781408	G	A	SNP	NONSYN	R241C
	MSSA +host +ox	six	782211	C		DEL	indel	

39 **Table S3.** Chi-square goodness-of-fit test results for comparison between the proportion of
40 blood/systemic infection-associated mutations *vs.* skin/nose/throat-associated mutations
41 against the expected proportion (*i.e.,* all BioSamples in the dataset). All degrees of freedom
42 equalled 1.

treatment	gene	chi-square	P-value
MRSA +HOST +OX	*pbpA*	21.626	< 0.001
MSSA +HOST +OX	*pbpA*	26.431	< 0.001
MRSA +HOST +OX	*saeRS*	4.8088	0.02832
MRSA +HOST +OX	*purR*	4.3463	0.03709
MRSA +HOST +OX	*pbpb*	11	0.000911
MRSA -HOST +OX	*pbpb*	8.2065	0.004174
MRSA -HOST -OX	*brnQ1*	27.748	< 0.001
MSSA +HOST -OX	*brnQ1*	27.748	< 0.001
MSSA -HOST -OX	*brnQ1*	19.773	< 0.001
MSSA -HOST +OX	*brnQ1*	23.097	< 0.001
MRSA +HOST +OX	*codY*	0.19574	0.6582
MSSA +HOST -OX	*codY*	1.6584	0.2914
MRSA +HOST +OX	*graSR*	0.19574	0.6582
MRSA +HOST +OX	*gdpP*	2.8047	0.09399
MRSA +HOST +OX	*agr*	0.3538	0.552
MRSA +HOST -OX	*agr*	0.35066	0.5537
MRSA -HOST +OX	*agr*	0.3538	0.552
MRSA -HOST -OX	*agr*	0.223	0.6368
MSSA +HOST -OX	*agr*	0.34068	
MSSA -HOST -OX	*agr*	0.2662	0.6059
MRSA -HOST -OX	*argR*	0.223	0.6368
MRSA -HOST +OX	*argR*	0.3538	0.552

Table S4. Categories for "isolation_source" terms from meta-data for BioSamples linked to
 our dataset of public *S. aureus* genomes in Figure 5A.

Category	"isolation_source" term
blood/systemic	blood
	bacteremia
	csf
	bronchial
	aortic
	osteomyelitis
	respiratory
	sputum
	bronch
	septicaemia
	pericar
	kidney
	liver
	spleen
	brain
	bone
	lung
	spine
	spinal
	periton
	pancreas
	joint
	trachea
	urethra
	uterus
	airway fluid
	bile
	ascitic fluid
	bsi
	bal
	bone marrow
	broncoscopy
	bronscoscopy
	cerebrospinal fluid
	endotracheal aspirate
	endotrracheal secretion
	trachael aspirate
	expectoration
	hematoma
	pleural fluid
	septic arthritis
	skeletal system
	synovial fluid
	thoracic cavity

	thymic pleural effusion tissue - aortic valve yolk sac infection urinary tract infection arthritis aspirates necrotizing "wound culture from necrotizing fasciitis patient"
skin/nose/throat	abcess abscess asbcess absess nare naso sinus nasal nostril cellulitis wound burn sore pus ulcer lesion swab screen abdom buttock eye groin leg arm back oral hand cheek chest elbow foot femur axilla knee neck ankle thigh cornea graft umbillicus

		umbilicus
		breast
		ear
		ulcus cruris
		darcocystitis
		decubitus ulcer
		empyema
		face
		fluid breast
		granuloma
		hip - left
		hip infection
		inguinal
		left hip aspiration
		oropharynx
		palm
		penis
		perineal
		perineum
		periodontal
		rectum
		sample from soft tissue
		soft tissue
		scapula
		secretion left hip
		ssti
general host-association		biological sample
		biological liquide
		aspirate
		aspiration
		biopsy
		excreted bodily substance
		tissue
		milk
		urine
		bodily fluid
		vaginal
		feces
		stool
		tissues
		body fluid
		faecal
		infection
		infection site
		mass
		post surgical secretion
		secretion surgical

		surgical site infection
		chronic
		colonisation
		colonization
		commensal
		community aquired
		clinical
		human
		patient
		male
		hcw
		host
animal		chimpanzee
		cloudrat
		corncrake
		cow
		sheep
		macaque
		lion
		bat
		chicken
		pork
		horse
		canine
		pig
		poultry
		meat
		capybara
		chaffinch
		meerkat
		mongoose
		monkey
		parrot
		pheasant
		porpoise
		rabbit
		seal
		sparrow
		tapir
		wildbird
		turkey
		cat
		goat
		great tit
		ground beef
		ground turkey
		guineapig

		claw
		paw
		tail
		diced chiken
		feline colonisation
		"rat feces from intestine (rattus norvegicus)"
		veal calf
		veterinary/diagnostic sample
		non-migratory seabirds
	environment	various material
		cryotube from air metagenome
		fish drying yard
		glove
		gown
		hardware
		heart valve
		hexachlorocyclohexane-contaminated soil
		household surface
		jp drainage
		catheter
		minibal
		pacemaker wire
		peg tube drainage
		peri-bypass anterior material
		pin tract
		surface veterinary clinic
		surgical ward
		suture
	others	culture
		lab
		physical
		food
		facility
		acute
		atcc
		isolate
		hospital
		in vitro derived
		in vitro evolution
		blops
		case
		drain
		drainage
		fluid
		icu
		index
		mara

		mrsa broth
		non-icu
		norway
		ny
		clinic
		pool
		pooled
		presumed outlier
		serial passagee experiment
		staphylococcus aureus usa300
		surface
		surgical site
		tip
missing		not applicable
		not collected
		not known
		unknown
		NA
		missing
		not avail
		unknonwn
		unknow
		unspecified

REFERENCES

1. Traber KE, Lee E, Benon S, Corrigan R, Cantera M, Shopsin B, et al. agr function in clinical Staphylococcus aureus isolates. Microbiology. 2008;154: 2265–2274. doi:10.1099/mic.0.2007/011874-0

2. Liang X, Yu C, Sun J, Liu H, Landwehr C, Holmes D, et al. Inactivation of a two-component signal transduction system, SaeRS, eliminates adherence and attenuates virulence of Staphylococcus aureus. Infect Immun. 2006;74: 4655–4665. doi:10.1128/IAI.00322-06

3. Majerczyk CD, Sadykov MR, Luong TT, Lee C, Somerville GA, Sonenshein AL. Staphylococcus aureus CodY negatively regulates virulence gene expression. J Bacteriol. 2008;190: 2257–2265. doi:10.1128/JB.01545-07

4. Falord M, Mäder U, Hiron A, Dbarbouillé M, Msadek T. Investigation of the Staphylococcus aureus GraSR regulon reveals novel links to virulence, stress response and cell wall signal transduction pathways. PLoS One. 2011;6. doi:10.1371/journal.pone.0021323

5. Müller A, Grein F, Otto A, Gries K, Orlov D, Zarubaev V, et al. Differential daptomycin resistance development in Staphylococcus aureus strains with active and mutated gra regulatory systems. Int J Med Microbiol. 2018;308: 335–348. doi:10.1016/j.ijmm.2017.12.002

6. Berti AD, Theisen E, Sauer JD, Nonejuie P, Olson J, Pogliano J, et al. Penicillin binding protein 1 is important in the compensatory response of Staphylococcus aureus to daptomycin-induced membrane damage and is a potential target for β-lactam-daptomycin synergy. Antimicrob Agents Chemother. 2016;60: 451–458. doi:10.1128/AAC.02071-15

7. Łęski TA, Tomasz A. Role of penicillin-binding protein 2 (PBP2) in the antibiotic susceptibility and cell wall cross-linking of Staphylococcus aureus: Evidence for the cooperative functioning of PBP2, PBP4, and PBP2A. J Bacteriol. 2005;187: 1815–1824. doi:10.1128/JB.187.5.1815-1824.2005

8. Sommer A, Fuchs S, Layer F, Schaudinn C, Weber RE, Richard H, et al. Mutations in the gdpp gene are a clinically relevant mechanism for β-lactam resistance in meticillin-resistant staphylococcus aureus lacking mec determinants. Microb Genomics. 2021;7: 1–13. doi:10.1099/MGEN.0.000623

9. Shore AC, Coleman DC. Staphylococcal cassette chromosome mec: Recent advances and new insights. Int J Med Microbiol. 2013;303: 350–359. doi:10.1016/j.ijmm.2013.02.002

10. Kaiser JC, Omer S, Sheldon JR, Welch I, Heinrichs DE. Role of BrnQ1 and BrnQ2 in branched-chain amino acid transport and virulence in Staphylococcus aureus. Infect Immun. 2015;83: 1019–1029. doi:10.1128/IAI.02542-14

11. Goncheva MI, Flannagan RS, Heinrichs DE. De Novo Purine Biosynthesis Is Required for Intracellular Growth of Staphylococcus aureus and for the Hypervirulence Phenotype of a purR Mutant. Infect Immun. 2020.

12. Radin JN, Kelliher JL, Solórzano PKP, Grim KP, Ramezanifard R, Slauch JM, et al. Metal-independent variants of phosphoglycerate mutase promote resistance to nutritional immunity and retention of glycolysis during infection. PLoS Pathog. 2019;15: 1–21. doi:10.1371/journal.ppat.1007971