TITLE
Host and antibiotic jointly select for greater virulence in *Staphylococcus aureus*

AUTHORS
Michelle Su*[1], Kim L. Hoang*[1], McKenna Penley[2], Michelle H. Davis[1], Jennifer D. Gresham[2,3], Levi T. Morran[2], Timothy D. Read[1]

*these authors contributed equally

AFFILIATIONS
1. Emory University School of Medicine, Division of Infectious Diseases, Atlanta, Georgia, USA
2. Department of Biology, Emory University, Atlanta, Georgia, USA
3. Science Department, Penn State Scranton, Dunmore, Pennsylvania, USA

CORRESPONDING AUTHOR
TDR: tread@emory.edu

ACKNOWLEDGEMENTS
The authors are grateful for insightful discussion with members of the Read and Morran labs, and Shaun Brinsmade for reading the manuscript. We thank Carmen Alvarez, Jason Chen, and Emily Stevens for help with experimental protocols.

FUNDING
MS was supported by the Antimicrobial Resistance and Therapeutic Discovery Training Program funded by NIAID T32 award AI106699. TDR was supported by National Institutes of Health (NIH) AI139188 and AI158452. KLH and TDR were supported by the Office of Advanced Molecular Detection, Centers for Disease Control and Prevention Cooperative Agreement Number CK22-2204 through contract 40500-050- 23234506 from the Georgia Department of Public Health. LTM was supported by a National Science Foundation Division of Environmental Biology (NSF-DEB) grant 1750553.

AUTHOR CONTRIBUTIONS
MS, LTM, and TDR conceptualized the project. MS, KLH, MP, MHD, and JDG conducted the experiments. KLH and MS analyzed the data and drafted the manuscript. All authors contributed to manuscript revision.

DATA AVAILABILITY
Raw sequences have been deposited in the NCBI Sequence Read Archive under the BioProject accession number PRJNA1162848, and phenotypic data deposited in the Figshare Repository (10.6084/m9.figshare.28745558).

CONFLICTS OF INTERESTS
The authors declare no conflicts of interests.

ABSTRACT

Widespread antibiotic usage has resulted in the rapid evolution of drug-resistant bacterial pathogens and poses significant threats to public health. Resolving how pathogens respond to antibiotics under different contexts is critical for understanding disease emergence and evolution going forward. The impact of antibiotics has been demonstrated most directly through *in vitro* pathogen passaging experiments. Independent from antibiotic selection, interactions with hosts have also altered the evolutionary trajectories and fitness landscapes of pathogens, shaping infectious disease outcomes. However, it is unclear how interactions between hosts and antibiotics impact the evolution of pathogen virulence. Here, we evolved and re-sequenced *Staphylococcus aureus,* a major bacterial pathogen, varying exposure to host and antibiotics to tease apart the contributions of these selective pressures on pathogen adaptation. After 12 passages, *S. aureus* evolving in *Caenorhabditis elegans* nematodes exposed to a sub-minimum inhibitory concentration of antibiotic (oxacillin) became highly virulent, regardless of whether the ancestral pathogen was methicillin-resistant (MRSA) or methicillin-sensitive (MSSA). Host and antibiotic exposure selected for reduced drug susceptibility in MSSA lineages while increasing MRSA total growth outside hosts. We identified mutations in genes involved in complex regulatory networks linking virulence and metabolism, including *codY*, *agr*, and *gdpP*, suggesting that rapid adaptation to infect hosts may have pleiotropic effects. In particular, MSSA populations under selection from host and antibiotic accumulated mutations in the global regulator gene *codY*, which controls biofilm formation in *S. aureus.* These populations had indeed evolved more robust biofilms—a trait linked to both virulence and antibiotic resistance—suggesting evolution of one trait can confer multiple adaptive benefits. Mutations that arose in these genes were also enriched in clinical isolates associated with systemic infections in humans. Despite evolving in similar environments, MRSA and MSSA populations proceeded on divergent evolutionary paths, with MSSA populations exhibiting more similarities across replicate populations. Our results underscore the importance of considering the host context as a critical driver of pathogen traits like virulence and antibiotic resistance.

KEYWORDS

Antibiotic resistance, virulence evolution, experimental evolution, *Staphylococcus aureus*, pathogen adaptation

INTRODUCTION

Antibiotic resistance is a global crisis [1]. Pathogens that evolve resistance and are able to survive antibiotic therapy can go on to cause disease and transmit to other hosts; hence antibiotic resistance and virulence are intertwined. Selective pressures like antibiotic usage and host defenses have independently been shown to alter the evolution of pathogen traits [1–5]. However, pathogens face multiple selective pressures in their environment, and selection by concurrent forces can interact synergistically to alter evolutionary rates and trajectories relative to individual pressures alone [6,7]. It is unclear how virulence evolves in the face of simultaneous selective pressures and whether this trait is constrained or facilitated by antibiotic resistance. These dynamics are even less understood in the early stages of disease emergence, where pathogens face a different suite of selection dynamics than pathogens having reached equilibrium [8,9]. Identifying conditions under which antibiotic resistance and virulence evolve in novel host populations will shed light on how diseases emerge and potential mitigation measures.

The strength of selection impacts pathogen evolution [10–13]. While antibiotic therapy administers concentrations that should kill all pathogen cells, low concentrations of antibiotics (sub-minimum inhibitory concentration, or sub-MIC) are pervasive in a variety of settings, including in natural environments (e.g., water bodies and soil) due to pollution and biological waste [1]. In the host, antibiotic concentrations will gradually decline after administration due to metabolism and excretion [14–16]. Sub-MICs allow susceptible populations to continue dividing, affording opportunities for mutations conferring greater resistance to emerge [15]. Ultimately, selection by sub-MICs can lead to highly resistant pathogens [10,17], which tend to incur less fitness costs compared to those under selection by high antibiotic concentrations [18]. Importantly, mutations not related to antibiotic resistance can arise, such as those involved in adaptation to the growth environment [19]. Low concentrations of antibiotics can also alter expression of virulence factors *in vitro* across a broad range of pathogens [20–23], suggesting that they can affect virulence during infection. Sub-MIC antibiotics therefore have the potential to alter the early stages of disease emergence in host populations, particularly for those pathogens that can spend parts of their life history outside the host. Taken together, exposure to low antibiotic concentrations likely alters how pathogens interact with their hosts, shaping evolution of both virulence and antibiotic resistance. However, there is a knowledge gap in the interaction between antibiotics and pathogen evolution *in vivo* [16].

The host can exert strong selection on pathogens, especially during the period when a new organism is infected (e.g., a zoonotic transition). Infection of new host individuals tends to bottleneck pathogen populations [24,25]. These bottlenecks can confer a fitness advantage to resistant bacteria in the face of low antibiotic concentrations [26]. In addition to defenses like the immune system, the host environment differs drastically in nutrient availability compared to rich laboratory media [16]. For example, *Staphylococcus aureus* passaged through macrophage cell lines had increased pathogen survival as well as resistance to antibiotics; these traits were lost when the pathogen was exposed to nutrient-rich media [27]. Similarly, the human defensive peptide b-defensin 3 can maintain reduced susceptibility of *S. aureus* to the antibiotic vancomycin [28]. Adapting to different niches within a host can also bring about drastic genomic changes. *Salmonella enterica* transitioning from an intestinal to a systemic lifestyle exhibited genome degradation in genes no longer necessary for inhabiting the gastrointestinal tract [24]. While sub-MIC antibiotic exposure and host factors have been shown to independently shape pathogen evolution, it is unclear how the collective actions of both selective pressures affect the evolution of virulence and antibiotic resistance.

Staphylococcus aureus is an opportunistic pathogen that commonly colonizes nares and skin in humans but can invade internal organs and blood, causing systemic infections and bacteremia

[29]. In 2017, more than 119,000 bacteremia infections caused by *S. aureus* occurred in the United
States with a mortality rate of 18% [30]. *S. aureus* colonizing host surfaces exhibit distinct genomic
signatures against those isolated during systemic infection, indicating that the pathogen employs
different strategies to adapt to different host sites [31]. For example, hemolytic strains tend to be
more abundant than non-hemolytic ones in murine systemic infection models, whereas the
opposite occurs for wound models [32]. Once inside the host, *S. aureus* evades detection by the
immune system by hindering or lysing immune cells [33,34]. Sub-MIC concentrations of beta-
lactam antibiotics—which disrupts bacterial cell wall synthesis—modify expression of virulence
factors in *S. aureus*, increasing the expression of alpha-hemolysin [35]. Treatment with beta-
lactam antibiotics systemically exposes *S. aureus* to low antibiotic levels at non-target host tissues
[36]. The pathogen can also survive for extended periods of time outside of hosts and have been
isolated from environments such as veterinary clinics and schools, as well as natural and
constructed settings such as wastewater and beaches [37–40]. Combined with the increasing
prevalence and concentrations of antibiotics in the environment, these factors likely increase
opportunities for exposure of *S. aureus* to sub-MIC antibiotics, subsequently affecting its
interaction with the host during infection.
142
Experimental evolution has been used to elucidate how different selective pressures impact
pathogen evolution. *In vitro* studies have yielded insights into the phenotypes and genetic loci
generated by longer-term antibiotic selection through passaging experiments lasting hundreds of
generations [10,18,28,41]. Conversely, *in vivo* studies have focused on transmitting pathogens
through individual hosts for a single to a handful of passages [26,42,43]. However, few systems
have been suitable to examine how antibiotics affect pathogen adaptation within a host context.
In this study, we take advantage of the suitability of a multicellular host, *Caenorhabditis elegans*
nematodes, to study designs of high replication across an appreciable temporal scale. The *S.*
aureus used here was originally isolated from a skin and soft tissue infection of a human inmate
[44] and thus is a novel pathogen to *C. elegans*. Nonetheless, *S. aureus* can kill nematodes by
colonizing the host intestine and lysing cells, inducing expression of defense genes in nematodes
with roles conserved in humans [45,46]. Virulence screens in *C. elegans* using *S. aureus*
transposon libraries also showed overlapping results to other infection models [47,48], with the
degree of *S. aureus* virulence in *C. elegans* reflecting disease severity of human infections
[49,50]. Experimentally evolving *S. aureus* in *C. elegans* thus allows us to track the early stages
of virulence and antibiotic resistance evolution in novel host populations with the potential to
identify clinically-relevant genomic regions underlying evolved traits.
160
Here, we directly test the impact of host and sub-MIC antibiotic exposure on pathogen evolution.
Selection exerted by two concurrent forces may impede the pathogen's response to one or both
forces [6]. Adaptation may require resources to be expended toward either virulence or antibiotic
resistance, leading to a trade-off between these traits [51]. Alternatively, weaker selection from
sub-MIC antibiotics may interact synergistically with hosts and facilitate the evolution or
maintenance of high virulence and antibiotic resistance. Sub-MIC antibiotics can favor no-cost
mutations [18], wherein pathogens can rapidly adapt without affecting other traits [8]. Because
emerging pathogens tend to be far from the optimum of the fitness landscape, we expect
pleiotropic or large-effect mutations to play a role [52]. We took advantage of the tractability of the
system to determine how virulence, fitness, and antibiotic resistance are connected and how
multiple selective pressures shape the evolutionary trajectory of two *S. aureus* isolates differing
only in their antibiotic susceptibility.
173
Carriage of Staphylococcal cassette chromosome *mec* (SCC*mec*), which encodes *mecA* and
provides resistance against beta-lactam antibiotics, is a major mechanism for existing antibiotic
resistance in *S. aureus*. We passaged two *S. aureus* isogenic strains, one with existing resistance

(*mecA+*, MRSA) and one with a transposon insertion in *mecA* (*mecA-*, MSSA) under selection by
C. elegans and a sub-MIC level of oxacillin, a beta-lactam antibiotic that has replaced methicillin
as a therapy for Staphylococcal infections. We selected for virulence in both strains to determine
whether antibiotic resistance can hinder the evolution of greater virulence. We then quantified the
ability of evolved pathogens to kill hosts as the metric for virulence. We also assessed whether
populations maintained their ability to hemolyze red blood cells—an indicator of virulence
expression in both humans and nematodes. Because growth outside the host is important for
transmission of opportunistic pathogens, we quantified the *in vitro* growth and MIC of evolved
pathogens. Finally, we identified mutations potentially underlying evolved traits and compared
them against a database of over 80,000 *S. aureus* genomes to ascertained whether these
mutations may be associated with adaptation to different human host sites.
188
RESULTS
Host and sub-MIC antibiotic exposure selected for the greatest virulence in both MRSA
and MSSA.
We experimentally evolved two *S. aureus* isogenic variants of USA300 JE2 [44], MRSA and
MSSA, with or without a host and sub-MIC antibiotic exposure. The ancestral MRSA and MSSA
isolates did not differ in terms of total growth without oxacillin, but exhibited a slight decline in sub-
MIC oxacillin (Figures 1A and 1B; χ^2_1 = 6.39, P = 0.01). For each ancestor, we passaged six
independently evolving populations under each condition 12 times, for a total of 48 evolved *S.*
aureus populations (Figure 1C).
198

C

Figure 1. Experimental evolution design. A) Growth curves and B) total growth of ancestral
MRSA and MSSA populations *in vitro* with and without sub-MIC oxacillin. Different letters
indicate significant differences. C) MRSA and MSSA were passaged 12 times with or without
hosts, in the presence or absence of a sub-MIC of the antibiotic oxacillin. Each treatment
consisted of six independently evolving replicate populations. Experimental evolution treatment
abbreviations are indicated in the purple and yellow boxes. Error bars indicate standard errors.

To assess the changes in virulence of evolved *S. aureus*, we measured the mortality of *C. elegans*
infected with evolved pathogens. While the MRSA and MSSA ancestors caused similar levels of
host mortality (dotted and dashed lines in Figure 2A), there was a significant treatment effect for
evolved pathogens (Figure 2A; χ^2_7 = 460.43, P < 0.001). For the MRSA genotype, host and
oxacillin exposure selected for the greatest virulence (P < 0.001 for all Tukey's post-hoc tests of
+host+ox *vs.* other treatments). Conversely, pathogens evolved in the absence of either pressure
exhibited attenuated virulence (-host-ox *vs.* +host-ox: P < 0.001; -host-ox *vs.* -host+ox: P < 0.001).
These populations also had significantly greater variance (Levene's test for homogeneity of
variance $F_{3,20}$ = 4.06, P = 0.021) compared to those under selection from host and oxacillin
(Tukey's post-hoc test P = 0.026), and those under solely host selection (P = 0.045). For MSSA,
host and oxacillin exposure similarly favored greater virulence over the other three conditions (P
< 0.001 for all Tukey's post-hoc tests of +host+ox *vs.* other treatments). There was no difference
between variances (Levene's test for homogeneity of variance $F_{3,20}$ = 0.99, P = 0.417).

We also characterized the hemolytic activity of evolved populations, which correlates with the
ability to secrete extracellular toxins and virulence [53]. At least one population from most
treatments had lost the ancestral ability to hemolyze sheep's blood (Figure 2B). Sub-MIC oxacillin
maintained hemolytic activity (χ^2_1 = 5.82, P = 0.016), suggesting that constant exposure to low
concentrations of oxacillin favored retention of extracellular toxicity. While hemolysis was not
necessary for increased host-killing, this ability was correlated with greater nematode mortality
(Figure 2C; Kruskal-Wallis χ^2_1 = 3.97, P = 0.046). As virulence was still heightened in pathogens
unable to destroy red blood cells (e.g., MSSA pathogens evolving without host or oxacillin caused

greater than 90% mean mortality despite none being hemolysis-positive), other virulence factors
may be compensating for the absence of hemolysis.

Natural populations of *S. aureus* exhibit variation in hemolytic ability [54], even within an individual
host [55]. We thus hypothesized that a significant selective pressure like oxacillin would favor little
variation of this trait within a population. We determined the hemolysis status of isolates from
each evolved population (10 isolates x 48 populations)—most populations were in consensus
(Figure S1). Furthermore, oxacillin still has a significant effect, where hemolysis was maintained
when under sub-MIC exposure selection (Figure 2D; χ^2_1 = 23.30, P < 0.001). Oxacillin favored
less hemolytic diversity in MSSA populations (Levene's test for homogeneity of variance $F_{1,22}$ =
26.06, P < 0.001), further demonstrating the interaction between antibiotics and virulence, and
that antibiotics can shape variation in virulence traits.

Figure 2. Evolution of virulence is facilitated by exposure to both host and sub-MIC antibiotic. A) Virulence in terms of *C. elegans* mortality. Dashed lines indicate respective ancestral virulence. Different letters indicate significant differences. B) Virulence in terms of the ability to hemolyze sheep's blood, assayed at the population level. The y-axis indicates the number of evolved populations for each category. C). Host mortality from A) grouped by hemolysis status in B). D) The proportion of colonies sampled from each evolved population that are able to hemolyze sheep's blood. Error bars indicate standard errors. *P < 0.05

Host and sub-MIC antibiotic promoted growth of MRSA outside hosts and reduced drug susceptibility in MSSA.

We measured the *in vitro* growth of evolved populations to evaluate how pathogen fitness outside the host had been impacted. We used rich media to replicate the conditions under which *S. aureus* evolved during the experiment. Importantly, rich media prevents introduction of additional selective pressures than those being tested. In media without oxacillin, there were no significant differences in total growth between treatments (Figure 3A; χ_7^2 = 9.24, P = 0.24). However, in sub-MIC oxacillin, MRSA populations under selection from both pressures exhibited the greatest growth (Figure 3B; χ_7^2 = 106.16, P < 0.001), with some achieving more growth than those in the absence of oxacillin. Similarly, MSSA populations exposed to host and sub-MIC oxacillin exhibited a moderate increase in growth compared to other combinations of selective pressures.

We then determined the MICs of evolved populations to ascertain how host and sub-MIC exposure affected the evolution and maintenance of antibiotic resistance. For the MRSA genotype, all but two populations retained the ancestral level of oxacillin resistance (16ug/mL oxacillin; Figure 3C). During evolution in hosts without antibiotic selection, two populations lost their resistance and exhibited similar susceptibility as the MSSA ancestor (0.25ug/mL). For the MSSA genotype, host and oxacillin exposure selected for decreased antibiotic sensitivity (Fisher's exact test P < 0.001; pairwise post-hoc P < 0.05 for +host+ox *vs.* all other treatments), up to 8-fold the ancestral MIC. These results suggested sub-MIC exposure combined with host factors potentiated the increase in antibiotic resistance in MSSA. One population under selection from solely the host also had an increased MIC, supporting previous evidence showing non-antibiotic selective pressures, such as a host in our study, can select for reduced antibiotic susceptibility [56]. There was no variation in antibiotic resistance within MRSA populations and little variation within MSSA populations across the most genetically diverse populations (Figure S2).

279
280 **Figure 3**. **Host and sub-MIC antibiotic selection facilitated pathogen growth in antibiotics**.
281 Pathogen *in vitro* growth A) without oxacillin and B) in sub-MIC oxacillin. Different letters
282 indicate significant differences. C). Oxacillin MIC of evolved populations. The y-axis indicates
283 the number of evolved populations for each category. Error bars indicate standard errors. **P <
284 0.01
285
Parallel evolution of regulators of virulence and antibiotic resistance across evolved
populations.
We conducted whole-genome sequencing of populations to identify mutations arisen from host
and antibiotic selection. Below we focus on mutations that had swept to fixation (Figure 4A, Table
S1); Figures S3 and S4 summarize mutations that were below 100% frequency in each
population. Populations evolved from the MSSA ancestor had significantly fewer mutations
(excluding those intergenic or synonymous) than MRSA populations (Figure S5; χ_1^2 = 5.28, P =
0.022), potentially due to the slightly reduced growth of MSSA in the presence of oxacillin (Figure
1B). Sub-MIC oxacillin selection also resulted in more mutations (χ_1^2 = 5.92, P = 0.015).
295
Between pathogens with the highest virulence—those under selection from both selective
pressures—populations arisen from MRSA had mutations in 25 genes and intergenic regions,
while those from MSSA had mutations in 12 genes and intergenic regions. Both treatments had
mutations in virulence regulators *codY* [57] and *saeRS* [58] suggesting some shared pathways to
heightened virulence among the two treatments, but mutations in these genes did not occur
across all replicate populations. These treatments also had mutations in *gdpP* and *pbpA*, which
have been implicated in resistance to beta-lactam antibiotics [59,60], with *gdpP* playing a role in
MSSA resistance to oxacillin in particular [61]. Five out of six populations evolved from the MSSA
ancestor had a mutation in either gene, which may have contributed to the reduced oxacillin
sensitivity in these populations.

306
Regulatory genes were enriched with mutations (Figure 4A; one-sample Poisson test P < 0.001
for MRSA and MSSA genotypes). These included *agr*, *saeRS,* and *codY*. *agr* is a quorum sensor
that regulates the expression of virulence factors [62]. All 22 mutations in *agr* were found in
populations that have lost the ability to hemolyze red blood cells. In contrast to the other two
regulators, *codY* is a transcriptional repressor found in gram-positive bacteria that controls
virulence by monitoring nutrient levels in the environment [57].
313
Nine populations experienced large scale deletions that included the entire SCC*mec* cassette.
Deletion of ACME has been shown to enhance the competitive fitness of *S. aureus* USA300 *in*
vivo [63]. The two MRSA populations that lost their resistance (Figure 3C) had SCC*mec* deletions,
suggesting that resistance could be more costly when evolving in a host. The loss of SCC*mec*
and ACME was associated with an increase in total growth from the ancestor outside the host
(Figure 4B; one-sample t-test t = 2.73, df = 8, P = 0.026).

A

323
Figure 4. Regulatory genes likely played an important role in pathogen adaptation. A)
Mutations swept to fixation, excluding intergenic and synonymous mutations, grouped by
general function. The size of each point indicates how many populations had acquired at least
one mutation in the gene. Colored shapes next to genes indicate whether these genes are
regulatory or have been implicated in virulence or antibiotic resistance in the literature (see
Table S1). B) *In vitro* growth of populations with SCC*mec* and ACME deletions with or without
sub-MIC oxacillin. C) Change from ancestral pathogen-induced mortality *vs.* mutations. D)

Hemolysis status *vs*. mutations. E) Oxacillin MIC *vs*. mutations. F) Biofilm production of evolved populations. Different letters indicate significant differences. The column widths in D and E corresponds to the number of mutations. Error bars indicate standard errors. All evolved populations were sequenced except for one population from the -host-ox treatment. *P < 0.05, **P < 0.01

Mutations in antibiotic resistance genes are associated with heightened virulence.
We correlated mutations arising in specific genes with measured traits to identify loci potentially underlying evolved phenotypes. We focused on genes where two or more mutations (excluding intergenic or synonymous mutations) had fixed during the experiment regardless of treatment (Figure 4C). Mutations in three genes were associated with significant increases in virulence from the ancestor: *codY* (one-sample t-test = 19.56, df = 7, P < 0.001), *gdpP* (one-sample t-test = 17.02, df = 3, P = 0.002), and *pbpA* (one-sample t-test = 7.54, df = 3, P = 0.012). Mutations in *agr* in general were not associated with changes in overall virulence, but greater host mortality was associated with *agr* mutations arising from the MSSA ancestor compared to the MRSA ancestor (Wilcoxon rank sum exact test P = 0.045).

Mutations in specific genes had a significant association with the ability to hemolyze red blood cells (Figure 4D; Fisher's exact test P < 0.001). There was a greater proportion of populations with *agr* mutations unable to hemolyze red blood cells compared to *alr* (P = 0.036)*, argR* (P = 0.006)*, codY* (P = 0.005)*, gdpP* (P = 0.006)*, and *purR* (P = 0.006). There were also significant differences between the mutations associated with oxacillin resistance in populations evolved from the MSSA ancestor (Figure 4E; Fisher's exact test P = 0.0015), where a greater proportion of populations with mutations in *codY* had reduced oxacillin susceptibility compared to *brnQ1* (P = 0.043). Exposure to oxacillin maintained hemolysis in evolved populations (Figure 2B), suggesting low concentrations of oxacillin exerted negative selection on the *agr* locus. By contrast, mutations in *agr* were associated with the loss of hemolytic activity, consistent with previous findings [32].

Because mutations in *codY* appeared to be important for both virulence (Figure 4C) and antibiotic resistance (Figure 4E), we hypothesized that traits controlled by *codY* were responsible for the traits observed in MSSA populations under selection from both pressures. A potential mechanism underlying changes in virulence and antibiotic sensitivity in MSSA may involve biofilm formation. Biofilm is implicated in *S. aureus* pathogenesis as well as in dampening the efficacy of antibiotic treatment [41]. In *C. elegans*, *S. aureus* biofilm enhances virulence and protects the pathogen from host innate immune defenses [64]. Both sub-MIC level of beta-lactam antibiotics and null-mutants of *codY* induce robust biofilm formation and hemolysis activity [14,65,66]. Taken together, exposure to sub-MIC level of oxacillin and acquisition of potentially deleterious mutations in *codY* may underlie both increased virulence and reduced antibiotic susceptibility. Our findings partially supported this hypothesis: biofilm formation did not significantly differ for the MRSA genotype (χ_3^2 = 7.24, P = 0.06). By contrast, biofilm production differed between treatments for the MSSA genotype (Figure 4F; χ_3^2 = 17.91, P < 0.001), with populations exposed to both selective pressures forming more biofilm than those evolved without host (Tukey's post-hoc test P < 0.001) or either pressure (P = 0.014), but not those evolved in solely sub-MIC oxacillin (P = 0.095). While host and sub-MIC oxacillin selection favored robust biofilms in MSSA, oxacillin by itself also increased biofilm formation, suggesting low antibiotic concentrations contributed significantly to the evolution of drug-sensitive populations. The loci underlying biofilm formation may be different in these two treatments, since mutations in *codY* did not appear in populations under solely oxacillin selection (biofilm is a polygenic trait in *S. aureus* [67]).

Mutations that arose during experimental evolution are associated with human systemic infections.

We then determined whether mutations that arose in our experiment exist in natural *S. aureus* isolates. We conducted BLAST searches against a dataset of 83,383 high quality public *S. aureus* whole genome assemblies [68] (Figure 5A) and found matches against the majority of our mutations. We hypothesized that these mutations may be associated with *S. aureus* adaptation to different environments. Indeed, for many genes implemented in virulence and antibiotic resistance, a greater proportion of natural isolates containing our mutations were found in blood and systemic infections compared to skin/nose/throat colonization than expected (Figure 5, statistics shown in Table S2). Mutations arisen in MRSA populations under selection from both host and sub-MIC oxacillin had the most matches to natural isolates (Figure 5B), further highlighting the importance of the interaction between these two selective forces. Overall, our results demonstrate that evolution experiments in an invertebrate model can give rise to clinically-relevant mutations in a major bacterial pathogen.

Figure 5. **Mutations that arose during experiment are enriched in isolates associated with systemic infection in humans**. A) Number of genomes in our public *S. aureus* genome dataset (see Methods) grouped by isolation source. General host-association refers to descriptions not specific enough to assign to other categories. We excluded samples that were ambiguously specified or missing information. B) to H) Number of genomes in the database containing mutations in the respective gene. I) Number of genomes in the dataset containing the

404 mutations arisen in *agr*. This gene is separate from the others to facilitate ease of visualization
405 due to the magnitude of the y-axis. Asterisks indicate significant difference in the proportion of
406 blood/systemic-associated genomes compared to the expected distribution in the dataset. *P <
407 0.05, **P < 0.01, ***P < 0.001
408
MRSA and MSSA exhibited divergent phenotypic and genomic evolution during
adaptation
We conducted principal component analysis on the evolved traits (virulence, growth in media with
or without sub-MIC oxacillin, and biofilm production) in MRSA and MSSA populations to determine
the impact of selective pressures on overall pathogen phenotypic evolution (Figures 6A and 6B).
We found a significant effect of treatment for the MRSA genotype (PERMANOVA $F_{(3,23)}$ = 8.38, r^2
= 0.557, P = 0.001), where populations exposed to host and sub-MIC oxacillin formed a distinct
cluster from all other treatments (P < 0.05 for all pairwise comparisons). The MSSA genotype had
a marginally non-significant effect of treatment (PERMANOVA $F_{(3,23)}$ = 2.20, r^2 = 0.248, P = 0.064).
Populations evolved from the MRSA ancestor had similar trajectories for growth in sub-MIC
oxacillin and virulence, whereas biofilm production and virulence were more correlated for
populations evolved from the MSSA ancestor.
421
To determine how host and sub-MIC antibiotic shaped pathogen genomic evolution, we built
phylogenies using mutations that arose from the MRSA and MSSA ancestors (Figures 6C and
D). We compared Euclidean distances calculated from the frequency of each mutation to
determine the genetic distance from the ancestor—how fast each population was evolving—and
the genetic distance between populations within each treatment—how much each population had
diverged from one another [69,70]. There was no significant difference between treatments for
MRSA populations in terms of genetic distance to the ancestor (Figure S6A; $F_{(3,20)}$ = 2.39, P:
0.099), suggesting that populations generally exhibited similar evolutionary rates. By contrast,
MSSA populations had a significant effect of treatment (Figure S6B; $F_{(3,19)}$ = 10.44, P < 0.001),
where populations under selection from solely the host had greater genetic distance from the
ancestor than populations under selection from solely oxacillin (Tukey's post-hoc test P < 0.001)
or under neither selective pressure (P = 0.004).
434
For the MSSA group, populations within each treatment tended to exhibit more genetic similarity
(Figures 6C and 6D). For MRSA, treatments varied in how divergent populations were (Figure
S6C; Kruskal-Wallis χ_3^2= 21.09, P < 0.001), where populations exposed to both selective
pressures diverged more from each other than populations in the other three treatments (Dunn's
post-hoc test P < 0.05 for all pairwise comparisons). Likewise, MSSA populations also differed in
how much they varied (Figure S6D; $F_{(3,51)}$ = 54.20, P < 0.001). Populations under solely host
selection had greater divergence from each other than all other treatments (Tukey's post-hoc test
P < 0.001 for all pairwise comparisons). By contrast, populations under selection solely from sub-
MIC oxacillin diverged the least (P < 0.05 for all pairwise comparisons). These findings altogether
suggest that even concentrations well below the MIC differentially impacts sensitive and resistant
pathogens.
446

448
449 **Figure 6**. **MRSA and MSSA underwent distinct evolutionary trajectories.** Principal
450 component analysis of traits evolved from A) MRSA and B) MSSA ancestors. Phylogenetic tree
451 constructed from frequencies of mutations in populations evolving from C) MRSA and D) MSSA
452 ancestors. Scale bar indicates Euclidean distance.

453
DISCUSSION
While antibiotics have existed for millions of years, the mass production and overuse of antibiotics
by humans in the last century have exerted an unprecedented level of selection on pathogens
[71,72]. In addition to antibiotics, pathogens face numerous selective pressures such as host
defenses and competitors [2,73,74]. Opportunistic pathogens are especially likely to encounter
diverse selective pressures due to their ability to proliferate outside a host. In this study, we
evolved *S. aureus* with or without pre-existing antibiotic resistance under different combinations
of host and sub-MIC antibiotics to explore the evolution of virulence and antibiotic resistance. We
determined whether increased virulence trades-off with antibiotic resistance by directly selecting
for virulence across all host-associated treatments. Nonetheless, we observed differing
evolutionary trajectories, where exposure to oxacillin resulted in pathogens causing the greatest
host mortality. Our experimental design allowed us to tease apart the contributions of the host,
sub-MIC antibiotics, and their interaction to demonstrate that adaptation to novel hosts did not
require trade-offs in pathogen traits. Indeed, selection from both host and antibiotics actually
resulted in rapid pathogen adaptation, potentially creating the conditions under which a pathogen
may emerge in new hosts.

Simultaneous selection from the host and sub-MIC antibiotic favored the greatest virulence and
growth in antibiotics, indicating that interactions between these pressures can shape both traits.
Sub-MIC antibiotics can facilitate adaptation to high drug concentrations and to the growth
environment [15,19]. Frequent exposure to low concentrations of antibiotics in *S. aureus* may
occur due to its ability to grow and survive in non-host environments [37–40], in addition to
infection of antibiotic-treated hosts, where there is uneven distribution of drugs across tissues
[36]. At the cellular level, sub-MIC antibiotics can affect *S. aureus* by directly binding to virulence
factors, modulating virulence regulators, and inducing the expression of toxins like hemolysin
[14,35]. Prior exposure to sub-MIC inside and outside hosts can thus mediate traits involved in
virulence upon host encounter. Our results further provided evidence for potential links between
responses to antibiotic selection and virulence. Mutations genes involved in resistance to
antibiotics were correlated with increased virulence, suggesting that adaptation to antibiotics may
allow for virulence to evolve. Furthermore, in the absence of oxacillin selection, populations lost
their ability to hemolyze red blood cells, supporting previous findings that antibiotics can influence
expression of virulence factors [14,15].

Likewise, host factors contributed to antibiotic resistance. All populations initially sensitive to
oxacillin exhibited increased MICs when under selection from both host and sub-MIC oxacillin.
One population evolving under selection from either solely sub-MIC antibiotics or solely the host
exhibited decreased oxacillin sensitivity, which is consistent with previous studies examining
hosts [26] and sub-MIC [10,19] exposure separately. The pathogens in these studies evolved for
much longer than our study, suggesting that having a host accelerated the evolution of antibiotic
resistance, potentially due to bottlenecks that favor resistant isolates [26]. Evolution in hosts also
contributed to pathogen growth outside hosts, an important trait for opportunistic pathogens as
persistence in the environment increases the likelihood of contact with new hosts. Ultimately,
exposure to hosts and antibiotics together could create a feedback loop where both evolutionary
pressures select for better adaptation in the other, resulting in highly virulent and resistant
pathogens that can persist outside hosts.

Several mutations arose in genes with primary roles in metabolism, such as *codY*. It is becoming
clear that pathogen metabolism plays a central role in its virulence [75,76] and antibiotic
resistance [77]. For example, mutations in chromosomal- and plasmid-encoded metabolic genes
are widely prevalent among *E. coli* strains and protect them against antibiotic challenges [78,79].
Furthermore, antibiotics can directly select for reduced metabolic rates to dampen the deleterious
effects of drugs [79], and increased resistance is more likely to evolve at high nutrient
concentrations [80]. Similarly, metabolic efficiency can alter virulence [75], while resource
limitations can hinder the evolution of virulence [81]. In *S. aureus*, CodY de-represses expression
of virulence factors under resource-depleted conditions in order to acquire nutrients through the
lysing of host cells [57]. Our genomic data were consistent with the observation that *codY* may be
mediating virulence through control of *agr*: MSSA populations under selection from both host and
sub-MIC oxacillin exhibited the greatest virulence and was one of the two treatments without
mutations in *agr*. These results suggested that the *codY* mutations in these populations may be
deleterious, rendering CodY less efficient in repressing *agr*-mediated virulence as a result. As
resources likely differ in abundance and type between hosts and media, our findings illustrated
the importance of taking into consideration the environment in which pathogens evolve when
evaluating traits of interest.

Regulatory genes in general acquired more mutations than expected by chance alone. This
finding supported previous work showing that regulatory elements were more likely to acquire
phenotype-altering mutations, especially in negative regulators (e.g., *codY*) [82]. Enrichment of
mutations in regulatory genes has also been found in *E. coli* under antibiotic selection [83].

Pleiotropic genes like regulators may increase the capacity to adapt to multiple environments.
Because *C. elegans* is a novel host to *S. aureus*, selection is more likely to act on genes that have
large initial gains in the fitness landscape, such as those that have pleiotropic effects [52]. Indeed,
mutations in several metabolic regulators (*codY, purR, gpmA*) and virulence regulators (*saeRS*)
were mainly associated with host infection. The resulting phenotypes of these mutations, such as
increased biofilm production, can then confer multiple benefits like increased virulence and
antibiotic resistance. Our experimental results revealed similar patterns to a comparative study
analyzing over 2000 *S. aureus* genomes, where mutations in genes with pleiotropic effects, such
as *purR*, may have underly pathogen adaptation during systemic infection [31]. Thus, changes in
genes of large effects like global regulators appear to be critical mediators between different traits
that altogether allow bacteria to rapidly adapt to new and stressful environments.

Antibiotic resistance did not impede the evolution of virulence. Antibiotic resistance tends to be
costly in the absence of antibiotics; this cost can subsequently constrain evolution of traits
requiring cellular resources, such as virulence [51]. However, mutations conferring resistance can
have little to no costs, depending on the pathogen species and drugs, and costs can be offset by
compensatory mutations [84]. Sub-MIC levels favor low-cost mutations [18], therefore virulence
and antibiotic resistance may not trade-off during disease emergence [8]. Virulence may
alternatively be costly in the presence of antibiotic resistance—populations evolved from the
MRSA ancestor without either selective pressure exhibited the lowest virulence. The
corresponding MSSA populations did not have a similar decline in virulence, suggesting that
relaxed selection attenuated virulence for populations with existing resistance. Genomic
differences between evolved MRSA and MSSA further demonstrate diverging evolutionary
pathways in antibiotic-resistant *versus* antibiotic-sensitive populations. Oxacillin-exposed MSSA
had no mutations in *agr* and retained their hemolytic ability, while some MRSA populations gained
agr mutations and lost the trait, suggesting that other pathways are utilized to express virulence.
Lastly, mutations in MRSA under selection from both host and antibiotics were significantly
enriched in blood and systemic *S. aureus* isolated from humans, suggesting that antibiotics may
facilitate MRSA invasive infection. Acquiring mutations in genes that impact antibiotic resistance,
such as *pbpA* and *pbpB*, may allow *S. aureus* to persist longer in hosts.

Recent work has shown that diverse genotypes of one pathogen species converge on similar
phenotypes when facing similar selective pressures [85]. Here, we showed that antibiotic
resistance status can lead otherwise isogenic pathogens to use different adaptive strategies.
Initially susceptible pathogens, facing novel selection from host and antibiotics, exhibited reduced
phenotypic and genomic variation over evolutionary time and thus may have had a limited number
of pathways to adapt to new environments. By contrast, resistant populations had greater
variance and thus may have had more routes to utilize. Overall, the potential for low-cost [8,18]
and pleiotropic mutations [52] may have created ideal conditions for a novel pathogen to establish
in host populations regardless of its susceptibility to antibiotics. Our work demonstrated that
responding to multiple pressures can result in rapid adaptation to new hosts and even evolution
of traits not under selection, emphasizing the importance of non-canonical pathways (e.g.,
metabolism) in shaping these traits. Pathogen evolution in a tractable animal model recapitulated
phenotypic and genomic findings from human studies, highlighting the utility of evolution
experiments to identify potential ecological and genetic mechanisms that give rise to pathogen
traits of interest. Our findings ultimately emphasize the importance of considering the host context
in the evolution of antibiotic resistance. Integrating multiple traits, such as virulence, antibiotic
resistance, and fitness may be critical in identifying the factors that facilitate host shifts and
persistence of drug-resistant pathogens.

REFERENCES

1.   Larsson DGJ, Flach CF. Antibiotic resistance in the environment. Nature Reviews
Microbiology. Nature Research; 2022. pp. 257–269. doi:10.1038/s41579-021-00649-x

2.   Hoang KL, Read TD, King KC. Incomplete immunity in a natural animal-microbiota
interaction selects for higher pathogen virulence. Curr Biol. 2024;34: 1–7.

3.   Kubinak JL, Ruff JS, Hyzer CW, Slev PR, Potts WK. Experimental viral evolution to
specific host MHC genotypes reveals fitness and virulence trade-offs in alternative MHC
types. Proc Natl Acad Sci U S A. 2012;109: 3422–3427. doi:10.1073/pnas.1112633109

4.   Herren CM, Baym M. Decreased thermal niche breadth as a trade-off of antibiotic
resistance. ISME J. 2022;16: 1843–1852. doi:10.1038/s41396-022-01235-6

5.   Toprak E, Veres A, Michel JB, Chait R, Hartl DL, Kishony R. Evolutionary paths to
antibiotic resistance under dynamically sustained drug selection. Nat Genet. 2012;44:
101–105. doi:10.1038/ng.1034

6.   Merlo LMF, Sprouffske K, Howard TC, Gardiner KL, Caulin AF, Blum SM, et al.
Application of simultaneous selective pressures slows adaptation. Evol Appl. 2020;13:
1615–1625. doi:10.1111/eva.13062

7.   Sharma M, Quader S, Guttal V, Isvaran K. The enemy of my enemy: multiple interacting
selection pressures lead to unexpected anti-predator responses. Oecologia. 2020;192:
1–12. doi:10.1007/s00442-019-04552-4

8.   Visher E, Evensen C, Guth S, Lai E, Norfolk M, Rozins C, et al. The three Ts of virulence
evolution during zoonotic emergence. Proc R Soc B Biol Sci. 2021;288.
doi:10.1098/rspb.2021.0900

9.   Day T, Kennedy DA, Read AF, Gandon S. Pathogen evolution during vaccination
campaigns. PLoS Biol. 2022;20: e3001804. doi:10.1371/journal.pbio.3001804

10.   Wistrand-Yuen E, Knopp M, Hjort K, Koskiniemi S, Berg OG, Andersson DI. Evolution of
high-level resistance during low-level antibiotic exposure. Nat Commun. 2018;9.
doi:10.1038/s41467-018-04059-1

11.   Oz T, Guvenek A, Yildiz S, Karaboga E, Tamer YT, Mumcuyan N, et al. Strength of
selection pressure is an important parameter contributing to the complexity of antibiotic
resistance evolution. Mol Biol Evol. 2014;31: 2387–2401. doi:10.1093/molbev/msu191

12.   Mahrt N, Tietze A, Künzel S, Franzenburg S, Barbosa C, Jansen G, et al. Bottleneck size
and selection level reproducibly impact evolution of antibiotic resistance. Nat Ecol Evol.
2021;53: 1275. doi:10.1038/s41588-021-00939-3

13.   Cisneros-Mayoral S, Graña-Miraglia L, Perez-Morales D, Peña-Miller R, Fuentes-
Hernández A. Evolutionary History and Strength of Selection Determine the Rate of
Antibiotic Resistance Adaptation. Mol Biol Evol. 2022;39. doi:10.1093/molbev/msac185

14.   Chen J, Zhou H, Huang J, Zhang R, Rao X. Virulence alterations in staphylococcus
aureus upon treatment with the sub-inhibitory concentrations of antibiotics. Journal of
Advanced Research. Elsevier B.V.; 2021. pp. 165–175. doi:10.1016/j.jare.2021.01.008

15.   Andersson DI, Hughes D. Microbiological effects of sublethal levels of antibiotics. Nature
Reviews Microbiology. Nature Publishing Group; 2014. pp. 465–478.
doi:10.1038/nrmicro3270

16.   Windels EM, Van Den Bergh B, Michiels J. Bacteria under antibiotic attack: Different
strategies for evolutionary adaptation. PLoS Pathog. 2020;16: 1–8.
doi:10.1371/journal.ppat.1008431

17.   Gullberg E, Cao S, Berg OG, Ilbäck C, Sandegren L, Hughes D, et al. Selection of
resistant bacteria at very low antibiotic concentrations. PLoS Pathog. 2011;7: 1–9.
doi:10.1371/journal.ppat.1002158

18.   Westhoff S, Van Leeuwe TM, Qachach O, Zhang Z, Van Wezel GP, Rozen DE. The
evolution of no-cost resistance at sub-MIC concentrations of streptomycin in
Streptomyces coelicolor. ISME J. 2017;11: 1168–1178. doi:10.1038/ismej.2016.194

19. Pereira C, Warsi OM, Andersson DI. Pervasive Selection for Clinically Relevant
Resistance and Media Adaptive Mutations at Very Low Antibiotic Concentrations. Mol
Biol Evol. 2023;40: 1–13.
20. Braga PC, Dal Sasso M, Sala MT. Sub-MIC concentrations of cefodizime interfere with
various factors affecting bacterial virulence. J Antimicrob Chemother. 2000;45: 15–25.
doi:10.1093/jac/45.1.15
21. El-Houssaini HH, Elnabawy OM, Nasser HA, Elkhatib WF. Influence of subinhibitory
antifungal concentrations on extracellular hydrolases and biofilm production by Candida
albicans recovered from Egyptian patients. BMC Infect Dis. 2019;19: 1–9.
doi:10.1186/s12879-019-3685-0
22. Khan F, Lee JW, Javaid A, Park SK, Kim YM. Inhibition of biofilm and virulence properties
of Pseudomonas aeruginosa by sub-inhibitory concentrations of aminoglycosides. Microb
Pathog. 2020;146. doi:10.1016/j.micpath.2020.104249
23. Haddadin RNS, Saleh S, Al-Adham ISI, Buultjens TEJ, Collier PJ. The effect of
subminimal inhibitory concentrations of antibiotics on virulence factors expressed by
Staphylococcus aureus biofilms. J Appl Microbiol. 2010;108: 1281–1291.
doi:10.1111/j.1365-2672.2009.04529.x
24. Klemm EJ, Gkrania-Klotsas E, Hadfield J, Forbester JL, Harris SR, Hale C, et al.
Emergence of host-adapted Salmonella Enteritidis through rapid evolution in an
immunocompromised host. Nat Microbiol. 2016;1: 1–6. doi:10.1038/nmicrobiol.2015.23
25. Bacigalupe R, Tormo-Mas MÁ, Penadés JR, Ross Fitzgerald J. A multihost bacterial
pathogen overcomes continuous population bottlenecks to adapt to new host species. Sci
Adv. 2019;5: 1–15. doi:10.1126/sciadv.aax0063
26. McVicker G, Prajsnar TK, Williams A, Wagner NL, Boots M, Renshaw SA, et al. Clonal
Expansion during Staphylococcus aureus Infection Dynamics Reveals the Effect of
Antibiotic Intervention. PLoS Pathog. 2014;10. doi:10.1371/journal.ppat.1003959
27. Alves J, Vrieling M, Ring N, Yebra G, Pickering A, Prajsnar TK, et al. Experimental
evolution of Staphylococcus aureus in macrophages: dissection of a conditional adaptive
trait promoting intracellular survival. Projan SJ, editor. MBio. 2024.
doi:10.1128/mbio.00346-24
28. Fait A, Andersson DI, Ingmer H. Evolutionary history of Staphylococcus aureus
influences antibiotic resistance evolution. Curr Biol. 2023;33: 3389-3397.e5.
doi:10.1016/j.cub.2023.06.082
29. Tsouklidis N, Kumar R, Heindl SE, Soni R, Khan S. Understanding the Fight Against
Resistance: Hospital-Acquired Methicillin-Resistant Staphylococcus Aureus vs.
Community-Acquired Methicillin-Resistant Staphylococcus Aureus. Cureus. 2020.
doi:10.7759/cureus.8867
30. Kourtis AP, Hatfield ; Kelly, Baggs J, Mu ; Yi, See I, Epson E, et al. Vital signs:
epidemiology and recent trends in methicillin-resistant and in methicillin-susceptible
Staphylococcus aureus bloodstream infections-United States. Morb Mortal Wkly Rep.
2019;68: 214–219. Available: https://www.cdc.gov/mmwr
31. Giulieri SG, Guérillot R, Duchene S, Hachani A, Daniel D, Seemann T, et al. Niche-
specific genome degradation and convergent evolution shaping Staphylococcus aureus
adaptation during severe infections. Elife. 2022;11: 1–33. doi:10.7554/eLife.77809
32. Schwan WR, Langhorne MH, Ritchie HD, Stover CK. Loss of hemolysin expression in
Staphylococcus aureus agr mutants correlates with selective survival during mixed
infections in murine abscesses and wounds. FEMS Immunol Med Microbiol. 2003;38: 23–
28. doi:10.1016/S0928-8244(03)00098-1
33. Thomer L, Schneewind O, Missiakas D. Pathogenesis of Staphylococcus aureus
Bloodstream Infections. Annu Rev Pathol Mech Dis. 2016;11: 343–364.
doi:10.1146/annurev-pathol-012615-044351

34. Kwiecinski JM, Horswill AR. Staphylococcus aureus bloodstream infections:
pathogenesis and regulatory mechanisms. Curr Opin Microbiol. 2020;53: 51–60.
doi:10.1016/j.mib.2020.02.005

35. Hodille E, Rose W, Diep BA, Goutelle S, Lina G, Dumitrescu O. The Role of Antibiotics in
Modulating Virulence in Staphylococcus aureus. 2017. doi:10.1128/CMR

36. Nix DE, Goodwin SD, Peloquin CA, Rotella DL, Schentag JJ. Antibiotic tissue penetration
and its relevance: Impact of tissue penetration on infection response. Antimicrob Agents
Chemother. 1991;35: 1953–1959. doi:10.1128/AAC.35.10.1953

37. Roberts MC, Soge OO, No D. Comparison of multi-drug resistant environmental
methicillin-resistant Staphylococcus aureus isolated from recreational beaches and high
touch surfaces in built environments. Front Microbiol. 2013;4: 1–8.
doi:10.3389/fmicb.2013.00074

38. Thapaliya D, Taha M, Dalman MR, Kadariya J, Smith TC. Environmental contamination
with Staphylococcus aureus at a large, Midwestern university campus. Sci Total Environ.
2017;599–600: 1363–1368. doi:10.1016/j.scitotenv.2017.05.080

39. Steadmon M, Ngiraklang K, Nagata M, Masga K, Kiana L. Frank. Effects of water turbidity
on the survival of Staphylococcus aureus in environmental fresh and brackish waters.
Water Environ Res. 2023.

40. Loeffler A, Boag AK, Sung J, Lindsay JA, Guardabassi L, Dalsgaard A, et al. Prevalence
of methicillin-resistant Staphylococcus aureus among staff and pets in a small animal
referral hospital in the UK. J Antimicrob Chemother. 2005;56: 692–697.
doi:10.1093/jac/dki312

41. Long DR, Penewit K, Lo HY, Almazan J, Holmes EA, Bryan AB, et al. In Vitro Selection
Identifies Staphylococcus aureus Genes Influencing Biofilm Formation. Infect Immun.
2023;91. doi:10.1128/iai.00538-22

42. Erler S, Cotter SC, Freitak D, Koch H, Palmer-Young EC, de Roode JC, et al. Insects'
essential role in understanding and broadening animal medication. Trends Parasitol.
2024;40: 338–349. doi:10.1016/j.pt.2024.02.003

43. Higazy D, Pham AD, van Hasselt C, Høiby N, Jelsbak L, Moser C, et al. In vivo evolution
of antimicrobial resistance in a biofilm model of Pseudomonas aeruginosa lung infection .
ISME J. 2024;18: 1–12. doi:10.1093/ismejo/wrae036

44. Diep BA, Gill SR, Chang RF, Phan TH Van, Chen JH, Davidson MG, et al. Complete
genome sequence of USA300, an epidemic clone of community-acquired meticillin-
resistant Staphylococcus aureus. Lancet. 2006;367: 731–739. doi:10.1016/S0140-
6736(06)68231-7

45. Irazoqui JE, Troemel ER, Feinbaum RL, Luhachack LG, Cezairliyan BO, Ausubel FM.
Distinct pathogenesis and host responses during infection of C. elegans by P. aeruginosa
and S. aureus. PLoS Pathog. 2010;6: 1–24. doi:10.1371/journal.ppat.1000982

46. Sifri CD, Begun J, Ausubel FM, Calderwood SB. Caenorhabditis elegans as a model host
for Staphylococcus aureus pathogenesis. Infect Immun. 2003;71: 2208–2217.
doi:10.1128/IAI.71.4.2208-2217.2003

47. Begun J, Sifri CD, Goldman S, Calderwood SB, Ausubel FM. Staphylococcus aureus
virulence factors identified by using a high-throughput Caenorhabditis elegans-killing
model. Infect Immun. 2005;73: 872–877. doi:10.1128/IAI.73.2.872-877.2005

48. Bae T, Banger AK, Wallace A, Glass EM, Åslund F, Schneewind O, et al. Staphylococcus
aureus virulence genes identified by bursa aurealis mutagenesis and nematode killing.
2004. Available: https://www.pnas.org

49. Wu K, Conly J, McClure JA, Elsayed S, Louie T, Zhang K. Caenorhabditis elegans as a
host model for community-associated methicillin-resistant Staphylococcus aureus. Clin
Microbiol Infect. 2010;16: 245–254. doi:10.1111/j.1469-0691.2009.02765.x

50. Wu K, Zhang K, McClure J, Zhang J, Schrenzel J, Francois P, et al. A correlative analysis

of epidemiologic and molecular characteristics of methicillin-resistant Staphylococcus
aureus clones from diverse geographic locations with virulence measured by a
Caenorhabditis elegans host model. Eur J Clin Microbiol Infect Dis. 2013;32: 33–42.
doi:10.1007/s10096-012-1711-x

51.   Ferenci T. Trade-off mechanisms shaping the diversity of bacteria. Trends Microbiol.
2016;24: 209–223. doi:10.1016/j.tim.2015.11.009

52.   Bomblies K, Peichel CL. Genetics of adaptation. Proc Natl Acad Sci U S A. 2022;119.
doi:10.1073/pnas.2122152119

53.   Cheung GYC, Duong AC, Otto M. Direct and synergistic hemolysis caused by
Staphylococcus phenol-soluble modulins: Implications for diagnosis and pathogenesis.
Microbes Infect. 2012;14: 380–386. doi:10.1016/j.micinf.2011.11.013

54.   King JM, Kulhankova K, Stach CS, Vu BG, Salgado-pabón W. Phenotypes and virulence
among Staphylococcus aureus USA100, USA200, USA300, USA400, and USA600 clonal
lineages. mSphere. 2016;1: 1–15. doi:10.1128/mSphere.00071-16.Editor

55.   McAdam PR, Holmes A, Templeton KE, Fitzgerald JR. Adaptive evolution of
Staphylococcus aureus during chronic endobronchial infection of a cystic fibrosis patient.
PLoS One. 2011;6: 1–5. doi:10.1371/journal.pone.0024301

56.   Koch G, Yepes A, Förstner KU, Wermser C, Stengel ST, Modamio J, et al. Evolution of
resistance to a last-resort antibiotic in staphylococcus aureus via bacterial competition.
Cell. 2014;158: 1060–1071. doi:10.1016/j.cell.2014.06.046

57.   Brinsmade SR. CodY, a master integrator of metabolism and virulence in Gram-positive
bacteria. Current Genetics. Springer Verlag; 2017. pp. 417–425. doi:10.1007/s00294-
016-0656-5

58.   Montgomery CP, Boyle-Vavra S, Daum RS. Importance of the global regulators agr and
SaeRS in the pathogenesis of CA-MRSA USA300 infection. PLoS One. 2010;5: 1–9.
doi:10.1371/journal.pone.0015177

59.   Bilyk BL, Panchal V V., Tinajero-Trejo M, Hobbs JK, Foster SJ. An interplay of multiple
positive and negative factors governs methicillin resistance in *Staphylococcus aureus*.
Microbiol Mol Biol Rev. 2022;86. doi:10.1128/mmbr.00159-21

60.   Poon R, Basuino L, Satishkumar N, Chatterjee A, Mukkayyan N, Buggeln E, et al. Loss of
GdpP function in *Staphylococcus aureus* leads to β-lactam tolerance and enhanced
evolution of β-lactam resistance. Antimicrob Agents Chemother. 2022;66.
doi:10.1128/AAC.01431-21

61.   Giulieri SG, Guérillot R, Kwong JC, Monk IR, Hayes AS, Daniel D, et al. Comprehensive
genomic investigation of adaptive mutations driving the low-level oxacillin resistance
phenotype in staphylococcus aureus. MBio. 2020;11: 1–18. doi:10.1128/mBio.02882-20

62.   Novick RP. Autoinduction and signal transduction in the regulation of staphylococcal
virulence. Mol Microbiol. 2003;48: 1429–1449. doi:10.1046/j.1365-2958.2003.03526.x

63.   Diep BA, Stone GG, Basuino L, Graber CJ, Miller A, Des Etages SA, et al. The arginine
catabolic mobile element and staphylococcal chromosomal cassette mec linkage:
Convergence of virulence and resistance in the USA300 clone of methicillin-resistant
Staphylococcus aureus. J Infect Dis. 2008;197: 1523–1530. doi:10.1086/587907

64.   Begun J, Gaiani JM, Rohde H, Mack D, Calderwood SB, Ausubel FM, et al.
Staphylococcal biofilm exopolysaccharide protects against Caenorhabditis elegans
immune defenses. PLoS Pathog. 2007;3: 526–540. doi:10.1371/journal.ppat.0030057

65.   Kuroda H, Kuroda M, Cui L, Hiramatsu K. Subinhibitory concentrations of β-lactam
induce haemolytic activity in Staphylococcus aureus through the SaeRS two-component
system. FEMS Microbiol Lett. 2007;268: 98–105. doi:10.1111/j.1574-6968.2006.00568.x

66.   Majerczyk CD, Sadykov MR, Luong TT, Lee C, Somerville GA, Sonenshein AL.
Staphylococcus aureus CodY negatively regulates virulence gene expression. J
Bacteriol. 2008;190: 2257–2265. doi:10.1128/JB.01545-07

67. Zapotoczna M, O'Neill E, O'Gara JP. Untangling the diverse and redundant mechanisms
of *Staphylococcus aureus* biofilm formation. PLoS Pathog. 2016;12: 1–6.
doi:10.1371/journal.ppat.1005671
68. Raghuram V, Petit RA, Karol Z, Mehta R, Weissman DB, Read TD. Average nucleotide
identity-based Staphylococcus aureus strain grouping allows identification of strain-
specific genes in the pangenome. mSystems. 2024;9. doi:10.1128/msystems.00143-24
69. Ford SA, Williams D, Paterson S, King KC. Co-evolutionary dynamics between a
defensive microbe and a pathogen driven by fluctuating selection. Mol Ecol. 2017;26:
1778–1789. doi:10.1111/mec.13906
70. Betts A, Gray C, Zelek M, Maclean RC, King KC. High parasite diversity accelerates host
adaptation and diversification. Science (80-). 2018;360: 907–911.
71. Spagnolo F, Trujillo M, Dennehy JJ. Why do antibiotics exist? MBio. 2021;12.
doi:10.1128/mBio.01966-21
72. Wright GD. The antibiotic resistome: The nexus of chemical and genetic diversity. Nat
Rev Microbiol. 2007;5: 175–186. doi:10.1038/nrmicro1614
73. Cobey S. Pathogen evolution and the immunological niche. Ann N Y Acad Sci.
2014;1320: 1–15. doi:10.1111/nyas.12493
74. Bashey F. Within-host competitive interactions as a mechanism for the maintenance of
parasite diversity. Philos Trans R Soc B Biol Sci. 2015;370. doi:10.1098/rstb.2014.0301
75. Lindsay RJ, Holder PJ, Talbot NJ, Gudelj I. Metabolic efficiency reshapes the seminal
relationship between pathogen growth rate and virulence. Ecol Lett. 2023;26: 896–907.
doi:10.1111/ele.14218
76. Feinbaum RL, Urbach JM, Liberati NT, Djonovic S, Adonizio A, Carvunis AR, et al.
Genome-wide identification of Pseudomonas aeruginosa virulence-related genes using a
Caenorhabditis elegans infection model. PLoS Pathog. 2012;8: 11.
doi:10.1371/journal.ppat.1002813
77. Zampieri M, Enke T, Chubukov V, Ricci V, Piddock L, Sauer U. Metabolic constraints on
the evolution of antibiotic resistance. Mol Syst Biol. 2017;13: 1–14.
doi:10.15252/msb.20167028
78. Palomino A, Gewurz D, DeVine L, Zajmi U, Moralez J, Abu-Rumman F, et al. Metabolic
genes on conjugative plasmids are highly prevalent in Escherichia coli and can protect
against antibiotic treatment. ISME J. 2023;17: 151–162. doi:10.1038/s41396-022-01329-
1
79. Lopatkin AJ, Bening SC, Manson AL, Stokes JM, Kohanski MA, Badran AH, et al.
Clinically relevant mutations in core metabolic genes confer antibiotic resistance. Science
(80-). 2021;371. doi:10.1126/science.aba0862
80. Windels EM, Cool L, Persy E, Swinnen J, Matthay P, Van Den Bergh B, et al. Antibiotic
dose and nutrient availability differentially drive the evolution of antibiotic resistance and
persistence. ISME J. 2024;18. doi:10.1093/ismejo/wrae070
81. Pak D, Kamiya T, Greischar MA. Proliferation in malaria parasites: How resource
limitation can prevent evolution of greater virulence. Evolution (N Y). 2024;78: 1287–
1301. doi:10.1093/evolut/qpae057
82. Lind PA, Farr AD, Rainey PB. Experimental evolution reveals hidden diversity in
evolutionary pathways. Elife. 2015;2015: 1–17. doi:10.7554/eLife.07074
83. Card KJ, Thomas MD, Graves JL, Barrick JE, Lenski RE. Genomic evolution of antibiotic
resistance is contingent on genetic background following a long-term experiment with
Escherichia coli. Proc Natl Acad Sci U S A. 2021;118: 1–9.
doi:10.1073/pnas.2016886118
84. Melnyk AH, Wong A, Kassen R. The fitness costs of antibiotic resistance mutations. Evol
Appl. 2015;8: 273–283. doi:10.1111/eva.12196
85. Filipow N, Mallon S, Shewaramani S, Kassen R, Wong A. The impact of genetic

background during laboratory evolution of Pseudomonas aeruginosa in a cystic fibrosis-
like environment . Evolution (N Y). 2023;78: 566–578. doi:10.1093/evolut/qpad189

86.   Stiernagle T. Maintenance of *C. elegans*. WormBook. 2006; 2006.
doi:10.1895/wormbook.1.101.1

87.   Vega NM, Gore J. Stochastic assembly produces heterogeneous communities in the
Caenorhabditis elegans intestine. PLoS Biol. 2017;15: 1–20.
doi:10.1371/journal.pbio.2000633

88.   Wahl LM, Gerrish PJ, Saika-Voivod I. Evaluating the impact of population bottlenecks in
experimental evolution. Genetics. 2002;162: 961–971. doi:10.1093/genetics/162.2.961

89.   Cordero OX, Wildschutte H, Kirkup B, Proehl S, Ngo L, Hussain F, et al. Ecological
populations of bacteria act as socially cohesive units of antibiotic production and
resistance. Science (80-). 2012;337: 1228–1231. doi:10.1126/science.1219385

90.   Paterson GK, Harrison EM, Murray GGR, Welch JJ, Warland JH, Holden MTG, et al.
Capturing the cloud of diversity reveals complexity and heterogeneity of MRSA carriage,
infection and transmission. Nat Commun. 2015;6. doi:10.1038/ncomms7560

91.   Mei M, Thomas J, Diggle SP. Heterogenous susceptibility to r-pyocins in populations of
pseudomonas aeruginosa sourced from cystic fibrosis lungs. MBio. 2021;12.
doi:10.1128/mBio.00458-21

92.   Launay A, Wu CJ, Dulanto Chiang A, Youn JH, Khil PP, Dekker JP. In vivo evolution of
an emerging zoonotic bacterial pathogen in an immunocompromised human host. Nat
Commun. 2021;12: 1–12. doi:10.1038/s41467-021-24668-7

93.   Ruiz-Bedoya T, Wang PW, Desveaux D, Guttman DS. Cooperative virulence via the
collective action of secreted pathogen effectors. Nat Microbiol. 2023;8: 640–650.
doi:10.1038/s41564-023-01328-8

94.   Azimi S, Roberts AEL, Peng S, Weitz JS, McNally A, Brown SP, et al. Allelic
polymorphism shapes community function in evolving Pseudomonas aeruginosa
populations. ISME J. 2020;14: 1929–1942. doi:10.1038/s41396-020-0652-0

95.   Penley M, Morran L. Assessment of Caenorhabditis elegans Competitive Fitness in the
Presence of a Bacterial Parasite. Bio-Protocol. 2018;8: 1–14.
doi:10.21769/bioprotoc.2971

96.   Blazanin M. gcplyr: an R package for microbial growth curve data analysis. bioRxiv. 2023;
2023.04.30.538883. Available:
https://www.biorxiv.org/content/10.1101/2023.04.30.538883v1%0Ahttps://www.biorxiv.org
/content/10.1101/2023.04.30.538883v1.abstract

97.   CLSI. Performance standards for antimicrobial susceptibility testing: CLSI approved
standard M100-S23 2013.

98.   Hudzicki J. Kirb-Bauer disk diffusion susceptibility test protocol. Am Soc Microbiol. 2009;
1–23. Available: https://asm.org/Protocols/Kirby-Bauer-Disk-Diffusion-Susceptibility-Test-
Pro

99.   Hoang KL, Read TD, King KC. Defense Heterogeneity in Host Populations Gives Rise to
Pathogen Diversity. Am Nat. 2024;204: 000–000. doi:10.1086/731996

100. Ford SA, Kao D, Williams D, King KC. Microbe-mediated host defence drives the
evolution of reduced pathogen virulence. Nat Commun. 2016;7: 1–9.
doi:10.1038/ncomms13430

101. Petit RA, Read TD. Bactopia: a Flexible Pipeline for Complete Analysis of Bacterial
Genomes. mSystems. 2020;5. doi:10.1128/msystems.00190-20

102. Deatherage DE, Barrick JE. Identification of mutations in laboratory-evolved microbes
from next-generation sequencing data using breseq. Engineering and Analyzing
Multicellular Systems: Methods and Protocols. 2014. pp. 165–188. doi:10.1007/978-1-
4939-0554-6

103. Souvorov A, Agarwala R, Lipman DJ. SKESA: Strategic k-mer extension for scrupulous

assemblies. Genome Biol. 2018;19: 1–13. doi:10.1186/s13059-018-1540-z

104. Lees JA, Kremer PHC, Manso AS, Croucher NJ, Ferwerda B, Serón MV, et al. Large
scale genomic analysis shows no evidence for pathogen adaptation between the blood
and cerebrospinal fluid niches during bacterial meningitis. Microb Genomics. 2017;3.
doi:10.1099/mgen.0.000103

105. Ibarra JA, Pérez-Rueda E, Carroll RK, Shaw LN. Global analysis of transcriptional
regulators in Staphylococcus aureus. BMC Genomics. 2013;14. doi:10.1186/1471-2164-
14-126

106. Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, et al. vegan:
Community Ecology Package. 2024.

107. Martinez Arbizu P. pairwiseAdonis: Pairwise multilevel comparison using adonis. 2020.

MATERIALS AND METHODS
Bacteria and nematode strains
Staphylococcus aureus USA300 JE2 (NR-46543) and USA300 JE2 *mecA*-tn (Transposon Mutant NE1868) from the Nebraska Transposon Mutant Library were acquired from BEI resources (https://www.beiresources.org/). To ensure true isogenic ancestors, the *mecA*-tn was transduced into the USA300 JE2 background, selecting for the erythromycin resistance marker on the Mariner transposon, and the insertion site was confirmed by polymerase chain reaction. Transposon maintenance was confirmed by testing for resistance to erythromycin (5µg/mL) or by genome sequencing when available. In cases where transposon loss was verified by genome sequencing, strains were tested and shown to maintain sensitivity to oxacillin. As the primary function of the transposon was to knock out *mecA* function, loss of the transposon should not confound subsequent analyses as antibiotic sensitivity was maintained.

Caenorhabditis elegans strain N2 Bristol and *Escherichia coli* strain OP50 were provided by the Caenorhabditis Genetics Center, which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440). Nematodes were maintained on Nematode Growth Medium Lite (US Biological, Swampscott, MA) according to WormBook protocols [86].

Experimental evolution
Selection for increased virulence of *S. aureus* (USA300 JE2 or USA300 JE2 *mecA-tn*) in *C. elegans* was performed by passaging *S. aureus* from dead hosts 24 hours post-infection. Half of the *S. aureus* populations were additionally subjected to antibiotic pressure during the passages before and after *C. elegans* selection with sub-MIC oxacillin exposure (0.03125µg/mL). This concentration was tested against both ancestral strains and did not substantially inhibit growth of the methicillin-sensitive USA300 JE2 *mecA*-tn strain.

For each passage of experimental evolution, *S. aureus* was grown in Brain Heart Infusion (BHI) broth with 4µg/mL colistin (to prevent *E. coli* contamination from the nematode intestine) ± 0.03125µg/mL oxacillin and incubated at 37˚C overnight, shaking at 250rpm. The next day, 200µL of the overnight *S. aureus* culture was plated onto BHI agar and incubated at 28˚C overnight. Approximately 1500 *C. elegans* were plated and allowed to consume *S. aureus* for 24 hours before 30 dead nematodes were picked from each plate. Nematodes were identified as dead by a lack of response to tapping by a platinum wire [2,69]. Bacterial extraction methods were adapted from [87]. In brief, picked dead nematodes were washed once with M9 buffer supplemented with 0.1% Triton X-100, once with M9 buffer, then subjected to a 1:1000 bleach-M9 buffer solution for 15 minutes. Nematodes were then washed twice with M9 buffer and ground with a motorized pestle to extract *S. aureus* from the host intestine. *S aureus* populations evolving without hosts were obtained from nematode-free bacterial lawns with a loop. Populations of *S. aureus* were grown as previously before freezing and storage at -80˚C. Twenty-five percent of each frozen culture was used as inoculum for the next round of experimental evolution. Preliminary experiments determined colony-forming units per milliliter dropped by approximately 50% after freezing. To maintain population diversity, the inoculum amount (25%), adjusted for freezing, was chosen to be within the dilution ratio presented in [88] that minimizes the chance that rare beneficial mutations are lost. Twelve rounds of passage were completed for 2 *S. aureus* genotypes x 2 host treatments x 2 antibiotic treatments x 6 replicates = 48 populations. Whole populations were frozen for subsequent analyses. We conducted all assays to evaluate changes in evolved *S. aureus* at the population level instead of single isolates to represent the context under which *S. aureus* evolved in our experiment and generally how pathogens exist in nature and in hosts [55,89–92]. Population level assays also allow for potential interactions between genotypes that give rise to phenotypes like virulence [93] and antibiotic resistance [89,94] that would otherwise not be captured in single-isolate assays.

Mortality assay

Nematodes were age-synchronized as in [95] and grown on *E.coli* OP50 until L4 larval stage (48 hours at 20˚C). Nematodes were subsequently washed off, counted, and approximately 200 individuals were plated on *S. aureus* lawns on BHI plates. Another set of nematodes were plated on *E. coli* OP50 to estimate the total number of nematodes. Plates of *S. aureus* were prepared 24 hours prior by plating 200µL of an overnight population culture on BHI agar and grown at 28˚C. After two days at 20˚C, plates were scored by counting live nematodes. Host mortality data were analyzed using a generalized linear model with a binomial distribution and followed by Tukey multiple-comparison tests to determine pairwise differences. To compare variances across treatments, we conducted Levene's test for homogeneity of variance. All statistical analyses, including those below, were conducted in R (version 4.2.0).

Hemolysis phenotype assay

Populations of *S. aureus* were streaked onto Trypticase Soy Agar II with 5% sheep's blood plates and incubated overnight at 37˚C. Bacteria were assessed as either hemolysis positive or negative depending on whether there was a complete clearing around the colonies. We did not cross-streak with RN4220, which produces beta-hemolysin, and allows for detection of delta-hemolysin. Therefore, our assay mainly detects alpha-toxin and phenol-soluble modulins. Hemolysis data were analyzed using a generalized linear model with a binomial distribution. To determine whether there was an association between hemolysis status and host mortality, we compared the host mortality means of hemolysis-positive vs. hemolysis-negative pathogens using a Kruskal–Wallis test.

To determine whether there was variation in hemolysis ability within populations, we streaked 10 colonies from each of the 48 evolved populations onto Trypticase Soy Agar II with 5% sheep's blood plates and incubated overnight at 37˚C. To compare the number of isolates able to hemolyze, we conducted a generalized linear model with a binomial distribution. To compare variances, we conducted Levene's test for homogeneity of variance.

***In vitro* growth assay**

We measured growth in BHI to simulate conditions under which pathogens were grown in our experiment. Importantly, rich media allowed us to determine growth when conditions are optimal for evolved pathogens. Measuring growth in minimal media, while being more similar to conditions that *S. aureus* may face in its environment, would introduce another selective pressure. Overnight cultures of each evolved population was diluted 1:1000 in BHI broth, then 200µl of each dilution was added to 96-well plates, covered with a Breathe-Easy sealing membrane (Sigma) to minimize evaporation. The optical density at 600nm at 37˚C with shaking at 282 cycles per minute was recorded every 10 minutes for 16 hours (1000 minutes) using a BioTek Synergy HTX Multimode Reader. We randomly assigned each of the 48 evolved populations to the inner wells of the 96-well plates, with at least three technical replicates per population. Each plate also included technical replicates of each ancestor. Each ancestral strain was assayed with four technical replicates per plate across four plates. These steps were repeated for media with 0.03125µg/mL oxacillin. We used the R package *gcplyr* [96] to calculate the area under the curve of density, a metric that quantifies the overall bacterial growth. We compared the growth of populations across treatments using the mean area under the curve of the technical replicates for each population. We used a linear model followed by Tukey multiple-comparison tests to determine pairwise differences between the ancestors, and linear mixed models for evolved populations grown with or without oxacillin.

Antimicrobial susceptibility assay

Antimicrobial susceptibility testing for oxacillin was done according to Clinical & Laboratory Standards Institute (CLSI) standard protocols for broth microdilution [97]. In brief, populations were streaked from frozen stock and re-streaked the next day. From the second day culture, populations were resuspended in normal saline to a 0.5 McFarland standard. Then, these cultures were diluted 1:20 in normal saline before 10μL was added to 90μL of CAMHB + 2% NaCl with the appropriate concentration of oxacillin. Plates were incubated at 35°C without shaking and read at 24 hours. For statistical analysis of populations evolved from the MSSA ancestor, we set a threshold such that anything above the ancestral susceptibility level (0.25ug/ml) was considered a decrease in oxacillin sensitivity. We then compared the proportion of populations exhibiting a decrease in oxacillin sensitivity with those that had not using Fisher's exact test with false discovery rate-adjusted (FDR) p-values.

To determine whether there was variation in antibiotic susceptibility within evolved populations, we sampled 10 colonies from each of the four populations with the most number of mutations (two from MRSA and two from MSSA), and two additional randomly selected populations. We argued that variation would be the most observable in populations with the most genetic diversity. We then followed the standard protocol for a Kirby-Bauer disk diffusion susceptibility test [98] using oxacillin. Briefly, we inoculated Mueller-Hinton agar plates (Hardy Diagnostics) of overnight cultures with standardized OD_{600} and placed an oxacillin-impregnated disk (HardyDisk AST Oxacillin, OX1) onto each plate. We measured the diameter of the zone of inhibition after incubation at 37°C overnight.

Biofilm assay

Overnight cultures of each population were diluted 1:40 in BHI broth containing 0.5% glucose, then 100μl of each dilution was added to the inner wells of 96-well plates (NEST Biotechnology, flat-bottom, tissue culture-treated). Plates were incubated for 24 hours at 20°C, the temperature that nematodes experience during infection. Each well was washed three times with 200μl of phosphate-buffered saline (PBS) to remove non-adherent cells. We then stained cells with 125μl of 0.1% crystal violet for 20 minutes, followed by washing each well twice with 200μl PBS. We then resuspended the crystal violet in 125μl of 30% acetic acid. After incubation for 10 minutes, we measured the optical density at 595nm using a BioTek Synergy HTX Multimode Reader. We randomly assigned each of the 48 evolved populations to the inner wells of the 96-well plates, with at least three technical replicates per population. Each plate also included technical replicates of each ancestor. We standardized the mean of technical replicates for each population against the respective ancestor within each plate to control for variation across plates. We used mixed linear models followed by Tukey multiple-comparison tests to assess pairwise differences between treatments.

Whole-genome shotgun sequencing and variant annotation

Bacterial genome DNA was extracted from whole populations using Wizard Genomic DNA Purification Kit (Promega). Population sequencing allows us to capture the genetic variation within populations, which is critical to understanding how pathogens evolve [2,12,69,94,99,100]. Genome sequencing of these DNA samples was performed at the M-Gen center using a paired-end library preparation method based on Illumina Nextera and sequenced on the NextSeq 550.

The raw data was quality trimmed and adapters removed using illumina-cleanup (https://github.com/rpetit3/illumina-cleanup) or with bactopia [101]. Mutations were called based on comparison to the reference USA300 JE2 genome (BioProject: PRJNA224116, BioSample: SAMN06677988, Assembly: GCF_002085525.1) using breseq [102]. The presence of large-scale deletions and other genetic changes were confirmed by analysis of de novo assembled genomes

performed by the SKESA assembler software [103]. The ancestral populations were almost
identical to the reference genome used except for three mutations that were not found in any
evolved populations. Another three mutations in *bioD1, cls* and *ccdC* were present in all
populations derived from the MSSA ancestor and likely occurred in the first passage from the
frozen culture before experimental evolution was performed. We therefore removed these six
mutations from the analysis.
1047
The number of mutations (excluding synonymous or intergenic regions) across treatments were
analyzed using a generalized linear model with a Poisson distribution. For mutation enrichment
analysis, we followed the steps as described previously [104]. Briefly, we calculated the "mutation
rate" for each ancestor by dividing the total number of mutations by the reference genome size.
We multiplied the mutation rate by the total length of regulatory genes [105] with mutations in our
study to generate the expected number of mutations for these genes. We then conducted single-
tailed Poisson tests to determine whether the actual number of mutations in regulatory genes was
greater than expected.
1056
1057 For phenotypes *vs.* mutations, we considered only mutations occurring at least twice throughout
1058 the experiment. For host mortality *vs.* mutations and growth rate *vs.* SCC*mec* and ACME, we
1059 conducted a one sample t-test for each evolved pathogen against its respective ancestor. P-
1060 values were adjusted using the FDR. For hemolysis status *vs.* mutations and oxacillin MIC *vs.*
1061 mutations, we conducted Fisher's exact tests with Bonferroni corrections.
1062
Comparing mutations in experiment against publicly available genomes
For each non-synonymous mutation in each gene, we created the FASTA file of the USA300 JE2
protein product and a cognate file of the mutant protein with the amino acid substitution edited
manually. We ran tBLASTn (version 2.12.0) searches of both proteins against 83,383 high quality
S. aureus whole genome assemblies [68]. We filtered out matches to genomes with less than
95% identity. We screened for cases where mutants had a better match to a genome than the
ancestral protein (higher bitscore).
1070
We collected meta-data for BioSamples accession linked to each genome from NCBI Entrez using
a Python script from Jason Stajich (https://github.com/stajichlab/biosample_metadata), then
manually curated the "isolation_source" column to either blood/systemic infection,
skin/nose/throat, general host-association, animal, environment, others, and missing (see Table
S3 for specific terms). For BioSamples matching mutants within each treatment, we compared
the proportion of blood/systemic infection *vs.* skin/nose/throat against the expected proportion (all
BioSamples in the dataset) using a chi-square goodness-of-fit test.
1078
1079 **Principal component analysis**
1080 We conducted a principal component analysis on the phenotypes quantified in evolved
1081 populations using the prcomp function in R for each *S. aureus* genotype. We performed
1082 permutational analysis of variances (PERMANOVA, 999 permutations) to test for differences
1083 between treatments using adonis2 of the *vegan* package [106] and pairwiseAdonis [107].
1084
Genetic distance
We generated a matrix of Euclidian genetic distances (the square root of pairwise differences)
using the frequencies of mutations that arose for each *S. aureus* genotype, from which we
constructed phylogenies [69,70]. We used pairwise differences between the ancestor and each
evolved populations to compare rates of evolution across treatments, and pairwise differences
between replicate populations within each treatment to compare the degree of divergence across
treatments. We compared rates of evolution for MRSA and MSSA treatments, and degree of

divergence for MSSA treatments, using linear models followed by Tukey multiple-comparison
tests. We compared the degree of divergence for MRSA treatments using a Kruskal–Wallis test
followed by Dunn's post-hoc test.