STAMBPL1 activates the GRHL3/HIF1A/VEGFA axis through interaction with FOXO1 to

promote angiogenesis in triple-negative breast cancer

Huan Fang[1,2#], Huichun Liang[1#], Chuanyu Yang[1], Dewei Jiang[1], Qianmei Luo[1], Wenming

Cao[3]*, Huifeng Zhang[4]*, Ceshi Chen[5,6,1]*

1. Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, Yunnan, China.

2. Kunming College of Life Sciences, University of Chinese Academy of Sciences, Kunming,

Yunnan, China.

3. The Department of Breast Medical Oncology, Zhejiang Cancer Hospital, Hangzhou

310022, China.

4. Department of Pharmacy, the First People's Hospital of Yunnan Province/the Affiliated

Hospital of Kunming University of Science and Technology, Kunming, Yunnan Province, China

5. Academy of Biomedical Engineering, Kunming Medical University, Kunming 650500, China.

6. The Third Affiliated Hospital, Kunming Medical University, Kunming 650118, China.

# These authors contributed equally to this work.

* Corresponding authors: Wenming Cao, E-mail: caowm@zjcc.org.cn; Huifeng Zhang, E-mail:

zhanghuifeng92@163.com; or Ceshi Chen, E-mail: chenc@kmmu.edu.cn.

Abstract

In the clinic, anti-tumor angiogenesis is commonly employed for treating recurrent, metastatic,

drug-resistant triple-negative and advanced breast cancer. Our previous research revealed

that the deubiquitinase STAMBPL1 enhances the stability of MKP-1, thereby promoting

cisplatin resistance in breast cancer. In this study, we discovered that STAMBPL1 could

upregulate the expression of the hypoxia-inducible factor HIF1α in breast cancer cells.

Therefore, we investigated whether STAMBPL1 promotes tumor angiogenesis. We

demonstrated that STAMBPL1 increased *HIF1A* transcription in a non-enzymatic manner,

thereby activating the HIF1α/VEGFA signaling pathway to facilitate TNBC angiogenesis.

Through RNA-seq analysis, we identified the transcription factor GRHL3 as a downstream

target of STAMBPL1 that is responsible for mediating *HIF1A* transcription. Furthermore, we

discovered that STAMBPL1 regulates *GRHL3* transcription by interacting with the

transcription factor FOXO1. These findings shed light on the role and mechanism of

STAMBPL1 in the pathogenesis of breast cancer, offering novel targets and avenues for the

treatment of triple-negative and advanced breast cancer.

Keywords Triple-negative breast cancer, tumor angiogenesis, transcription regulation

Introduction

Triple-negative breast cancer (TNBC) is a subtype of breast cancer characterized by a lack of expression of estrogen receptor (ER), progesterone receptor (PR), and human epidermal growth factor receptor 2 (HER2). Despite representing only 10–15% of all breast cancer cases, it is associated with a greater risk of recurrence, metastasis, and resistance to chemotherapy, leading to a poorer prognosis than other types of breast cancer[1]. Therefore, understanding the pathogenesis of this subtype of breast cancer and identifying effective treatment targets are key areas of research focus and challenge. In the clinic, in addition to surgery, radiotherapy, and chemotherapy, treatment approaches for TNBC also include the use of epidermal growth factor receptor (EGFR) inhibitors, poly (ADP-ribose) polymerase (PARP) inhibitors, immune checkpoint inhibitors, and anti-angiogenic therapies[2,3].

For solid tumors, the survival, proliferation, and invasion of cancer cells rely on surrounding blood vessels to provide nutrients and oxygen. Cancer cells can increase the synthesis and secretion of the vascular endothelial growth factor A (VEGFA), activate endothelial cells in neighboring blood vessels through paracrine pathways, and stimulate tumor angiogenesis[4]. Therefore, investigating the upstream molecular mechanisms that activate VEGFA in cancer cells offers new targets for inhibiting TNBC angiogenesis via disruption of this signaling pathway.

The deubiquitinase STAMBPL1 (also known as AMSH-LP) belongs to the AMSH family and cleaves K63-linked polyubiquitin chains[5]. STAMBPL1 stabilizes XIAP to inhibit apoptosis in prostate cancer cells[6] and confers resistance to honokiol-induced apoptosis in various cancer cell types by stabilizing Survivin and c-FLIP[7]. Our previous research demonstrated that

STAMBPL1 enhances cisplatin resistance in TNBC cells through the stabilization of MKP-1[8].

In this study, we discovered that, independent of its deubiquitinating enzyme activity,

STAMBPL1 upregulates HIF1α expression in TNBC cells. We aimed to investigate the

molecular mechanism by which it upregulates HIF1α and explore its role in tumor

angiogenesis in TNBC.

FOXO1, a member of the forkhead box transcription factor family, regulates various

physiological and pathological processes, such as cell proliferation, apoptosis, autophagy,

and oxidative stress[9]. In breast cancer, FOXO1 enhances the stemness of cancer cells by

promoting *SOX2* transcription[10] and confers resistance to chemotherapy drugs in basal-like

breast cancer cells by activating *KLF5* transcription[11]. Furthermore, FOXO1 plays a crucial

role in angiogenesis by upregulating VEGFA transcription[12]. However, the specific mechanism

by which FOXO1 regulates VEGFA transcription remains unclear, and its role in tumor

angiogenesis in TNBC requires further investigation.

In this study, we discovered that STAMBPL1 facilitates the transcriptional regulation of

GRHL3 by interacting with FOXO1, consequently enhancing *HIF1A* transcription through

GRHL3 to activate the HIF1α/VEGFA pathway. This results in increased endothelial cell

activity via paracrine signaling, thereby promoting tumor angiogenesis in TNBC.

Methods

Cell lines and reagents

All cell lines used in this study, including HCC1806, HCC1937, and HEK293T cells, were purchased from ATCC (American Type Culture Collection, Manassas, VA, USA) and validated by short tandem repeat (STR) analysis, and these cell lines tested negative for mycoplasma contamination. HCC1806 and HCC1937 cells were cultured in RPMI 1640 medium supplemented with 5% FBS. HEK293T cells were cultured in DMEM (Thermo Fisher, Grand Island, USA) with 5% FBS at 37°C with 5% CO_2. Primary human umbilical vein endothelial cells (HUVECs) were maintained in an EGM-2 Bullet Kit (CC-3162, Lonza, USA). AS1842856 (Cat#HY-100596) and apatinib (Cat#HY-13342S) were purchased from MCE (New Jersey, USA).

Plasmid construction and stable STAMBPL1, GRHL3 and FOXO1 overexpression

We constructed the full-length *STAMBPL1/GRHL3/FOXO1* genes and then subcloned them into the pCDH lentiviral vector. The packaging plasmids (pMDLg/pRRE, pRSV-Rev, and pCMV-VSV-G) and the pCDH-STAMBPL1/GRHL3/FOXO1 expression plasmid were cotransfected into HEK293T cells (2 ×10^6 in 10 cm plates) to produce lentiviruses. Following transfection for 48 hours, the lentivirus was collected and used to infect HCC1806 and HCC1937 cells. Forty-eight hours later, puromycin (2 µg/ml) was used to screen the cell populations.

Stable knockdown of STAMBPL1 and GRHL3

The pSIH1-H1-puro shRNA vector was used to express STAMBPL1, GRHL3 and luciferase (LUC) shRNAs. *STAMBPL1* shRNA#1, 5'-GGAGCATCAGAGATTGATA-3'; *STAMBPL1*

shRNA#2, 5'-GCTGCTATGCCTGACCATA-3'; *GRHL3* shRNA#1, 5'-

CCTTGAGCTTCCTCTATGA-3'; *GRHL3* shRNA#2, 5'-AGAGGAAGATGCGCGATGA-3';

Luciferase shRNA, 5'-CUUACGCUGAGUACUUCGA-3'; HCC1806 and HCC1937 cells were

infected with lentivirus. Stable populations were selected via the use of 1 to 2 mg/mL

puromycin. The knockdown effect was evaluated by Western blotting.

RNA interference

The siRNA target sequences used in this study were as follows: *STAMBPL1* siRNA#1, 5'-

GGAGCATCAGAGATTGATA-3'; *STAMBPL1* siRNA#2, 5'-GCTGCTATGCCTGACCATA-3';

GRHL3 siRNA#1, 5'-CCTTGAGCTTCCTCTATGA-3'; *GRHL3* siRNA#2, 5'-

AGAGGAAGATGCGCGATGA-3'; *HIF1α* siRNA#1, 5'-AAGAGGTGGATATGTCTGG-3'; *HIF1α*

siRNA#2, 5'-CGTCGAAAAGAAAAGTCTCTT-3'; *FOXO1* siRNA#1, 5'-

CCCAGAUGCCUAUACAAAC-3'; and *FOXO1* siRNA#2, 5'-CTCAAATGCTAGTACTATTAG-3'.

All siRNAs were synthesized by RiboBio (RiboBio, China) and transfected at a final

concentration of 50 nM.

Western blotting (WB) and antibodies

The WB procedure was described in our previous study[13]. Anti-STAMBPL1 (sc-376526) and

anti-GAPDH (sc-25778) antibodies were purchased from Santa Cruz Biotechnology (Santa

Cruz, CA, USA). Anti-FOXO1 (#2880S) and anti-HIF1α (#36169S) antibodies were purchased

from CST. Anti-β-actin (A5441) and anti-GST (G7781) antibodies were purchased from

Sigma–Aldrich (St. Louis, MO, USA). The anti-Flag (ab205606) antibody was purchased from

Abcam.

Real-time polymerase chain reaction (RT–qPCR)

Total RNA was extracted via TRIzol (Invitrogen, 15596026). One microgram of total RNA was

reverse transcribed to cDNA according to the manufacturer's instructions for the HiScript II

QRT SuperMix for qPCR Kit (Vazyme, R223-01). For quantitative PCR (RT–qPCR), the

SYBR Green Select Master Mix system (Applied Biosystems, 4472908, USA) was used on an

ABI-7900HT system (Applied Biosystems, 4351405). The primers used for PCR were as

follows: 18S forward, 5′-CTCAACACGGGAAACCTCAC-3′; 18S reverse, 5′-

CGCTCCACCAACTAAGAACG-3′; HIF1α forward, 5′-AAGTCTGCAACATGGAAGGTAT-3′;

HIF1α reverse, 5′-TGAGGAATGGGTTCACAAATC-3′; VEGFA forward, 5′-

TGAGGAATGGGTTCACAAATC-3′; VEGFA reverse, 5′-ATCTGCATGGTGATGTTGGA-3′;

GLUT1 forward, 5′-TCGTCGGCATCCTCATCGCC-3′; GLUT1 reverse, 5′-

CCGGTTCTCCTCGTTGCGGT-3′; GRHL3 forward, 5′-GGGGCTGAGGAATGCGATCT-3′;

and GRHL3 reverse, 5′-AATTTTGCCGTCCAGCTCCC-3′.

Cell proliferation and migration assays

To detect the proliferation of HUVECs, we used the Click-iT™ EdU Alexa Fluor™ 647 Imaging

Kit (Invitrogen) according to the manufacturer's protocol. Briefly, HUVECs were seeded on

coverslips (BD Biosciences) at 0.5×10^5 cells per well. The next day, the supernatants were

discarded, and the cells were cultured with conditioned medium (CM). Six hours later, the

cells were incubated with EdU in CM for 4 hours, followed by fixation and staining. For each

sample, three random fields were observed via fluorescence microscopy, and the total

numbers of cells and EdU-positive cells were counted. To detect the migration of HUVECs,

we performed a wound-healing assay. Twenty-four hours after seeding, the supernatants of

the HUVECs were discarded, and the cells were scratched and cultured with CM for 24 hours.

Wound closure was imaged via microscopy. For each image, the gap width was analyzed via

ImageJ.

Tube formation assays

HUVECs (1 × 10^4) in CM were seeded onto Matrigel (BD Biosciences)-coated μ-Slide

angiogenesis plates (ibidiGmbH, Munich, Germany). At 6 hours after seeding, images were

taken via microscopy and then analyzed with ImageJ. The total branch length was measured.

Chromatin immunoprecipitation assays

After the sample preparation was completed, the subsequent experimental steps were

performed according to the instructions of the Simple ChIP (R) Plus Kit (CST, # 9005). The

PCR primers for amplifying the region of interest on the HIF1α gene promoter were as

follows: 5′-GACTGACAGGCTTGAAGTTTATGC-3′ and 5′-

TGTTGCTGTAAACTTCAAGGGAAA-3′, and the PCR primers for amplifying the region of

interest on the GRHL3 gene promoter were as follows: 5′-TTCTATCCCTTCTGTGCTGACCA-

3′ and 5′-TGTGCCAGACCCTACTCTGGG-3′.

Dual-luciferase reporter assays

The DNA fragments HIF1α and GRHL3 were amplified from MCF10A cell genomic DNA via

PCR template and then cloned and inserted into the pGL3-Basic vector. HEK293T cells were

seeded in 24-well plates and transfected with pCDH-GRHL3-3×Flag or pCDH-FOXO1-3×Flag

and pGL3 luciferase reporter plasmids (both 600 ng/well) together with the pCMV-Renilla

control (5 ng/well). After transfection for 48 h, the cell lysates were collected, and the

luciferase activities were detected via the dual-luciferase reporter assay system (Promega,

USA). For the WT HIF1α promoter (with the GRHL3 binding motif), the primers used for PCR

were as follows: forward, 5′-gctagcccgggctcgagatctCCACTGCGCTCCAGCCTG-3′; reverse,

5′-cagtaccggaatgccaagcttCCTCAGACGAGGCAGCACTG-3′. For the mutant HIF1α promoter

(without the GRHL3 binding motif), the primers used for PCR were as follows: forward, 5′-

TCTTTCCCTGAGGCCTTCCTATATGCTTAT-3′; reverse, 5′-

ATAAGCATATAGGAAGGCCTCAGGGAAAGA-3′. For the WT GRHL3 promoter (with the

FOXO1 binding motif), the primers used for PCR were as follows: forward, 5′-

gctagcccgggctcgagatctATTAACAAGGGTGACTGAAGAGGG-3′; reverse, 5′-

cagtaccggaatgccaagcttTGGAGGTATACCTCAACAGGTGC-3′. For the mutant GRHL3

promoter (without the FOXO1 binding motif), the primers used for PCR were as follows:

forward, 5′-CTCCCCCACCAAACAAAGAAGGAGAACACCCC-3′; reverse, 5′-

GGGGTGTTCTCCTTCTTTGTTTGGTGGGGGAG-3′.

Immunofluorescence staining

HEK293T cells plated on cell culture slides were transfected with the STAMBPL1 and FOXO1

expression plasmids. Two days after transfection, the cells were fixed in 3.7%

polyformaldehyde at 4°C overnight. After being blocked with 5% BSA at room temperature for

1 hour, the cells were stained with anti-STAMBPL1 (mouse) and anti-FOXO1 (rabbit)

antibodies at 4°C overnight. The cells were subsequently stained with both an Alexa

Fluor647-labeled anti-mouse secondary antibody and a FITC-labeled anti-rabbit secondary

antibody (ZSGB-Bio, Beijing, China) at room temperature for 1 hour. Nuclei were stained with

DAPI (Biosharp, BL739A) for 15 min. Images were captured via a confocal microscope.

Immunoprecipitation and GST pull-down

For endogenous protein interaction, cell lysates were first incubated with anti-FOXO1

antibodies or rabbit IgG (2729S; Cell Signaling Technology) and then incubated with protein

A/G magnetic beads (HY-K0202; MCE). For the GST pull-down assay, the cell lysates were

directly incubated with Glutathione Sepharose 4B (10312185; Cytiva) overnight at 4°C. For

the IP-Flag assay, cell lysates were directly incubated with anti-Flag magnetic beads (HY-

K0207; MCE) overnight at 4°C. The precipitates were washed four times with 1 ml of lysis

buffer, boiled for 10 minutes with 1×SDS sample buffer, and subjected to WB analysis.

Xenograft experiments

We purchased 5- to 6-week-old female BALB/c nude mice from SLACCAS (Changsha,

China). Animal feeding and experiments were approved by the animal ethics committee of

Kunming Institute of Zoology, Chinese Academy of Sciences. HCC1806-shLuc, HCC1806-

shSTAMBPL1, or HCC1806-shGRHL3 cells and HCC1806-PCDH-Vector or HCC1806-

PCDH-STAMBPL1 cells (1×10^6 in Matrigel (BD Biosciences, NY, USA)) were implanted into

the mammary fat pads of the mice. When the tumor volume reached approximately 50 mm^3,

the nude mice were randomly assigned to the control or treatment groups (n = 4/group). The

control group was given vehicle alone, and the treatment group received the FOXO1 inhibitor

AS1842856 (10 mg/kg), the VEGFR inhibitor apatinib (50 mg/kg), and the FOXO1 inhibitor

AS1842856 combined with the VEGFR inhibitor apatinib via intragastric administration every

two days for 20 days. The tumor volume was calculated as follows: tumor volume was

calculated by the formula (π × length × width^2)/6. The maximal tumor size permitted by the

animal ethics committee of Kunming Institute of Zoology, Chinese Academy of Sciences, and

the maximal tumor size in this study was not exceeded.

Immunohistochemical staining

The xenograft tumor tissues were fixed in 3.7% formalin solution. The immunohistochemistry

was performed on 4-µm-thick paraffin sections after pressure-cooking for antigen retrieval. An

anti-CD31 primary antibody (1:400, Abcam, ab28364) was used. After 12 h, the slides were

washed three times with PBS and incubated with secondary antibodies (hypersensitive

enzyme-labeled goat anti-mouse/rabbit IgG polymer (OriGene, China) at room temperature

for 20 min, DAB concentrated chromogenic solution (1:200 dilution of concentrated DAB

chromogenic solution)), counterstained with 0.5% hematoxylin, dehydrated with graded

concentrations of ethanol for 3 min each (70%–80%–90%–100%), and finally stained with

dimethyl benzene. The immunostained slides were evaluated via light microscopy, and the

number of microvessels with positive CD31 expression was counted.

Statistical analysis

All the graphs were created via GraphPad Prism software version 8.0. Comparisons between

two independent groups were assessed via two-tailed Student's t tests. One-way analysis of

variance with least significant differences was used for multiple group comparisons. P values

of < 0.05, 0.01 or 0.001 were considered to indicate statistically significant results, and

comparisons that were significant at the 0.05 level are indicated by *, those at the 0.01 level

are indicated by **, and those at the 0.001 level are indicated by ***.

Results

STAMBPL1 upregulates HIF1α in a non-enzymatic manner in TNBC cells.

Our previous research revealed that STAMBPL1, a deubiquitinase, promotes cisplatin

resistance in TNBC by stabilizing MKP1[8]. To further investigate the role and mechanism of

STAMBPL1 in TNBC, we discovered that the knockdown of STAMBPL1 can inhibit hypoxia-

induced HIF1α expression (Fig. 1A) and suppress the transcription of the HIF1α downstream

genes *VEGFA* and *GLUT1* (Fig. 1B-E). Under normoxic conditions, both STAMBPL1 and its

enzymatically mutated variant (E292A) can upregulate the protein expression of HIF1α (Fig.

1F). These findings suggest that STAMBPL1 enhances the expression of HIF1α in breast

cancer cells through a non-DUB enzyme activity mechanism.

STAMBPL1 promotes *HIF1A* transcription and activates the HIF1α/VEGFA axis.

To elucidate how STAMBPL1 upregulates HIF1α, we examined the transcription levels of

HIF1A. These results indicate that knocking down STAMBPL1 significantly inhibits the

transcription of *HIF1A* (Fig. 2A-C). STAMBPL1 overexpression increased the transcription of

HIF1A (Fig. 4E and SFig. 3E) and *VEGFA* (Fig. 2E and 2G). The ability of STAMBPL1 to

induce *VEGFA* transcription was blocked by HIF1α knockdown (Fig. 2D-G). These findings

suggested that STAMBPL1 activated the HIF1α/VEGFA axis through enhancing the

transcription of *HIF1α*.

STAMBPL1 in TNBC cells enhances the activity of HUVECs and promotes TNBC

angiogenesis.

The conditioned medium (CM) from TNBC cells with STAMBPL1 knockdown inhibited the

proliferation (Fig. 3A-B, and SFig. 1A-B), migration (Fig. 3C-D, and SFig. 1C-D), and tube

formation (Fig. 3E-F, and SFig. 1E-F) of HUVECs. When STAMBPL1 was overexpressed in

TNBC cells, the conditioned medium of HCC1806 and HCC1937 cells promoted the ability of

HUVECs to proliferate (SFig. 2A and 2D), migrate (SFig. 2B and 2E), and form tubes (SFig.

2C and 2F), which could be reversed by knocking down HIF1α in TNBC cells. These findings

suggest that STAMBPL1 activates the HIF1α/VEGFA axis in TNBC cells, leading to enhanced

abilities of HUVECs through a paracrine pathway. Knocking down STAMBPL1 inhibited the

growth of HCC1806 xenografts in mice (Fig. 3G-I) and decreased the number of microvessels

in tumor tissue (Fig. 3J-K), indicating that STAMBPL1 promoted tumor angiogenesis.

STAMBPL1 promotes *HIF1A* transcription via the upregulation of GRHL3.

By silencing the *STAMBPL1* gene in HCC1806 cells subjected to 10 hours of hypoxia, we

performed RNA-seq analysis to investigate the mechanism by which STAMBPL1 promotes

HIF1A transcription. We found that silencing of STAMBPL1 resulted in the downregulation of

27 genes (of which only 18 were annotated), with *GRHL3* being the sole gene encoding a

transcription factor (Fig. 4A and SFig. 3A). Silencing of STAMBPL1 inhibited *GRHL3*

transcription in both HCC1806 and HCC1937 cells (Fig. 4B and SFig. 3B). Conversely, the

overexpression of STAMBPL1 promoted *GRHL3* transcription (Fig. 4D and SFig. 3D).

Furthermore, the stimulatory effects of STAMBPL1 on the HIF1α/VEGFA axis (Fig. 4C-F and

SFig. 3C-F) and on HUVECs (Fig. 4G-I and SFig. 3G-I) were reversed by GRHL3 knockdown.

These findings suggest that STAMBPL1 activates the HIF1α/VEGFA axis by upregulating

GRHL3.

GRHL3 enhances *HIF1A* transcription by binding to its promoter.

On the basis of the GRHL binding motif (Fig. 5A), a GRHL binding sequence was identified in

the *HIF1A* promoter (Fig. 5B). ChIP and luciferase assays revealed that GRHL3 bound to the

HIF1A promoter (Fig. 5D) and increased its activity (Fig. 5E). Knockdown of GRHL3 (Fig. 5G

and SFig. 4B) resulted in decreased expression of HIF1α at both the mRNA (Fig. 5H and

SFig. 4C) and protein levels (Fig. 5F and SFig. 4A), leading to suppressed *VEGFA*

transcription (Fig. 5I and SFig. 4D), indicating that GRHL3 promotes *HIF1A* transcription by

binding to its promoter.

Furthermore, conditioned medium from TNBC cells with GRHL3 knockdown inhibited HUVEC

proliferation, migration, and tube formation (Fig. 5J-L and SFig. 4E-G). The overexpression of

GRHL3 in TNBC cells activated the HIF1α/VEGFA axis (Fig. 6A-B and SFig. 4H-J), resulting

in increased proliferation, migration, and tube formation in HUVECs. This effect was reversed

by knocking down HIF1α in TNBC cells (Fig. 6C-E and SFig. 4K-M). Knockdown of GRHL3

(Fig. 6F) also inhibited tumor growth in HCC1806 xenograft mice (Fig. 6G-I) and reduced the

number of microvessels in tumor tissues (Fig. 6J-K).

STAMBPL1 mediates *GRHL3* transcription by interacting with FOXO1.

Our previous study demonstrated that STAMBPL1 is localized in the nucleus[8]. This protein

may promote the transcription of GRHL3 through interactions with other transcription factors.

Previous studies have indicated that FOXO1 acts as an upstream transcription factor of

GRHL3[14]. Therefore, we aimed to investigate whether STAMBPL1 promotes GRHL3

transcription via FOXO1. Our experimental data revealed that knockdown of FOXO1 in TNBC

cells not only inhibited the GRHL3/HIF1α/VEGFA axis (Fig. 7A-D and SFig. 5A-D) but also

reversed the stimulatory effects of STAMBPL1 on this axis (Fig. 7E-H and SFig. 5E-H).

Furthermore, FOXO1 was found to bind to the promoter region of GRHL3 (Fig. 7I and SFig.

5I) and enhance its transcriptional activity (Fig. 7J). STAMBPL1 was shown to increase the

transcriptional activation of FOXO1 at the GRHL3 promoter (Fig. 7K), and it also bound to the

GRHL3 promoter; however, this interaction could be disrupted by FOXO1 knockdown (Fig.

7L).

To investigate the mechanism by which STAMBPL1 promotes GRHL3 transcription through

FOXO1, we conducted immunofluorescence and immunoprecipitation assays. We observed

the colocalization of STAMBPL1 and FOXO1 in the nucleus (Fig. 7M) and identified an

interaction between them in HCC1937 cells (Fig. 7N). Through the generation of a series of

Flag-FOXO1/GST-fused STAMBPL1 deletion mutants, we mapped the regions of the proteins

responsible for this interaction (SFig. 5F–G). The results of the GST pulldown assay indicated

that the N-terminus (1–140 aa) of STAMBPL1 interacted with FOXO1 (SFig. 5H). Additionally,

an immunoprecipitation assay revealed an interaction between the FOXO1 protein (250–596

aa) and STAMBPL1 (SFig. 5I). These findings suggest that STAMBPL1 promotes GRHL3

transcription by interacting with FOXO1.

The combination of VEGFR and FOXO1 inhibitors synergistically suppresses TNBC

xenograft growth.

Through analysis of the TCGA database, we determined that *STAMBPL1*, *FOXO1*, and

GRHL3 exhibited high expression levels in TNBC tumors compared with non-TNBC tumors

(Fig. 8A-C). To assess the potential therapeutic efficacy of cotreatment with the FOXO1

inhibitor AS1842856 and the VEGFR inhibitor apatinib *in vivo*, we conducted animal

experiments in nude mice. HCC1806 cells overexpressing STAMBPL1 were orthotopically

implanted into the mammary fat pads of 6-week-old female mice (n=16/group). Once the

tumor volume reached approximately 50 mm^3, the mice were divided into four subgroups to

receive apatinib (50 mg/kg, once every two days), AS1842856 (10 mg/kg, once every two

days), a combination of both drugs, or vehicle control for 20 days. Our findings indicated that

STAMBPL1 overexpression enhanced breast cancer cell growth *in vivo*. While the individual

inhibitory effects of the FOXO1 and VEGFR inhibitors on tumor growth were not significant,

the combined treatment markedly suppressed tumor growth in nude mice (Fig. 8E-G).

Importantly, the drug treatments did not affect the body weights of the mice (Fig. 8H). These

results suggest that the combined administration of AS1842856 and apatinib effectively

inhibits tumor growth in nude mice.

Discussion

In this study, we discovered that STAMBPL1 promotes angiogenesis in TNBC by activating the HIF1α/VEGFA pathway independently of its enzymatic activity. Through RNA-seq analysis, we revealed that STAMBPL1 positively regulates the transcription of *GRHL3*. Furthermore, we demonstrated that GRHL3 binds to the promoter of the *HIF1A* gene, thereby increasing its transcription. Additionally, we found that STAMBPL1 interacts with FOXO1 to facilitate the transcription of GRHL3/HIF1α/VEGFA. Importantly, we confirmed that the combination of the FOXO1 inhibitor AS1842856 and the VEGFR inhibitor apatinib effectively inhibited tumor growth in nude mice. These findings suggest that both STAMBPL1 and FOXO1 may be potential therapeutic targets for inhibiting angiogenesis in TNBC. This study is the first to uncover the role of STAMBPL1 in promoting angiogenesis through the FOXO1/GRHL3/HIF1α axis. Furthermore, a novel transcription factor, GRHL3, which regulates the transcription of HIF1α, was identified. These findings provide valuable insights for developing new therapeutic strategies to target TNBC.

The role of STAMBPL1 in tumors has not been fully recognized. Recent studies have reported its involvement in the epithelial–mesenchymal transition of various cancers, and its absence has been shown to affect the mesenchymal phenotype of lung and breast cancer[15]. Our previous research demonstrated that STAMBPL1 can stabilize MKP1 through deubiquitination and that the deletion of STAMBPL1 and MKP1 increases the sensitivity of breast cancer cells to cisplatin[16], suggesting that STAMBPL1 may be a potential therapeutic target for breast cancer. However, the mechanism by which STAMBPL1 activates the transcriptional activity of FOXO1 has not been elucidated, as STAMBPL1 does not stabilize the FOXO1 protein

through its deubiquitinating enzyme activity. We speculate that STAMBPL1 may recruit other

coactivators to FOXO1 transcription complexes. It will be important to develop a *Stambpl1*

knockout mouse model to investigate the exact role of STAMBPL1 in TNBC. Additionally, we

need to develop HIF1α and FOXO1 antibodies suitable for immunohistochemistry to detect

their expression in TNBC clinical samples.

Several studies have reported that FOXO1 inhibits tumor angiogenesis[17-21]. Studies have

shown that M2 macrophage-derived exosomal miR-942 promotes the migration and invasion

of lung adenocarcinoma cells and facilitates angiogenesis by binding to FOXO1 to alleviate

the inhibition of β-catenin, in which the upregulation of FOXO1 induces a decrease in cell

invasion and angiogenesis *in vitro*[17]. FOXO1 inhibits gastric cancer growth and angiogenesis

under hypoxic conditions via inactivation of the HIF1α-VEGF pathway, possibly in association

with SIRT1[18]. Cancer-associated fibroblast (CAF)-derived extracellular vesicles (EVs) deliver

miR-135b-5p into colorectal adenocarcinoma cells to downregulate FOXO1 and promote

HUVEC proliferation, migration, and angiogenesis[19]. Colorectal cancer cell-derived exosomes

overexpressing miR-183-5p promote the proliferation, migration and tube formation of HMEC-

1 (human microvascular endothelial cells) cells through the inhibition of FOXO1[20]. Bladder

cancer cell-derived exosomal miR-1247-3p facilitates angiogenesis by inhibiting FOXO1

expression[21]. However, the role of FOXO1 in breast cancer angiogenesis has not been

studied, and our study revealed that FOXO1 can promote the expression of HIF1α and

VEGFA, suggesting that it may play a role in promoting angiogenesis in breast cancer.

Studies have demonstrated that the AKT–FOXO1 signaling pathway regulates the expression

of GRHL3. To investigate this, the researchers utilized a previously published ChIP-seq

dataset for FOXO1 from human endometrial stromal cells. They reported that FOXO1

occupied BRD4-bound enhancers near the *GRHL3* gene. The authors subsequently

confirmed that the removal of EGF and insulin from the growth medium significantly increased

the expression of GRHL3, but the administration of the FOXO1 inhibitor AS1842856 partially

blocked the induced expression of GRHL3[14]. These results suggest that FOXO1 plays a

regulatory role upstream of GRHL3, and our study also confirmed that FOXO1 promotes the

transcription of *GRHL3* by binding to its promoter.

GRHL3 is a highly conserved epidermal-specific developmental transcription factor that has

recently gained attention in the field of cancer research[22]. To date, only a few genes regulated

by GRHL3 have been identified. Studies have shown that GRHL3 levels are significantly

reduced in human skin and head and neck squamous cell carcinomas and suggest that

GRHL3 is a key tumor suppressor pathway in squamous cell carcinomas[23]. GRHL3 was also

found to be induced by TNF-α in the mammary carcinoma cell line MCF-7 and was identified

as a TNFα-induced endothelial cell migration factor with promigratory activity as high as that

of VEGF[24]. However, the specific role of GRHL3 in breast cancer is unclear. Studies have

confirmed that GRHL3 strongly stimulates endothelial cell migration, which is consistent with

an angiogenic, protumorigenic function[24]. In our study, we demonstrated that GRHL3

promotes TNBC angiogenesis. HIF1α, a known regulator of angiogenesis, is regulated by

growth factors[25,26], cytokines[27] and mitogens[28,29]. The EGFR/Akt pathway is a known positive

regulator of HIF1α, potentially through mTOR[25] or independent of it[30,31]. While the

mechanisms regulating HIFα protein expression have been extensively studied, those

modulating HIF1α transcriptional activity remain unclear. Our study reveals for the first time

that GRHL3 promotes transcriptional activity by binding to the promoter of the *HIF1α* gene.

Conclusions

In summary, STAMBPL1 promoted TNBC angiogenesis by activating the GRHL3/HIF1A

pathway via FOXO1 in a non-enzymatic manner. These findings highlight the significant role

of STAMBPL1 in TNBC angiogenesis and suggest that targeting the

STAMBL1/FOXO1/GRHL3/HIF1α/VEGFA axis could be a potential therapeutic strategy to

inhibit angiogenesis in TNBC.

List of abbreviations

TNBC: triple-negative breast cancer

ER: estrogen receptor

PR: progesterone receptor

HER2: human epidermal growth factor receptor 2

EGFR: epidermal growth factor receptor

PARP: poly (ADP-ribose) polymerase

VEGFA: vascular endothelial growth factor A

HUVECs: primary human umbilical vein endothelial cells

CM: conditioned medium

Supplementary information

Supplementary Data (Supplementary Material 1) are available at JECCR Online.

Acknowledgements

We sincerely thank the team members for their dedication to this study.

Authors' contributions

Ceshi Chen, Huifeng Zhang, Wenmin Cao and Huichun Liang: conceptualization; Huan Fang, Huichun Liang and Ceshi Chen: data curation; Huan Fang and Huichun Liang: formal analysis, methodology and writing-original draft with help from Chuanyu Yang, Dewei Jiang and Qianmei Luo; Ceshi Chen, Huichun Liang, Huifeng Zhang, and Wenmin Cao: funding acquisition, project administration, validation, supervision and writing-review & editing.

Funding

This work was supported by National Key Research and Development Program of China (2023YFA1800500, 2020YFA0112300), National Natural Science Foundation of China (U2102203 to CC, 82203413 to HL), Biomedical Projects of Yunnan Key Science and Technology Program (202302AA310046 to CC), Yunnan Fundamental Research Projects (202301AS070050 to CC, 202201AT070290 to HL), Yunnan Revitalization Talent Support Program (Yunling Shcolar Project to CC), Yunnan (Kunming) Academician Expert Workstation (grant No. YSZJGZZ-2020025 to CC), Kunming University of Science and Technology-The First People's Hospital of Yunnan Province Joint Major Project (No. KUST-KH2022005Z), and Center of Clinical Pharmacy of the First People's Hospital of Yunnan Province Open Project (No. 2023YJZX-YX04).

Declarations

Ethics approval and consent to participate

All experimental procedures involving animals were performed in accordance to the Animal Welfare Law of China and the Guidelines for Animal Experiment of Kunming Institute of Zoology, Chinese Academy of Sciences.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the

corresponding author on reasonable request.

Competing interests

The authors declare that they have no conflicts of interest.

References

1.   Geyer, F.C., Pareja, F., Weigelt, B., Rakha, E., Ellis, I.O., Schnitt, S.J., and Reis-Filho, J.S.
(2017). The Spectrum of Triple-Negative Breast Disease: High- and Low-Grade Lesions.
Am J Pathol *187*, 2139-2151. 10.1016/j.ajpath.2017.03.016.
2.   Zagami, P., and Carey, L.A. (2022). Triple negative breast cancer: Pitfalls and progress.
NPJ Breast Cancer *8*, 95. 10.1038/s41523-022-00468-0.
3.   Garcia, J., Hurwitz, H.I., Sandler, A.B., Miles, D., Coleman, R.L., Deurloo, R., and Chinot,
O.L. (2020). Bevacizumab (Avastin(R)) in cancer treatment: A review of 15 years of clinical
experience and future outlook. Cancer Treat Rev *86*, 102017. 10.1016/j.ctrv.2020.102017.
4.   Detmar, M. (2000). Tumor angiogenesis. J Investig Dermatol Symp Proc *5*, 20-23.
10.1046/j.1087-0024.2000.00003.x.
5.   Sato, Y., Yoshikawa, A., Yamagata, A., Mimura, H., Yamashita, M., Ookata, K., Nureki, O.,
Iwai, K., Komada, M., and Fukai, S. (2008). Structural basis for specific cleavage of Lys 63-
linked polyubiquitin chains. Nature *455*, 358-362. 10.1038/nature07254.
6.   Chen, X., Shi, H., Bi, X., Li, Y., and Huang, Z. (2019). Targeting the deubiquitinase
STAMBPL1 triggers apoptosis in prostate cancer cells by promoting XIAP degradation.
Cancer Lett *456*, 49-58. 10.1016/j.canlet.2019.04.020.
7.   Woo, S.M., Seo, S.U., Kubatka, P., Min, K.J., and Kwon, T.K. (2019). Honokiol Enhances
TRAIL-Mediated Apoptosis through STAMBPL1-Induced Survivin and c-FLIP
Degradation. Biomolecules *9*. 10.3390/biom9120838.
8.   Liu, R., Yang, G., Bao, M., Zhou, Z., Mao, X., Liu, W., Jiang, X., Zhu, D., Ren, X., Huang, J.,
and Chen, C. (2022). STAMBPL1 promotes breast cancer cell resistance to cisplatin
partially by stabilizing MKP-1 expression. Oncogene *41*, 2265-2274. 10.1038/s41388-
022-02252-7.
9.   Xing, Y.Q., Li, A., Yang, Y., Li, X.X., Zhang, L.N., and Guo, H.C. (2018). The regulation of
FOXO1 and its role in disease progression. Life Sci *193*, 124-131.
10.1016/j.lfs.2017.11.030.
10.   Yu, J.M., Sun, W., Wang, Z.H., Liang, X., Hua, F., Li, K., Lv, X.X., Zhang, X.W., Liu, Y.Y., Yu,
J.J., et al. (2019). TRIB3 supports breast cancer stemness by suppressing FOXO1

degradation and enhancing SOX2 transcription. Nat Commun *10*, 5720.
10.1038/s41467-019-13700-6.

11.　Cui, Q., Sun, J., Yuan, J., Li, J., Yang, C., Du, G., Zhou, C., Qiu, P., Gao, J., Zhang, Y., et al.
(2024). DNA damage chemotherapeutic drugs suppress basal-like breast cancer growth
by down-regulating the transcription of the FOXO1-KLF5 axis. Genes Dis *11*, 91-94.
10.1016/j.gendis.2023.03.028.

12.　Jeon, H.H., Yu, Q., Lu, Y., Spencer, E., Lu, C., Milovanova, T., Yang, Y., Zhang, C.,
Stepanchenko, O., Vafa, R.P., et al. (2018). FOXO1 regulates VEGFA expression and
promotes angiogenesis in healing wounds. J Pathol *245*, 258-264. 10.1002/path.5075.

13.　Chen, C., Sun, X., Ran, Q., Wilkinson, K.D., Murphy, T.J., Simons, J.W., and Dong, J.T.
(2005). Ubiquitin-proteasome degradation of KLF5 transcription factor in cancer and
untransformed epithelial cells. Oncogene *24*, 3319-3327. 10.1038/sj.onc.1208497.

14.　Nagarajan, S., Bedi, U., Budida, A., Hamdan, F.H., Mishra, V.K., Najafova, Z., Xie, W., Alawi,
M., Indenbirken, D., Knapp, S., et al. (2017). BRD4 promotes p63 and GRHL3 expression
downstream of FOXO in mammary epithelial cells. Nucleic Acids Res *45*, 3130-3145.
10.1093/nar/gkw1276.

15.　Ambroise, G., Yu, T.T., Zhang, B., Kacal, M., Hao, Y., Queiroz, A.L., Ouchida, A.T., Lindskog,
C., Norberg, E., and Vakifahmetoglu-Norberg, H. (2020). Systematic analysis reveals a
functional role for STAMBPL1 in the epithelial-mesenchymal transition process across
multiple carcinomas. *123*, 1164-1177. 10.1038/s41416-020-0972-x.

16.　Liu, R., Yang, G., Bao, M., Zhou, Z., Mao, X., Liu, W., Jiang, X., Zhu, D., Ren, X., and Huang,
J. (2022). STAMBPL1 promotes breast cancer cell resistance to cisplatin partially by
stabilizing MKP-1 expression. *41*, 2265-2274. 10.1038/s41388-022-02252-7.

17.　Wei, K., Ma, Z., Yang, F., Zhao, X., Jiang, W., Pan, C., Li, Z., Pan, X., He, Z., Xu, J., et al.
(2022). M2 macrophage-derived exosomes promote lung adenocarcinoma progression
by delivering miR-942. Cancer letters *526*, 205-216. 10.1016/j.canlet.2021.10.045.

18.　Kim, S.Y., Ko, Y.S., Park, J., Choi, Y., Park, J.W., Kim, Y., Pyo, J.S., Yoo, Y.B., Lee, J.S., and
Lee, B.L. (2016). Forkhead Transcription Factor FOXO1 Inhibits Angiogenesis in Gastric
Cancer in Relation to SIRT1. Cancer research and treatment *48*, 345-354.
10.4143/crt.2014.247.

19.　Dai, X., Xie, Y., and Dong, M. (2022). Cancer-associated fibroblasts derived extracellular
vesicles promote angiogenesis of colorectal adenocarcinoma cells through miR-135b-
5p/FOXO1 axis. *23*, 76-88. 10.1080/15384047.2021.2017222.

20.　Shang, A., Wang, X., Gu, C., Liu, W., Sun, J., Zeng, B., Chen, C., Ji, P., Wu, J., Quan, W., et
al. (2020). Exosomal miR-183-5p promotes angiogenesis in colorectal cancer by
regulation of FOXO1. Aging *12*, 8352-8371. 10.18632/aging.103145.

21.　Liu, Z., Du, D., and Zhang, S. (2024). Tumor-derived exosomal miR-1247-3p promotes
angiogenesis in bladder cancer by targeting FOXO1. *25*, 2290033.
10.1080/15384047.2023.2290033.

22.　Darido, C., and Jane, S.M. (2010). Grhl3 and GEF19 in the front rho. Small GTPases *1*,
104-107. 10.4161/sgtp.1.2.13620.

23.　Darido, C., Georgy, S.R., Wilanowski, T., Dworkin, S., Auden, A., Zhao, Q., Rank, G.,
Srivastava, S., Finlay, M.J., Papenfuss, A.T., et al. (2011). Targeting of the tumor
suppressor GRHL3 by a miR-21-dependent proto-oncogenic network results in PTEN

512   loss and tumorigenesis. Cancer cell *20*, 635-648. 10.1016/j.ccr.2011.10.014.

513   24.   Guardiola-Serrano, F., Haendeler, J., Lukosz, M., Sturm, K., Melchner, H., and Altschmied,
514   J. (2008). Gene trapping identifies a putative tumor suppressor and a new inducer of cell
515   migration. Biochemical and biophysical research communications *376*, 748-752.
516   10.1016/j.bbrc.2008.09.070.

517   25.   Fukuda, R., Hirota, K., Fan, F., Jung, Y.D., Ellis, L.M., and Semenza, G.L. (2002). Insulin-like
518   growth factor 1 induces hypoxia-inducible factor 1-mediated vascular endothelial
519   growth factor expression, which is dependent on MAP kinase and phosphatidylinositol
520   3-kinase signaling in colon cancer cells. The Journal of biological chemistry *277*, 38205-
521   38211. 10.1074/jbc.M203781200.

522   26.   Zhong, H., Chiles, K., Feldser, D., Laughner, E., Hanrahan, C., Georgescu, M.M., Simons,
523   J.W., and Semenza, G.L. (2000). Modulation of hypoxia-inducible factor 1alpha
524   expression by the epidermal growth factor/phosphatidylinositol 3-
525   kinase/PTEN/AKT/FRAP pathway in human prostate cancer cells: implications for tumor
526   angiogenesis and therapeutics. Cancer research *60*, 1541-1545.

527   27.   Hellwig-Bürgel, T., Stiehl, D.P., Wagner, A.E., Metzen, E., and Jelkmann, W. (2005).
528   Review: hypoxia-inducible factor-1 (HIF-1): a novel transcription factor in immune
529   reactions. Journal of interferon & cytokine research : the official journal of the
530   International Society for Interferon and Cytokine Research *25*, 297-310.
531   10.1089/jir.2005.25.297.

532   28.   Kasuno, K., Takabuchi, S., Fukuda, K., Kizaka-Kondoh, S., Yodoi, J., Adachi, T., Semenza,
533   G.L., and Hirota, K. (2004). Nitric oxide induces hypoxia-inducible factor 1 activation that
534   is dependent on MAPK and phosphatidylinositol 3-kinase signaling. The Journal of
535   biological chemistry *279*, 2550-2558. 10.1074/jbc.M308197200.

536   29.   Li, J., Davidson, G., Huang, Y., Jiang, B.H., Shi, X., Costa, M., and Huang, C. (2004). Nickel
537   compounds act through phosphatidylinositol-3-kinase/Akt-dependent, p70(S6k)-
538   independent pathway to induce hypoxia inducible factor transactivation and Cap43
539   expression in mouse epidermal Cl41 cells. Cancer research *64*, 94-101. 10.1158/0008-
540   5472.can-03-0737.

541   30.   Mazure, N.M., Chen, E.Y., Laderoute, K.R., and Giaccia, A.J. (1997). Induction of vascular
542   endothelial growth factor by hypoxia is modulated by a phosphatidylinositol 3-
543   kinase/Akt signaling pathway in Ha-ras-transformed cells through a hypoxia inducible
544   factor-1 transcriptional element. Blood *90*, 3322-3331.

545   31.   Pore, N., Jiang, Z., Shu, H.K., Bernhard, E., Kao, G.D., and Maity, A. (2006). Akt1 activation
546   can augment hypoxia-inducible factor-1alpha expression by increasing protein
547   translation through a mammalian target of rapamycin-independent pathway. Molecular
548   cancer research : MCR *4*, 471-479. 10.1158/1541-7786.mcr-05-0234.

 Figures and legends

Figure 1. STAMBPL1 upregulates HIF1α in a non-enzymatic manner in TNBC cells. (A)

After STAMBPL1 was knocked down in HCC1806 and HCC1937 cells and then subjected to

hypoxia for 10 h, STAMBPL1 knockdown downregulated the protein level of HIF1α. **(B-C)**

RNA samples were collected after 10 h of hypoxia treatment following the knockdown of

STAMBPL1 in HCC1806 cells. RT–qPCR experiments revealed that the knockdown of

STAMBPL1 downregulated the mRNA levels of the HIF1α downstream targets *VEGFA* and

GLUT1. **(D-E)** RNA samples were collected after 10 hours of hypoxia treatment following the

knockdown of STAMBPL1 in HCC1937 cells. RT–qPCR experiments revealed that the

knockdown of STAMBPL1 downregulated the mRNA levels of the HIF1α downstream targets

VEGFA and *GLUT1*. **(F)** Overexpression of STAMBPL1/STAMBPL1-E292A upregulated

HIF1α protein expression in HCC1806 and HCC1937 cells stably overexpressing

STAMBPL1/STAMBPL1-E292A under normoxia. *: $P<0.05$, **: $P<0.01$, ***: $P<0.001$, *t* test.

Figure 2. STAMBPL1 promotes *HIF1A* transcription and activates the HIF1α/VEGFA

axis. (A) Western blot analysis was performed to confirm the knockdown of STAMBPL1 in

HCC1806 and HCC1937 cells. **(B)** RNA samples were collected after 10 h of hypoxia

treatment following the knockdown of STAMBPL1 in HCC1806 cells. RT–qPCR experiments

revealed that the knockdown of STAMBPL1 could downregulate the mRNA levels of HIF1α.

(C) RNA samples were collected after 10 hours of hypoxia treatment following the knockdown

of STAMBPL1 in HCC1937 cells. RT–qPCR experiments revealed that the knockdown of

STAMBPL1 downregulated the mRNA levels of HIF1α. **(D)** HCC1806 cells stably

overexpressing STAMBPL1 were used to knock down HIF1α via siRNA. The protein was then

collected and subjected to Western blotting experiments. The results showed that the

overexpression of STAMBPL1 promoted the expression of HIF1α, which could be rescued by

the knockdown of HIF1α. **(E)** HIF1α was knocked down in HCC1806 cells stably

overexpressing STAMBPL1 via siRNA. The RNA was then collected and subjected to RT–

qPCR experiments. The results showed that the overexpression of STAMBPL1 promoted the

expression of the VEGFA mRNA, which could be rescued by the knockdown of HIF1α. **(F)** In

HCC1937 cells stably overexpressing STAMBPL1, HIF1α was knocked down via siRNA, and

protein was collected. Western blotting experiments revealed that the overexpression of

STAMBPL1 promoted HIF1α expression, which could be rescued by the knockdown of HIF1α.

(G) In HCC1937 cells stably overexpressing STAMBPL1, HIF1α was knocked down via

siRNA, and RNA was collected. RT–qPCR experiments revealed that the overexpression of

STAMBPL1 promoted VEGFA mRNA expression, which could be rescued by the knockdown

of HIF1α. *: $P<0.05$, **: $P<0.01$, ***: $P<0.001$, t test.

Figure 3. STAMBPL1 in TNBC cells enhances the activity of HUVECs and promotes

TNBC angiogenesis. (A) In HCC1806 cells, STAMBPL1 was knocked down via siRNA, and

the cells were then subjected to hypoxia for 24 hours. The conditioned medium (CM)

collected from these cells was used to treat HUVECs. EdU assays revealed that the

conditioned medium after STAMBPL1 knockdown inhibited the proliferation of HUVECs. **(B)**

Statistical analysis of the EdU assay results. **(C)** In HCC1806 cells, STAMBPL1 was knocked

down via siRNA, and the cells were then subjected to hypoxia for 24 hours. The CM collected

from these cells was used to treat HUVECs. Wound healing assays demonstrated that the

conditioned medium after STAMBPL1 knockdown inhibited the migration of HUVECs. **(D)**

Statistical analysis of the wound healing assay results. **(E)** In HCC1806 cells, STAMBPL1 was

knocked down via siRNA, and the cells were then subjected to hypoxia for 24 hours. The CM

collected from these cells was used to treat HUVECs. A tube formation assay revealed that

the conditioned medium after STAMBPL1 knockdown inhibited the tube formation of

HUVECs. **(F)** Statistical analysis of the tube formation assay results. **(G)** The growth of TNBC

xenograft tumors was evaluated by photographing the tumors in nude mice to assess the role

of STAMBPL1. Knockdown of STAMBPL1 significantly inhibited tumor growth. **(H-I)** The

growth and weight of the transplanted tumors in the nude mice were statistically analyzed,

and the STAMBPL1sh#1 and STAMBPL1sh#2 groups were compared with the shLuc group.

(J) An immunohistochemical assay was used to detect the expression of the angiogenesis

marker CD31 in xenograft tumors. **(K)** The number of microvessels in the

immunohistochemical experiments was statistically analyzed. *: $P<0.05$, **: $P<0.01$, ***:

$P<0.001$, t test.

Figure 4. STAMBPL1 promotes *HIF1A* transcription via the upregulation of GRHL3. (A)

Venn diagram showing the overlap of differentially expressed genes identified via RNA-seq

analysis. **(B)** RT–qPCR experiments demonstrated that the knockdown of STAMBPL1

resulted in decreased mRNA levels of GRHL3 in HCC1806 cells. **(C)** In HCC1806 cells stably

overexpressing STAMBPL1, GRHL3 was knocked down via siRNA, after which the protein

was collected. Western blotting experiments revealed that the overexpression of STAMBPL1

promoted the expression of HIF1α, which could be rescued by the knockdown of GRHL3. **(D-**

F) In HCC1806 cells stably overexpressing STAMBPL1, GRHL3 was knocked down via

siRNA, and RNA was subsequently collected. RT–qPCR experiments revealed that the

overexpression of STAMBPL1 upregulated the mRNA expression of GRHL3, HIF1α and its

downstream target VEGFA, which could be rescued by the knockdown of GRHL3. **(G)**

Statistical analysis of the EdU assay results. **(H)** Statistical analysis of the wound healing

assay results. **(I)** Statistical analysis of the tube formation assay results. *: $P<0.05$, **: $P<0.01$,

***: $P<0.001$, t test.

Figure 5. GRHL3 enhances *HIF1A* transcription by binding to its promoter. (A) The

JASPAR website was used to predict the potential binding sequence of the transcription factor

GRHL3 to the HIF1α promoter. **(B)** Mutation pattern of the HIF1α promoter binding sequence.

(C) Overexpression of GRHL3 in HCC1806 cells stably overexpressing GRHL3 resulted in

increased protein expression of HIF1α. **(D)** ChIP–PCR experiments conducted in HCC1806

cells stably overexpressing GRHL3 revealed that GRHL3 could bind to the promoter region of

the target gene *HIF1A*. **(E)** A luciferase assay performed in HEK293T cells indicated that

GRHL3 could bind to the promoter sequence of *HIF1A* and increase its transcriptional activity.

(F) In HCC1806 cells, GRHL3 was knocked down by siRNA, and the cells were subjected to

hypoxia for 4 hours. Western blotting experiments revealed that the knockdown of GRHL3

downregulated the protein level of HIF1α. **(G-I)** In HCC1806 cells, GRHL3 was knocked down

by siRNA, followed by hypoxia treatment for 4 hours, after which RNA samples were

collected. RT–qPCR experiments revealed that GRHL3 knockdown was effective and that the

knockdown of GRHL3 could downregulate the mRNA levels of HIF1α and its downstream

target VEGFA. **(J)** In HCC1806 cells, GRHL3 was knocked down via siRNA, and the cells

were then subjected to hypoxia for 24 hours. The conditioned medium was collected and

used to treat HUVECs. EdU assays revealed that the conditioned medium after GRHL3

knowdown inhibited the proliferation of HUVECs. Statistical analysis of the EdU assay results

was performed. **(K)** In HCC1806 cells, GRHL3 was knocked down via siRNA, and the cells

were then subjected to hypoxia for 24 hours. The conditioned medium was collected and

used to treat HUVECs. Wound healing assays revealed that the conditioned medium after

GRHL3 knockdown inhibited the migration of HUVECs. Statistical analysis of the wound

healing assay results was performed. **(L)** In HCC1806 cells, GRHL3 was knocked down via

siRNA, and the cells were then subjected to hypoxia for 24 hours. The conditioned medium

was collected and used to treat HUVECs. The results of the tube formation assay revealed

that conditioned medium from GRHL3-knockdown HUVECs inhibited tube formation.

Statistical analysis of the tube formation assay results was performed. *: P<0.05, **: P<0.01,

***: P<0.001, t test.

Figure 6. GRHL3 enhances *HIF1A* transcription by binding to its promoter. (A) In

HCC1806 cells stably overexpressing GRHL3, HIF1α was knocked down by siRNA, and the

protein was subsequently collected. Western blotting experiments revealed that the

overexpression of GRHL3 promoted the expression of HIF1α, which could be rescued by the

knockdown of HIF1α. **(B)** In HCC1806 cells stably overexpressing GRHL3, HIF1α was

knocked down by siRNA, and then the RNA was collected. RT–qPCR experiments revealed

that the overexpression of GRHL3 promoted the expression of the VEGFA mRNA, which

could be rescued by the knockdown of HIF1α. **(C)** HCC1806 cells stably overexpressing

GRHL3 were used for the knockdown of HIF1α via siRNA. The conditioned medium was then

collected and used to treat HUVECs. EdU assays revealed that the overexpression of GRHL3

promoted the proliferation of HUVECs, which could be rescued by the knockdown of HIF1α.

(D) HCC1806 cells stably overexpressing GRHL3 were used for the knockdown of HIF1α via

siRNA. The conditioned medium was then collected and used to treat HUVECs. A wound

healing assay revealed that the overexpression of GRHL3 promoted the migration of

HUVECs, which could be rescued by the knockdown of HIF1α. **(E)** HCC1806 cells stably

overexpressing GRHL3 were used for the knockdown of HIF1α via siRNA. The conditioned

medium was then collected and used to treat HUVECs. The tube formation assay revealed

that the overexpression of GRHL3 promoted the tube formation of HUVECs, which could be

rescued by the knockdown of HIF1α. **(F)** RT–qPCR was used to detect the knockdown of

GRHL3 in HCC1806 cells. **(G)** Photographs of xenograft tumors from nude mice were taken

to evaluate the role of GRHL3 in the growth of TNBC xenograft tumors. The results showed

that the knockdown of GRHL3 significantly inhibited tumor growth. **(H)** The weights of the

transplanted tumors from the nude mice were statistically analyzed, and the shGRHL3 1# and

shGRHL3 2# groups were compared with the shLuc group. **(I)** The growth of transplanted

tumors in nude mice was statistically analyzed, and the shGRHL3 1# and shGRHL3 2#

groups were compared with the shLuc group. **(J)** An immunohistochemical assay was

performed to detect the expression of the angiogenesis marker CD31 in xenograft tumors. **(K)**

Statistical analysis of the number of microvessels in the immunohistochemical experiments. *:

P<0.05, **: *P*<0.01, ***: *P*<0.001, *t* test.

Figure 7. STAMBPL1 mediates *GRHL3* transcription by interacting with FOXO1. (A)

Western blotting experiments revealed that the knockdown of FOXO1 in HCC1806 cells

followed by hypoxia treatment for 4 hours inhibited HIF1α protein expression. **(B-D)** RT–

qPCR experiments revealed that the knockdown of FOXO1 in HCC1806 cells followed by

hypoxia treatment for 4 hours decreased the mRNA levels of GRHL3/HIF1α/VEGFA. **(E)** In

HCC1806 cells stably overexpressing STAMBPL1, protein samples were collected after

FOXO1 was knocked down via siRNA. Western blotting experiments revealed that the

overexpression of STAMBPL1 promoted the expression of HIF1α, which could be rescued by

the knockdown of FOXO1. **(F-H)** In HCC1806 cells stably overexpressing STAMBPL1, RNA

samples were collected after the knockdown of FOXO1 via siRNA. RT–qPCR experiments

revealed that the overexpression of STAMBPL1 promoted the mRNA expression of GRHL3,

HIF1α and VEGFA, which could be rescued by the knockdown of FOXO1. **(I)** An endogenous

ChIP–PCR assay was performed using an anti-FOXO1 antibody in HCC1806 cells. **(J)** A

luciferase assay performed in HEK293T cells revealed that FOXO1 was able to bind to the

promoter sequence of GRHL3 and increase its transcriptional activity. **(K)** A luciferase assay

conducted in HEK293T cells revealed that STAMBPL1 enhanced the activation of the GRHL3

promoter by FOXO1. **(L)** ChIP–PCR experiments were performed in HCC1806 cells stably

overexpressing STAMBPL1 after knocking down FOXO1 via siRNA. **(M)** PCDH-STAMBPL1-

3×Flag and PCDH-FOXO1-3×Flag plasmids were cotransfected into HEK293T cells, and

then, immunofluorescence experiments were performed. Red represents STAMBPL1

staining, green represents FOXO1 staining, and blue represents DAPI staining. **(N)**

Endogenous STAMBPL1 protein was immunoprecipitated from HCC1937 cells via an anti-

FOXO1 antibody. Immunoglobulin G (IgG) served as the negative control. Endogenous

STAMBPL1 was detected via western blotting. *: $P<0.05$, **: $P<0.01$, ***: $P<0.001$, t test.

Figure 8. The combination of VEGFR and FOXO1 inhibitors synergistically suppresses

TNBC xenograft growth. (A-C) TCGA database analysis revealed high expression levels of

STAMBPL1, FOXO1, and GRHL3 in TNBC. **(D)** The effect of STAMBPL1 overexpression in

HCC1806 cells was detected by Western blotting. **(E)** The effect of drug treatment on

transplanted tumors in nude mice was evaluated by photographing the tumors. The

overexpression of STAMBPL1 promoted tumor growth, whereas the combination of the

FOXO1 inhibitor AS1842856 (10 mg/kg, every two days) and the VEGFR inhibitor apatinib (50

mg/kg, every two days) significantly inhibited tumor growth. **(F-G)** The weight and growth of

the transplanted tumors in the nude mice were statistically analyzed. The vector control group

and the STAMBPL overexpression group were compared. The nondrug group and the

combined drug group in the vector control group were compared. The nondrug group and the

combined drug group in the STAMBPL overexpression group were compared. **(H)** The final

weights of the nude mice were statistically analyzed. *: $P<0.05$, **: $P<0.01$, ***: $P<0.001$, t

test.

Figure 9. The working model of this study. STAMBPL1 interacts with FOXO1 to promote

TNBC angiogenesis by activating the GRHL3/HIF1α/VEGFA axis.

Additional material

Supplementary Material 1

Figure S1. STAMBPL1 in TNBC cells enhances the activity of HUVECs. (A) HCC1937

cells were transfected with STAMBPL1 siRNA and subjected to hypoxia for 24 hours. The

conditioned medium collected from these cells was then used to treat HUVECs. EdU assay

revealed that the conditioned medium after STAMBPL1 knockdown inhibited the proliferation

of HUVECs. **(B)** Statistical analysis of the EdU assay results. **(C)** HCC1937 cells were

transfected with STAMBPL1 siRNA and subjected to hypoxia for 24 hours. The conditioned

medium collected from these cells was used to treat HUVECs. Wound healing assay

demonstrated that the conditioned medium after STAMBPL1 knockdown inhibited the

migration of HUVECs. **(D)** Statistical analysis of the wound healing assay results. **(E)**

HCC1937 cells were transfected with STAMBPL1 siRNA and subjected to hypoxia for 24

hours. The conditioned medium collected from these cells was used to treat HUVECs. Tube

formation assay showed that the conditioned medium after STAMBPL1 knockdown inhibited

tube formation of HUVECs. **(F)** Statistical analysis of the tube formation assay results. *:

P<0.05, **: *P*<0.01, ***: *P*<0.001, *t* test.

Figure S2. STAMBPL1 in TNBC cells enhances the activity of HUVECs. (A and D)

HCC1806 and HCC1937 cells were stably overexpressing STAMBPL1, and HIF1α was

knocked down using siRNA. The conditioned medium was then collected and used to treat

HUVECs. EdU assay results demonstrated that the overexpression of STAMBPL1 promoted

the proliferation of HUVECs, which could be rescued by knockdown of HIF1α. Statistical

analysis of the EdU assay results was performed. **(B and E)** HCC1806 and HCC1937 cells

stably overexpressing STAMBPL1 were used to knock down HIF1α using siRNA. The

conditioned medium was collected and used to treat HUVECs. Wound healing assay results

showed that the overexpression of STAMBPL1 promoted the migration of HUVECs, which

could be rescued by knockdown of HIF1α. Statistical analysis of the wound healing assay

results was performed. **(C and F)** HCC1806 and HCC1937 cells stably overexpressing

STAMBPL1 were used to knock down HIF1α using siRNA. The conditioned medium was

collected and used to treat HUVECs. Tube formation assay results showed that the

overexpression of STAMBPL1 promoted the tube formation of HUVECs, which could be

rescued by knockdown of HIF1α. Statistical analysis of the tube formation assay results was

performed. *: $P<0.05$, **: $P<0.01$, ***: $P<0.001$, t test.

Figure S3. STAMBPL1 promotes *HIF1A* transcription via upregulating GRHL3. (A) RNA-

seq result: the list of 18 genes downregulated after STAMBPL1 knockdown in HCC1806 cells.

(B) RT-qPCR experiments showed that knockdown of STAMBPL1 inhibited GRHL3 mRNA

levels in HCC1937 cells. **(C)** In HCC1937 cells stably overexpressing STAMBPL1, GRHL3

was knocked down using siRNA, and then the protein was collected. Western blotting

experiments showed that overexpression of STAMBPL1 promoted the expression of HIF1α,

which could be rescued by knockdown of GRHL3. **(D-F)** In HCC1937 cells stably

overexpressing STAMBPL1, GRHL3 was knocked down using siRNA and then the RNA was

collected. RT-qPCR experiments showed that overexpression of STAMBPL1 promoted the

mRNA expression of GRHL3, HIF1α and its downstream target VEGFA, which could be

rescued by knockdown of GRHL3. **(G)** Statistical analysis of EdU assay. **(H)** Statistical

analysis of wound healing assay. **(I)** Statistical analysis of tube formation assay. *: $P<0.05$, **:

$P<0.01$, ***: $P<0.001$, t test.

Figure S4. GRHL3 enhances *HIF1A* transcription by binding to its promoter. (A) In

HCC1937 cells, knockdown of GRHL3 using siRNA and treated with hypoxia for 4 hours.

Western blotting experiments showed that knockdown of GRHL3 could down-regulate the

protein level of HIF1α. **(B-D)** In HCC1937 cells, GRHL3 was knocked down by siRNA,

followed by hypoxia treatment for 4 hours, and then RNA samples were collected. RT-qPCR

experiments showed that GRHL3 knockdown was effective, and knockdown of GRHL3 could

down-regulate the mRNA levels of HIF1α and its downstream target VEGFA. **(E)** In HCC1937

cells, GRHL3 was knocked down by siRNA and then treated with hypoxia for 24 hours, and

then the conditioned medium was collected to treat HUVECs, EdU assay showed that the

conditioned medium after knocking down GRHL3 inhibited the proliferation of HUVECs. **(F)** In

HCC1937 cells, GRHL3 was knocked down by siRNA and then treated with hypoxia for 24

hours, and then the conditioned medium was collected to treat HUVECs, wound healing

assay showed that the conditioned medium after knocking down GRHL3 inhibited the

migration of HUVECs. **(G)** In HCC1937 cells, GRHL3 was knocked down by siRNA and then

treated with hypoxia for 24 hours, and then the conditioned medium was collected to treat

HUVECs, the tube formation assay showed that the conditioned medium after knocking down

GRHL3 inhibited tube formation of HUVECs. **(H)** In HCC1937 cells stably overexpressing

GRHL3, HIF1α was knocked down using siRNA, and then the protein was collected. Western

blotting experiments showed that overexpression of GRHL3 promoted the expression of

HIF1α, which could be rescued by knockdown of HIF1α. **(I)** In HCC1937 cells stably

overexpressing GRHL3, HIF1α was knocked down using siRNA and then the RNA was

collected. RT-qPCR experiments showed that overexpression of GRHL3 promoted the mRNA

expression of VEGFA, which could be rescued by knockdown of HIF1α. **(J)** Statistical

analysis of EdU assay results. **(K)** Statistical analysis of wound healing assay results. **(L)**

Statistical analysis of tube formation assay results. *: $P<0.05$, **: $P<0.01$, ***: $P<0.001$, *t* test.

Figure S5. STAMBPL1 mediates *GRHL3* transcription by interacting with FOXO1. (A)

Western blotting experiments showed that knockdown of FOXO1 in HCC1937 cells followed

by hypoxia treatment for 4 hours inhibited HIF1α protein expression. **(B-D)** RT-qPCR

experiments showed that knockdown of FOXO1 in HCC1937 cells followed by hypoxia

treatment for 4 hours inhibited the mRNA levels of GRHL3/HIF1α/VEGFA. **(E)** The JASPAR

website predicts the possible binding sequence of transcription factor FOXO1 to the GRHL3

promoter and its mutation pattern map. **(F)** The truncated structure diagram of STAMBPL1.

(G) The truncated structure diagram of FOXO1. **(H)** The full-length and truncated plasmids of

pEBG-STAMBPL1-GST and pCDH-FOXO1-no tag were transfected into HEK293T cells, and

the proteins were collected for GST-pull down assay at 48 hours after transfection. **(I)** The full-

length and truncated plasmids of pCDH-FOXO1-3×Flag and pEBG-STAMBPL1-GST were

transfected into HEK293T cells, and the proteins were collected for IP-Flag assay at 48 hours

after transfection. *: $P<0.05$, **: $P<0.01$, ***: $P<0.001$, t test.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A. GRHL binding motif

B. HIF1α Promoter
-2000 ———————— 0

WT -875 TGA**AATTGGTT**TATA -860
TATA**AACCAATT**TCA

Mutant -875 TGA**GGCCTTCC**TATA -860
TATA**GGAAGGCC**TCA

C.
GRHL3-Flag − +
Vector + − KDa
HIF1α-L 120
HIF1α-S 120
Flag 70
β-Actin 42
HCC1806

D.
Marker | Input | IP-Flag
Vector | GRHL3-Flag | Vector | GRHL3-Flag
250bp
HCC1806
HIF1α Promoter

E. HEK293T
Relative Luciferase Activity
***
***
Vector+PGL3
Vector+PGL3-HIF1α-WT
GRHL3+PGL3
GRHL3+PGL3-HIF1α-WT
GRHL3+PGL3-HIF1α-Mut

F.
siGRHL3 2# − − + − − +
siGRHL3 1# − + − − + −
siCon + − − + − −
Hypoxia(h) 0 0 0 4 4 4 KDa
HIF1α 120
β-Actin 42
HCC1806

G. HCC1806 *GRHL3*
Relative mRNA Expression (Fold Change)
*** ***
siCon-Normoxia
siCon-Hypoxia
siGRHL3 1#-Hypoxia
siGRHL3 2#-Hypoxia

H. HCC1806 *HIF-1α*
Relative mRNA Expression (Fold Change)
* **
siCon-Normoxia
siCon-Hypoxia
siGRHL3 1#-Hypoxia
siGRHL3 2#-Hypoxia

I. HCC1806 *VEGFA*
Relative mRNA Expression (Fold Change)
*** *** ***
siCon-Normoxia
siCon-Hypoxia
siGRHL3 1#-Hypoxia
siGRHL3 2#-Hypoxia

J. HUVECs in HCC1806 CM
EdU positive cells (%)
Normoxia | Hypooxia
* ** ***
siCon | siGRHL3 1# | siGRHL3 2# | siCon | siGRHL3 1# | siGRHL3 2#

K. HUVECs in HCC1806 CM
Wound closure (%)
Normoxia | Hypooxia
** * **
siCon | siGRHL3 1# | siGRHL3 2# | siCon | siGRHL3 1# | siGRHL3 2#

L. HUVECs in HCC1806 CM
Total branch length
Normoxia | Hypooxia
** ** **
siCon | siGRHL3 1# | siGRHL3 2# | siCon | siGRHL3 1# | siGRHL3 2#

Figure 6

Figure 7

Figure 8

Figure 9