1

4   The size-weight illusion and beyond: a new model of perceived weight

7   Veronica Pisu[1*], Erich W. Graf[1], Wendy J. Adams[1]

11   [1] School of Psychology, University of Southampton, Southampton, United Kingdom

14   * Corresponding author

15   Email: Veronica.Pisu@soton.ac.uk (VP)

Abstract

In the size-weight illusion (SWI), the smaller of two same-weight, same apparent material objects is perceived as heavier. The SWI has proved difficult to explain via traditional Bayesian models, which predict the opposite effect: expected weight from size (smaller = lighter) should be integrated with felt weight, such that the smaller object should be perceptually lighter. Other authors have proposed that weight and density are combined according to Bayesian principles, or that Bayesian models incorporating efficient coding can predict the SWI via 'likelihood repulsion'. These models, however, have been evaluated only under the narrow conditions of typical SWI stimuli. Here we establish a general model of perceived weight for pairs of objects that differ in weight and / or density and / or size by varying amounts. In a visuo-haptic task, participants (N = 30) grasped and lifted pairs of cubes, and reported their perceived heaviness. We report that the SWI occurs even at very small density differences, repudiating the idea that the illusion requires a significant conflict between expected and felt weight. Across all object pairs, perceived weight was well explained by a model (R^2 = .98) that includes a positive influence of both objects' weights and the judged object's density, but a negative influence of the other object's density. Critically, the influence of both densities on perceived weight is strongly modulated by weight difference, being three times as large for zero / small weight differences than large differences. Thus, it is only under the unusual conditions of typical SWI studies that we confuse weight with density to a substantial extent. Unlike existing models, that are inconsistent with our more comprehensive dataset, our model provides a quantitative, accurate and generalised account of weight perception for pairs of objects across various weight and size conditions.

Author summary

Knowing how heavy an object is allows us to grasp and lift it efficiently and without mishaps. Surprisingly, humans make systematic errors when judging the weight of objects of different

size. For example, when two objects differ in size but have identical weight, the smaller object

feels heavier. This is the 'size-weight illusion'. The illusion is strong and reliable and occurs

even when we know that the two objects actually have the same weight. The size-weight illusion

demonstrates that the human perceptual system doesn't work like a weighing scale, but

instead takes into account other object features such as size or density, alongside weight. In

this paper, we present a model that allows us to predict perceived weight in the size-weight

illusion and across a wide range of objects of different weight / size / density, where previous

models have failed.

1. Introduction

In the well-known size-weight illusion (SWI) (1), participants are typically presented with

one large and one small object, apparently of the same material, but in fact of equal weight. On

lifting, the smaller object is perceived as heavier than the larger one (Figure 1A). The illusion is

robust, persisting even when the lifter knows that the objects have equal weight (2,3), and after

grasping forces have adapted to the actual weight of the objects (4).

Figure 1: The size weight illusion. (A) Typical procedure for measuring the size-weight illusion. The subject lifts two objects of equal weight, but different size. The smaller object is perceived to be heavier than the larger one. (B) The SWI is 'anti-Bayesian'. Stimulus pairs are plotted in a two-dimension stimulus space: log volume ratio (x-axis) and log weight ratio (y-axis). A value of zero indicates that the two objects have equal volume (or weight). The green markers show SWI stimuli (equal weight but different volume) and arrows show the empirical bias, i.e. the perceived weight ratio, for the same stimuli (data taken from the current experiment). The solid red line represents an equal density prior (log density ratio=0). The dotted red line indicates another plausible prior based on object statistics (5): the larger object is heavier but slightly less dense. In the SWI illusion, perceived weight is biased *away* from these priors. (C) SWI and three hypothetical priors ($R_1 - R_3$) over log volume ratio and log weight ratio, as suggested by Peters and colleagues (6). Each prior corresponds to a different density / volume relationship. R_2 (smaller is denser and heavier: dashed black line) would predict the SWI. R_1 = equal-density prior (red line); R_3 = smaller is less dense and lighter (dotted blue line).

Although the SWI has long been investigated as a means to understanding human

heaviness perception, its mechanism remains largely unexplained (for reviews, see (3,7,8)).

Early descriptive accounts suggested that the SWI is caused by a contrast / repulsion effect,

where the felt weights of the two objects conflict with those predicted by assuming equal

density. Other accounts have suggested that the lifter confounds / combines felt weight with

other physical features (generally density: (9–11); but also rotational inertia: (12,13)).

In 'contrast' or 'violated expectation' accounts it is noted that object size is used as a

predictive feature for object weight (14–16). Smaller objects are expected to be lighter than

larger objects, particularly when they are apparently made of the same material – that is, of

equal density (see Figure 1B). Upon lifting, the smaller object is heavier than expected, and this

disparity between expectation and sensory evidence is suggested to cause the illusion.

Evidence for this conflict between expected and actual weight comes from the forces used to

grip and lift the two objects. On initial lifts, participants use more force to lift the larger

(expected to be heavier) object than to lift the smaller one (4,17). These misapplied forces have

thus been suggested to drive the illusion: the larger object is lifted faster and more easily than

expected, and vice versa for the smaller object, producing a mismatch between the expected

and actual sensory feedback (7,17). However, it has been widely reported that grasping forces

adjust to the weight of the object after a few lifts, while the illusion persists (4,18–20)).

Because the illusion is at odds with the perceiver's prior expectations (that the smaller

object should be lighter) the SWI has been labelled 'anti-Bayesian' (21,22). The Bayesian

framework has successfully described numerous perceptual phenomena in terms of the

integration (generally a weighted averaging) of prior expectations with current sensory cues (23–

28). In the SWI, however, the Bayesian prediction is that the smaller object will be perceived as

lighter than the larger object, given an assumption of equal density for the two objects. Figure 1

illustrates this: Figure 1B represents the stimulus space, with different possible combinations

of two stimuli (their weight, volume and density log ratios). Two theoretically plausible,

hypothetical priors are shown on these axes: (i) the larger object is heavier and same-density

(solid line), or (ii) the larger object is heavier but slightly less dense (dashed line, reflecting

measured object statistics (5)). Both of these priors predict the opposite of the SWI (green dots

show stimuli, arrowheads give perceived log weight ratio, taken from current study). The SWI

involves biases *away* from the theoretically sensible priors, hence the 'anti-Bayesian' label. The

dashed black line in Figure 1C shows a hypothetical prior (smaller = heavier) that would predict

the SWI. However, this prior is implausible, given environmental statistics in which size and

weight are positively correlated.

Nonetheless, Bayesian models of the SWI have been proposed (6,29). Peters and

colleagues (6) proposed a Bayesian model that includes three sub-priors over the relationship

between the log volume ratio of two objects and their log density ratio (shown in Figure 1C).

Each sub-prior represents a categorical density relationship; R_1: objects have equal density, R_2:

smaller objects are denser and heavier than larger ones and R_3: smaller objects are less dense

(much lighter) than larger objects. When a typical SWI stimulus pair (e.g. log weight ratio = 0, log

volume ratio = 0.5) is encountered, the 'smaller is denser and heavier' prior (R_2 in Figure 1C)

dominates, resulting in an illusion. It is not clear whether the model can account for weight

perception more broadly as it was only evaluated on pairs of equal-weight objects (i.e., typical

SWI stimuli); predicted biases in weight perception depend on the volume and density ratios of

the stimuli and thus which sub-prior dominates perception.

Wei and Stocker (30,31) have suggested that contrast effects analogous to the SWI, i.e.

biases away from a prior, can be predicted by Bayesian models incorporating efficient coding of

stimulus properties (32,33). They note that when efficient coding maximises Fisher information

(i.e., to reflect the stimulus distribution in the environment), this results in likelihoods that are

systematically skewed with a long tail away from the peak of the prior. If a symmetric loss

function is also employed (selecting the mean rather than the maximum of the posterior: L2

loss), estimates will be systematically biased away from the prior, relative to the peak of the

likelihood ('likelihood repulsion'), see 'Modelling' section for more detail. Recently, Bays (29)

has proposed a Bayesian model of the SWI that incorporates efficient coding and a prior

reflecting a positive association between size (volume) and weight. However, Bays' model has

only been tested on Peters and colleagues (6) dataset.

Other authors suggest that we are unable to fully separate density and weight (34), or

proactively combine them (11). Under these models, denser objects are systematically

perceived as heavier. Indeed, weight and density estimation appear to be closely related:

perceived weight difference increases with increasing density difference (34); objects of greater

weight are reported as denser (35), and weight discrimination is improved when density varies

in the same direction as weight (36). Within typical SWI stimuli, weight is kept constant such

that density is negatively correlated with volume. Thus, size contributes to the illusion because

it is used to infer density (9,11,37).

Wolf and colleagues (10,11) proposed a cue-integration model in which perceived

weight is a weighted average of two heaviness estimates, one from the object's felt weight and

one derived from the object's inferred density. Because denser objects tend to be heavier, the

authors suggest that it is adaptive / Bayesian to combine estimates of weight and density when

estimating weight, to obtain more reliable (less noisy) weight estimates. It is not clear, however,

why we should combine weight and density information in this way, when volume information is

available to condition the density-weight relationship. Not only is density a poor cue to weight

(when not conditioned on size) but the two estimates will also have correlated noise (thus

negating some benefits of integration).

It is well known that the SWI magnitude increases with volume (density) difference

(6,8,34,38,39) and this relationship is predicted by all of the above models. SWI demonstrations

tend to use fairly large, obvious differences in object volume, in order to obtain a reliable

illusion. However, under small conflicts between size-based expectations of weight and direct

sensory evidence, these estimates might be integrated; we know that integration occurs for

small conflicts but breaks down under conditions of large conflict (40,41) . If integration (the

opposite of the SWI) occurs under small conflicts, while the SWI only occurs with large

conflicts, this would be broadly in line with the idea that the SWI is driven by violated weight

expectations. Integration under small conflicts would be consistent with Peters and

colleagues' model (6), with integration driven by their equal density sub-prior. However,

integration under small density differences would be at odds with models in which weight and

density are confounded / integrated in general (10,11). Furthermore, it is unclear whether

humans conflate weight and density in all situations (such as when a smaller object is slightly

less dense), or only in the specific and non-accidental conditions of SWI stimuli, when weight is

held constant and volume varies.

Here we asked: can we develop a generalised model of weight perception for varied

combinations of size, weight and density? To this end we measured perceived weight for pairs

of cubes from a large set in which volume, and / or weight, and / or density were varied in small

steps. Size and density were uncorrelated in the full set to prevent learning of unusual size -

weight associations. To preview the main findings: 1) The SWI occurred even at very small size

(density) differences, and thus does not rely on large conflicts. 2) Perceived weight was

positively correlated with the true weight of both the judged and other stimulus, in addition to

the judged object's density. 3) The effect of both objects' densities on perceived weight was

strongly modulated by the weight difference between the two objects: three times larger at

small / zero weight differences than at large weight differences. In fact, the other object's

density had an effect only when the two objects had similar weight, when it was negatively

associated with the judged object's perceived weight. Thus, the perceptual system does

conflate weight and density, but this effect is much larger in the specific and peculiar

conditions of typical studies of the SWI. We provide a model (R^2 = .98) that captures biases in

perceived weight in the SWI and across the full stimulus set. In addition, we compare this

model to existing models of the SWI, and to a novel efficient-coding Bayesian model of the SWI.

2. Materials and Methods

2.1 Participants

Thirty naïve participants (age M 21.4, SD 5.6 years) from the University of Southampton Psychology student community took part in the experiment; undergraduate students were compensated for their participation with course credits. All reported normal or corrected-to-normal eyesight and no upper limb impairment. The experiment was approved by the University of Southampton Psychology Ethics Committee; all participants gave prior informed consent.

2.2 Stimuli

Stimuli were 15 plastic cubes (3D-printed in black polylactic acid (Figure 2A–B). Each cube was filled with sand, polystyrene pellets, and / or brass to create the desired volume, weight and density values for the different stimulus sets, depicted in Figure 2A. The centre of mass of the cubes remained at the geometric centre and the filling did not move inside the cubes during lifting. A panel concealed at the bottom kept the appearance homogeneous. Weight was measured with a 0.1 g resolution.

Figure 2: Stimulus set. (A) Full stimulus set, with 500 cm^3 stimulus for scale on the left. (B) Stimulus volume as a function of weight. (C) Stimulus density as a function of weight.

The complete stimulus set included three subsets (Figure 2B–C), corresponding to different volume / weight relationships: 1) Equal density (red): weight increased linearly with volume (as for 'normal' same-material objects); 2) Equal weight (green): density decreased with increasing size (typical SWI stimuli); 3) Increasing density with increasing size (blue): both weight and density increased with size. The middle cube (volume, weight, density: 750 cm^3, 375 g, 0.5 g / cm^3) was common to all three subsets. Volumes within each set increased in 100 cm^3 steps, with the addition of 50 cm^3 steps below and above the middle stimulus in the fixed

density set. Minimum (maximum) volume, weight, density differences across any stimulus pair

were: 50 (250) cm^3, 25 (200) g, 0.02 (0.18) g / cm^3, respectively. Density ratios for stimulus pairs

were in the range 0.47–0.97 (less dense / denser stimulus), 0.58–0.89 g / cm^3 for equal weight

(SWI) stimulus pairs. This contrasts with previous work, where smaller ratios (bigger

differences) in density have been explored (largest ratios for classic SWI pairs: 0.67 (42), 0.64

(43), 0.51 (44); a larger ratio of 0.87 was tested by Wolf and colleagues (11) using same-volume

pairs).

2.3 Task and trials

Stimuli were presented in pairs. In each trial, participants simultaneously grasped and

lifted both cubes in full visuo-haptic conditions, and verbally reported both objects' weights

using magnitude estimation (as in Figure 1A). This paired presentation (common in SWI

experiments) allowed us to model the perceived weight of each object in addition to the pair's

perceived weight difference or weight ratio. Experimental trials included all possible stimulus

pairs (same- and different-subset pairs), excluding pairs of identical stimuli. This gave 105

unique pairs and 133 pairs in total (trials including the middle / common stimulus were

repeated due to a coding error). Each stimulus pair was presented twice, with left and right

position switched between the two trials (266 trials in total). Trials were randomised for each

participant.

2.4 Setup

The participant and the experimenter sat at opposite sides of a table covered in thick

felt to muffle auditory cues to the objects' weights. Two numbers (1, 2) marked the positions of

the stimuli on the table (centre-to-centre distance between stimuli, centre of stimuli to table

edge: both 20 cm) and served to remind participants the order in which to report their

estimates. A frame-mounted black curtain was used to control stimulus presentation, such

that the participant only ever saw the two stimuli of the current trial. A desktop computer

running MATLAB (The MathWorks Inc., 2020) was used to randomise stimulus presentation and

record participant responses.

2.5 Procedure

The experiment was completed in two identical laboratory sessions on separate days,

each lasting 60–90 minutes. At the beginning of each session, participants were presented with

the middle cube (750 cm^3, 375 g) as the reference stimulus, and told that this had a weight of

'50' (arbitrary units). The reference was presented again every 20 trials. Participants were

instructed to look at the stimuli while lifting. Exploration mode was not restricted: participants

were free to grasp and lift the cubes as they thought best; they were not restricted to use index

and thumb only. This was done to provide the most reliable size estimates (11). Trials were not

timed, but participants were encouraged to give an answer if they waited more than 30 seconds

(approx.).

3. Results

Participants' magnitude estimates were transformed into grams and then normalised to

remove individual differences in how the response scale was used: Each participant's

estimates were multiplied by the ratio of their individual grand mean across stimuli to the grand

mean across all participants (thus, after normalisation, each observer had the same mean

response). Outliers per condition were then removed iteratively: for each object within each

stimulus pair, values more than 5 SDs, then 4 SDs, and 3.5 SDs from the mean across

observers were removed (65 datapoints in total, or 0.4%). Data were re-normalised after each

iteration. All further analyses were conducted on the resulting dataset.

Weight estimates for within-subset pairs (i.e., both objects from the same subset) are

shown in Figure 3, alongside model fits (see below, Equations 1–2). Data have been collapsed

across repetitions within participants and *SE* bars show inter-observer variation. We present

the perceived weight of the judged object (i.e., the object within each pair whose weight is being

reported) in terms of its own properties (weight / density) and the properties (weight / density) of

the other object in the pair. Each object in turn was considered the judged object.

For equal density, i.e. 'normal' object pairs (Figure 3A–B), weight estimates were highly

correlated with the true weight of the judged object, as expected. However, they were also

positively correlated with the weight of the other object (denoted by symbol size in Figure 3A;

see the vertical spread corresponding to weight of other object); note that this is an averaging,

rather than a contrast effect. Perceived weight difference between the two objects was linearly

related to stimulus weight difference (Figure 3B; data are symmetrical in all difference plots (B,

D, F) as each object in each stimulus pair serves as 'judged').

Figure 3 Perceived weight (A, C, E) and perceived weight difference (B, D, F) for within-subset
stimulus pairs, averaged across participants, error bars show ± 1 SEM. Solid lines / open
symbols show the model predictions (described below, Equations 1 and 2). The dashed lines
indicate veridical perception. (A–B) Equal density pairs: (A) Weight estimates as a function of
stimulus weight; symbol size indicates the weight of the other stimulus in the pair. (B) Data re-
expressed as perceived weight difference. (C, D) Equal weight (SWI) pairs: (C) Weight estimates
as a function of stimulus density; symbol size indicates the density of the other stimulus in the
pair. (D) Perceived weight difference as a function of density difference; symbol size indicates
the average stimulus density. (E, F): Increasing density pairs: (E) Symbol size indicates the other
stimulus' weight; (F) Symbol size indicates the average density.

Weight estimates for the equal weight (SWI) pairs are shown as a function of the judged

object's stimulus density (x-axis) and the other object's density (symbol size, Figure 3C). In line

with the size-weight illusion, perceived weight was positively correlated with object density:

denser (and smaller) stimuli were perceived as heavier (overall positive slope of the data).

Perceived weight was also negatively correlated with the other object's density, the judged

object appearing heavier when judged next to less dense (larger) objects (see the vertical

spread of data points in 3C). The data are re-expressed as perceived weight difference as a

function of density difference in Figure 3D. The magnitude of the SWI (i.e. deviation from

veridical perception) increases with density difference, as expected, but still occurs with small

density (or volume) differences. Thus, we find no evidence for integration of expected and

sensory weight estimates under small conflicts.

Lastly, in the increasing density pairs (Figure 3E), weight estimates were positively

correlated with the object's weight, but also correlated with the other object's weight (as in the

equal-density pairs; see vertical spread of data in 3E). Figure 3F shows that perceived weight

difference increases linearly with true weight difference. In this set, weight (or weight

difference) is correlated with density (or density difference); the effect of density contributes to

the increased slope of the data, relative to veridical perception (identity line).

Figure 4A summarises results for all trials, including across-subset pairs (i.e. objects

within a pair are from different stimulus subsets), in a format that illustrates the variable

influence of density on perceived weight. For every stimulus pair (represented by a dot),

perceived weight difference (y-axis) is presented as a function of stimulus weight difference (x-

axis) and stimulus density difference (colour). Lines show the model fit (see Equation 2 below).

The overall slope of the data shows the strong relationship between true and perceived weight

difference. The vertical spread of the different colours shows the effect of density difference on

perceived weight difference. The two are not independent: there is a central 'bulge', where the

effect of density is larger (more vertical spread). In other words, the effect of density (or density

difference) on perceived weight (or perceived weight difference) is strongly modulated by

stimulus weight difference, being much greater for small / zero weight differences. Figure 4B

summarises the influence of the two objects' densities on perceived weight, as captured by the

model's density coefficients (see below). The conflation of weight and density is dramatically

increased under the conditions of typical size-weight stimuli, where weight difference is zero.

Figure 4 (A) Perceived weight difference as a function of weight difference and density
difference in the full dataset. Markers show the data, solid lines show the model predictions
(see Equation 2) at equally spaced density differences. The vertical spread of the data with
density difference (marker colours) shows the effect of density in typical SWI stimuli (i.e., equal
weight: vertical dotted line) and across the full set. The dashed diagonal line shows veridical
perception. (B) The influence of density (i.e., density coefficients) as a function of weight
difference in the model.

4. Modelling

4.1 Model of perceived weight.

Our model captures perceived weight $(\hat{w}_1)$ for all stimulus combinations, as a function

of the weight and density of the judged (w_1, d_1) and other object (w_2, d_2). It is presented as a

schematic in Figure 5A, which shows the 5 components of the model, and described formally

by Equation 1; perceived weight is positively correlated with both objects' weights, and the

judged object's density. When the two objects' weights are similar (as in the SWI), the two

objects' densities have an additional influence (dictated by a Gaussian over weight difference).

The model (Equation 1) explains 98% of the variance in the full dataset:

$$\hat{w}_1 = C + \beta_1 w_1 + \beta_2 d_1 + \beta_3 w_2 + (\beta_4 d_1 + \beta_5 d_2)N(|w_1 - w_2|, 0, \sigma^2) \qquad (1)$$

Figure 5B shows how the model fit improves with each additional component (i.e.,

components 1-5, as in Figure 5A) of the model.

327
Figure 5 Model of perceived weight. (A) Schematic of the model's 5 components. The perceived
weight of an object is positively correlated with both objects' weights (components 1 and 3),
and the judged object's density (component 2). When the objects' weights are similar (captured
by a Gaussian over their difference) the two objects' densities have an additional influence
(components 4 and 5). The importance of each component is conveyed by the thickness of the
corresponding arrow. (B) Variance in perceived weight explained by the addition of each
component in the model (bars) and corresponding SSR (magenta lines and markers) to the
constant (C). The order of components follows the maximum increase in R^2 with each addition
(see SI, Table S1). X = $N(|w_1 - w_2|, 0, \sigma^2)$.
337
Accordingly, the perceived weight difference ($\hat{w}_1$ - $\hat{w}_2$) between two stimuli can be

predicted via the following equation:

$$\hat{w}_{diff} = (\beta_1 - \beta_3)(w_1 - w_2) + \beta_2(d_1 - d_2) + (\beta_4 - \beta_5)(d_1 - d_2) N(|w_1 - w_2|, 0, \sigma^2) \quad (2)$$

Perceived weight is positively correlated with the object's weight (β_1 = 1.23) and its

density (β_2 = 126.43), and with the other object's weight (β_3 = 0.13). However, when the two

objects have similar weight, the effect of the object's density is magnified, and the other

object's density also has an effect. This interaction between weight difference and the effect of

density is captured by a Gaussian centred on a weight difference of 0. The density of the judged

object positively influences its perceived weight (β_4 = 27814), whereas the other object's

density has a negative influence (β_5 = -17919). Sigma for the Gaussian component is 76.44.

Figure 4B shows the influence of both objects' densities as a function of weight difference: the

combined influence of the two densities on perceived weight is approximately three times

larger for zero weight differences than for large weight differences. All analyses were done in

MATLAB and optimal coefficients (minimising the summed squared residuals, SSR) were

identified via gradient descent. Alternative candidate models are presented in Supplementary

Table 1 alongside model comparison statistics. Models including the objects' volumes (v_1, v_2)

instead of density were also tested, see Supplementary Table 2. To prevent over-fitting, we used

leave-one-out cross-validation on the normalised dataset, leaving out each participant in turn.

This method ensures that every component in the model improves its ability to predict data

from new observers.

Whilst our model of perceived weight provides an excellent quantitative account of

perceived weight, it does not explain *why* our sensory system computes perceived weight in

this way. Below we explore whether a Bayesian model incorporating efficient coding might

provide a more theoretically appealing account of biases in weight perception. We then

compare our models to those previously proposed to explain biases in weight perception.

4.2 Efficient coding Bayesian ratio model

Traditionally, Bayesian models have been employed to account for perceptual biases

towards a prior (i.e. towards the most frequent or probable stimuli). Examples in the literature

include biases for light-from above (45), isotropy assumptions when interpreting foreshortening

(40), global convexity of surfaces (46), and 'slow and smooth' motion (47). However, as

described by Wei and Stocker (30,31), Bayesian models incorporating efficient coding of

stimulus features can also predict repulsion biases, i.e. biases *away* from the prior. Given that

the SWI (and not integration) is observed even for very small density differences, we asked

whether such a model might provide a quantitative, Bayesian model of the SWI as a 'repulsion'

effect, where perception is biased away from a prior for equal density. Wei and Stocker (30,31)

show that the expected perceptual bias $b(\theta_0)$ for a stimulus value θ_0 from a stimulus

distribution θ can be predicted from the prior, $p(\theta)$ (in fact from the slope of its squared

reciprocal), as in Equation 3:

$$b(\theta_0) \approx C(1/p(\theta_0)^2)' \qquad\qquad (3)$$

We present a simple model that includes a prior over the relationship between two

objects' weights and volumes: i.e. the two dimensions of log weight ratio and log volume ratio,

as depicted in Figure 1. Such a prior, if aligned with the log weight ratio = log volume ratio unity

line (solid red line in Figure 1B), represents the assumption that the two objects have equal

density. Perceptual biases *away* from this prior would be consistent with the 'contrast' or

'repulsion' explanation of the SWI that has prevailed for decades (7,18). For pairs of objects,

considering ratios of weight, volume and / or density (rather than their differences) is a natural

choice: any two of these ratios define the third, as $(w_1/w_2) / (v_1/v_2) = d_1/d_2$. Taking the log ratio

gives a useful symmetry: $\log(w_1/w_2) = -\log(w_2/w_1)$; allowing us to predict the same magnitude of

bias, irrespective of which object in a pair is designated as object 1. Our prior is defined by a

mixture of three bi-variate Gaussians centred on [0, 0], i.e. equal weight and volume (Figure 6A).

This mixture-model allowed us to explore variations in the shape of the prior; there is no reason

to assume that it should be a perfect bivariate Gaussian. Model parameters and model

comparisons given priors of different complexity are provided in SI, Table S3.

Figure 6 Efficient coding Bayesian ratio model. (A) Bi-variate prior over log weight and log
volume ratio, optimised to minimise errors in predicted log weight ratio. Red asterisk and
dashed black line indicate an example stimulus condition and corresponding slice through the
prior that are shown in part B. The solid red line is the identity line (i.e. equal density), and the
dashed red line shows the orientation of the fitted prior (plotted over the stimulus range) as
defined by its peak values. (B) Likelihood repulsion predicted by Wei and Stocker (30) for one
stimulus condition with equal weight, unequal volume stimuli. Schematic of a median
(unbiased / noise-free) observer likelihood (red) and posterior (blue) for an example stimulus
pair (red asterisk in A), given the 1D slice through prior (solid black line). Efficient coding of
stimulus properties results in skewed likelihoods with a long tail away from the peak of the
prior. The posterior is also skewed. When a symmetric loss function is chosen, e.g. the mean of
the posterior (L2, blue dashed line), a bias away from the prior is predicted, i.e. the posterior
estimate is shifted away from the prior peak, relative to the peak of the likelihood.
Measurement noise (expected to be skewed in stimulus space) further increases the predicted
repulsive bias (30). (C-D) Biases in perceived log weight ratio. Contour lines show the prior;
black dots show the stimulus pairs; arrows show the magnitude and direction of the bias for
each pair, with SWI conditions in green; the solid red line shows equal density. (C) Empirical
biases; (D) predicted biases.

Similarly to Bays (29), we employ the simplifying assumption that perceptual estimates

of volume ratio are relatively precise, and thus not significantly affected by the prior. Indeed,

there is no clear evidence that objects with unexpected weights (e.g. typical SWI objects) cause

biases in *volume* perception (i.e., a 'weight-size illusion' (48,49)). Thus, we estimated the

expected bias in perceived log weight ratio by taking a slice through the prior at the true value of

log volume ratio (e.g., vertical dashed line in Figure 6A) and finding the slope of this 1D prior

with respect to log weight ratio (Figure 6B). If volume ratio judgements were also available, one

could drop the simplifying assumption, and simultaneously fit biases in log weight ratio and log

volume ratio. The resultant fitted prior would have a slightly different shape.

We fitted the model via gradient descent to optimally predict biases in perceived log

weight ratio, averaged across participants, for all unique stimulus pairs (N = 105). (For this

model, we calculated each participant's ratio of weight estimates on each trial). Figure 6A

shows the best fitting prior, whose orientation is close to an equal density prior (solid red,

identity line), but is in fact rotated slightly towards larger objects being heavier and slightly less

dense (dashed red line) in keeping with the object statistics reported by Peters and colleagues

(5). The empirical biases in perceived log weight ratio are shown in (C) and those predicted by

the fitted prior are shown in (D). The model accounts for perceived log weight ratio well, with 12

parameters (R^2 = .988; see SI S2. Efficient-coding Bayesian ratio model; Table S3). However,

goodness-of-fit is poorer than our descriptive model (Equation 1, 7 parameters), when

parameters are optimised to minimise the SSR in perceived log weight ratio (R^2 = .991, see also

cross-validation errors, SI Table S1). Nonetheless, this efficient-coding Bayesian model

provides a theoretically appealing account of the SWI; biases in perceived log weight ratio are

minimal close to the $d_1=d_2$ identity line (solid red lines in Figure 6 C-D) and increase with log

density ratio, broadly in line with repulsion away from an equal-density prior. Indeed, biases in

log weight ratio correlate quite strongly with log density ratio (R^2 = .691).

Tempting as it is to break out the champagne and toast our Bayesian brains, this may be

premature. Closer inspection yields some difficulties with this model. First, the model is

agnostic regarding how the bias in log weight ratio is split across the two objects. A reasonable

assumption is that the prior has equal and opposite effects on the perceived weights of the two

objects. However, this is not observed (see Figure S1); there is only a weak correlation (R^2 =

.283) between biases in estimates of log weight ratio and biases in estimates of $\log w_1$ and log

w_2. In fact, biases in estimates of $\log w_1$ and $\log w_2$ are largely independent of the two objects'

relative densities (R^2 = .215), in contrast to what we would expect given repulsion from an equal

density prior. The pattern of bias in log weight ratio is fairly well approximated by a model that

predicts each object's perceived weight from its weight and density, that is, independently of

any interaction between the two objects (R^2, biases in log weight ratio =. 847 see SI, Table S1).

This suggests that, although empirical biases are consistent with repulsion from an equal-

density prior (Figure 6B), this might not be the underlying mechanism. It is worth noting that any

pattern of bias can be explained with a complex enough prior. Observed and predicted biases in

judgements of $\log w_1$ and $\log w_2$ are shown in supplementary Figure S1.

Second, a key feature of Wei & Stocker (30,31) is that the model gives predictions for

both biases *and* reliability (discrimination thresholds or estimate variability). According to the

efficient coding model, more frequently observed stimuli are represented more precisely, such

that discrimination thresholds are proportional to the reciprocal of the prior. At the individual

observer level, our data provide poor estimates of discrimination (only two estimates per pair).

However, at the group level, a systematic pattern is clear (see SI, Figure S2). Empirical response

variance is well predicted by the absolute log weight ratio (R^2 = .532); observers' judgements of

relative weight are most reliable when the two objects' weights are similar. Log volume ratio

adds little explanatory power (R^2 = .557, for regression model predicting response variance from

log weight and log volume ratios). Predictions from the efficient coding Bayesian model follow

the opposite pattern: modelled response variance depends primarily on log volume ratio (R^2 =

.765) and far less on log weight ratio (R^2 = .86, for regression model including both log volume

and log weight ratios, see SI, Figure S1). In fact, the empirical pattern of response variance is

better predicted by log weight ratio than by the modelled pattern of response variance (R^2 = .532

vs R^2 = .444). For a variety of perceptual tasks, Wei & Stocker's approach explains both the

pattern of bias *and* the pattern of response variance (or discrimination thresholds) with a single

fitted prior (31). Unfortunately, that is not the case for perceived weight, and is therefore

unlikely to represent the underlying mechanism of the SWI. (We also considered, and rejected,

Bayesian models of perceptual bias involving a 2D prior over *differences* in weight and

differences in volume or density, see SI, S2. Efficient-coding Bayesian ratio model).

4.3 Other models

Here we explore how well existing models of weight perception by Bays (29), Peters and

colleagues (6), and Wolf and colleagues (11) (see Introduction) explain our data.

Bays (29) proposed a Bayesian model of the SWI, which, as above, predicts biases away

from the prior, given efficient coding and an L2 loss function (30). However, rather than

considering object pairs, Bays' model employs a bivariate Gaussian prior over the perceived

weight and volume of single objects, i.e., a positive association between weight and size.

Supplementary Figure S3 shows the model's fit to our data. We extended the model to

incorporate a mixture of Gaussians prior (see SI, Table S4 for model parameters and model

comparisons). Whilst the model captures the mean bias for each stimulus reasonably well, it

cannot capture the way in which perceived weight is modulated by the other object in the pair,

nor is it consistent with the pattern of response variance (see SI Figure S2).

As described in the Introduction, Peters and colleagues (6) assume a prior over weight

and density ratio that is comprised of three sub-priors, representing three different classes of

weight-density relationships (here re-expressed as weight-volume relationships, see Figure

1C). Figure 7 compares the empirical biases in perceived log weight ratio (A) to the predicted

biases from the best-fitting model parameters (B). Sub-priors (represented by dashed lines) are

encoded as bivariate Gaussians (See SI, Table S5 for model parameters and model

comparisons). Unfortunately, the model cannot account for the set of biases in perceived log

weight ratio across all stimulus pairs (R^2 = 0.413). It predicts a negative influence of log density

ratio for some stimulus pairs; the predicted effect depends on where the stimuli lie relative to

the three sub-priors. For example, in pairs of equal-volume (different weight, density) stimuli,

log weight ratio is predicted to be underestimated as log density ratio, log weight ratio increase,

whereas empirically it is overestimated.

Figure 7: Peters and colleagues' (6) model of perceived weight. (A) Empirical biases in perceived
log weight ratio. Dotted lines show the optimal three sub-priors from Peters and colleagues'
model when fitted to our data. Black dots show the stimulus pairs; arrows show the magnitude
and direction of the bias, with SWI conditions in green. (B) Predicted biases according to Peters
and colleagues' model.

Finally, we evaluated Wolf and colleagues' (11) model fit to our data. In this model,

perceived weight is a weighted average of weight and density estimates, weighted according to

their relative reliabilities. Estimates are modelled as power functions of the corresponding

physical feature. As we didn't manipulate the reliability of size information, the model reduces

to:

$$\hat{w} = \beta_1 w^x + \beta_2 d^y \qquad (4)$$

The model provides a good account of perceived weight (R^2 = .96). Figure S5 (SI), shows

the fit to our data, for comparison with Figure 3; for model parameters and model comparisons

see SI Table S6). However, like Bays' (29), it does not capture the contextual effects of the other

object, including the observation the influence of density is strongly modulated by weight

difference (as shown in Figure 4A).

We note that Wolf and colleagues model perceived weight and density as power

functions of their physical counterparts. However, additional exponent parameters for weight

or density were not statistically supported by our data. Wolf and colleagues had a 5:1 ratio of

heaviest to lightest stimuli, compared to our 3.3:1 ratio. It is possible that, given an increased

range of stimulus weights / densities, our descriptive model would need to similarly include

power functions of weight and / or density, in tandem an intercept of 0.

5. Discussion

We investigated perceived weight in a set of stimuli which varied in weight and / or

volume, and / or density in small steps. We report several novel findings: 1) The SWI occurs

even at very small density (size) differences; 2) Across all stimuli, an object's weight and density

are confounded / integrated to estimate weight; 3) Perceived weight is affected by the weight of

the other object in a pair but is largely invariant to the other object's density. 4) However, the

influences of both objects' densities on perceived weight are strongly modulated by weight

difference. The negative, or 'contrast' effect of the other object's density only appears for pairs

of similar weight.

Our descriptive model captures the above effects, providing an excellent account of

perceived weight (R^2 = .98). It captures the SWI as well as biases in perceived weight across a

range of variations in pairings of stimulus weight and density. Additionally, we evaluated a novel

efficient-coding Bayesian model of perceived weight ratio, alongside previously proposed

models of the SWI. Here we discuss our findings within the existing literature.

The magnitude of the SWI increased, as expected, with increasing density (size)

difference (6,34,38,39). We extended this, by measuring the SWI with smaller density

differences than those previously tested. We hypothesised that under small conflicts, the SWI

might reverse (i.e., to show integration: smaller stimuli would be perceived as lighter). However,

the SWI was clearly present for all equal-weight pairs, at odds with the idea that the SWI is

driven by a (substantial) conflict between expected and perceived weight.

Our results are inconsistent with Peters and colleagues' (6) Bayesian mixture model.

Also, at a conceptual level, the subdivision of the continuous world of size and volume/density

into three relatively narrow categories is difficult to reconcile with environmental statistics. The

best fitting parameters for our data assign a prior probability of near 0 to equal-density objects

(R_1, see Figure 7, and SI, Table S5). The other two sub-priors encapsulate the expectation that

smaller objects are either much heavier and denser (R_2) or much lighter and less dense than

larger ones (R_3), both at odds with reported natural statistics for everyday liftable objects (5).

The efficient-coding Bayesian ratio model provides a theoretically appealing account of

the SWI, predicting a repulsion away from a plausible equal-density prior. However, neither our

efficient-coding Bayesian ratio model, nor that recently proposed by Bays (29) satisfactorily

account for our pattern of data. Whilst our efficient-coding model provides a reasonable fit to

the empirical biases in perceived log weight ratio, it fails to account for the observed pattern of

response variance, and does not explain biases in perceived weight for individual objects. As

noted above, Bays' model (29) does not account for contextual bias effects found in our data

(i.e., influence of the other object's weight and, in certain conditions, its density), and neither is

it consistent with empirical patterns of response variance.

So, is the SWI Bayesian after all? We don't think so; the SWI does not appear to be

caused by repulsion from an equal-density prior. It is reasonable to assume that the perceptual

system does expect same-material objects to be same-density. Indeed, the sensorimotor

system may rely on an object's size, together with density expectations for classes of objects /

materials to infer the weight of a stimulus prior to lifting (14). Such expectations presumably

drive the initial (erroneous) forces used when lifting SWI objects (50). However, the illusion

persists after forces quickly adapt to the true weights, and the SWI is just as large when same-

density cues are reduced, i.e., when the colour or surface material of the two stimuli differs

(51,52). Thus, while we may employ an equal-density prior, this does not seem to underlie the

perceptual SWI.

In general, perceived weight was modulated by density; denser objects are perceived as

heavier than less dense objects, consistent with previous literature (11). In line with Wolf and

colleagues (10,11) weight does appear to be routinely confounded / integrated with density

information. However, our data reveal that the effect of density is substantially increased when

equal-weight objects are encountered, as in the typical size-weight illusion paradigm. Thus, like

Bays' (29) model, Wolf and colleagues' (11) model fails to capture the contextual effects of the

other object. Rather, these models predict biases in perceived weight that occur in single-

object judgements (see, e.g., (53)). Obviously, object perception occurs in various contexts, not

only object pairs. Our model could be generalised to judgements of single objects, by replacing

the other object's properties with a weighted combination of previously encountered stimuli.

Wolf and colleagues (11) also showed that when the reliability of (visual or haptic) size

information is reduced, density has less effect on the final heaviness estimate (11). In the

limiting case, when size information is completely absent (objects lifted via strings / a handle,

and no vision), the illusion disappears (11,39,54). Our model of weight perception does not

encapsulate the additional effect of varying the reliability of size estimation; this would be

accommodated by reducing the coefficients associated with density to reflect the availability of

visual-haptic size cues (β_2, β_4, β_5, Equation 1).

6. Conclusions

Our study situates the SWI in the broader context of weight perception under varied conditions (combinations of volume, weight, and density) allowing us to evaluate different models of weight perception, including our own.

Our explorations reveal that, whilst the SWI magnitude increases with size / density difference, it occurs even for small density differences. Second, we showed that typical SWI experiments have inflated the apparent role of density on weight perception more generally. The effect of density on perceived weight is substantially smaller when two objects of different weights are encountered. Our simple model provides an excellent account of perceived weight under various combinations of size, weight, and density. However, we still don't know why weight is estimated in this way. Why do we conflate weight and density, and why is this effect magnified when weight differences are small?

We explored different avenues to reconcile the SWI within the Bayesian framework, given the appeal of finding a model that provides a more explanatory account of weight perception in the SWI and beyond. However, none of the Bayesian models we tested, including efficient-coding models that do predict biases away from a prior, could successfully account for our data.

Acknowledgments

We thank Megan A. K. Peters for helpful discussions about modelling.

References

1. Charpentier A. Analyse experimentale de quelques elements de la sensation de poids [Experimental study of some aspects of weight perception]. Archieves de Physiologie Normales et Pathologiques [Internet]. 1891 [cited 2021 Sep 13];3:122–35. Available from: http://ci.nii.ac.jp/naid/10018698834/en/

2. Flournoy Th. Illusions de poids. Annee Psychol [Internet]. 1894;1(1):198–208. Available from: https://www.persee.fr/doc/psy_0003-5033_1894_num_1_1_1051

3. Buckingham G. Getting a grip on heaviness perception: A review of weight illusions and their probable causes. Exp Brain Res. 2014;232(6):1623–9.

4. Flanagan JR, Beltzner MA. Independence of perceptual and sensorimotor predictions in the size- weight illusion. Nat Neurosci. 2000;3(7):737–41.

5. Peters MAK, Balzer J, Shams L. Smaller = denser, and the brain knows it: Natural statistics of object density shape weight expectations. PLoS One. 2015 Mar 13;10(3).

6. Peters MAK, Ma WJ, Shams L. The Size-Weight Illusion is not anti-Bayesian after all: A unifying Bayesian account. PeerJ. 2016;2016(6).

7. Dijker AJM. The role of expectancies in the size-weight illusion: A review of theoretical and empirical arguments and a new explanation. Psychon Bull Rev. 2014 Nov 15;21(6):1404–14.

8. Saccone EJ. A meta-analysis of the size-weight and material-weight illusions. 2019;

9. Thouless RH. Individual differences in phenomenal regression. British Journal of Psychology. 1932;22:216–241.

10. Wolf C, Drewing K. The size-weight illusion comes along with improved weight discrimination. PLoS One. 2020 Jul 1;15(7 July).

11. Wolf C, Bergmann Tiest WM, Drewing K. A mass-density model can account for the size-weight illusion. PLoS One. 2018 Feb 1;13(2).

12. Amazeen EL. The Effects of Volume on Perceived Heaviness by Dynamic Touch: With and Without Vision. Ecological Psychology. 1997;9(4):245–63.

13. Amazeen EL, Turvey MT. Weight perception and the haptic size–weight illusion are functions of the inertia tensor. Vol. 22, Journal of Experimental Psychology: Human Perception and Performance. US: American Psychological Association; 1996. p. 213–32.

14. Cole KJ. Lifting a familiar object: Visual size analysis, not memory for object weight, scales lift force. Exp Brain Res. 2008 Jul;188(4):551–7.

15. Gordon AM, Forssberg H, Johansson RS, Westling G. Visual size cues in the programming of manipulative forces during precision grip. Vol. 83, Exp Brain Res. 1991.

16. Gordon AM, Forssberg H, Johansson RS, Westling G. Integration of sensory information during the programming of precision grip: comments on the contributions of size cues. Vol. 85, Exp Brain Res. 1991.

17. Davis CM, Roberts W. Lifting movements in the size-weight illusion. Vol. 20, Perception & Psychophysics. 1976.

18. Buckingham G. Getting a grip on heaviness perception: A review of weight illusions and their probable causes. Vol. 232, Experimental Brain Research. Springer Verlag; 2014. p. 1623–9.

19. Grandy MS, Westwood DA. Opposite perceptual and sensorimotor responses to a size-weight illusion. J Neurophysiol. 2006 Jun;95(6):3887–92.

20. Mon-Williams M, Murray AH. The size of the visual size cue used for programming manipulative forces during precision grip. Exp Brain Res. 2000;135(3):405–10.

21. Brayanov JB, Smith MA. Bayesian and "Anti-Bayesian" biases in sensory integration for action and perception in the size-weight illusion. J Neurophysiol. 2010 Mar;103(3):1518–31.

655   22.   Ernst MO. Perceptual Learning: Inverting the Size–Weight Illusion. Current Biology
656   [Internet]. 2009 Jan 13 [cited 2019 Dec 2];19(1):R23–5. Available from:
657   https://www.sciencedirect.com/science/article/pii/S0960982208014085?via%3Dihub
658   23.   Adams WJ, Graf EW, Ernst MO. Experience can change the "light-from-above" prior. Nat
659   Neurosci. 2004;7(10):1057–8.
660   24.   Adams WJ, Mamassian P. Bayesian combination of ambiguous shape cues. Vol. 4,
661   Journal of Vision. Adams, Wendy J.: Department of Psychology, University of
662   Southampton, Southampton, United Kingdom, S017 1BJ, w.adams@soton.ac.uk: Assn
663   for Research in Vision & Ophthalmology (ARVO); 2004. p. 921–9.
664   25.   Ernst MO. A Bayesian View on Multimodal Cue Integration. Human body perception from
665   the inside out: Advances in visual cognition. New York, NY, US: Oxford University Press;
666   2006. p. 105–31. (Advances in visual cognition.).
667   26.   Ernst MO, Bülthoff HH. Merging the senses into a robust percept. Trends Cogn Sci.
668   2004;8(4):162–9.
669   27.   Kersten D, Mamassian P, Yuille A. Object Perception as Bayesian Inference. Annu Rev
670   Psychol. 2004;55(1):271–304.
671   28.   Knill DC, Richards W. Perception as Bayesian Inference [Internet]. Cambridge:
672   Cambridge University Press; 1996. Available from:
673   https://www.cambridge.org/core/product/0442F577F5E4CD874FA6819978574C8F
674   29.   Bays PM. Weight illusions explained by efficient coding based on correlated natural
675   statistics. bioRxiv [Internet]. 2024 Jan 1;2024.02.09.579650. Available from:
676   http://biorxiv.org/content/early/2024/05/02/2024.02.09.579650.abstract
677   30.   Wei XX, Stocker AA. A Bayesian observer model constrained by efficient coding can
678   explain "anti-Bayesian" percepts. Nat Neurosci. 2015 Oct 28;18(10):1509–17.
679   31.   Wei XX, Stocker AA. Lawful relation between perceptual bias and discriminability. Proc
680   Natl Acad Sci U S A. 2017 Sep 19;114(38):10244–9.
681   32.   Attneave F. Some informational aspects of visual perception. Vol. 61, Psychological
682   Review. US: American Psychological Association; 1954. p. 183–93.
683   33.   Barlow HB. Possible Principles Underlying the Transformations of Sensory Messages. In:
684   Rosenblith WA, editor. Sensory Communication [Internet]. The MIT Press; 1961. p. 216–
685   34. Available from: https://doi.org/10.7551/mitpress/9780262518420.003.0013
686   34.   Ross J, Di Lollo V. Differences in heaviness in relation to density and weight. Percept
687   Psychophys. 1970;7(3):161–2.
688   35.   Merken L, van Polanen V. Density estimation is influenced more by mass when objects
689   are denser [Internet]. Vol. 12272 LNCS, Lecture Notes in Computer Science (including
690   subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics).
691   Springer International Publishing; 2020. 87–95 p. Available from:
692   http://dx.doi.org/10.1007/978-3-030-58147-3_10
693   36.   Kawai S. Heaviness perception: II. Contributions of object weight, haptic size, and
694   density to the accurate perception of heaviness or lightness. Exp Brain Res.
695   2002;147(1):23–8.
696   37.   Harshfield SP, DeHardt DC. Weight judgment as a function of apparent density of
697   objects. Psychon Sci. 1970;20(6):365–6.
698   38.   Cross D V., Rotkin L. The relation between size and apparent heaviness. Percept
699   Psychophys. 1975;18(2):79–87.
700   39.   Ellis RR, Lederman SJ. The role of haptic versus visual volume cues in the size-weight
701   illusion. Percept Psychophys. 1993;53(3):315–24.
702   40.   Knill DC. Robust cue integration: A Bayesian model and evidence from cue-conflict
703   studies with stereoscopic and figure cues to slant. J Vis. 2007;7(7):1–24.
704   41.   Landy MS, Maloney LT, Johnston EB, Young M. Measurement and modeling of depth cue
705   combination: in defense of weak fusion. Vision Res. 1995;35(3):389–412.

706   42.   Plaisier MA, Smeets JBJ. Object size can influence perceived weight independent of
707   visual estimates of the volume of material. Sci Rep [Internet]. 2015;5(February):1–9.
708   Available from: http://dx.doi.org/10.1038/srep17719
709   43.   Stevens JC, Rubin LL. Psychophysical scales of apparent heaviness and the size-weight
710   illusion. Percept Psychophys. 1970;8(4):225–30.
711   44.   Walker P, Scallon G, Francis B. Cross-Sensory Correspondences: Heaviness is Dark and
712   Low-Pitched. Perception. 2017 Jul 1;46(7):772–92.
713   45.   Adams WJ. A common light-prior for visual search, shape, and reflectance judgments. J
714   Vis. 2007 Aug 31;7(11):1–7.
715   46.   Langer MS, Bülthoff HH. A prior for global convexity in local shape-from-shading.
716   Perception. 2001;30(4):403–10.
717   47.   Weiss Y, Simoncelli EP, Adelson EH. Motion illusions as optimal percepts. Nat Neurosci.
718   2002 Jun 1;5(6):598–604.
719   48.   Hirsiger S, Pickett K, Konczak J. The integration of size and weight cues for perception
720   and action: Evidence for a weight-size illusion. Exp Brain Res. 2012 Nov;223(1):137–47.
721   49.   Smeets JBJ, Vos K, Abbink E, Plaisier M. Size, weight, and expectations. Perception.
722   2022;51(5):344–53.
723   50.   Flanagan JR, Beltzner MA. Independence of perceptual and sensorimotor predictions in
724   the size-weight illusion [Internet]. 2000. Available from: http://neurosci.nature.com
725   51.   Buckingham G, Goodale MA, White JA, Westwood DA. Equal-magnitude size-weight
726   illusions experienced within and between object categories. J Vis. 2016;16(3).
727   52.   Vicovaro M, Burigana L. Contribution of surface material and size to the expected versus
728   the perceived weight of objects. Atten Percept Psychophys [Internet]. 2017;79(1):306–
729   19. Available from: http://dx.doi.org/10.3758/s13414-016-1212-6
730   53.   Ross HE. When is a weight not illusory? Q J Exp Psychol. 1969;21(4):346–55.
731   54.   Plaisier MA, Kuling IA, Brenner E, Smeets JBJ. When Does One Decide How Heavy an
732   Object Feels While Picking It Up? Psychol Sci. 2019 Jun 1;30(6):822–9.