Extinction potential from invasive alien species

Martin Philippe-Lesaffre[1], Ugo Arbieu[1], Alok Bang[2], Morelia Camacho[3], Ross N. Cuthbert[4], Piero Genovesi[5,6], Sabrina Kumschick[6,7], Arman Pili[8], Hanno Seebens[9], Shengyu Wang[1], Guillaume Latombe[10,*]

[1]Université Paris-Saclay, CNRS, AgroParisTech, Ecologie Systématique Evolution, 91190, Gif-sur-Yvette, France
[2]Biology Group, Azim Premji University, Bhopal 462022, India
[3]Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, 04510, Mexico City, Mexico
[4]Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast, BT9 5DL, Belfast, United Kingdom
[5]Institute for Environmental Protection and Research, and Chair IUCN SSC Invasive Species Specialist Group, Via B. Brancati 48, 00144, Rome, Italy
[6]Centre for Invasion Biology, Botany and Zoology Department, Stellenbosch University, South Africa
[7]South African National Biodiversity Institute, Kirstenbosch Research Centre, Cape Town, South Africa
[8]Institute of Biochemistry and Biology, University of Potsdam, 14469 Potsdam, Germany
[9]Department of Animal Ecology & Systematics, Justus-Liebig-University Giessen, Giessen, Germany
[10]Institute of Ecology and Evolution, The University of Edinburgh, King's Buildings, Edinburgh EH9 3FL, United Kingdom

*Corresponding author: glatombe@ed.ac.uk

Abstract

Biological invasions pose significant threats to biodiversity, while impacting ecosystem services, human health, and cultural heritage. Despite these far-ranging effects, their impacts are generally underappreciated by both the public and policymakers, resulting in insufficient management and inadequate conservation outcomes. Recognizing the gap in effective quantitative measurement tools, we introduce the Extinction Potential Metric (EPM) and its derivative, EPM for Unique species (EPM-U; adjusted for phylogenetic uniqueness) to quantify the ecological damage caused by invasive alien species (IAS). These metrics estimate the number of current and projected extinct species within a 50-year horizon under a business-as-usual scenario due to specific IAS.

We applied EPM and EPM-U to assess threats to native terrestrial vertebrates from IAS, examining impacts on 2178 amphibians, 920 birds, 865 reptiles, and 473 mammals. The analysis identified that damage mostly stems from a limited number of IAS, notably two pathogenic fungi affecting amphibians (up to 380 equivalent extinct species) and primarily cats (139 equivalent extinct species) and rats (50 equivalent extinct species) impacting other groups, through mechanisms such as predation, disease, and reduced reproductive success in birds.

The proposed metrics not only provide a standardised measure of ecological impacts but are sufficiently versatile to be tailored for specific spatial and temporal scales or taxonomic groups. Furthermore, EPM could serve as a model for developing unified indicators to monitor global biodiversity targets, such as those defined in the Kunming-Montreal Global Biodiversity Framework (GBF), by assessing the ecological effects of various individual and combined anthropogenic stresses. Also, EPM and EPM-U could support the enforcement of Target 6 of the GBF, by establishing lists of IAS requiring urgent prevention and control.

60   Thus, EPM and EPM-U offer critical tools for improving the management of biological

61   invasions and enhancing global conservation strategies.

64   **Keywords:** Conservation, Extinction, Biological invasion, Ecological impact, Metric,

65   Phylogeny, Red List

Introduction

Biological invasions are a major threat to biodiversity and cause impacts across all aspects of the natural and human world, including reduced ecosystem services, increased human health hazards, and loss of indigenous and cultural practices (Roy et al., 2024). The number of newly introduced alien species has been steadily increasing since the 1800's (Seebens et al., 2017), and the number of established alien species is expected to increase by 36% between 2005 and 2050 under a business-as-usual scenario (Seebens et al., 2021). The subset of these species that have negative impacts, so-called invasive alien species (IAS), have contributed to 60% of known species extinctions, cost hundreds of billions of $US each year (increasing four-fold every decade and reaching $423 billion in 2019), and adversely impact quality of life in 85% of cases (Roy et al., 2024). Despite these figures, the impacts of biological invasions remain underestimated and poorly understood by both the public and policy-makers, leading to hindered public support and participation, inadequate management actions, and hampered conservation outcomes (Courchamp et al., 2017).

The recent *InvaCost* database, which compiles the economic costs of IAS (including management- and damage-related costs), provides a quantitative metric documenting the economic impacts of IAS (i.e. monetary costs in a standardised currency and year; Diagne et al., 2020, 2021). As such, it succeeded in facilitating public and policy awareness (Ahmed et al., 2023). For ecological impacts, multiple metrics have also been proposed, primarily based on impact categories. For example, the Environmental Impact Classification for Alien Taxa scheme (EICAT; Blackburn et al., 2014; Hawkins et al., 2015) categorises the maximum impact an IAS has had globally into four categories above minimal (undetected) impact, and has been adopted by the IUCN (IUCN, 2020b, 2020a). These EICAT scores range from impacts on individual fitness to the local extinction of species and irreversible changes in

community composition (see Bernardo-Madrid et al., 2022 for an assessment of the

consistency of seven impact classification schemes).

However, to our knowledge, there is no fully quantitative, continuous impact metric that

would allow for comparison of the ecological impacts of IAS across taxa and scales. Such a

metric would allow for more granular impact comparisons among contexts and taxa with

more detailed analyses and predictions, such as spatial and temporal trends of derived

indicators, quantitative links with invasion drivers and alien species traits, or disentangling

the demographic and per-unit or per-capita components of impact, therefore allowing for

more targeted management actions (Latombe et al., 2022). Since managing biological

invasions is often the most effective conservation action (Langhammer et al., 2024), robust

and comparable quantitative metrics would permit prioritisation of action as requested from

Target 6 of the Kunming-Montreal Global Biodiversity Framework (GBF), which aims to

"eliminate, minimise, reduce and or mitigate the impacts of invasive alien species on

biodiversity and ecosystem services" (CBD, 2022). It would also underpin indicators

appropriate for risk identification and prioritisation efforts to meet global biodiversity targets

(Butchart et al., 2005; Vicente et al., 2022). The lack of such standardised metrics capable of

capturing the severity of IAS impacts to ecosystems and their services has been a major

impediment, not only to action, but also to effective communication between scientists,

stakeholders and policy-makers.

Here, we propose the Extinction Potential Metric (EPM) as a measure of impact of individual

IAS. The EPM of an IAS can be conceptualised as the number of species that have already

been led to extinction plus those that are expected to go extinct within a 50-year time frame

under a business-as-usual scenario because of this IAS. It can also be seen as the

equivalent number of extinct species, allowing for the comparison of IAS with different

current levels of impacts on multiple native species, from complete extinction to population

decrease or range contraction. The EPM uses the existing IUCN Red List of Threatened

Species framework (IUCN, 2024), quantifying the threats posed by IAS to native fauna to

compute a risk of extinction within 50 years for each native species threatened by IAS.

Moreover, since native species extinctions are unequal in terms of evolutionary

distinctiveness among taxa, we introduce an additional metric, namely EPM-U, representing

the number of 'unique' native species that have and would go extinct within a 50-year time

frame. Species uniqueness accounts for the evolutionary history of the impacted native

species. EPM-U therefore compares IAS with impacts on phylogenetically related vs

unrelated species. Below, we present the EPM and EPM-U metrics and their computations

in detail. We showcase both metrics with a proof of concept on native terrestrial vertebrate

species threatened by IAS (amphibians, reptiles, birds and mammals). We also show how

EPM and EPM-U can be used to identify which impact mechanisms are most detrimental to

native species. EPM and EPM-U complement other methodologies to measure the impact of

IAS, and in particular EICAT (Blackburn et al., 2014; Hawkins et al., 2015).

Method

Data: Native species impacted by identified invasive alien species

We utilised the IUCN Red List data (IUCN, 2024) to catalogue terrestrial vertebrate species

impacted by IAS, i.e. those associated with Threat 8.1 (Invasive non-native/alien

species/diseases). We identified 2178 amphibians, 920 birds, 865 reptiles, and 473

mammals in categories Least Concern (LC), Near-Threatened (NT), Vulnerable (VU),

Critically Endangered (CR), Extinct in the Wild (EW), Extinct (EX), and Data Deficient (DD)

for which at least one IAS was listed as a threat. DD species were discarded from the

analyses. Invasive organisms threatening these native species were identified at the species

level when possible (73%), but some were only identified at the genus (7%), family (1%), or

order and class (0.6%) level. For 19% of the threats, the invasive organism was not

identified and was recorded as "Unspecified species". We refer to entries in the database as

IAS hereafter for simplicity, and to genus, family, or order when specifically addressing these

categories. For each of the native species previously identified, we also distinguished

between species endemic to islands and other species, using data from Marino et al. (2022).

We also extracted information on the scope (proportion of population affected) and severity

(speed of population decline) of the IAS threat to these native species from the database.

Those with over 50% of their population witnessing a very rapid (>30% over 10 years or

three generations; whichever is the longer), rapid (20–30% over 10 years or three

generations; whichever is the longer), or slow and significant (<20% over 10 years or three

generations; whichever is the longer) decline due to IAS were deemed to experience a high

impact, while the other species were considered to experience a low impact (definitions from

IUCN & CMP, 2012a). When neither the extent nor the severity was available, impact level

was considered as 'not available' (NA). In addition, we extracted the mechanism by which

each native species was affected by IAS, using the IUCN Red List stress classification

scheme (IUCN & CMP, 2012b).

In addition to tabular information summarising the threat details on native species described

above, the IUCN Red List website provides more detailed textual descriptions of the threats

affecting native species, including from IAS, interlinking data with the Global Invasive

Species Database implemented by the IUCN Invasive Species Specialist Group.

Examination of these textual descriptions revealed that species included in the description

are often missing from the tabular information. To assess the effects of this difference of

information content, we manually examined the description for the 473 mammals in the LC,

NT, VU, CR, EW, and EX categories, and generated an extended database including all

species mentioned both in the text and in the table. Comparison between the extended

dataset and the tabular dataset for mammals showed that the tabular information could miss

up to 10 native species impacted by a given IAS with a mean value of 0.89±1.12 (Supp. Fig.

1).

Computing the probability of extinction of each native species

To convert IUCN Red List categories into probabilities of extinction, we followed the EDGE2

and FUSE approaches (Gumbs et al., 2023; Pimiento et al., 2023) (Fig. 1). These

approaches first assign the following species extinction probabilities within 50 years based

on their IUCN Red List category: 0.06 for LC, 0.12 for NT, 0.24 for VU, 0.49 for EN, and 0.97

for Critically Endangered CR. EX and EW were attributed a probability of 1. To mitigate

potential biases resulting from the discrete categorisation of species extinction risk, we then

applied probabilistic draws for VU to CR species. These five IUCN Red List categories were

first transformed into ordinal values from 1 to 5, and a quartic distribution was then applied to

link these ordinal values and the extinction probabilities using the `polyfit` function from the

pracma R package V 2.4.4 (Borchers, 2023) (Figure 1). For each species, a value was

randomly chosen from a uniform distribution spanning the range of its ordinal category value

+/- 0.5, using the `runif` function in R. We then computed its probability of extinction based

on the previously established quartic distribution (Figure 1). This process was replicated 10

times for each native species, generating a set of 10 extinction probabilities per species

according to their IUCN Red List category, enabling us to generate confidence intervals and

account for uncertainty.

Translating the extinction risks of native species into future number of species

extinct due to IAS

We calculated the extinction potential of each IAS by summing the probabilities of extinction

(including EX and EW species) for all native species impacted by this specific IAS. However,

the contribution of an IAS to the extinction risk or to the past extinction of a native species

can vary. According to the IUCN Red List, the impact of IAS on a native species can be

classified as high or low (Fig. 1), but these data can also be missing (hereafter NA). To

account for these different levels of impact of IAS on the decline of native species population

globally (low, high, or NA—as described above), we divided the probability of extinction of

native species by a factor 2 for low impacts. For IAS whose impacts were classified as NA

on some native species, we performed two sets of analyses to account for different

uncertainty scenarios. First, assuming that NAs potentially reflect very small impacts, we

excluded the corresponding native species in the computation of EPM values, providing a

conservative value. As a result, 10 EPM values were computed for each IAS, based on the

10 extinction probabilities generated for each native species, as described above, adjusted

for the level of impact of IAS.

Second, assuming NAs may simply represent a gap in how impacts were reported rather

than a level of impact, we randomly selected between low and high impact categories, and

repeated this process 100 times to assess the uncertainty linked to data deficiency. As a

result, for each original set of 10 probabilities of extinction generated for each native species,

we calculated 100 scores when the impact level of an IAS was NA, corresponding to their

probability of extinction adjusted for the impact level. Consequently, to each of these IAS

corresponded 1000 sets of extinction probabilities for the native species they impacted (10

extinction probabilities drawn from the quartic function x 100 potential impact levels). For

each IAS, we therefore obtained a distribution of the number of native extinct species. We

then computed the EPM for all native species together, but also for each native taxonomic

group separately (i.e. each IAS had an impact distribution for amphibians, birds, mammals,

and reptiles).

Considering unspecified IAS

As some IAS were only identified at the genus and family levels in the IUCN Red List

database, we also used information on unspecified species to compute the maximum

possible EPM value for an IAS based on the available data, by assuming unspecified

species included all species in the reported genus or family. In other words, we considered a

given IAS to be responsible for all the threats for which it is identified, but also for all the

threats imposed by unspecified IAS of the same genus or family. For example, the maximum

EPM of *Rattus rattus* was computed by aggregating the EPM of *Rattus rattus*, the EPM of

Unspecified Rodentia and the EPM of Unspecified *Rattus*. The maximum EPM of the other

Rattus species, i.e. *Rattus norvegicus*, and *Rattus exulans*, was computed in the same way.

Conversely, doing so implies that native species impacted by unspecified *Rattus* species

were considered as affected by all *Rattus* species (i.e. *Rattus rattus*, *Rattus norvegicus*, and

Rattus exulans).

Accounting for evolutionary distinctiveness

Evolutionary distinctiveness is an important component of biodiversity that can be

incorporated into our approach by following an approach similar to that used to compute the

EDGE2 metric (Gumbs et al., 2023). The EDGE2 score of a species represents the number

of millions of years of evolutionary history lost if a species goes extinct. It is the product of

the heightened evolutionary distinctiveness (ED2) of a species and of its risk of extinction

(GE2, for global endangerment). ED2 accounts for the phylogenetic distance of a species

from other species (its "raw" evolutionary distinctiveness ED), but also the extinction risk of

these other species. A species with an intermediate ED whose phylogenetically related

species are endangered will therefore have a higher ED2, as the risk of losing the whole

clade is then higher.

To account for evolutionary history in measuring the impact of IAS, we propose an additional

metric, the Extinction Potential Metric for Unique species due to an IAS (EPM-U), which

incorporates ED2 scores into EPM. As ED2 values are only available for mammals (Gumbs

et al. 2023), we only computed EPM-U of IAS on native mammals. To do so, we propose to

use the maximum ED2 score of all mammals to determine the most unique species in the

database, with a value of 1. We then rescaled all ED2 values by dividing them by this

maximum value, to generate a "uniqueness score" for each impacted native species in [0,1].

In addition, Gumbs et al. (2023) account for uncertainty in the calculations of ED2 values

and provide a median value, with the corresponding lower and upper interquartile ranges for

each species. We sampled 10 ED2 values per native species from a normal distribution

where the mean is the median ED2 value and the standard deviation is the lower

interquartile interval divided by 1.35. These 10 ED2 values for each native species were

rescaled between 0 and 1 after dividing them by the maximum ED2 value. We then

computed the number of unique extinct species due to an IAS as the sum of the probabilities

of extinction for each native species impacted by the IAS multiplied by their uniqueness

score. Following the same methods as for the EPM, we generated 100 scores per IAS.

Linking impact strength to impact mechanisms

We assessed the importance of the 11 impact mechanisms listed in the IUCN Red List

through which IAS affect native species, using EPM values. For each of these 11

mechanisms—competition, ecosystem conversion, ecosystem degradation, hybridisation,

inbreeding, indirect ecosystem effects, reduced reproductive success, skewed sex ratios,

species disturbance, species mortality, and species stresses—we extracted the native

species impacted by any IAS (identified or not). We then computed the corresponding EPM

score based on the IUCN Red List category of the native species, the severity, and the

scope of the threats, generating an EPM score for each impact mechanism following the

same framework as detailed above. This was done for all native species and for each

taxonomic group separately.

Results

Global impacts of IAS on native taxa

Considering all native species for which the impact level of IAS was known, we observed

from the 196 IAS that median EPM values (all EPM values hereafter correspond to median

values) ranged from 0.02 (*Pavo cristatus*) to 89.8 (*Batrachochytrium dendrobatidis*) extinct

species, with a mean value of 2.4±9.64 (hereafter corresponding to the standard deviation).

EPM values followed a log-normal distribution, with a few species having high values and

most having low values and 47 IAS with a value above 1 extinct species (Fig. 2.A, Supp.

Table 1). When including native species with an impact level defined as 'NA' (i.e. randomly

assigned to low or high 100 times), 357 IAS were included, and mean EPM values increased

by 38% with 80 IAS above 1 extinct species. The EPM values of IAS with the greatest

impact were particularly affected, with *B. dendrobatidis* rising from 89.8 to 380.7, *Felis catus*

from 87.2 to 138.8, and *Batrachochytrium salamandrivorans* from 15.1 to 88.5 (Supp. Fig.

2.A).

When computing EPM values of IAS for the different native taxonomic groups separately

(i.e. the equivalent number of extinct native amphibians, birds, mammals or reptiles) we

observed different rankings across taxa (Fig. 2.B, Supp. Table 2). Five IAS had an EPM

higher than 1 extinct amphibian species. Two pathogenic fungi were particularly impactful, *B.*

dendrobatidis (the pathogen causing chytridiomycosis in frogs; EPM = 89.8.0), followed by

B. salamandrivorans (EPM = 15.1). Twenty-six IAS had an EPM above 1 extinct bird

species, and three IAS above 10: *Felis catus* with (EPM = 40.1), *Rattus rattus* (EPM = 27.2),

and *Canis familiaris* (EPM = 10.7) (Fig. 2.B). Five IAS had an EPM score above one extinct

mammal species, but one was particularly impactful: *Felis catus* (EPM = 22.6) (Fig. 2.B).

Finally, 15 IAS had an EPM score above 1 extinct reptile species, and 3 had a score above

10: *F. catus* (EPM = 25.0), *Wasmannia auropunctatus* (EPM = 12.2), and *Herpestes*

auropunctatus (EPM = 11.3) (Fig. 2.B). When we included native species with an impact

defined as 'NA', EPM values increased particularly for amphibians, for which the number of

IAS with a score above 1 rose from 5 to 19 (a 280% increase). The number of IAS above 1

rose by 30% for birds (34 *vs* 26), 100% for mammals (10 *vs* 5) and 60% for reptiles (25 *vs*

15) (Supp. Fig. 2.B, Supp. Table 3). Similarly, we observed substantial increases in EPM

values for the most impacting IAS for amphibians (*B. dendrobatidis* EPM = 380.5 and *B.*

salamandrivorans = 87.0) but also for reptiles (*F. catus* EPM = 51.8, *R. rattus* EPM = 21.3,

and *H. auropunctatus* EPM = 19.8) and to a lesser extent for birds (*F. catus* with EPM =

54.1, *R. rattus* EPM = 32.3, and *C. familiaris* EPM = 14.4) and mammals (*F. catus* EPM =

31.7) (Supp. Fig. 2.B, Supp. Table 4).

Results were similar for maximum EPM scores considering unspecified genera or families,

with *Rattus* species seeing the main increase, due to many impacts being reported for

unspecified *Rattus* and rodents in the IUCN Red List database (Supp. Fig. 3). *Mus musculus*

and *H. auropunctatus* had a substantive increase in their score when considering

unspecified rodents and unspecified *Herpestes*.

Insular endemic species accounted for 82.0±25.7% of the previously computed EPM scores

across all native species. Specifically, the impact of IAS was highest for island endemic

amphibians and reptiles, with mean changes of -92.4±27.2% and -88.7±16.6%, respectively.

The effect was slightly less pronounced in birds, at -80.6±27.8%, and was the weakest in

mammals, at -62.0±37.0% (Fig. 3). Adding native species with 'NA' impact did not broadly

affect these results (Supp. Fig. 4).

Impacts of IAS on mammal species: tabular vs. textual information and number of

unique extinct species

When considering the extended database comprising the tabular (high, low, and 'NA'

threats) and textual information (only 'NA' threats), 29 IAS had an EPM score of more than 1

extinct mammal species (Fig. 4). *F. catus* had the largest impacts (EPM = 64.2), followed by

C. familiaris (EPM = 30.6), *R. rattus* (EPM = 20.0), and *Vulpes vulpes* (EPM = 19.3). When

considering the evolutionary distinctiveness of impacted mammals, the same 4 species had

the highest EPM values (*F. catus* = 3.9, *C. familiaris* = 2.2, *V. vulpes* = 1.9, and *R. rattus* =

0.98) (Supp. Fig. 6.B). The evolutionary distinctiveness of impacted mammals had little

effect on the ranking of the four most impactful IAS, but had important effects on rankings for

other IAS (Fig. 5).

Impacts caused through different mechanisms

When considering all taxa, 'Species mortality' was associated with the largest EPM value of

396.9, more than three times the second and third highest EPM value from 'reduced

reproductive success' (EPM = 122.5) and 'ecosystem degradation' (EPM = 109.6) (Fig. 6.A).

Furthermore, 'Species mortality' was the most consistent mechanism, showing a significant

association with higher EPM scores for amphibians, birds, mammals, and reptiles (Fig. 6.B).

For native birds, it was closely followed by 'reduced reproductive success,' and for native

birds and reptiles, 'ecosystem degradation' also had a substantial effect (Fig. 6.B). Adding

native species with 'NA' impact did not affect 'species mortality' which stayed largely the

most impactful mechanism with EPM = 1018.7, but 'ecosystem degradation' became the

second most important mechanisms with EPM = 235.3 above 'reduced reproductive

success' EPM = 158.4 when considering all taxa (Supp. Fig. 6).

Discussion

We introduced and applied a new metric, the Extinction Potential Metric, which computes the

ecological impacts of IAS as the equivalent number of native species that IAS have

contributed to driving to extinction. The EPM uses the IUCN Red List assessments to

compute extinction probabilities of native species impacted by IAS within 50 years, in

addition to those species already extinct. A quantitative EPM value is then attributed to each

IAS based on the number of species it is expected to drive to extinction, allowing for detailed

comparison of the impacts of different IAS. We also introduced an alternative metric, EPM-U,

the Extinction Potential Metric for Unique species, to account for differences in evolutionary

history of impacted native species, and complement the information provided by EPM.

Most impacts on native tetrapods result from a few IAS

Birds, mammals, and reptiles were all mostly impacted by cats (*F. catus*), followed by dogs

(*C. familiaris*) and rats (especially the black rat, *R. rattus*, but also other rat species, often

unspecified or including *Rattus norvegicus* and *Rattus exulans* for birds), foxes (*V. vulpes*),

the small Indian mongoose (*H. auropunctatus*), and domestic goats (*Capra hircus*). All these

species belong to the IUCN list of 100 of the worst invasive species (Lowe et al., 2000;

Luque et al., 2014). Results are nonetheless different for amphibians, which are mostly

affected by two invasive pathogenic fungi, *B. dendrobatidis* and *B. salamandrivorans* (Fig.

2), with the former also appearing in the 100 of the worst list. The number of amphibian

species affected by these two species is higher than the number of all tetrapod species

affected by any other IAS (see also Bellard, Genovesi, et al., 2016). An important part of

these impacts is recorded for species that are endemic to islands (Fig. 3), indicating the

disproportionate impacts IAS have on species that are potentially evolutionarily distinct

within insular habitats (Millien, 2006). In addition, for most IAS, the numbers of impacted

native species in each Red List category are of the same order of magnitude (Fig. 7, Supp.

Fig. 7). This means that EPM scores are mostly driven by past extinctions (EX and EW

categories) or by decreases in native species populations and ranges that have driven them

to the brink of extinction (CR category), rather than by expected future extinctions (LC to VU

categories).

The evidence presented above confirms that controlling a limited number of IAS could

effectively prevent and mitigate the impacts of biological invasions on biodiversity. This is

consistent with the conclusions of the IPBES assessment that highlighted that, with sufficient

resources, political will and long-term commitment, preventing and controlling IAS are

attainable goals that will yield significant long-term benefits for nature (Roy et al., 2024).

Also, the results of our analysis support the need to identify priority IAS for prevention and

control, as requested from Target 6 of the GBF.

Considering the evolutionary distinctiveness of impacted native mammals, did not

fundamentally change the ranking of the most impactful IAS (Fig. 5), but changed the score

distribution, with a more gradual increase in EPM-U compared to EPM values (Supp. Fig. 5).

Evolutionary distinctiveness, and more generally phylogenetic diversity, is a core component

of conservation as it captures multiple concepts (Gumbs et al., 2023; Lean & Maclaurin,

2016; Winter et al., 2013). Evolutionarily distinct species may be considered to have a high

intrinsic value on the basis of their uniqueness. Phylogenetic diversity (maximised by

conserving the most unique species) has also been advanced as a proxy for functional

diversity and evolutionary potential, whose maximisation is important for ecosystem services

and for maintaining the benefits humans draw from nature, although these claims have been

questioned (Mazel et al., 2018; Winter et al., 2013). The difference in EPM and EPM-U score

distribution has implications for the design of management actions and IAS prioritisation.

Although assessing the risk caused by IAS is insufficient to prioritise which IAS to manage,

which should also account for management feasibility and costs (Robertson et al., 2021), it is

an essential component of this process. Here, using EPM scores alone would give a greater

weight to the few species with a disproportionately high score, whereas EPM-U scores

would suggest considering a wider range of species.

"Species mortality" is the main mechanism through which IAS impact native species (Fig. 6).

This occurs through predation for mammals, birds, and reptiles, and through two diseases

for amphibians. Ranking of the other mechanisms varied substantially across the four

taxonomic groups. Interestingly, birds were the only group for which reduced reproductive

success was an important mechanism, ranking second. However, this is likely due to the fact

that IAS consume native bird eggs before hatching, which is therefore similar to predation in

the sense that this is linked to direct resource consumption. Aggregating the EPM scores of

these two impact mechanisms for native birds would generate a profile similar to that of the

other taxonomic groups. Other important but secondary, non-lethal impact mechanisms

include ecosystem degradation, species disturbance, and competition. The predominance of

direct predation over other impact mechanisms suggests that appropriate IAS management

requires the removal of interactions between IAS predators and their prey. It may also

suggest that increased attention is needed towards more subtle threat mechanisms which

are less conspicuous than those driving direct mortality. While lethal control of predators

may seem to be the most obvious approach, it may not be effective or even achievable,

especially on the mainland (García-Díaz et al., 2021). IAS populations can recover quickly,

preventing eradication success, and lethal control may need to be performed at levels that

are not attainable with current techniques, or are financially unsustainable to see positive

effects. Efficacy is also context-dependent, with studies on various native taxa showing

effects of IAS control ranging from positive to negligible and even negative effects (García-

Díaz et al., 2021). Context-specific alternative approaches to lethal control have been

proposed, including IAS exclusion, for example using fences (D. Smith et al., 2020), nest

cages to prevent egg consumption (Buzuleciu et al., 2015; R. K. Smith et al., 2011),

translocation of native species (Miskelly & Powlesland, 2013), or prey training and

harnessing natural selection to combat prey naivete (Moseby et al., 2016), although their

efficacy has been questioned. Nevertheless, IAS eradications have had high success rates

on islands (Spatz et al., 2022), which contributed the majority of extinction risk in those

habitats.

The results presented here should be taken with some level of caution. First, the IUCN Red

List, despite being a major conservation resource, is not fully comprehensive, and not all

native nor invasive species have been assessed or are up-to-date (Cazalis et al., 2022,

2023). This is exemplified by the large shares of IAS with impacts classified as 'NA' towards

particular native species in the current study. The resource is also more likely to report

impacts from IAS when these have been long standing, and often uses observed and

inferred impact through expert opinion rather than comprehensive, published evidence (e.g.

Gula et al., 2023; Van der Colff et al., 2021). The figures we present are therefore

conservative. The IUCN Red List is also of limited use in identifying future problematic IAS,

given time lags to invasion and impact (Essl et al., 2011). Finally, and importantly, the IUCN

Red List data is open access and can easily be extracted from the IUCN Red List website as

text files containing data in a tabular format. However, examination of more detailed textual

information provided on the webpage of each assessed native mammal revealed that this

tabular information was incomplete, with up to 10 IAS missing in the list of threats (Supp.

Fig. 1). Computing EPM scores for the extended database comprising the textual information

revealed that impacts were largely under-estimated, even for the most impactful species,

and that ranking differed for many species. We therefore advocate for an in-depth curation of

the IUCN Red List database, to harmonise the information content on IAS threats on native

species between tabular and textual information, to ensure consistency across taxa.

Fostering knowledge on IAS impacts on native species is also paramount to improve these

figures and develop an even more accurate application of EPM.

EPM as a measure of ecological impacts

The new metric we propose provides a bespoke quantitative measure of ecological impact of

IAS and has many advantages compared to existing, mostly qualitative metrics. Using

probabilities, the EPM links impacts at the population and species level. It uses species as a

unit, which is the fundamental unit of conservation, as exemplified by the IUCN Red List of

Threatened Species—arguably the most influential global conservation tool to date. As the

IUCN Red List is open access, EPM assessments are transparent, readily applicable and

reproducible. The EPM score of an IAS can be interpreted as the number of species that it

will have driven to extinction in 50 years under a business-as-usual scenario, including past

and future extinctions. Thus, EPM is easily conceptualised by policy-makers and the civil

society, and will therefore foster communication about the impacts of biological invasions

and the design of management actions, such as native conservation prioritisation.

In addition, EPM offers the possibility to be temporally and spatially downscaled. Red List

statuses are ideally updated every few years for each species, which will enable us to

generate estimates and track temporal changes in the impacts of IAS. EPM, as presented

here, assesses ecological impacts of IAS at the global scale, because it is the scale for

which the IUCN most commonly provides Red List categories and with the most complete

level of details. However, Red List categories can also be assessed at national and regional

levels (IUCN, 2012). EPM could therefore also be spatially explicit if the same level of

information on the status and threats on native species is assessed at multiple spatial

scales. This downscaling is particularly pertinent given that the impacts of IAS can differ

substantially among populations occurring in different regions over time, undermining the

efficacy of global species 'watch lists'.

As demonstrated in our analyses, EPMs can be computed for different groups of impacted

native species, allowing for the capture of multiple dimensions of impacts caused by IAS,

rather than generating a single value or category for an IAS. Here, we showed that IAS

impacts varied across four taxonomic groups of native tetrapods. In addition, although we

applied the EPM at the IAS level, it would be straightforward to upscale the metric to the

genus level or beyond, or for any group of IAS of interest. For example, here we also

quantified the importance of each mechanism through which IAS impact native species, by

computing EPM scores after grouping all IAS impacting native species through these threat

mechanisms. Although the most important impacts occur through direct mortality, it also

revealed that other, non-lethal impacts (e.g. competition or species disturbance, such as the

disruption of critical lifecycle stages) or changes in the environment (e.g. ecosystem

degradation) can also have important impacts. Doing so therefore enables us to capture the

broader ecological impacts of biological invasions for different components of ecosystems,

and how they cascade to *in fine* affect native species survival.

Note, however, that native species impacted by biological invasions are often threatened by

multiple IAS simultaneously, as well as other anthropogenic pressures, and their effects can

combine non-additively. Disentangling the relative impacts of different IAS on a single native

species is not possible with the current granularity of IUCN data. It is consequently not

possible to assess if an IAS could lead a native species to extinction in the absence of

another IAS (e.g. cats are often associated with red foxes in Australia, but have led many

native species to extinction by themselves in other regions). In the absence of a clearly

identifiable rule to determine the contribution of each IAS, here we took a conservative,

objective and reproducible approach, attributing the same probability to cause extinction to

different IAS affecting a native species with the same level of impact. This follows

precautionary principles, an approach already used by other impact classification schemes,

including EICAT (Kumschick et al., 2024), to avoid irreversible impacts or high costs to the

environment and society that would result from inaction.

The versatility of EPM offers avenues to derive indicators assessing the ecological impacts

of biological invasions to inform international policy initiatives, in particular Target 6 of the

GBF. This framework is lacking appropriate indicators that would allow tracking of policy

performance regarding this target (Henriksen et al., 2024). Such indicators should capture

spatial and temporal changes, be species-specific and allow for identification of the

mechanisms through which IAS impact native species (Henriksen et al., 2024), which are all

readily included in EPM. Different indicators have been developed, but are either qualitative

(e.g. Genovesi et al., 2012), categorical (e.g. Kumschick & Nentwig, 2010), do not provide

information on the IAS and their impact mechanisms (e.g. Butchart, 2008; Butchart et al.,

2005), or focus on species extinction (i.e., extreme 'end-points') rather than combining

multiple levels of earlier-stage impacts on different native species (e.g. Bellard, Cassey, et

al., 2016). Indicators derived from EPM could be combined with other indicators of impacts

(Van der Colff et al., 2021) but also with trends or projections in IAS numbers and ranges,

trends in introduction and spread mechanisms, and trends in policy responses (McGeoch et

al., 2006, 2010; Rabitsch et al., 2016), to provide a comprehensive description of the issue

and analysis of its drivers.

Finally, the EPM could be used to assess the ecological impacts of other anthropogenic

pressures listed as threats under the IUCN Red List classification scheme (e.g. residential &

commercial development, agriculture & aquaculture, pollution, climate change & severe

weather, etc., as per the IUCN Red List Threat Classification scheme V3.3; IUCN & CMP,

2012a). Doing so could lead to a set of harmonised indicators to assess progress towards

different targets of the GBF (e.g. Target 7: Reduce Pollution to Levels That Are Not Harmful

to Biodiversity; CBD, 2022). It could also permit analyses of synergies among global change

drivers, by identifying which processes combine to create the highest threats, compared to

where they occur individually. Indeed, specific native species are often impacted by multiple

anthropogenic pressures simultaneously, which can reinforce each other.

Conclusion

We introduced the Extinction Potential Metric, a new quantitative metric to assess the

ecological impacts of invasive alien species. The EPM computes the equivalent number of

native species driven to extinction by an IAS using the IUCN Red List categories to assign

probabilities of extinction within 50 years. Doing so enables us to incorporate IAS impacts at

the population and species levels into a single metric, and also to account for evolutionary

distinctiveness between impacted native species. Applying this metric to native terrestrial

vertebrates shows that a few IAS disproportionately affect native species populations

through direct mortality, suggesting that controlling a limited number of IAS could effectively

prevent and mitigate the impacts of biological invasions on biodiversity. The metric is

versatile and transparent, and can be applied to different groups of native species and IAS,

to permit prioritisation of conservation action and to provide a detailed and more

comprehensive assessment of IAS ecological impacts than existing approaches. We argue it

can be the basis to derive spatio-temporal indicators that will improve our capacity to assess

the efficacy of global and regional conservation policies.

576
577

Acknowledgements

HS acknowledges funding by the Deutsche Forschungsgemeinschaft (DFG, German
Research Foundation) (grant no. 521529463). SK acknowledges the support of the Centre
for Invasion Biology (CIB) at Stellenbosch University, the South African Department of
Forestry, Fisheries and the Environment (DFFE), and the B3 (Biodiversity Building Blocks for
Policy) project, which receives funding from the European Union's Horizon Europe Research
and Innovation Programme (through grant no. 584101059592).

586
587

Data availability

The data on which this paper is based are freely available on the IUCN Red List website
(www.redlist.org). A list of the native and invasive alien species used in the analyses is given
in the electronic supplementary material.

593
594 **References**
595
Ahmed, D. A., Haubrock, P. J., Cuthbert, R. N., Bang, A., Soto, I., Balzani, P., Tarkan, A. S.,

Macêdo, R. L., Carneiro, L., Bodey, T. W., Oficialdegui, F. J., Courtois, P.,

Kourantidou, M., Angulo, E., Heringer, G., Renault, D., Turbelin, A. J., Hudgins, E. J.,

Liu, C., … Courchamp, F. (2023). Recent advances in availability and synthesis of

the economic costs of biological invasions. *BioScience*, *73*(8), 560–574.

https://doi.org/10.1093/biosci/biad060

Bellard, C., Cassey, P., & Blackburn, T. M. (2016). Alien species as a driver of recent

extinctions. *Biology Letters*, *12*(2), 20150623–20150623.

https://doi.org/10.1098/rsbl.2015.0623

Bellard, C., Genovesi, P., & Jeschke, J. M. (2016). Global patterns in threats to vertebrates

by biological invasions. *Proceedings of the Royal Society B: Biological Sciences*,

283(1823), 20152454. https://doi.org/10.1098/rspb.2015.2454

Bernardo-Madrid, R., González-Moreno, P., Gallardo, B., Bacher, S., & Vilà, M. (2022).

Consistency in impact assessments of invasive species is generally high and

depends on protocols and impact types. *NeoBiota*, *76*, 163–190.

https://doi.org/10.3897/neobiota.76.83028

Blackburn, T. M., Essl, F., Evans, T., Hulme, P. E., Jeschke, J. M., Kühn, I., Kumschick, S.,

Marková, Z., Mrugała, A., Nentwig, W., Pergl, J., Pyšek, P., Rabitsch, W., Ricciardi,

A., Richardson, D. M., Sendek, A., Vilà, M., Wilson, J. R. U., Winter, M., … Bacher,

S. (2014). A unified classification of alien species based on the magnitude of their

environmental impacts. *PLoS Biology*, *12*(5), e1001850–e1001850.

https://doi.org/10.1371/journal.pbio.1001850

Borchers, H. (2023). *_pracma: Practical Numerical Math Functions_. R package version*

2.4.4, <https://CRAN.R-project.org/package=pracma>. [Computer software].

Butchart, S. H. M. (2008). Red List Indices to measure the sustainability of species use and

    impacts of invasive alien species. *Bird Conservation International*, *18*(S1), S245–

    S262. https://doi.org/10.1017/S095927090800035X

Butchart, S. H. M., Stattersfield, A. J., Baillie, J., Bennun, L. A., Stuart, S. N., Akçakaya, H.

    R., Hilton-Taylor, C., & Mace, G. M. (2005). Using Red List Indices to measure

    progress towards the 2010 target and beyond. *Philosophical Transactions of the*

    Royal Society B: Biological Sciences, *360*(1454), 255–268.

    https://doi.org/10.1098/rstb.2004.1583

Buzuleciu, S. A., Spencer, M. E., & Parker, S. L. (2015). Predator Exclusion Cage for Turtle

    Nests: A Novel Design. *Chelonian Conservation and Biology*, *14*(2), 196–201.

    https://doi.org/10.2744/CCB-1163.1

Cazalis, V., Di Marco, M., Butchart, S. H., Akçakaya, H. R., González-Suárez, M., Meyer, C.,

    Clausnitzer, V., Böhm, M., Zizka, A., & Cardoso, P. (2022). Bridging the research-

    implementation gap in IUCN Red List assessments. *Trends in Ecology & Evolution*,

    37(4), 359–370.

Cazalis, V., Santini, L., Lucas, P. M., González-Suárez, M., Hoffmann, M., Benítez-López,

    A., Pacifici, M., Schipper, A. M., Böhm, M., Zizka, A., Clausnitzer, V., Meyer, C.,

    Jung, M., Butchart, S. H. M., Cardoso, P., Mancini, G., Akçakaya, H. R., Young, B.

    E., Patoine, G., & Di Marco, M. (2023). Prioritizing the reassessment of data-deficient

    species on the IUCN Red List. *Conservation Biology*, *37*(6), e14139.

    https://doi.org/10.1111/cobi.14139

CBD. (2022). *Decision adopted by the conference of the parties to the convention on*

    biological diversity 15/4. Kunming-montreal global biodiversity framework.

Courchamp, F., Fournier, A., Bellard, C., Bertelsmeier, C., Bonnaud, E., Jeschke, J. M., &

    Russell, J. C. (2017). Invasion biology: Specific problems and possible solutions.

    Trends in Ecology & Evolution, *32*(1), 13–22.

    https://doi.org/10.1016/j.tree.2016.11.001

647   Diagne, C., Leroy, B., Gozlan, R. E., Vaissière, A.-C., Assailly, C., Nuninger, L., Roiz, D.,

648   Jourdain, F., Jarić, I., & Courchamp, F. (2020). InvaCost, a public database of the

649   economic costs of biological invasions worldwide. *Scientific Data*, *7*(1), 1–12.

650   https://doi.org/10.1038/s41597-020-00586-zgas

651   Diagne, C., Leroy, B., Vaissière, A.-C., Gozlan, R. E., Roiz, D., Jarić, I., Salles, J.-M.,

652   Bradshaw, C. J. A., & Courchamp, F. (2021). High and rising economic costs of

653   biological invasions worldwide. *Nature*, *592*, 571–576.

654   https://doi.org/10.1038/s41586-021-03405-6

655   Essl, F., Dullinger, S., Rabitsch, W., Hulme, P. E., Hülber, K., Jarošík, V., Kleinbauer, I.,

656   Krausmann, F., Kühn, I., & Nentwig, W. (2011). Socioeconomic legacy yields an

657   invasion debt. *Proceedings of the National Academy of Sciences*, *108*(1), 203–207.

658   https://doi.org/10.1073/pnas.1011728108

659   García-Díaz, P., Cassey, P., Norbury, G., Lambin, X., Montti, L., Pizarro, J. C., Powell, P. A.,

660   Burslem, D. F. R. P., Cava, M., Damasceno, G., Fasola, L., Fidelis, A., Huerta, M. F.,

661   Langdon, B., Linardaki, E., Moyano, J., Núñez, M. A., Pauchard, A., Phimister, E., …

662   Tomasevic, J. A. (2021). Management Policies for Invasive Alien Species:

663   Addressing the Impacts Rather than the Species. *BioScience*, *71*(2), 174–185.

664   https://doi.org/10.1093/biosci/biaa139

665   Genovesi, P., Carnevali, L., Alonzi, A., & Scalera, R. (2012). Alien mammals in Europe:

666   Updated numbers and trends, and assessment of the effects on biodiversity.

667   *Integrative Zoology*, *7*(3), 247–253. https://doi.org/10.1111/j.1749-4877.2012.00309.x

668   Gula, J., Sundar, K. S. G., Willows-Munro, S., & Downs, C. T. (2023). The state of stork

669   research globally: A systematic review. *Biological Conservation*, *280*, 109969.

670   https://doi.org/10.1016/j.biocon.2023.109969

671   Gumbs, R., Gray, C. L., Böhm, M., Burfield, I. J., Couchman, O. R., Faith, D. P., Forest, F.,

672   Hoffmann, M., Isaac, N. J. B., Jetz, W., Mace, G. M., Mooers, A. O., Safi, K., Scott,

673   O., Steel, M., Tucker, C. M., Pearse, W. D., Owen, N. R., & Rosindell, J. (2023). The

674   EDGE2 protocol: Advancing the prioritisation of Evolutionarily Distinct and Globally

Endangered species for practical conservation action. *PLOS Biology*, *21*(2),

e3001991. https://doi.org/10.1371/journal.pbio.3001991

Hawkins, C. L., Bacher, S., Essl, F., Hulme, P. E., Jeschke, J. M., Kühn, I., Kumschick, S.,

Nentwig, W., Pergl, J., Pyšek, P., Rabitsch, W., Richardson, D. M., Vilà, M., Wilson,

J. R. U., Genovesi, P., & Blackburn, T. M. (2015). Framework and guidelines for

implementing the proposed IUCN Environmental Impact Classification for Alien Taxa

(EICAT). *Diversity and Distributions*, *21*(11), 1360–1363.

Henriksen, M. V., Arlé, E., Pili, A., Clarke, D. A., García-Berthou, E., Groom, Q., Lenzner, B.,

Meyer, C., Seebens, H., Tingley, R., Winter, M., & McGeoch, M. A. (2024). Global

indicators of the environmental impacts of invasive alien species and their

information adequacy. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *379*(1902), 20230323. https://doi.org/10.1098/rstb.2023.0323

IUCN. (2012). *Guidelines for Application of IUCN Red List Criteria at Regional and National*

Levels: Version 4.0. (p. iii + 41pp).

IUCN. (2020a). *Guidelines for using the IUCN Environmental Impact Classification for Alien*

Taxa (EICAT) Categories and Criteria): First edition. Version 1.1. IUCN.

IUCN. (2020b). *IUCN EICAT Categories and Criteria. The Environmental Impact*

Classification for Alien Taxa (EICAT) First edition. IUCN.

https://doi.org/10.2305/IUCN.CH.2020.05.en

IUCN. (2024). *The IUCN Red List of Threatened Species. Version 2024-1.* [Dataset].

https://www.iucnredlist.org

IUCN, & CMP. (2012a). *IUCN-CMP Unified Classification of Direct Threats Version 3.3.*

https://www.iucnredlist.org/resources/threat-classification-scheme

IUCN, & CMP. (2012b). *IUCN-CMP Unified Classification of Stresses Version 1.1.*

https://nc.iucnredlist.org/redlist/content/attachment_files/dec_2012_guidance_stresse

s_classification_scheme.pdf

Kumschick, S., Bertolino, S., Blackburn, T. M., Brundu, G., Costello, K. E., de Groot, M.,

Evans, T., Gallardo, B., Genovesi, P., Govender, T., Jeschke, J. M., Lapin, K.,

Measey, J., Novoa, A., Nunes, A. L., Probert, A. F., Pyšek, P., Preda, C., Rabitsch,

W., … Bacher, S. (2024). Using the IUCN Environmental Impact Classification for

Alien Taxa to inform decision-making. *Conservation Biology*, *38*(2), e14214.

https://doi.org/10.1111/cobi.14214

Kumschick, S., & Nentwig, W. (2010). Some alien birds have as severe an impact as the

most effectual alien mammals in Europe. *Biological Conservation*, *143*(11), 2757–

2762. https://doi.org/10.1016/j.biocon.2010.07.023

Langhammer, P. F., Bull, J. W., Bicknell, J. E., Oakley, J. L., Brown, M. H., Bruford, M. W.,

Butchart, S. H. M., Carr, J. A., Church, D., Cooney, R., Cutajar, S., Foden, W.,

Foster, M. N., Gascon, C., Geldmann, J., Genovesi, P., Hoffmann, M., Howard-

McCombe, J., Lewis, T., … Brooks, T. M. (2024). The positive impact of conservation

action. *Science*, *384*(6694), 453–458. https://doi.org/10.1126/science.adj6598

Latombe, G., Catford, J. A., Essl, F., Lenzner, B., Richardson, D. M., Wilson, J. R. U., &

McGeoch, M. A. (2022). GIRAE: a generalised approach for linking the total impact of

invasion to species' range, abundance and per-unit effects. *Biological Invasions*,

24(10), 3147–3167. https://doi.org/10.1007/s10530-022-02836-0

Lean, C., & Maclaurin, J. (2016). The value of phylogenetic diversity. *Biodiversity*

Conservation and Phylogenetic Systematics: Preserving Our Evolutionary Heritage in

an Extinction Crisis, 19–37.

Lowe, S., Browne, M., Boudjelas, S., & De Poorter, M. (2000). *100 of the world's worst*

invasive alien species: A selection from the global invasive species database (Vol.

12). Invasive Species Specialist Group Auckland.

Luque, G. M., Bellard, C., Bertelsmeier, C., Bonnaud, E., Genovesi, P., Simberloff, D., &

Courchamp, F. (2014). The 100th of the world's worst invasive alien species.

Biological Invasions, *16*(5), 981–985. https://doi.org/10.1007/s10530-013-0561-5

Marino, C., Leclerc, C., & Bellard, C. (2022). Profiling insular vertebrates prone to biological

invasions: What makes them vulnerable? *Global Change Biology*, *28*(3), 1077–1090.

https://doi.org/10.1111/gcb.15941

Mazel, F., Pennell, M. W., Cadotte, M. W., Diaz, S., Dalla Riva, G. V., Grenyer, R., Leprieur,

F., Mooers, A. O., Mouillot, D., Tucker, C. M., & Pearse, W. D. (2018). Prioritizing

phylogenetic diversity captures functional diversity unreliably. *Nature*

Communications, *9*(1), 2888. https://doi.org/10.1038/s41467-018-05126-3

McGeoch, M. A., Butchart, S. H. M., Spear, D., Marais, E., Kleynhans, E. J., Symes, A.,

Chanson, J., & Hoffmann, M. (2010). Global indicators of biological invasion: Species

numbers, biodiversity impact and policy responses. *Diversity and Distributions*, *16*(1),

95–108.

McGeoch, M. A., Chown, S. L., & Kalwij, J. M. (2006). A Global Indicator for Biological

Invasion. *Conservation Biology*, *20*(6), 1635–1646. https://doi.org/10.1111/j.1523-

1739.2006.00579.x

Millien, V. (2006). Morphological Evolution Is Accelerated among Island Mammals. *PLOS*

Biology, *4*(10), e321. https://doi.org/10.1371/journal.pbio.0040321

Miskelly, C. M., & Powlesland, R. G. (2013). Conservation translocations of New Zealand

birds, 1863–2012. *Notornis*, *60*(1), 3–28.

Moseby, K. E., Blumstein, D. T., & Letnic, M. (2016). Harnessing natural selection to tackle

the problem of prey naïveté. *Evolutionary Applications*, *9*(2), 334–343.

https://doi.org/10.1111/eva.12332

Pimiento, C., Albouy, C., Silvestro, D., Mouton, T. L., Velez, L., Mouillot, D., Judah, A. B.,

Griffin, J. N., & Leprieur, F. (2023). Functional diversity of sharks and rays is highly

vulnerable and supported by unique species and locations worldwide. *Nature*

Communications, *14*(1), 7691. https://doi.org/10.1038/s41467-023-43212-3

Rabitsch, W., Genovesi, P., Scalera, R., Biała, K., Josefsson, M., & Essl, F. (2016).

Developing and testing alien species indicators for Europe. *Journal for Nature*

Conservation, *29*, 89–96. https://doi.org/10.1016/j.jnc.2015.12.001

Robertson, P. A., Mill, A. C., Adriaens, T., Moore, N., Vanderhoeven, S., Essl, F., & Booy, O.

(2021). Risk Management Assessment Improves the Cost-Effectiveness of Invasive

Species Prioritisation. *Biology*, *10*(12). https://doi.org/10.3390/biology10121320

Roy, H. E., Pauchard, A., Stoett, P., Renard Truong, T., Bacher, S., Galil, B. S., Hulme, P.

E., Ikeda, T., Sankaran, K., McGeoch, M. A., Meyerson, L. A., Nuñez, M. A.,

Ordonez, A., Rahlao, S. J., Schwindt, E., Seebens, H., Sheppard, A. W., & Vandvik,

V. (2024, January). *IPBES Invasive Alien Species Assessment: Summary for*

Policymakers (Version 3). Zenodo. https://doi.org/10.5281/zenodo.10521002

Seebens, H., Bacher, S., Blackburn, T. M., Capinha, C., Dawson, W., Dullinger, S.,

Genovesi, P., Hulme, P. E., van Kleunen, M., Kühn, I., Jeschke, J. M., Lenzner, B.,

Liebhold, A. M., Pattison, Z., Pergl, J., Pyšek, P., Winter, M., & Essl, F. (2021).

Projecting the continental accumulation of alien species through to 2050. *Global*

Change Biology, *27*(5), 970–982. https://doi.org/10.1111/gcb.15333

Seebens, H., Essl, F., & Blasius, B. (2017). The intermediate distance hypothesis of

biological invasions. *Ecology Letters*, *20*(2), 158–165.

Smith, D., King, R., & Allen, B. L. (2020). Impacts of exclusion fencing on target and non-

target fauna: A global review. *Biological Reviews*, *95*(6), 1590–1606.

https://doi.org/10.1111/brv.12631

Smith, R. K., Pullin, A. S., Stewart, G. B., & Sutherland, W. J. (2011). Is nest predator

exclusion an effective strategy for enhancing bird populations? *Biological*

Conservation, *144*(1), 1–10. https://doi.org/10.1016/j.biocon.2010.05.008

Spatz, D. R., Holmes, N. D., Will, D. J., Hein, S., Carter, Z. T., Fewster, R. M., Keitt, B.,

Genovesi, P., Samaniego, A., Croll, D. A., Tershy, B. R., & Russell, J. C. (2022). The

global contribution of invasive vertebrate eradication as a key island restoration tool.

Scientific Reports, *12*(1), 13391. https://doi.org/10.1038/s41598-022-14982-5

Van der Colff, D., Kumschick, S., Foden, W., & Wilson, J. R. U. (2021). Comparing the

IUCN's EICAT and Red List to improve assessments of the impact of biological

invasions. *NeoBiota*, *62*, 509–523. https://doi.org/10.3897/neobiota.62.52623

Vicente, J. R., Vaz, A. S., Roige, M., Winter, M., Lenzner, B., Clarke, D. A., & McGeoch, M.

A. (2022). Existing indicators do not adequately monitor progress toward meeting

invasive alien species targets. *Conservation Letters*, *15*(5), e12918.

https://doi.org/10.1111/conl.12918

Winter, M., Devictor, V., & Schweiger, O. (2013). Phylogenetic diversity and nature

conservation: Where are we? *Trends in Ecology & Evolution*, *28*(4), 199–204.

FIGURES

Figure 1. Workflow used to compute the Extinction Potential Metric. Species uniqueness was computed by rescaling ED2 scores from Gumbs et al. (2023); although they are indicated for all species here to show the general principle, ED2 scores were only available for native mammals. For impact magnitude, NAs were either considered as no impact (x 0) or randomly drawn as low or high impacts. Rug plots depict random draws along the quartic function for extinction probabilities and between lower and upper interquartiles from Gumbs et al. (2023) for ED2 scores.

Figure 2. Extinction Potential Metric (EPM) scores of invasive alien species (IAS) impacting native tetrapods globally. Only IAS with a score above 1 are displayed. Panel A shows EPM scores based on the impacts of IAS on all native tetrapods combined. Panel B shows EPM scores based on the impacts of IAS on each taxonomic group (amphibians, birds, mammals, and reptiles) separately.

Figure 3. Contribution of insular endemic species to the EPM scores. EPM scores
reflect the impact of IAS on native tetrapods globally, considering insular endemic species.
Panel A shows the change in EPM scores (in %) when excluding insular endemic species for
each of the four native taxonomic groups. Panel B highlights IAS with a difference in score of
more than 50% when excluding insular endemic specie

822
Figure 4. EPM scores of IAS impacting native mammals globally using textual and tabular data from the IUCN Red List. EPM scores were computed using both textual and tabular data from the IUCN Red List, including all impacts on native species, even when the severity or scope of the impact was not assessed.

835
Figure 5. Comparison of EPM-U and EPM scores of IAS according to rank. The lower an IAS's rank, the higher its score, and vice versa. Species close to the diagonal have similar ranks for both approaches. IAS whose absolute difference between the two metrics is greater than 50 are displayed with their scientific name.

Figure 6. EPM scores of threat mechanisms impacting native tetrapods globally. Only IAS with a score above 1 are displayed. Panel A shows EPM scores based on the impacts of mechanisms on all native tetrapods combined. Panel B shows EPM scores based on the impacts of mechanisms on each taxonomic group (amphibians, birds, mammals, and reptiles) separately.

846
Figure 7. IUCN categories of native tetrapods impacted by IAS globally and their
associated EPM scores. Only IAS with an EPM score above 1 are displayed. Panel A
shows EPM median measure (blue diamonds) and the number of native species per IUCN
category, based on the impacts of IAS on all native tetrapods combined. Panel B shows
EPM median measure (blue diamonds) and the number of native species per IUCN
category, based on the impacts of IAS on each taxonomic group (amphibians, birds,
mammals, and reptiles) separately.