The fate of pollen in two morphologically contrasting buzz-pollinated

Solanum flowers

Christian A. Vasquez-Castro [1,2,†]

Elodie Morel[1,3,†]

Bernardo Garcia-Simpson[1,4,†]*

Mario Vallejo-Marín[1]

[1] Department of Ecology and Genetics, Uppsala University, Uppsala, 752 36, Sweden

[2] Department of Biology, Faculty of Science, Universidad Nacional Agraria La Molina, Lima, 150 24, Peru

[3] Institute of Biology, University of Neuchâtel, 2000, Neuchâtel, Switzerland

[4] Círculo de Investigación en Taxonomía, Florística y Ecología Vegetal, Department of Biology, Faculty of

Science, Universidad Nacional Agraria La Molina, Lima, 150 24, Peru

*Author for correspondence. Email: bernicereus3000@gmail.com

† Co-first authors

23 Manuscript received _______; revision accepted _______

25 Running title: Pollen fates in buzz-pollinated *Solanum*

ABSTRACT

Study premise: Pollen transfer efficiency (PTE) and pollen deposition patterns on a pollinator's body significantly influence plant reproductive success. However, studies on pollen fates (i.e., the destination of pollen grains after being released) in animal-pollinated species, particularly those offering pollen as the sole reward, are limited. Here, we investigated pollen fates in two nectarless, buzz-pollinated *Solanum* species with contrasting floral morphology.

Methods: We conducted experimental trials involving one pollen donor and four recipient flowers of *Solanum rostratum* and *S. dulcamara*, using captive *Bombus terrestris* as pollinator. After each trial, we assessed the amount of pollen remaining in the anthers, deposited on stigmas, placed on the pollinator, and falling to the ground. We then estimated pollen fates and PTE, and modelled their pollen deposition curves.

Key results: We found that *S. rostratum* produced more pollen but had a more restricted pollen dispensing schedule compared to *S. dulcamara*. Although PTE was similar between species (0.72% vs. 1.07%, for *S. rostratum* and *S. dulcamara,* respectively), pollen loss mainly occurred due to falling to the ground in *S. rostratum* and bee grooming in *S. dulcamara*, potentially explained by their different floral architectures. Both species exhibited a typical exponential decay pattern in pollen deposition, with the first visited flowers receiving the most outcross pollen.

Conclusions: Our results suggest that PTE in pollen-rewarding, buzz-pollinated species is quantitatively similar to that in nectar-rewarding flowers with single pollen units (PTE = 1-2%), and that different buzz-pollinated flower architectures achieve similar PTE although through different pollen-loss pathways.

Keywords: Anther cone; buzz pollination; floral morphology; heteranthery; pollen carryover; pollen transfer and deposition; *Solanum* (Solanaceae)

INTRODUCTION

Pollen transfer from anthers to stigmas is a major determinant of plant fitness (Minnaar et al., 2019; Christopher et al., 2020). Plants produce a large amount of pollen per flower (Cruden, 2000), yet very few pollen grains produced in anthers ever make it to their destination. For instance, it is estimated that 2.4% of pollen grains produced by animal-pollinated plants with pollen grains released as single units (monads) reach the stigma of conspecific individuals (Harder and Thomson, 1989; Johnson and Harder, 2023). Pollen fates can be decomposed in three phases: pollen production and presentation, pollen

transfer, and pollen germination (Minnaar et al., 2019). Among these, pollen transfer, the pathway

between pollen release and pollen deposition, is highly stochastic and mediated by the behavior of

pollinators in animal-pollinated plants (Thomson and Plowright, 1980; Richards et al., 2009) (Figure 1).

However, despite its importance for plant reproductive success, studies of pollen transfer are scarce and

often focused on species with pollen packed in pollinia such as in orchids (Minnaar et al., 2019). The

pathway followed by pollen is especially hazardous for plants giving pollen as the only reward, as floral

visitors actively collect pollen grains during floral visitation and might further reduce the pool of pollen

remaining for fertilizing other flowers (Luo et al., 2008; Vallejo-Marín et al., 2009).

Pollen transfer efficiency (PTE) measures the proportion of pollen removed from the anthers

that successfully reaches conspecific stigmas and is therefore available for potential ovule fertilization

(Johnson and Harder, 2023). In animal-pollinated species, PTE usually decreases as the amount of pollen

released per visit increases (Harder and Thomson, 1989; Harder, 1990), giving plants with gradual pollen

release or pollen packaging strategies a fitness advantage. The Pollen Presentation Theory (Thomson,

2006) suggests that natural selection via male fitness should favor traits that regulate pollen dispensing

and/or restrict pollen removed per visit (Thomson, 2006).

In addition to PTE, a key characteristic affecting the fate of pollen and its consequences for plant

fitness is the amount of pollen deposited on each stigma in a consecutive sequence of recipient flowers,

i.e., the shape of its pollen deposition curve (pollen carryover curve; Thomson and Plowright 1980).

Pollen deposition curves tend to show an exponential decay pattern, but most studies have found

longer tails than expected under an exponential model (i.e., leptokurtic distribution) due to long

dispersal events (Lertzman, 1981; Price and Waser, 1982; Morris et al., 1994; Santa-Martinez et al.,

2021). The slope of the curve varies with pollinator type (Castellanos et al., 2003; Santa-Martinez et al.,

2021), type of flower, and reward availability in the recipient flower (Thomson and Plowright, 1980).

Deposition curves have been studied in hummingbird pollinated *Ipomopsis* and *Castilleja* (e.g., Lertzman,

1981; Price and Waser, 1982; Morris et al., 1994), and in bee-pollinated species (e.g., Thomson and

Plowright, 1980, Castellanos et al., 2003; Santa-Martinez et al., 2021). These studies have shown that

grooming behavior plays a crucial role in determining the steepness of the curve. However, pollen

deposition curves have rarely been estimated for nectarless plants that use pollen as the only reward to

floral visitors. Given that in these plants pollen serves both as a vehicle for male gametes and

simultaneously as a reward to potential pollinators (Vallejo-Marín et al., 2009), pollen deposition curves

could have a particularly steep slope if pollinators are efficient at grooming pollen following a visit.

Here, we characterize and compare the removal, dispersal, and transfer efficiency of pollen

(pollen fates) in two species of nectarless, bee-pollinated species of *Solanum* (Solanaceae). As most of

the other ~1,200 described *Solanum* species (Hilgenhof et al., 2023)*,* the species studied here are buzz

pollinated, i.e., they are pollinated by pollen-foraging bees that use vibrations to harvest pollen from

their tube-like poricidal anthers (Buchmann, 1983; Vallejo-Marín and Russell, 2024). To characterise

pollen fates in these two species, we exposed experimental flowers to visits by a generalist buzz-

pollinating bee, *Bombus terrestris* (Apidae). The specific objectives of our research were: (1) To quantify

pollen fates and pollen transfer efficiency of two *Solanum* species with contrasting size and floral

architecture (*Solanum rostratum* and *S. dulcamara*), and (2) to estimate the pollen deposition curves for

both species. Our study provides a starting point for establishing pollen fates in nectarless, buzz

pollinated species, and to investigate how floral architecture may influence pollen removal and transfer

efficiency.

MATERIALS AND METHODS

Plants

As most other *Solanum* species, the ones studied here have bisexual and nectarless flowers, with five

poricidal stamens (anthers that open through small apical pores). However, the species studied here

differ markedly in their stamen shape and architecture (i.e., the shape and arrangement of anthers

within the flower). *Solanum dulcamara* is a perennial vine native to Eurasia and introduced into North

America (Knapp et al., 2013). Its flowers are actinomorphic, relatively small (1.5-2 cm diameter) and

have monomorphic anthers that are fused into an anther cone that surrounds the centrally-placed pistil

(Figure 2A) (Glover et al., 2004). In Europe, *S. dulcamara* is visited by different species of bumblebees

(Free, 1970), it is self-compatible, but its mating system is unknown. Conversely, *S. rostratum* is an

annual herb native to Mexico and perhaps southern USA, and introduced in disturbed habitats around

the world (Zhao et al., 2013). Flowers are medium-sized (2.3-3.5 cm diameter) and have two

morphologically and functionally distinct sets of stamens in the same flower (i.e., heteranthery): one set

is composed of four smaller, bright yellow 'feeding' anthers centrally located, and the other set includes

a single larger, yellow to brown 'pollinating' anther that is laterally displaced, opposite to the style

(Figure 2B) (Vallejo-Marín et al., 2009). *Solanum rostratum* is also enantiostylous, i.e., each individual

produces flowers with the style deflected either to the right- or left-hand side; this type of enantiostyly

is thought to increase pollen export between flowers of reciprocal morphs (Jesson and Barrett, 2002). In

its native and introduced range, *S. rostratum* is visited by different types of bees, including bumblebees

(*Bombus spp.*) (Bowers, 1975; Solís-Montero et al., 2015). Genetic estimation of outcrossing rates,

indicates that *S. rostratum* is predominantly outcrossing (t = 0.70) in both native and introduced ranges

(Vallejo-Marín et al., 2013; Zhang et al., 2017). For our study, plants of *S. rostratum and S. dulcamara*

were grown from field-collected seeds at the glasshouses in Uppsala University. For *S. rostratum,* we

used seeds from accessions 10s71-5, 10s74, 10s77, 10s79, and 10s80 (San Miguel de Allende,

Guanajuato, Mexico), and 11s182, 11s188, 11s191, 11s205, and 11s207 (Dolores, Guanajuato, Mexico).

For *S. dulcamara,* we used seeds from accessions 22-UPP-1 and 22-UPP-2 collected at Uppsala. Seeds

were germinated as described in Vallejo-Marín et al. (2014), grown at 16h/20°C light and 8h/16°C dark

cycles, and fertilised with general purpose liquid fertiliser.

Pollinators

For evaluating pollen transfer, we used *Bombus terrestris*, a generalist, buzz-pollinating bumblebee that

is widely distributed in Europe and which is commonly used to supplement pollination of soft-fruit

crops. *Bombus terrestris* visits *S. dulcamara* in the native range and is similar in size and behaviour to

other bumblebees visiting *S. rostratum* in the Americas (Bowers, 1975; M. Vallejo-Marín, Uppsala

University, personal observation). We used individuals from two commercial colonies (Koppert

Biological Systems, Helsingborg, Sweden; and Biobest Group, Westerlo, Belgium). Colonies were

maintained on 2 M solution of sucrose available from artificial feeders within enclosed foraging arenas

(82×60×60 cm), follow Russell et al. (2017). Twice a week, bees were also provided with approximately

10g of "pollen dough" made of pulverised honeybee-collected pollen (Koppert Biological Systems) mixed

with sucrose solution and placed inside the colony. The foraging arenas were kept at room temperature

and illuminated with an LED ceiling panel set to a 14 h:10 h light:dark cycle. Before the experiment, we

conditioned the bees to visit *Solanum* flowers by allowing them to forage freely on flowers placed in the

foraging arena. This conditioning phase was never done on the same day of the experimental trials to

avoid bees carrying over pollen grain collected during the conditioning phase to the experimental trials.

Experimental trials

Before each trial, fresh flowers were collected from the greenhouse each morning before experiments

and carefully transported in floral foam (OASIS Floral Products, Washington, UK) within a closed plastic

box. Each trial consisted of one donor flower from which pollen could be removed by the bee, and four

recipients (R1-R4) that were experimentally prevented from releasing pollen by sealing the anther pores

using a small drop of PVA glue (Elmer's Products, Ohio, USA) under a stereoscopic microscope. A similar

amount of glue was applied just below the anther pore of the donor flowers as a sham treatment. After

the glue dried, we mounted each flower on a bamboo stick with Blu-Tack adhesive gum (Bostik,

Staffordshire, UK) and placed it in a plastic tube containing floral foam. A black plate was placed under

the donor flower to recover pollen falling from the flower (dislodged pollen). To control for differences

in pollen transfer within and between morphs of enantiostylous *S. rostratum*, we set all recipients to

have the reciprocal mirror-image morph of the donor flower (i.e., the left-hand side morph for all

recipients if the donor was a right-hand flower, and vice versa).

For each trial, one donor flower was placed in the centre of the foraging arena and a single bee

was allowed to enter the arena. Immediately after the bee landed on the donor flower, we placed the

first recipient flower (R1) next to it in the arena and removed the donor flower from the arena as soon

as the bee departed from it. We proceeded like so with the consecutive recipients (R2, R3 and R4). If

after 10 minutes the bee had not landed on the donor flower, we ended the trial and removed the bee.

All trials were video recorded using a camera (Canon EOS M50, Tokyo, Japan) placed right outside of the

arena. Recordings were used to calculate the duration of each flower visit. Temperature, humidity of the

arena and time of the day were recorded.

Pollen sampling and counting

After each trial, the stigmas of the four recipients were individually cut and placed on fuchsin jelly cubes

(32 mm3) (Kearns and Inouye, 1993). Dislodged pollen was collected from the black plate using 1-2

cubes of jelly, and anthers from the donor flower were cut and preserved in 1ml of 70% ethanol. Under

a stereoscopic microscope, we removed the pollen in the corbiculae of the bee and placed it on a jelly

cube and collected pollen on the body of the bee with a separate jelly cube. Each sample was stored in

individual 1.5 mL Eppendorf tubes. Each fuchsin jelly cube containing a stigma sample was mounted on a

slide and examined under a light microscope at 400x magnification for pollen grain counting. To process

the remaining pollen samples, 500 µL of 70% ethanol was added to tubes containing pollen from the

ground, corbiculae, and body. Donor anthers were subjected to sonication using an Elmasonic S40H

ultrasonic cleaner (Elma Schmidbauer GmbH, Singen, Germany), followed by staining pollen grains with

200 µL of lactoglycerol-aniline blue (Kearns and Inouye, 1993). For pollen grain counting, an aliquot was

taken from each tube and placed onto a hemocytometer chamber until 200 pollen grains were counted.

To estimate the number of pollen grains, we considered the combined volumes of ethanol, jelly, and

aniline used in their respective cases. After the trial concluded, we estimated: (C) the number of pollen

grains remaining of the donor anthers, (D) number of pollen grains that fell on the ground, (F) the

number of pollen grains on the bee's corbiculae and (G) on the bee's body, and (H) the number of pollen

grains deposited on the stigma of each recipient. From these variables we also calculated the following

derived variables: total pollen production (A = D + F + G + H), pollen released (B = F + G + H + D), pollen

collected (E = F + G + H), and pollen lost from the pollination process (I = D + F).

Statistical analyses

Proportion of pollen grains deposited on the body or the stigma were calculated from the total pollen

production and from the amount of pollen released. To determine if the proportions of pollen grains

deposited on body and stigmas varied between species, we first verified the assumptions of normality

and homoscedasticity using the Shapiro-Wilk and Levene's tests from *stats* (R Core Team, 2023) and *car*

(Fox and Weisberg, 2019) packages, respectively. These assumptions were tested using absolute values

and the log transformations from both species separately. In all cases, at least one of the assumptions

was not met. Thus, we compared each variable using a non-parametric Wilcoxon test (R Core Team,

2023).

Pollen deposition curves (Thomson and Plowright, 1980) were analyzed using Generalized Linear

Mixed-effects Models (GLMM) with a Poisson error (Log-link) using the *glmer* function (*lme4*) (Bates et

al., 2014). First, we built a model using the number of pollen grains deposited in each recipient flower

was the response variable, recipient order (R1-R4), species, and visit time were considered as fixed

effects, and trial number as a random effect. Second, we analyzed a similar GLMM model as above but

adding the number of removed pollen grains as an offset variable. This allowed us to compare pollen

deposition curves independently of the absolute number of pollen grains removed from the donor. Back

transformation and visualization were conducted using the *sjPlot* package (Lüdecke et al., 2018). We

assessed overdispersion and zero inflation using the *DHARMa* package (Hartig, 2018). All statistical

analyses were done in R v.4.1.0 (R Core Team, 2021) using *RStudio* (RStudio Team, 2021).

RESULTS

Pollen fates and pollen transfer efficiency

The absolute numbers of pollen grains in each stage of pollen transfer are presented in Figure 3. We found that *Solanum rostratum* flowers produce on average 27,59% more pollen grains per flower than *S. dulcamara* flowers (Figure 4). However, after a single visit, a bee removed on average 37.5% of the flower's total production of pollen grains in *S. dulcamara*, and only 8.91% in a *S. rostratum*. This is equivalent to the removal of about four times as many pollen grains in *S. dulcamara* than in *S. rostratum* in absolute numbers (Table 1)*.* From the pollen released by the flowers, between 19.12% (*S. dulcamara*) and 67.93% (*S. rostratum*) fell to the ground. In addition to the pollen wasted on the ground, bees collected in their corbiculae, on average, 65.82% of released pollen grains for *S. dulcamara* and 24.03% for *S. rostratum*. The stigmas of recipient flowers received $1\ 860 \pm 370$ (mean $\pm$ se; *S. dulcamara*) and 356 ± 93 (*S. rostratum*) pollen grains. Thus, only approximately 8.03% (*S. rostratum*) to 15.06% (*S. dulcamara*) of released pollen grains remain available for pollination (pollen on stigmas and body) (Table 1).

The pollen transfer efficiency (PTE; i.e., the percentage of released pollen grains that reach recipient's stigmas) was 1.07% for *S. dulcamara* and 0.72% for *S. rostratum* (Figure 5). The percentage of pollen remaining on the body was not significantly different between the two species N=19, U=44, P = 0.968). Most of the pollen was lost on the ground (67.93% in *S. rostratum*) or in the corbiculae (65.82% in *S. dulcamara*) (Figure 5). There was a significant difference in the percentage of pollen lost in the corbiculae and the dislodged pollen between species (N=19, U = 77, P = 0.008; N=19, U = 5, P < 0.001, respectively). However, the percentage of all the pollen lost (from released pollen) was not significantly different between both species (N=19, U = 43, P = 0.9).

Pollen deposition curves

As expected, most pollen from the donor was deposited in the first recipient flower visited, strongly decreasing with the increase of recipient order (Figure 6 and 7B; Table 2, Z = -43.19, P < 0.001 and Z = -42.53, P < 0.001, for analyses with and without using pollen released as an offset, respectively). We found a statistically significant difference in the average number of pollen grains per recipient stigma between *Solanum* species with fewer pollen grains deposited on *S. rostratum* than in *S. dulcamara* (Figure 6; Table 2, Z = -3.65, P < 0.001). However, our analysis of the relative rates of pollen deposition

(using the number of pollen grains released as an offset variable) suggests that both species deposit

approximately the same proportion of pollen grains (Figure 7B; Table 2, Z = 0.01, P = 0.996). The

relationship between recipient order and pollen deposited differed between species, with *S. rostratum*

exhibiting a steeper pollen deposition curve, as indicated by the statistically significant interaction

between recipient order and flower species (Table 2, Z = -7.23, P < 0.001 and Z = -7.89, P < 0.001, for

analyses with and without using pollen released as an offset, respectively). We also found that longer

visits resulted in more pollen deposited per stigma (Figure 7A; Table 2, Z = 30.24, P < 0.001 and Z =

30.02, P < 0.001, with and without pollen released as an offset, respectively).

DISCUSSION

Our study's main findings can be summarised as: (1) Both buzz-pollinated species of *Solanum* studied

here have a similar or lower values of pollen transfer efficiency (PTE) as those reported for other non-

buzz pollinated plants that disperse pollen in individual pollen grain units (monads), (2) pollen

deposition curves follow a classic exponential decay pattern, as in other animal-pollinated plants, and

(3) although both *Solanum* species studied showed a similar percentage of pollen loss, the pathway to

this loss differs and is either lost on the ground (*S. rostratum*) or in the bees' corbiculae (*S. dulcamara*),

potentially reflecting the functional effects of different flower architectures.

Pollen transfer efficiency in buzz-pollinated flowers

Buzz pollinated species are thought to produce large numbers of pollen grains compared to nectar-

producing relatives as a mechanism to offset the costs of active pollen collection by floral visitors (Vogel,

1978; De Luca and Vallejo-Marín, 2013; Vallejo-Marín and Russell, 2024). The average pollen production

per flower is indeed high in *S. dulcamara* (>600,000) and *S. rostratum* (>1,000,000), and comparable to

other nectarless plant species (Vallejo-Marín et al., 2009; Vallejo-Marín et al., 2014; Vallejo-Marín et al.,

2022). In our experiment, visitation by a generalist, buzz-pollinating bumblebee resulted in a pollen

transfer efficiency (PTE) of between 0.72 and 1.07%, which is towards the lower end of the 1-2% PTE

range reported for other species with monads (Johnson and Harder, 2023). Pollen transfer efficiency has

only been rarely quantified for buzz pollinated plants, but low PTE characterizes two previously studied

species. *Rhexia virginica* (Melastomataceae), an herbaceous species distributed in eastern North

America, has a PTE of 0.92% (Supplementary material from Johnson and Harder, 2023), while

Senna reticulata a neotropical tree, has a much lower PTE (0.05%) (Supplementary material from

Johnson and Harder, 2023). These relatively low values of PTE in buzz pollinated species might suggest

that active pollen collection indeed impacts the realized efficiency of pollen transfer, and that higher

rates of total pollen production are selected for to ensure delivery of sufficiently high numbers of pollen

grains to fertilize all ovules. However, further work comparing total pollen production and PTE in buzz

pollinated taxa and their closest non-buzz pollinated relatives are required to establish if this is a general

pattern. Furthermore, PTE can vary among different species of floral visitors, and this is an area that

requires additional study.

Although both *Solanum* species studied here achieve a similar PTE, the pathways of pollen

production, release, and loss differ (Figure 5). Percentage of pollen released in other buzz pollinated

species ranges widely from 10-60% (*Rhexia virginica*: 10–43%; Larson and Barrett, 1999;

Senna reticulata: 25%; Snow and Roubik, 1987; *Erica tetralix* and *Vaccinum myrtillus:* 28–60%; Moquet

et al., 2017), and is similar or lower than in non-buzz-pollinated species such as *Aconitum septentrionale*

(44-61%; Thøstesen and Olesen, 1996), *Malus domestica* and *Prunus dulcis* (40-50%; Thomson and

Goodell, 2001), and *Calluna vulgaris* (≤ 88%; Moquet et al., 2017). The small-flowered *S. dulcamara*

produces fewer pollen grains than *S. rostratum* but releases a higher number (and percentage) of pollen

grains after a single visit by a bumblebee. If all pollen produced in the flower is available for removal,

about 11 visits would be required to empty a flower of *S. rostratum* (8.9% of total pollen removed per

visit), while in *S. dulcamara* fewer than three visits would be needed (37.5% pollen released per visit).

However, pollen inside the anther may not all be available to be removed at first, and it appears that

some buzz pollinated flowers present a staggered release of pollen as the flower ages (Harder and

Barclay, 1994; Kemp and Vallejo-Marín, 2021). The extent to which pollen becomes available

simultaneously or in a staggered fashion in buzz pollinated species is not well understood. Regardless of

the mechanism, the high rates of pollen release observed in *S. dulcamara,* characterize other small-

flowered, non-heterantherous taxa in *Solanum* Section *Androceras* and this has been interpreted as a

consequence to a shift of mating system towards more selfing (Kemp and Vallejo-Marín, 2021). The

mating system of *S. dulcamara* is presently unknown, but this self-compatible species sometimes sets

fruit autonomously (M. Vallejo-Marín, Uppsala University, personal observation). In contrast, although

S. rostratum is also self-compatible, it has a high outcrossing rate (0.70-0.90) (Vallejo-Marín et al., 2013;

Morra-Carrera et al., 2019) and rarely sets fruits autonomously, at least in the native range. Thus, the

difference in the proportion of pollen removed per visit could be associated to different floral

morphologies and mating systems in *S. dulcamara* and *S. rostratum*. For outcrossing species visited

regularly by pollinators, a more restricted pollen schedule might allow them to dispense pollen more

efficiently or to more flowers (Harder and Thomson, 1989; Harder and Wilson, 1994; Kemp and Vallejo-

Marín, 2021).

Although both species end up exporting similar proportion of pollen grains to stigmas, the

pathway to pollen loss differs: In *S. dulcamara*, pollen is mostly loss to bees collecting it and placing it in

their corbiculae, while in *S. rostratum,* pollen is mostly loss by falling on to the ground, meaning that this

removed pollen does not contribute to the fitness of either bee or plant. What is the reason for these

contrasting pollen fates in these two species? With only two species studied, it is difficult to generalize,

but in *S. dulcamara* the bees can curl their body in close proximity of the anther pores while buzzing the

flower. The fused anther cone of *S. dulcamara* then works as a single pollen dispensing unit that places

pollen grains in the ventral region of the bee where it can be groomed and placed in the corbiculae

(Nevard and Vallejo-Marín, 2022). In contrast, in heterantherous *S. rostratum*, the bee cannot work

simultaneously both feeding and pollinating anthers, and it usually focuses its attention on the feeding

anthers. The feeding anthers contact the bee on the ventral side, while the pollinating anthers contact

the bee on the dorsal side of the abdomen, where it is harder to groom by the bee. The body of the bee

often contacts the terminal pores of the feeding anthers, but the distance between the bee's body and

the terminal pore of the pollinating anther depends both on the positioning of the bee and on the

relative size of bee and flower. Because approximately 50% of the total pollen production in these

flowers resides on the single pollinating anther (Vallejo-Marín et al., 2009; Vallejo-Marín et al., 2014),

flower handling and bee-flower size matching can then affect significantly whether the pollen reaches

the bee's body or falls to the ground. Although we did not quantify this in our experiment, we expect

that much of the pollen loss occurring during buzz pollination in *S. rostratum* is due to mismatches

between the size of the bee and the flower. All else being equal, a bee of similar characteristics to the

ones studied here might benefit more from a single visit to a species like *S. dulcamara* than one like

S. rostratum. Of course, differences in total pollen production, and number of flowers per plant might

help offset the increased difficulty of gathering pollen from flowers with a more complex morphology as

in *S. rostratum*.

Pollen deposition curves

Our results demonstrate that the pollen deposition curves (i.e., the pattern of pollen deposition on the

stigmas of consecutive recipients) for the buzz-pollinated *Solanum* species studied follow a typical

exponential decay pattern, as reported for other species (Thomson and Plowright, 1980; Mitchell et al.,

2013; Santa-Martinez et al., 2021). Specifically, the amount of pollen deposited on stigmas decreased by

27.6% and 36.9% with each successive recipient for *S. dulcamara* and *S. rostratum*, respectively. These

percentages are on the higher end of the spectrum for pollen carryover curves, which ranges from 50%

to 0.5% (see Table 1 in Robertson, 1992). This indicates a relatively steep decline in deposited pollen

with each successive recipient. Generally, grooming pollinators such as bees exhibit steeper deposition

curves, indicating less extensive pollen carryover, compared to pollinators like hummingbirds and

hawkmoths (Robertson, 1992; Castellanos, 2003; Mitchell et al., 2013). However, this pattern has not

been studied in other buzz-pollinated plants, highlighting a gap in our understanding of these systems.

Pollen deposition curves are influenced by both the amount of pollen that lands on the

pollinator's body and the intensity of its grooming before visiting the next flower. Buzz-pollinated

Solanum flowers, which produce and gradually release large amounts of pollen as a reward for

pollinators, tend to encourage more intense grooming behavior compared to nectar-producing, non-

buzz-pollinated flowers (Harder, 1990; Johnson and Harder, 2023). This intense grooming results in

steeper pollen deposition curves (Holmquist et al., 2012), potentially influencing gene flow patterns

(Santa-Martinez et al., 2021). Consequently, if buzz-pollinated flowers generally exhibit steeper pollen

deposition curves, they might experience more restricted gene flow compared to non-buzz-pollinated

species offering other rewards, such as nectar. While insect-pollinated species often display higher levels

of population genetic structure (Dellinger et al., 2022; Gamba and Muchhala, 2020; Gamba and

Muchhala, 2022), the specific impact of buzz pollination on genetic structure remains underexplored

CONCLUSIONS

Empirical studies of pollen fates, although experimentally complex and time-consuming, provide an

opportunity to illuminate a key process affecting the reproductive success of plants. Our study found

significant differences in pollen fates between *S. dulcamara* and *S. rostratum*, highlighting the impact of

floral architecture on pollen transfer, even among species visited by the same buzz-pollinating bee.

Specifically, we observed distinct pollen deposition curves and pathways of pollen loss between species.

S. rostratum primarily loses pollen through dislodging due to its restricted dispensing schedule, while *S.*

dulcamara experiences significant pollen loss from bee grooming. Despite these differences, both

species achieve comparable pollen transfer efficiency (PTE). These findings highlight the role of floral

morphology on pollen transfer dynamics and suggest that different floral architectures can lead to

similar reproductive outcomes. Future research should encompass a wider range of species with diverse

morphological traits and reward types and explore how different pollen pathways affect realized female

and male fitness.

ACKNOWLEDGMENTS

356 We thank D. Scaccabarozzi and L. van Kolfschoten for their technical support and advice throughout the

357 development of this work. We are also grateful to the Research Experience for Peruvian Undergraduates

358 (REPU) Program and the University of Lausanne exchange program for enabling A.V.C., E.M., and B.G.S.

359 to carry out this project at Uppsala University. This work was partly supported by start-up funds from

360 the Plant Ecology and Evolution program at Uppsala University, awarded to M.V.M.

AUTHOR CONTRIBUTIONS

362 A.V.C., E.M., B.G.S., and M.V.M conceived the study. A.V.C., E.M. and B.G.S. conducted the experiments,

363 collected the data and performed the analyses. A.V.C., E.M., B.G.S., and M.V.M wrote and revised the

364 manuscript. M.V.M supervised the study and acquired funding.

DATA AVAILABILITY STATEMENT

366 Pollen transfer data used for the analyses and the R script will be made available upon acceptance.

LITERATURE CITED

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using

lme4. *Journal of Statistical Software* 67: 1–48. https://doi.org/10.18637/jss.v067.i01

Bowers, K. A. 1975. The pollination ecology of Solanum rostratum (Solanaceae). *American Journal of*

Botany 62: 633-638. https://doi.org/10.1002/j.1537-2197.1975.tb14094.x

Buchmann, S. L. 1983. Buzz pollination in angiosperms. *In* C. E. Jones and R. J. Little [eds.], Handbook of

experimental pollination biology, 73-113. Scientific and Academic Editions, New York, USA.

Castellanos, M. C., P. Wilson, and J. D. Thomson. 2003. Pollen transfer by hummingbirds and

bumblebees, and the divergence of pollination modes in *Penstemon*. *Evolution* 57: 2742–2752.

https://doi.org/10.1111/j.0014-3820.2003.tb01516.x

Christopher, D. A., R. J. Mitchell, and J. D. Karron. 2020. Pollination intensity and paternity in flowering

plants. *Annals of Botany* 125: 1-9. https://doi.org/10.1093/aob/mcz159

Cruden, R. W. 2000. Pollen grains: Why so many? *In* A. Dafni, M. Hesse, and E. Pacini [eds.], Pollen and

Pollination, 143–165. Springer, Berlin, Germany. https://doi.org/10.1007/978-3-7091-6306-1_8

De Luca, P. A., and M. Vallejo-Marín. 2013. What's the 'buzz' about? The ecology and evolutionary

significance of buzz-pollination. *Current Opinion in Plant Biology* 16: 429–435.

https://doi.org/10.1016/j.pbi.2013.05.002

Dellinger, A. S., O. Paun, J. Baar, E. M. Temsch, D. Fernández-Fernández, and J. Schönenberger. 2022.

Population structure in Neotropical plants: Integrating pollination biology, topography and climatic

niches. *Molecular Ecology* 31: 2264-2280. https://doi.org/10.1111/mec.16403

Fox, J., and S. Weisberg. 2019. car: Companion to Applied Regression. R package version 3.1-2. Website:

https://CRAN.R-project.org/package=car [accessed 09 September 2024].

Free, J. B. 1970. The Flower Constancy of Bumblebees. *Journal of Animal Ecology* 39: 395–402.

https://doi.org/10.2307/2978

Gamba, D., and N. Muchhala. 2020. Global patterns of population genetic differentiation in seed plants.

Molecular Ecology 29: 3413–3428. https://doi.org/10.1111/mec.15575

Gamba, D., and N. Muchhala. 2023. Pollinator type strongly impacts gene flow within and among plant

populations for six Neotropical species. *Ecology 104*: e3845. https://doi.org/10.1002/ecy.3845

Glover, B. J., S. Bunnewell, and C. Martin. 2004. Convergent evolution within the genus *Solanum*: The

specialised anther cone develops through alternative pathways. *Gene* 331: 1–7.

https://doi.org/10.1016/j.gene.2004.01.027

Harder, L. D., and R. M. R. Barclay. 1994. The functional significance of poricidal anthers and buzz

pollination: controlled pollen removal from *Dodecatheon*. *Functional Ecology* 8: 509-517.

https://doi.org/10.2307/2390076

Harder, L. D. 1990. Behavioral responses by bumble bees to variation in pollen availability. *Oecologia* 85:

41–47. https://doi.org/10.1007/BF00317341

Harder, L. D., and W. G. Wilson. 1994. Floral evolution and male reproductive success: Optimal

dispensing schedules for pollen dispersal by animal-pollinated plants. *Evolutionary Ecology* 8: 542–

559. https://doi.org/10.1007/BF01238257

Harder, L. D., and W. G. Wilson. 1998. Theoretical consequences of heterogeneous transport conditions

for pollen dispersal by animals. *Ecology* 79: 2789–2807. https://doi.org/10.1890/0012-

9658(1998)079[2789:TCOHTC]2.0.CO;2

Harder, L. D., and J. D. Thomson 1989. Evolutionary Options for Maximizing Pollen Dispersal of Animal-

Pollinated Plants. *The American Naturalist* 133: 323–344. https://doi.org/10.1086/284922

Hartig, F. 2022. DHARMa: residual diagnostics for hierarchical (multi-Level / mixed) regression models. R

package version 0.4.6. Website: https://CRAN.R-project.org/package=DHARMa [accessed 09

September 2024].

Hilgenhof, R., E. Gagnon, S. Knapp, X. Aubriot, E. J. Tepe, L. Bohs, L. L. Giacomin, et al. 2023.

Morphological trait evolution in *Solanum* (Solanaceae): Evolutionary lability of key taxonomic

characters. *Taxon* 72: 811–847. https://doi.org/10.1002/tax.12990

Holmquist, K. G., R. J. Mitchel, and J. D. Karron. 2012. Influence of pollinator grooming on pollen-

mediated gene dispersal in *Mimulus ringens* (Phrymaceae). *Plant Species Biology* 27: 77–85.

https://doi.org/10.1111/j.1442-1984.2011.00329.x

Jesson, L. K., and S. C. H. Barrett. 2002. Solving the puzzle of mirror-image flowers. *Nature 417*: 707–707.

https://doi.org/10.1038/417707a

Johnson, S. D., and L. D. Harder. 2023. The economy of pollen dispersal in flowering plants. *Proceedings*

of the Royal Society B: Biological Sciences 290: 20231148. https://doi.org/10.1098/rspb.2023.1148

Kearns, C. A., and D. W. Inouye. 1993. Techniques for pollination biologists. University press of Colorado,

Boulder, Colorado, USA.

Kemp, J. E., and M. Vallejo-Marín. 2021. Pollen dispensing schedules in buzz-pollinated plants:

Experimental comparison of species with contrasting floral morphologies. *American Journal of*

Botany 108: 993–1005. https://doi.org/10.1002/ajb2.1680

Knapp, S. 2013. A revision of the Dulcamaroid Clade of *Solanum* L. (Solanaceae). *PhytoKeys 22*: 1–428.

https://doi.org/10.3897/phytokeys.22.4041

Larson, B. M., and S. C. Barrett. 1999. The ecology of pollen limitation in buzz-pollinated *Rhexia virginica*

(Melastomataceae). *Journal of Ecology 87*: 371-381. https://doi.org/10.1046/j.1365-

2745.1999.00362.x

Lertzman, K. P. 1981. Pollen transfer: processes and consequences. University of British Columbia,

Vancouver, Canada.

Lüdecke, D., A. Bartel, C. Schwemmer, C. Powell, A. Djalovski, and J. Titz. 2023. sjPlot: data visualization

for statistics in social science. R package version 2.8.14. Website: https://CRAN.R-

project.org/package=sjPlot [accessed 09 September 2024].

Luo, Z., D. Zhang, and S. S. Renner. 2008. Why two kinds of stamens in buzz-pollinated flowers?

Experimental support for Darwin's division-of-labour hypothesis. *Functional Ecology* 22: 794–800.

https://doi.org/10.1111/j.1365-2435.2008.01444.x

Minnaar, C., B. Anderson, M. L. de Jager, and J. D. Karron. 2019. Plant–pollinator interactions along the

pathway to paternity. *Annals of Botany* 123: 225–245. https://doi.org/10.1093/aob/mcy167

Mitchell, R. J., W. G. Wilson, K. G. Holmquist, and J. D. Karron. 2013. Influence of pollen transport

dynamics on sire profiles and multiple paternity in flowering plants. *PLoS One* 8: e76312.

https://doi.org/10.1371/journal.pone.0076312

Moquet, L., L. Bruyère, B. Pirard, and A. L. Jacquemart. 2017. Nectar foragers contribute to the

pollination of buzz-pollinated plant species. *American Journal of Botany* 104: 1451–1463.

https://doi.org/10.3732/ajb.1700090

Mora-Carrera, E., M. Castañeda-Zárate, J. Fornoni, K. Boege, and C. A. Domínguez. 2019. On the

adaptive value of monomorphic versus dimorphic enantiostyly in *Solanum rostratum*. *Annals of*

Botany 123: 205–212. https://doi.org/10.1093/aob/mcy162

Morris, W. F., M. V. Price, N. M. Waser, J. D. Thomson, B. Thomson, and D. A. Stratton. 1994. Systematic

increase in pollen carryover and its consequences for geitonogamy in plant populations. *Oikos* 71:

431–440. https://doi.org/10.2307/3545831

Nevard, L., and M. Vallejo-Marín. 2022. Floral orientation affects outcross-pollen deposition in buzz-

pollinated flowers with bilateral symmetry. *American Journal of Botany* 109: 1568–1578.

https://doi.org/10.1002/ajb2.16078

Price, M. V., and N. M. Waser. 1982. Experimental studies of pollen carryover: Hummingbirds and

Ipomopsis aggregata. *Oecologia* 54. 353–358. https://doi.org/10.1007/BF00380004

Richards, S. A., N. M. Williams, and L. D. Harder. 2009. Variation in Pollination: Causes and

Consequences for Plant Reproduction. *The American Naturalist* 174: 382–398.

https://doi.org/10.1086/603626

Robertson, A. W. 1992. The relationship between floral display size, pollen carryover and geitonogamy

in *Myosotis colensoi* (Kirk) Macbride (Boraginaceae). *Biological Journal of the Linnean Society* 46:

333-349. https://doi.org/10.1111/j.1095-8312.1992.tb00868.x

Russell, A. L., S. L. Buchmann, and D. R. Papaj. 2017. How a generalist bee achieves high efficiency of

pollen collection on diverse floral resources. *Behavioral Ecology* 28: 991–1003.

https://doi.org/10.1093/beheco/arx058

Santa-Martinez, E., C. Cardoso Castro, A. Flick, M. Sullivan, H. Riday, M. K. Clayton, and J. Brunet. 2021.

Bee species visiting *Medicago sativa* differ in pollen deposition curves with consequences for gene

flow. *American Journal of Botany* 108: 1016–1028. https://doi.org/10.1002/ajb2.1683

Snow, A. A., and D. W. Roubik. 1987. Pollen Deposition and Removal by Bees Visiting Two Tree Species in

Panama. *Biotropica* 19: 57–63. https://doi.org/10.2307/2388460

Solís-Montero, L., C. H. Vergara, and M. Vallejo-Marín. 2015. High incidence of pollen theft in natural

populations of a buzz-pollinated plant. *Arthropod-Plant Interactions 9*: 599–611.

https://doi.org/10.1007/s11829-015-9397-5

Thomson, J. D. 2006. Tactics for male reproductive success in plants: Contrasting insights of sex

allocation theory and pollen presentation theory. *Integrative and Comparative Biology* 46: 390–

397. https://doi.org/10.1093/icb/icj046

Thomson, J. D., and K. Goodell. 2001. Pollen Removal and Deposition by Honeybee and Bumblebee

Visitors to Apple and Almond Flowers. *Journal of Applied Ecology* 38: 1032-1044.

https://www.jstor.org/stable/827241

Thomson, J. D., and R. C. Plowright. 1980. Pollen carryover, nectar rewards, and pollinator behavior with

special reference to *Diervilla lonicera*. *Oecologia* 46: 68–74. https://doi.org/10.1007/BF00346968

Thøstesen, A. M., and J. M. Olesen. 1996. Pollen Removal and Deposition by Specialist and Generalist

Bumblebees in *Aconitum septentrionale*. *Oikos* 77: 77–84. https://doi.org/10.2307/3545587

Vallejo-Marín, M., J. S. Manson, J. D. Thomson, and S. C. H. Barrett. 2009. Division of labour within

flowers: Heteranthery, a floral strategy to reconcile contrasting pollen fates. *Journal of Evolutionary*

Biology 22: 828–839. https://doi.org/10.1111/j.1420-9101.2009.01693.x

Vallejo-Marín, M., C. E. Pereira Nunes, and A. L. Russell. 2022. Anther cones increase pollen release in

buzz-pollinated *Solanum* flowers. *Evolution* 76: 931–945. https://doi.org/10.1111/evo.14485

Vallejo-Marín, M., and A. L. Russell. 2024. Harvesting pollen with vibrations: towards an integrative

understanding of the proximate and ultimate reasons for buzz pollination. *Annals of Botany* 133:

379–398. https://doi.org/10.1093/aob/mcad189

Vallejo-Marín, M., L. Solís-Montero, D. Souto Vilaros, and M. Y. Q. Lee. 2013. Mating system in Mexican

populations of the annual herb *Solanum rostratum* Dunal (Solanaceae). *Plant Biology* 15: 948–954.

https://doi.org/10.1111/j.1438-8677.2012.00715.x

Vallejo-Marín, M., C. Walker, P. Friston-Reilly,L. Solís-Montero, and B. Igic. 2014. Recurrent modification

of floral morphology in heterantherous Solanum reveals a parallel shift in reproductive strategy.

Philosophical Transactions of the Royal Society B: Biological Sciences 369: 20130256.

https://doi.org/10.1098/rstb.2013.0256

Vogel, S. 1978. Evolutionary shifts from reward to deception in pollen flowers. *In* A. J. Richards [eds], The

pollination of flowers by insects, 89-96. Academic Press for the Linnean Society of London, London,

UK.

Zhang, L., L. Yu, and A. Lou. 2017. No evolutionary change in the mating system of Solanum rostratum

(Solanaceae) during its invasion in China. *Scientific Reports* 7: 17698.

https://doi.org/10.1038/s41598-017-17881-2

Zhao, J., L. Solís-Montero, A. Lou, and M. Vallejo-Marín. 2013. Population Structure and Genetic

Diversity of Native and Invasive Populations of *Solanum rostratum* (Solanaceae). *PLOS ONE* 8:

e79807. https://doi.org/10.1371/journal.pone.0079807

 TABLES

513 **Table 1**. Estimated number of pollen grains (mean $\pm$ se) at different stages of the pollination process in

514 *Solanum dulcamara* and *S. rostratum*.

	S. dulcamara		*S. rostratum*	
	N	Number of pollen grains	N	Number of pollen grains
Total pollen production (A)	9	641 780 $\pm$ 92 979	10	1 065 712 $\pm$ 116 142
Pollen release phase				
Pollen released (B)	9	243 323 $\pm$ 49 331	10	86 732 $\pm$ 21 423
Remaining pollen (C)	10	378 642 $\pm$ 65 596	10	1 000 891 $\pm$ 121 491
Pollen transfer phase				
Pollen on ground/dislodged (D)	9	41 177 + 19 734	10	53 561 + 17 369
Pollen collected (E)	9	202 146 + 51 589	10	29 358 + 9 222
Pollen on the pollinator				
Pollen in corbiculae (F)	9	188 601 $\pm$ 53 757	10	24 121 $\pm$ 8 634
Pollen on body (G)	9	11 685 + 4466	10	4 892 + 2198
Pollen deposition phase				
Pollen deposited on stigmas (H)	9	1860 + 370	10	356 + 93
Pollen loss				
Pollen on corbiculae and on the ground (I)	9	229 779 + 51 778	10	81 495 + 21 477

516 **Table 2.** Statistical analysis of the effect of recipient order, flower species (*S. dulcamara, S. rostratum*)

517 and duration of visit (s) on the number of pollen grains deposited on stigmas. Model estimates were

518 obtained from GLMMs with trial number as a random effect and recipient order, flower species and

519 duration of visit as fixed effects. (A) N = 73, residual df = 67; (B) N = 66, residual df = 60.

Fixed effects	Coefficient	Standard error	Effect size (Z)	p-value
(A) Pollen deposition model				
Intercept	5.801	± 0.205	28.33	<0.001
Recipient order	-0.303	± 0.007	-42.53	<0.001
Flower species (*S. rostratum*)	-1.053	± 0.289	-3.65	<0.001
Log (duration of visit (s))	0.324	± 0.011	30.02	<0.001
Recipient order: flower species (*S. rostratum*)	-0.147	± 0.019	-7.89	<0.001
(B) Pollen deposition model with offset variable				
Intercept	-6.248	± 0.317	-19.67	<0.001
Recipient order	-0.323	± 0.007	-43.19	<0.001
Flower species (*S. rostratum)*	0.002	± 0.449	0.01	0.996
Log (duration of visit (s))	0.332	± 0.011	30.24	<0.001
Recipient order: flower species (*S. rostratum*)	-0.138	± 0.019	-7.24	<0.002

FIGURE LEGENDS

Figure 1. Pathways depicting the fates of pollen in a bee pollinated species such as the ones studied here. The diagram shows pollen contained inside the anthers (yellow); pollen released that can be potentially transferred to conspecific stigmas (blue); pollen lost from the fertilization process, including dislodged pollen that falls to the ground and pollen stored by the bee (e.g., in corbiculae) and thus unable to contribute to fertilization (red); and pollen deposited on stigmas (green).

Figure 2. (A) Flower of *S. dulcamara*, arrow indicates fused anthers (anther cone). (B) Flower of *S. rostratum*, arrows indicate the feeding (above) and pollinating anthers (below). Photo credits: M. Vallejo-Marín.

Figure 3. Diagram showing the fates of pollen of (A) *S. rostratum* and (B) *S. dulcamara*. The number of pollen grains for each stage of the pollen dispersal process is specified (mean ± SE).

Figure 4. Percentage (from the total pollen production) of pollen in different paths after one visit on a donor flower in *S. dulcamara* and *S. rostratum*.

Figure 5. Percentage (from the amount of pollen released from the anthers) in different paths after one visit on a donor flower in *S. dulcamara* and *S. rostratum*.

Figure 6. Number of pollen grains deposited on recipient flower's stigmas as influenced by flower species and recipient order. Plotted lines indicate estimated means and shaded regions indicate standard errors.

Figure 7. Number of pollen grains deposited on recipient flower stigmas as influenced by (A) duration of the visit in seconds, and (B) flower species and recipient order with released pollen as offset variable. Plotted lines indicate estimated means and shaded regions indicate standard

FIGURES

Figure 1. Pathways depicting the fates of pollen in a bee pollinated species such as the ones studied here. The diagram shows pollen contained inside the anthers (yellow); pollen released that can be potentially transferred to conspecific stigmas (blue); pollen lost from the fertilization process, including dislodged pollen that falls to the ground and pollen stored by the bee (e.g., in corbiculae) and thus unable to contribute to fertilization (red); and pollen deposited on stigmas (green).

551 **Figure 2**. (A) Flower of *S. dulcamara*, arrow indicates fused anthers (anther cone). (B) Flower of *S.*

552 *rostratum*, arrows indicate the feeding (above) and pollinating anthers (below). Photo credits: M.

553 Vallejo-Marín.

Figure 3. Diagram showing the fates of pollen of (A) *S. rostratum* and (B) *S. dulcamara*. The number of pollen grains for each stage of the pollen dispersal process is specified (mean ± SE).

559   **Figure 4.** Percentage (from the total pollen production) of pollen in different paths after one visit on a

560   donor flower in *S. dulcamara* and *S. rostratum*.

563   **Figure 5.** Percentage (from the amount of pollen released from the anthers) in different paths after one

564   visit on a donor flower in *S. dulcamara* and *S. rostratum*.

Figure 6. Number of pollen grains deposited on recipient flower stigmas as influenced by flower species
and recipient order. Plotted lines indicate estimated means and shaded regions indicate standard errors.

Figure 7. Number of pollen grains deposited on recipient flower stigmas as influenced by (A) duration of
the visit in seconds, and (B) flower species and recipient order with released pollen as offset variable.
Plotted lines indicate estimated means and shaded regions indicate standard errors.

A

Total pollen production
1065712 ± 116142

Remaining pollen
1000891 ± 121491

Pollen load
29358 ± 9222

Pollen on body
4892 ± 2198

Pollen in corbiculae
24121 ± 8634

Pollen deposited
356 ± 93

Dislodged pollen
53561 ± 17369

Pollen released
86732 ± 21423

Pollen loss
81495 ± 21477

Pollinator visit

B

Total pollen production
641780 ± 92979

Remaining pollen
378642 ± 65596

Pollen load
202146 ± 51589

Pollen on body
11685 ± 4466

Pollen in corbiculae
188601 ± 53757

Pollen deposited
1860 ± 370

Dislodged pollen
41177 ± 19734

Pollen released
243323 ± 49331

Pollen loss
229779 ± 51778

Pollinator visit