1 **Mating strategy explains sex-biased infections in an emerging fungal disease**
2

3 Macy J. Kailing[1], Joseph R. Hoyt[1], J. Paul White[2], Jennifer A. Redell[2], Heather M. Kaarakka[2],
4 and Kate E. Langwig[1]
5
1 Department of Biological Sciences, Virginia Polytechnic Institute, Blacksburg VA 24061
2 Wisconsin Department of Natural Resources, Madison WI 53707

Abstract
1) Mating dynamics can govern species impacts from rapid global change by influencing
population rates of growth and adaptation, as well as individual traits that affect mortality
risks from novel pressures.
2) Here, we examined sex differences in mating phenology of *Myotis lucifugus* to
understand how reproductive strategies contribute to disease impacts as the mating
season coincides with exposure to a lethal fungal pathogen, *Pseudogymnoascus
destructans*. We expected differences in activity among sexes to modify seasonal disease
dynamics as *P. destructans* can replicate only at the cool temperatures at which bats
hibernate.
3) We used passive antenna systems installed at the entrances of hibernacula and PIT tags to
characterize activity patterns of bats impacted by white-nose syndrome. We also
measured pathogen loads on bats during autumn mating and early hibernation to assess
how infection severity changed according to host phenology.
4) We found that females spent fewer days active during autumn, arrived after males, and
were primarily active on the warmest nights. Males remained highly active throughout
the mating period and later in autumn than females. Importantly, these differences in
mating phenology corresponded to more severe infections in females during early
hibernation as male activity, and thus warm body temperatures, inhibit pathogen growth.
5) Differences in activity between sexes and in the transition from swarm to hibernation
likely reflects males maximizing their mating opportunities while females conserve
energy to meet the cost of spring migration and reproduction. More broadly, our results
show how mating phenology can contribute to sex-biased impacts of a novel disease and
highlight the value of understanding species' mating systems to anticipate the impacts of
environmental change.

34
1. Introduction

Global change has imposed novel pressures that threaten many species with extinction,

including habitat loss, pollution, overutilization and emerging infectious disease (Dirzo et al.,

2014; Mazor et al., 2018; Otto, 2018; Tilman et al., 2017). The resilience of populations to global

changes can vary based on the species' life history strategies, including the mating system

(Capdevila et al., 2022; Péron, 2013; Plesnar-Bielak et al., 2012; Sæther et al., 2004). Mating

system dynamics govern how quickly populations recover and adapt by influencing demographic

and evolutionary processes, such as rates of recruitment and genetic mixing (Bessa-Gomes,

Legendre, & Clobert, 2004; Legendre et al., 1999). Accordingly, traits associated with mating

systems are often shared within demographic classes (e.g., sex, age, social status, etc.) and can

result in mortality on only some demographic classes that can restructure or destabilize

populations (Clutton-Brock et al., 2002; Grüebler et al., 2008; Martin et al., 2022; Oro, Álvarez,

& Velando, 2018; Rosa et al., 2019; Wells et al., 2017). Therefore, understanding how aspects of

the mating system influence interactions between individuals and novel pressures should

improve the ability to predict responses of populations to rapid changes.

Sexes use different strategies for maximizing their fitness that give rise to variation in

morphology, behaviors, or physiology (Kolata, 1977; Promislow, Montgomerie, & Martin, 1992;

Trivers, 1972), which collectively can affect their mortality risk (Promislow, 1997). In many

species where sexual selection leads to dimorphic traits between sexes, males invest in quantity

of offspring through copious mating, whereas females, who are more limited by gametic

production, invest in ensuring survival of few high-quality offspring (Bateman, 1948; Lehtonen,

Parker, & Schärer, 2016). Accordingly, males may be more likely to engage in risky behaviors to

increase their competitive mating advantages whereas females, whose fitness relies on surviving

to carry and care for young, may be more risk-averse (Bateman, 1948; Bondurlansky et al., 2008;

Promislow, 1997). Additionally, asynchrony between sexes in their reproductive investments can

also contribute to differences in how females and males respond to the same environmental

pressures (Cornwallis & Uller, 2010; Forsgren et al., 2004; Kasumovic et al., 2008), including

when their peak vulnerability to stressors occurs (Williams et al., 2022). While sex specific

responses to novel pressures are expected, the proximate factors that influence differences in

impacts between females and males are overlooked for many novel threats (Connallon, Débarre,

& Li, 2018; Connallon & Hall, 2018).

Novel pressures like infectious diseases frequently interact with mating systems resulting

in differential impacts to sexes (Herrera & Nunn, 2019; Stephenson, 2019; Stephenson et al.,

2016; Zuk, 2009; Zuk & McKean, 1996) that can influence the severity of epidemics and

coevolution between hosts and pathogens (Hall & Mideo, 2018). Typically, male behaviors or

physiological states optimize fitness at the costs of higher pathogen exposure or growth,

including territory defense or aggression, dispersal, and reduced immunity (Ferrari et al., 2003;

Grear, Luong, & Hudson, 2012; Luong et al., 2010; Sauer et al., 2024; Stephenson, 2019). For

example, adult male badgers had higher contacts with conspecifics and higher risks of acquiring

tuberculosis compared to females (Delahay et al., 2006; Graham et al., 2013; McDonald et al.,

2014). Male-biased disease is more frequently documented, but female-biased disease has been

shown to arise seasonally (Cattadori et al., 2005; Sweeny et al., 2021). For example, during

parturition adult female European rabbits had the sharpest peak in nematode intensities relative

to non-reproductive females and males (Cattadori et al., 2005), illustrating that the magnitude

and direction of sex-biases can vary temporally to affect the dynamics between hosts and

pathogens (Cattadori et al., 2005; Sweeny et al., 2021). However, the behavioral and

physiological mechanisms contributing to sex-biases in impacts for introduced pathogens are

often unclear, thus limiting opportunities for management.

White-nose syndrome (WNS) is a recently introduced disease of bats that is caused by the

psychrophilic fungal pathogen, *Pseudogymnoascus destructans*, and has led to population

declines exceeding 95% (Frick et al., 2015; Langwig et al., 2012; Langwig et al., 2016).

Euthermic bats that are active on the landscape maintain higher body temperatures (37-41°C)

(O'Shea et al., 2014) that are above the upper critical limit of *P. destructans* growth (20°C)

(Verant et al., 2012). However, when bats reduce their body temperature during torpor (short

periods of inactivity that comprise hibernation) to near ambient (1-16°C) (Perry, 2013; Webb,

Speakman, & Racey, 1996) the pathogen is able to grow, making it dependent on seasonal

hibernation cycles (Jackson, Willcox, & Bernard, 2022; Jackson et al., 2022; Langwig et al.,

2015; Langwig et al., 2021). Pathogen exposure occurs during autumn when bats return annually

to contaminated hibernation sites (Hoyt et al., 2020; Laggan et al., 2023) but infection and

mortality peak over winter (Hoyt, Kilpatrick, & Langwig, 2021; Langwig et al., 2015; Reeder et

al., 2012).

The mating period for temperate bats impacted by WNS, known as swarm, occurs during

autumn when large numbers of individuals aggregate near hibernacula entrances, then enter the

site to form smaller groups (typically pairs or triads), and copulate after landing on walls or in

crevices (Fenton, 1969; Fraser & McGuire, 2023; Thomas, Brock Fenton, & Barclay, 1979; Wai-

Ping & Fenton, 1988). While females mate during this period, they store sperm through

hibernation until spring emergence, when they migrate to summer maternity colonies, become

pregnant, and rear pups cooperatively with other females (Fenton & Robert, 1980; Kurta &

Kunz, 1987; Wimsatt, 1942, 1944, 1945). Males segregate from females during parturition and

pup rearing (Davis & Hitchcock, 1965; Fenton & Robert, 1980). To that end, females'

reproductive investments are primarily during spring and summer, and are more extensive in

magnitude and duration than males, whose reproductive investments are highest during autumn

(Racey & Entwistle, 2000; Fig 1). While autumn mating is critical for both sexes, asynchronous

allocation of energy for reproduction between females and males could lead to extensive

behavioral and physiological differences that contribute to disease patterns and impacts.

Importantly, the congregation of sexes during autumn mating and hibernation provide a unique

opportunity to explore how sex-specific mating dynamics and phenology influence infection

between sexes.

Figure 1. Hypothesized reproductive investment of females and males with different

reproductive strategies developed from Fenton & Robert, 1980, Racey & Entwistle, 2000, and

Wimsatt, 1945. Males are expected to invest most towards their reproductive fitness during

autumn when they mate polygynously. Females likely require less energy during autumn mating,

but considerable investment is made following hibernation as they singly parent their young in

maternity colonies and require energy for gestation, parturition, and lactation. Light blue points

correspond to the tagging and sampling dates of autumn mating and early hibernation as

described in Methods.

Temperate bats reduce activity seasonally and use torpor to conserve energy even outside of

winter hibernation (Alston et al., 2022). However, females and males balance torpor and activity

differently throughout the year (Grinevitch, Holroyd, & Barclay, 1995; Hoff et al., 2023). These

sex specific differences in activity and torpor could strongly influence WNS dynamics if sex-

specific torpor use influences growth of the cool-temperature dependent fungus. Yet, empirical

investigations are limited by the challenge of obtaining individually based data on mating

dynamics through time. Previous research has shown that females can be more infected and

suffer greater impacts (Grieneisen et al., 2015; Kailing et al., 2023), but see (Johnson et al.,

2014). However, the mechanisms driving the difference between sexes remain unknown. We

hypothesize that autumn mating activity drives highly disparate infections in early hibernation

since the pathogen is unable to grow on bats when they are active. As such, we investigated how

traits associated with mating vary between females and males during seasons important for

reproduction (autumn mating) and disease (autumn mating and early hibernation) and contribute

to sex-biased disease. Specifically, we aimed to characterize sex-specific mating periods,

examine environmental and host factors associated with mating activity, and explore links

between phenology and disease.

2.   Methods

2.1 Field monitoring

We studied the activity of *Myotis lucifugus* in Wisconsin, USA during autumn mating

(i.e., July 01 to November 01) each year from 2020 to 2023 at three large hibernacula within

Pierce and Dodge counties, consisting of 10 total entrances. At every entrance, we installed an

RFID system consisting of an antenna with an automated data logger (IS10001; Biomark) that

recorded a time-stamped datapoint each time a passive integrated transponder (PIT) tag was in

proximity. Each system was scheduled to run continuously by a direct or solar source of power.

Systems were programmed to record unique tags every minute to help to maximize detections as

>1 tag cannot be simultaneously recorded and bats sitting on top of antenna would otherwise

block detections of all other individuals. In addition, systems recorded test tags at least once

every 12 hours enabling us to determine when systems were operational.

2.2 PIT tagging and measuring infection

We captured bats at each site during autumn mating (between August 15 and September

02, Fig. 1) using a harp trap (Harptrap; Faunatech Austbat) and mist-nets (Polyester 38mm mesh;

Avinet) near the entrances where individuals were actively swarming. For every individual, we

weighed, injected a PIT tag, categorized sex based on external morphology, and attached an

aluminum-lipped band (2.9 mm; Porzana, Icklesham, UK) as secondary identification. Capture

efforts at each site concluded after one night, unless the target number of bats (approximately

100) was not achieved, in which we trapped a second night.

We also sampled bats in early hibernation (between November 01 and November 17, Fig.

1) at the same sites by removing them from their roosting locations on the walls and ceilings of

the hibernacula. We walked sections of the hibernacula to collect a stratified sample of up to 20

bats and targeted previously banded individuals. As in autumn, we recorded sex based on

external morphology, PIT tag ID, and band number of every individual. We collected an

epidermal swab from every bat sampled in autumn and early hibernation with a standardized

swabbing technique that is used widely to measure infection severity across seasons (Frick et al.,

2015; Langwig et al., 2015). Immediately upon collection, we stored swabs in RNAlater. We

tested swab samples for *P. destructans* using Qiagen DNeasy Blood and Tissue Kits (Qiagen,

USA) to extract fungal DNA followed by qPCR to quantify fungal loads (Langwig et al., 2015;

Muller et al., 2013). All bat handling procedures followed the approved IACUC protocols

[IACUC Protocol #20-150, #23-197] and decontamination followed USFWS as well state

recommendations.

2.3 Characterizing individual activity and climate variables

We determined individuals' mating phenology from PIT tag detections. An individual

was considered active if it was detected at any entrance to the site at least once in a night because

bats commonly visit multiple entrances throughout the night. We classified bats that were

detected between the hours of 00:00 and 07:00 as active on the previous date because bats are

nocturnal. We only included bats detected on at least two nights to properly categorize the

detections as being the bats' first or last within the same year to capture the apparent swarm

period. For each bat observed within a year, we assigned a start swarm date as the first date

detected after July 01 and an end swarm date as the last date detected before Nov 01. Start dates

were only assigned for individuals that returned from a previous year as we could not confirm

when bats first arrived at hibernacula if they were untagged. We were able to assign an end

swarm date to all bats, including those tagged in the same year, because the last detection always

followed the date they were captured and tagged. *Myotis lucifugus* can only be reliably aged as

juveniles (young of the year) and adults and all bats that return from a previous year are

considered adults. Therefore, we only included adults in all analyses.

To assess environmental factors associated with mating phenology, we extracted climate

data using the R package 'daymetr' (Hufkens et al., 2018) and GPS coordinates of hibernation

sites, which provided continuous, grid-based estimates of daily weather variables based on

location. To reflect bat ecology, and because mean temperatures were not available in the climate

data, we calculated a mid-point temperature (Jackson et al., 2022) for each date as: (minimum

temperature on the following day + maximum temperature of current day)/2.

2.4 Statistical analyses

2.4.1 Summary

We tested our hypotheses using generalized (i.e., GLMM) and linear (i.e., LMM) mixed

effects models to characterize autumn mating activity and disease by sex with R packages 'lme4'

(Bates et al., 2015) and 'glmmTMB' (Brooks et al., 2017). For each model, we tested variables

based on *a priori* hypotheses and evaluated their significance using Anova tables for statistical

model objects from the 'car' package (Fox & Weisberg, 2019). Furthermore, we compared

candidate and null models using Akaike Information Criterion (AIC) to evaluate relative quality

(difference in AIC scores |> 2|; See Appendix 4.6.2). We also visually checked model

assumptions, such as the residual distribution of linear models, and conducted k-fold cross-

validation of logistic models to check predictive accuracy (Appendix 4.6.3) using the

'performance' package (Lüdecke et al., 2021). Because many more males were detected than

females, we also generated a balanced dataset between sexes to check that the results informed

by our male-biased dataset were consistent and not due to the sampling bias. We truncated our

full dataset by randomly selecting the number of male observations equivalent to the number of

female observations from all the male observations (see Supplemental Methods). We extracted

model estimates using R package 'emmeans' (Lenth, 2022) to report final models. See

Supplemental Table 1 for overview of all analyses as well as Appendix with code for all

statistical models, output, and model validation.

2.4.2 Dates of autumn mating activity

We determined individually based activity periods using LMMs. First, we estimated

when individuals of each sex started autumn mating using the first date (in monthly units then

scaled to start at 0) an individual was detected at the hibernacula as a continuous response

variable, with sex as a fixed effect (F|M), and site-year and individual as random effects. Then,

we used the same model structure, but with last date detected as a response to estimate when

individuals of each sex ended mating activity.

220   *2.4.3 Effects of temperature and body condition on nightly activity*

221   To test whether environmental conditions differently affected nightly activity between

sexes, we used a GLMM with a binomial distribution to test the probability of an individual

being active (=1) or undetected (=0) with temperature and sex as interactive fixed effects and

individual and site-year as random effects. For the response variable, we treated each bat as a

series of binomial trials using all the possible dates between the first and last detection where an

individual was detected (=1) or undetected (=0). We excluded nights where the minimum

temperature was less than 10°C to minimize outlier effects as bat activity is greatly reduced

(Gorman et al., 2021).

229   We also tested the effect of body condition on nightly activity to assess whether

energetics might contribute to differences in activity between males and females. As we could

only associate mass to bats at capture, for this dataset, we used the activity of bats in their first

year tagged. We used a GLMM with a binomial distribution to test the probability of a bat being

active. Like the above analysis, our response variable was detection, with each bat treated as a

series of binomial trials where it was either detected or undetected. We used sex, mass, and

temperature as interactive fixed effects and included individual as a random effect.

236   *2.4.4 Effects of length of active period and temperature on last detection*

We examined how the number of days since arriving to the hibernaculum and

temperature affected whether sexes transition from the fall mating/activity period to hibernation.

First, we used a GLMM with a binomial distribution that predicted whether individuals stopped

activity for the year. For the response variable, detections of each bat were treated as a series of

binomial trials where an individual was detected for the last time (=1) or was detected on a

subsequent date (=0). We included interactive effects of sex and time since their first detection

(to standardize differences in start date) with site-year and individual as random effects. For days

since first detection, we binned dates to 3-day intervals and counted the number of intervals that

passed since their arrival to the hibernacula. Accordingly, we only included individuals that we

were able to assign start dates (i.e., returned from a previous year). Date intervals were used to

overcome sparsity in the number of females that are active each night.

To examine the effect of temperature on mating phenology, we used a GLMM with a

binomial distribution to test a similar response as above where each bats' detections were treated

as a series of binomial trials and an individual was detected for the last night (=1) or was

detected on a subsequent night (=0). We included sex and temperature as interacting fixed effects

and random effects of site-year and individual.

2.4.5 Seasonality and mating phenology effects on infection severity

To understand disease dynamics according to mating phenology, we tested differences in

pathogen loads between sexes and across seasons using a LMM with $\log_{10}$ *P. destructans* loads

as our response variable, with sex and date (in monthly units as above) as fixed effects, and site-

year as a random effect. We explored associations between phenology and disease further by

testing the effect of swarm activity on early hibernation infection severity. We used a LMM to

test $\log_{10}$ *P. destructans* loads as a response variable with median last date of activity as a fixed

effect. We included site-year as a random effect. We calculated the median date of last detection

in autumn for both sexes for each site and year (i.e., the end of swarm activity and assumed start

of hibernation) and assigned it to individuals that we sampled for infection in early hibernation

of the respective site and year. We used the median date of last detection to minimize the weight

of outlier individuals that may leave the hibernacula late in autumn but are not representative of

swarm activity. We also performed this analysis at the population level using the mean early

hibernation fungal loads from each site and year as the response. We used these approaches

because the ability to connect an individual bat across our target seasons (i.e., sample and

quantify infection from a bat in early hibernation that was previously detected in autumn) is

challenging.

3. Results

We characterized autumn activity of 148 (by site: 37, 40, 71) female and 608 (by site:

225, 155, 228) male *M. lucifugus* and measured infection during capture on 168 (by site: 56, 63,

49) female and 416 (by site: 151, 130, 135) males between 2020 and 2023. See Supplemental

Table 1 for overview of all analyses and Appendix 4.7 for full sample lists used in each analysis

by site, years, and sex. All statistical results are reported with males as the intercept or reference

level and thus coefficient $\pm$ SE are in reference to the value above or below the male references.

Females started swarm activity later (Female coef $\pm$ SE = 0.249 $\pm$ 0.08, t = 3.25, p-value = 0.001;

Fig. 2, top panel; Appendix 1.1) but also ended activity earlier than males (-0.387 $\pm$ 0.06, t = -

6.29, p-value < 0.001; Fig. 2, top panel; Appendix 1.3). Male activity fully encompassed the

activity period of females (Fig. 2, bottom panel; Appendix 1.3) and the number of individuals

detected was overwhelmingly male biased throughout autumn, given their higher abundance and

activity overall (Supplemental Fig. 1; Appendix 4.1), and was consistent between sampling

events (Supplemental Table 2; Appendix 4.6.1).

Within their respective mating periods, the probability of being active in a night or not for

each sex was differently affected by temperature (detected=0|1; Female*temperature $= 0.151 \pm$

0.03, t $= 4.97$, p-value < 0.001, Fig. 3A; Appendix 2.1), such that the probability of females

being active was lower overall but increased more quickly with increasing temperatures

(Temperature slope estimated using 'emtrends': Female slope $= 0.152 \pm 0.03$ vs Male slope

0.001 ± 0.01, Fig. 3A; Appendix 2.1), notably above 16°C. An interaction between sex and

temperature was also evident when using a balanced dataset of males and females (see

Supplemental Statistical Methods, Supplemental Fig. 2A; Appendix 4.2.1). We found that the

effect of body condition on activity differed between males and females, with light females less

active at cooler temperatures and more active at warmer temperatures, whereas males maintained

high activity despite large differences in temperatures (Female coef*mass*temperature $\pm$ SE $= -$

0.122 ± 0.06, z $= -2.07$, p-value $= 0.039$, Supplemental Fig. 3, Appendix 4.3), We also found

support for an interaction between the length of the active period and sex on the probability of

being detected for the last time (Date intervals since arrival*female $= 0.367 \pm 0.07$, z $= 5.49$, p-

value < 0.001, Fig. 3B; Appendix 2.2), as well as an interactive effect of temperature and sex on

the probability of being detected for the last time, such that females were significantly more

likely to end their active period as temperatures decreased than males (Temperature*female $= -$

0.164 ± 0.08, z $= -2.13$, p-value $= 0.033$, Fig. 3C; Appendix 2.3).

There was no clear sex bias in pathogen loads throughout autumn (Female $= -0.014 \pm$

0.10, t $= -0.14$; p-value $= 0.887$, Appendix 3.1), however, following autumn mating, female

pathogen loads increased more rapidly, and females became significantly more infected than

males by early hibernation (Female*date = 0.340 ± 0.08, t = 4.50, p-value = < 0.001; Fig. 4A;

Appendix 3.2). Ending of autumn activity of males and females generally varied among sites and

years and earlier end dates were associated with higher pathogen loads on individuals (Median

last date of activity slope on pathogen loads: -1.574 ± 0.41, t = -3.87, p-value < 0.001; Fig. 4B;

Appendix 3.3) as well as site-level pathogen loads in early hibernation (Median last date of

activity slope on mean pathogen loads: -1.562 ± 0.33, t = -4.75, p-value < 0.001; Supplemental

Fig. 4; Appendix 4.4).

Figure 2. Top: Differences in the autumn mating period by sex. Each point represents the date

of starting (first detection) and ending (last detection) activity during autumn mating for an

individual bat. Black circles represent model estimated dates with standard errors shown with

horizontal bars. **Bottom:** Distribution of activity by sex. Each density plot shows the proportion

of all detections grouped by sex that occurred on each date. An individual bat can be represented

on multiple nights if activity occurred. Asymmetry in the peak of male activity (based on the

total number of observations) relative to their estimated active period in the top panel illustrates

visually that while most males arrive early in autumn, they maintain activity into later swarm

dates.

Figure 3A. Effect of temperature on the probability that an individual was active in a night. Each

point represents the proportion of individuals from a site and year that were active at each

temperature by sex. Females are shown in purple and males in orange and lines represent the

model predicted probabilities of being active across the temperature gradient. **Figure 3B.**

Relationship between length of the active period in days and the probability of last annual

detection for females in purple and males in orange. Lines denote the model predicted

probabilities of being detected for the last time within a year. Actual data are binary (1|0) such

that 1 = ends activity and 0 = continues to be active. For visualization, points show the

proportions of bats of each sex that were detected for the last time within the year, binned across

a three-day interval. **Figure 3C**. Relationship between temperature and probability that bats

ended their autumn activity. Each point summarizes the proportion of bats that were detected for

the last time at a given temperature for each site and year. Females are shown in purple and

males are in orange. Solid lines denote the model predicted probability of being detected for the

last time for each sex. Size of points show the number of bats represented in proportion

calculations. Notably, males do not end their swarm activity even at temperatures that are

relatively low. In each panel, shaded grey regions show standard errors of model predictions.

Figure 4A: Pathogen loads measured on adults of each sex throughout the autumn mating and

early hibernation periods. Each point represents an individual bat with females in purple and

males in orange. Solid lines denote model predicted fungal loads over time for females and males

respectively, with grey shaded regions displaying the standard error. Dashed lines represent the

model predicted end dates of activity as estimated in Fig. 2. There was no clear difference in

infection between sexes during the active mating period in autumn, but female infections were

nearly an order of magnitude more severe than males by early hibernation. **Figure 4B.**

Association between the median date of last detection at a site and early hibernation pathogen

loads on individuals. Each point represents an individual bat sampled in early hibernation

colored by sex as in panel A and are jittered slightly for visualization. Median last date of

autumn activity was calculated for both sexes during each sampling event (i.e., site-year) and

used to minimize the outlier effects of a few individuals with exceptionally early or late last

active dates. Black line shows model predicted pathogen loads with standard errors displayed

with the grey shading. In each panel, date units have been converted to calendar days for clarity.

4. Discussion

Sex specific reproductive strategies are common across species, yet how aspects of

mating systems give rise to sex-biased disease impacts are often unexplored. Our results suggest

that asynchronous phenology between sexes may be a key driver of seasonally sex-biased

infections, which are known to affect pathogen transmission and population impacts in other

host-pathogen systems (Ferrari et al., 2003; Graham et al., 2013; Grear, Luong, & Hudson, 2012;

Perkins, Ferrari, & Hudson, 2008; Sauer et al., 2024; Silk et al., 2018; Skorping & Jensen, 2004).

We show female *M. lucifugus* begin swarm activity later and end earlier, spend fewer nights

active after arriving to swarm, and have a more pronounced increase in activity on warmer nights

compared to males. Importantly, pathogen loads do not differ between sexes until after the active

period of autumn mating, when bats transition to hibernation, and earlier end dates of autumn

activity were associated with higher pathogen loads by early hibernation. Earlier use of torpor by

females likely contributes to female-biased disease as it allows an extended period for the

pathogen to replicate as opposed to males that remain active and can minimize pathogen

replication through euthermia.

In exploring within activity period differences, we found that females are far less active

during their shorter swarm period compared to males. This is probably best explained by the

need for females to conserve enough energy stores to survive hibernation. Females store sperm

overwinter and then give birth and raise offspring following emergence (Czenze, Jonasson, &

Willis, 2017; Jonasson & Willis, 2011; Wimsatt, 1945). We provide two new lines of

individually derived evidence that aligns with females using energy more conservatively than

males. First, we found a clear effect of temperature differently affecting the probability that

females and males are active on a given night within their swarm period, and the relationship

changed depending on the body condition of females but not males (Supplemental Fig. 3,

Appendix 4.3). This suggests that females may be prioritizing the limited days they spend active

in autumn to forage on warmer nights that average nearly 20°C (Supplemental Fig. 2B,

Appendix 4.2.2) likely because thermoregulating is more efficient at warmer temperatures

(Fjelldal et al., 2023; Rubalcaba et al., 2022) and prey resources are more active (Grüebler,

Morand, & Naef-Daenzer, 2008; Taylor, 1963). Second, we found that females on average,

regardless of their arrival date, begin transitioning into hibernation (i.e., remained in the

site/stopped activity) after four days of activity, whereas males maintained nightly activity until

late in autumn regardless of the date they arrived to swarm, and averaged over 13 active nights

(Supplemental Figs 1 and 5, Appendix 4.1 and 4.5). Males that are less constrained by

reproductive energy budgets were active nearly four times as many nights as females

(Supplemental Figs 1 and 5; Appendix 4.1 and 4.5), when temperatures are relatively low (Figs

3A and 3C, Appendix 2.1 and 2.3), and activity was independent of body condition (in contrast

to females, Supplemental Fig. 3), suggesting that males are optimizing mating opportunities over

energy conservation and supports previous hypotheses (Burns & Broders, 2015; Glover &

Altringham, 2008; Rivers, Butlin, & Altringham, 2006; Willis, 2017). Importantly, the disparate

reproductive costs between sexes are likely contributing to how each sex interacts with the same

novel pressure of an introduced pathogen.

The earlier arrival of males to mating locations than females, termed protandry, is

observed across many taxa, and two hypotheses have been posed to justify its ubiquity (Morbey

& Ydenberg, 2001). First, the 'rank advantage hypothesis' states the earliest arriving individuals

have access to the best resources or territories, providing a competitive advantage in acquiring

females. Second, the 'mate opportunity hypothesis' suggests that earlier arriving males can

increase the number of females mated, resulting in greater annual reproductive success. In

systems under strong sexual selection and in cases where females also have resource-based

advantages of arriving early, mate opportunity has more reliably explained protandry than rank

advantage, particularly in species exhibiting sperm competition and highly male-biased sex ratios

(Kokko et al., 2006), which is known of temperate bats (Racey et al., 1987; Racey & Entwistle,

2000). Sperm storage by females can also generate advantages for males that more closely

synchronize copulations with female receptivity, such that males that mate earlier can secure

storage sites in uteri that improve fertilization success (Scialli, 1993). Thus, we hypothesize that

the longer swarm period of males is driven by the advantage of increasing copulations with more

females in addition to elevating fertilization potential, as suggested by other studies of temperate

bats (Burns & Broders, 2015; Pfeiffer & Mayer, 2013). Other environmentally mediated

mechanisms likely function in concert with mating advantages. Females migrate from summer

maternity colonies and, because foraging to store fat for hibernation is essential for bats in

autumn (Kunz, Wrazen, & Burnett, 1998), timing the departure from diminishing resources in

summer habitats to optimal resource conditions at swarm locations is likely a more important

balance for females that have greater annual energetic constraints than males.

As in other disease systems, sex-specific differences in behavior or physiology during

periods of pathogen exposure can result in sex-biased disease impacts and govern host-pathogen

dynamics (Guerra-Silveira & Abad-Franch, 2013). We showed that during the mating season

when bats are initially exposed to *P. destructans* after having cleared prior infections (Langwig

et al., 2015), activity differs markedly between sexes, however it did not promote differences in

infection within that period (Appendix 3.1) as we expected based on other disease systems

(Ferrari et al., 2003; Graham et al., 2013; Guerra-Silveira & Abad-Franch, 2013; Klein, 2000).

Furthermore, infections during the autumn mating period or differences in tolerance between

sexes are not expected to influence activity patterns because the fungal loads during the active

period are likely too low to initiate behaviors or physiological responses symptomatic of disease

(Warnecke et al., 2012). Rather, reproductive demands that influence broader phenological

patterns are more likely to be shaping sex-biased disease through physiological mechanisms.

Female infections rapidly increased almost two orders of magnitude within the two months

between the end of their active period and early hibernation (Fig. 4A, Appendix 3.2), suggesting

the timing of exposure and pathogen growth have temporally decoupled the effects on sex-

specific disease in this host-pathogen system. We found that ending activity earlier was

associated with higher pathogen loads in early hibernation, which aligns with females being the

earliest to hibernate and develop the highest pathogen loads (Fig. 4B, Appendix 3.3). Ending

activity is significant in this system because the pathogen can only replicate once bats begin

hibernation, and hosts that exceed an infection threshold earlier have lower over-winter survival

(Langwig et al., 2016). Consequently, females are likely predisposed to have higher disease than

males because of their need to conserve energy to achieve reproductive fitness.

Phenology evolves to synchronize important biological events, such as reproduction, to

optimal conditions and resources (Visser & Gienapp, 2019). However, rapid changes in the

biotic or abiotic environment can modify which traits are beneficial or maladaptive (Schlaepfer,

Runge, & Sherman, 2002). For bats affected by WNS, if swarm periods evolved to maximize

survival and recruitment, disease-induced mortality of early arriving and less active individuals

may be selecting for traits that are beneficial for surviving infection but potentially maladaptive

for recruitment. Alternatively, if phenology is a plastic trait in bats as it is in other taxa (de

Villemereuil et al., 2020), a shift to a later swarm period may minimize disease while still

supporting recruitment, thus ameliorating population impacts. Given the clear, primarily additive

effect of sex on periods of mating activity (Fig. 2; Supplemental Table 2; Appendix 4.6.1), we

expect that any shifts in activity of one sex will be paralleled by the other. Because females have

a more restricted activity period and energy budgets, we expect females to be less flexible in

their phenology than males. As such, females will likely need to adapt alternative mechanisms to

overcome novel disease pressure or compensatory mechanisms to maintain recruitment if a later

swarm window is suboptimal for fitness.

Environmental conditions may also alleviate impacts on females throughout their

geographic range or between years. Female bats in milder climates, such as at lower latitudes or

years with warmer autumns, may not face the same restriction of suboptimal temperatures that

cause bats to begin hibernation. In such instances, females may stay active longer to avoid

developing infections severe enough to cause mortality. Similarly, female-biased infections are

observed across most host species (Kailing et al., 2023) and share similar reproductive cycles as

in Fig. 1. However, more subtle interspecific differences in mating phenology, like those

explored in this study, are largely unknown for most species impacted by WNS but may

plausibly contribute to the known variation in early hibernation infections and impacts among

species (Langwig et al., 2016; Langwig et al., 2015). Compared to species that mate earlier,

species with later swarm activity and delayed hibernation may have lower infections overall that

allow more females to survive winter. Previous work has implicated female-biased disease in an

initial reduction in the proportion of females following WNS invasion (Kailing et al., 2023),

however, many colonies where the pathogen invaded earliest are recovering (Grimaudo et al.,

2021; Hoyt, Kilpatrick, & Langwig, 2021; Langwig et al., 2017). Therefore, long-term

ecological and evolutionary responses of each sex to disproportionate disease impacts remain to

be explored but are warranted, particularly at spatial and temporal scales beyond this study, as

they may contribute to how populations persist.

While sex-based differences in disease have received considerable attention, the

mechanisms are often unknown. We demonstrate that in the absence of clear behaviorally

mediated transmission differences, sex-specific physiology associated with phenological

transitions following exposure can facilitate sex-biased disease, providing a unique example in

which physiological asynchronies decouple impacts of the timing of acquiring a pathogen and

disease progression between sexes. Importantly, our findings suggest that asynchronous

reproductive energy allocation contribute to sex-biased infections, revealing a potentially

significant influence of mating dynamics on the trajectory of epidemics and pathways of host

persistence or extinction. Other novel pressures that similarly act more strongly on female

mating dynamics may impede population recoveries if sex-ratios are disrupted or if traits

beneficial for survival are antagonistic for recruitment. Given the importance of demographic

structure and recruitment in population impacts, determining how mating strategies influence

responses to novel stressors will be critical for conserving species in the face of increasing global

change.

Acknowledgements
We thank Kirsten Fuller and Steffany Yamada for data curation support, and Alexander T.
Grimaudo, Nichole A. Laggan, Ariel E. Leon, and Tonie E. Rocke for support in field data
collection. We also thank University of Wisconsin Milwaukee Field Station and Gretchen Meyer
for site access.
492
493   **Conflict of interest**
494   All authors declare no conflict of interest.
495
Author contributions
Macy J. Kailing, Kate E. Langwig, and Joseph R. Hoyt conceived the ideas and designed
methodology; Macy J. Kailing, Joseph R. Hoyt, J. Paul White, Jennifer A. Redell, Heather M.
Kaarakka, and Kate E. Langwig collected the data; Macy J. Kailing analyzed data with input
from Kate E. Langwig and Joseph R. Hoyt. Macy J. Kailing wrote the original manuscript with
Kate E. Langwig and Joseph R. Hoyt. All authors contributed critically to the drafts and gave
final approval for publication.
503
504   **Data availability statement**
505   The datasets and code generated in this study will be deposited in Dryad Digital Repository upon
506   final submission. Exact site locations are not disclosed to protect endangered species and
507   landowners.
508

References
Alston, J. M., Dillon, M. E., Keinath, D. A., Abernethy, I. M., & Goheen, J. R. (2022). Daily
torpor reduces the energetic consequences of microhabitat selection for a widespread bat.
Ecology, 103(6), e3677. https://doi.org/10.1002/ecy.3677
Bateman, A. J. (1948). Intra-sexual selection in Drosophila. *Heredity, 2*(3), 349-368.
https://doi.org/10.1038/hdy.1948.21
Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software, 67*(1), 1 - 48.
https://doi.org/10.18637/jss.v067.i01
Bessa-Gomes, C., Legendre, S., & Clobert, J. (2004). Allee effects, mating systems and the
extinction risk in populations with two sexes. *Ecology Letters, 7*(9), 802-812.
https://doi.org/10.1111/j.1461-0248.2004.00632.x

Bonduriansky, R., Maklakov, A., Zajitschek, F., & Brooks, R. (2008). Sexual Selection, Sexual
Conflict and the Evolution of Ageing and Life Span. *Functional Ecology, 22*(3), 443-
453. https://doi.org/10.1111/j.1365-2435.2008.01417.x
Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A.,
Skaug, H.J., Maechler, M., Bolker, B.M. (2017). "glmmTMB Balances Speed and
Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling." *The
R Journal*, *9*(2), 378–400. https://doi.org/10.32614/RJ-2017-066.
Brunet-Rossinni, A. K., & Wilkinson, G. S. (2009). Methods for Age Estimation and the Study
of Senescense in Bats. In T. H. Kunz & S. Parsons (Eds.), *Ecological and Behavioral*
Methods for the Study of Bats. Baltimore: The Johns Hopkins University Press.
Burns, L. E., & Broders, H. G. (2015). Maximizing mating opportunities: higher autumn
swarming activity in male versus female Myotis bats. *Journal of Mammalogy, 96*(6),
1326-1336. https://doi.org/10.1093/jmammal/gyv141
Capdevila, P., Stott, I., Cant, J., Beger, M., Rowlands, G., Grace, M., & Salguero-Gómez, R.
(2022). Life history mediates the trade-offs among different components of demographic
resilience. *Ecology Letters, 25*(6), 1566-1579. https://doi.org/10.1111/ele.14004
Cattadori, I. M., Boag, B., Bjørnstad, O. N., Cornell, S. J., & Hudson, P. J. (2005). Peak shift and
epidemiology in a seasonal host-nematode system. *Proceedings of the Royal Society B:*
Biological Sciences, 272(1568), 1163-1169. https://doi.org/10.1098/rspb.2004.3050
Clutton-Brock, T. H., Coulson, T. N., Milner-Guiland, E. J., Thomson, D., & Armstrong, H. M.
(2002). Sex differences in emigration and mortality affect optimal management of deer
populations. *Nature, 415*(6872), 633-637. https://doi.org/10.1038/415633a
Connallon, T., Débarre, F., & Li, X.-Y. (2018). Linking local adaptation with the evolution of
sex differences. *Philosophical Transactions of the Royal Society B: Biological Sciences,*
373(1757), 20170414. https://doi.org/10.1098/rstb.2017.0414
Connallon, T., & Hall, M. D. (2018). Environmental Changes and Sexually Antagonistic
Selection. In *Encyclopedia of Life Sciences* (pp. 1-7).
Cornwallis, C. K., & Uller, T. (2010). Towards an evolutionary ecology of sexual traits. *Trends*
in Ecology & Evolution, 25(3), 145-152. https://doi.org/10.1016/j.tree.2009.09.008
Czenze, Z. J., Jonasson, K. A., & Willis, C. K. R. (2017). Thrifty females, frisky males: Winter
energetics of hibernating bats from a cold climate. *Physiological and Biochemical*
Zoology, 90(4), 502-511. https://doi.org/10.1086/692623
Davis, W. H., & Hitchcock, H. B. (1965). Biology and migration of the bat, Myotis lucifugus, in
New England. *Journal of Mammalogy, 46*(2), 296-313.
de Villemereuil, P., Charmantier, A., Arlt, D., Bize, P., Brekke, P., Brouwer, L., Cockburn, A.,
Côté, S. D., Dobson, F. S., Evans, S. R., Festa-Bianchet, M., Gamelon, M., Hamel, S.,
Hegelbach, J., Jerstad, K., Kempenaers, B., Kruuk, L. E. B., Kumpula, J., Kvalnes, T.,
McAdam, A. G., McFarlane, S. E., Morrissey, M. B., Pärt, T., Pemberton, J. M.,
Qvarnström, A., Røstad, O. W., Schroeder, J., Senar, J. C., Sheldon, B. C., van de Pol,
M., Visser, M. E., Wheelwright, N. T., Tufto, J., & Chevin, L.-M. (2020). Fluctuating
optimum and temporally variable selection on breeding date in birds and mammals.
Proceedings of the National Academy of Sciences, 117(50), 31969-31978.
https://doi.org/10.1073/pnas.2009003117
Delahay, R. J., Walker, N. J., Forrester, G. J., Harmsen, B., Riordan, P., Macdonald, D. W.,
Newman, C., & Cheeseman, C. L. (2006). Demographic correlates of bite wounding in

Eurasian badgers, Meles meles L., in stable and perturbed populations. *Animal Behaviour, 71*(5), 1047-1055. https://doi.org/10.1016/j.anbehav.2005.07.018

Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J. B., & Collen, B. (2014). Defaunation in the Anthropocene. *Science, 345*(6195), 401-406. https://doi.org/10.1126/science.1251817

Fenton, M. B. (1969). Summer activity of Myotis lucifugus (Chiroptera:Vespertilionidae) at hibernacula in Ontario and Quebec. *Canadian Journal of Zoology, 47*(4), 597-602. https://doi.org/10.1139/z69-103

Fenton, M. B., & Robert, M. R. B. (1980). Myotis lucifugus. *Mammalian Species*(142), 1-8. https://doi.org/10.2307/3503792

Ferrari, N., Cattadori, I. M., Nespereira, J., Rizzoli, A., & Hudson, P. J. (2003). The role of host sex in parasite dynamics: field experiments on the yellow-necked mouse Apodemus flavicollis. *Ecology Letters, 7*(2), 88-94. https://doi.org/10.1046/j.1461-0248.2003.00552.x

Fjelldal, M. A., Muller, A. S., Ratikainen, I. I., Stawski, C., & Wright, J. (2023). The small-bat-in-summer paradigm: Energetics and adaptive behavioural routines of bats investigated through a stochastic dynamic model. *Journal of Animal Ecology, 92*(10), 2078-2093. https://doi.org/10.1111/1365-2656.13999

Forsgren, E., Amundsen, T., Borg, Å. A., & Bjelvenmark, J. (2004). Unusually dynamic sex roles in a fish. *Nature, 429*(6991), 551-554. doi:10.1038/nature02562

Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression* (3rd ed.). Thousand Oaks, CA: Sage.

Fraser, E. E., & McGuire, L. P. (2023). Prehibernation swarming in temperate bats: a critical transition between summer activity and hibernation. *Canadian Journal of Zoology, 101*(6), 408-422. https://doi.org/10.1139/cjz-2022-0129

Frick, W. F., Puechmaille, S. J., Hoyt, J. R., Nickel, B. A., Langwig, K. E., Foster, J. T., Barlow, K. E., Bartonička, T., Feller, D., Haarsma, A.-J., Herzog, C., Horáček, I., Kooij, J., Mulkens, B., Petrov, B., Reynolds, R., Rodrigues, L., Stihler, C. W., Turner, G. G., & Kilpatrick, A. M. (2015). Disease alters macroecological patterns of North American bats. *Global Ecology & Biogeography, 24*(7), 741-749. https://doi.org/10.1111/geb.12290

Glover, A. M., & Altringham, J. D. (2008). Cave selection and use by swarming bat species. *Biological Conservation, 141*(6), 1493-1504. https://doi.org/10.1016/j.biocon.2008.03.012

Gorman, K. M., Barr, E. L., Ries, L., Nocera, T., & Ford, W. M. (2021). Bat activity patterns relative to temporal and weather effects in a temperate coastal environment. *Global Ecology and Conservation, 30*, e01769. https://doi.org/10.1016/j.gecco.2021.e01769

Graham, J., Smith, G. C., Delahay, R. J., Bailey, T., McDonald, R. A., & Hodgson, D. (2013). Multi-state modelling reveals sex-dependent transmission, progression and severity of tuberculosis in wild badgers. *Epidemiology and Infection, 141*(7), 1429-1436. https://doi.org/10.1017/S0950268812003019

Grear, D. A., Luong, L. T., & Hudson, P. J. (2012). Sex-biased transmission of a complex life-cycle parasite: Why males matter. *Oikos, 121*(9), 1446-1453. https://doi.org/10.1111/j.1600-0706.2012.20358.x

Grieneisen, L. E., Brownlee-Bouboulis, S. A., Johnson, J. S., & Reeder, D. M. (2015). Sex and hibernaculum temperature predict survivorship in white-nose syndrome affected little

brown myotis (Myotis lucifugus). *Royal Society Open Science, 2*.
https://doi.org/10.1098/rsos.140470
Grimaudo, A. T., Hoyt, J. R., Yamada, S. A., Herzog, C. J., Bennett, A. B., & Langwig, K. E.
(2021). Host traits and environment interact to determine persistence of bat populations
impacted by white-nose syndrome. *Ecology Letters, 25*(2), 483-497.
https://doi.org/10.1111/ele.13942
Grinevitch, L., Holroyd, S. L., & Barclay, R. M. R. (1995). Sex differences in the use of daily
torpor and foraging time by big brown bats (Eptesicus fuscus) during the reproductive
season. *Journal of Zoology, 235*(2), 301-309. https://doi.org/10.1111/j.1469-
7998.1995.tb05146.x
Grüebler, M. U., Morand, M., & Naef-Daenzer, B. (2008). A predictive model of the density of
airborne insects in agricultural environments. *Agriculture, Ecosystems & Environment,*
123(1), 75-80. https://doi.org/10.1016/j.agee.2007.05.001
Grüebler, M. U., Schuler, H., Müller, M., Spaar, R., Horch, P., & Naef-Daenzer, B. (2008).
Female biased mortality caused by anthropogenic nest loss contributes to population
decline and adult sex ratio of a meadow bird. *Biological Conservation, 141*(12), 3040-
3049. https://doi.org/10.1016/j.biocon.2008.09.008
Guerra-Silveira, F., & Abad-Franch, F. (2013). Sex Bias in Infectious Disease Epidemiology:
Patterns and Processes. *PLOS ONE, 8*(4). https://doi.org/10.1371/journal.pone.0062390
Hall, M. D., & Mideo, N. (2018). Linking sex differences to the evolution of infectious disease
life-histories. *Philosophical Transactions of the Royal Society B: Biological Sciences,*
373(1757). https://doi.org/10.1098/rstb.2017.0431
Herrera, J., & Nunn, C. L. (2019). Behavioural ecology and infectious disease: implications for
conservation of biodiversity. *Philosophical Transactions of the Royal Society B:*
Biological Sciences, 374(1781), 20180054. https://doi.org/10.1098/rstb.2018.0054
Hoff, S., Pendergast, C., Johnson, L., Olson, E., O'Dell, D., Dowling, Z. R., Gorman, K. M.,
Herzog, C., & Turner, W. C. (2023). Seasonal roost characteristics and fall behavior of
coastal populations of Northern Myotis (Myotis septentrionalis). *Journal of Mammalogy*,
gyad102. https://doi.org/10.1093/jmammal/gyad102
Hoyt, J. R., Kilpatrick, A. M., & Langwig, K. E. (2021). Ecology and impacts of white-nose
syndrome on bats. *Nature Reviews Microbiology, 19*(3), 196-210.
https://doi.org/10.1038/s41579-020-00493-5
Hoyt, J. R., Langwig, K. E., Sun, K., Parise, K. L., Li, A., Wang, Y., Huang, X., Worledge, L.,
Miller, H., White, J. P., Kaarakka, H. M., Redell, J. A., Görföl, T., Boldogh, S. A., Fukui,
D., Sakuyama, M., Yachimori, S., Sato, A., Dalannast, M., Jargalsaikhan, A., Batbayar,
N., Yovel, Y., Amichai, E., Natradze, I., Frick, W. F., Foster, J. T., Feng, J., & Kilpatrick,
A. M. (2020). Environmental reservoir dynamics predict global infection patterns and
population impacts for the fungal disease white-nose syndrome. *Proceedings of the*
National Academy of Sciences, 201914794. https://doi.org/10.1073/pnas.1914794117
Hufkens, K., Basler, D., Milliman, T., Melaas, E. K., & Richardson, A. D. (2018). An integrated
phenology modelling framework in r. *Methods in Ecology and Evolution, 9*(5), 1276-
1285. https://doi.org/10.1111/2041-210X.12970
Jackson, R. T., Willcox, E. V., & Bernard, R. F. (2022). Winter torpor expression varies in four
bat species with differential susceptibility to white-nose syndrome. *Scientific Reports,*
12(1), 5688. https://doi.org/10.1038/s41598-022-09692-x

Jackson, R. T., Willcox, E. V., Zobel, J. M., & Bernard, R. F. (2022). Emergence activity at hibernacula differs among four bat species affected by white-nose syndrome. *Ecology and Evolution, 12*(7), e9113. https://doi.org/10.1002/ece3.9113

Johnson, J. S., Reeder, D. M., McMichael, J. W., Meierhofer, M. B., Stern, D. W. F., Lumadue, S. S., Sigler, L. E., Winters, H. D., Vodzak, M. E., Kurta, A., Kath, J. A., & Field, K. A. (2014). Host, pathogen, and environmental characteristics predict white-nose syndrome mortality in captive little brown myotis (Myotis lucifugus). *PLOS ONE, 9*(11). https://doi.org/10.1371/journal.pone.0112502

Jonasson, K. A., & Willis, C. K. R. (2011). Changes in Body Condition of Hibernating Bats Support the Thrifty Female Hypothesis and Predict Consequences for Populations with White-Nose Syndrome. *PLOS ONE, 6*(6). https://doi.org/10.1371/journal.pone.0021061

Kailing, M. J., Hoyt, J. R., White, J. P., Kaarakka, H. M., Redell, J. A., Leon, A. E., Rocke, T. E., DePue, J. E., Scullon, W. H., Parise, K. L., Foster, J. T., Kilpatrick, A. M., & Langwig, K. E. (2023). Sex-biased infections scale to population impacts for an emerging wildlife disease. *Proceedings of the Royal Society B: Biological Sciences, 290*(1995), 20230040. https://doi.org/10.1098/rspb.2023.0040

Kasumovic, M. M., Bruce, M. J., Andrade, M. C. B., & Herberstein, M. E. (2008). Spatial and temporal demographic variation drives within-season fluctuations in sexual selection. *Evolution, 62*(9), 2316-2325. https://doi.org/10.1111/j.1558-5646.2008.00446.x

Klein, S. L. (2000). The effects of hormones on sex differences in infection: From genes to behavior. *Neuroscience & Biobehavioral Reviews, 24*(6), 627-638. https://doi.org/10.1016/S0149-7634(00)00027-0

Kokko, H., Gunnarsson, T. G., Morrell, L. J., & Gill, J. A. (2006). Why do female migratory birds arrive later than males? *Journal of Animal Ecology, 75*(6), 1293-1303. https://doi.org/10.1111/j.1365-2656.2006.01151.x

Kolata, G. B. (1977). Sexual Dimorphism and Mating Systems: How Did They Evolve? *Science, 195*(4276), 382-383.

Kunz, T. H., Wrazen, J. A., & Burnett, C. D. (1998). Changes in body mass and fat reserves in pre-hibernating little brown bats (Myotis lucifugus). *Ecoscience, 5*(1), 8-17. https://doi.org/10.1080/11956860.1998.11682443

Kurta, A., & Kunz, T. H. (1987). Size of bats at birth and maternal investment during pregnancy. *Symposia of the Zoological Society of London, 57*, 79-106.

Laggan, N. A., Parise, K. L., White, J. P., Kaarakka, H. M., Redell, J. A., DePue, J. E., Scullon, W. H., Kath, J., Foster, J. T., Kilpatrick, A. M., Langwig, K. E., & Hoyt, J. R. (2023). Host infection and disease-induced mortality modify species contributions to the environmental reservoir. *Ecology, 104*(10), e4147. https://doi.org/10.1002/ecy.4147

Langwig, K. E., Frick, W. F., Bried, J. T., Hicks, A. C., Kunz, T. H., & Marm Kilpatrick, A. (2012). Sociality, density-dependence and microclimates determine the persistence of populations suffering from a novel fungal disease, white-nose syndrome. *Ecology Letters, 15*(9), 1050-1057. https://doi.org/10.1111/j.1461-0248.2012.01829.x

Langwig, K. E., Frick, W. F., Hoyt, J. R., Parise, K. L., Drees, K. P., Kunz, T. H., Foster, J. T., & Kilpatrick, A. M. (2016). Drivers of variation in species impacts for a multi-host fungal disease of bats. *Philosophical Transactions of the Royal Society B: Biological Sciences, 371*(1709). https://doi.org/10.1098/rstb.2015.0456

Langwig, K. E., Frick, W. F., Reynolds, R., Parise, K. L., Drees, K. P., Hoyt, J. R., Cheng, T. L., Kunz, T. H., Foster, J. T., & Marm Kilpatrick, A. (2015). Host and pathogen ecology

drive the seasonal dynamics of a fungal disease, white-nose syndrome. *Proceedings of*
the Royal Society B: Biological Sciences, 282(1799).
https://doi.org/10.1098/rspb.2014.2335
Langwig, K. E., Hoyt, J. R., Parise, K. L., Frick, W. F., Foster, J. T., & Kilpatrick, A. M. (2017).
Resistance in persisting bat populations after white-nose syndrome invasion.
Philosophical Transactions of the Royal Society B: Biological Sciences, 372(1712),
20160044. https://doi.org/10.1098/rstb.2016.0044
Langwig, K. E., Paul White, J., Parise, K. L., Kaarakka, H. M., Redell, J. A., DePue, J. E.,
Scullon, W. H., Foster, J. T., Marm Kilpatrick, A., Hoyt, J. R., & Kate Langwig, C. E.
(2021). Mobility and infectiousness in the spatial spread of an emerging fungal pathogen.
J Anim Ecol, 90, 1134-1141. https://doi.org/10.1111/1365-2656.13439
Legendre, S., Clobert, J., Møller, A. P., & Sorci, G. (1999). Demographic Stochasticity and
Social Mating System in the Process of Extinction of Small Populations: The Case of
Passerines Introduced to New Zealand. *The American Naturalist, 153*(5), 449-463.
https://doi.org/10.1086/303195
Lehtonen, J., Parker, G. A., & Schärer, L. (2016). Why anisogamy drives ancestral sex roles.
Evolution, 70(5), 1129-1135.
Lenth, R. V. (2024). emmeans: Estimated Marginal Means, aka Least-Squares Means (Version R
package version 1.10.6). Retrieved from https://CRAN.R-project.org/package=emmeans
Lüdecke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P., & Makowski, D. (2021). performance:
An R Package for Assessment, Comparison and Testing of Statistical Models. *Journal of*
Open Source Software, 6(60). https://doi.org/10.21105/joss.03139
Luong, L. T., Perkins, S. E., Grear, D. A., Rizzoli, A., & Hudson, P. J. (2010). The relative
importance of host characteristics and co-infection in generating variation in
Heligmosomoides polygyrus fecundity. *Parasitology, 137*(6), 1003-1012.
https://doi.org/10.1017/S0031182009991892
Martin, A. M., Hogg, J. T., Manlove, K. R., LaSharr, T. N., Shannon, J. M., McWhirter, D. E.,
Miyasaki, H., Monteith, K. L., & Cross, P. C. (2022). Disease and secondary sexual
traits: effects of pneumonia on horn size of bighorn sheep. *The Journal of Wildlife*
Management, 86(1), e22154. https://doi.org/10.1002/jwmg.22154
Mazor, T., Doropoulos, C., Schwarzmueller, F., Gladish, D. W., Kumaran, N., Merkel, K., Di
Marco, M., & Gagic, V. (2018). Global mismatch of policy and research on drivers of
biodiversity loss. *Nature Ecology & Evolution, 2*(7), 1071-1074.
https://doi.org/10.1038/s41559-018-0563-x
McDonald, J. L., Smith, G. C., McDonald, R. A., Delahay, R. J., & Hodgson, D. (2014).
Mortality trajectory analysis reveals the drivers of sex-specific epidemiology in natural
wildlife–disease interactions. *Proceedings of the Royal Society B: Biological Sciences,*
281(1790). https://doi.org/10.1098/rspb.2014.0526
Morbey, Y. E., & Ydenberg, R. C. (2001). Protandrous arrival timing to breeding areas: a
review. *Ecology Letters, 4*(6), 663-673. https://doi.org/10.1046/j.1461-
0248.2001.00265.x
Muller, L. K., Lorch, J. M., Lindner, D. L., O'Connor, M., Gargas, A., & Blehert, D. S. (2013).
Bat white-nose syndrome: a real-time TaqMan polymerase chain reaction test targeting
the intergenic spacer region of Geomyces destructans. *Mycologia, 105*(2), 253-259.
https://doi.org/10.3852/12-242

O'Shea, T., Cryan, P., Cunningham, A., Fooks, A., Hayman, D. T. S., Luis, A., Peel, A., Plowright, R., & Wood, J. L. N. (2014). Bat Flight and Zoonotic Viruses. *Emerging Infectious Disease journal, 20*(5), 741. https://doi.org/10.3201/eid2005.130539

Oro, D., Álvarez, D., & Velando, A. (2018). Complex demographic heterogeneity from anthropogenic impacts in a coastal marine predator. *Ecological Applications, 28*(3), 612-621. https://doi.org/10.1002/eap.1679

Otto, S. P. (2018). Adaptation, speciation and extinction in the Anthropocene. *Proceedings of the Royal Society B: Biological Sciences, 285*(1891), 20182047. https://doi.org/10.1098/rspb.2018.2047

Perkins, S. E., Ferrari, M. F., & Hudson, P. J. (2008). The effects of social structure and sex-biased transmission on macroparasite infection. *Parasitology, 135*(13), 1561-1569. https://doi.org/10.1017/S0031182008000449

Péron, G. (2013). Compensation and additivity of anthropogenic mortality: life-history effects and review of methods. *Journal of Animal Ecology, 82*(2), 408-417. https://doi.org/10.1111/1365-2656.12014

Perry, R. W. (2013). A review of factors affecting cave climates for hibernating bats in temperate North America. *Environmental Reviews 21: 28-39, 21*, 28-39. https://doi.org/10.1139/er-2012-0042

Pfeiffer, B., & Mayer, F. (2013). Spermatogenesis, sperm storage and reproductive timing in bats. *Journal of Zoology, 289*(2), 77-85. https://doi.org/10.1111/j.1469-7998.2012.00970.x

Pikula, J., Amelon, S. K., Bandouchova, H., Bartonička, T., Berkova, H., Brichta, J., Hooper, S., Kokurewicz, T., Kolarik, M., Köllner, B., Kovacova, V., Linhart, P., Piacek, V., Turner, G. G., Zukal, J., & Martínková, N. (2017). White-nose syndrome pathology grading in Nearctic and Palearctic bats. *PLOS ONE, 12*(8), e0180435. https://doi.org/10.1371/journal.pone.0180435

Plesnar-Bielak, A., Skrzynecka, A. M., Prokop, Z. M., & Radwan, J. (2012). Mating system affects population performance and extinction risk under environmental challenge. *Proceedings of the Royal Society B: Biological Sciences, 279*(1747), 4661-4667. https://doi.org/10.1098/rspb.2012.1867

Promislow, D. E. (1997). Costs of sexual selection in natural populations of mammals. *Proceedings of the Royal Society of London. Series B: Biological Sciences, 247*(1320), 203-210. https://doi.org/10.1098/rspb.1992.0030

Promislow, D. E., Montgomerie, R., & Martin, T. E. (1992). Mortality Costs of Sexual Dimorphism in Birds. *Proceedings: Biological Sciences, 250*(1328), 143-150.

Racey, P., Uchida, T., Mori, T., Avery, M., & Fenton, M. (1987). Sperm-epithelium relationships in relation to the time of insemination in little brown bats (Myotis lucifugus). *Journal of reproduction and fertility, 80*(2), 445-454. https://doi.org/10.1530/jrf.0.0800445

Racey, P. A., & Entwistle, A. C. (2000). *9-Life-history and Reproductive Strategies of Bats*. London: Academic Press.

Reeder, D. M., Frank, C. L., Turner, G. G., Meteyer, C. U., Kurta, A., Britzke, E. R., Vodzak, M. E., Darling, S. R., Stihler, C. W., Hicks, A. C., Jacob, R., Grieneisen, L. E., Brownlee, S. A., Muller, L. K., & Blehert, D. S. (2012). Frequent Arousal from Hibernation Linked to Severity of Infection and Mortality in Bats with White-Nose Syndrome. *PLOS ONE, 7*(6), e38920. https://doi.org/10.1371/journal.pone.0038920

Rivers, N. M., Butlin, R. K., & Altringham, J. D. (2006). Autumn swarming behaviour of Natterer's bats in the UK: Population size, catchment area and dispersal. *Biological Conservation, 127*(2), 215-226. https://doi.org/10.1016/j.biocon.2005.08.010

Rosa, G. M., Bosch, J., Martel, A., Pasmans, F., Rebelo, R., Griffiths, R. A., & Garner, T. W. J. (2019). Sex-biased disease dynamics increase extinction risk by impairing population recovery. *Animal Conservation, 22*(6), 579-588. https://doi.org/10.1111/acv.12502

Rubalcaba, J. G., Gouveia, S. F., Villalobos, F., Cruz-Neto, A. P., Castro, M. G., Amado, T. F., Martinez, P. A., Navas, C. A., Dobrovolski, R., Diniz-Filho, J. A. F., & Olalla-Tárraga, M. Á. (2022). Physical constraints on thermoregulation and flight drive morphological evolution in bats. *Proceedings of the National Academy of Sciences, 119*(15), e2103745119. https://doi.org/10.1073/pnas.2103745119

Sæther, B.-E., Engen, S., Lande, R., Møller, A. P., Bensch, S., Hasselquist, D., Beier, J., & Leisler, B. (2004). Time to Extinction in Relation to Mating System and Type of Density Regulation in Populations with Two Sexes. *Journal of Animal Ecology, 73*(5), 925-934. https://doi.org/10.1111/j.0021-8790.2004.00869.x

Sauer, E. L., Connelly, C., Perrine, W., Love, A. C., & DuRant, S. E. (2024). Male pathology regardless of behaviour drives transmission in an avian host–pathogen system. *Journal of Animal Ecology, 93*(1), 36-44. https://doi.org/10.1111/1365-2656.14026

Schlaepfer, M. A., Runge, M. C., & Sherman, P. W. (2002). Ecological and evolutionary traps. *Trends in Ecology & Evolution, 17*(10), 474-480. https://doi.org/10.1016/S0169-5347(02)02580-6

Scialli, A. R. (1993). A comparative overview of mammalian fertilization: Dunbar BS, O'Rand MG. New York: Plenum Press; 1991. In: Pergamon.

Silk, M. J., Weber, N. L., Steward, L. C., Hodgson, D. J., Boots, M., Croft, D. P., Delahay, R. J., & McDonald, R. A. (2018). Contact networks structured by sex underpin sex-specific epidemiology of infection. *Ecology Letters, 21*(2), 309-318. https://doi.org/10.1111/ele.12898

Skorping, A., & Jensen, K. H. (2004). Disease dynamics: All caused by males? *Trends in Ecology and Evolution, 19*(5), 219-220. https://doi.org/10.1016/j.tree.2004.02.006

Stephenson, J. F. (2019). Parasite-induced plasticity in host social behaviour depends on sex and susceptibility. *Biology Letters, 15*(11). https://doi.org/10.1098/rsbl.2019.0557

Stephenson, J. F., Kinsella, C., Cable, J., & van Oosterhout, C. (2016). A further cost for the sicker sex? Evidence for male-biased parasite-induced vulnerability to predation. *Ecology and Evolution, 6*(8), 2506-2515. https://doi.org/10.1002/ece3.2049

Sweeny, A. R., Albery, G. F., Venkatesan, S. V., Fenton, A., & Pedersen, A. B. (2021). Spatiotemporal variation in drivers of parasitism in a wild wood mouse population. *Functional Ecology*, 1365-2435.13786. https://doi.org/10.1111/1365-2435.13786

Taylor, L. R. (1963). Analysis of the Effect of Temperature on Insects in Flight. *Journal of Animal Ecology, 32*(1), 99-117. https://doi.org/10.2307/2520

Thomas, D. W., Brock Fenton, M., & Barclay, R. M. R. (1979). Social behavior of the little brown bat, Myotis lucifugus. *Behavioral Ecology and Sociobiology, 6*(2), 129-136. https://doi.org/10.1007/BF00292559

Tilman, D., Clark, M., Williams, D. R., Kimmel, K., Polasky, S., & Packer, C. (2017). Future threats to biodiversity and pathways to their prevention. *Nature, 546*(7656), 73-81. https://doi.org/10.1038/nature22900

Trivers, R. L. (1972). Parental Investment and Sexual Selection. In B. G. Campbell (Ed.), *Sexual*
selection and the descent of man: 1871-1971. (Vol. 1871-1971). Chicago, IL: Aldine.
Verant, M. L., Boyles, J. G., Waldrep, W., Wibbelt, G., & Blehert, D. S. (2012). Temperature-
Dependent Growth of Geomyces destructans, the Fungus That Causes Bat White-Nose
Syndrome. *PLOS ONE, 7*(9), 46280. https://doi.org/10.1371/journal.pone.0046280
Visser, M. E., & Gienapp, P. (2019). Evolutionary and demographic consequences of
phenological mismatches. *Nature Ecology & Evolution, 3*(6), 879-885.
https://doi.org/10.1038/s41559-019-0880-8
Wai-Ping, V., & Fenton, M. B. (1988). Nonselective Mating in Little Brown Bats (Myotis
lucifugus). *Journal of Mammalogy, 69*(3), 641-645. https://doi.org/10.2307/1381364
Warnecke, L., Turner, J.M., Bollinger, T.K., Lorch, J.M., Misra, V., Cryan, P.M., Wibbelt, G.,
Blehert, D.S., & Willis, C.K.R. (2012). Inoculation of bats with European Geomyces
destructans supports the novel pathogen hypothesis for the origin of white-nose
syndrome, *Proceedings of the National Academy of Sciences*, *109*(18), 6999-7003,
https://doi.org/10.1073/pnas.1200374109
Webb, P. I., Speakman, J. R., & Racey, P. A. (1996). How hot is a hibernaculum? A review of
the temperatures at which bats hibernate. *Canadian Journal of Zoology, 74*(4), 761-765.
https://doi.org/10.1139/z96-087
Wells, K., Hamede, R. K., Kerlin, D. H., Storfer, A., Hohenlohe, P. A., Jones, M. E., &
McCallum, H. I. (2017). Infection of the fittest: devil facial tumour disease has greatest
effect on individuals with highest reproductive output. *Ecology Letters*.
https://doi.org/10.1111/ele.12776
Williams, C. T., Chmura, H. E., Deal, C. K., & Wilsterman, K. (2022). Sex-Differences in
Phenology: A Tinbergian Perspective. *Integrative and Comparative Biology, 62*(4), 980-
997. https://doi.org/10.1093/icb/icac035
Willis, C. K. R. (2017). Trade-offs Influencing the Physiological Ecology of Hibernation in
Temperate-Zone Bats. *Integrative and Comparative Biology, 57*(6), 1214-1224.
https://doi.org/10.1093/icb/icx087
Wimsatt, W. A. (1942). Survival of spermatozoa in the female reproductive tract of the bat. *The*
Anatomical Record, 83(2), 299-307. https://doi.org/10.1002/ar.1090830208
Wimsatt, W. A. (1944). Further studies on the survival of spermatozoa in the female
reproductive tract of the bat. *The Anatomical Record, 88*(2), 193-204.
https://doi.org/10.1002/ar.1090880206
Wimsatt, W. A. (1945). Notes on Breeding Behavior, Pregnancy, and Parturition in Some
Vespertilionid Bats of the Eastern United States. *Journal of Mammalogy, 26*(1), 23.
https://doi.org/10.2307/1375029
Zuk, M. (2009). The Sicker Sex. *PLoS Pathogens, 5*(1), e1000267-e1000267.
https://doi.org/10.1371/journal.ppat.1000267
Zuk, M., & McKean, K. A. (1996). Sex Differences in Parasite Infections: Patterns and
Processes. *International Journal for Parasitology, 26*(10), 1009-1024.
https://doi.org/10.1016/S0020-7519(96)80001-4