Activity patterns during the mating season predict sex-biased infections in an emerging

fungal disease

Macy J. Kailing[1], Joseph R. Hoyt[1], J. Paul White[2], Jennifer A. Redell[2], Heather M. Kaarakka[2],

and Kate E. Langwig[1]

1 Department of Biological Sciences, Virginia Polytechnic Institute, Blacksburg VA 24061

2 Wisconsin Department of Natural Resources, Madison WI 53707

<u>Abstract</u>

1) Mating dynamics can govern species impacts from rapid global change by influencing

population rates of growth and adaptation, as well as individual traits that affect mortality

risks from novel pressures.

2) Here, we examined sex differences in activity of *Myotis lucifugus* during their mating

season, which coincides with exposure to the lethal fungal pathogen (*Pseudogymnoascus*

destructans) that causes white-nose syndrome. We expected differences in activity

between sexes to modify seasonal *P. destructans* dynamics as the pathogen can replicate

only at the cool temperatures at which bats hibernate.

3) We used passive antenna systems installed at the entrances of hibernacula and PIT tags to

characterize activity patterns of bats. We also measured pathogen loads on bats during the

autumn mating and early hibernation periods to assess how infection changed according

to host phenology.

4) We found that females spent fewer days active during autumn, arrived after males, and

were primarily active on the warmest nights. Males remained highly active throughout

the mating season and later in autumn than females. Importantly, differences in

phenology corresponded to higher pathogen loads on females during early hibernation as

male activity, and thus warm body temperatures, inhibit pathogen growth.

5) Differences in activity between sexes and in the transition from swarm to hibernation

likely reflects males maximizing their mating opportunities while females conserve

energy to meet the cost of spring migration and reproduction. More broadly, our results

show how activity during the mating season and phenology can contribute to sex-biased

impacts of a novel disease and highlight the value of understanding species' mating

systems to anticipate the impacts of environmental change.

Key-words: sex-biased infection, mating, phenology, white-nose syndrome, bats,

seasonality, fungal disease, life history

1. Introduction

Global change has imposed novel pressures that threaten many species with extinction,

including habitat loss, pollution, overutilization and emerging infectious disease (Dirzo et al.,

2014; Mazor et al., 2018; Otto, 2018; Tilman et al., 2017). The resilience of populations to global

changes can vary based on the species' life history, including the mating system (Capdevila et

al., 2022; Péron, 2013; Plesnar-Bielak et al., 2012; Sæther et al., 2004). Mating systems can

govern how quickly populations recover from and adapt to novel stressors by influencing

demographic and evolutionary processes, such as rates of recruitment and genetic mixing (Bessa-

Gomes, Legendre, & Clobert, 2004; Legendre et al., 1999). Further, mating strategies,

characterized by the collection of activities and behaviors used by individuals to monopolize and

secure mates, can vary within species and be a substantial source of trait variation (Bradbury &

Vehrencamp, 1977; Clutton-Brock, 1989; Shuster & Wade 2003). Accordingly, traits expressed

during mating periods are often shared within demographic classes (e.g., sex, age, social status,

etc.) and can result in selective mortality that can restructure or destabilize populations (Clutton-

Brock et al., 2002; Grüebler et al., 2008; Martin et al., 2022; Oro, Álvarez, & Velando, 2018;

Rosa et al., 2019; Wells et al., 2017). Therefore, understanding how aspects of the mating system

influence interactions between individuals and novel pressures should improve the ability to

predict responses of populations to rapid changes.

Sexes use different strategies for maximizing their fitness that give rise to variation in

morphology, behaviors, or physiology (Kolata, 1977; Promislow, Montgomerie, & Martin, 1992;

Trivers, 1972), which collectively can affect their mortality risk (Promislow, 1997). In many

species where sexual selection leads to dimorphic traits between sexes, males invest in quantity

of offspring through copious mating, whereas females, who are more limited by gametic

production, invest in ensuring survival of few high-quality offspring (Bateman, 1948; Lehtonen,

Parker, & Schärer, 2016). Accordingly, males may be more likely to engage in risky behaviors to

increase their competitive mating advantages whereas females, whose fitness relies on surviving

to carry and care for young, may be more risk-averse (Bateman, 1948; Bonduriansky et al., 2008;

Promislow, 1997). Additionally, asynchrony between sexes in their reproductive investments can

also contribute to differences in how females and males respond to the same environmental

pressures (Cornwallis & Uller, 2010; Forsgren et al., 2004; Kasumovic et al., 2008), including

when their peak vulnerability to stressors occurs (Williams et al., 2022). While sex specific

responses to novel pressures are expected, the proximate factors that influence differences in

impacts between females and males are overlooked for many novel threats (Connallon, Débarre,

& Li, 2018; Connallon & Hall, 2018).

Novel pressures like infectious diseases frequently interact with mating systems resulting

in differential impacts to sexes (Herrera & Nunn, 2019; Stephenson, 2019; Stephenson et al.,

2016; Zuk, 2009; Zuk & McKean, 1996) that can influence the severity of epidemics and

coevolution between hosts and pathogens (Hall & Mideo, 2018). Typically, male behaviors or

physiological states optimize fitness at the costs of higher pathogen exposure or growth,

including territory defense or aggression, dispersal, and reduced immunity (Ferrari et al., 2003;

Grear, Luong, & Hudson, 2012; Luong et al., 2010; Sauer et al., 2024; Stephenson, 2019). For

example, adult male badgers had higher contacts with conspecifics and higher risks of acquiring

tuberculosis compared to females (Delahay et al., 2006; Graham et al., 2013; McDonald et al.,

2014). Male-biased disease is more frequently documented, but female-biased disease has been

shown to arise seasonally (Cattadori et al., 2005; Sweeny et al., 2021). For example, during

parturition adult female European rabbits had the sharpest peak in nematode intensities relative

to non-reproductive females and males (Cattadori et al., 2005), illustrating that the magnitude

and direction of sex-biases can vary temporally to affect the dynamics between hosts and

pathogens (Cattadori et al., 2005; Sweeny et al., 2021). However, the behavioral and

physiological mechanisms contributing to sex-biases in impacts from introduced pathogens are

often unclear, thus limiting opportunities for management.

White-nose syndrome (WNS) is a recently introduced disease of bats that is caused by the

psychrophilic fungal pathogen, *Pseudogymnoascus destructans*, and has led to population

declines exceeding 95% (Frick et al., 2015; Langwig et al., 2012; Langwig et al., 2016).

Euthermic bats that are active on the landscape maintain higher body temperatures (37-41°C)

(O'Shea et al., 2014) that are above the upper critical limit of *P. destructans* growth (20°C)

(Verant et al., 2012). However, when bats reduce their body temperature during torpor (short

periods of inactivity that comprise hibernation) to near ambient (1-16°C) (Perry, 2013; Webb,

Speakman, & Racey, 1996) the pathogen is able to grow, making it dependent on seasonal

hibernation cycles (Jackson, Willcox, & Bernard, 2022; Jackson et al., 2022; Langwig et al.,

2015; Langwig et al., 2021). Pathogen exposure occurs during autumn when bats return annually

to contaminated hibernation sites (Hoyt et al., 2020; Laggan et al., 2023) but infection and

mortality peak over winter (Hoyt, Kilpatrick, & Langwig, 2021; Langwig et al., 2015; Reeder et

al., 2012).

The mating system for temperate bats impacted by WNS is polygynandrous and

described as a swarm (Fenton 1969, Bradbury & Vehrencamp 1977). Swarming occurs during

autumn when large numbers of individuals aggregate near hibernacula entrances, then enter the

site to form smaller groups (typically pairs or triads), and copulate after landing on walls or in

crevices (Fenton, 1969; Fraser & McGuire, 2023; Thomas, Brock Fenton, & Barclay, 1979; Wai-

Ping & Fenton, 1988). While females mate during this period, they store sperm through

hibernation until spring emergence, when they migrate to summer maternity colonies, become

pregnant, and rear pups cooperatively with other females (Fenton & Robert, 1980; Kurta &

Kunz, 1987; Wimsatt, 1942, 1944, 1945). Males segregate from females during parturition and

pup rearing (Davis & Hitchcock, 1965; Fenton & Robert, 1980). To that end, females'

reproductive investments are primarily during spring and summer, and are more extensive in

magnitude and duration than males, whose reproductive investments are highest during autumn

(Racey & Entwistle, 2000; Fig 1). While both sexes mate during autumn, asynchronous

reproductive investments between females and males could lead to extensive behavioral and

physiological traits that contribute to disease impacts. Specifically, the congregation of sexes at

contaminated hibernacula during autumn and hibernation provide a unique opportunity to

explore how sex differences in activity during the mating and pathogen exposure period

influences infection intensity between sexes.

Figure 1. Hypothesized reproductive investment of females and males with different

reproductive strategies developed from Fenton & Robert, 1980, Racey & Entwistle, 2000, and

Wimsatt, 1945. Males are expected to invest most towards their reproductive fitness during

autumn when they mate with many females. Females likely require less energy during the

autumn swarm, but considerable investment is made following hibernation as they singly parent

their young in maternity colonies and require energy for gestation, parturition, and lactation.

Light blue points correspond to the tagging and sampling dates during the seasons of autumn

mating and early hibernation as described in Methods.

Temperate bats reduce activity seasonally and use torpor to conserve energy even outside of

winter hibernation (Alston et al., 2022). However, females and males balance torpor and activity

differently throughout the year (Grinevitch, Holroyd, & Barclay, 1995; Hoff et al., 2023). These

sex specific differences in activity and torpor could strongly influence WNS dynamics if sex-

specific torpor use influences growth of the cool-temperature dependent fungus. Yet, empirical

investigations are limited by the challenge of obtaining individually based data on activity

patterns through time. Previous research has shown that females can be more infected and suffer

greater impacts (Grieneisen et al., 2015; Kailing et al., 2023), but see Johnson et al., 2014.

However, the mechanisms driving the difference between sexes remain unknown. We

hypothesize that activity during the autumn mating season drives highly disparate infections in

early hibernation since the pathogen is unable to grow on bats when they are active. As such, we

investigated how traits associated with mating strategy, evaluated here based on activity during

the mating season, vary between females and males and contribute to sex-biased infection

intensity during seasons important for reproduction (autumn mating) and disease (autumn mating

and early hibernation). Specifically, we aimed to characterize the active period at swarm

locations (i.e., hibernacula), examine environmental and host factors associated with activity

during the mating season, and explore links between phenology and infection intensity.

2. Methods

2.1 Field monitoring

We studied the activity of *Myotis lucifugus* in Wisconsin, USA during the autumn mating

season each year from 2020 to 2023 at three large hibernacula within Pierce and Dodge counties,

consisting of 10 total entrances. We considered July 01 to November 01 as the autumn mating

period as activity patterns at the hibernacula are likely to be motivated by breeding. We used this

relatively broad timeframe to avoid missing the earliest arriving or latest swarming bats. At

every entrance, we installed an RFID system consisting of an antenna with an automated data

logger (IS10001; Biomark) that recorded a time-stamped datapoint each time a passive integrated

transponder (PIT) tag was in proximity. Each system was scheduled to run continuously by a

direct or solar source of power. Systems were programmed to record unique tags every minute to

help to maximize detections as >1 tag cannot be simultaneously recorded and bats sitting on top

of antenna would otherwise block detections of all other individuals. In addition, systems

recorded test tags at least once every 12 hours enabling us to determine when systems were

operational.

2.2 PIT tagging and measuring pathogen quantity

We captured bats at each site during autumn (between August 15 and September 02, Fig.

1) using a harp trap (Harptrap; Faunatech Austbat) and mist-nets (Polyester 38mm mesh; Avinet)

near the entrances where individuals were active. For every individual, we weighed, injected a

PIT tag, categorized sex based on external morphology, and attached an aluminum-lipped band

(2.9 mm; Porzana, Icklesham, UK) as secondary identification. Capture efforts at each site

concluded after one night, unless the target number of bats (approximately 100) was not

achieved, in which we trapped a second night.

We also sampled bats in early hibernation (between November 01 and November 17, Fig.

1) at the same sites by removing them from their roosting locations on the walls and ceilings of

the hibernacula. We walked sections of the hibernacula to collect a stratified sample of up to 20

bats. As in autumn, we recorded sex based on external morphology of every individual and

attached a band. We collected an epidermal swab from all the bats that were sampled in autumn

or early hibernation with a standardized swabbing technique that is used widely to assess

pathogen presence across seasons (Frick et al., 2015; Langwig et al., 2015). Immediately upon

collection, we stored swabs in RNAlater. We tested swab samples for *P. destructans* using

Qiagen DNeasy Blood and Tissue Kits (Qiagen, USA) to extract fungal DNA followed by qPCR

to quantify fungal loads (Langwig et al., 2015; Muller et al., 2013; McGuire et al., 2016). All bat

handling procedures were granted ethical approval by the Institutional Animal Care and Use

Committee at Virginia Tech (IACUC Protocols #20-150, #23-197), field work was conducted

under permits granted by the Wisconsin Department of Natural Resources (Endangered and

Threatened Species Permits #886, #882), and decontamination procedures followed United

States Fish and Wildlife recommendations.

2.3 Characterizing individual activity and climate variables

We determined individuals' active periods (i.e., the timing in which mating could occur

based on a bat being active at the swarm location) from PIT tag detections. An individual was

considered active if it was detected at any entrance to the site at least once in a night because bats

commonly visit multiple entrances throughout the night. We classified bats that were detected

between the hours of 00:00 and 07:00 as active on the previous date because bats are nocturnal,

and thus, their active date begins and ends at sunset and sunrise, respectively. We only included

bats detected on at least two nights per year to properly categorize the detections as being the

bats' first or last active date within the same year to capture the apparent swarm period. For each

bat observed within a year, we assigned a start swarm date as the first date detected after July 01

and an end swarm date as the last date detected before Nov 01. Start dates were only assigned for

individuals that returned from a previous year as we could not confirm when bats first arrived at

hibernacula if they were untagged. We were able to assign an end swarm date to all bats,

including those tagged in the same year, because the last detection always followed the date they

were captured and tagged. *Myotis lucifugus* can only be reliably aged as juveniles (young of the

year) or adults and all bats that return from a previous year are considered adults. Therefore, we

only included adults in all analyses.

To assess environmental factors associated with activity during the mating period, we

extracted climate data using the R package 'daymetr' (Hufkens et al., 2018) and GPS coordinates

of hibernation sites, which provided continuous, grid-based estimates of daily weather variables

based on location. Because most bats begin nightly activity at sunset before temperatures reach

an overnight low, the warm temperatures prior to sunset are relevant for activity. Therefore, we

calculated a mid-point temperature (Jackson et al., 2022) for each date as: (minimum

temperature on the following day + maximum temperature of current day)/2 to incorporate the

warm and cool temperatures that could influence bat activity and because mean temperatures

were not available in the climate data.

2.4 Statistical analyses

2.4.1 Summary

We tested our hypotheses using generalized (i.e., GLMM) and linear (i.e., LMM) mixed

effects models to characterize activity throughout the autumn mating period and pathogen

quantity by sex with R packages 'lme4' (Bates et al., 2015) and 'glmmTMB' (Brooks et al.,

2017). For each model, we tested variables based on *a priori* hypotheses and evaluated their

significance using ANOVA tables for statistical model objects from the 'car' package (Fox &

Weisberg, 2019). We combined site and year of sampling (site-year) to use as a random effect in

all models and randomize the intercepts of annual sampling events. Condensing to site-year was

more relevant for our analyses than using site and year separately because our sites are relatively

far apart and years vary in time since pathogen arrival. In analyses where individuals were

observed repeatedly in the same site-year, we included band or PIT tag ID as an additional

random effect. We compared candidate and null models using Akaike Information Criterion

(AIC) to evaluate relative quality (difference in AIC scores $|> 2|$; See Appendix 4.6.2). We also

visually checked model assumptions, such as the residual distribution of linear models, and

conducted k-fold cross-validation of logistic models to check predictive accuracy (Appendix

4.6.3) using the 'performance' package (Lüdecke et al., 2021). We also used the 'performance'

package (Lüdecke et al., 2021) and model outputs to visualize normality and verify the structure

of our random effects. Because many more males were detected than females, we also generated

a balanced dataset between sexes to check that the results informed by our male-biased dataset

were consistent and not due to the sampling bias. We truncated our full dataset by randomly

selecting the number of male observations equivalent to the number of female observations from

all the male observations (see Supplemental Methods). We extracted model estimates using R

package 'emmeans' (Lenth, 2022) to report final models. See Supplemental Table 1 for overview

of all analyses as well as Appendix with code for all statistical models, output, and model

validation.

238   *2.4.2 Dates of autumn mating activity*

We determined individually based active periods at swarm locations using LMMs. First,

we estimated when individuals of each sex started activity during autumn using the first date an

individual was detected at the hibernacula as a continuous response variable, with sex as a fixed

effect (F|M), and site-year and individual as random effects. Then, we used the same model

structure, but with last date detected as a response to estimate when individuals of each sex

ended activity. Prior to analyses, we converted dates to continuous monthly units with decimal

days by dividing the day of each month by 31 and adding it to numeric month. We then scaled

the dates to start at 0 on July 1. For example, July 15 corresponded to 0.5.

2.4.3 Effects of temperature and body condition on nightly activity

To test whether environmental conditions differently affected nightly activity between

sexes, we used a GLMM with a binomial distribution to test the probability of an individual

being active (=1) or undetected (=0) with temperature and sex as interactive fixed effects and

individual and site-year as random effects. For the response variable, we treated each bat as a

series of binomial trials using all the possible dates between the first and last detection where an

individual was detected (=1) or undetected (=0). We excluded nights where the minimum

temperature was less than 10°C to minimize outlier effects as bat activity is greatly reduced

(Gorman et al., 2021).

We also tested the effect of body condition on nightly activity to assess whether

energetics might contribute to differences in activity between males and females. As we could

only associate mass to bats at capture, for this dataset, we used the activity of bats in their first

year tagged. We used a GLMM with a binomial distribution to test the probability of a bat being

active. Like the above analysis, our response variable was detection probability, with each bat

treated as a series of binomial trials where it was either detected or undetected. We used sex,

mass, and temperature as interactive fixed effects and included site-year and individual as a

random effect.

2.4.4 Effects of length of active period and temperature on last detection

We examined how the number of days since arriving to the hibernaculum and

temperature affected whether sexes transition from being active to hibernating to understand how

host factors (time spent active) or environmental conditions (temperature) influence phenology.

First, we used a GLMM with a binomial distribution that predicted whether individuals stopped

activity for the year. For the response variable, detections of each bat were treated as a series of

binomial trials where an individual was detected for the last time (=1) or was detected on a

subsequent date (=0). We included interactive effects of sex and time since their first detection

(to standardize differences in start date) with site-year and individual as random effects. For days

since first detection, we binned dates to 3-day intervals and counted the number of intervals that

passed since their arrival to the hibernacula. Accordingly, we only included individuals that we

were able to assign start dates (i.e., returned from a previous year). Date intervals were used to

overcome sparsity in the number of females that are active each night.

To also examine the effect of temperature on the conclusion of autumn activity, we used

a GLMM with a binomial distribution to test a similar response as above where each bats'

detections were treated as a series of binomial trials and an individual was detected for the last

night (=1) or was detected on a subsequent night (=0). We included sex and temperature as

interacting fixed effects and random effects of site-year and individual. We tested length of

swarm period and temperature interacting with sex in separate models given collinearity between

the two predictors.

284   *2.4.5 Seasonality and host phenology effects on pathogen quantity*

To understand infection dynamics according to host phenology, we tested differences in

pathogen quantity between sexes and across seasons using a LMM with $\log_{10}$ *P. destructans*

loads as our response variable, with sex and date (in monthly units as above) as fixed effects, and

site-year as a random effect. Because we did not resample the same bats between autumn and

early hibernation within the site-year, we did not include individual ID as a random effect. We

explored associations between phenology and infection intensity further by testing the effect of

the active period on early hibernation pathogen quantity. We used a LMM to test $\log_{10} P.$

destructans loads as a response variable with median last date of activity as a fixed effect. We

included site-year as a random effect. We calculated the median date of last detection in autumn

for both sexes for each site and year (i.e., the end of swarm activity and assumed start of

hibernation) and assigned it to individuals that we sampled for pathogen quantity in early

hibernation of the respective site and year. We used the median date of last detection to minimize

the weight of outlier individuals that may leave the hibernacula late in autumn but are not

representative of swarm activity. We also performed this analysis at the population level using

the mean early hibernation pathogen quantity from each site and year as the response. We used

these approaches because genetic analyses indicate that swarming and hibernating populations at

a site are generally comprised of the same individuals (van Schaik et al., 2015) but relocating and

sampling the same individuals across seasons is rarely achieved, in part due to the complexity of

passages and high ceilings of hibernacula. We also evaluated autumn temperature as a predictor

of pathogen quantity.

3. Results

We characterized autumn activity of 148 (by site: 37, 40, 71) female and 608 (by site:

225, 155, 228) male *M. lucifugus* and measured pathogen quantity during capture on 168 (by

site: 56, 63, 49) female and 416 (by site: 151, 130, 135) males between 2020 and 2023. See

Supplemental Table 1 for overview of all analyses and Appendix 4.7 for full sample lists used in

each analysis by site, years, and sex. We also provide a visual representation of the full dataset in

Supplemental Fig. 1 (Appendix 4.1). All statistical results are reported with males as the

intercept or reference level and thus coefficient ± SE are in reference to the value above or below

the male references. Females started swarm activity later (Female coef ± SE = 0.249 ± 0.08, t =

3.25, p-value = 0.001; Fig. 2, top panel; Appendix 1.1) but also ended activity earlier than males

(-0.387 ± 0.06, t = -6.29, p-value < 0.001; Fig. 2, top panel; Appendix 1.2). Male activity fully

encompassed the active period of females (Fig. 2, bottom panel; Appendix 1.3) and the number

of individuals detected was overwhelmingly male biased throughout autumn, given their higher

abundance and activity overall (Supplemental Fig. 1; Appendix 4.1), and was consistent between

sampling events (Supplemental Table 2; Appendix 4.6.1).

Within their respective active periods, the probability of being active in a night or not for

each sex was differently affected by temperature (detected=0|1; Female*temperature = 0.151 ±

0.03, t = 4.97, p-value < 0.001, Fig. 3A; Appendix 2.1), such that the probability of females

being active was lower overall but increased more quickly with increasing temperatures

(Temperature slope estimated using 'emtrends': Female slope = 0.152 ± 0.03 vs Male slope

0.001 ± 0.01, Fig. 3A; Appendix 2.1), notably above 16°C. An interaction between sex and

temperature was also evident when using a balanced dataset of males and females (see

Supplemental Statistical Methods, Supplemental Fig. 2A; Appendix 4.2.1). We found that the

effect of body condition on activity differed between males and females, with light females less

active at cooler temperatures and more active at warmer temperatures, whereas all males

maintained high activity despite large differences in temperatures (Female

coef*mass*temperature ± SE = -0.121 ± 0.06, z = -2.07, p-value = 0.039, Supplemental Fig. 3,

Appendix 4.3).

We also found support for an interaction between the length of the active period and sex

on the probability of being detected for the last time (Date intervals since arrival*female = 0.367

$\pm\ 0.07$, $z = 5.49$, p-value < 0.001, Fig. 3B; Appendix 2.2), as well as an interactive effect of

temperature and sex on the probability of being detected for the last time, such that females were

significantly more likely to end their active period as temperatures decreased than males

(Temperature*female $= -0.164 \pm 0.08$, $z = -2.13$, p-value $= 0.033$, Fig. 3C; Appendix 2.3).

Observationally, we also found that within females' approximate 15-day active period, they were

active an average of 4 nights, whereas within males' 45-day active period they averaged 15

active nights (Supplemental Figure 5, Appendix 4.5).

There was no clear sex bias in pathogen quantity throughout autumn (Female $= -0.014 \pm$

0.10, $t = -0.14$; p-value $= 0.887$, Appendix 3.1), however, following autumn swarm, female

pathogen loads increased more rapidly, and females had significantly higher pathogen loads than

males by early hibernation (Female*date $= 0.340 \pm 0.08$, $t = 4.50$, p-value < 0.001; Fig. 4A;

Appendix 3.2). Ending of autumn activity of males and females generally varied among sites and

years and earlier end dates were associated with higher pathogen loads on individuals (Median

last date of activity slope on pathogen loads: -1.574 ± 0.41, $t = -3.87$, p-value < 0.001; Fig. 4B;

Appendix 3.3) as well as site-level pathogen quantities in early hibernation (Median last date of

activity slope on mean pathogen quantity: -1.562 ± 0.33, $t = -4.75$, p-value < 0.001;

Supplemental Fig. 4; Appendix 4.4). Autumn temperature generally lacked support as an

important predictor of pathogen quantity in early hibernation.

Figure 2. Top: Differences in the autumn active period by sex. Each point represents the date of starting (first detection) and ending (last detection) activity during autumn mating for an individual bat. Black circles represent model estimated dates with standard errors shown with horizontal bars. **Bottom:** Distribution of activity by sex. Each density plot shows the proportion of all detections grouped by sex that occurred on each date. An individual bat can be represented on multiple nights if activity occurred. Asymmetry in the peak of male activity (based on the total number of observations) relative to their estimated active period in the top panel illustrates visually that while most males arrive early in autumn, they maintain activity into later swarm dates.

Figure 3A. Effect of temperature on the probability that an individual was active in a night

throughout the swarm period. Each point represents the proportion of individuals from a site and

year that were active at each temperature by sex. Females are shown in purple and males in

orange and lines represent the model predicted probabilities of being active across the

temperature gradient. **Figure 3B.** Relationship between length of the active period in days and

the probability of last annual detection for females in purple and males in orange. Lines denote

the model predicted probabilities of being detected for the last time within a year. Actual data are

binary (1|0) such that 1 = ends activity and 0 = continues to be active. For visualization, points

show the proportions of bats of each sex that were detected for the last time within the year,

binned across a three-day interval. **Figure 3C**. Relationship between temperature and probability

that bats ended their autumn activity. Each point summarizes the proportion of bats that were

detected for the last time at a given temperature for each site and year. Females are shown in

purple and males are in orange. Solid lines denote the model predicted probability of being

detected for the last time for each sex. Size of points show the number of bats represented in

proportion calculations. Notably, males do not end their swarm activity even at temperatures that

are relatively low. In each panel, shaded grey regions show standard errors of model predictions.

Figure 4A: Pathogen quantity measured on adults of each sex throughout the autumn mating and

early hibernation periods. Each point represents an individual bat with females in purple and

males in orange. Solid lines denote model predicted fungal loads over time for females and males

respectively, with grey shaded regions displaying the standard error. Dashed lines represent the

model predicted end dates of activity as estimated in Fig. 2. There was no clear difference in

pathogen quantity between sexes during the active mating season in autumn, but female

pathogen loads were nearly an order of magnitude higher than male pathogen loads by early

hibernation. **Figure 4B.** Association between the median date of last detection at a site and early

hibernation pathogen quantities on individuals. Each point represents an individual bat sampled

in early hibernation colored by sex as in panel A and are jittered slightly for visualization.

Median last date of autumn activity was calculated for both sexes during each sampling event

(i.e., site-year) and used to minimize the outlier effects of a few individuals with exceptionally

early or late last active dates. Black line shows model predicted pathogen loads with standard

errors displayed with the grey shading. In each panel, date units have been converted to calendar

days for clarity.

4. Discussion

Sex specific reproductive strategies are common across species, yet how aspects of

mating systems give rise to sex-biased disease impacts are often unexplored. Our results suggest

that asynchronous swarm phenology between sexes may be a key driver of seasonally sex-biased

infection intensity, which are known to affect pathogen transmission and population impacts in

other host-pathogen systems (Ferrari et al., 2003; Graham et al., 2013; Grear, Luong, & Hudson,

2012; Perkins, Ferrari, & Hudson, 2008; Sauer et al., 2024; Silk et al., 2018; Skorping & Jensen,

2004). We show female *M. lucifugus* begin swarm activity later and end earlier, spend fewer

nights active after arriving to swarm, and have a more pronounced increase in activity on warmer

nights compared to males. Importantly, pathogen quantity did not differ between sexes until after

the active period of autumn mating, when bats transition to hibernation, and earlier end dates of

autumn activity were associated with higher pathogen quantities by early hibernation. Earlier use

of torpor by females likely contributes to female-biased infection intensity in early hibernation as

it allows an extended period for the pathogen to replicate as opposed to males that remain active

and can minimize pathogen replication through euthermia.

In exploring within activity period differences, we found that females are far less active

during their shorter swarm period compared to males. This is probably best explained by the

need for females to conserve enough energy stores to survive hibernation. Females store sperm

overwinter and then give birth and raise offspring following emergence (Czenze, Jonasson, &

Willis, 2017; Jonasson & Willis, 2011; Wimsatt, 1945). We provide two new lines of

individually derived evidence that aligns with females using energy more conservatively than

males. First, we found a clear effect of temperature differently affecting the probability that

females and males are active on a given night within their swarm period, and the relationship

changed depending on the body condition of females but not males (Supplemental Fig. 3,

Appendix 4.3). This suggests that females may be prioritizing the limited days they spend active

in autumn to mate and forage on warmer nights that average nearly 20°C (Supplemental Fig. 2B,

Appendix 4.2.2) likely because thermoregulating is more efficient at warmer temperatures

(Fjelldal et al., 2023; Rubalcaba et al., 2022) and prey resources are more active (Grüebler,

Morand, & Naef-Daenzer, 2008; Taylor, 1963). Second, we found that females on average,

regardless of their arrival date, begin transitioning into hibernation (i.e., remained in the

site/stopped activity) after being active on four nights within their estimated 15 day period,

whereas males maintained nightly activity until late in autumn regardless of the date they arrived

to swarm, and averaged over 13 active nights (Supplemental Figs 1 and 5, Appendix 4.1 and 4.5)

within their 45 day active period. Males that are less constrained by reproductive energy budgets

were active more than three times as many nights as females (Supplemental Figs 1 and 5;

Appendix 4.1 and 4.5), when temperatures are relatively low (Figs 3A and 3C, Appendix 2.1 and

2.3), and activity was independent of body condition (in contrast to females, Supplemental Fig.

3), suggesting that males are optimizing mating opportunities over energy conservation and

supports previous hypotheses (Burns & Broders, 2015; Glover & Altringham, 2008; Rivers,

Butlin, & Altringham, 2006; Willis, 2017). Importantly, the disparate reproductive costs between

sexes are likely contributing to how each sex interacts with the same novel pressure of an

introduced pathogen.

The earlier arrival of males to mating locations than females is observed across many

taxa, and two hypotheses have been posed to justify its ubiquity (Morbey & Ydenberg, 2001).

First, the 'rank advantage hypothesis' states the earliest arriving individuals have access to the

best resources or territories, providing a competitive advantage in acquiring females. Second, the

'mate opportunity hypothesis' suggests that earlier arriving males can increase the number of

females mated, resulting in greater annual reproductive success. In systems under strong sexual

selection and in cases where females also have resource-based advantages of arriving early, mate

opportunity has more reliably explained the early arrival of males than rank advantage,

particularly in species exhibiting sperm competition and highly male-biased sex ratios (Kokko et

al., 2006), which is known of temperate bats (Racey et al., 1987; Racey & Entwistle, 2000).

Sperm storage by females can also generate advantages for males that more closely synchronize

copulations with female receptivity, such that males that mate earlier can secure storage sites in

uteri that improve fertilization success (Scialli, 1993). Thus, we hypothesize that the longer

swarm period of males is driven by the advantage of increasing copulations with more females in

addition to elevating fertilization potential, as suggested by other studies of temperate bats

(Burns & Broders, 2015; Pfeiffer & Mayer, 2013). Other environmentally mediated mechanisms

likely function in concert with mating advantages. Females migrate from summer maternity

colonies and, because foraging to store fat for hibernation is essential for bats in autumn (Kunz,

Wrazen, & Burnett, 1998), timing the departure from diminishing resources in summer habitats

to optimal resource conditions near swarm locations is likely a more important balance for

females that have greater annual energetic constraints than males.

As in other disease systems, sex-specific differences in behavior or physiology during

periods of pathogen exposure can result in sex-biased disease impacts and govern host-pathogen

dynamics (Guerra-Silveira & Abad-Franch, 2013). We showed that during the season that bats

mate and are initially exposed to *P. destructans* after having cleared prior infections (Langwig et

al., 2015), activity differs markedly between sexes, however it did not correspond to differences

in pathogen quantity within that period (Appendix 3.1) as we expected based on other disease

systems (Ferrari et al., 2003; Graham et al., 2013; Guerra-Silveira & Abad-Franch, 2013; Klein,

2000). Furthermore, infections during the autumn mating season or differences in tolerance

between sexes are not expected to influence activity patterns because the pathogen quantity on

hosts during the active period are likely too low to initiate behaviors or physiological responses

symptomatic of disease (Warnecke et al., 2012). Rather, reproductive demands that influence

broader phenological patterns are more likely to be shaping sex-biased disease through

physiological mechanisms. Female infections rapidly increased almost two orders of magnitude

within the two months between the end of their active period and early hibernation (Fig. 4A,

Appendix 3.2), suggesting the timing of exposure and pathogen growth likely have temporally

decoupled effects on sex-specific disease (Gagnon et al., 2025) in this host-pathogen system. We

found that ending activity earlier was associated with higher pathogen loads in early hibernation,

which aligns with females being the earliest to hibernate and develop the highest pathogen loads

(Fig. 4B, Appendix 3.3). Ending activity is significant in this system because the pathogen can

only replicate once bats begin hibernation, and hosts that exceed an infection threshold earlier

have lower over-winter survival (Langwig et al., 2016). Consequently, females may be

predisposed to have higher disease than males because they initiate hibernation earlier, likely

because of their need to conserve energy to achieve reproductive fitness (Czenze et al., 2017;

Gagnon et al., 2025).

Phenology evolves to synchronize important biological events, such as reproduction, to

optimal conditions and resources (Visser & Gienapp, 2019). However, rapid changes in the

biotic or abiotic environment can modify which traits are beneficial or maladaptive (Schlaepfer,

Runge, & Sherman, 2002). For bats affected by WNS, if swarm periods evolved to maximize

survival and recruitment, disease-induced mortality of early arriving and less active individuals

may be selecting for traits that are beneficial for surviving infection but potentially maladaptive

for recruitment. Alternatively, if phenology is a plastic trait in bats as it is in other taxa (de

Villemereuil et al., 2020), a shift to a later swarm period may minimize disease while still

supporting recruitment, thus ameliorating population impacts. Given the clear, primarily additive

effect of sex on the timing of activity during the mating period (Fig. 2; Supplemental Table 2;

Appendix 4.6.1), we expect that any shifts in activity of one sex will be paralleled by the other.

Because females have a more restricted activity period and energy budgets, we expect females to

be less flexible in their phenology than males. As such, females will likely need to adapt

alternative mechanisms to overcome novel disease pressure, like the reduction in fungal invasion

of wing tissue found by Gagnon et al., 2025, or compensatory mechanisms to maintain

recruitment if a later swarm window is suboptimal for fitness.

Environmental conditions may also alleviate impacts on females throughout their

geographic range or between years. Female bats in milder climates, such as at lower latitudes or

years with warmer autumns, may not face the same restriction of suboptimal temperatures that

cause bats to begin hibernation. In such instances, females may stay active longer to avoid

developing infections severe enough to cause mortality. Similarly, female-biased infections are

observed across most host species (Kailing et al., 2023) that share similar reproductive cycles as

in Fig. 1. However, more subtle interspecific differences in phenology, like those explored in this

study, are largely unknown for most species impacted by WNS but may plausibly contribute to

the known variation in early hibernation infections and impacts among species (Langwig et al.,

2016; Langwig et al., 2015). Compared to species that mate earlier, species with later swarm

activity and delayed hibernation may have lower infections overall that allow more females to

survive winter. Previous work has implicated female-biased disease in an initial reduction in the

proportion of females following WNS invasion (Kailing et al., 2023), however, many colonies

where the pathogen invaded earliest are recovering (Grimaudo et al., 2021; Hoyt, Kilpatrick, &

Langwig, 2021; Langwig et al., 2017). Therefore, long-term ecological and evolutionary

responses of each sex to disproportionate disease impacts warrant further exploration (Gagnon et

al 2025), particularly at spatial and temporal scales beyond this study, as they may contribute to

how populations persist.

While sex-based differences in disease have received considerable attention, the

mechanisms are often unknown. We demonstrate that sex-specific phenological transitions

following exposure can facilitate sex-biased infection intensity, providing a unique example in

which physiological asynchronies decouple impacts of the timing of acquiring a pathogen and

disease progression between sexes. Importantly, our findings reveal a potentially significant

influence of mating dynamics on the trajectory of epidemics and pathways of host persistence or

extinction. Other novel pressures that similarly act more strongly on female mating dynamics

may impede population recoveries if sex-ratios are disrupted or if traits beneficial for survival are

antagonistic for recruitment. Given the importance of demographic structure and recruitment in

population impacts, determining how mating strategies influence responses to novel stressors

will be critical for conserving species in the face of increasing global change.

Acknowledgements

We thank Kirsten Fuller and Steffany Yamada for data curation support, and Alexander T.

Grimaudo, Nichole A. Laggan, Ariel E. Leon, and Tonie E. Rocke for support in field data

collection. We also thank University of Wisconsin Milwaukee Field Station and Gretchen Meyer

for site access.

Conflict of interest

All authors declare no conflict of interest.

Author contributions

Macy J. Kailing, Kate E. Langwig, and Joseph R. Hoyt conceived the ideas and designed

methodology; Macy J. Kailing, Joseph R. Hoyt, J. Paul White, Jennifer A. Redell, Heather M.

Kaarakka, and Kate E. Langwig collected the data; Macy J. Kailing analyzed data with input

from Kate E. Langwig and Joseph R. Hoyt. Macy J. Kailing wrote the original manuscript with

Kate E. Langwig and Joseph R. Hoyt. All authors contributed critically to the drafts and gave

final approval for publication.

Data availability statement

The datasets generated in this study are available from the Dryad Digital Repository:

https://doi.org/10.5061/dryad.rjdfn2zq6 (Kailing et al 2025). Exact site locations are not

disclosed to protect endangered species and landowners.

<u>References</u>

Alston, J. M., Dillon, M. E., Keinath, D. A., Abernethy, I. M., & Goheen, J. R. (2022). Daily

torpor reduces the energetic consequences of microhabitat selection for a widespread bat.

Ecology, 103(6), e3677. https://doi.org/10.1002/ecy.3677

Bateman, A. J. (1948). Intra-sexual selection in Drosophila. *Heredity, 2*(3), 349-368.

https://doi.org/10.1038/hdy.1948.21

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software, 67*(1), 1 - 48.

https://doi.org/10.18637/jss.v067.i01

Bessa-Gomes, C., Legendre, S., & Clobert, J. (2004). Allee effects, mating systems and the

extinction risk in populations with two sexes. *Ecology Letters, 7*(9), 802-812.

https://doi.org/10.1111/j.1461-0248.2004.00632.x

Bonduriansky, R., Maklakov, A., Zajitschek, F., & Brooks, R. (2008). Sexual Selection, Sexual

Conflict and the Evolution of Ageing and Life Span. *Functional Ecology, 22*(3), 443

Bradbury, J. W., & Vehrencamp, S. L. (1977). Social organization and foraging in emballonurid

bats. *Behavioral Ecology and Sociobiology, 2*(1), 1-17. doi:10.1007/BF00299284

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A.,

Skaug, H.J., Maechler, M., Bolker, B.M. (2017). "glmmTMB Balances Speed and

Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling." *The*

R Journal, *9*(2), 378–400. https://doi.org/10.32614/RJ-2017-066.

Brunet-Rossinni, A. K., & Wilkinson, G. S. (2009). Methods for Age Estimation and the Study

of Senescense in Bats. In T. H. Kunz & S. Parsons (Eds.), *Ecological and Behavioral*

Methods for the Study of Bats. Baltimore: The Johns Hopkins University Press.

Burns, L. E., & Broders, H. G. (2015). Maximizing mating opportunities: higher autumn

swarming activity in male versus female Myotis bats. *Journal of Mammalogy, 96*(6),

1326-1336. https://doi.org/10.1093/jmammal/gyv141

Capdevila, P., Stott, I., Cant, J., Beger, M., Rowlands, G., Grace, M., & Salguero-Gómez, R.

(2022). Life history mediates the trade-offs among different components of demographic

resilience. *Ecology Letters, 25*(6), 1566-1579. https://doi.org/10.1111/ele.14004

Cattadori, I. M., Boag, B., Bjørnstad, O. N., Cornell, S. J., & Hudson, P. J. (2005). Peak shift and

epidemiology in a seasonal host-nematode system. *Proceedings of the Royal Society B:*

Biological Sciences, 272(1568), 1163-1169. https://doi.org/10.1098/rspb.2004.3050

Clutton-Brock, T. H. (1997). Review Lecture: Mammalian mating systems. *Proceedings of the*

Royal Society of London. B. Biological Sciences, 236(1285), 339-372.

doi:10.1098/rspb.1989.0 027

Clutton-Brock, T. H., Coulson, T. N., Milner-Guiland, E. J., Thomson, D., & Armstrong, H. M.

(2002). Sex differences in emigration and mortality affect optimal management of deer

populations. *Nature, 415*(6872), 633-637. https://doi.org/10.1038/415633a

Connallon, T., Débarre, F., & Li, X.-Y. (2018). Linking local adaptation with the evolution of

sex differences. *Philosophical Transactions of the Royal Society B: Biological Sciences,*

373(1757), 20170414. https://doi.org/10.1098/rstb.2017.0414

Connallon, T., & Hall, M. D. (2018). Environmental Changes and Sexually Antagonistic

Selection. In *Encyclopedia of Life Sciences* (pp. 1-7).

Cornwallis, C. K., & Uller, T. (2010). Towards an evolutionary ecology of sexual traits. *Trends*

in Ecology & Evolution, 25(3), 145-152. https://doi.org/10.1016/j.tree.2009.09.008

Czenze, Z. J., Jonasson, K. A., & Willis, C. K. R. (2017). Thrifty females, frisky males: Winter

energetics of hibernating bats from a cold climate. *Physiological and Biochemical*

Zoology, 90(4), 502-511. https://doi.org/10.1086/692623

Davis, W. H., & Hitchcock, H. B. (1965). Biology and migration of the bat, Myotis lucifugus, in

New England. *Journal of Mammalogy, 46*(2), 296-313.

de Villemereuil, P., Charmantier, A., Arlt, D., Bize, P., Brekke, P., Brouwer, L., Cockburn, A.,

Côté, S. D., Dobson, F. S., Evans, S. R., Festa-Bianchet, M., Gamelon, M., Hamel, S.,

Hegelbach, J., Jerstad, K., Kempenaers, B., Kruuk, L. E. B., Kumpula, J., Kvalnes, T.,

McAdam, A. G., McFarlane, S. E., Morrissey, M. B., Pärt, T., Pemberton, J. M.,

Qvarnström, A., Røstad, O. W., Schroeder, J., Senar, J. C., Sheldon, B. C., van de Pol,

M., Visser, M. E., Wheelwright, N. T., Tufto, J., & Chevin, L.-M. (2020). Fluctuating

optimum and temporally variable selection on breeding date in birds and mammals.

Proceedings of the National Academy of Sciences, 117(50), 31969-31978.

https://doi.org/10.1073/pnas.2009003117

Delahay, R. J., Walker, N. J., Forrester, G. J., Harmsen, B., Riordan, P., Macdonald, D. W.,

Newman, C., & Cheeseman, C. L. (2006). Demographic correlates of bite wounding in

Eurasian badgers, Meles meles L., in stable and perturbed populations. *Animal*

Behaviour, 71(5), 1047-1055. https://doi.org/10.1016/j.anbehav.2005.07.018

Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J. B., & Collen, B. (2014).

Defaunation in the Anthropocene. *Science, 345*(6195), 401-406.

https://doi.org/10.1126/science.1251817

Fenton, M. B. (1969). Summer activity of Myotis lucifugus (Chiroptera:Vespertilionidae) at

hibernacula in Ontario and Quebec. *Canadian Journal of Zoology, 47*(4), 597-602.

https://doi.org/10.1139/z69-103

Fenton, M. B., & Robert, M. R. B. (1980). Myotis lucifugus. *Mammalian Species*(142), 1-8.

https://doi.org/10.2307/3503792

Ferrari, N., Cattadori, I. M., Nespereira, J., Rizzoli, A., & Hudson, P. J. (2003). The role of host

sex in parasite dynamics: field experiments on the yellow-necked mouse Apodemus

flavicollis. *Ecology Letters, 7*(2), 88-94. https://doi.org/10.1046/j.1461-

0248.2003.00552.x

Fjelldal, M. A., Muller, A. S., Ratikainen, I. I., Stawski, C., & Wright, J. (2023). The small-bat-

in-summer paradigm: Energetics and adaptive behavioural routines of bats investigated

through a stochastic dynamic model. *Journal of Animal Ecology, 92*(10), 2078-2093.

https://doi.org/10.1111/1365-2656.13999

Forsgren, E., Amundsen, T., Borg, Å. A., & Bjelvenmark, J. (2004). Unusually dynamic sex

roles in a fish. *Nature, 429*(6991), 551-554. doi:10.1038/nature02562

Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression* (3rd ed.). Thousand

Oaks, CA: Sage.

Fraser, E. E., & McGuire, L. P. (2023). Prehibernation swarming in temperate bats: a critical

transition between summer activity and hibernation. *Canadian Journal of Zoology,*

101(6), 408-422. https://doi.org/10.1139/cjz-2022-0129

Frick, W. F., Puechmaille, S. J., Hoyt, J. R., Nickel, B. A., Langwig, K. E., Foster, J. T., Barlow,

K. E., Bartonička, T., Feller, D., Haarsma, A.-J., Herzog, C., Horáček, I., Kooij, J.,

Mulkens, B., Petrov, B., Reynolds, R., Rodrigues, L., Stihler, C. W., Turner, G. G., &

Kilpatrick, A. M. (2015). Disease alters macroecological patterns of North American

bats. *Global Ecology & Biogeography, 24*(7), 741-749. https://doi.org/10.1111/geb.12290

Glover, A. M., & Altringham, J. D. (2008). Cave selection and use by swarming bat species.

Biological Conservation, 141(6), 1493-1504.

https://doi.org/10.1016/j.biocon.2008.03.012

Gorman, K. M., Barr, E. L., Ries, L., Nocera, T., & Ford, W. M. (2021). Bat activity patterns

relative to temporal and weather effects in a temperate coastal environment. *Global*

Ecology and Conservation, 30, e01769. https://doi.org/10.1016/j.gecco.2021.e01769

Graham, J., Smith, G. C., Delahay, R. J., Bailey, T., McDonald, R. A., & Hodgson, D. (2013).

Multi-state modelling reveals sex-dependent transmission, progression and severity of

tuberculosis in wild badgers. *Epidemiology and Infection, 141*(7), 1429-1436.

https://doi.org/10.1017/S0950268812003019

Grear, D. A., Luong, L. T., & Hudson, P. J. (2012). Sex-biased transmission of a complex life-

cycle parasite: Why males matter. *Oikos, 121*(9), 1446-1453.

https://doi.org/10.1111/j.1600-0706.2012.20358.x

Grieneisen, L. E., Brownlee-Bouboulis, S. A., Johnson, J. S., & Reeder, D. M. (2015). Sex and

hibernaculum temperature predict survivorship in white-nose syndrome affected little

brown myotis (Myotis lucifugus). *Royal Society Open Science, 2*.

https://doi.org/10.1098/rsos.140470

Grimaudo, A. T., Hoyt, J. R., Yamada, S. A., Herzog, C. J., Bennett, A. B., & Langwig, K. E.

(2021). Host traits and environment interact to determine persistence of bat populations

impacted by white-nose syndrome. *Ecology Letters, 25*(2), 483-497.

https://doi.org/10.1111/ele.13942

Grinevitch, L., Holroyd, S. L., & Barclay, R. M. R. (1995). Sex differences in the use of daily

torpor and foraging time by big brown bats (Eptesicus fuscus) during the reproductive

season. *Journal of Zoology, 235*(2), 301-309. https://doi.org/10.1111/j.1469-

7998.1995.tb05146.x

Grüebler, M. U., Morand, M., & Naef-Daenzer, B. (2008). A predictive model of the density of

airborne insects in agricultural environments. *Agriculture, Ecosystems & Environment,*

123(1), 75-80. https://doi.org/10.1016/j.agee.2007.05.001

Grüebler, M. U., Schuler, H., Müller, M., Spaar, R., Horch, P., & Naef-Daenzer, B. (2008).

Female biased mortality caused by anthropogenic nest loss contributes to population

decline and adult sex ratio of a meadow bird. *Biological Conservation, 141*(12), 3040-

3049. https://doi.org/10.1016/j.biocon.2008.09.008

Guerra-Silveira, F., & Abad-Franch, F. (2013). Sex Bias in Infectious Disease Epidemiology:

Patterns and Processes. *PLOS ONE, 8*(4). https://doi.org/10.1371/journal.pone.0062390

Hall, M. D., & Mideo, N. (2018). Linking sex differences to the evolution of infectious disease

life-histories. *Philosophical Transactions of the Royal Society B: Biological Sciences,*

373(1757). https://doi.org/10.1098/rstb.2017.0431

Herrera, J., & Nunn, C. L. (2019). Behavioural ecology and infectious disease: implications for

conservation of biodiversity. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, 374(1781), 20180054. https://doi.org/10.1098/rstb.2018.0054

Hoff, S., Pendergast, C., Johnson, L., Olson, E., O'Dell, D., Dowling, Z. R., Gorman, K. M.,

Herzog, C., & Turner, W. C. (2023). Seasonal roost characteristics and fall behavior of

coastal populations of Northern Myotis (Myotis septentrionalis). *Journal of Mammalogy*,

gyad102. https://doi.org/10.1093/jmammal/gyad102

Hoyt, J. R., Kilpatrick, A. M., & Langwig, K. E. (2021). Ecology and impacts of white-nose

syndrome on bats. *Nature Reviews Microbiology, 19*(3), 196-210.

https://doi.org/10.1038/s41579-020-00493-5

Hoyt, J. R., Langwig, K. E., Sun, K., Parise, K. L., Li, A., Wang, Y., Huang, X., Worledge, L.,

Miller, H., White, J. P., Kaarakka, H. M., Redell, J. A., Görföl, T., Boldogh, S. A., Fukui,

D., Sakuyama, M., Yachimori, S., Sato, A., Dalannast, M., Jargalsaikhan, A., Batbayar,

N., Yovel, Y., Amichai, E., Natradze, I., Frick, W. F., Foster, J. T., Feng, J., & Kilpatrick,

A. M. (2020). Environmental reservoir dynamics predict global infection patterns and

population impacts for the fungal disease white-nose syndrome. *Proceedings of the*

National Academy of Sciences, 201914794. https://doi.org/10.1073/pnas.1914794117

Hufkens, K., Basler, D., Milliman, T., Melaas, E. K., & Richardson, A. D. (2018). An integrated

phenology modelling framework in r. *Methods in Ecology and Evolution, 9*(5), 1276-

1285. https://doi.org/10.1111/2041-210X.12970

Jackson, R. T., Willcox, E. V., & Bernard, R. F. (2022). Winter torpor expression varies in four

bat species with differential susceptibility to white-nose syndrome. *Scientific Reports,*

12(1), 5688. https://doi.org/10.1038/s41598-022-09692-x

Jackson, R. T., Willcox, E. V., Zobel, J. M., & Bernard, R. F. (2022). Emergence activity at

hibernacula differs among four bat species affected by white-nose syndrome. *Ecology*

and Evolution, 12(7), e9113. https://doi.org/10.1002/ece3.9113

Johnson, J. S., Reeder, D. M., McMichael, J. W., Meierhofer, M. B., Stern, D. W. F., Lumadue,

S. S., Sigler, L. E., Winters, H. D., Vodzak, M. E., Kurta, A., Kath, J. A., & Field, K. A.

(2014). Host, pathogen, and environmental characteristics predict white-nose syndrome

mortality in captive little brown myotis (Myotis lucifugus). *PLOS ONE, 9*(11).

https://doi.org/10.1371/journal.pone.0112502

Jonasson, K. A., & Willis, C. K. R. (2011). Changes in Body Condition of Hibernating Bats

Support the Thrifty Female Hypothesis and Predict Consequences for Populations with

White-Nose Syndrome. *PLOS ONE, 6*(6). https://doi.org/10.1371/journal.pone.0021061

Kailing, M. J., Hoyt, J. R., White, J. P., Kaarakka, H. M., Redell, J. A., Leon, A. E., Rocke, T.

E., DePue, J. E., Scullon, W. H., Parise, K. L., Foster, J. T., Kilpatrick, A. M., &

Langwig, K. E. (2023). Sex-biased infections scale to population impacts for an emerging

wildlife disease. *Proceedings of the Royal Society B: Biological Sciences, 290*(1995),

20230040. https://doi.org/10.1098/rspb.2023.0040

Kailing, M. J., Hoyt, J. R., White, J. P., Redell, J. A., Kaarakka, H. M., & Langwig, K. E. (2025).

Data for: Activity patterns during the mating season predict sex-biased infections in an

emerging fungal disease. *Dryad Digital Repository.*

https://doi.org/10.5061/dryad.rjdfn2zq6

Kasumovic, M. M., Bruce, M. J., Andrade, M. C. B., & Herberstein, M. E. (2008). Spatial and

temporal demographic variation drives within-season fluctuations in sexual selection.

Evolution, 62(9), 2316-2325. https://doi.org/10.1111/j.1558-5646.2008.00446.x

Klein, S. L. (2000). The effects of hormones on sex differences in infection: From genes to

behavior. *Neuroscience & Biobehavioral Reviews, 24*(6), 627-638.

https://doi.org/10.1016/S0149-7634(00)00027-0

Kokko, H., Gunnarsson, T. G., Morrell, L. J., & Gill, J. A. (2006). Why do female migratory

birds arrive later than males? *Journal of Animal Ecology, 75*(6), 1293-1303.

https://doi.org/10.1111/j.1365-2656.2006.01151.x

Kolata, G. B. (1977). Sexual Dimorphism and Mating Systems: How Did They Evolve? *Science,*

195(4276), 382-383.

Kunz, T. H., Wrazen, J. A., & Burnett, C. D. (1998). Changes in body mass and fat reserves in

pre-hibernating little brown bats (Myotis lucifugus). *Ecoscience, 5*(1), 8-17.

https://doi.org/10.1080/11956860.1998.11682443

Kurta, A., & Kunz, T. H. (1987). Size of bats at birth and maternal investment during pregnancy.

Symposia of the Zoological Society of London, 57, 79-106.

Laggan, N. A., Parise, K. L., White, J. P., Kaarakka, H. M., Redell, J. A., DePue, J. E., Scullon,

W. H., Kath, J., Foster, J. T., Kilpatrick, A. M., Langwig, K. E., & Hoyt, J. R. (2023).

Host infection and disease-induced mortality modify species contributions to the

environmental reservoir. *Ecology, 104*(10), e4147. https://doi.org/10.1002/ecy.4147

Langwig, K. E., Frick, W. F., Bried, J. T., Hicks, A. C., Kunz, T. H., & Marm Kilpatrick, A.

(2012). Sociality, density-dependence and microclimates determine the persistence of

populations suffering from a novel fungal disease, white-nose syndrome. *Ecology Letters,*

15(9), 1050-1057. https://doi.org/10.1111/j.1461-0248.2012.01829.x

Langwig, K. E., Frick, W. F., Hoyt, J. R., Parise, K. L., Drees, K. P., Kunz, T. H., Foster, J. T., &

Kilpatrick, A. M. (2016). Drivers of variation in species impacts for a multi-host fungal

disease of bats. *Philosophical Transactions of the Royal Society B: Biological Sciences,*

371(1709). https://doi.org/10.1098/rstb.2015.0456

Langwig, K. E., Frick, W. F., Reynolds, R., Parise, K. L., Drees, K. P., Hoyt, J. R., Cheng, T. L.,

Kunz, T. H., Foster, J. T., & Marm Kilpatrick, A. (2015). Host and pathogen ecology

drive the seasonal dynamics of a fungal disease, white-nose syndrome. *Proceedings of*

the Royal Society B: Biological Sciences, 282(1799).

https://doi.org/10.1098/rspb.2014.2335

Langwig, K. E., Hoyt, J. R., Parise, K. L., Frick, W. F., Foster, J. T., & Kilpatrick, A. M. (2017).

Resistance in persisting bat populations after white-nose syndrome invasion.

Philosophical Transactions of the Royal Society B: Biological Sciences, 372(1712),

20160044. https://doi.org/10.1098/rstb.2016.0044

Langwig, K. E., Paul White, J., Parise, K. L., Kaarakka, H. M., Redell, J. A., DePue, J. E.,

Scullon, W. H., Foster, J. T., Marm Kilpatrick, A., Hoyt, J. R., & Kate Langwig, C. E.

(2021). Mobility and infectiousness in the spatial spread of an emerging fungal pathogen.

J Anim Ecol, 90, 1134-1141. https://doi.org/10.1111/1365-2656.13439

Legendre, S., Clobert, J., Møller, A. P., & Sorci, G. (1999). Demographic Stochasticity and

Social Mating System in the Process of Extinction of Small Populations: The Case of

Passerines Introduced to New Zealand. *The American Naturalist, 153*(5), 449-463.

https://doi.org/10.1086/303195

Lehtonen, J., Parker, G. A., & Schärer, L. (2016). Why anisogamy drives ancestral sex roles.

Evolution, 70(5), 1129-1135.

Lenth, R. V. (2024). emmeans: Estimated Marginal Means, aka Least-Squares Means (Version R

package version 1.10.6). Retrieved from https://CRAN.R-project.org/package=emmeans

Lüdecke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P., & Makowski, D. (2021). performance:

An R Package for Assessment, Comparison and Testing of Statistical Models. *Journal of*

Open Source Software, 6(60). https://doi.org/10.21105/joss.03139

Luong, L. T., Perkins, S. E., Grear, D. A., Rizzoli, A., & Hudson, P. J. (2010). The relative

importance of host characteristics and co-infection in generating variation in

Heligmosomoides polygyrus fecundity. *Parasitology, 137*(6), 1003-1012.

https://doi.org/10.1017/S0031182009991892

Martin, A. M., Hogg, J. T., Manlove, K. R., LaSharr, T. N., Shannon, J. M., McWhirter, D. E.,

Miyasaki, H., Monteith, K. L., & Cross, P. C. (2022). Disease and secondary sexual

traits: effects of pneumonia on horn size of bighorn sheep. *The Journal of Wildlife*

Management, 86(1), e22154. https://doi.org/10.1002/jwmg.22154

Mazor, T., Doropoulos, C., Schwarzmueller, F., Gladish, D. W., Kumaran, N., Merkel, K., Di

Marco, M., & Gagic, V. (2018). Global mismatch of policy and research on drivers of

biodiversity loss. *Nature Ecology & Evolution, 2*(7), 1071-1074.

https://doi.org/10.1038/s41559-018-0563-x

McDonald, J. L., Smith, G. C., McDonald, R. A., Delahay, R. J., & Hodgson, D. (2014).

Mortality trajectory analysis reveals the drivers of sex-specific epidemiology in natural

wildlife–disease interactions. *Proceedings of the Royal Society B: Biological Sciences,*

281(1790). https://doi.org/10.1098/rspb.2014.0526

McGuire, L. P., Turner, J. M., Warnecke, L., McGregor, G., Bollinger, T. K., Misra, V., Foster,

J. T., Frick, W. F., Kilpatrick, A. M., & Willis, C. K. R. (2016). White-nose syndrome

disease severity and a comparison of diagnostic methods. *EcoHealth, 13*(1), 60-71.

doi:10.1007/s10393-016-1107-y

Morbey, Y. E., & Ydenberg, R. C. (2001). Protandrous arrival timing to breeding areas: a

review. *Ecology Letters, 4*(6), 663-673. https://doi.org/10.1046/j.1461-

0248.2001.00265.x

Muller, L. K., Lorch, J. M., Lindner, D. L., O'Connor, M., Gargas, A., & Blehert, D. S. (2013).

Bat white-nose syndrome: a real-time TaqMan polymerase chain reaction test targeting

the intergenic spacer region of Geomyces destructans. *Mycologia, 105*(2), 253-259.

https://doi.org/10.3852/12-242

O'Shea, T., Cryan, P., Cunningham, A., Fooks, A., Hayman, D. T. S., Luis, A., Peel, A.,

Plowright, R., & Wood, J. L. N. (2014). Bat Flight and Zoonotic Viruses. *Emerging*

Infectious Disease journal, 20(5), 741. https://doi.org/10.3201/eid2005.130539

Oro, D., Álvarez, D., & Velando, A. (2018). Complex demographic heterogeneity from

anthropogenic impacts in a coastal marine predator. *Ecological Applications, 28*(3), 612-

621. https://doi.org/10.1002/eap.1679

Otto, S. P. (2018). Adaptation, speciation and extinction in the Anthropocene. *Proceedings of the*

Royal Society B: Biological Sciences, 285(1891), 20182047.

https://doi.org/10.1098/rspb.2018.2047

Perkins, S. E., Ferrari, M. F., & Hudson, P. J. (2008). The effects of social structure and sex-

biased transmission on macroparasite infection. *Parasitology, 135*(13), 1561-1569.

https://doi.org/10.1017/S0031182008000449

Péron, G. (2013). Compensation and additivity of anthropogenic mortality: life-history effects

and review of methods. *Journal of Animal Ecology, 82*(2), 408-417.

https://doi.org/10.1111/1365-2656.12014

Perry, R. W. (2013). A review of factors affecting cave climates for hibernating bats in temperate

North America. *Environmental Reviews 21: 28-39, 21*, 28-39.

https://doi.org/10.1139/er-2012-0042

Pfeiffer, B., & Mayer, F. (2013). Spermatogenesis, sperm storage and reproductive timing in

bats. *Journal of Zoology, 289*(2), 77-85. https://doi.org/10.1111/j.1469-

7998.2012.00970.x

Pikula, J., Amelon, S. K., Bandouchova, H., Bartonička, T., Berkova, H., Brichta, J., Hooper, S.,

Kokurewicz, T., Kolarik, M., Köllner, B., Kovacova, V., Linhart, P., Piacek, V., Turner,

G. G., Zukal, J., & Martínková, N. (2017). White-nose syndrome pathology grading in

Nearctic and Palearctic bats. *PLOS ONE, 12*(8), e0180435.

https://doi.org/10.1371/journal.pone.0180435

Plesnar-Bielak, A., Skrzynecka, A. M., Prokop, Z. M., & Radwan, J. (2012). Mating system

affects population performance and extinction risk under environmental challenge.

Proceedings of the Royal Society B: Biological Sciences, 279(1747), 4661-4667.

https://doi.org/10.1098/rspb.2012.1867

Promislow, D. E. (1997). Costs of sexual selection in natural populations of mammals.

Proceedings of the Royal Society of London. Series B: Biological Sciences, 247(1320),

203-210. https://doi.org/10.1098/rspb.1992.0030

Promislow, D. E., Montgomerie, R., & Martin, T. E. (1992). Mortality Costs of Sexual

Dimorphism in Birds. *Proceedings: Biological Sciences, 250*(1328), 143-150.

Racey, P., Uchida, T., Mori, T., Avery, M., & Fenton, M. (1987). Sperm-epithelium relationships

in relation to the time of insemination in little brown bats (Myotis lucifugus). *Journal of*

reproduction and fertility, 80(2), 445-454. https://doi.org/10.1530/jrf.0.0800445

Racey, P. A., & Entwistle, A. C. (2000). *9-Life-history and Reproductive Strategies of Bats*.

London: Academic Press.

Reeder, D. M., Frank, C. L., Turner, G. G., Meteyer, C. U., Kurta, A., Britzke, E. R., Vodzak, M.

E., Darling, S. R., Stihler, C. W., Hicks, A. C., Jacob, R., Grieneisen, L. E., Brownlee, S.

A., Muller, L. K., & Blehert, D. S. (2012). Frequent Arousal from Hibernation Linked to

Severity of Infection and Mortality in Bats with White-Nose Syndrome. *PLOS ONE,*
7(6), e38920. https://doi.org/10.1371/journal.pone.0038920

Rivers, N. M., Butlin, R. K., & Altringham, J. D. (2006). Autumn swarming behaviour of
Natterer's bats in the UK: Population size, catchment area and dispersal. *Biological
Conservation, 127*(2), 215-226. https://doi.org/10.1016/j.biocon.2005.08.010

Rosa, G. M., Bosch, J., Martel, A., Pasmans, F., Rebelo, R., Griffiths, R. A., & Garner, T. W. J.
(2019). Sex□biased disease dynamics increase extinction risk by impairing population
recovery. *Animal Conservation, 22*(6), 579-588. https://doi.org/10.1111/acv.12502

Rubalcaba, J. G., Gouveia, S. F., Villalobos, F., Cruz-Neto, A. P., Castro, M. G., Amado, T. F.,
Martinez, P. A., Navas, C. A., Dobrovolski, R., Diniz-Filho, J. A. F., & Olalla-Tárraga,
M. Á. (2022). Physical constraints on thermoregulation and flight drive morphological
evolution in bats. *Proceedings of the National Academy of Sciences, 119*(15),
e2103745119. https://doi.org/10.1073/pnas.2103745119

Sæther, B.-E., Engen, S., Lande, R., Møller, A. P., Bensch, S., Hasselquist, D., Beier, J., &
Leisler, B. (2004). Time to Extinction in Relation to Mating System and Type of Density
Regulation in Populations with Two Sexes. *Journal of Animal Ecology, 73*(5), 925-934.
https://doi.org/10.1111/j.0021-8790.2004.00869.x

Sauer, E. L., Connelly, C., Perrine, W., Love, A. C., & DuRant, S. E. (2024). Male pathology
regardless of behaviour drives transmission in an avian host–pathogen system. *Journal of
Animal Ecology, 93*(1), 36-44. https://doi.org/10.1111/1365-2656.14026

Schlaepfer, M. A., Runge, M. C., & Sherman, P. W. (2002). Ecological and evolutionary traps.
Trends in Ecology & Evolution, 17(10), 474-480. https://doi.org/10.1016/S0169-
5347(02)02580-6

Scialli, A. R. (1993). A comparative overview of mammalian fertilization: Dunbar BS, O'Rand

MG. New York: Plenum Press; 1991. In: Pergamon.

Shuster, S. M., & Wade, M. J. (2003). *Mating Systems and Strategies* (Vol. 61). Princeton

University Press. https://doi.org/10.2307/j.ctvs32sqk

Silk, M. J., Weber, N. L., Steward, L. C., Hodgson, D. J., Boots, M., Croft, D. P., Delahay, R. J.,

& McDonald, R. A. (2018). Contact networks structured by sex underpin sex-specific

epidemiology of infection. *Ecology Letters, 21*(2), 309-318.

https://doi.org/10.1111/ele.12898

Skorping, A., & Jensen, K. H. (2004). Disease dynamics: All caused by males? *Trends in

Ecology and Evolution, 19*(5), 219-220. https://doi.org/10.1016/j.tree.2004.02.006

Stephenson, J. F. (2019). Parasite-induced plasticity in host social behaviour depends on sex and

susceptibility. *Biology Letters, 15*(11). https://doi.org/10.1098/rsbl.2019.0557

Stephenson, J. F., Kinsella, C., Cable, J., & van Oosterhout, C. (2016). A further cost for the

sicker sex? Evidence for male-biased parasite-induced vulnerability to predation. *Ecology

and Evolution, 6*(8), 2506-2515. https://doi.org/10.1002/ece3.2049

Sweeny, A. R., Albery, G. F., Venkatesan, S. V., Fenton, A., & Pedersen, A. B. (2021).

Spatiotemporal variation in drivers of parasitism in a wild wood mouse population.

Functional Ecology, 1365-2435.13786. https://doi.org/10.1111/1365-2435.13786

Taylor, L. R. (1963). Analysis of the Effect of Temperature on Insects in Flight. *Journal of

Animal Ecology, 32*(1), 99-117. https://doi.org/10.2307/2520

Thomas, D. W., Brock Fenton, M., & Barclay, R. M. R. (1979). Social behavior of the little

brown bat, Myotis lucifugus. *Behavioral Ecology and Sociobiology, 6*(2), 129-136.

https://doi.org/10.1007/BF00292559

Tilman, D., Clark, M., Williams, D. R., Kimmel, K., Polasky, S., & Packer, C. (2017). Future

threats to biodiversity and pathways to their prevention. *Nature, 546*(7656), 73-81.

https://doi.org/10.1038/nature22900

Trivers, R. L. (1972). Parental Investment and Sexual Selection. In B. G. Campbell (Ed.), *Sexual*

selection and the descent of man: 1871-1971. (Vol. 1871-1971). Chicago, IL: Aldine.

van Schaik, J., Janssen, R., Bosch, T., Haarsma, A-J., Dekker, J. J. A., & Kranstauber, B. (2015).

Bats Swarm Where They Hibernate: Compositional Similarity between Autumn

Swarming and Winter Hibernation Assemblages at Five Underground Sites. *PLOS ONE,*

10(7), e0130850. doi:10.1371/journal.pone.0130850

Verant, M. L., Boyles, J. G., Waldrep, W., Wibbelt, G., & Blehert, D. S. (2012). Temperature-

Dependent Growth of Geomyces destructans, the Fungus That Causes Bat White-Nose

Syndrome. *PLOS ONE, 7*(9), 46280. https://doi.org/10.1371/journal.pone.0046280

Visser, M. E., & Gienapp, P. (2019). Evolutionary and demographic consequences of

phenological mismatches. *Nature Ecology & Evolution, 3*(6), 879-885.

https://doi.org/10.1038/s41559-019-0880-8

Wai-Ping, V., & Fenton, M. B. (1988). Nonselective Mating in Little Brown Bats (Myotis

lucifugus). *Journal of Mammalogy, 69*(3), 641-645. https://doi.org/10.2307/1381364

Warnecke, L., Turner, J.M., Bollinger, T.K., Lorch, J.M., Misra, V., Cryan, P.M., Wibbelt, G.,

Blehert, D.S., & Willis, C.K.R. (2012). Inoculation of bats with European Geomyces

destructans supports the novel pathogen hypothesis for the origin of white-nose

syndrome, *Proceedings of the National Academy of Sciences*, *109*(18), 6999-7003,

https://doi.org/10.1073/pnas.1200374109

Webb, P. I., Speakman, J. R., & Racey, P. A. (1996). How hot is a hibernaculum? A review of

the temperatures at which bats hibernate. *Canadian Journal of Zoology, 74*(4), 761-765.

https://doi.org/10.1139/z96-087

Wells, K., Hamede, R. K., Kerlin, D. H., Storfer, A., Hohenlohe, P. A., Jones, M. E., &

McCallum, H. I. (2017). Infection of the fittest: devil facial tumour disease has greatest

effect on individuals with highest reproductive output. *Ecology Letters*.

https://doi.org/10.1111/ele.12776

Williams, C. T., Chmura, H. E., Deal, C. K., & Wilsterman, K. (2022). Sex-Differences in

Phenology: A Tinbergian Perspective. *Integrative and Comparative Biology, 62*(4), 980-

997. https://doi.org/10.1093/icb/icac035

Willis, C. K. R. (2017). Trade-offs Influencing the Physiological Ecology of Hibernation in

Temperate-Zone Bats. *Integrative and Comparative Biology, 57*(6), 1214-1224.

https://doi.org/10.1093/icb/icx087

Wimsatt, W. A. (1942). Survival of spermatozoa in the female reproductive tract of the bat. *The*

Anatomical Record, 83(2), 299-307. https://doi.org/10.1002/ar.1090830208

Wimsatt, W. A. (1944). Further studies on the survival of spermatozoa in the female

reproductive tract of the bat. *The Anatomical Record, 88*(2), 193-204.

https://doi.org/10.1002/ar.1090880206

Wimsatt, W. A. (1945). Notes on Breeding Behavior, Pregnancy, and Parturition in Some

Vespertilionid Bats of the Eastern United States. *Journal of Mammalogy, 26*(1), 23.

https://doi.org/10.2307/1375029

Zuk, M. (2009). The Sicker Sex. *PLoS Pathogens, 5*(1), e1000267-e1000267.

https://doi.org/10.1371/journal.ppat.1000267

Zuk, M., & McKean, K. A. (1996). Sex Differences in Parasite Infections: Patterns and

Processes. *International Journal for Parasitology, 26*(10), 1009-1024.

https://doi.org/10.1016/S0020-7519(96)80001-4

<u>Supporting information</u>

Additional supporting information may be found in the online version of this article.

Methods, Tables, Figures S1 File detailing supplemental methods, tables, and figures

Appendix S1 File containing code and statistical output used throughout the manuscript