Nanoscale visualization of *Drosophila* E-cadherin ectodomain fragments and their interactions using DNA origami nanoblocks

Hiroki Oda[1,2,†], Shigetaka Nishiguchi[3,†,‡], Chihong Song[4,5,6,‡], Kazuyoshi Murata[4,5,6], Takayuki Uchihashi[3,7], Yuki Suzuki[8]

[1] Laboratory of Cell and Developmental Biology, JT Biohistory Research Hall, 1-1 Murasaki-cho, Takatsuki, Osaka 569-1125, Japan
[2] Department of Biological Sciences, Graduate School of Science, Osaka University, 1-1 Machikaneyama-cho, Toyonaka, Osaka 560-0043, Japan
[3] Exploratory Research Center on Life and Living Systems, National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan
[4] Exploratory Research Center on Life and Living Systems, National Institutes of Natural Sciences, 38 Nishigonaka, Myodaiji, Okazaki, Aichi 444-8585, Japan
[5] National Institute for Physiological Sciences, National Institutes of Natural Sciences, 38 Nishigonaka, Myodaiji, Okazaki, Aichi, 444-8585, Japan
[6] Department of Physiological Sciences, School of Life Science, The Graduate University for Advanced Studies (SOKENDAI), 38 Nishigonaka, Myodaiji, Okazaki, Aichi, 444-8585, Japan
[7] Department of Physics, Graduate School of Science, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan
[8] Department of Applied Chemistry, Graduate School of Engineering, Mie University, 1577 Kurimamachiya-cho, Tsu, Mie 514–8507, Japan

[†] These authors were equally contributed to this work.
Correspondence to: Hiroki Oda (hoda@brh.co.jp)

[‡] Present address (S.N.): Department of Biotechnology, Graduate School of Engineering, Osaka University, Yamadaoka 2-1, Suita, Osaka 565-0871, Japan
[‡] Present address (C.S.): Department of Convergence Medicine, School of Medicine, Pusan National University, Yangsan 50612, Republic of Korea

Keywords: cell-cell adhesion, cadherin, DNA origami, *Drosophila*, transmission electron microscopy, high-speed atomic force microscopy

Highlights:
• DNA origami technology was applied to perform a structure-function study of cadherin.

• DNA origami nanoblocks decorated with DE-cadherin ectodomains were observed by TEM/HS-AFM.

• A hinge-like site that separated the membrane-distal and -proximal portions of the DE-cadherin ectodomain was revealed.

• An adhesion-like process was mimicked via pairing two nanoblocks using DE-cadherin ectodomain interactions.

• Homophilic associations of DE-cadherin ectodomains between the nanoblocks were visualized at the nanoscale level.

Abstract

The adhesive function of cell surface proteins can be assessed visually; however, the underlying structures that mediate adhesion typically remain invisible at the nanoscale level. This hinders knowledge on the diversity of molecular architectures responsible for cell-cell adhesion. *Drosophila* E-cadherin (DE-cadherin), a classical cadherin with a unique domain structure, demonstrates adhesive function; however, it lacks a structural model that explains its adhesion mechanism. In this study, we present a novel application of DNA origami technology to create a cell-free, flat environment in which full DE-cadherin ectodomains are anchored using SNAP-tags and biotin-streptavidin interactions. DNA origami was assembled into a 120 nm long block, bearing 5 or 14 biotin:streptavidin sites that were evenly spaced on one lateral face. DE-cadherin ectodomain fragments were attached via biotinylated SNAP-tags. These decorated DNA origami nanoblocks were subjected to transmission electron and high-speed atomic force microscopy, which revealed a hinge-like site that separated the membrane-distal and -proximal portions of the DE-cadherin ectodomain, suggesting a role in mechanical flexibility. We also observed interactions between DE-cadherin ectodomains via their membrane-distal portions on single DNA origami nanoblocks. We reconstituted an adhesion-like process via pairing DNA origami nanoblocks using DE-cadherin ectodomain interactions. Homophilic associations of functional DE-cadherin ectodomains between the paired DNA origami nanoblocks were visualized at the nanoscale, displaying strand-like molecular configurations, likely representing the extracellular cadherin repeats without regular arrays of structural elements. This study introduces a DNA origami-based platform for reconstituting and visualizing cadherin ectodomain interactions, with potential applications for a broader range of adhesion molecules.

Introduction

The metazoan genome contains numerous genes that encode cell surface proteins that can mediate cell-cell adhesion (Hynes and Zhao, 2000; Putnam et al., 2007; Srivastava et al., 2010). A major category of cell-cell adhesion molecules is the cadherin superfamily, whose members are involved in various collective cell behaviors, including epithelial organization, pattern formation, morphogenesis, tissue homeostasis, and cancer development (Cavallaro and Christofori, 2004; Janiszewska et al., 2020). The cadherin superfamily is characterized by the presence of repeated extracellular cadherin domains (ECs), each consisting of approximately 110 amino acid residues with specific structural characteristics. However, its members represent considerable diversity in sizes and domain organizations (Nichols et al., 2012; Oda and Takeichi, 2011; Gul et al., 2017). The structural models proposed for adhesion vary among different cadherin types and subtypes (Gul et al., 2017). Ancestral and derived features in cadherin superfamily members have attracted attention for understanding metazoan evolution (Nichols et al., 2012; Oda, 2012).

Classical cadherins, a family found across metazoans but not in non-metazoans, are distinguished by their conserved cytoplasmic domains, which interact with the actin cytoskeleton via catenins and other proteins (Nichols et al., 2012; Oda and Takeichi, 2011; Clarke et al., 2016). These cadherins exhibit Ca^{2+}-dependent homophilic adhesion properties through *trans* (between proteins on opposing membranes)- and *cis* (between proteins on same membranes)-interactions with their ectodomains (Brasch et al., 2012)

and act as dynamic clusters at the cell-cell contact sites (Yap et al., 2015). Their crucial role in adherens junction assembly and morphogenesis are conserved across metazoans (Nathaniel Clarke et al., 2019). Despite their functional significance and evolutionary conservation, ectodomains of classical cadherins have diversified and acquired lineage-specific domain organizations through stepwise reductive transitions (Oda et al., 2005; Hulpiau and Van Roy, 2010; Sasaki et al., 2017). Typical classical cadherins in vertebrates possess a derived ectodomain comprising five ECs. Structural studies primarily focus on these 5-EC cadherins, among which the N-terminal-most ECs (EC1s) are major interfaces of *trans*-interactions, mediating adhesion with specificity (Boggon et al., 2002). This 5-EC configuration is conserved in desmosomal cadherins and restricted to vertebrates and urochordates (Gul et al., 2017). In contrast, epithelial classical cadherins in insects, such as *Drosophila* E-cadherin (DE-cadherin), exhibit an independently derived ectodomain with a distinct arrangement of seven ECs, a non-chordate cadherin domain (NC), a cysteine-rich EGF-like domain (CE), and a laminin-G domain (LG) (Oda and Tsukita, 1999; Oda et al., 2005; Sasaki et al., 2017). This cadherin type and other non-chordate classical cadherins lack a structural model that explains its adhesion mechanism. Domain-swapping experiments involving insect E-cadherin ectodomains, however, have identified membrane-distal four ECs, none of which correspond directly to vertebrate EC1 (Oda et al., 2005; Sasaki et al., 2017), as major determinants of homophilic adhesion specificities (Nishiguchi et al., 2016). In contrast to a rod-like configuration of the vertebrate classical cadherin ECs (Pokutta et al., 1994), these four ECs exhibit a folded configuration, as observed via high-speed atomic force microscopy (HS-AFM) (Nishiguchi et al., 2016; Nishiguchi and Oda, 2021). These studies suggest that the core mechanisms of adhesion at adherens junctions fundamentally differ between vertebrates and insects.

The adhesive function of cell surface proteins, such as classical cadherins, can be assessed using aggregation assays using cells that express the protein or microbeads coated with protein fragments (Chappuis-Flament et al., 2001; Nagafuchi et al., 1987). Protein-coated glass surfaces (Kovacs et al., 2002), functionalized cantilevers (Baumgartner et al., 2000), protein-anchored liposomes (Lambert et al., 2005), and microfluidic chambers (Eslami Amirabadi et al., 2019) have also been used to study adhesive interactions of cadherin ectodomains. Despite using these techniques and observing tangible functional outcomes, the underlying structures and processes that mediate adhesion remain unknown. Traditional techniques in structural biology, such as X-ray crystallography, cryo-electron microscopy, and HS-AFM, are powerful tools for analyzing cadherin structures and cadherin-cadherin interactions. These methods have been instrumental in modeling classical cadherins and other cadherin superfamily members (Boggon et al., 2002; Maker et al., 2022; Nishiguchi et al., 2022; Cooper et al., 2016). However, visualizing adhesion molecules at the nanoscale level in binding and unbinding states or processes remains challenging because of difficulties associated with reconstructing adhesion assemblies and preparing samples for microscopy. A promising approach involves using liposomes in conjunction with cryo-electron microscopy. Vertebrate classical cadherin ectodomains anchored to liposomes were crystallized to display highly ordered arrays of structural components between opposing liposome membranes, reminiscent of desmosomes (Lambert et al., 2005; Harrison et al., 2011). This method revealed the junction-like outcome; however, the dynamic states of adhesion processes remained inaccessible. Challenges, such as sample thickness and

uniformity issues, must also be addressed for wider applicability. The lipid bilayer environment is not essential for *in vitro* reconstruction of classical cadherin ectodomain adhesive interactions (Chappuis-Flament et al., 2001). Involving membrane-free scaffolds may facilitate cooperative interactions of cadherin molecules and help visualize the adhesive and dynamic nature of molecules at the nanoscale level, which could complement the techniques using liposomes.

In this study, we applied DNA origami technology to reconstitute and visualize DE-cadherin ectodomain-adhesive interactions. A key feature of DNA origami is the designability of nanostructures with functionalized sites (Dey et al., 2021). Utilizing this feature, we created a flat environment wherein cadherin ectodomain adhesive interactions occurred and were visualized at the nanoscale level. Transmission electron microscopy (TEM) and HS-AFM was used to visualize DNA origami nanostructures decorated with DE-cadherin ectodomains, which revealed a hinge-like site that separated the membrane-distal and -proximal portions of the DE-cadherin ectodomain, suggesting a role in mechanical flexibility. We mimicked cell-cell interface formation via pairing two DNA origami nanoblocks through DE-cadherin ectodomain interactions. The interfaces between the paired DNA origami nanoblocks showed strand-like molecular structures, likely representing the repeated ECs, although no regular arrays of structural elements were identified. The presented DNA origami−based platform allows the simultaneous realization of adhesion reconstitution and nanoscale visualization, which may have potential applications for structural and functional studies of a broader range of adhesion molecules.

Results
Preparing DE-cadherin ectodomain fragments that are biotinylated using a SNAP-tag
We prepared three constructs that each had a whole (denoted DEectW) or partial (denoted DEectD) DE-cadherin ectodomain (Fig. 1A). All of them had a V5/6×His-tag at the C-terminal end, and two of them had a SNAP-tag between the DE-cadherin region and the V5/6×His-tag (DEectW-S and DEectD-S; Supplementary Fig. 1). DEectW is identical to the previously described construct DEEXf (Nishiguchi and Oda, 2021). The deleted region in DEectD-S corresponds to EC1−EC4 (Fig. 1A; Supplementary Fig. 1), which was previously shown to be a major determinant of homophilic adhesion specificity (Nishiguchi et al., 2016). Using the *Drosophila* S2 cell culture system, we expressed these constructs, which were secreted in the culture media. Proteins were purified from the culture supernatants using cobalt NTA resin. The SNAP-tagged proteins were reacted with biotin-conjugated benzylguanine (BG-bio) (Fig. 1B), followed by removal of free BG-bio using dialysis. The resultant biotinylated DEectW-S and DEectD-S products were referred to as DEectW-Sbio and DEectD-Sbio, respectively.

The matured DE-cadherin ectodomain consists of two polypeptides, resulting from the removal of a prodomain and proteolytic cleavage between Gly^{1010} and Ser^{1011} in the NC, which are referred to as membrane-distal polypeptide (MD) and -proximal polypeptide (MP) (Oda and Tsukita, 1999). Reasonable sizes of the MD and MP for DEectW-Sbio and DEectD-Sbio were observed, as compared with those for DEectW (Fig. 1C, D), although no antibody was available for detecting the MD of DEectD-Sbio. Native PAGE mobility shift assay confirmed DEectW-Sbio and DEectD-Sbio could

whereas DEectW-S could not bind to streptavidin (SA) (Fig. 1D). No traces of non-shifted products were observed for DEectW-Sbio and DEectD-Sbio upon SA addition, indicating a complete reaction of the SNAP-tag with BG-bio.

Functional characterization of DEectW-Sbio using SA-conjugated microbeads
To test the capability of DEectW-Sbio in mediating homophilic adhesion, we established an aggregation assay system using SA-conjugated microbeads (hereafter referred to as SA beads) (Fig. 2A). DEectD-Sbio served as a negative control construct in functional assays. SA beads were incubated with 50 nM DEectW-Sbio or DEectD-Sbio for 5 min in HEPES-based Ca^{2+}- and Mg^{2+}-containing buffer (HCM buffer), followed by adding excess biotin (250 μM) to fill all open biotin-binding sites of the SAs. The beads were then shaken in 4-mm wells at 1,500 rpm for 5 min. After shaking, the microwell was left still for 10 min and the microwell bottom where the beads were sitting was examined by an inverted microscope to determine the degree of bead aggregation, which was defined to be the average size of the top 10% largest bead particles (pixel). Results showed that the degree of bead aggregation clearly differed between DEectW-Sbio and DEectD-Sbio, indicating that DEectW-Sbio could induce bead aggregates, but DEectD-Sbio could not (Fig. 1C, D). Two independent modifications, as additional negative controls, were applied to the assay procedure. First, the initial incubation step was performed with neither DEectW-Sbio nor DEectD-Sbio added. Second, SA bead incubation with DEectW-Sbio was preceded by the excess biotin addition step. Neither of these modifications resulted in bead aggregation (Fig. 1C, D). Overall, these results suggest that DEectW-Sbio molecules anchored to SA beads via the biotin-SA binding system mediate bead aggregation. Additional time-course analysis showed that the degree of bead aggregation mediated by DEectW-Sbio increased with time during the 5 min of shaking (Fig. 1E). The assays performed in the presence of 4 mM $CaCl_2$ or 1 mM EGTA indicated that the adhesion capability of DEectW-Sbio was dependent on Ca^{2+} ions (Fig 2F, G). Assays using serially lowered concentrations of DEectW-Sbio (50, 13, and 3 nM) showed that the adhesion activity of DEectW-Sbio was detectable at 50 and 13 nM but not at 3 nM (Fig. 2H).

To roughly estimate the density of DEectW-Sbio molecules on SA beads, we semi-quantified the number of DEectW-Sbio molecules bound to two aliquots of SA beads (an aliquot of SA beads is 5×10^5 beads, which was used in each microwell in the aggregation assays) in the same conditions as when using DEectW-Sbio at 50 nM and 13 nM in bead aggregation assays. This quantification showed that each aliquot of SA beads bound ~20 and ~6 fmol DEectW-Sbio molecules in the 50 and 13 nM conditions, respectively (Fig. 2I). Based on the assumption that all DEectW-Sbio molecules were uniformly distributed as monomers on the SA bead surface, the average distance between neighboring DEectW-Sbio molecules was estimated to be ~19 and 35 nm in the 50 and 13 nM conditions, respectively.

Constructing and visualizing DNA origami nanoblocks decorated with DEectW-Sbio molecules
To create a flat environment that was conducive to protein fragment anchoring and TEM/HS-AFM observation, we designed a block-shaped, three-dimensional DNA origami nanostructure (~120 nm × ~20 nm × ~10 nm) that had five biotinylated sites at intervals of ~21.5 nm on one of the long lateral faces (Fig. 3A). The DNA origami

nanoblock orientation was marked by lacking one of the corners. This DNA origami nanoblock bearing five biotin sites will be hereafter referred to as DO-5×bio. DO-5×bio was decorated with SAs, referred to as DO-5×bio:SA, and these decorated objects were purified using agarose gel electrophoresis (Fig. 3B; Supplementary Fig. 2). TEM observation revealed that the nanostructure of DO-5×bio:SA was formed as designed, with one of the largest faces adsorbed onto the support film (Fig. 3C).

We examined the binding of DEectW-Sbio to DO-5×bio:SA. DO-5×bio:SA (final concentration, 0.08 nM) was mixed with DEectW-Sbio at different concentrations (400, 40, 8, and 2 nM) and with 400 nM DEectW-S, followed by 10-min's incubation and agarose gel electrophoresis. The addition of DEectW-Sbio resulted in differentially shifted peaks of DO-5×bio:SA in a concentration-dependent manner; however, DEectW-S addition did not affect the DO-5×bio:SA migratory behavior (Fig. 3B). The peak profiles indicated the presence of at least four states in the binding between DO-5×bio:SA and DEectW-Sbio, which may reflect the multivalency of DO-5×bio:SA. Although DNA origami objects were prepared in Tris-based Folding buffer, which contained 1 mM EDTA (with no Ca^{2+} ions), we ensured that in the Folding buffer supplemented with 4 mM $CaCl_2$, DEectW-Sbio exhibited adhesive activity at the same level as that in the HCM buffer (Supplementary Fig. 3).

We used negative staining and TEM to observe DO-5×bio:SA nanoblocks decorated with DEectW-Sbio molecules. DO-5×bio:SA (final concentration, 0.08 nM) was mixed with DEectW-Sbio at an approximate molar ratio of 1:100 in the presence of 4mM $CaCl_2$ and then incubated for 2 min, followed by the negative staining procedure. TEM observations identified a total of 70 DO-5×bio:SA blocks in which one or more of the five bio:SA sites were recognizably occupied by DEectW-Sbio molecules (Fig. 3D-I), from three independent preparations. These decorated DNA origami nanoblocks in calcium-containing buffer will be referred to as DO-5×bio:SA+DEectW-Sbio:CA (CA denotes Ca^{2+} ions). Substantial variations were observed in the visualized shape and size of cadherin fragments. However, although each SA on the DNA origami block had theoretically three biotin-binding pockets vacant, only one of the three pockets, which was likely farthest from the DNA origami surface, appeared to have been used for the DEectW-Sbio molecule binding in most cases, possibly because of steric hindrance.

To grasp specific aspects of the structural properties of DEectW-Sbio molecules with Ca^{2+} ions, we examined DEectW-Sbio and DEectD-Sbio molecules in the absence and presence of Ca^{2+} ions, respectively. A total of 49 DNA origami nanoblocks decorated with DEectW-Sbio in EDTA-containing buffer without Ca^{2+} (Fig. 3J, K), referred to as DO-5×bio:SA+DEectW-Sbio:ED (ED denotes EDTA), and of a total of 37 DNA origami nanoblocks for DEectD-Sbio with Ca^{2+} (Fig. 3L, M), referred to as DO-5×bio:SA+DEectD-Sbio:CA, were identified from two independent preparations for each. Comparisons of visualized cadherin fragments bound to bio:SA sites of the DNA origami nanoblocks confirmed that specifically structured DE-cadherin ectodomain monomers, featured by folded configurations of the membrane-distal multiple ECs, existed in the sample type DO-5×bio:SA+DEectW-Sbio:CA (Fig. 3D, D', E, E'), consistent with our previous observations of DE-cadherin ectodomains using HS-AFM (Nishiguchi et al., 2016; Nishiguchi and Oda, 2021). However, we found exceptions: the membrane-distal ECs were in an unfolded or opened state (Fig. 3F, F'). We also found a novel structural feature of the DE-cadherin ectodomain—the presence of a specific hinge-like site that separated the membrane-distal and -proximal portion

(Fig. 3D'-F'). Flexibility of the hinge angle appeared to allow various bent configurations of the DE-cadherin ectodomain. To understand the flexibility further, we performed two-dimensional (2D) class averaging of the negatively stained TEM images of DEectW molecules, which revealed various shapes of DE-cadherin ectodomain, consistent with the folded, unfolded, and bent configurations of DEectW-Sbio molecules (Supplementary Fig. 4). The negative-staining TEM images (Fig. 3D–M) and 2D class averaging of DEectW molecules (Fig. 3N) allowed us to propose a structural arrangement for DEectW on SNAP, SA, and DO-5×bio (Fig. 3O), revealing its spatial configuration and potential functional interactions.

Additionally, some molecular images from the sample type DO-5×bio:SA+DEectW-Sbio:CA displayed potential dimers, exhibiting symmetric shapes (Fig. 3G–I, G'–I'). At least one of them indicated an interaction or close proximity of the membrane-distal portions of two DEectW-Sbio molecules rooted at neighboring bio:SA sites (Fig. 3I, I').

HS-AFM revealed the dynamic nature of the DE-cadherin ectodomain
To examine the structures and dynamics of the DE-cadherin ectodomain on DNA origami scaffold in solution, we visualized DO-5×bio:SA nanostructures decorated with DEectW-Sbio molecules using HS-AFM. DO-5×bio:SA nanoblocks, DEectW-Sbio molecules, and the combined were imaged (Fig. 4A–C), showing consistencies with the TEM observations (Fig. 3). HS-AFM showed that the membrane-distal and -proximal polarity of the DE-cadherin ectodomain could be recognized (Fig. 4B) as the membrane-proximal portion of the DE-cadherin ectodomain was larger in size than that of the folded membrane-distal ECs (Nishiguchi and Oda, 2021). However, when the DEectW-Sbio molecule was anchored to the bio:SA site of the DNA origami nanoblock, the membrane-proximal portion of the DE-cadherin ectodomain was not separately visualized from the bio:SA portion, owing to limited resolution of HS-AFM (Fig. 4C). HS-AFM still revealed the dynamics of the membrane-distal portion of DEectW-Sbio molecules. Extending and shrinking dynamics of the DEectW-Sbio molecule were visualized at a time resolution of 0.3 s/frame, showing flexibility mainly derived from the membrane-distal portion of the DEectW-Sbio (Fig. 4D and Movie S1). HS-AFM also showed binding and unbinding dynamics of two DEectW-Sbio molecules via their membrane-distal portions on same DNA origami scaffold (Fig. 4E and Movie S2). A dimer-like structure formed between a DEectW-Sbio molecule on the DNA origami scaffold and another DEectW-Sbio molecule free from the scaffold were visualized, showing flexibility of the membrane-distal portions of the DEectW-Sbio molecules while retaining binding between protomers (Fig. 4F and Movie S3).

Induction and visualization of DE-cadherin ectodomain adhesive interactions between the DNA origami nanoblocks
DE-cadherin ectodomain-mediated adhesion was suggested to depend on the molecule density (Fig. 2H, I). To facilitate interactions between DE-cadherin ectodomains on different DNA origami nanoblocks, we modified the DO-5×bio design by replacing nine more staple oligonucleotides with biotinylated ones to produce a DO-14×bio nanoblock, which had two sets of seven biotin sites in line at intervals of ~7.2 nm on one of the long lateral faces (Fig. 5A). DO-14×bio nanoblocks decorated with SAs (DO-14×bio:SA) were examined using negative staining and TEM, confirming the

binding of SAs at many, if not all, biotin sites (Fig. 5A). Agarose gel electrophoresis showed that DO-14×bio:SA could form a larger complex with DEectW-Sbio molecules than with DO-5×bio:SA (Fig. 5B).

A mixture of DO-14×bio:SA blocks and DEectW-Sbio molecules was incubated in the presence of Ca^{2+} ions and then shaken, followed by negative staining and TEM (Fig. 5C–F). This sample type will be referred to as DO-14×bio:SA+DEectW-Sbio:CA. TEM observations showed that a small proportion of DO-14×bio:SA nanoblocks were found as pairs in which the facing nanoblocks were bridged via DEectW-Sbio molecule interactions (Fig. 5E, F; see data in the repository); most DO-14×bio:SA blocks remained single, with the bio:SA sites bearing DEectW-Sbio molecules (Fig. 5D; see data in the repository). A total of 47 DO-14×bio:SA+DEectW-Sbio:CA block pairs were identified and imaged from two independent preparations (repository). For comparison, we additionally tested the same combination in the absence of Ca^{2+} ions (DO-14×bio:SA+DEectW-Sbio:ED; Fig. 5G–I), and a mixture of DO-14×bio:SA blocks and DEectD-Sbio molecules in the presence of Ca^{2+} ions (DO-14×bio:SA+DEectD-Sbio:CA; Fig. 5J–L). In both sample types, most DO-14×bio:SA nanoblocks were observed as singles, with the bio:SA sites bearing cadherin fragments (Fig. 5G, J). Even in these control sample types, however, paired DO-14×bio:SA blocks that were bridged through interactions of cadherin fragments were found, with lower frequencies than in the case of DO-14×bio:SA+DEectW-Sbio:CA. A total of 22 DO-14×bio:SA+DEectW-Sbio:ED block pairs and 23 DO-14×bio:SA+DEectD-Sbio:CA block pairs were identified and imaged from two independent preparations (ref, repository).

To compare the dimension of space occupied by the interacting cadherin ectodomain fragments between the three sample types, we measured the distance (D_{b-b}) and angle between the paired DO-14×bio:SA blocks (Fig. 5M), which showed that 75% or more of the DNA origami block pairs for each sample type were at a 20° or smaller angle (Fig. 5N). The median D_{b-b} values for the DO-14×bio:SA+DEectW-Sbio:CA block pairs was 32.5 nm, in contrast to 19.7 and 19.5 nm for the DO-14×bio:SA+DEectW-Sbio:ED and DO-14×bio:SA+DEectD-Sbio:CA block pairs, respectively (Fig. 5N, O). The DO-14×bio:SA+DEectW-Sbio:CA block pairs had larger D_{b-b} values than the DO-14×bio:SA+DEectW-Sbio:ED and DO-14×bio:SA+DEectD-Sbio:CA block pairs (Fig. 5O), suggesting that homophilic associations of functional DE-cadherin ectodomains were induced and visualized between the facing DNA origami blocks.

Moreover, we examined the morphological aspects of DE-cadherin ectodomain fragments between the facing origami blocks. Strand-like molecular configurations that likely reflected the ECs of DE-cadherin were observed in some of the DO-14×bio:SA+DEectW-Sbio:CA block pairs (Fig. 5F'). Thick strands of protein materials were observed in some of the DO-14×bio:SA+DEectW-Sbio:CA block pairs with the largest D_{b-b} values (Supple fig. 4), contrasting with the other two sample types (Fig. 5I', L'; Supple fig. 4). However, as the recognized molecular shapes and arrangements varied, interpreting the individual interactions of DE-cadherin ectodomains was difficult.

Discussion

We applied DNA origami technology to perform a structure-function study of the DE-cadherin ectodomain. The SNAP-tag-biotinylated membrane-proximal side of

the DE-cadherin ectodomain showed that it recapitulated Ca^{2+}-dependent adhesive properties in a cell and membrane-free environment using SA microbeads, despite the tagging and biotinylation. The strong biotin:SA interactions non-covalently immobilized cadherin fragments to the DNA origami nanoblock at a low concentration, making it compatible with subsequent TEM/AFM observations of molecules without purification. In this application, the tetrameric SA organization may have complicated the anchoring interactions of biotinylated cadherin fragments to the DNA origami object. However, this did not occur based on the peak shift assay of decorated DNA origami (Fig. 3B) and TEM/AFM observations of anchored molecules (Fig. 3D–I; Fig. 4C), which suggested that each bio:SA site on the origami block appeared to serve as a single link to the biotinylated cadherin fragment, presumably because of steric hindrance. The linkage angle was vertical (or near-vertical) to the DNA origami nanoblock surface. These situations facilitate the interpretation of visualized molecule structures and polarities, even when their total shapes vary or change dynamically.

The oriented immobilization of cadherin fragments to the DNA origami nanoblock provides a uniform environment in which adhesion molecules function and are visualized. The designable DNA origami allows immobilized molecules to be constantly placed to be viewed from the lateral side, and their spacing can be easily controlled. The estimated density of DE-cadherin ectodomain fragments in mediating bead aggregation (~1000–3500 molecules per μm^2; Fig. 2I) was larger, yet roughly comparable to that of DE-cadherin molecules at the adherens junction in epithelial cells of the stage-9 *Drosophila* embryo (~300 molecules per μm^2) (Truong Quang et al., 2013). The designed spacing of cadherin fragments on the DNA origami nanoblock was comparable to these functional cadherin densities. Although an apparent demerit of using DNA origami is lack of fluidity in its architecture, this extreme circumstance enables exploring the dynamic nature of cadherin ectodomains, on their own, without considering dynamics of scaffolding materials.

Deploying the DNA origami-based platform has increased our understanding of the structural properties of DE-cadherin ectodomains. A previous study used HS-AFM, in which cadherin fragments that were directly adsorbed on mica substrates were observed; the study suggested that the membrane-distal four ECs of DE-cadherin were folded to form a terminal knot (Nishiguchi et al., 2016). In this study, this structural feature was reproduced not only by HS-AFM but also by TEM. However, as we found some exceptions, the structure of the N-terminal ECs could have multiple states, such as closed and open states (Fig. 3F'). This study also discovered that the DE-cadherin ectodomain had a hinge-like linker site between the membrane-proximal and -distal portions. TEM observations captured various hinge angles of this linker site, whereas HS-AFM observations captured extending and shrinking motions of the DE-cadherin ectodomain around the linker site. Overall, these observations suggest that the hinge-like linker site is flexible and facilitates mechanical motions in the DE-cadherin ectodomain, at least in a monomeric state. The precise amino acid residues responsible for the flexible linker site could not be determined because of limitations in resolution of our TEM/HS-AFM measurements.

We observed dimer-like DE-cadherin ectodomains on single DNA origami nanoblocks. Dimers of DE-cadherin ectodomains have not been observed in previous studies using HS-AFM; hence, we considered that the interactions between individual DE-cadherin fragments could be weak (Nishiguchi et al., 2016; Nishiguchi and Oda,

2021). Immobilization of cadherin fragments to the DNA origami nanoblock may affect the thermodynamic stability of interactions, increasing chances for dimer observation. The observed mirror-symmetric shapes of some dimer-like molecules suggested that the DE-cadherin ectodomain dimerization appeared to be mediated by their membrane-distal portions, which were demonstrated to be determinants of homophilic binding specificity (Nishiguchi et al., 2016). HS-AFM directly showed transient and stable interactions between DE-cadherin ectodomains, likely via their membrane-distal portions. Considering the mechanical flexibility of the hinge-like linker site and various motions of the paired molecules, however, we could not determine the mode (*cis-* or *trans-*) of the visualized interaction between the dimer-like molecules.

We reconstituted an adhesion-like process by pairing DNA origami nanoblocks via DE-cadherin ectodomain interactions. Induced adhesion between DNA origami nanoblocks yielded thin, flat architectures that were suitable for TEM sample preparation, providing opportunities to observe homophilic associations of functional DE-cadherin ectodomains at the nanoscale level. The space between the facing DNA origami nanoblocks included two bio：SA layers, which were each ~5 nm in width (Fan et al., 2019). Considering the distribution of the measured D_{b-b} values (Fig. 5O), the net space available for homophilic associations of DEectW-Sbio molecules between the DNA origami blocks was ~16–27 nm in width. This space size is comparable to that between the facing cell membranes at the adherens junctions in which adhesion is mediated by DE-cadherin in epithelial cells of *Drosophila* embryos and tissues (Tepass and Hartenstein, 1994). In the presented reconstitution strategy, however, even non-anchored DEectW-Sbio molecules can participate in DE-cadherin ectodomain assemblies forming between the DNA origami nanoblocks. This was particularly probable in DNA origami nanoblock pairs with large D_{b-b} values (Supplementary Fig. 5A). Based on this situation and strand-like configurations of the likely ECs with no regular arrays of structural elements, DE-cadherin ectodomains may relatively loosely interact with one another to act as protein condensates.

Unlike cadherin molecules sitting in the lipid bilayer, DEectW-Sbio ectodomains anchored onto the DNA origami nanoblock or magnetic bead may be prevented from behaving flexibly toward minimizing their interaction energies. This restrictive circumstance could have not allowed structurally organized assemblies of DE-cadherin ectodomains to be observed, in contrast to junction-like assemblies of vertebrate cadherin ectodomains formed between liposomes in a previous study (Lambert et al., 2005; Harrison et al., 2011). If the DNA origami nanoblocks are allowed to flexibly slide via setting them on a lipid bilayer (Suzuki et al., 2015), we could observe dynamic processes of adhesion formation and stabilization of DE-cadherin ectodomains using HS-AFM. If the movement of the DNA origami nanoblocks is manipulated by applied external forces, such as using magnetic devices and optical tweezers (Marjoram et al., 2016; Bartsch et al., 2019), we could observe the detaching and/or force-responsive processes of DE-cadherin adhesions. These challenges should be addressed in the future.

DE-cadherin ectodomains anchored onto the DNA origami nanoblock were in a linear spatial arrangement, unlike the microbeads. This circumstance may limit the functional performance of the DE-cadherin ectodomains. DNA origami nanostructures are designable; therefore, the structural design could be improved toward optimizing the performance of the anchored DE-cadherin fragments, which should be based on

elucidating the structural mechanisms of DE-cadherin adhesion. DE-cadherin, similar to vertebrate classical cadherins, has been suggested to act as a nanocluster (Cavey et al., 2008; Strale et al., 2015; Truong Quang et al., 2013; Yap et al., 2015). DE-cadherin ectodomains could have multiple binding interfaces or stepwise regulations involved in forming adhesion assemblies. Further development of the DNA origami-based platform could contribute to visualizing such DE-cadherin ectodomain interactions and adhesion processes.

In conclusion, we applied DNA origami technology for the nanoscale visualization of DE-cadherin ectodomains and their interactions. This technology is designable and can be easily handled; therefore, it has the potential to be applied to various adhesion molecules. For example, the structural, functional, and mechanical aspects of the molecules could be better understood. Homophilic binding and selective sorting properties of adhesion molecules (Nishiguchi et al., 2016) could be beneficial for artificial cell engineering and molecular robotics, which aim to integrate molecular devices into an autonomous system.

Methods
DNA construction
To construct the plasmid for expressing cadherin fragments in *Drosophila* S2 cells, we utilized the pAc5.1 V5-His A vector (Thermo Fisher Scientific, MA, USA). The plasmid was designed for DEectW and named as pAcHis-DEectW. It was identical to the previously described pAcHis-DEEXf (Nishiguchi and Oda, 2021). Integrating a DNA fragment that encoded the SNAP-tag (New England Biolabs, MA, USA) into the *Xba*I site of pAcHis-DEectW yielded the newly generated construct, pAcHis-DEectW-S. Two distinct regions of DE-cadherin cDNA, corresponding to amino acid (aa) 1–70 and aa 523–981, were amplified via polymerase chain reaction and seamlessly fused using In-Fusion Cloning technology (Takara Bio, Shiga, Japan) to produce pAcHis-DEectD. The HindIII-BamHI region of pAcHis-DEectD was replaced with that of pAcHis-DEectW-S to produce pAcHis-DEectD-S. Primers used for the plasmid construction are listed in Supplementary Table 1.

Preparation of cadherin fragments
Drosophila S2 cells, originally obtained from the *Drosophila* community over 2 decades ago (Schneider, 1972), were cultured at 25°C in Schneider's medium (Thermo Fisher Scientific) and supplemented with 10% heat-inactivated fetal bovine serum (Biowest, Nuaille, France). In comparison to commercially available S2 cells (R69007; Thermo Fisher Scientific), these S2 cells exhibited enhanced cadherin fragment production when transfected with pAcHis-DEEXf-G (Nishiguchi and Oda, 2021). For a standard scale of cadherin fragment expression and purification, S2 cells were seeded at a density of 1.5×10^6 cells/ml in two 90-mm dishes (10 ml medium/dish) approximately 5 h before transfection. In each dish, S2 cells were transfected with 10 µg of plasmid DNA using *Trans*IT-Insect Transfection Reagent (MIR 6100: Mirus Bio, WI, USA). Following 3.5 d of incubation at 25°C, conditioned media were collected from the two dishes and centrifuged at 1,690 g for 15 min at 4°C, resulting in approximately 20 mL of supernatant. This supernatant was concentrated to 1 mL or less using Amicon Ultra-15 Centrifugal Filter, 100 kDa MWCO (UFC910008; Merck KGaA, Darmstadt, Germany). The concentrated supernatant was then diluted by adding

Equilibration/Wash buffer (50 mM sodium phosphate pH 7.4, 300 mM NaCl, 10 mM imidazole) to a volume of 10 mL. The resulting solution was applied to 0.5 mL of TALON Metal Affinity Resin (#635501; Takara Bio), washed with Equilibration/Wash buffer, and eluted with Elution buffer (50 mM sodium phosphate pH 7.4, 300 mM NaCl, 150 mM imidazole). The protein concentration of each eluted fraction was measured and peak fractions were combined, followed by concentration to 40 µL or less using vivaspin 500-30K (Cytiva, Uppsala, Sweden). Typical yields of the proteins were 50–100 µg. For biotinylation, 5 µM purified SNAP-tagged proteins were reacted with 10 µM biotin-conjugated benzylguanine (BG-bio) (S9110S; New England Biolabs) for 30 min at 37°C. The reacted solution was subsequently dialyzed over 3 d using the *Xpress* Micro Dialyzer (MD100, MWCO 12–15 kDa; Scienova, Jena, Germany) to remove free biotin and exchange the buffer (twice with 40 ml of HE buffer [20 mM HEPES pH 7.35, 1 mM EDTA] and four times with 40 ml of HCM buffer [20 mM HEPES pH 7.35, 4 mM $CaCl_2$, 10 mM $MgCl_2$]). DEectW-Sbio used for docking on DNA origami in the absence of Ca^{2+} was subjected to dialysis using HE buffer over six exchanges. The prepared cadherin fragments were stored at 4°C until use.

Protein analyses
The size, quantity, and quality of purified proteins were assessed using sodium dodecyl-sulfate polyacrylamide gel electrophoresis (SDS-PAGE, 7.5 or 6% gels) and Coomassie Brilliant Blue staining. For western blotting, the following antibodies were used: rat ant-DE-cadherin antibody (DCAD2 [1:100]; (Oda et al., 1994)), mouse anti-V5 tag antibody (#46-0705 [1:1000]; Thermo Fisher Scientific), and horseradish peroxidase-conjugated anti-rat (NA935V [1:1000]) and anti-mouse (NA931V [1:1000]) IgG antibodies (GE Healthcare, IL, USA). Signals were detected using WesternSure Premium Chemiluminescent Substrate (#926-95010) and the C-DiGit Blot Scanner (LI-COR; NE, USA). Native PAGE, conducted without SDS and a reducing agent, was performed using a 6% gel. Protein quantity was determined using the Qubit Protein Assay Kit and Qubit 2.0 fluorometer (Thermo Fisher Scientific).

Bead aggregation assays
Magnetic microbeads coated with streptavidin (J-MS-S160S; Medical and Biological Laboratories, Tokyo, Japan) were prepared as a suspension in HCM-BSA buffer (HCM buffer containing 0.01% bovine serum albumin [BSA; A-2143; Sigma Aldrich, MO, USA]) at a density of approximately 5×10^5 beads/µL. For experiments involving Ca^{2+}, 1 µL of the SA beads suspension was mixed with 20 µL HCM buffer, followed by a 5-min incubation with either DEectW-Sbio or DEectD-Sbio at appropriate concentrations (50, 13, or 3 nM), after which it was incubated with excess biotin (250 µM; 12987-14; Nacalai Tesque, Kyoto, Japan) for 1 min. Following the SA-masking step, the bead suspension was transferred to a 4-mm well in µ-Slide 15 Well 3D Glass Bottom (#81507; ibidi, Gräfelfing, Germany), which had been pretreated with HCM-BSA buffer and shaken at 1,500 rpm (mixing radius = 1.5 mm) using the Eppendorf ThermoMixer C (Eppendorf, Hamburg, Germany) for 5 min at 25°C. In Ca^{2+}-free conditions, HEM-BSA buffer (20 mM HEPES pH 7.35, 1 mM EGTA, 10 mM $MgCl_2$, 0.01% BSA) was used throughout the procedure, except for the initial preparation of the DEectW-Sbio stock in HCM buffer for bead aggregation assays. In additional bead aggregation assays, a Tris-based folding buffer (for preparing DNA origami) was used

instead of the HEPES-based buffers. After shaking, the multi-well slide was left undisturbed for 10 min. It was then carefully transferred to an inverted microscope (IX71, Evident, Tokyo, Japan), equipped with a cooled CCD camera (CoolSNAP HQ; Roper Scientific, AZ, USA). Three different regions of the well bottom were photographed using a 20× objective lens for each well. The obtained images were then systematically cropped into squares of 800×800 pixels, equivalent to a dimension of 256×256 µm. These images were analyzed using the 'Analyze Particles' function in Image J 1.53t software (National Institutes of Health, MD, USA; Schindelin et al., 2012). During this analysis, areas of individual bead aggregates (pixels) were measured, with a cut-off value of 10 pixels. Measured bead areas were sorted by size to determine the average top 10% bead areas, which were a measure of the aggregation degree (pixel). Each sample/condition type was typically subjected to four independent repetitions of the experiment, utilizing cadherin fragments from at least two separate purifications.

To quantify the DEectW-Sbio protein quantity bound to SA microbeads under the same conditions as those used in the aggregation assays, the beads were isolated using centrifugation immediately after incubating the mixture for 5 min. The molecules bound to the beads were then separated via boiling in SDS sample buffer.

Preparing DNA origami structures

DNA origami nanostructures were designed using the caDNAno2 software (Douglas et al., 2009)(Supplementary Fig 6; Supplementary Tables 2 and 3). A total of 239 oligonucleotide species were synthesized (Eurofins Genomics Inc., Tokyo, Japan) and used as staple DNAs for preparing rectangle-shaped origami. To generate the DNA origami-5×bio and -14×bio (DO-5×bio and -14×bio), 5 and 14 specific species of oligonucleotides, respectively, were replaced with corresponding oligonucleotides modified by adding TT-Biotin-TEG when preparing the staple DNA mixtures. The origami unit was assembled in a 20-µL solution containing 10 nM single-stranded, circular scaffold DNA p8064 (tilibit nanosystems, Munich, Germany), 32 nM staple DNAs, 5 mM Tris buffer (pH 8.0), 1 mM EDTA, and 15 mM $MgCl_2$. The mixture of scaffold and staple DNAs was annealed by reducing the temperature from 85 to 65°C for over 1 h, followed by a reduction from 65°C to 44°C at a rate of 1°C/h, and finally to 25 °C. After annealing was completed, the origami products were precipitated by adding an equal volume of 15% PEG6000, 5 mM Tris pH 8.0, 1 mM EDTA, and 505 mM NaCl (Stahl et al., 2014). The precipitate was then resolved in 20 µL of Folding buffer (5 mM Tris pH 8.0, 1 mM EDTA, and 15 mM $MgCl_2$). For decoration, a 20-µL solution containing biotinylated DNA origami was mixed with 2 µL of 500 mM SA (32243-11; Nacalai Tesque) and incubated for 20 min at room temperature. The decorated DNA origami products were purified using agarose gel electrophoresis and Freeze N' Squeeze DNA Gel Extraction Spin Columns (#7326165; Bio-Rad Laboratories, Hercules, CA, USA), immediately followed by dialysis with Folding buffer. Electrophoresis of the DNA origami was performed using 0.7 or 1% agarose gel (#161-3106; Bio-Rad Laboratories, CA, USA) and 0.5× Tris/Borate/EDTA buffer containing 5 mM $MgCl_2$. To stain the DNA materials in agarose gels, SYBR Gold (S11494; Thermo Fisher Scientific) was used. The DNA origami quantity (8,052 bp) was determined based on the staining intensity relative to the 2,322 bp and 2,027 bp fragments of the lambda DNA/HindIII marker on the agarose gel.

Docking of cadherin fragments onto DNA origami nanostructures
DEectW-Sbio and DEectD-Sbio, maintained in HCM buffer, were diluted to a
concentration of 40 fmol/µL in Folding buffer containing 20 mM $CaCl_2$. To prepare
DNA origami with cadherin fragments anchored, 4 µL of 1.0 fmol/µl DO-5×bio:SA
was combined with 1 µL of the diluted DEectW-Sbio or DEectD-Sbio solution via
pipetting in a 1.5 mL Eppendorf tube (#022431081). The mixture was then incubated
for 2 min at room temperature, immediately followed by negative staining for TEM
observation. To pair DNA origami nanoblocks via cadherin fragments, 4 µL of 0.75–1.0
fmol/µL DO-14×bio:SA was mixed with 1 µL of 40 fmol/µl DEectW-Sbio or DEectD-
Sbio via pipetting in a 1.5 mL Eppendorf tube (#022431081) and incubated for 5 h on
ice. The mixture was then shaken at 1,500 rpm on the Eppendorf ThermoMixer C
(Eppendorf, Hamburg, Germany) for 5 min at 25°C, immediately followed by negative
staining for TEM observation. To establish the Ca^{2+}-free condition for DEectW-Sbio
docking onto DNA origami, DEectW-Sbio that was initially prepared in HE buffer was
diluted in Folding buffer, which contained 1 mM EDTA.

Negative-staining electron microscopy
To observe mixtures of DNA origami and cadherin fragments, the sample was absorbed
onto a glow-discharged carbon support film and stained with 2% uranyl acetate. The
stained samples were analyzed using an electron microscope (Hitachi H-7600, JOEL
JEM-2010). TEM observations for each sample/condition type were conducted using at
least two independent preparations of DNA origami objects and cadherin fragments.

For 2D class averaging of DEectW molecules, the sample (2.5 µL) was applied
onto a carbon-coated copper grid that was glow-discharged beforehand. After removing
the excess sample solution with a filter paper, specimens were stained with a 2% uranyl
acetate solution for 30 s. The grids were dried in air after removing the staining
solution with a filter paper. EM images were recorded on a DE-20 camera (Direct
electron LP) at a nominal magnification of 40,000 using a JEM2200FS electron
microscope (JEOL Ltd.) operated at 200 kV accelerating voltage. The energy slit
width was adjusted at 20 eV. The image pixel size was 1.4 Å on the camera. A 2–
3 µm under-focused condition was selected to enhance the contrast of EM images.
Image analysis was performed with Relion 2.0 (Scheres, 2012) motion correcting the
images using the DE_process_frames.py script provided by the manufacturer. A total of
1,655 particles of DEectW were automatically picked from the images after individually
correcting the contrast transfer function. The particles were classified at 2D in Relion
2.0.

HS-AFM
A mica substrate with 1.5 mm diameter and 0.1 mm thickness (Furuuchi Chemical,
Tokyo, Japan) was fixed using glue on a glass stage. A 2 µL 0.04% 3-
aminopropyltriethoxysilane (APTES) solution was added on a freshly cleaved mica
substrate and incubated for 3 min. The APTES-coated mica substrate was washed twice
with 80 µL Milli-Q water. A 2 µL DO-5×bio:SA solution (1.0 fmol/µL) was added on
the APTES-coated mica substrate for 3 min. A 2 µL DEectW-Sbio solution that was
diluted using HCM buffer was then added on the APTES-coated mica substrate for 3
min. The DEectW-Sbio solution concentration was adjusted based on the pilot
observation. DO-5×bio:SA and the DEectW-Sbio were individually observed separately

using the method described in the previous section. HS-AFM was performed in HCM buffer at room temperature (25°C) using a laboratory-built HS-AFM instrument operated in tapping mode (Ando et al., 2001). BL-AC10DS-A2 cantilevers (Olympus Corporation, Tokyo, Japan) with a length, width, and thickness of 9, 2, and 0.13 μm, respectively, a spring constant of ~0.1 Nm−1, and resonant frequencies of 400–500 kHz were used in solution. The carbon tip at the end of the cantilever was grown via electron beam deposition and sharpened using argon gas etching to a tip radius of 2–5 nm (Wendel et al., 1995). The free oscillation amplitude of the cantilever was set to 1–2 nm and the amplitude setpoint for feedback control was ~80% of the free amplitude during scanning. Two or more independent protein purifications, DNA origami nanoblock preparation, and HS-AFM were performed. HS-AFM images were processed using the laboratory-made analysis software FalconViewer based on Igor Pro-8 (WaveMetrics, OR, USA).

Image processing
Images were processed using Image J, Photoshop 2023, or Illustrator 2023. The structural model of the designed DNA origami block was generated using the Web-based service offered by the Aksimentiev group (https://bionano.physics.illinois.edu/origami-structure). Molecular models were visualized and edited using the open-source version of PyMOL version 2.5.0 (https://www.pymol.org/). Schematic illustrations were created using either PowerPoint (Microsoft, WA, USA) or Illustrator (Adobe, CA, USA). Plots, processed images, and illustrations were assembled using PowerPoint.

Statistical analysis
OriginPro 2021 (OriginLab) was used for graphing and statistical analyses. One-tailed Welch's t-test was used to compare the mean of two groups in the bead aggregation assays. One-tailed Mann–Whitney U test was used to compare the mean ranks of two groups for measuring the distance between paired DNA origami blocks.

Data and materials availability
Raw data on TEM and HS-AFM of DNA origami objects and cadherin fragments obtained in this study are available on figshare (Oda et al., 2024). Additional data and materials will be available from the authors upon request.

CrediT author contribution statement
Hiroki Oda: Writing-review & editing, Writing-original draft, Funding acquisition, Visualization, Investigation, Methodology, Conceptualization
Shigetaka Nishiguchi: Writing-review & editing, Visualization, Investigation, Methodology, Conceptualization
Chihong Song: Investigation
Kazuyoshi Murata: Investigation, Resources, Funding acquisition
Takayuki Uchihashi: Methodology, Resources
Yuki Suzuki: Methodology, Investigation, Conceptualization

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments
The TEM work on the DNA origami objects was performed at the Hanaichi Ultrastructure Research Institute (Okazaki, Aichi, Japan). We would like to thank Ibuki Kawamata for help with installing caDNAno2; Christian Ganser for technical advice of HS-AFM; and Akiko Noda for technical assistance.

Source of funding
This work was supported in part by JSPS KAKENHI (22K05686 to HO) and the Cooperative Study Program (18NIPS232 to HO) of National Institute for Physiological Sciences.

Figure legends

Figure 1. Biotinylated whole or partial *Drosophila* E (DE)-cadherin ectodomain fragment preparation

(A) Schematic representation of the DE-cadherin ectodomain organization, featuring three DE-cadherin ectodomain constructs: DEectW, DEectW-S, and DEectD-S. Nt, N-terminal region; EC, extracellular cadherin domain; NC, non-chordate classical cadherin domain; CE, cysteine-rich EGF-like domain; LG, laminin globular domain. Proteolytic cleavage in the NC domain results in a membrane-distal polypeptide (MD) and a membrane-proximal polypeptide (MP), non-covalently bound to each other (Oda and Tsukita, 1999). (B) Schematic representation of DEectW-S biotinylation via the enzymatic SNAP-tag reaction with BG-biotin, resulting in DEectW-Sbio. (C, D) Coomassie Brilliant Blue (CBB) staining (C) and western blot immunodetection (D) of DEectW, DEectW-Sbio, and DEectD-Sbio separated using odium dodecyl-sulfate polyacrylamide gel electrophoresis. Approximately 5 pmol of the protein was loaded in each lane in C. The MD fragment was detected with DCAD2, whereas the MP fragment was detected with anti-V5 tag antibody. However, DEectD-Sbio lacked the DCAD2 epitope region (within EC2; (Nishiguchi et al., 2016)). (E) Native PAGE separation and CBB staining of DEectW-Sbio, DEectW-S, and DEectD-Sbio, pre-incubated with streptavidin (SA) at an approximate molar ratio of 1:5 for 10 min, or without SA. Approximately 10 pmol of the cadherin fragment was loaded on each lane.

Figure 2. Bead aggregation assays checking for specific adhesive properties of DEectW-Sbio

(A) Outline of the bead aggregation assay procedure. (B) Schematic illustration of an streptavidin (SA)-conjugated microbead capturing the DEect-Sbio molecule. (C) Images showing aggregate formation of DeectW-Sbio bound microbeads, with corresponding patterns of microbeads obtained for three negative controls, as indicated (DEectD-Sbio used instead of DeectW-Sbio; cadherin fragment omission; SA microbeads pre-incubated with excess biotin used). The boxed region in each panel is magnified below. (D) Quantification of the aggregation degree for assay results shown in C. (E) Images and plots showing increased aggregation of DEectW-Sbio binding microbeads with shaking duration time. (F) Images showing the Ca^{2+}-dependency of DEectW-Sbio

activity mediating SA microbead aggregation. (G) Quantification of aggregation degree for assay results shown in E. (H) Images and plots showing the concentration dependency of DEectW-Sbio activity. (I) Comparative measurement to assess the DEectW-Sbio protein quantity bound to two aliquots of SA microbeads at concentrations of 50 and 13 nM, with rough estimates of the average distance of DEectW-Sbio molecules on the SA microbead surface. For reference, one aliquot of SA microbeads is defined as the number of microbeads used in each well during aggregation assays, approximately 5×10^5 microbeads. Assuming that the DEectW-Sbio molecules were distributed evenly on the microbead surface, the relationship between the quantity of bound DEectW-Sbio molecules and average distance between the neighboring molecules was calculated. Error bars in D and F indicate standard deviation (s.d.), whereas bars in G and H indicate the mean and 95% confidence intervals. *, $p < 0.01$ (Welch's t-test). Scale bars; 10 µm.

Figure 3. Transmission electron microscopy (TEM) observation of DEectW-Sbio molecules docked onto the DNA origami structure via streptavidin (SA)

(A) Schematic representation of the DO-5×bio:SA nanostructure, characterized by a long block shape, an indentation at a corner, and five evenly spaced, biotinylated sites where SA molecules are selectively bound. The designed configuration places these biotinylated sites at intervals of ~21.5 nm along one long side of the DNA origami. (B) Agarose gel electrophoresis (AGE) analysis checking for DO-5×bio:SA docking with DEectW-SNAP-bio molecules. DO-5×bio:SA was mixed with DectW-S at 400 nM or with DEectW-Sbio at different concentrations (400, 40, 8, and 2 nM) and then incubated for 10 min. DNA migration pattern (left) and DNA stain intensity profiles (right) reveal a concentration-dependent increase in bio:SA site occupation by DEectW-Sbio molecules. (C) Structure prediction of DO-5×bio:SA (left) and TEM observation of negatively stained DO-5×bio:SA (right). (D–M) TEM observation of DO-5×bio:SA nanoblocks onto which DectW-Sbio (D–K) or DEectD-Sbio (L, M) molecules were docked. The docking was performed in the presence of 4 mM CaCl₂ (D–I, L, M) or in the absence of Ca²⁺ ions (J, K). Areas boxed in D–I are magnified in E'–I', respectively. Arrows pointing to the hinge-like site separating the membrane-distal (di) and -proximal (pr) portions of the *Drosophila* E (DE)-cadherin ectodomain. Closed arrowheads indicate folded configurations of the membrane-distal multiple ECs, whereas open arrowhead indicate an unfolded configuration of the membrane-distal ECs. Broken lines indicate symmetry axes in visualized molecular shapes. (N) A 2D class averaged image of negative-staining TEM images of DEectW molecules from 1,655 particles. The particles were classified into 15 classes and the major class #1 image is presented here (see supplementary Fig. 3 for details). (O) Proposed structural arrangement of DEectW, SNAP, SA, and DO-5×bio. Scale bars; 10 nm.

Figure 4. High-speed atomic force microscopy observation of DEectW-Sbio molecules docked onto DNA origami nanoblocks

(A) Two representative images of DO-5×bio:SA structures. Arrows point to the streptavidin (SA)s, which are evenly spaced, on the long side of the rectangle opposite to that with an asymmetric indentation shown with "Slope". (B) Two representative images of DEectW-Sbio molecules. Membrane-distal and -proximal portions are shown with "Distal" and "Proximal". (C) Two representative images of DEectW-Sbio docked

onto DNA origami nanoblocks. White arrowheads point to the DEectW-Sbio. (D) Sequential images showing extending and shrinking dynamics of membrane-distal potions of the single DEectW-Sbio on the DNA origami nanoblock. Membrane-distal tip region of DEectW-Sbio is indicated by a blue arrowhead. (E) Sequential images showing interactions between membrane-distal portions of two DEectW-Sbio on the same DNA origami nanoblock. Membrane-distal tip regions of two DEectW-Sbio are shown with blue arrowheads. (F) Sequential images showing structural dynamics of the dimer-like structure formed between DEectW-Sbio on DNA origami nanoblock and DNA origami-free DEectW-Sbio. A flexible feature of the membrane-distal portions of DEectW-Sbio with retaining dimer are shown. See Movies S1 to S3. Scale bars; 10 nm.

Figure 5. Induction and visualization of *Drosophila* E (DE)-cadherin ectodomain interactions assisted by DNA origami
(A) Schematic representation and transmission electron microscopy image of the DO-14×bio:SA nanoblock. Two sets of seven evenly spaced bio:SA sites, at intervals of ~7.2 nm, are placed in a line on a long, lateral face of the block. Bar, 10 nm. (B) Agarose gel electrophoresis analysis comparatively showing DO-5×bio:SA and DO-14×bio:SA docking with DEectW-Sbio molecules. (C) DNA origami nanoblock pairing procedure. (D–K) Images of single (D, G, J) and paired (E, F, H, I, K, L) DNA origami blocks, formed by combining DO-14×bio:SA with DEectW-Sbio in the presence or absence of Ca^{2+} ions and with DEectD-Sbio in the presence of 4 mM Ca^{2+} ions. Areas boxed in E, H, and K are magnified in E', H', and K', respectively. Bars, 10 nm. (M) Schematic explanation of the defined distance (D_{b-b}) and angle between paired DNA origami blocks. (N, O) Scatter plots of D_{b-b} and angle values (N) and box plots of D_{b-b} values (O) for DNA origami block pairs obtained from the three sample types of the DNA origami pairing are indicated in different colors (gray, DEectW-Sbio:CA; red, DEectW-Sbio:ED; and blue, DEectD:CA).

Figure S1. Amino acid sequence and domain composition of DEectW-S
The entire amino acid sequence of DEectW-S, consisting of 12 defined domains (Pro, NT, EC1 to EC7, NC, CE, and LG) and two tag regions (SNAP [S] and V5/6×His tags). DEectD-S is the same as DEectW-S except that the region indicated by asterisks is missing in DEectD-S. DEectW is the same as DEectW-S except that the SNAP- tag is not present in DEectW. Pro, including signal peptide, is removed for protein maturation. The cleavage in NC occurs between the Gly and Ser residues indicated by "><".

Figure S2. Bead aggregation assays using DEectW-Sbio in DNA origami folding buffer without or with Ca^{2+} ions
Assays were performed in the same way as in Fig. 1 except that DNA origami folding buffer (Tris-based, EDTA-containing) was used with (right) or without (left) adding 4 mM $CaCl_2$. Error bars indicate standard deviation (s.d.). *, p < 0.01 (Welch's t-test).

Figure S3. Agarose gel electrophoresis analysis showing the preparation of DO-5×bio and its binding to streptavidin
A 0.7% gel was used.

Figure S4. Transmission electron microscopy (TEM) analysis of DEectW molecules
(A) A TEM image of negative-stained, DEectW molecules. (B) 2D classification using 1,655 particles. Scale bars; 100 nm (A), 20 nm (B)

Figure S5. Extreme examples of paired DO-14×bio:SA nanoblocks
(A–C) Transmission electron microscopy images of paired DO-14×bio:SA nanoblocks having the five largest $D_{b\text{-}b}$ values, from each of the three sample types related to Fig. 5. (A) DO-14×bio:SA + DEectW-Sbio:CA. Arrows indicate string-like configurations of *Drosophila* E (DE)-cadherin ectodomain fragments contributing to bridging the apposed DNA origami nanoblocks. (B) DO-14×bio:SA + DEectW-Sbio:ED. (C) DO-14×bio:SA + DEectD-Sbio:CA.

Figure S6. DNA origami block design
Annealing patterns of scaffold (p8064; light blue) and staple DNAs (239 species; arbitrary multiple colors) are shown. Positions of biotinylated sites for DO-5×bio and DO-14×bio are indicated. See supplementary tables 1 and 2 for staple DNA details.

Movie S1. Extending and shrinking motion of the membrane-distal portion of DEectW-Sbio on DO-5×bio. The images were captured at a scanning speed of 0.3 s/frame. Image size: 60×60 nm^2 (60×60 pixels).

Movie S2. Binding and unbinding of neighboring DEectW-Sbio on DO-5×bio. The images were captured at a scanning speed of 0.3 s/frame. Image size: 60×60 nm^2 (60×60 pixels).

Movie S3. Flexible motion of dimer-like structure formed between DEectW-Sbio on DO-5×bio and DO-5×bio-free DEectW-Sbio. The images were captured at a scanning speed of 0.3 s/frame. Image size: 60×60 nm^2 (60×60 pixels).

Table S1. Polymerase chain reaction (PCR) primers used for plasmid construction.

Table S2. Staple sequences for the DNA origami block.

Table S3. Biotinylated staple DNAs.

References
Ando, T., Kodera, N., Takai, E., Maruyama, D., Saito, K., Toda, A., 2001. A high-speed atomic force microscope for studying biological macromolecules. Proc Natl Acad Sci U A 98, 12468–72. https://doi.org/10.1073/pnas.211400898
Bartsch, T.F., Hengel, F.E., Oswald, A., Dionne, G., Chipendo, I.V., Mangat, S.S., El Shatanofy, M., Shapiro, L., Müller, U., Hudspeth, A.J., 2019. Elasticity of individual protocadherin 15 molecules implicates tip links as the gating springs for hearing. Proc Natl Acad Sci U A 116, 11048–11056. https://doi.org/10.1073/pnas.1902163116

Baumgartner, W., Hinterdorfer, P., Ness, W., Raab, A., Vestweber, D., Schindler, H., Drenckhahn, D., 2000. Cadherin interaction probed by atomic force microscopy. Proc Natl Acad Sci U A 97, 4005–10. https://doi.org/10.1073/pnas.070052697

Boggon, T.J., Murray, J., Chappuis-Flament, S., Wong, E., Gumbiner, B.M., Shapiro, L., 2002. C-cadherin ectodomain structure and implications for cell adhesion mechanisms. Science 296, 1308–13. https://doi.org/10.1126/science.1071559

Brasch, J., Harrison, O.J., Honig, B., Shapiro, L., 2012. Thinking outside the cell: how cadherins drive adhesion. Trends Cell Biol. 22, 299–310. https://doi.org/10.1016/j.tcb.2012.03.004

Cavallaro, U., Christofori, G., 2004. Cell adhesion and signalling by cadherins and Ig-CAMs in cancer. Nat. Rev. Cancer 4, 118–132. https://doi.org/10.1038/nrc1276

Cavey, M., Rauzi, M., Lenne, P.-F.F.-. F.F., Lecuit, T., 2008. A two-tiered mechanism for stabilization and immobilization of E-cadherin. Nature 453, 751–6. https://doi.org/10.1038/nature06953

Chappuis-Flament, S., Wong, E., Hicks, L.D., Kay, C.M., Gumbiner, B.M., 2001. Multiple cadherin extracellular repeats mediate homophilic binding and adhesion. J Cell Biol 154, 231–43.

Clarke, D.N., Miller, P.W., Lowe, C.J., Weis, W.I., Nelson, W.J., 2016. Characterization of the Cadherin-Catenin Complex of the Sea Anemone Nematostella vectensis and Implications for the Evolution of Metazoan Cell-Cell Adhesion. Mol Biol Evol 33, 2016–29. https://doi.org/10.1093/molbev/msw084

Cooper, S.R., Jontes, J.D., Sotomayor, M., 2016. Structural determinants of adhesion by Protocadherin-19 and implications for its role in epilepsy. eLife 5, e18529. https://doi.org/10.7554/eLife.18529

Dey, S., Fan, C., Gothelf, K.V., Li, J., Lin, C., Liu, L., Liu, N., Nijenhuis, M.A.D., Saccà, B., Simmel, F.C., Yan, H., Zhan, P., 2021. DNA origami. Nat. Rev. Methods Primer 1, 1–24. https://doi.org/10.1038/s43586-020-00009-8

Douglas, S.M., Marblestone, A.H., Teerapittayanon, S., Vazquez, A., Church, G.M., Shih, W.M., 2009. Rapid prototyping of 3D DNA-origami shapes with caDNAno. Nucleic Acids Res. 37, 5001–5006. https://doi.org/10.1093/nar/gkp436

Eslami Amirabadi, H., Tuerlings, M., Hollestelle, A., SahebAli, S., Luttge, R., van Donkelaar, C.C., Martens, J.W.M., den Toonder, J.M.J., 2019. Characterizing the invasion of different breast cancer cell lines with distinct E-cadherin status in 3D using a microfluidic system. Biomed. Microdevices 21, 101. https://doi.org/10.1007/s10544-019-0450-5

Fan, X., Wang, J., Zhang, X., Yang, Z., Zhang, J.-C., Zhao, L., Peng, H.-L., Lei, J., Wang, H.-W., 2019. Single particle cryo-EM reconstruction of 52 kDa streptavidin at 3.2 Angstrom resolution. Nat. Commun. 10, 2386. https://doi.org/10.1038/s41467-019-10368-w

Gul, I.S., Hulpiau, P., Saeys, Y., van Roy, F., 2017. Evolution and diversity of cadherins and catenins. Exp Cell Res 358, 3–9. https://doi.org/10.1016/j.yexcr.2017.03.001

Harrison, O.J., Jin, X., Hong, S., Bahna, F., Ahlsen, G., Brasch, J., Wu, Y., Vendome, J., Felsovalyi, K., Hampton, C.M., Troyanovsky, R.B., Ben-Shaul, A., Frank, J., Troyanovsky, S.M., Shapiro, L., Honig, B., 2011. The extracellular architecture

of adherens junctions revealed by crystal structures of type I cadherins. Structure 19, 244–56. https://doi.org/10.1016/j.str.2010.11.016

Hulpiau, P., Van Roy, F., 2010. New Insights into the Evolution of Metazoan Cadherins. Mol Biol Evol 28, 647–57. https://doi.org/10.1093/molbev/msq233

Hynes, R.O., Zhao, Q., 2000. The evolution of cell adhesion. J Cell Biol 150, F89-96.

Janiszewska, M., Primi, M.C., Izard, T., 2020. Cell adhesion in cancer: Beyond the migration of single cells. J. Biol. Chem. 295, 2495–2505. https://doi.org/10.1074/jbc.REV119.007759

Kovacs, E.M., Goodwin, M., Ali, R.G., Paterson, A.D., Yap, A.S., 2002. Cadherin-Directed Actin Assembly: E-Cadherin Physically Associates with the Arp2/3 Complex to Direct Actin Assembly in Nascent Adhesive Contacts. Curr. Biol. 12, 379–382. https://doi.org/10.1016/S0960-9822(02)00661-9

Lambert, O., Taveau, J.-C., Him, J.L.K., Al Kurdi, R., Gulino-Debrac, D., Brisson, A., 2005. The basic framework of VE-cadherin junctions revealed by cryo-EM. J. Mol. Biol. 346, 1193–1196. https://doi.org/10.1016/j.jmb.2004.12.053

Maker, A., Bolejack, M., Schecterson, L., Hammerson, B., Abendroth, J., Edwards, T.E., Staker, B., Myler, P.J., Gumbiner, B.M., 2022. Regulation of multiple dimeric states of E-cadherin by adhesion activating antibodies revealed through Cryo-EM and X-ray crystallography. PNAS Nexus 1, pgac163. https://doi.org/10.1093/pnasnexus/pgac163

Marjoram, R.J., Guilluy, C., Burridge, K., 2016. Using magnets and magnetic beads to dissect signaling pathways activated by mechanical tension applied to cells. Methods, Experimental Approaches in Mechanotransduction 94, 19–26. https://doi.org/10.1016/j.ymeth.2015.09.025

Nagafuchi, A., Shirayoshi, Y., Okazaki, K., Yasuda, K., Takeichi, M., 1987. Transformation of cell adhesion properties by exogenously introduced E-cadherin cDNA. Nature 329, 341–3. https://doi.org/10.1038/329341a0

Nathaniel Clarke, D., Lowe, C.J., James Nelson, W., 2019. The cadherin-catenin complex is necessary for cell adhesion and embryogenesis in Nematostella vectensis. Dev Biol 447, 170–181. https://doi.org/10.1016/j.ydbio.2019.01.007

Nichols, S.A., Roberts, B.W., Richter, D.J., Fairclough, S.R., King, N., 2012. Origin of metazoan cadherin diversity and the antiquity of the classical cadherin/β-catenin complex. Proc Natl Acad Sci U A 109, 13046–51. https://doi.org/10.1073/pnas.1120685109

Nishiguchi, S., Furuta, T., Uchihashi, T., 2022. Multiple dimeric structures and strand-swap dimerization of E-cadherin in solution visualized by high-speed atomic force microscopy. Proc. Natl. Acad. Sci. 119, e2208067119. https://doi.org/10.1073/pnas.2208067119

Nishiguchi, S., Oda, H., 2021. Structural variability and dynamics in the ectodomain of an ancestral-type classical cadherin revealed by AFM imaging. J. Cell Sci. 134, jcs258388. https://doi.org/10.1242/jcs.258388

Nishiguchi, S., Yagi, A., Sakai, N., Oda, H., 2016. Divergence of structural strategies for homophilic E-cadherin binding among bilaterians. J Cell Sci 129, 3309–19. https://doi.org/10.1242/jcs.189258

Oda, H., 2012. Evolution of the cadherin-catenin complex. Subcell Biochem 60, 9–35. https://doi.org/10.1007/978-94-007-4186-7_2

Oda, H., Nishiguchi, S., Suzuki, Y. Data for: Nanoscale visualization of of *Drosophila* E-cadherin ectodomain fragments and their interactions using DNA origami nanoblocks. figshare. 2024. https://figshare.com/s/310b6b21dbd5d6c7607a

Oda, H., Tagawa, K., Akiyama-Oda, Y., 2005. Diversification of epithelial adherens junctions with independent reductive changes in cadherin form: identification of potential molecular synapomorphies among bilaterians. Evol Dev 7, 376–89. https://doi.org/10.1111/j.1525-142X.2005.05043.x

Oda, H., Takeichi, M., 2011. Evolution: Structural and functional diversity of cadherin at the adherens junction. J Cell Biol 193, 1137–46. https://doi.org/10.1083/jcb.201008173

Oda, H., Tsukita, S., 1999. Nonchordate classic cadherins have a structurally and functionally unique domain that is absent from chordate classic cadherins. Dev Biol 216, 406–22. https://doi.org/10.1006/dbio.1999.9494

Oda, H., Uemura, T., Harada, Y., Iwai, Y., Takeichi, M., 1994. A Drosophila homolog of cadherin associated with armadillo and essential for embryonic cell-cell adhesion. Dev Biol 165, 716–26. https://doi.org/10.1006/dbio.1994.1287

Pokutta, S., Herrenknecht, K., Kemler, R., Engel, J., 1994. Conformational changes of the recombinant extracellular domain of E-cadherin upon calcium binding. Eur J Biochem 223, 1019–26.

Putnam, N.H., Srivastava, M., Hellsten, U., Dirks, B., Chapman, J., Salamov, A., Terry, A., Shapiro, H., Lindquist, E., Kapitonov, V.V., Jurka, J., Genikhovich, G., Grigoriev, I.V., Lucas, S.M., Steele, R.E., Finnerty, J.R., Technau, U., Martindale, M.Q., Rokhsar, D.S., 2007. Sea anemone genome reveals ancestral eumetazoan gene repertoire and genomic organization. Science 317, 86–94. https://doi.org/10.1126/science.1139158

Sasaki, M., Akiyama-Oda, Y., Oda, H., 2017. Evolutionary origin of type IV classical cadherins in arthropods. BMC Evol. Biol. 17, 142. https://doi.org/10.1186/s12862-017-0991-2

Scheres, S.H.W., 2012. RELION: Implementation of a Bayesian approach to cryo-EM structure determination. J. Struct. Biol. 180, 519–530. https://doi.org/10.1016/j.jsb.2012.09.006

Schneider, I., 1972. Cell lines derived from late embryonic stages of Drosophila melanogaster. J Embryol Exp Morphol 27, 353–65.

Srivastava, M., Simakov, O., Chapman, J., Fahey, B., Gauthier, M.E.A., Mitros, T., Richards, G.S., Conaco, C., Dacre, M., Hellsten, U., Larroux, C., Putnam, N.H., Stanke, M., Adamska, M., Darling, A., Degnan, S.M., Oakley, T.H., Plachetzki, D.C., Zhai, Y., Adamski, M., Calcino, A., Cummins, S.F., Goodstein, D.M., Harris, C., Jackson, D.J., Leys, S.P., Shu, S., Woodcroft, B.J., Vervoort, M., Kosik, K.S., Manning, G., Degnan, B.M., Rokhsar, D.S., 2010. The Amphimedon queenslandica genome and the evolution of animal complexity. Nature 466, 720–6. https://doi.org/10.1038/nature09201

Stahl, E., Martin, T.G., Praetorius, F., Dietz, H., 2014. Facile and scalable preparation of pure and dense DNA origami solutions. Angew Chem Int Ed Engl 53, 12735–40. https://doi.org/10.1002/anie.201405991

Strale, P.-O., Duchesne, L., Peyret, G., Montel, L., Nguyen, T., Png, E., Tampé, R., Troyanovsky, S., Hénon, S., Ladoux, B., Mège, R.-M., 2015. The formation of

ordered nanoclusters controls cadherin anchoring to actin and cell–cell contact fluidity. J. Cell Biol. 210, 333–346. https://doi.org/10.1083/jcb.201410111

Suzuki, Y., Endo, M., Sugiyama, H., 2015. Lipid-bilayer-assisted two-dimensional self-assembly of DNA origami nanostructures. Nat Commun 6, 8052. https://doi.org/10.1038/ncomms9052

Tepass, U., Hartenstein, V., 1994. The development of cellular junctions in the Drosophila embryo. Dev Biol 161, 563–96. https://doi.org/10.1006/dbio.1994.1054

Truong Quang, B.-A., Mani, M., Markova, O., Lecuit, T., Lenne, P.-F., 2013. Principles of E-Cadherin Supramolecular Organization In Vivo. Curr. Biol. 23, 2197–2207. https://doi.org/10.1016/j.cub.2013.09.015

Wendel, M., Lorenz, H., Kotthaus, J.P., 1995. Sharpened electron beam deposited tips for high resolution atomic force microscope lithography and imaging. Appl. Phys. Lett. 67, 3732–3734. https://doi.org/10.1063/1.115365

Yap, A.S., Gomez, G.A., Parton, R.G., 2015. Adherens Junctions Revisualized: Organizing Cadherins as Nanoassemblies. Dev. Cell 35, 12–20. https://doi.org/10.1016/j.devcel.2015.09.012

A

Incubate SA beads with DEectW-Sbio for 5 min

⇩

Add excess biotin to the beads and wait for 1 min

⇩

Shake the beads in 4-mm wells at 1,500 rpm

⇩

Let the wells stand still for 10 min

⇩

Take photos of beads sunk on the well bottom

⇩

Determine the degree of aggregation for each well

B

DEectW-Sbio

Magnetic bead coated with SA

~1.5 μm

● Biotin (bio)
◆ Streptavidin (SA)
▢ SNAP-tag (S)

C

DEectW-Sbio   DEectD-Sbio   No cadherin added   Biotin pre-incubated

D

Degree of aggregation (pix)

DEectW-Sbio   DEectD-Sbio   No cadherin added   Biotin pre-incubated

*   *   *

E

DEectW-Sbio
+Ca²⁺   +EGTA

F

Degree of aggregation (pix)

*

+Ca²⁺   +EGTA
DEectW-Sbio

G

Degree of aggregation (pix)

50 nM   13 nM   3 nM

*   *   ns

50   13   3   0
DEectW-Sbio concentration (nM)

H

0 min   1 min   2 min   3 min   4 min   5 min

Degree of aggregation (pix)

0   1   2   3   4   5
Shaking duration (min)

I

1: 2× Proteins bound to SA beads/well at 50 nM
2: 2× Proteins bound to SA beads/well at 13 nM
(5 × 10⁵ beads/well)

DEectW-Sbio						
Quantity indicator (fmol)	40	20	16	12	8	4
D_m (nm)	19	27	30	35	42	60

D_m = Estimated average distance between molecules

Average surface area = 7 × 10⁶ nm²

D_m Bead

A

DNA origami-5 × bio:SA
(DO-5 × bio:SA)

- ● Biotin (bio)
- ◆ Streptavidin (SA)
- ▢ SNAP-tag (S)

~21.5 nm

DEectW-Sbio

C

Prediction   TEM

B

M: λ DNA/HindIII + ΦX174 DNA/HaeIII markers
1: DO-5 × bio:SA (0.08 nM, 8052 bp)
2: DO-5 × bio:SA + DEectW-S (400 nM)
3: DO-5 × bio:SA + DEectW-Sbio (400 nM)
4: DO-5 × bio:SA + DEectW-Sbio (40 nM)
5: DO-5 × bio:SA + DEectW-Sbio (8 nM)
6: DO-5 × bio:SA + DEectW-Sbio (2 nM)

Markers (bp): 23130, 9416, 6557, 4361, 2322, 2027, 1353, 1078, 872, 603

Marker (bp): 9416, 6557, 4361, 2322, 2027

Intensity (a.u.)

Migration distance (cm)

DO-5 × bio:SA + DEectW-Sbio:CA

D **E** **F** **G** **H** **I**

D' di, pr **E'** **F'** **G'** **H'** **I'**

N **O** SNAP, SA

DO-5 × bio:SA + DEectW-Sbio:ED

J **K**

DO-5 × bio:SA + DEectD-Sbio:CA

L **M**

A DNA origami-14 × bio:SA
(DO-14 × bio:SA)

SNAP
bio
SA

~7.2 nm

DEectW-Sbio

TEM

B

Markers (bp)
23130 —
9416 —
6557 —
4361 —
2322 —
2027 —
1353 —
1078 —
872 —
603 —

1 2 3 4

C

Mix 0.08 nM DO-14 × bio:SA with 8 nM DEectW-Sbio

⬇

Incubate the mixture for 5 h on ice

⬇

Shake the mixture at 1,500 rpm for 5 min

⬇

Negative staining and TEM observation

1: DO-5 × bio:SA
(0.08 nM)

2: DO-5 × bio:SA + DEectW-Sbio
(0.08 nM)   (400 nM)

3: DO-14 × bio:SA
(0.08 nM)

4: DO-14 × bio:SA + DEectW-Sbio
(0.08 nM)   (400 nM)

DO-14 × bio:SA + DEectW-Sbio:Ca

D **E** D_{b-b} = 33.1 nm **F** D_{b-b} = 33.8 nm **F'**

DO-14 × bio:SA + DEectW-Sbio:ED

G **H** D_{b-b} = 19.5 nm **I** D_{b-b} = 20.5 nm **I'**

DO-14 × bio:SA + DEectD-Sbio:CA

J **K** D_{b-b} = 19.5 nm **L** D_{b-b} = 20.9 nm **L'**

M

$$D_{b-b} = (d_{ij} + d_{ji}) / 2$$

m_i

Block i

d_{ij}

Block j

Angle

N

D_{b-b} (nm)

Angle (degree)

- DEectW-Sbio:CA
- DEectW-Sbio:ED
- DEectD-Sbio:CA

O

D_{b-b} (nm)

** **

n = 47   n = 22   n = 23

DEectW-Sbio:CA   DEectW-Sbio:ED   DEectD-Sbio:CA