Unraveling genetic load dynamics during biological invasion: insights from two invasive insect species

Eric Lombaert[1], Aurélie Blin[1], Barbara Porro[1,2], Thomas Guillemaud[1], Julio S. Bernal[3], Gary Chang[4], Natalia Kirichenko[5,6,7], Thomas W. Sappington[8,9], Stefan Toepfer[10,11] and Emeline Deleury[1]

[1] INRAE, Université Côte d'Azur, ISA, Sophia-Antipolis, France
[2] Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Univ Evry, Université Paris-Saclay, Evry, France
[3] Department of Entomology, Texas A&M University, College Station, TX 77843-2475, USA
[4] Biology Department, Gonzaga University, WA, USA
[5] Sukachev Institute of Forest, Siberian Branch of the Russian Academy of Sciences, Federal Research Center "Krasnoyarsk Science Center SB RAS", Akademgorodok 50/28, 660036 Krasnoyarsk, Russia
[6] Institute of Ecology and Geography, Siberian Federal University, Svobodny pr. 79, 660041 Krasnoyarsk, Russia
[7] All-Russian Plant Quarantine Center, Krasnoyarsk branch, Krasnoyarsk, Zhelyabova Str., 6/6, 660020, Russia
[8] USDA-Agricultural Research Service, Corn Insects & Crop Genetics Research Unit, Ames, Iowa, USA
[9] Iowa State University, Department of Plant Pathology, Entomology, and Microbiology, Ames, Iowa, USA
[10] CABI Switzerland, c/o Plant Protection and Soil Conservation Directorate, Hodmezovasarhely, Hungary
[11] MARA-CABI Joint Laboratory for Biosafety, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, China

Corresponding author:
Eric Lombaert – Institut Sophia Agrobiotech – 400, route des chappes – BP 167 – 06903 Sophia-Antipolis Cedex - France
E-mail: eric.lombaert@inrae.fr
Tel: +33 4 92 38 64 81

Keywords: target enrichment, non-model organism, population genomics, pool-sequencing, *Harmonia axyridis*, *Diabrotica virgifera virgifera*, genetic load

Running title: Evolution of genetic load in two invasive species

Abstract

Many invasive species undergo a significant reduction in genetic diversity, i.e. a genetic bottleneck, in the early stages of invasion. However, this reduction does not necessarily prevent them from achieving considerable ecological success and becoming highly efficient colonizers. Here we investigated the purge hypothesis, which suggests that demographic bottlenecks may facilitate conditions (e.g., increased homozygosity and inbreeding) under which natural selection can purge deleterious mutations, thereby reducing genetic load. We used a transcriptome-based exome capture protocol to identify thousands of SNPs in coding regions of native and invasive populations of two highly successful invasive insect species, the western corn rootworm (Chrysomelidae: *Diabrotica virgifera virgifera*) and the harlequin ladybird (Coccinelidae: *Harmonia axyridis*). We categorized and polarized SNPs to investigate changes in genetic load between invasive populations and their sources. Our results differed between species. In *D. virgifera virgifera*, although there was a general reduction in genetic diversity in invasive populations, including that associated with genetic load, we found no clear evidence for purging of genetic load, except marginally for highly deleterious mutations in one European population. Conversely, in *H. axyridis*, the reduction in genetic diversity was minimal, and we detected signs of genetic load fixation in invasive populations. These findings provide new insights into the evolution of genetic load during invasions, but do not offer a definitive answer to the purge hypothesis. Future research should include larger genomic datasets and a broader range of invasive species to further elucidate these dynamics.

Introduction

Biological invasions represent a significant aspect of global change, profoundly impacting biodiversity through the alteration of species distributions worldwide, particularly in recent times due to the significant increase in human-assisted dispersal (Seebens et al. 2017). The key factors determining the success of invasive species are not yet fully understood, although numerous hypotheses have been proposed (Enders et al. 2018; Sherpa and Després 2021; Daly et al. 2023). Most proposals fail to account for one or more consistent characteristics of successful invasions: (i) the rarity of successful invasions resulting from introductions (Williamson and Fitter 1996), (ii) the lag time between initial introduction and invasion (Sakai et al. 2001), (iii) the frequent reduction in genetic diversity due to demographic bottlenecks (Nei et al. 1975), and (iv) the prevalence of multiple invasions originating from an initial invasive population (i.e. the bridgehead effect; Lombaert et al. 2010). These characteristics suggest that the success of invasions may stem from partially stochastic biological processes spanning multiple generations and combining both demographic and genetic mechanisms.

A hypothesis that aligns with these characteristics is that genetic load is purged in the initial stages of invasions, i.e. deleterious alleles that cause inbreeding depression are eliminated from the introduced population (Estoup et al. 2016). During the demographic bottleneck following introduction of a few individuals into a new environment, genetic drift intensifies and homozygosity increases. While genetic drift may randomly eliminate some deleterious alleles, thereby reducing part of the genetic load, it also contributes to the transformation of the masked load (i.e., the load which may become express in future generations; Bertorelle et al. 2022) into a realized load (i.e., the load which reduces fitness in the current generation; Bertorelle et al. 2022). This shift may result in a "mutational meltdown" (Lynch et al. 1995; Simberloff 2009), where expression of deleterious mutations increases risk of extinction, ultimately leading to failure of the introduced population to establish. Conversely, the increased homozygosity resulting from bottlenecks exposes recessive deleterious alleles to selection. In the short term, this exposure can reduce mean fitness, but if the population persists, purifying selection may progressively remove some highly deleterious alleles over multiple generations, potentially reducing genetic load (Dussex et al. 2023). Theoretically, such purges can occur under specific demographic (i.e., moderate reduction of population size) and genetic (i.e., highly deleterious, recessive alleles) conditions, optimizing exposure to natural selection (Crow 1970; Charlesworth et al. 1990; Glémin 2003; Robinson et al. 2023). The potential for such purging in introduced populations is particularly interesting for its role in facilitating successful invasions by reducing inbreeding depression and enabling inbred individuals to maintain high fitness levels.

Purging has been demonstrated empirically by measuring the evolution of various traits in artificially bottlenecked populations (Crnokrak and Barrett 2002; Avila et al. 2010). However, measuring life history traits can be challenging in the context of naturally occurring bottlenecks during biological invasions, and evidence of purging has been documented in only a few invasive species (Parisod et al. 2005; Mullarkey et al. 2013; Fountain et al. 2014; Marchini et al. 2016). One notable formal test of this hypothesis was conducted on the invasive Asian ladybird *Harmonia axyridis*, where measurement of life history traits revealed that invasive populations showed no evidence of the inbreeding depression observed in native ones, suggesting that deleterious alleles were purged during the invasion process (Facon et al. 2011). Overall, case studies focusing on the dynamics of genetic load during biological

invasions have either examined a few life history traits (e.g. Facon et al. 2011) or used a single
locus approach (e.g. Zayed et al. 2007), making it difficult to generalize the results.

Advances in population genomics over the past decade have provided promising
avenues for investigating genetic load on large scales. Initially explored in humans (Lohmueller
et al. 2008; Henn et al. 2015), these approaches have been applied in the fields of
domestication (e.g. Schubert et al. 2014; Marsden et al. 2016; Makino et al. 2018; Wang et al.
2021) and conservation (Xue et al. 2015; Robinson et al. 2016; Grossen et al. 2020; Dussex et
al. 2021; Ochoa and Gibbs 2021). These studies typically involve SNP-calling within coding
regions, determination of ancestral SNP states, categorization of the severity of fitness
reduction caused by derived alleles, and population comparisons while accounting for genetic
drift using synonymous and/or intergenic polymorphisms. High quality genomic resources are
essential for such studies, which may explain the limited application of these methods to
invasive species, which are mostly non-model organisms. However, advances in genome
sequencing technologies and bioinformatics have significantly reduced costs and improved
accessibility, making these methods increasingly routine and affordable (Bertorelle et al.
2022).

In this study, we examined the dynamics of genetic load during the invasion of two
successful insect species, namely the western corn rootworm, *Diabrotica virgifera virgifera*,
and the harlequin ladybird, *Harmonia axyridis*, by directly measuring/estimating genetic load
using genomic data from feral native and invasive populations. Importantly, our study was not
designed to assess the instrumental role of purging in invasion success, but rather to
investigate its occurrence during these invasions. A broader study of both successful and failed
invasions across many species would be required to draw conclusions on purging's role in
invasion success. We used a pool-seq transcriptome-based exome capture protocol previously
developed for non-model species (Deleury et al. 2020) to identify SNPs within coding
sequences and categorize them as synonymous, moderately deleterious, or highly
deleterious. Our results offer insights into the fate of the genetic load and provide valuable
perspectives on the purge hypothesis in the context of biological invasions.

Methods

Species choice

The two species, *Diabrotica virgifera virgifera* (hereafter DVV) and *Harmonia axyridis* (HA), are
good candidates for testing the purging of genetic load hypothesis. Both species are highly
successful invaders with extensive invasive ranges (Gray et al. 2009; Roy et al. 2016).
Additionally, their invasion routes are well-documented and supported by robust analyses
using diverse datasets and methodologies (Miller et al. 2005; Ciosi et al. 2008; Lombaert et al.
2010, 2011, 2014, 2018). Both species exhibit bridgehead effects, meaning that certain
invasive populations have played a crucial role in global dissemination (Guillemaud et al.
2011). Additionally, purging of genetic load was detected in HA in a laboratory study focusing
on life-history traits (Facon et al. 2011).

Design of exome capture probes for target enrichment

150 Design of the exome capture probes was performed as described in Deleury et al. (2020). We
151 searched for peptide-coding sequences using FRAMEDP (v1.2.2; Gouzy et al. 2009) on the *de*

novo transcriptomes described in Coates et al. (2021) and Vogel et al. (2017) for DVV and HA, respectively. BLASTX results (e-value ≤ 1e^{-7}) of transcripts were used against the insect proteomes of *Tribolium castaneum*, *Anoplophora glabripennis*, *Dendroctonus ponderosae*, *Drosophila melanogaster* and the SwissProt database (v2016-02-17) for training. From the obtained coding sequences (CDS), we eliminated (i) *Wolbachia* and other putative endosymbiont sequences, (ii) CDS with > 1% missing nucleotide bases (Ns) or with more than four consecutive Ns, (iii) CDS with a GC% below 25 or above 75 and (iv) CDS with lengths < 400 bp or > 3500 bp. From the remaining 7,132 and 12,739 CDS for DVV and HA respectively, we drew at random c.a. 5.5 Mb for each species.

Probes based on the selected CDS were designed and manufactured by NimbleGen. In the case of DVV, repetitiveness of the probes was assessed based on the highest 15-mer frequency among the genomes of *Tribolium castaneum* (GCA_000002335.3), *Dendroctonus ponderosae* (GCA_000355655.1 and GCA_000346045.2) and DVV (the one available at the time, GCA_003013835.2). Probes with more than five close matches in the DVV genome were discarded. Close matches were defined as no more than five single-base insertions, deletions or substitutions using the SSAHA algorithm (Ning et al. 2001). Probes that matched sequences in the mitochondrial genome were also discarded. Random nucleotides were used to replace residual Ns in target sequences. For HA, the method used to ensure probe uniqueness is described in Deleury *et al.* (2020).

Overall, the final probe sets corresponded to a total of 4,151 CDS (5,282,603 bases across 12,017 regions of overlapping probes) and 5,717 CDS (5,347,461 bases across 6,400 regions of overlapping probes; Deleury et al. 2020) for DVV and HA respectively. This final set of probes was manufactured in the form of biotinylated DNA oligomers. One capture reaction contained 2,100,000 overlapping probes of 50 to 99 bp in length (mean length of 73.86 ± 4.46 bp for DVV, and 74.71 ± 4.92 bp for HA).

Sample collection

For both species, the choice of populations to be sampled was based on previously known invasion routes (Figure S1; Miller et al. 2005; Lombaert et al. 2010, 2018), so that each invasive population could be compared to its source. We sampled adult DVV at 6 sites: two in the native area (Mexico), two in North America (Colorado and Pennsylvania, respectively corresponding to the core and front of the first invasive population; Lombaert et al. 2018), and two in Europe (Hungary, referred to the Central-Southeastern European population, and north-western Italy, which are independently derived from the same source area of eastern North America; Miller et al. 2005). Adult HA were sampled at four sites: two in the native area (Russia [Siberia] and China), and two in North America (Pennsylvania and Washington, corresponding to two independent outbreaks from the native area; Lombaert et al. 2010, 2014). None of the selected invasive populations resulted from multiple introductions. The east North American population of HA was previously hypothesized to be an admixture between two populations (Lombaert et al. 2011), but our analysis using ABC (approximate Bayesian computation; Beaumont et al. 2002) with synonymous SNPs from the current dataset indicates that this is unlikely (see Appendix S1 for details).

To determine the ancestral and derived alleles for each SNP (i.e. to polarize the alleles), outgroup species were also sampled. For DVV, we selected the closely related species *Diabrotica adelpha*, as well as the more phylogenetically distant chrysomelid *Cerotoma trifurcata* (Eben and de los Monteros 2013). In the case of HA, from phylogenetically closest

to farthest (Tomaszewska et al. 2021), the sampled outgroup species were *Harmonia yedoensis*, *Harmonia conformis* and *Harmonia quadripunctata*. Complete information about samples is provided in Table 1 and Table S1.

Species	Population code name	Status (source)	Sampling site (Lat.; long.)	Sampling date	Haploid sample size
Diabrotica virgifera virgifera					
	D-N-MX1	Native	Canatlán, Mexico (24.554; -104.741)	Oct. 2015	80
	D-N-MX2	Native	Durango, Mexico (23.999; -104.614)	Oct. 2015	80
	D-I-COL	Invasive (D-N-MX1)	Fort Morgan, CO, USA (40.218; -103.869)	Aug. 2015	80
	D-I-PEN	Invasive (D-I-COL)	Landisville, PA, USA (40.119; -76.432)	Aug. 2015	80
	D-I-HUN	Invasive (D-I-PEN)	Kondoros, Hungary (46.736; 20.816)	July 2015	80
	D-I-ITA	Invasive (D-I-PEN)	Cuneo, Italy (44.463; 7.571)	July 2015	80
Harmonia axyridis					
	H-N-CHI	Native	Beijing, China (40.057; 116.540)	Oct. 2015	72
	H-N-RUS	Native	Krasnoyarsk, Russia (55.992; 92.757)	Sept. 2015	72
	H-I-PEN	Invasive (H-N-CHI)	Landisville, PA, USA (40.119; -76.432)	Aug. 2015	72
	H-I-WAS	Invasive (H-N-CHI)	Spokane, WA, USA (47.665; -117.403)	Nov. 2015	72
Diabrotica adelpha					
	D-OG-ADE	Outgroup DVV	Oaxaca State, Mexico (15.926; 97.151)	Jan. 2017	4
Cerotoma trifurcata					
	D-OG-TRI	Outgroup DVV	Weldon, CA, USA (35.660; -118.330)	Oct. 2016	4
Harmonia yedoensis					
	H-OG-YED	Outgroup HA	Beijing, China (40.031; 116.265)	Feb. 2015	2
Harmonia conformis					
	H-OG-CON	Outgroup HA	Mouans-Sartoux, France (43.620; 6.934)	Oct. 2009	2
Harmonia quadripunctata					
	H-OG-QUA	Outgroup HA	Rearing, Valbonne, France (NA)	July 2007	2

Table 1: Description of *Diabrotica virgifera virgifera* (DVV) and *Harmonia axyridis* (HA) samples, including outgroup species (see Table S1 for BioSample accessions). Sources of invasive populations were determined based on known invasion routes (Figure S1; Miller et al. 2005; Lombaert et al. 2010, 2018), except for H-I-PEN for which we retraced invasion routes in this study (see Appendix S1 for details). Because the two native DVV populations are genetically almost indistinguishable, we selected D-N-MX1 as the source of D-I-COL based on the smallest pairwise F_{ST} value between native and invasive populations (see Results section).

 DNA extraction, exome capture and pool sequencing

For each population, 4 legs from each of 40 individuals (DVV) or 2 legs from each of 36
individuals (HA) were pooled for DNA extraction with the Qiagen DNeasy Blood & Tissue kit,
in accordance with the manufacturer's recommendations. For the outgroup species, the same
kit was used to extract DNA from a single individual each for the three *Harmonia* species, and
from a pool of two individuals each for the *Diabrotica* and *Cerotoma* species.
Genomic libraries were prepared using NimbleGen SeqCap EZ HyperCap Library v2.0
and NimbleGen SeqCap EZ Library v5.0 for DVV and HA, respectively. In brief, for each of the
population and outgroup samples, DNA (2 µg in 100 µl) was mechanically sheared to an
average size of 200 bp using a Covaris S2 E210 device (6 cycles of 30 seconds each). In the case
of DVV, the fragmented DNA was then divided into three technical replicates for each
population. Subsequently, the fragments were subjected to end-repair, A-tailing and indexing
(with one unique index per sample) using the KAPA Library Preparation kit designed for
Illumina platforms. Following the ligation of Illumina adapters and indexes, only fragments
falling within the size range of 250 to 450 bp were retained. A PCR amplification step consisting
of 7 cycles was performed using standard Illumina paired-end primers, and the resulting
amplicons were purified using AMPure XP beads (Beckman). The length, quality, and
concentration of the prepared DNA fragments were assessed using a BioAnalyzer with Agilent
High Sensitivity DNA Assay, along with a Qubit.
For each capture (five for DVV and two for HA), we used a total of approximately 1 µg
of amplified DNA for exome enrichment, combining multiple samples in proportions that
allowed for equimolarity between population samples (see Table S2 and Table S3). This
enrichment was performed using the capture probes described above, following the
guidelines of either the NimbleGen SeqCap EZ HyperCap Library Protocol v2.0 or the SeqCap
EZ Library Protocol v5.0. Following each capture, we conducted two parallel PCRs, each
comprising 14 cycles, on the elution solution. In the case of HA, the resulting PCR products
were combined. Subsequently, all PCR products underwent purification using AMPure XP
beads. The length, quality, and concentration of the final DNA fragments were assessed using
a BioAnalyzer equipped with the Agilent High Sensitivity DNA Assay and a Qubit fluorometer.
For sequencing, we used one lane of an Illumina HiSeq3000 sequencer per species,
following the manufacturer's instructions, in paired-end mode for 150 cycles. After
sequencing, the data were demultiplexed and exported as FastQ files, and the libraries were
processed independently. For DVV, the FastQ files of technical replicates were merged prior
to subsequent analysis.

Mapping, SNP calling and annotation

Sequence quality assessment was conducted using FastQC v0.11.5 (Andrews 2010).
Subsequently, adapter sequences were removed, and low-quality base pairs were eliminated
using *Trimmomatic* v0.35 (Bolger et al. 2014), with the following parameter settings:
ILLUMINACLIP:TruSeq-file.fa:2:30:10 LEADING:25 TRAILING:25 SLIDINGWINDOW:5:30
MINLEN:75. We used almost the same parameters for the outgroup species sequences, but
with less stringency on quality of reads (SLIDINGWINDOW:5:20).
Filtered reads were mapped onto the genome assemblies PGI_DIABVI_V3a (GenBank
identifier: GCA_917563875.2) and icHarAxyr1.1 (GenBank identifier: GCA_914767665.1), for
DVV and HA respectively, using default options of the *bwa-mem* aligner v0.7.15 (Li 2013). We

used the *SAMtools* v1.15.1 software package (Li et al. 2009) and its *fixmate*, *view*, and
markdup tools to perform the following operations sequentially: (1) removal of unmapped
reads and secondary alignments, (2) filtering out read alignments with a mapping quality
Phred-score <20 and improperly paired reads, and (3) identification and removal of PCR
duplicates. Processing filtered reads for the outgroup species followed the same steps,
including mapping to the genome of the corresponding focal species, except that PCR
duplicates were not removed.
For each species, we conducted variant calling on the resulting *bam* alignment files
using Freebayes v1.3.6 software (Garrison and Marth 2012), with a focus on exonic regions
(options *-t* exons.bed *--pooled-continuous --min-alternate-count* 1 *--min-alternate-fraction*
0.001 *--min-alternate-total* 1 *--min-coverage* 80 *--use-best-n-alleles* 3). The resulting *vcf* file
was subsequently filtered using the *view* tool within *bcftools* v1.13 software (Danecek et al.
2021) and an in-house script to retain only true bi-allelic SNPs. Annotation was then
performed with the *SnpEff* program v5.0 (Cingolani et al. 2012b).
272
273   *Allele polarization*
274
SNP positions were extracted from both *vcf* files using the bcftools query tool. Subsequently,
for each outgroup species, we identified the nucleotides (or absence of typing) at positions
matching those in their respective focal species using the *samtools mpileup* tool on the
previously generated *bam* alignment files.
To polarize the SNPs, we used *est-sfs* software v2.03 (Keightley and Jackson 2018) with
the Kimura 2-parameter model. The input files contained data from one native population of
the focal species, together with the corresponding outgroup species, ensuring a consistent
phylogenetic topology for the software. Given that *est-sfs* computes the probability of the
most frequent allele being ancestral solely for polymorphic SNPs within the native population,
we used probabilities from a parallel analysis for monomorphic loci, incorporating an extra
haplotype for each allele. The complete process was repeated twice for each focal species,
considering the availability of two native populations for each. This resulted in two *est-sfs*
probabilities per focal species, both reflecting the likelihood of the most frequent allele being
the ancestral one.
To consider a SNP as polarized, we applied the following rules. If both probabilities
were above 0.5 for the same allele, and at least one of the probabilities exceeded 0.75, we
considered the most frequent allele as the ancestral one. If both probabilities were below 0.5
for the same allele, with at least one below 0.25, and if there were no more than two distinct
nucleotides present in total across the focal and outgroup species combined, we considered
the less frequent allele to be the ancestral one. SNPs that did not meet these criteria were
considered as not polarized but were retained for computations that did not require
polarization. Note that, because the est-sfs probabilities were predominantly close to 1 (Figure
S2), the threshold value had minimal impact on the number of polarized SNPs (Figure S3).
298
299   *Additional filters and categorization of deleterious mutations*
300
Key information was extracted from the *vcf* file using the *SnpSift* program v5.0 (Cingolani et
al. 2012a), including SNP coordinates, reference and alternative alleles, allele depths,
predicted effects, impact annotations, and protein-level changes. Additional filtering was
performed with an in-house R script (R Core Team 2021). First, we retained only biallelic SNPs

with coverage greater than 50 reads, falling below the 95th percentile of overall coverage in
each pool, and exhibiting a minor allele frequency exceeding 0.01 in at least one population.
Second, we retained SNPs located in coding regions with unambiguous annotations. Finally,
we excluded SNPs located on the X-chromosome in the specific case of HA, for which this
information was available.

SnpEff annotations were used to categorize the severity of fitness loss due to
mutations. Synonymous variants were used as proxies for neutral polymorphism. Non-
synonymous mutations were categorized as either "missense" (considered as potentially
moderately deleterious, involving amino acid changes) or "LoF" (loss-of-function, considered
as potentially highly deleterious, involving gain or loss of stop codons).

Genetic diversity and genetic load analyses

We computed several descriptive statistics on the whole set of SNPs. At the inter-population
level, we assessed genetic differentiation by calculating pairwise F_{ST} values using the R package
poolfstat v2.1.1 (Gautier et al. 2022). At the intra-population level, we used in-house R scripts
to compute the synonymous expected heterozygosity H_{eS}, as a measure of diversity, and the
ratio of non-synonymous to synonymous expected heterozygosity H_{eN}/H_{eS}, which provides an
indirect measure of the efficacy of selection. We used a 100-block jackknife resampling
approach to estimate means and standard errors. Additionally, we estimated bottleneck
intensities for all invasive populations of both species using ABC analyses (See Appendix S1 for
details).

Using polarized SNPs, we reported for each population and each severity category the
proportion of positions for which the derived allele is absent (frequency $f = 0$), rare ($f > 0$ and
$f \leq 0.1$), common ($f > 0.1$ and $f < 1$), or fixed ($f = 1$). The proportion of rare derived alleles
serves as a proxy for the masked load, whereas common and fixed alleles serve as a proxy for
the realized load. Additionally, within each severity category and population, we calculated
the mean derived allele frequency and estimated standard errors via a 100-block jackknife
resampling approach. Within each species and severity category, mean derived allele
frequencies were compared among populations using z-tests with false discovery rate (FDR;
Benjamini and Hochberg 1995) correction for multiple comparisons.

Finally, polarized SNPs were used to compute the R_{XY} statistic, which allows assessment
of relative excess or deficit of derived alleles within specific categories of deleterious
mutations in one population compared to another (Appendix S2; Do et al. 2015; Xue et al.
2015). This statistic effectively corrects for variation due to demography using neutral
mutations (synonymous SNPs in our case) as a reference point. In our analyses, population X
is an invasive population, and population Y is its source population (provided in Table 1). An
R_{XY} value of 1 indicates an equivalent relative genetic load in both populations, whereas values
greater or less than 1 suggest an excess (including fixation) or a deficit (including purge) of the
genetic load in the invasive population X, respectively. We assessed the significance of any
deviation from 1 using a z-score two-tailed test based on the 100-block jackknife resampling
method.

Results

SNP-calling and genomic variation

In the target enrichment experiments, we obtained mean numbers of raw read sequences of 50,196,834 and 63,360,676 for DVV and HA, respectively (BioProject PRJNA1079689; Table S1). After trimming, 85.0% and 93.8% of the sequences were retained for DVV and HA respectively. For outgroup species, the total number of raw reads was highly variable, with a mean of 2,631,775 of which 97.8% were conserved after trimming (see detailed information for all libraries, including outgroup species, in Table S4). Mapping, calling and filtering identified 66,274 SNPs (of which 62,034 could be polarized) and 169,755 SNPs (of which 169,102 could be polarized) within coding sequences for DVV and HA respectively (see Table S5 for details).

In the case of DVV, pairwise F_{ST} computed on the full set of SNPs showed patterns very close to those expected from results of previous analyses of microsatellite datasets (Figure S4). The differentiation between the two native Mexican populations was almost negligible with a mean F_{ST} estimate below 0.001. Given that the F_{ST} estimates between the invasive populations and the native sample D-N-MX1 (mean F_{ST} of 0.121) were consistently lower than those between the invasive populations and D-N-MX2 sample (mean F_{ST} of 0.128), D-N-MX1 was therefore considered the source of the D-I-COL sample (see Table 1). The Pennsylvania and Colorado populations had a low pairwise F_{ST} of 0.012, consistent with an Eastern expansion from Colorado (Figure S1), with plausible continuous gene flow. Conversely, all other F_{ST} estimates were relatively high, particularly between the native and European populations (mean F_{ST} value = 0.152). The highest value was observed between the two European populations at 0.169, confirming previous results indicating these populations originated from two independent introductions (Miller et al. 2005).

In the case of HA, F_{ST} estimates were moderate and fairly homogeneous, ranging from 0.020 to 0.050 (Figure S4). The smallest values were found between the eastern native population H-N-CHI and both invasive populations. The F_{ST} between invasive populations was slightly larger, consistent with previous results indicating independent origins for the two North American outbreaks (Lombaert et al. 2010). The highest values were observed between the western native population H-N-RUS and all other populations.

Within each species, invasive populations were characterized by reduced synonymous heterozygosity compared to their native counterparts, reflecting demographic bottlenecks encountered during the invasion process (Figure 1, x axis). One exception was the western native population of HA (H-N-RUS), which exhibited the lowest diversity within this species. Overall, the loss of diversity was more pronounced in DVV invasive populations, consistent with our ABC results, which indicate that bottlenecks were generally more intense in DVV than in HA (see Appendix S1). The ratio of non-synonymous to synonymous expected heterozygosity H_{eN}/H_{eS} is typically expected to increase in populations experiencing pronounced drift, as the reduced efficacy of selection allows more non-synonymous mutations, including potentially deleterious ones, to persist. In line with this expectation, we observed a higher ratio in invasive populations (Figure 1, y axis), and a negative correlation between this ratio and synonymous expected heterozygosity within each species (Pearson's $r = -0.95$, $P < 10^{-2}$ for DVV; Pearson's $r = -0.90$, $P < 10^{-1}$ for HA; Figure 1). Overall, the differences between native and invasive populations were more pronounced in DVV, with invasive ranges showing lower diversities and higher H_{eN}/H_{eS} ratios. This could suggest a

395 reduced efficacy of selection, although other factors may also contribute to the observed
396 patterns.
397

398
399
400 **Figure 1:** Ratio of non-synonymous to synonymous expected heterozygosity H_{eN}/H_{eS} vs. the
401 synonymous expected heterozygosity H_{eS} of each *Diabrotica virgifera virgifera* (DVV) and *Harmonia*
402 *axyridis* (HA) population. Means and standard errors (error bars) were determined from 100-block
403 jackknife resampling. See Table 1 for population codes (underlined population codes correspond to
404 the native populations).
405
406 *Genetic load on polarized SNPs*
407
In all populations studied and for each species, derived alleles were mostly rare (with
frequencies below 0.1) or absent (Figure 2). The highest derived allele frequencies consistently
showed a higher prevalence in synonymous positions compared to non-synonymous
positions. In DVV, we observed a substantial loss of non-synonymous derived alleles in
invasive populations compared to native ones. Although this was characterized by a reduction
in both masked load (proportion of non-synonymous SNPs with rare derived alleles ranging
from 0.496 to 0.523 in native populations and from 0.145 to 0.188 in invasive populations)
and realized load (proportion of non-synonymous SNPs with frequent or fixed derived alleles
ranging from 0.075 to 0.077 in native populations and from 0.046 to 0.051 in invasive
populations), it also came with slightly higher proportions of fixed derived alleles. This trend
was less pronounced for mutations inferred to be highly deleterious (LoF) than for those
inferred to be moderately deleterious (missense). For HA, the disparities between native and

invasive populations were considerably less pronounced (Figure 2). Compared to their native
Chinese source, HA invasive populations displayed a slight reduction in the proxy for masked
load (proportion of non-synonymous SNPs with rare derived alleles equal to 0.455 in the
Chinese population and from 0.347 to 0.381 in invasive populations) and a slight increase in
the proxy for realized load (proportion of non-synonymous SNPs with frequent or fixed
derived alleles equal to 0.019 in the Chinese population and from 0.024 to 0.027 in invasive
populations). Again, this trend was less pronounced for highly deleterious mutations (Figure
2).
428

429
430
431 **Figure 2**: Proportion of derived alleles found in each population of *Diabrotica virgifera virgifera* (DVV)
432 and *Harmonia axyridis* (HA) within categories of putative severity, classified by their frequency. See
433 Table 1 for population codes (underlined population codes correspond to the native populations).
434

As expected, the mean derived allele frequencies within populations decline with increasing putative severity in both species (Figure 3). For DVV, synonymous and missense mean derived allele frequencies were consistently significantly lower in the invasive populations than in the native populations ($P < 10^{-4}$ for all comparisons after FDR correction), whereas no significant differences were observed between native populations or between invasive populations ($P > 0.3$ for all comparisons after FDR correction). Within each of the three severity categories, mean derived allele frequencies were virtually identical in the four HA populations (P > 0.02 for all comparisons after FDR correction). No differences in LoF derived allele frequencies were significant ($P > 0.2$ for all comparisons after FDR correction), for either species. Overall, derived allele frequencies were lower in HA than in DVV.

Figure 3: Mean derived allele frequencies in all *Diabrotica virgifera virgifera* (DVV) and *Harmonia axyridis* (HA) populations in all three severity categories. Means and standard errors (error bars) were determined from 100-block jackknife resampling. See Table 1 for population codes (underlined population codes correspond to the native populations).

Finally, the R_{XY} ratio revealed no significant differences in relative frequencies of missense derived alleles between invasive populations and their respective sources in DVV (Figure 4). For putatively highly deleterious loss-of-function alleles, R_{XY} values sometimes deviate sharply from 1, but the number of SNPs is small (Table S5), and most differences are not statistically significant. The only exception is found in the population from Hungary (D-I-HUN), which shows reduced relative frequencies within the putatively highly deleterious category (LoF) of alleles (Figure 4). Regarding HA, slightly higher loads were significant in three of four comparisons across severity categories between the two invasive populations (H-I-PEN and H-I-WAS) and the native Chinese population (Figure 4).

464
465
Figure 4: Relative mutation load estimated as the R_{XY} ratio of derived alleles for moderate (missense)
and high (LoF) severity categories in populations of *Diabrotica virgifera virgifera* (D) and *Harmonia*
axyridis (H). A ratio below or above 1 indicates, respectively, a relative frequency deficit or excess of
derived alleles of indicated severity category in the invasive population (listed first) compared to its
population source (in parentheses). Means and standard errors (error bars) were determined from
100-block jackknife resampling. A star indicates that the corresponding R_{XY} value is significantly
different from 1 (z-score two-tailed test). For population codes, refer to Table 1 (underlined population
codes correspond to native populations).

474
475 ## Discussion
476
Exploring the evolution of genetic load using genomic data has long been restricted to model
species. Here, we successfully identified putatively deleterious derived alleles within coding
regions in two non-model invasive insect species using a pool-seq transcriptome-based exome
capture protocol (Deleury et al. 2020). By comparing native and invasive populations, we
tested the hypothesis that genetic load was purged during the invasion process, which may
enhance invasion success of certain populations (Sherpa and Després 2021; Daly et al. 2023).
Given that bottlenecks amplify genetic drift and weaken selection effectiveness (Crow 1970;
Glémin et al. 2003), the likelihood of purging is expected to increase for recessive, highly
deleterious mutations, while moderately deleterious mutations may still experience some
degree of drift, leading to higher frequencies (Agrawal and Whitlock 2011). Our study
exploring the dynamics of the genetic load in DVV and HA yielded mixed results, underscoring
the nuanced interplay of evolutionary forces in invasive species.
489
490 *Evolution of the genetic load in* Diabrotica virgifera virgifera
491
DVV exhibits marked genetic differences between native and invasive populations. Consistent
with previous studies (Ciosi et al. 2008; Lombaert et al. 2018), invasive populations, including

the long-established population in Colorado, display sharp declines in genetic diversity,
consistent with the strong bottlenecks inferred from ABC analyses. Compared to their native
counterparts, all invasive populations generally exhibit lower frequencies of derived alleles,
regardless of their potential fitness impacts, thus confirming the substantial role of genetic
drift in the invasion process of the species. While this overall reduction in genetic load within
invasive populations is apparent, it is worth emphasizing the presence of a slightly larger fixed
load.

In North America, despite a strong initial loss of genetic diversity, invasive populations
showed no significant relative deficit or excess of deleterious mutations compared to their
source, although a trend toward an excess of loss-of-function variants was observed.

Interestingly, the oldest European outbreak in Central-Southeastern Europe (here a
sample from Hungary) exhibited a significant deficit of "loss-of-function" deleterious
mutations compared to its source population, while "missense" deleterious mutations did not
differ significantly. This suggests that the demographic and selective constraints in this
population were effective at reducing highly deleterious mutations through a combination of
genetic drift and purifying selection, while moderately deleterious mutations have been less
affected. Such differences in evolutionary trajectories according to mutation severity are
theoretically expected (Whitlock 2002; Caballero et al. 2017; Dussex et al. 2023) and have
been observed in other species experiencing severe bottlenecks (Xue et al. 2015; Khan et al.
2021; Ochoa and Gibbs 2021; Wang et al. 2023).

Finally, the invasive northwestern Italian population, despite originating from an
independent introduction (Miller et al. 2005), showed patterns similar to that of the Central-
Southeastern European population. However, it did not exhibit significant signals of deficit or
excess of deleterious alleles, regardless of the mutation severity considered.

Evolution of the genetic load in Harmonia axyridis

In HA, the overall loss of genetic diversity in both invasive populations was relatively low, in
agreement with previous findings (Lombaert et al. 2010, 2011) and consistent with our ABC
results suggesting that bottlenecks were relatively mild in HA invasive populations. While the
putative masked genetic load decreased slightly, the putative realized load, although
consistently low in all populations, showed no sign of decreasing and even exhibited a slight
increase. This observation aligns with a significant trend, although subtle, towards a relative
excess of deleterious mutations in invasive populations compared to their native source,
regardless of putative fitness impact. This tendency to fix alleles contributing to genetic load,
likely driven by drift, is significant across all populations and severity categories, except for the
highly deleterious mutations in western North America (sampled in Washington).

These findings diverge from a previous study based on life history traits, which
demonstrated that invasive populations of HA experienced minimal inbreeding depression
while maintaining fitness comparable to outbred native populations (Facon et al. 2011). That
study has greatly contributed to the hypothesis of purging genetic load as a key mechanism
driving successful biological invasions (Estoup et al. 2016; Sherpa and Després 2021; Daly et
al. 2023). Given the markedly different methodologies used in the study by Facon et al. (2011)
and our study, it is challenging to ascertain why our results do not align. It is conceivable that
a few key genes could have disproportionate effects on invasion success, an aspect difficult to
quantify using our approach, particularly in the context of the overall low genetic load
previously described in this species (Tayeh et al. 2013). This issue is further complicated by the

imperfect nature of our categorization, where missense mutations likely cover a broad
spectrum of severity levels, potentially obscuring the influence of key genes. Moreover, very
highly deleterious mutations, which are critical contributors to inbreeding depression, are
expected to segregate at extremely low frequencies in large populations and may thus escape
detection in our sampling due to their rarity. Finally, our exome capture protocol only targets
a fraction of the exome, approximately one-third, and a broader approach may yield
qualitatively different results.
548
549 *Conclusions, limits and perspectives*
550
We tested the hypothesis that genetic load was purged during the invasion of two insect
species, a crop pest (DVV) and a predator (HA). At first glance, our results offer a nuanced
perspective. In the case of DVV, we were unable to detect any significant evolution of the
genetic load, except for the purge of highly deleterious mutations in the invasive Central-
Southeastern European population. For HA, we observed subtle evidence suggesting a
tendency toward fixation, rather than toward purge, of the genetic load. Although these
results are not entirely contradictory, they may stem from differences in initial genetic
composition, varying demographic history, and divergent ecological niches. For instance, the
two species experienced markedly different bottleneck intensities, yet our data do not allow
us to establish a direct link between these differences and the observed dynamics of genetic
load.
While population genomics approaches to assess the evolution of genetic load have
become increasingly popular in conservation and domestication biology (Moyers et al. 2018;
Bertorelle et al. 2022), our study is, to our knowledge, the first to use these methods to
specifically investigate the evolution of genetic load in the context of biological invasions.
However, it does not yet provide a comprehensive and definitive answer to the question
addressed. One limitation lies in our reliance on allelic frequencies derived from pool-seq data,
which precludes access to genotypic frequencies and therefore limits our ability to precisely
estimate inbreeding depression and explore the non-linear effects of genetic load (Bataillon
and Kirkpatrick 2000). While we used complementary statistics to mitigate this issue, future
studies based on individual-level sequencing would allow for more accurate assessments of
genetic variation and its relationship to genetic load. Additionally, to further broaden our
findings and enhance our statistical power, particularly for highly deleterious mutations, it will
be essential to consider the entire exome rather than only a fraction. Another promising
avenue involves categorizing mutation severity by quantifying evolutionary constraints within
a phylogenetic framework, which could enhance our understanding of genetic load evolution,
including in non-coding regions (Davydov et al. 2010). Furthermore, extending this research
to a wider range of species would enhance our ability to identify potential universal patterns
in the evolution of genetic load during biological invasions. Finally, to test not only the
occurrence of purging in invasions but also its role in the success of invasions, we must
compare the load dynamics in introductions that fail with those that succeed – a comparison
that is rarely feasible (Zenni and Nuñez 2013), except in controlled invasions such as in classical
biological control (Fauvergue et al. 2012).
584

Acknowledgments

We thank our colleagues Pamela Bruno, Charlyne Martignier-Jaccard, Pascal Maignet and Wang Su for *Diabrotica adelpha*, *Harmonia quadripunctata* and *Harmonia yedoensis* samples. We also thank Emmanuelle Murciano-Germain for administrative assistance. Sequencing was performed at the GENOTOUL GeT platform (https://get.genotoul.fr/en/).

Funding

This work was funded by grants from ANR project GENLOADICS, and INRAe SPE department. Sampling in Russia was supported by the basic project of the Sukachev Institute of Forest SB RAS (no. FWES-2024-0029).

Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the content of this article. TG and EL are both recommenders at PCI Evolutionary Biology and PCI Ecology. TG is co-founder of Peer Community In.

Data availability

Target capture pool sequencing files and all sequencing data have been deposited in Sequence Read Archive, National Center for Biotechnology Information, under project PRJNA1079689 (see also Table S1): https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1079689. All scripts used in this study have been deposited at Data INRAE: https://doi.org/10.57745/ESQFDB

Author contributions

EL and ED designed the study. JB, GC, NK, TS and ST managed the collection of samples. AB and ED prepared the samples and the libraries. ED and BP developed the bioinformatics pipelines. EL and ED analyzed the data. EL and TG wrote the paper. All authors have revised and approved the final manuscript.

References

Agrawal AF, Whitlock MC (2011) Inferences about the distribution of dominance drawn from yeast gene knockout data. Genetics 187:553–566. https://doi.org/10.1534/genetics.110.124560

Andrews S (2010) FastQC: a quality control tool for high throughput sequence data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Avila V, Amador C, Garcia-Dorado A (2010) The purge of genetic load through restricted panmixia in a Drosophila experiment. J Evol Biol 23:1937–1946. https://doi.org/10.1111/j.1420-9101.2010.02058.x

Bataillon T, Kirkpatrick M (2000) Inbreeding depression due to mildly deleterious mutations in finite populations: Size does matter. Genet Res 75:75–81.

https://doi.org/10.1017/S0016672399004048

Beaumont MA, Zhang WY, Balding DJ (2002) Approximate Bayesian computation in
population genetics. Genetics 162:2025–2035

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate - a practical and powerful
approach to multiple testing. J R Stat Soc Ser B-Methodological 57:289–300

Bertorelle G, Raffini F, Bosse M, et al (2022) Genetic load: genomic estimates and
applications in non-model animals. Nat Rev Genet 23:492–503.
https://doi.org/10.1038/s41576-022-00448-x

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: A flexible trimmer for Illumina sequence
data. Bioinformatics 30:2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Caballero A, Bravo I, Wang J (2017) Inbreeding load and purging: implications for the short-
term survival and the conservation management of small populations. Heredity (Edinb)
118:117–185. https://doi.org/10.1038/hdy.2016.80

Charlesworth D, Morgan MT, Charlesworth B (1990) Inbreeding Depression, Genetic Load,
and the Evolution of Outcrossing Rates in a Multilocus System With No Linkage.
Evolution (N Y) 44:1469–1489. https://doi.org/10.1111/j.1558-5646.1990.tb03839.x

Cingolani P, Patel VM, Coon M, et al (2012a) Using *Drosophila melanogaster* as a model for
genotoxic chemical mutational studies with a new program, SnpSift. Front Genet 3:1–9.
https://doi.org/10.3389/fgene.2012.00035

Cingolani P, Platts A, Wang LL, et al (2012b) A program for annotating and predicting the
effects of single nucleotide polymorphisms, SnpEff. Fly (Austin) 6:80–92.
https://doi.org/10.4161/fly.19695

Ciosi M, Miller NJ, Kim KS, et al (2008) Invasion of Europe by the western corn rootworm,
Diabrotica virgifera virgifera: Multiple transatlantic introductions with various
reductions of genetic diversity. Mol Ecol 17:3614–3627. https://doi.org/10.1111/j.1365-
294X.2008.03866.x

Coates BS, Deleury E, Gassmann AJ, et al (2021) Up-regulation of apoptotic- and cell survival-
related gene pathways following exposures of western corn rootworm to *B.
thuringiensis* crystalline pesticidal proteins in transgenic maize roots. BMC Genomics
22:1–27. https://doi.org/10.1186/s12864-021-07932-4

Crnokrak P, Barrett SCH (2002) Purging the genetic load: A review of the experimental
evidence. Evolution (N Y) 56:2347–2358. https://doi.org/10.1554/0014-
3820(2002)056[2347:pptgla]2.0.co;2

Crow JF (1970) Genetic loads and the cost of natural selection. In: Krickeberg K, Lewontin RC,
Neyman J, Schreiber M (eds) Mathematical topics in population genetics. Springer
Berlin Heidelberg, Berlin, pp 128–177

Daly EZ, Chabrerie O, Massol F, et al (2023) A synthesis of biological invasion hypotheses
associated with the introduction–naturalisation–invasion continuum. Oikos e09645:1–
29. https://doi.org/10.1111/oik.09645

Danecek P, Bonfield JK, Liddle J, et al (2021) Twelve years of SAMtools and BCFtools.
Gigascience 10:1–4. https://doi.org/10.1093/gigascience/giab008

Davydov E V., Goode DL, Sirota M, et al (2010) Identifying a high fraction of the human
genome to be under selective constraint using GERP++. PLoS Comput Biol 6:1–13.
https://doi.org/10.1371/journal.pcbi.1001025

Deleury E, Guillemaud T, Blin A, Lombaert E (2020) An evaluation of pool-sequencing
transcriptome-based exon capture for population genomics in non-model species.
bioRxiv 10.1101/58:v7 peer-reviewed and recommended by PCI Genomics.

677    https://doi.org/10.1101/583534

678 Do R, Balick D, Li H, et al (2015) No evidence that selection has been less effective at
679    removing deleterious mutations in Europeans than in Africans. Nat Genet 47:126–131.
680    https://doi.org/10.1038/ng.3186

681 Dussex N, Morales HE, Grossen C, et al (2023) Purging and accumulation of genetic load in
682    conservation. Trends Ecol Evol 38:961–969. https://doi.org/10.1016/j.tree.2023.05.008

683 Dussex N, van der Valk T, Morales HE, et al (2021) Population genomics of the critically
684    endangered kākāpō. Cell Genomics 100002:1–10.
685    https://doi.org/10.1016/j.xgen.2021.100002

686 Eben A, de los Monteros AE (2013) Tempo and mode of evolutionary radiation in
687    Diabroticina beetles (genera *Acalymma*, *Cerotoma*, and *Diabrotica*). Zookeys 332:207–
688    231. https://doi.org/10.3897/zookeys.332.5220

689 Enders M, Hutt MT, Jeschke JM (2018) Drawing a map of invasion biology based on a
690    network of hypotheses. Ecosphere 9:. https://doi.org/10.1002/ECS2.2146

691 Estoup A, Ravigné V, Hufbauer R, et al (2016) Is There A Genetic Paradox of Biological
692    Invasion? Annu Rev Ecol Evol Syst 47:51–72. https://doi.org/10.1146/annurev-ecolsys-
693    121415

694 Facon B, Hufbauer RA, Tayeh A, et al (2011) Inbreeding depression is purged in the invasive
695    insect *Harmonia axyridis*. Curr Biol 21:424–427.
696    https://doi.org/10.1016/j.cub.2011.01.068

697 Fauvergue X, Vercken E, Malausa T, Hufbauer RA (2012) The biology of small, introduced
698    populations, with special reference to biological control. Evol Appl 5:424–443.
699    https://doi.org/10.1111/j.1752-4571.2012.00272.x

700 Fountain T, Duvaux L, Horsburgh G, et al (2014) Human-facilitated metapopulation dynamics
701    in an emerging pest species, *Cimex lectularius*. Mol Ecol 23:1071–1084.
702    https://doi.org/10.1111/mec.12673

703 Garrison E, Marth G (2012) Haplotype-based variant detection from short-read sequencing.
704    arXiv 1207.3907:1–9. https://doi.org/10.48550/arXiv.1207.3907

705 Gautier M, Vitalis R, Flori L, Estoup A (2022) f-Statistics estimation and admixture graph
706    construction with Pool-Seq or allele count data using the R package poolfstat. Mol Ecol
707    Resour 22:1394–1416. https://doi.org/10.1111/1755-0998.13557

708 Glémin S (2003) How are deleterious mutations purged? Drift versus nonrandom mating.
709    Evolution (N Y) 57:2678–2687. https://doi.org/10.1111/j.0014-3820.2003.tb01512.x

710 Glémin S, Ronfort J, Bataillon T (2003) Patterns of Inbreeding Depression and Architecture of
711    the Load in Subdivided Populations. Genetics 165:2193–2212.
712    https://doi.org/10.1093/genetics/165.4.2193

713 Gouzy J, Carrere S, Schiex T (2009) FrameDP: Sensitive peptide detection on noisy matured
714    sequences. Bioinformatics 25:670–671. https://doi.org/10.1093/bioinformatics/btp024

715 Gray ME, Sappington TW, Miller NJ, et al (2009) Adaptation and Invasiveness of Western
716    Corn Rootworm: Intensifying Research on a Worsening Pest. Annu Rev Entomol 54:303–
717    321

718 Grossen C, Guillaume F, Keller LF, Croll D (2020) Purging of highly deleterious mutations
719    through severe bottlenecks in Alpine ibex. Nat Commun 11:1001.
720    https://doi.org/10.1038/s41467-020-14803-1

721 Guillemaud T, Ciosi M, Lombaert E, Estoup A (2011) Biological invasions in agricultural
722    settings: Insights from evolutionary biology and population genetics. C R Biol 334:237–
723    246. https://doi.org/10.1016/j.crvi.2010.12.008

Henn BM, Botigué LR, Bustamante CD, et al (2015) Estimating the mutation load in human
genomes. Nat Rev Genet 16:333–343. https://doi.org/10.1038/nrg3931

Keightley PD, Jackson BC (2018) Inferring the Probability of the Derived vs. the Ancestral
Allelic State at a Polymorphic Site. Genetics 209:897–906. https://doi.org/10.1007/978-
1-59745-494-0_7

Khan A, Patel K, Shukla H, et al (2021) Genomic evidence for inbreeding depression and
purging of deleterious genetic variation in Indian tigers. Proc Natl Acad Sci
118:e2023018118. https://doi.org/10.1073/pnas.2023018118

Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv 1303.3997:1–3. https://doi.org/10.48550/arXiv.1303.3997

Li H, Handsaker B, Wysoker A, et al (2009) The Sequence Alignment/Map format and
SAMtools. Bioinformatics 25:2078–2079.
https://doi.org/10.1093/bioinformatics/btp352

Lohmueller KE, Indap AR, Schmidt S, et al (2008) Proportionally more deleterious genetic
variation in European than in African populations. Nature 451:994–997.
https://doi.org/10.1038/nature06611

Lombaert E, Ciosi M, Miller NJ, et al (2018) Colonization history of the western corn
rootworm (*Diabrotica virgifera virgifera*) in North America: insights from random forest
ABC using microsatellite data. Biol Invasions 20:665–677.
https://doi.org/https://doi.org/10.1007/s10530-017-1566-2

Lombaert E, Guillemaud T, Cornuet JM, et al (2010) Bridgehead effect in the worldwide
invasion of the biocontrol harlequin ladybird. PLoS One 5:e9743. https://doi.org/e9743
10.1371/journal.pone.0009743

Lombaert E, Guillemaud T, Lundgren J, et al (2014) Complementarity of statistical treatments
to reconstruct worldwide routes of invasion: the case of the Asian ladybird *Harmonia*
axyridis. Mol Ecol 23:5979–5997. https://doi.org/10.1111/mec.12989

Lombaert E, Guillemaud T, Thomas CE, et al (2011) Inferring the origin of populations
introduced from a genetically structured native range by approximate Bayesian
computation: case study of the invasive ladybird *Harmonia axyridis*. Mol Ecol 20:4654–
4670. https://doi.org/10.1111/j.1365-294X.2011.05322.x

Lynch M, Conery IJ, Burger R (1995) Mutation Accumulation and the Extinction of Small
Populations. Am Nat 146:489–518. https://doi.org/10.1086/285812

Makino T, Rubin C-J, Carneiro M, et al (2018) Elevated proportions of deleterious genetic
variation in domestic animals and plants. Genome Biol Evol 10:276–290.
https://doi.org/10.1093/gbe/evy004/4794728

Marchini GL, Sherlock NC, Ramakrishnan AP, et al (2016) Rapid purging of genetic load in a
metapopulation and consequences for range expansion in an invasive plant. Biol
Invasions 18:183–196. https://doi.org/10.1007/s10530-015-1001-5

Marsden CD, Ortega-Del Vecchyo D, O'Brien DP, et al (2016) Bottlenecks and selective
sweeps during domestication have increased deleterious genetic variation in dogs. Proc
Natl Acad Sci 113:152–157. https://doi.org/10.1073/pnas.1512501113

Miller N, Estoup A, Toepfer S, et al (2005) Multiple transatlantic introductions of the western
corn rootworm. Science (80-) 310:992. https://doi.org/10.1126/science.1115871

Moyers BT, Morrell PL, McKay JK (2018) Genetic costs of domestication and improvement. J
Hered 109:103–116. https://doi.org/10.1093/jhered/esx069

Mullarkey A a., Byers DL, Anderson RC (2013) Inbreeding depression and partitioning of
genetic load in the invasive biennial *Alliaria petiolata* (Brassicaceae). Am J Bot 100:509–

518. https://doi.org/10.3732/ajb.1200403

Nei M, Maruyama T, Chakraborty R (1975) Bottleneck Effect and Genetic Variability in Populations. Evolution (N Y) 29:1–10. https://doi.org/10.2307/2407137

Ning Z, Cox AJ, Mullikin JC (2001) SSAHA: a fast search method for large DNA databases. - Abstract - UK PubMed Central. 1725–1729. https://doi.org/10.1101/gr.194201.1

Ochoa A, Gibbs HL (2021) Genomic signatures of inbreeding and mutation load in a threatened rattlesnake. Mol Ecol 30:5454–5469. https://doi.org/10.1111/mec.16147

Parisod C, Trippi C, Galland N (2005) Genetic variability and founder effect in the pitcher plant *Sarracenia purpurea* (Sarraceniaceae) in populations introduced into Switzerland: From inbreeding to invasion. Ann Bot 95:277–286. https://doi.org/10.1093/aob/mci023

R Core Team (2021) R: A language and environment for statistical computing

Robinson J, Kyriazis CC, Yuan SC, Lohmueller KE (2023) Deleterious Variation in Natural Populations and Implications for Conservation Genetics. Annu Rev Anim Biosci 11:93–114. https://doi.org/10.1146/annurev-animal-080522-093311

Robinson JA, Ortega-Del Vecchyo D, Fan Z, et al (2016) Genomic Flatlining in the Endangered Island Fox. Curr Biol 26:1183–1189. https://doi.org/10.1016/j.cub.2016.02.062

Roy HE, Brown PMJ, Adriaens T, et al (2016) The harlequin ladybird, *Harmonia axyridis*: global perspectives on invasion history and ecology. Biol Invasions 18:997–1044. https://doi.org/10.1007/s10530-016-1077-6

Sakai AK, Allendorf FW, Holt JS, et al (2001) The population biology of invasive species. Annu Rev Ecol Syst 32:305–332. https://doi.org/10.1146/annurev.ecolsys.32.081501.114037

Schubert M, Jónsson H, Chang D, et al (2014) Prehistoric genomes reveal the genetic foundation and cost of horse domestication. Proc Natl Acad Sci U S A 111:E5661-9. https://doi.org/10.1073/pnas.1416991111

Seebens H, Blackburn TM, Dyer E, et al (2017) No saturation in the accumulation of alien species worldwide. Nat Commun 8:14435. https://doi.org/10.1038/ncomms14435

Sherpa S, Després L (2021) The evolutionary dynamics of biological invasions: A multi-approach perspective. Evol Appl 1–23. https://doi.org/10.1111/eva.13215

Simberloff D (2009) The Role of Propagule Pressure in Biological Invasions. Annu Rev Ecol Evol Syst 40:81–102

Tayeh A, Estoup A, Hufbauer RA, et al (2013) Investigating the genetic load of an emblematic invasive species: the case of the invasive harlequin ladybird *Harmonia axyridis*. Ecol Evol 3:864–871. https://doi.org/10.1002/ece3.490

Tomaszewska W, Escalona HE, Hartley D, et al (2021) Phylogeny of true ladybird beetles (Coccinellidae: Coccinellini) reveals pervasive convergent evolution and a rapid Cenozoic radiation. Syst Entomol 46:611–631. https://doi.org/10.1111/syen.12479

Vogel H, Schmidtberg H, Vilcinskas A (2017) Comparative transcriptomics in three ladybird species supports a role for immunity in invasion biology. Dev Comp Immunol 67:452–456. https://doi.org/10.1016/j.dci.2016.09.015

Wang MS, Zhang JJ, Guo X, et al (2021) Large-scale genomic analysis reveals the genetic cost of chicken domestication. BMC Biol 19:1–16. https://doi.org/10.1186/s12915-021-01052-x

Wang X, Peischl S, Heckel G (2023) Demographic history and genomic consequences of 10,000 generations of isolation in a wild mammal. Curr Biol 33:2051-2062.e4. https://doi.org/10.1016/j.cub.2023.04.042

Whitlock MC (2002) Selection, load and inbreeding depression in a large metapopulation. Genetics 160:1191–1202. https://doi.org/10.1093/genetics/160.3.1191

Williamson M, Fitter A (1996) The varying success of invaders. Ecology 77:1661–1666

Xue Y, Prado-martinez J, Sudmant PH, et al (2015) Mountain gorilla genomes reveal the
impact of long-term population decline and inbreeding. Science (80-) 348:242–245.
https://doi.org/10.1126/science.aaa4484

Zayed A, Constantin S a., Packer L (2007) Successful biological invasion despite a severe
genetic load. PLoS One 2:. https://doi.org/10.1371/journal.pone.0000868

Zenni RD, Nuñez MA (2013) The elephant in the room: The role of failed invasions in
understanding invasion biology. Oikos 122:801–815. https://doi.org/10.1111/j.1600-
0706.2012.00254.x

828