# Unraveling genetic load dynamics during biological invasion: insights from two invasive insect species

Eric Lombaert[1], Aurélie Blin[1], Barbara Porro[1,2], Thomas Guillemaud[1], Julio S. Bernal[3], Gary Chang[4], Natalia Kirichenko[5,6,7], Thomas W. Sappington[8], Stefan Toepfer[9,10] and Emeline Deleury[1]

[1] INRAE, Université Côte d'Azur, ISA, Sophia-Antipolis, France
[2] Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Univ Evry, Université Paris-Saclay, Evry, France
[3] Department of Entomology, Texas A&M University, College Station, TX 77843-2475, USA
[4] Biology Department, Gonzaga University, WA, USA
[5] Sukachev Institute of Forest, Siberian Branch of the Russian Academy of Sciences, Federal Research Center "Krasnoyarsk Science Center SB RAS", Akademgorodok 50/28, 660036 Krasnoyarsk, Russia
[6] Institute of Ecology and Geography, Siberian Federal University, Svobodny pr. 79, 660041 Krasnoyarsk, Russia
[7] All-Russian Plant Quarantine Center, Krasnoyarsk branch, Krasnoyarsk, Zhelyabova Str., 6/6, 660020, Russia
[8] USDA-Agricultural Research Service, Corn Insects & Crop Genetics Research Unit, Genetics Laboratory, Iowa State University, Ames, Iowa, USA
[9] CABI Switzerland, c/o Plant Protection and Soil Conservation Directorate, Hodmezovasarhely, Hungary
[10] MARA-CABI Joint Laboratory for Biosafety, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, China

**Corresponding author:**
Eric Lombaert – Institut Sophia Agrobiotech – 400, route des chappes – BP 167 – 06903 Sophia-Antipolis Cedex - France
E-mail: eric.lombaert@inrae.fr
Tel: +33 4 92 38 64 81

**Keywords:** target enrichment, non-model organism, population genomics, pool-sequencing, *Harmonia axyridis*, *Diabrotica virgifera virgifera*, genetic load

**Running title:** Evolution of genetic load in two invasive species

## Abstract

Many invasive species undergo a significant reduction in genetic diversity, i.e. a genetic bottleneck, in the early stages of invasion. However, this reduction does not necessarily prevent them from achieving considerable ecological success and becoming highly efficient colonizers. Here we tested the so-called purge hypothesis, which suggests that demographic bottlenecks may indeed help to purge deleterious mutations, thereby reducing genetic load. We used a transcriptome-based exome capture protocol to identify thousands of SNPs in coding regions of native and invasive populations of two highly successful invasive insect species, the western corn rootworm (Chrysomelidae: *Diabrotica virgifera virgifera*) and the harlequin ladybird (Coccinelidae: *Harmonia axyridis*). We categorized and polarized SNPs to investigate changes in genetic load between invasive populations and their sources. Our results varied between species. In *D. virgifera virgifera*, although there was a general reduction in genetic diversity in invasive populations, including that associated with genetic load, we found no clear evidence for genetic load purging, except marginally for strongly deleterious mutations in one European population. Conversely, in *H. axyridis*, the reduction in genetic diversity was minimal, and we detected signs of genetic load fixation in invasive populations. These findings provide new insights into the evolution of genetic load during invasions, but do not offer a definitive answer to the purge hypothesis. Future research should include larger genomic datasets and a broader range of invasive species to further elucidate these dynamics.

## Introduction

Biological invasions represent a significant aspect of global change, profoundly impacting biodiversity through the alteration of species distributions worldwide, particularly in recent times due to the significant increase in human-assisted dispersal (Seebens et al. 2017). The key factors determining the success of invasive species remain largely hypothetical. While numerous hypotheses have been proposed to explain invasion success (Sherpa and Després 2021; Daly et al. 2023), most fail to simultaneously account for several persistent characteristics of invasions: (i) the rarity of successful invasions resulting from introductions (Williamson and Fitter 1996), (ii) the lag time between initial introduction and invasion (Sakai et al. 2001), (iii) the frequent reduction in genetic diversity due to demographic bottlenecks (Nei et al. 1975) and (iv) the prevalence of multiple invasions originating from an initial invasive population (i.e. the bridgehead effect; Lombaert et al. 2010). These characteristics suggest that the success of invasions may stem from partially stochastic biological processes spanning multiple generations and combining both demographic and genetic mechanisms.

An intriguing hypothesis that aligns with these characteristics is that the genetic load is purged in the initial stages of invasions, i.e. deleterious alleles that cause inbreeding depression are eliminated from the introduced population (Estoup et al. 2016). During the demographic bottleneck induced by the introduction of a limited number of individuals into a new environment, genetic drift intensifies and homozygosity increases. This situation often results in a "mutational meltdown" (Lynch et al. 1995; Simberloff 2009), culminating in failure of the introduced population to establish. Conversely, bottlenecks may also purge deleterious alleles, thereby increasing the mean fitness of the introduced individuals. Theoretically, such purges can occur under specific demographic (i.e., reduction of population size of intermediate intensity) and genetic (i.e., strongly deleterious, recessive alleles) conditions, optimizing exposure to natural selection (Crow 1970; Charlesworth et al. 1990; Glémin 2003; Robinson et al. 2023). Such purging processes within introduced populations are particularly interesting for their potential to facilitate successful invasions by sharply decreasing inbreeding depression so that even inbred individuals may exhibit high fitness levels.

Purging has been demonstrated empirically by measuring the evolution of various traits in artificially bottlenecked populations (Crnokrak and Barrett 2002; Avila et al. 2010). However, in the context of naturally occurring bottlenecks during biological invasions, measuring life history traits can be challenging, and evidence of purging has been documented in only a few invasive species (Parisod et al. 2005; Mullarkey et al. 2013; Fountain et al. 2014; Marchini et al. 2016). One notable formal test of this hypothesis was conducted on the invasive Asian ladybird *Harmonia axyridis*, where measurement of life history traits revealed that invasive populations did not experience the inbreeding depression suffered by native ones, suggesting that deleterious alleles were purged during the invasion process (Facon et al. 2011). Overall, case studies focusing on the evolution of genetic load during biological invasions have either examined a few life history traits (e.g. Facon et al. 2011) or used a single locus approach (e.g. Zayed et al. 2007), making it difficult to generalize the results.

Advances in population genomics over the last decade have provided promising avenues for investigating genetic load on large scales. Initially explored in humans (Lohmueller et al. 2008; Henn et al. 2015), these approaches have been expanded into the fields of domestication (e.g. Schubert et al. 2014; Marsden et al. 2016; Makino et al. 2018; Wang et al. 2021) and conservation (Xue et al. 2015; Robinson et al. 2016; Grossen et al. 2020; Dussex et al. 2021; Ochoa and Gibbs 2021). These studies typically involve SNP-calling within coding

regions, determination of ancestral SNP states, categorization of the severity of fitness losses due to derived alleles, and population comparisons while accounting for genetic drift using synonymous and/or intergenic polymorphisms. A good knowledge of the genome is essential for such studies, which may explain the limited application of these methods to invasive species, which are mostly non-model organisms. However, this limitation is gradually diminishing as progress in genome sequencing and computational techniques make these methods more accessible and cost-effective (Bertorelle et al. 2022).

In this study, we investigated whether deleterious mutations were purged in two successful insect invaders, namely the western corn rootworm, *Diabrotica virgifera virgifera*, and the harlequin ladybird, *Harmonia axyridis*, by directly assessing genetic load using genomic data from feral native and invasive populations. Importantly, the purpose of this study did not include testing the instrumental role of purging in invasion success. We used a pool-seq transcriptome-based exome capture protocol previously developed for non-model species (Deleury et al. 2020) to identify SNPs within coding sequences and categorize them as synonymous, moderately deleterious, or strongly deleterious. Our results offer insights into the fate of the genetic load and provide valuable perspectives on the purge hypothesis on a broader scale.

## Methods

*Species choice*

The two species, *Diabrotica virgifera virgifera* (hereafter called DVV) and *Harmonia axyridis* (HA), are good candidates for testing the genetic load purging hypothesis. Both species are highly successful invaders with extensive invasive ranges (Gray et al. 2009; Roy et al. 2016). Additionally, their invasion routes are well-documented (Ciosi et al. 2008; Lombaert et al. 2014, 2018) and they exhibit bridgehead effects, meaning that certain invasive populations have played a crucial role in global dissemination (Miller et al. 2005; Lombaert et al. 2010). Additionally, genetic load purging was detected in HA through a laboratory study focusing on life-history traits (Facon et al. 2011).

*Design of exome capture probes for target enrichment*

Design of the exome capture probes was performed as described in Deleury et al. (2020). We searched for peptide-coding sequences using FRAMEDP (v1.2.2; Gouzy et al. 2009) on the *de novo* transcriptomes described in Coates et al. (2021) and Vogel et al. (2017) for DVV and HA, respectively. BLASTX results (e-value ≤ 1e$^{-7}$) of transcripts were used against the insect proteomes of *Tribolium castaneum*, *Anoplophora glabripennis*, *Dendroctonus ponderosae*, *Drosophila melanogaster* and the SwissProt database (v2016-02-17) for training. From the obtained coding sequences (CDS), we eliminated (i) *Wolbachia* and other putative endosymbiont sequences, (ii) CDS with > 1% missing nucleotide bases (Ns) or with more than four consecutive Ns, (iii) CDS with a GC% below 25 or above 75 and (iv) CDS with lengths < 400 bp or > 3500 bp. From the remaining 7,132 and 12,739 CDS for DVV and HA respectively, we drew at random c.a. 5.5 Mb for each species.

Probes based on the selected CDS were designed and manufactured by NimbleGen. In the case of DVV, repetitiveness of the probes was assessed based on the highest 15-mer frequency among the genomes of *Tribolium castaneum* (GCA_000002335.3), *Dendroctonus*

*ponderosae* (GCA_000355655.1 and GCA_000346045.2) and DVV (the one available at the
time, GCA_003013835.2). Probes with more than five close matches, defined as no more than
five single-base insertions, deletions or substitutions using the SSAHA algorithm (Ning et al.
2001), in the DVV genome were then discarded. Probes that matched sequences in the
mitochondrial genome were also discarded. Random nucleotides were used to replace
residual Ns in target sequences. For HA, the method used to ensure probe uniqueness is
described in Deleury *et al.* (2020).
Overall, the final probe sets corresponded to a total of 4,151 CDS (5,282,603 bases
across 12,017 regions of overlapping probes) and 5,717 CDS (5,347,461 bases across 6,400
regions of overlapping probes; Deleury et al. 2020) for DVV and HA respectively. This final set
of probes was manufactured in the form of biotinylated DNA oligomers. One capture reaction
contained 2,100,000 overlapping probes of 50 to 99 bp in length (mean length of 73.86 ± 4.46
bp for DVV, and 74.71 ± 4.92 bp for HA).
165
166    *Sample collection*
167
For both species, the choice of populations to be sampled was based on previously known
invasion routes (Figure S1; Miller et al. 2005; Lombaert et al. 2010, 2018), so that each invasive
population could be compared to its source. We sampled adult DVV at 6 sites: two in the
native area (Mexico), two in North America (Colorado and Pennsylvania, respectively
corresponding to the core and front of the first invasive population; Lombaert et al. 2018),
and two in Europe (Hungary, referred to the Central-Southeastern European population, and
north-western Italy, which are independently derived from the same source area of eastern
North America; Miller et al. 2005). Adult HA were sampled at four sites: two in the native area
(Russia (Siberia) and China), and two in North America (Pennsylvania and Washington,
corresponding to two independent outbreaks from the native area; Lombaert et al. 2010). Due
to mixed results regarding the origin of the east North American population of HA in a previous
study using 18 microsatellite markers (Lombaert et al. 2011), we chose to identify its source
through an ABC (approximate Bayesian computation; Beaumont et al. 2002) analysis using
synonymous SNPs from the current study (see Appendix S1 for details).
To determine the ancestral and derived alleles for each SNP (i.e. to polarize the alleles),
outgroup species were also sampled. For DVV, we selected the closely related species
*Diabrotica adelpha*, as well as the more phylogenetically distant *Cerotoma trifurcata* (Eben
and de los Monteros 2013). In the case of HA, from phylogenetically closest to farthest
(Tomaszewska et al. 2021), the sampled outgroup species were *Harmonia yedoensis*,
*Harmonia conformis* and *Harmonia quadripunctata*. Complete information about samples is
provided in Table 1 and Table S1.
189

| Species | Population code name | Status (source) | Sampling site (Lat.; long.) | Sampling date | Haploid sample size |
|---|---|---|---|---|---|
| *Diabrotica virgifera virgifera* | | | | | |
| | D-N-MX1 | Native | Canatlán, Mexico (24.554; -104.741) | Oct. 2015 | 80 |
| | D-N-MX2 | Native | Durango, Mexico (23.999; -104.614) | Oct. 2015 | 80 |
| | D-I-COL | Invasive (D-N-MX1) | Fort Morgan, CO, USA (40.218; -103.869) | Aug. 2015 | 80 |
| | D-I-PEN | Invasive | Landisville, PA, USA | Aug. 2015 | 80 |

| | | | | |
|---|---|---|---|---|
| | (D-I-COL) | (40.119; -76.432) | | |
| D-I-HUN | Invasive (D-I-PEN) | Kondoros, Hungary (46.736; 20.816) | July 2015 | 80 |
| D-I-ITA | Invasive (D-I-PEN) | Cuneo, Italy (44.463; 7.571) | July 2015 | 80 |
| *Harmonia axyridis* | | | | |
| H-N-CHI | Native | Beijing, China (40.057; 116.540) | Oct. 2015 | 72 |
| H-N-RUS | Native | Krasnoyarsk, Russia (55.992; 92.757) | Sept. 2015 | 72 |
| H-I-PEN | Invasive (H-N-CHI) | Landisville, PA, USA (40.119; -76.432) | Aug. 2015 | 72 |
| H-I-WAS | Invasive (H-N-CHI) | Spokane, WA, USA (47.665; -117.403) | Nov. 2015 | 72 |
| *Diabrotica adelpha* | | | | |
| D-OG-ADE | Outgroup DVV | Oaxaca State, Mexico (15.926; 97.151) | Jan. 2017 | 4 |
| *Cerotoma trifurcata* | | | | |
| D-OG-TRI | Outgroup DVV | Weldon, CA, USA (35.660; -118.330) | Oct. 2016 | 4 |
| *Harmonia yedoensis* | | | | |
| H-OG-YED | Outgroup HA | Beijing, China (40.031; 116.265) | Feb. 2015 | 2 |
| *Harmonia conformis* | | | | |
| H-OG-CON | Outgroup HA | Mouans-Sartoux, France (43.620; 6.934) | Oct. 2009 | 2 |
| *Harmonia quadripunctata* | | | | |
| H-OG-QUA | Outgroup HA | Rearing, Valbonne, France (NA) | July 2007 | 2 |

**Table 1:** Description of *Diabrotica virgifera virgifera* (DVV) and *Harmonia axyridis* (HA) samples, including outgroup species (see Table S1 for BioSample accessions). Sources of invasive populations were determined based on known invasion routes (Figure S1; Miller et al. 2005; Lombaert et al. 2010, 2018), except for H-I-PEN for which we retraced invasion routes in this study (see Appendix S1 for details). Because the two native DVV populations are genetically almost indistinguishable, we selected D-N-MX1 as the source of D-I-COL based on the smallest pairwise $F_{ST}$ value between native and invasive populations (see Results section).

*DNA extraction, exome capture and pool sequencing*

For each population, 4 legs from each of 40 individuals (DVV) or 2 legs from each of 36 individuals (HA) were pooled for DNA extraction with the Qiagen DNeasy Blood & Tissue kit, in accordance with the manufacturer's recommendations. For the outgroup species, the same kit was used to extract DNA from a single individual each for the three *Harmonia* species, and from a pool of two individuals each for the *Diabrotica* and *Cerotoma* species.

Genomic libraries were prepared using NimbleGen SeqCap EZ HyperCap Library v2.0 and NimbleGen SeqCap EZ Library v5.0 for DVV and HA, respectively. In brief, for each of the population and outgroup samples, DNA (2 µg in 100 µl) was mechanically sheared to an average size of 200 bp using a Covaris S2 E210 device (6 cycles of 30 seconds each). In the case of DVV, the fragmented DNA was then divided into three technical replicates for each population. Subsequently, the fragments were subjected to end-repair, A-tailing and indexing

(with one unique index per sample) using the KAPA Library Preparation kit designed for
Illumina platforms. Following the ligation of Illumina adapters and indexes, only fragments
falling within the size range of 250 to 450 bp were retained. A PCR amplification step consisting
of 7 cycles was performed using standard Illumina paired-end primers, and the resulting
amplicons were purified using AMPure XP beads (Beckman). The length, quality, and
concentration of the prepared DNA fragments were assessed using a BioAnalyzer with Agilent
High Sensitivity DNA Assay, along with a Qubit.

For each capture (five for DVV and two for HA), we used a total of approximately 1 µg
of amplified DNA for exome enrichment, combining multiple samples in proportions that
allowed for equimolarity between population samples (see Table S2 and Table S3). This
enrichment was performed using the capture probes described above, following the
guidelines of either the NimbleGen SeqCap EZ HyperCap Library Protocol v2.0 or the SeqCap
EZ Library Protocol v5.0. Following each capture, we conducted two parallel PCRs, each
comprising 14 cycles, on the elution solution. In the case of HA, the resulting PCR products
were combined. Subsequently, all PCR products underwent purification using AMPure XP
beads. The length, quality, and concentration of the final DNA fragments were assessed using
a BioAnalyzer equipped with the Agilent High Sensitivity DNA Assay and a Qubit fluorometer.

For sequencing, we used one lane of an Illumina HiSeq3000 sequencer per species,
following the manufacturer's instructions, in paired-end mode for 150 cycles. After
sequencing, the data were demultiplexed and exported as FastQ files, and the libraries were
processed independently. For DVV, the FastQ files of technical replicates were merged prior
to subsequent analysis.

*Mapping, SNP calling and annotation*

Sequence quality assessment was conducted using FastQC v0.11.5 (Andrews 2010).
Subsequently, adapter sequences were removed, and low-quality base pairs were eliminated
using *Trimmomatic* v0.35 (Bolger et al. 2014), with the following parameter settings:
ILLUMINACLIP:TruSeq-file.fa:2:30:10 LEADING:25 TRAILING:25 SLIDINGWINDOW:5:30
MINLEN:75. We used almost the same parameters for the outgroup species sequences, but
with less stringency on quality of reads (SLIDINGWINDOW:5:20).

Filtered reads were mapped onto the genome assemblies PGI_DIABVI_V3a (GenBank
identifier: GCA_917563875.2) and icHarAxyr1.1 (GenBank identifier: GCA_914767665.1), for
DVV and HA respectively, using default options of the *bwa-mem* aligner v0.7.15 (Li 2013). We
used the *SAMtools* v1.15.1 software package (Li et al. 2009) and its *fixmate*, *view*, and
*markdup* tools to perform the following operations sequentially: (1) removal of unmapped
reads and secondary alignments, (2) filtering out read alignments with a mapping quality
Phred-score <20 and improperly paired reads, and (3) identification and removal of PCR
duplicates. Processing filtered reads for the outgroup species followed the same steps,
including mapping to the genome of the corresponding focal species, except that PCR
duplicates were not removed.

For each species, we conducted variant calling on the resulting *bam* alignment files
using Freebayes v1.3.6 software (Garrison and Marth 2012), with a focus on exonic regions
(options *-t* exons.bed *--pooled-continuous --min-alternate-count* 1 *--min-alternate-fraction*
0.001 *--min-alternate-total* 1 *--min-coverage* 80 *--use-best-n-alleles* 3). The resulting *vcf* file
was subsequently filtered using the *view* tool within *bcftools* v1.13 software (Danecek et al.

2021) and an in-house script to retain only true bi-allelic SNPs. Annotation was then
performed with the *SnpEff* program v5.0 (Cingolani et al. 2012b).
259
260  *Allele polarization*
261
SNP positions were extracted from both *vcf* files using the bcftools query tool. Subsequently,
for each outgroup species, we identified the nucleotides (or absence of typing) at positions
matching those in their respective focal species using the *samtools mpileup* tool on the
previously generated *bam* alignment files.
To polarize the SNPs, we used *est-sfs* software v2.03 (Keightley and Jackson 2018) with
the Kimura 2-parameter model. The input files contained data from one native population of
the focal species, together with the corresponding outgroup species, ensuring a consistent
phylogenetic topology for the software. Given that *est-sfs* computes the probability of the
most frequent allele being ancestral solely for polymorphic SNPs within the native population,
we used probabilities from a parallel analysis for monomorphic loci, incorporating an extra
haplotype for each allele. The complete process was repeated twice for each focal species,
considering the availability of two native populations for each. This resulted in two *est-sfs*
probabilities per focal species, both reflecting the likelihood of the most frequent allele being
the ancestral one.
To consider a SNP as polarized, we applied the following rules. If both probabilities
were above 0.5 for the same allele, and at least one of the probabilities exceeded 0.75, we
considered the most frequent allele as the ancestral one. If both probabilities were below 0.5
for the same allele, with at least one below 0.25, and if there were no more than two distinct
nucleotides present in total across the focal and outgroup species combined, we considered
the less frequent allele to be the ancestral one. SNPs that did not meet these criteria were
considered as not polarized but were retained for computations that did not require
polarization.
284
285  *Additional filters and categorization of deleterious mutations*
286
After extracting essential information from the *vcf* file using the *SnpSift* program v5.0
(Cingolani et al. 2012a), we applied additional filters through an in-house R script (R Core Team
2021). First, we selected only biallelic SNPs with coverage greater than 50 reads, falling below
the 95th percentile of overall coverage in each pool, and exhibiting a minor allele frequency
exceeding 0.01 in at least one population. Second, we retained SNPs that were located in
coding regions and had unambiguous annotations. Finally, we excluded SNPs located on the
X-chromosome in the specific case of HA, for which this information was available.
*SnpEff* annotations were used to categorize the severity of fitness loss due to
mutations. Synonymous variants were used as proxies for neutral polymorphism. Non-
synonymous mutations were categorized as either "missense" (considered as potentially
moderately deleterious, involving amino acid changes) or "LoF" (loss-of-function, considered
as potentially highly deleterious, involving gain or loss of stop codons).
299
300  *Genetic diversity and genetic load analyses*
301
302  We computed several descriptive statistics on the whole set of SNPs. At the inter-population
303  level, we assessed genetic differentiation by calculating pairwise $F_{ST}$ values using the R package

*poolfstat* v2.1.1 (Gautier et al. 2022). At the intra-population level, we used in-house R scripts to compute the synonymous expected heterozygosity $H_{eS}$, as a measure of diversity, and the ratio of non-synonymous to synonymous expected heterozygosity $H_{eN}/H_{eS}$, as a measure of the efficacy of selection. We used a 100-block jackknife resampling approach to estimate means and standard errors.

Using polarized SNPs, we reported for each population and each severity category the proportion of positions for which the derived allele is absent (frequency $f = 0$), rare ($f > 0$ and $f \leq 0.1$), common ($f > 0.1$ and $f < 1$), or fixed ($f = 1$). The proportion of rare derived alleles serves as a proxy for the masked load (i.e. the load which may become express in future generations; Bertorelle et al. 2022), whereas common and fixed alleles serve as a proxy for the realized load (i.e. the load which reduces the fitness in the current generation; Bertorelle et al. 2022). Additionally, within each severity category and population, we calculated the mean derived allele frequency and estimated standard errors via a 100-block jackknife resampling approach. Within each species and severity category, mean derived allele frequencies were compared among populations using z-tests with false discovery rate (FDR; Benjamini and Hochberg 1995) correction for multiple comparisons.

Finally, polarized SNPs were used to compute the $R_{XY}$ statistic, which allows assessment of relative excess or deficit of derived alleles within specific categories of deleterious mutations in one population compared to another (Appendix S2; Do et al. 2015; Xue et al. 2015). This statistic effectively corrects for variation due to demography using neutral mutations (synonymous SNPs in our case) as a reference point. In our analyses, population $X$ is an invasive population, and population $Y$ is its source population (provided in Table 1). An $R_{XY}$ value of 1 indicates an equivalent relative genetic load in both populations, whereas values greater or less than 1 suggest an excess (including fixation) or a deficit (including purge) of the genetic load in the invasive population $X$, respectively. We assessed the significance of any deviation from 1 using a z-score two-tailed test based on the 100-block jackknife resampling method.

## Results

*SNP-calling and genomic variation*

In the target enrichment experiments, we obtained mean numbers of raw read sequences of 50,196,834 and 63,360,676 for DVV and HA, respectively (BioProject PRJNA1079689; Table S1). After trimming, 85.0% and 93.8% of the sequences were retained for DVV and HA respectively. For outgroup species, the total number of raw reads was highly variable, with a mean of 2,631,775 of which 97.8% were conserved after trimming (see detailed information for all libraries, including outgroup species, in Table S4). Mapping, calling and filtering identified 66,274 SNPs (of which 62,034 could be polarized) and 169,755 SNPs (of which 169,102 could be polarized) within coding sequences for DVV and HA respectively (see Table S5 for details).

In the case of DVV, pairwise $F_{ST}$ computed on the full set of SNPs showed patterns very close to those expected from results of previous analyses of microsatellite datasets (Figure S2). The differentiation between the two native Mexican populations was almost negligible with a mean $F_{ST}$ estimate below 0.001. Given that the $F_{ST}$ estimates between the invasive populations and the native sample D-N-MX1 were consistently lower (mean $F_{ST}$ of 0.121) than those with the sample D-N-MX2 (mean $F_{ST}$ of 0.128), D-N-MX1 was therefore considered the

source of the D-I-COL sample (see Table 1). The Pennsylvania and Colorado populations had a
low pairwise $F_{ST}$ of 0.012, consistent with an Eastern expansion from Colorado (Figure S1),
with plausible continuous gene flow. Conversely, all other $F_{ST}$ estimates were relatively high,
particularly between the native and European populations (mean $F_{ST}$ value = 0.152). The
highest value was observed between the two European populations at 0.169, confirming
previous results indicating that these populations originated from two independent
introductions (Miller et al. 2005).
In the case of HA, $F_{ST}$ estimates were moderate and fairly homogeneous, ranging from
0.020 to 0.050 (Figure S2). The smallest values were found between the eastern native
population H-N-CHI and both invasive populations, while the $F_{ST}$ between invasive populations
was slightly larger, consistent with previous results indicating independent origins for the two
North American outbreaks (Lombaert et al. 2010). The highest values were observed between
the western native population H-N-RUS and all other populations.
Within each species, invasive populations were characterized by reduced synonymous
heterozygosity compared to their native counterparts, reflecting demographic bottlenecks
encountered during the invasion process (Figure 1, x axis). One exception was the western
native population of HA (H-N-RUS), which stood out for having the lowest diversity within this
species. The ratio of non-synonymous to synonymous expected heterozygosity $H_{eN}/H_{eS}$ is
typically expected to increase in populations experiencing pronounced drift, as the reduced
efficacy of selection allows more non-synonymous mutations, including potentially
deleterious ones, to persist. In line with this expectation, we observed a higher ratio in invasive
populations (Figure 1, y axis), and a negative correlation between this ratio and synonymous
expected heterozygosity within each species (Pearson's $r$ = -0.95, $P < 10^{-2}$ for DVV; Pearson's
$r$ = -0.90, $P < 10^{-1}$ for HA; Figure 1). Overall, the differences between native and invasive
populations were more pronounced in DVV, characterized by lower diversities and reduced
efficacy of selection in the invasive ranges.
377

**Figure 1:** Ratio of non-synonymous to synonymous expected heterozygosity $H_{eN}/H_{eS}$ vs. the synonymous expected heterozygosity $H_{eS}$ of each *Diabrotica virgifera virgifera* (DVV) and *Harmonia axyridis* (HA) population. Means and standard errors (error bars) were determined from 100-block jackknife resampling. See Table 1 for population codes.

*Genetic load on polarized SNPs*

In all populations studied for each species, derived alleles were mostly rare (with frequencies below 0.1) or absent (Figure 2). The highest derived allele frequencies consistently showed a higher prevalence in synonymous positions compared to non-synonymous positions. In DVV, we observed a substantial loss of non-synonymous derived alleles in invasive populations compared to native ones. Although this was characterized by a reduction in both masked load (proportion of non-synonymous SNPs with rare derived alleles ranging from 0.496 to 0.523 in native populations and from 0.145 to 0.188 in invasive populations) and realized load (proportion of non-synonymous SNPs with frequent or fixed derived alleles ranging from 0.075 to 0.077 in native populations and from 0.046 to 0.051 in invasive populations), it also came with slightly higher proportions of fixed derived alleles. This trend was less pronounced for mutations inferred to be strongly deleterious (LoF) than for those inferred to be moderately deleterious (missense). For HA, the disparities between native and invasive populations were considerably less pronounced (Figure 2). Compared to their native Chinese source, HA invasive populations displayed a slight reduction in masked load (proportion of non-synonymous SNPs with rare derived alleles equal to 0.455 in the Chinese population and from 0.347 to 0.381 in invasive populations) and a slight increase in realized load (proportion of non-synonymous SNPs with frequent or fixed derived alleles equal to 0.019 in the Chinese population and from 0.024 to 0.027 in invasive populations). Again, this trend was less pronounced for strongly deleterious mutations (Figure 2).

**Figure 2**: Proportion of derived alleles found in each population of *Diabrotica virgifera virgifera* (DVV) and *Harmonia axyridis* (HA) within categories of putative severity, classified by their frequency. See Table 1 for population codes.

413   As expected, the mean derived allele frequencies within populations decline with
414 increasing putative severity in both species (Figure 3). For DVV, synonymous and missense
415 mean derived allele frequencies were consistently significantly lower in the invasive
416 populations than in the native populations ($P < 10^{-4}$ for all comparisons after FDR correction),
417 whereas no significant differences were observed between native populations or between
418 invasive populations ($P > 0.3$ for all comparisons after FDR correction). Within each of the
419 three severity categories, the mean derived allele frequencies were virtually identical in the
420 four HA populations ($P > 0.02$ for all comparisons after FDR correction). No differences in LoF
421 derived allele frequencies were significant ($P > 0.2$ for all comparisons after FDR correction),
422 for either species. Overall, derived allele frequencies were lower in HA than in DVV.

**Figure 3**: Mean derived allele frequencies in all *Diabrotica virgifera virgifera* (DVV) and *Harmonia axyridis* (HA) populations at all three severity categories. Means and standard errors (error bars) were determined from 100-block jackknife resampling. See Table 1 for population codes.

431   Finally, the $R_{XY}$ ratio revealed no significant differences in relative frequencies of
432 missense derived alleles between invasive populations and their respective sources in DVV
433 (Figure 4). For putatively highly deleterious loss-of-function alleles, $R_{XY}$ values sometimes
434 deviate sharply from 1, but the number of SNPs is small (Table S5), and most differences are
435 not statistically significant. The only exception is found in the population from Hungary (D-I-
436 HUN), which shows reduced relative frequencies within the putatively highly deleterious
437 category (LoF) of alleles (Figure 4). Regarding HA, slightly higher loads were significant in three
438 out of four comparisons across severity categories between the two invasive populations (H-
439 I-PEN and H-I-WAS) and the native Chinese population (Figure 4).

**Figure 4**: Relative mutation load estimated as the $R_{XY}$ ratio of derived alleles for moderate and strong severity categories in populations of *Diabrotica virgifera virgifera* (D) and *Harmonia axyridis* (H). A ratio below or above 1 indicates, respectively, a relative frequency deficit or excess of derived alleles of indicated severity category in the invasive population (listed first) compared to its population source (in parentheses). Means and standard errors (error bars) were determined from 100-block jackknife resampling. A star indicates that the corresponding $R_{XY}$ value is significantly different from 1 (z-score two-tailed test). For population codes, refer to Table 1.

## Discussion

Exploring the evolution of genetic load using genomic data has long been restricted to model species. Here, we successfully identified putatively deleterious derived alleles within coding regions in two non-model invasive insect species using a pool-seq transcriptome-based exome capture protocol (Deleury et al. 2020). By comparing native and invasive populations, we were thereby able to test the hypothesis that the genetic load may have been purged during the invasion process, which may enhance invasion success of certain populations (Sherpa and Després 2021; Daly et al. 2023). Given that bottlenecks amplify genetic drift and weaken selection effectiveness (Crow 1970; Glémin et al. 2003), the likelihood of purging is expected to increase for recessive, strongly deleterious mutations, while moderately deleterious mutations may still experience some degree of drift, leading to higher frequencies (Agrawal and Whitlock 2011). Our study exploring the fate of the genetic load in DVV and HA yielded mixed results, underscoring the nuanced interplay of evolutionary forces in invasive species.

*Evolution of the genetic load in* Diabrotica virgifera virgifera

DVV exhibits marked genetic differences between native and invasive populations. Consistent with previous studies (Ciosi et al. 2008; Lombaert et al. 2018), invasive populations, including the long-established population in Colorado, display sharp declines in genetic diversity. Compared to their native counterparts, all invasive populations generally exhibit lower

frequencies of derived alleles, regardless of their potential fitness impacts, thus confirming
the substantial role of genetic drift in the invasion process of the species. While this overall
reduction in genetic load within invasive populations is apparent, it is worth emphasizing the
presence of a slightly larger fixed load.

In North America, despite a strong initial loss of genetic diversity, no significant relative
deficit or excess of deleterious mutations was detected in invasive populations compared to
their source. This pattern includes the absence of any quantifiable evolution of genetic load
during DVV's geographic expansion eastward to Pennsylvania, differing from expectations
associated with the concept of "expansion load". Under this concept, rare deleterious alleles
may "surf" along the expansion front and ultimately reach high frequencies (Klopfstein et al.
2006; Travis et al. 2007; Peischl and Excoffier 2015).

Interestingly, the oldest European outbreak in Central-Southeastern Europe (here a
sample from Hungary) exhibited a significant deficit of "loss-of-function" deleterious
mutations compared to its source population, while "missense" deleterious mutations did not
differ significantly. This suggests that the demographic and selective constraints in this
population were effective at purging highly deleterious mutations but perhaps not moderately
deleterious ones. Such differences in evolutionary trajectories according to mutation severity
are theoretically expected (Whitlock 2002; Caballero et al. 2017; Dussex et al. 2023) and have
been observed in other species experiencing significant bottlenecks (Xue et al. 2015; Khan et
al. 2021; Ochoa and Gibbs 2021; Wang et al. 2023).

Finally, the invasive Italian population, despite originating from an independent
introduction (Miller et al. 2005), showed patterns similar to that of the Central- Southeastern
European population. However, it did not exhibit significant signals of purging or fixation of
alleles that contribute to genetic load, regardless of the mutation severity considered.

*Evolution of the genetic load in* Harmonia axyridis

In HA, the overall loss of genetic diversity in both invasive populations is relatively low, in
agreement with previous findings (Lombaert et al. 2010, 2011). While the putative masked
genetic load has decreased slightly, the putative realized load, although consistently low in all
populations, showed no sign of decreasing and even exhibits a slight increase. This observation
aligns with a trend towards a relative excess of deleterious mutations in invasive populations
compared to their native source, regardless of the putative fitness impact. This tendency to
fix alleles contributing to genetic load, although subtle, is significant across all populations and
severity categories, except for the strongly deleterious mutations in Western North America
(sampled in Washington).

These findings diverge from a previous study based on life history traits, which
demonstrated that invasive populations of HA experienced minimal inbreeding depression
while maintaining fitness comparable to outbred native populations (Facon et al. 2011). That
study has greatly contributed to the hypothesis of purging genetic load as a key mechanism
driving successful biological invasions (Estoup et al. 2016; Sherpa and Després 2021; Daly et
al. 2023). Given the markedly different methodologies used in the study by Facon et al. (2011)
and our study, it is challenging to ascertain why our results do not align. It is conceivable that
a few key genes could play decisive roles in invasion success, an aspect difficult to quantify
using our approach, particularly in the context of an overall low genetic load, as previously
described in this species (Tayeh et al. 2013). This issue is further complicated by the imperfect
nature of our categorization, where missense mutations likely cover a broad spectrum of

severity levels, potentially obscuring the influence of these key genes. Finally, our exome capture protocol only targets a fraction of the exome, approximately one-third, and a broader approach may yield qualitatively different results.

*Conclusions and perspectives*

We tested the hypothesis of the purge of the genetic load during the invasion of two insect species, a crop pest and a predator. At first glance, our results offer a nuanced perspective. In the case of DVV, we were unable to detect any significant evolution of the genetic load, except for the purge of strongly deleterious mutations in the invasive Central- Southeastern European population. For HA, we observed subtle evidence suggesting a tendency toward fixation, rather than toward purge, of the genetic load. Although these results are not entirely contradictory, they may stem from differences in initial genetic composition, varying demographic history, and divergent ecological niches.

While population genomics approaches to assess the evolution of genetic load have become increasingly popular in conservation and domestication biology (Moyers et al. 2018; Bertorelle et al. 2022), our study is one of the first in the context of biological invasions. However, it does not yet provide a comprehensive and definitive answer to the question addressed. To broaden our results and enhance our statistical power, particularly for strongly deleterious mutations, it would be essential to consider the entire exome rather than only a fraction. Additionally, categorizing mutation severity by quantifying evolutionary constraints within a phylogenetic framework could enhance our understanding of genetic load evolution, including in non-coding regions (Davydov et al. 2010). Furthermore, extending this research to a wider range of species would enhance our ability to identify potential patterns in the evolution of the genetic load during biological invasions. Finally, to test not only the occurrence of purging in invasions but also its role in the success of invasions, we must compare the load dynamics in introductions that fail with those that succeed – a comparison that is rarely feasible (Zenni and Nuñez 2013), except in controlled invasions such as in classical biological control (Fauvergue et al. 2012).

# Acknowledgments

We thank our colleagues Pamela Bruno, Charlyne Martignier-Jaccard, Pascal Maignet and Wang Su for *Diabrotica adelpha*, *Harmonia quadripunctata* and *Harmonia yedoensis* samples. We also thank Emmanuelle Murciano-Germain for administrative assistance. Sequencing was performed at the GENOTOUL GeT platform (https://get.genotoul.fr/en/).

# Funding

This work was funded by grants from ANR project GENLOADICS, and INRAe SPE department. Sampling in Russia was supported by the basic project of the Sukachev Institute of Forest SB RAS (no. FWES-2024-0029).

# Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the content of this article. TG and EL are both recommenders at PCI Evolutionary Biology and PCI Ecology. TG is co-founder of Peer Community In.

## Data availability

Target capture pool sequencing files and all sequencing data have been deposited in Sequence Read Archive, National Center for Biotechnology Information, under project PRJNA1079689 (see also Table S1): https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1079689. All scripts used in this study have been deposited at Data INRAE: https://doi.org/10.57745/ESQFDB

## Author contributions

EL and ED designed the study. JB, GC, NK, TS and ST managed the collection of samples. AB and ED prepared the samples and the libraries. ED and BP developed the bioinformatics pipelines. EL and ED analyzed the data. EL and TG wrote the paper. All authors have revised and approved the final manuscript.

## References

Agrawal AF, Whitlock MC (2011) Inferences about the distribution of dominance drawn from yeast gene knockout data. Genetics 187:553–566. https://doi.org/10.1534/genetics.110.124560

Andrews S (2010) FastQC: a quality control tool for high throughput sequence data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Avila V, Amador C, Garcia-Dorado A (2010) The purge of genetic load through restricted panmixia in a Drosophila experiment. J Evol Biol 23:1937–1946. https://doi.org/10.1111/j.1420-9101.2010.02058.x

Beaumont MA, Zhang WY, Balding DJ (2002) Approximate Bayesian computation in population genetics. Genetics 162:2025–2035

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate - a practical and powerful approach to multiple testing. J R Stat Soc Ser B-Methodological 57:289–300

Bertorelle G, Raffini F, Bosse M, et al (2022) Genetic load: genomic estimates and applications in non-model animals. Nat Rev Genet 23:492–503. https://doi.org/10.1038/s41576-022-00448-x

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: A flexible trimmer for Illumina sequence data. Bioinformatics 30:2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Caballero A, Bravo I, Wang J (2017) Inbreeding load and purging: implications for the short-term survival and the conservation management of small populations. Heredity (Edinb) 1–9. https://doi.org/10.1038/hdy.2016.80

Charlesworth D, Morgan MT, Charlesworth B (1990) Inbreeding Depression, Genetic Load, and the Evolution of Outcrossing Rates in a Multilocus System With No Linkage. Evolution (N Y) 44:1469–1489. https://doi.org/10.1111/j.1558-5646.1990.tb03839.x

Cingolani P, Patel VM, Coon M, et al (2012a) Using *Drosophila melanogaster* as a model for genotoxic chemical mutational studies with a new program, SnpSift. Front Genet 3:1–9. https://doi.org/10.3389/fgene.2012.00035

Cingolani P, Platts A, Wang LL, et al (2012b) A program for annotating and predicting the
effects of single nucleotide polymorphisms, SnpEff. Fly (Austin) 6:80–92.
https://doi.org/10.4161/fly.19695

Ciosi M, Miller NJ, Kim KS, et al (2008) Invasion of Europe by the western corn rootworm,
*Diabrotica virgifera virgifera*: multiple transatlantic introductions with various
reductions of genetic diversity. Mol Ecol 17:3614–3627. https://doi.org/10.1111/j.1365-
294X.2008.03866.x

Coates BS, Deleury E, Gassmann AJ, et al (2021) Up-regulation of apoptotic- and cell survival-
related gene pathways following exposures of western corn rootworm to *B.*
*thuringiensis* crystalline pesticidal proteins in transgenic maize roots. BMC Genomics
22:1–27. https://doi.org/10.1186/s12864-021-07932-4

Crnokrak P, Barrett SCH (2002) Purging the genetic load: A review of the experimental
evidence. Evolution (N Y) 56:2347–2358. https://doi.org/10.1554/0014-
3820(2002)056[2347:pptgla]2.0.co;2

Crow JF (1970) Genetic loads and the cost of natural selection. In: Krickeberg K, Lewontin RC,
Neyman J, Schreiber M (eds) Mathematical topics in population genetics. Springer
Berlin Heidelberg, Berlin, pp 128–177

Daly EZ, Chabrerie O, Massol F, et al (2023) A synthesis of biological invasion hypotheses
associated with the introduction–naturalisation–invasion continuum. Oikos e09645:1–
29. https://doi.org/10.1111/oik.09645

Danecek P, Bonfield JK, Liddle J, et al (2021) Twelve years of SAMtools and BCFtools.
Gigascience 10:1–4. https://doi.org/10.1093/gigascience/giab008

Davydov E V., Goode DL, Sirota M, et al (2010) Identifying a high fraction of the human
genome to be under selective constraint using GERP++. PLoS Comput Biol 6:1–13.
https://doi.org/10.1371/journal.pcbi.1001025

Deleury E, Guillemaud T, Blin A, Lombaert E (2020) An evaluation of pool-sequencing
transcriptome-based exon capture for population genomics in non-model species.
bioRxiv 10.1101/58:v7 peer-reviewed and recommended by PCI Genomics.
https://doi.org/10.1101/583534

Do R, Balick D, Li H, et al (2015) No evidence that selection has been less effective at
removing deleterious mutations in Europeans than in Africans. Nat Genet 47:126–131.
https://doi.org/10.1038/ng.3186

Dussex N, Morales HE, Grossen C, et al (2023) Purging and accumulation of genetic load in
conservation. Trends Ecol Evol 38:961–969. https://doi.org/10.1016/j.tree.2023.05.008

Dussex N, van der Valk T, Morales HE, et al (2021) Population genomics of the critically
endangered kākāpō. Cell Genomics 100002:1–10.
https://doi.org/10.1016/j.xgen.2021.100002

Eben A, de los Monteros AE (2013) Tempo and mode of evolutionary radiation in
Diabroticina beetles (genera *Acalymma*, *Cerotoma*, and *Diabrotica*). Zookeys 332:207–
231. https://doi.org/10.3897/zookeys.332.5220

Estoup A, Ravigné V, Hufbauer R, et al (2016) Is There A Genetic Paradox of Biological
Invasion? Annu Rev Ecol Evol Syst 47:51–72. https://doi.org/10.1146/annurev-ecolsys-
121415

Facon B, Hufbauer RA, Tayeh A, et al (2011) Inbreeding depression is purged in the invasive
insect *Harmonia axyridis*. Curr Biol 21:424–427.
https://doi.org/10.1016/j.cub.2011.01.068

Fauvergue X, Vercken E, Malausa T, Hufbauer RA (2012) The biology of small, introduced

populations, with special reference to biological control. Evol Appl 5:424–443.
https://doi.org/10.1111/j.1752-4571.2012.00272.x

Fountain T, Duvaux L, Horsburgh G, et al (2014) Human-facilitated metapopulation dynamics
in an emerging pest species, *Cimex lectularius*. Mol Ecol 23:1071–1084.
https://doi.org/10.1111/mec.12673

Garrison E, Marth G (2012) Haplotype-based variant detection from short-read sequencing.
arXiv 1207.3907:1–9. https://doi.org/10.48550/arXiv.1207.3907

Gautier M, Vitalis R, Flori L, Estoup A (2022) f-Statistics estimation and admixture graph
construction with Pool-Seq or allele count data using the R package poolfstat. Mol Ecol
Resour 22:1394–1416. https://doi.org/10.1111/1755-0998.13557

Glémin S (2003) How are deleterious mutations purged? Drift versus nonrandom mating.
Evolution (N Y) 57:2678–2687. https://doi.org/10.1111/j.0014-3820.2003.tb01512.x

Glémin S, Ronfort J, Bataillon T (2003) Patterns of Inbreeding Depression and Architecture of
the Load in Subdivided Populations. Genetics 165:2193–2212.
https://doi.org/10.1093/genetics/165.4.2193

Gouzy J, Carrere S, Schiex T (2009) FrameDP: Sensitive peptide detection on noisy matured
sequences. Bioinformatics 25:670–671. https://doi.org/10.1093/bioinformatics/btp024

Gray ME, Sappington TW, Miller NJ, et al (2009) Adaptation and Invasiveness of Western
Corn Rootworm: Intensifying Research on a Worsening Pest. Annu Rev Entomol 54:303–
321

Grossen C, Guillaume F, Keller LF, Croll D (2020) Purging of highly deleterious mutations
through severe bottlenecks in Alpine ibex. Nat Commun 11:1001.
https://doi.org/10.1038/s41467-020-14803-1

Henn BM, Botigué LR, Bustamante CD, et al (2015) Estimating the mutation load in human
genomes. Nat Rev Genet 16:333–343. https://doi.org/10.1038/nrg3931

Keightley PD, Jackson BC (2018) Inferring the Probability of the Derived vs. the Ancestral
Allelic State at a Polymorphic Site. Genetics 209:897–906. https://doi.org/10.1007/978-
1-59745-494-0_7

Khan A, Patel K, Shukla H, et al (2021) Genomic evidence for inbreeding depression and
purging of deleterious genetic variation in Indian tigers. Proc Natl Acad Sci
118:e2023018118. https://doi.org/10.1073/pnas.2023018118

Klopfstein S, Currat M, Excoffier L (2006) The fate of mutations surfing on the wave of a
range expansion. Mol Biol Evol 23:482–490. https://doi.org/10.1093/molbev/msj057

Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv 1303.3997:1–3. https://doi.org/10.48550/arXiv.1303.3997

Li H, Handsaker B, Wysoker A, et al (2009) The Sequence Alignment/Map format and
SAMtools. Bioinformatics 25:2078–2079.
https://doi.org/10.1093/bioinformatics/btp352

Lohmueller KE, Indap AR, Schmidt S, et al (2008) Proportionally more deleterious genetic
variation in European than in African populations. Nature 451:994–997.
https://doi.org/10.1038/nature06611

Lombaert E, Ciosi M, Miller NJ, et al (2018) Colonization history of the western corn
rootworm (*Diabrotica virgifera virgifera*) in North America: insights from random forest
ABC using microsatellite data. Biol Invasions 20:665–677.
https://doi.org/https://doi.org/10.1007/s10530-017-1566-2

Lombaert E, Guillemaud T, Cornuet JM, et al (2010) Bridgehead effect in the worldwide
invasion of the biocontrol harlequin ladybird. PLoS One 5:e9743. https://doi.org/e9743

10.1371/journal.pone.0009743

Lombaert E, Guillemaud T, Lundgren J, et al (2014) Complementarity of statistical treatments
to reconstruct worldwide routes of invasion: the case of the Asian ladybird *Harmonia*
*axyridis*. Mol Ecol 23:5979–5997. https://doi.org/10.1111/mec.12989

Lombaert E, Guillemaud T, Thomas CE, et al (2011) Inferring the origin of populations
introduced from a genetically structured native range by approximate Bayesian
computation: case study of the invasive ladybird *Harmonia axyridis*. Mol Ecol 20:4654–
4670. https://doi.org/10.1111/j.1365-294X.2011.05322.x

Lynch M, Conery IJ, Burger R (1995) Mutation Accumulation and the Extinction of Small
Populations. Am Nat 146:489–518. https://doi.org/10.1086/285812

Makino T, Rubin C-J, Carneiro M, et al (2018) Elevated proportions of deleterious genetic
variation in domestic animals and plants. Genome Biol Evol 10:276–290.
https://doi.org/10.1093/gbe/evy004/4794728

Marchini GL, Sherlock NC, Ramakrishnan AP, et al (2016) Rapid purging of genetic load in a
metapopulation and consequences for range expansion in an invasive plant. Biol
Invasions 18:183–196. https://doi.org/10.1007/s10530-015-1001-5

Marsden CD, Ortega-Del Vecchyo D, O'Brien DP, et al (2016) Bottlenecks and selective
sweeps during domestication have increased deleterious genetic variation in dogs. Proc
Natl Acad Sci 113:152–157. https://doi.org/10.1073/pnas.1512501113

Miller N, Estoup A, Toepfer S, et al (2005) Multiple transatlantic introductions of the western
corn rootworm. Science (80- ) 310:992. https://doi.org/10.1126/science.1115871

Moyers BT, Morrell PL, McKay JK (2018) Genetic costs of domestication and improvement. J
Hered 109:103–116. https://doi.org/10.1093/jhered/esx069

Mullarkey A a., Byers DL, Anderson RC (2013) Inbreeding depression and partitioning of
genetic load in the invasive biennial *Alliaria petiolata* (Brassicaceae). Am J Bot 100:509–
518. https://doi.org/10.3732/ajb.1200403

Nei M, Maruyama T, Chakraborty R (1975) Bottleneck Effect and Genetic Variability in
Populations. Evolution (N Y) 29:1–10. https://doi.org/10.2307/2407137

Ning Z, Cox AJ, Mullikin JC (2001) SSAHA: a fast search method for large DNA databases. -
Abstract - UK PubMed Central. 1725–1729. https://doi.org/10.1101/gr.194201.1

Ochoa A, Gibbs HL (2021) Genomic signatures of inbreeding and mutation load in a
threatened rattlesnake. Mol Ecol 30:5454–5469. https://doi.org/10.1111/mec.16147

Parisod C, Trippi C, Galland N (2005) Genetic variability and founder effect in the pitcher
plant *Sarracenia purpurea* (Sarraceniaceae) in populations introduced into Switzerland:
From inbreeding to invasion. Ann Bot 95:277–286. https://doi.org/10.1093/aob/mci023

Peischl S, Excoffier L (2015) Expansion load: Recessive mutations and the role of standing
genetic variation. Mol Ecol 24:2084–2094. https://doi.org/10.1111/mec.13154

R Core Team (2021) R: A language and environment for statistical computing

Robinson J, Kyriazis CC, Yuan SC, Lohmueller KE (2023) Deleterious Variation in Natural
Populations and Implications for Conservation Genetics. Annu Rev Anim Biosci 11:93–
114. https://doi.org/10.1146/annurev-animal-080522-093311

Robinson JA, Ortega-Del Vecchyo D, Fan Z, et al (2016) Genomic Flatlining in the Endangered
Island Fox. Curr Biol 26:1183–1189. https://doi.org/10.1016/j.cub.2016.02.062

Roy HE, Brown PMJ, Adriaens T, et al (2016) The harlequin ladybird, *Harmonia axyridis*:
global perspectives on invasion history and ecology. Biol Invasions 18:997–1044.
https://doi.org/10.1007/s10530-016-1077-6

Sakai AK, Allendorf FW, Holt JS, et al (2001) The population biology of invasive species. Annu

Rev Ecol Syst 32:305–332. https://doi.org/10.1146/annurev.ecolsys.32.081501.114037

Schubert M, Jónsson H, Chang D, et al (2014) Prehistoric genomes reveal the genetic
foundation and cost of horse domestication. Proc Natl Acad Sci U S A 111:E5661-9.
https://doi.org/10.1073/pnas.1416991111

Seebens H, Blackburn TM, Dyer E, et al (2017) No saturation in the accumulation of alien
species worldwide. Nat Commun 8:14435. https://doi.org/10.1038/ncomms14435

Sherpa S, Després L (2021) The evolutionary dynamics of biological invasions: A multi-
approach perspective. Evol Appl 1–23. https://doi.org/10.1111/eva.13215

Simberloff D (2009) The Role of Propagule Pressure in Biological Invasions. Annu Rev Ecol
Evol Syst 40:81–102

Tayeh A, Estoup A, Hufbauer RA, et al (2013) Investigating the genetic load of an emblematic
invasive species: the case of the invasive harlequin ladybird *Harmonia axyridis*. Ecol Evol
3:864–871. https://doi.org/10.1002/ece3.490

Tomaszewska W, Escalona HE, Hartley D, et al (2021) Phylogeny of true ladybird beetles
(Coccinellidae: Coccinellini) reveals pervasive convergent evolution and a rapid
Cenozoic radiation. Syst Entomol 46:611–631. https://doi.org/10.1111/syen.12479

Travis JMJ, Munkemuller T, Burton OJ, et al (2007) Deleterious mutations can surf to high
densities on the wave front of an expanding population. Mol Biol Evol 24:2334–2343.
https://doi.org/10.1093/molbev/msm167

Vogel H, Schmidtberg H, Vilcinskas A (2017) Comparative transcriptomics in three ladybird
species supports a role for immunity in invasion biology. Dev Comp Immunol 67:452–
456. https://doi.org/10.1016/j.dci.2016.09.015

Wang MS, Zhang JJ, Guo X, et al (2021) Large-scale genomic analysis reveals the genetic cost
of chicken domestication. BMC Biol 19:1–16. https://doi.org/10.1186/s12915-021-
01052-x

Wang X, Peischl S, Heckel G (2023) Demographic history and genomic consequences of
10,000 generations of isolation in a wild mammal. Curr Biol 33:2051-2062.e4.
https://doi.org/10.1016/j.cub.2023.04.042

Whitlock MC (2002) Selection, load and inbreeding depression in a large metapopulation.
Genetics 160:1191–1202. https://doi.org/10.1093/genetics/160.3.1191

Williamson M, Fitter A (1996) The varying success of invaders. Ecology 77:1661–1666

Xue Y, Prado-martinez J, Sudmant PH, et al (2015) Mountain gorilla genomes reveal the
impact of long-term population decline and inbreeding. Science (80- ) 348:242–245.
https://doi.org/10.1126/science.aaa4484

Zayed A, Constantin S a., Packer L (2007) Successful biological invasion despite a severe
genetic load. PLoS One 2:. https://doi.org/10.1371/journal.pone.0000868

Zenni RD, Nuñez MA (2013) The elephant in the room: The role of failed invasions in
understanding invasion biology. Oikos 122:801–815. https://doi.org/10.1111/j.1600-
0706.2012.00254.x

791