# HLA-E and NKG2A Mediate Resistance to *M. bovis* BCG Immunotherapy in Non-Muscle-Invasive Bladder Cancer

D. Ranti[1,2,3,4,5*], H. Yu[1,2,3,4,5*], Y.A. Wang[1,2,3,4,5], C. Bieber[1,2,3,4,5], T. Strandgaard[6,7], B. Salomé[1,2,3,4], Sean Houghton[8], J. Kim[8], H. Ravichandran[8], I. Okulate[1,2,3,4,5], E. Merritt[3,9], S. Bang[9], A. Demetriou[1,2,3,4,5], Z. Li[1,2,3,4], S. V. Lindskrog[6,7], D.F. Ruan[1,2,3,4], J. Daza[1,2,3,4,5], R. Rai[4], E. Hegewisch-Solloa[10], E.M. Mace[10], R., Fernandez-Rodriguez[2,4], S. Izadmehr[3,9], G. Doherty[11,12], A. Narasimhan[11,12], A.M. Farkas[3,4,11], P. Cruz-Encarnacion[1,2], S. Shroff[7,13], F. Patel[3,14], M. Tran[3,4,11], S.J. Park[1,2,3,4], J. Qi[15], M. Patel[15], D. Geanon[15], G. Kelly[15], R.M.de Real[15], B. Lee[15], K. Nie[15], S. Miake-Iye[15], K. Angeliadis[15], E. Radkevich[15], T.H. Thin[16], M. Garcia-Barros[16], H. Brown[16], B. Martin[16], A. Mateo[16], A. Soto[16], R. Sussman[16], S. Shiwlani[16], S. Francisco-Simon[16], K.G. Beaumont[9,13], Y. Hu[15], Y-C. Wang[7,13], L. Wang[9], R.P. Sebra[9,13,17], S. Smith[18], M. Skobe[2,4], E. Clancy-Thompson[19], D. Palmer[19], S. Hammond[19], B. D. Hopkins[15], P. Wiklund[5], J. Zhu[9,13], J.J. Bravo-Cordero[11,12], R. Brody[16], Hopkins, B., Z. Chen[1,3,15], S. Kim-Schulze[1,3,15], L. Dyrskjøt[6,7], O. Elemento[8], A. Tocheva[3,9], W-M. Song[9], N. Bhardwaj[3,4,11], M.D. Galsky[4,11], J.P. Sfakianos[5#] & A. Horowitz[1,2,3,4#]

[1]Department of Immunology and Immunotherapy, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[2]Department of Oncological Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[3]The Marc and Jennifer Lipschultz Precision Immunology Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[4]Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[5]Department of Urology, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[6]Department of Molecular Medicine, Aarhus University Hospital, Aarhus, Denmark
[7]Department of Clinical Medicine, Aarhus University, Aarhus, Denmark
[8]Caryl and Israel Englander Institute for Precision Medicine, Weill Cornell Medicine, New York, NY, USA
[9]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[10]Department of Pediatrics, Vagelos College of Physicians and Surgeons, Columbia University, New York NY, USA
[11]Division of Hematology and Medical Oncology, Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[12]Microscopy and Advanced Bioimaging Core, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[13]Icahn Institute for Data Science and Genomics Technology, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[14]Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[15]Human Immune Monitoring Center, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[16]Department of Pathology, Molecular and Cell-Based Medicine, Icahn School of Medicine at Mount Sinai, New York, NY
[17]Black Family Stem Cell Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[18]Center for Inflammation research and Translational Medicine, Brunel University London, London, UK
[19]AstraZeneca, Oncology R & D Unit, Gaithersburg, Maryland, USA
[20]Department of Physiology and Biophysics, Weill Cornell Medicine, New York, NY, USA
* These authors contributed equally
# Co-Senior author

**Running title.** HLA-E and NKG2A in BCG Resistance

49    **Keywords.** Non-muscle-invasive bladder cancer; BCG-unresponsive; NK cells;
50    Immunotherapy.
51
52

**Corresponding Author:**
Amir Horowitz, PhD
Associate Professor of Immunology & Immunotherapy and Oncological Sciences
Lipschultz Precision Immunology Institute and Tisch Cancer Institute
Icahn School of Medicine at Mount Sinai
1425 Madison Ave., Box 1630
New York, New York 10029
Tel: (212) 659-9391 | Fax: (646) 537-9577
Email: amir.horowitz@mssm.edu
62
John P. Sfakianos, MD
Associate Professor of Urology
Icahn School of Medicine at Mount Sinai
1425 Madison Ave., L6-58, Box 1272
New York, New York 10029
Tel: (212) 659-9375
Email: john.sfakianos@mountsinai.org
70
71
72

73    **Conflict of Interest Disclosure Statement:** The authors declare no potential conflicts of interest.
74
75    **Word Count (excluding methods and references):**
76    **Fig. Count: 7**
77    **Table Count: 1**
78

**Abstract:**

*Mycobacterium bovis* Bacillus Calmette-Guerin (BCG) is the primary treatment for non-muscle-invasive bladder cancer (NMIBC), known to stimulate inflammatory cytokines, notably interferon (IFN)-γ. We observed that prolonged IFN-γ exposure fosters adaptive resistance in recurrent tumors, aiding immune evasion and tumor proliferation. We identify HLA-E and NKG2A, part of a novel NK and T cell checkpoint pathway, as key mediators of resistance in BCG-unresponsive NMIBC. IFN-γ enhances HLA-E and PD-L1 expression in recurrent tumors, with an enrichment of intra-tumoral NKG2A-expressing NK and CD8 T cells. CXCL9$^+$ macrophages and dendritic cells and CXCL12-expressing stromal cells likely recruit CXCR3/CXCR4-expressing NK and T cells and CXCR7$^+$ HLA-E$^{HIGH}$ tumor cells. NK and CD8 T cells remain functional within BCG-unresponsive tumors but are inhibited by HLA-E and PD-L1, providing a framework for combined NKG2A and PD-L1 blockade strategy for bladder-sparing treatment of BCG-unresponsive NMIBC.

**Introduction**

Intravesical *Mycobacterium bovis* Bacillus Calmette-Guerin (BCG) is the only FDA-approved, first-line treatment for high risk non-muscle-invasive bladder cancer (NMIBC), and is hypothesized to act in part via the release of inflammatory cytokines, notably interferon (IFN)-γ [1].Unfortunately, tumor recurrence following BCG treatment is common, with local recurrence rates between 32.6 to 42.1% and progression rates between 9.5 to 13.4% [2]. Radical cystectomy is the only definitive treatment option for BCG-unresponsive disease, leaving patients at high risk for complications and the need for urinary diversion which can lead to diminished quality of life [3,4]. Currently, there is substantial interest and numerous ongoing clinical trials to identify novel therapeutic options for BCG-unresponsive NMIBC.

Despite the use of BCG as the first-line immunotherapy for NMIBC for over 40 years, the mechanisms underlying resistance to BCG's therapeutic benefits are poorly understood. This lack of understanding has further hindered progress in developing novel approaches to treat tumors resistant to BCG. The six-dose weekly induction regimen was devised anecdotally based on rudimentary urine studies performed in the 1970's [5]. Only a single dosing study has ever been performed to assess the immunologic response with this dosing regimen, which concluded that six doses stimulated an immune response, but did not investigate the ramifications of the magnitude nor the duration of that immune response [6]. Contributing to the difficulty in predicting BCG response is the variety of immune lineages implicated in both its success and failure, including neutrophils, monocytes, macrophages, dendritic cells (DCs), T cells, and natural killer (NK) cells [7].

Adaptive immune resistance, in which cancers are recognized by immune cells, but accommodative changes prevent effective clearance, is an important mechanism of therapeutic resistance [8]. IFN-γ has been shown to have both pro- and anti-tumorigenic functions across an array of tumor types[9]. The equilibrium established by IFN-γ signaling may be critical in preventing therapeutic resistance. In melanoma and lung cancer, dysfunctional and overactive IFN-γ signaling is known to drive T cell inhibition via upregulation of PD-L1 [10]. In muscle-invasive and metastatic urothelial carcinoma, IFN-γ associates with improved as well as poor responses to PD-1/PD-L1 blockade and exists as part of a balance (coined 2IR score) between pro- and anti-tumorigenic pathways expressed in the TME [11]. The impact of long-term intravesical BCG, which drives chronic immune stimulation via IFN-γ production, and its role in immune dysregulation and exhaustion in BCG unresponsive NMIBC is emerging but still unclear [12, 13].

Previous work comparing BCG-unresponsive to treatment-naïve tumors showed a significant increase in PD-L1 expression, but only in a minority of unresponsive samples [14]. Immunotherapeutic strategies, especially those engaging the PD-1/PD-L1 axis to unleash exhausted immune cells, have

modestly improved NMIBC treatment. Unfortunately, response rates to PD-1 blockade monotherapies in NMIBC remain low, at ~19% durable response [15,16,17,18]. Further, PD-L1 expression levels alone have failed to accurately predict response to immunotherapy in NMIBC, suggesting that alternative mechanisms of immune suppression are engaged [19].

The HLA-E/NKG2A axis has been identified as a potent immune checkpoint regulating both CD8 T and NK cells [20,21,22,23,24]. Abrogation of HLA-E interactions with NKG2A on NK cells increased effector activity against tumor cells and potentiated anti-tumor function. Further, blocking NKG2A on CD8 T cells in a murine model increased tumor vaccine response, even in a PD-1 resistant setting [20,22]. Additionally, the combination of anti-NKG2A antibody (monalizumab) and antiPD-L1 antibody (durvalumab) has been studied in a Phase 2 clinical trial in patients with nonresectable, stage III NSCLC. The combination demonstrated an objective response rate of 35% vs 17.9% with durvalumab alone, as well as improved progression-free survival (PFS) with durvalumab and monalizumab (72.7%) compared to durvalumab alone (33.9%) [25].

Most recently, BCG was shown to directly infect a proportion of bladder tumor cells in NMIBC patients driving impairment in autophagy flux resulting in downregulation of HLA class I expression along with a shift towards epithelial-to-mesenchymal transitioning (EMT) by BCG-infected tumor cells [26]. In a cohort of BCG-treated NMIBC patients with tumor recurrence, the authors observed that abundance of class I HLA-negative tumor cells associated with disease progression and metastasis and overall survival. This study highlighted two different tumor cell types within BCG-unresponsive settings and suggested that a tumor phenotype defined by high HLA class I expression correlated with improved long-term survival post-cystectomy.

In this study, we confirm the presence of both tumor cell phenotypes but focus on their interactions with stromal and immune cells. We present a novel mechanism of resistance, whereby HLA class I-positive tumor cells are activated in response to NK and CD8 T cell-derived IFN-γ and recruited to CXCL12-expressing stromal-rich regions of the tumor where myeloid cells, regulatory T cells (Tregs), and immune effector cells (NK, CD8 T cells) are enriched. Consequently, class I HLA-negative tumor cells can subvert immune cell trafficking. We pose a novel hypothesis that BCG more broadly induces changes in the bladder tumor microenvironment (TME), promoting expression of NKG2A and PD-1 on CD8$^+$ T cells, NKG2A on NK cells, and HLA-E, PD-L1, and CXCR7 on recurring, cytokine-activated tumor cells, and these phenotypic changes are mechanistically linked to treatment resistance and immune evasion by recurring tumors. This study investigates HLA-E and NKG2A as a novel axis of resistance induced by chronic inflammation and provides a mechanistic framework for a bladder-sparing combination immunotherapy strategy of NKG2A and PD-L1 blockade for overcoming resistance to BCG immunotherapy for treating high-grade NMIBC.

## Results

**Clinical cohort description.**

Cohort descriptions of patient samples used for the experiments below are described in Table 1.
Patient samples were collected and processed at The Mount Sinai Hospital, and the validation cohort was
collected and processed at Aarhus University Hospital, Denmark. Median patient age ranged from 64 to
70.7 years old; median months to recurrence ranged from 1.7 to 2.9 months; median months to progression
ranged from 4.43 to 23.28 (**Table 1**).

**IFN-γ initiates adaptive resistance including increased tumor expression of HLA-E and PD-L1.**

To investigate potential mechanisms underlying BCG resistance in NMIBC, we used tumor
specimens from BCG-naïve patients, prior to initiating induction intravesical BCG therapy and BCG-
unresponsive tumors, with a recurrence after at least an induction course of intravesical BCG, collected
from a prospective cohort of NMIBC patients. Targeted *in situ* hybridization and *mRNA* sequencing and
gene-set enrichment analysis (GSEA) comparing BCG-unresponsive tumors to pre-treatment timepoints
were performed (**Fig. 1A***)* [27]. Significant gene sets are shown ranked by their normalized enrichment score
(NES) (**Fig. 1A**, **Supplemental Table S1***)*. The top enriched gene-sets comparing BCG-unresponsive to
BCG-naïve tumors (n=36) included, but were not limited to, IFN-γ response, cytotoxic lymphocytes, T cell
checkpoint modulation, tumor inflammation, TH1 response, and lymphocyte trafficking (**Supplemental**
**Table S1**). Of note, the "Lymphocyte trafficking" pathway was largely driven by a combination of genes
encoding activating cytokines and receptors as well as inhibitory immune checkpoint genes (*PDCD1* (PD-
1), *CD274* (PD-L1), *PDCD1LG2* (PD-L2), *CTLA4*, *TIGIT*, *KLRC1* (NKG2A), *HLA-E*) indicating a balance
between antitumor and pro-tumorigenic features in the TME. Gene sets downregulated in the BCG-
unresponsive state and upregulated in the BCG naïve state included TGF beta signaling, hypoxia,
peroxisome, oxidative phosphorylation, P53 pathway, DNA repair, and stress toxicity (**Fig. 1A**).

To better understand the inflammatory state observed in BCG-unresponsive tumors, we used cell-
free urine supernatants from a prospective cohort of BCG-treated NMIBC patients and performed an
analysis using the OLINK Proteomics™ "inflammation" panel, a 92-analyte protein proximity extension
assay, to profile urine cytokines reflective of the urothelial microenvironment [28, 29]. Independent of clinical
response to BCG therapy, all patients experienced elevated cytokine levels in their urine. Significantly
upregulated cytokines of interest at the sixth dose compared with the BCG-naïve state included IFN-γ,
Flt3L, IL2RB, CCL8, CCL13, CCL19, CCL23, CXCL1, CXCL9, CXCL10, CXCL11, and TWEAK (**Fig.**

**1B**; all p<0.05). Importantly, urine cytokine levels kept increasing with each consecutive dose of BCG,
where even after a six-week treatment interruption (time of first cystoscopy,) urine cytokine levels never
equilibrated back to baseline, although the differences were not significant. The inflammatory signature
was validated in an independent cohort, "Aarhus" (Aarhus University Hospital, Denmark), consisting of
BCG-treated patients with long term clinical follow-up. Urine samples from 66 patients were collected at
the BCG-naïve time point and compared to patient-matched urine samples collected within four months
after completion of BCG induction course. CXCL9, CXCL10, CXCL11, CCL19, and CCL23 had lasting
upregulation at the post-BCG timepoint (p < 0.05) (**Fig. 1C**). Additionally, urinary protein profiles at these
timepoints were compared between responsive and unresponsive patients considering whether visible
bladder tumors were present at the time of urine sampling (**Fig. 1D**). With exception of CXCL5, when
assessing urine from BCG-unresponsive patients (before the start of BCG induction therapy) we identified
significantly elevated levels of cytokines and chemokines only when tumors were present at time of
sampling, where the presence of CCL3, CCL20, CXCL11, IL-8, MMP7 and MMP12 were attributable to
bladder tumors rather than in response to BCG. Patients unresponsive to BCG showed a distinct urinary
protein profile after BCG treatment with a greater magnitude of inflammatory cytokine and chemokine
production as well as Caspase-8 (a marker of apoptosis) and TGF-β1 in its pre-processed inactive state
interacting with its latency associated peptide (LAP) (**Fig. 1D**).

To better understand the effects of IFN-γ on recurring bladder tumors, primary tumor cells from
NMIBC patients (n=10) and urothelial tumor lines were cultured for 24 hours in the presence or absence of
recombinant human IFN-γ (**Fig. 1E**) and HLA-E and PD-L1 expression were measured via fluorescence
flow cytometry. Tumor cells expressed significantly higher levels of PD-L1 and HLA-E following
stimulation with IFN-γ, which was also observed across the collection of bladder tumor lines (n=11) (**Fig.
1E**). We saw no effects of IFN-γ on wild type (WT) K562 tumors (lack expression of HLA class I, including
HLA-E). HLA-E-stably–transduced K562 cells were used as a positive control for maximal HLA-E
expression. Overall, these results suggest adaptive resistance by recurring tumor cells after BCG therapy
that respond to IFN-γ by increasing HLA-E and PD-L1 expression and in a microenvironment replete with
immune effector cells.

**Selective upregulation of tumor HLA-E in regions near NKG2A-expressing NK and CD8 T cells**

To validate *in situ* hybridization *mRNA* analyses and to better understand the spatial relationship
between tumor cells and tumor infiltrating lymphocytes (TILs), we performed multiplex imaging analyses
using both immunofluorescence (IF) as well as immunohistochemistry (IHC) with whole slide scanning for
targeted high-resolution analyses on a cohort BCG naïve (N=17) and BCG unresponsive (N=24) NMIBC
patients (**Fig. 2**). We first performed a subset analysis of NMIBC sections (N=4) using a PhenoCycler™

(formerly CODEX) imaging system (Akoya Biosciences). Representative imaging for one BCG-

unresponsive NMIBC section highlights S100A4$^+$ tumor cells intermixed with stroma (Vimentin) (**Fig.**

**2A**). We observed a large abundance of NKp46$^+$ NK cells and CD8 T cells with variegated expression of

NKG2A and/or PD-1 (**Figs. 2B** and **2C**). While the analysis was underpowered for robust analyses, the data

demonstrated substantially greater abundance of NKG2A-expressing NK and CD8 T cells in BCG-

unresponsive sections and in regions of tumor tissue with high HLA-E expression by tumor cells and

stroma.

To validate the observation, we stained the cohort of 41 NMIBC sections by multiplexed IHC for Pan-

cytokeratin (pan-CK), HLA-E, CD3,  and NKG2A (**Fig. 2D**). Among tumors from the cohort, we analyzed

840,348 tumor cells in total (277,315 for BCG-naïve and 563,033 for BCG-unresponsive).  We used Halo

digital pathology software (Indica Labs, Albuquerque, NM) to annotate cell lineages, and to measure tumor

HLA-E expression within the for density and spatial relationships with stromal and T/NK cell infiltration

and proximity (**Fig. 2E-2I**). Here, we confirmed that HLA-E$^{BRIGHT}$ tumor cells were significantly more

abundant in BCG-unresponsive tumors (p= 0.0002) and represented 43.6%% of total tumor cells compared

to 10.3% in BCG-naïve tumors (**Fig. 2G**). Further, we identified NKG2A$^+$ CD3$^-$ NK cells and NKG2A$^+$

CD3$^+$ T cells (**Fig. 2H**) and determined they were in significantly closer proximity to HLA-E$^{BRIGHT}$ tumor

cells (p<0.00001)(**Fig. 2I**). Finally, we performed a proximity ligation assay by immunofluorescence using

antibodies targeting HLA-E and NKG2A to quantify the numbers of interactions between NKG2A$^+$

NK/CD8 T cells and HLA-E$^+$ tumor cells and found a significantly greater number of interactions in BCG-

unresponsive tumors (p = 0.009) (**Fig. 2J** and **2K**).

**Chronic activation of tumor-infiltrating NK and CD8 T cells in BCG-unresponsive tumors**

To better understand the composition of the tumor microenvironment and to characterize the

potential drivers of adaptive resistance, single-cell RNA sequencing (scRNA-seq) of nine tumors (BCG-

naïve (n=3) and unresponsive (n=6)), was performed. In total, 65,324 tumor-derived cells containing 26,520

gene features passed quality control and were used for subsequent analyses, with a median of 958 genes

detected per cell.

Twenty-two identifiable cell populations, including CD4 and CD8A T cells, regulatory T cells

(Tregs), NK cells, dendritic cells (DCs), monocytes and macrophages, granulocytes, B cells, plasma cells,

and nonhematopoietic stromal, and tumor cells were identified after graph-based clustering (**Fig. 3A** and

**3B**, **Supplementary Table S3 and S4**). To identify the cell origin(s) of IFN-γ and related pro-inflammatory

gene signatures, we examined the expression of *IFNG* in each of the cell populations. NK cells, CD8 T

cells, and a subset comprised of both CD4 and CD8 T cells defined by cell cycle genes (T cell-cycle)

demonstrated the highest levels of *IFNG* expression among all cell clusters and had distinct IFN-γ$^{high}$ cell

populations (**Extended data Fig. S1A).** Additionally, we profiled the median gene expression across all

major canonical checkpoint receptors and observed variegated expression on T cell subsets. However,

tumor-derived NK cells mostly expressed *KLRC1*, and to a lesser extent *HAVCR2* (Tim3) and *TIGIT*

(**Extended data Fig. S1B).**

281           To better understand the functional significance associated with NKG2A expression, we stratified

NK cells, CD8 T and T cell-cycle cells by low and high *KLRC1* expression and performed a differential

expression of genes (DEG) analysis (**Fig. 3C**). Our analysis revealed a balance between cytokine and

cytolytic activity and expression of inhibitory checkpoint receptors. *KLRC1$^{high}$* NK cells expressed

significantly higher CCL5, CSF1, *XCL2,* IL2RB, and *FCER1G*, but additionally expressed higher levels of

*TIGIT*. Tumor-infiltrating NK cells also expressed significantly higher levels of *CD2*, *ITGA1* (CD49a),

*ITGAE* (CD103) and *RGS1* suggesting they are tissue-resident. Additionally, we observed that *KLRC1$^{high}$*

CD8 T cells expressed elevated levels of CCL3, CSF1, *GNLY*, *GZMA*, *GZMB*, and *PRF1* (perforin), but

they additionally expressed higher levels of *CTLA4*, *ENTPD1* (CD39), *HAVCR2*, *LAG3*, *KIR2DL4* and

*TIGIT*. *KLRC1$^{high}$* T cell-cycle cells showed significant upregulation in *CCL5, CSF1, CXCL13, NKG7,*

*GNLY*, *GZMA*, *GZMB*, but additionally expressed *LAG3*, *KIR2DL4,* and *TIGIT*.  Collectively, the data

suggest that BCG therapy drives strong immune activation and that many repeated doses of treatment

sustain activation over prolonged period driving upregulation of inhibitory checkpoint pathways.

**Functionally distinct states of bladder tumor-derived NK and CD8 T cells in BCG-unresponsive**

**NMIBC**

297           NK cells are very heterogenous and remain poorly described in solid tumors with virtually nothing

known in human bladder cancer. We performed unbiased clustering on all NK cells to derive five major

clusters (**Fig. 3D**), which were divided across conventional definitions of CD56$^{bright}$ and CD56$^{dim}$ NK cells

(**Fig. 3E**). A pathway analysis revealed significant enrichment in CD56$^{bright}$ NK cells for genes involved in

metabolism, IL-2 signaling, and immune 'stress' (ROS and DNA repair) (**Fig. 3F**). Conversely, CD56$^{dim}$

NK cell enriched pathways included response to TNFα and IFNα and apoptosis, MAPK activity but also

reflect immune 'stress' (hypoxia, complement regulation). Building on two recent seminal resources for

transcriptional reference mapping of human NK cells in solid tissues[30, 31], we integrated the newly defined

gene signatures to annotate 'functional states' to bladder tumor-derived NK cells (**Fig. 3G**, **Extended data**

**Fig. S2**). Our analysis confirmed that clusters 1 and 2 are very similar to Group 2 typical CD56$^{bright}$ NK

cells, and cluster 5 strongly correlates with Group 1 "stressed"  CD56$^{bright}$ NK cells, described by Netskar

et al.[30]. Collectively clusters 1,2, and 5 reflecting Groups 1 and 2 NK cells represent ~50% of the total

tumor-infiltrating NK cells. Additionally, the remaining fraction of CD56$^{dim}$ NK cells were distributed

among groups 3-6 NK cells, defined by effector functions (Gr. 3), adaptive NKG2C[+] NK cells (Gr. 4),

activated (Gr. 5), and typical (Gr. 6) NK cells.

We similarly applied gene signatures for NK1, NK2, and NK3 subsets as recently demonstrated by

Rebuffet et al[31, 32], and these definitions more closely reflect a developmental or ontogeny trajectory of NK

cells across human tissues, NK1 and NK-intermediate cells represent immature CD56[bright] NK cells and

NK3 cells include terminally mature and adaptive NK cells (**Extended data Fig. S2**). Our comparative

analysis revealed an enrichment of NK-intermediate cells representing the original clusters 1 and 2, and

NK1B cells representing the original clusters 3 and 4. Both NK1B and NK-intermediate signatures were

shared among the original cluster 5 suggesting transitional properties of this NK cell subset. Finally, we

profiled tumor-infiltrating NK cells according to the six functional states (Gr. 1-6) (and CD8 T cells and T

cell-cycle cells according to KLRC1[+/-]) for expression of tissue residency genes as well as

cytokines/chemokines and cytokine/chemokine receptors to understand how NK cells and T cells are

trafficking to the bladder, and we confirmed they are functionally competent (**Fig. 3H** and **3I**). We also

observed clear evidence suggesting functional specialization within the NK cells and within the T cells. For

instance, *KLRC1[+]* CD8 T cells express high levels of *IL2RB, TNFRFS9* (4-1BB), *CSF1 and FASLG*

compared to their *KLRC1[-]* counterparts. However, *KLRC1[-]* CD8 T cells expressed higher levels of *IFNG*,

and *FLT3LG* (**Fig. 3I**).

**BCG-unresponsive tumors are characterized by a segregated microenvironment, where activated**

**HLA-E[HIGH] tumor cells are recruited to stromal regions replete with immune cells**.

We next sought to understand the composition of tumor cells in tumors from both treatment-naïve

(N=14,852) and BCG-unresponsive (N=29,858) NMIBC patients. UMAP clustering of scRNAseq analysis

revealed seven major subsets of tumor cells (**Fig. 4A**), where clusters B1 and B3 had significantly higher

HLA-E expression (**Fig. 4B**) and were significantly more responsive to cytokines and inflammatory signals

(**Fig. 4C**). We profiled tumor clusters for expression of known inhibitory and activating ligands as well as

for cytokines/chemokines and cytokine/chemokine receptors (**Figs. 4D** and **4E**). Importantly, of all

chemokine receptors, we observed that *HLA-E[HIGH]* bladder tumor cells had significantly higher *ACKR3*

(CXCR7) expression, and pseudo-bulking the scRNAseq data revealed that HLA-E and CXCR7 tumor

expression were strongly correlated (**Fig. 4F**). Additionally, we profiled the abundance of each tumor

cluster according to BCG treatment status and observed cluster B2 was uniquely enriched in BCG-naïve

tumors (**Fig. 4G**). The other tumor clusters were either evenly distributed between naïve and unresponsive

tumors or they were disproportionately skewed towards unresponsive.

Collectively, our analyses of NK cells, T cells and tumor cells suggest immune crosstalk within the

TME. Further, the data are supported by our observations that NKG2A[+] NK and CD8 T cells are

significantly closer in proximity to HLA-E$^{HIGH}$ tumor cells (**Fig. 2**). NK cells and T cells expressed a

variegated array of *CXCR3, CXCR4*, and *CCR7* (**Fig. 3I**) and *HLA-E*$^{HIGH}$ tumor cells uniquely express

*ACKR3* (CXCR7) (and to lesser extent *CXCR4*) (**Fig. 4E**), indicating an important axis with CXCL12.

Additionally, *KLRC1*-expressing NK and CD8 T cells expressed significantly higher *AREG* (amphiregulin)

which signals through EGFR. To test the hypothesis that chemokine signaling explains why NKG2A$^+$ NK

and CD8 T cells are biased towards HLA-E$^{HIGH}$ tumor cells, we measured across all major clusters

expression of chemokines and their receptors: a) CCL19, CCL21 → CCR7; b) CXCL9, CXCL10, CXCL11

→ CXCR3; c) CXCL12 → CXCR4 and CXCR7; and d) amphiregulin → EGFR (**Fig. 4H**). We observed

*HLA-E*$^{HIGH}$ tumor cells expressed higher levels of EGFR compared to *HLA-E*$^{DIM/NEG}$ tumor cells, whereas

*AREG* was expressed by NK cells as well as dendritic cells (DCs), monocytes, mast cells, and even *HLA-*

*E*$^{HIGH}$ tumor cells at low levels. CXCL12 was mostly expressed by stromal cells, whereas CXCR4 was

expressed by NK cells, B and T cells (including Tregs), and myeloid cells. CXCR3 was mostly relegated

to NK and T cells and DC1 cells, and CXCL9, CXCL10, and CXCL11 were mostly expressed by

monocytes, macrophages and DCs. Finally, CCL19 was very specific for mregDCs and CCL21 by

endothelial cells suggesting important interactions with CCR7-expressing B and T cells.

In a recent study of BCG-unresponsive NMIBC by Rouanne et al[26], the authors demonstrated that

BCG can directly infect bladder tumor lines driving epithelial to mesenchymal transitioning (EMT),

autophagy flux impairment, a loss of class I HLA expression, and increased cell cycling. The study focused

on BCG-unresponsive NMIBC and found that patients with a dominant tumor phenotype showing these

hallmarks experienced significantly poorer metastasis-free survival. In our study, we next sought to stratify

tumor cells according to their HLA-E expression and measure average gene expression across the hallmark

pathways for EMT, autophagy, and cell cycle (**Extended data Fig. S3**). We observed a remarkably similar

phenotype of HLA-E-negative tumors as *in vitro* BCG-infected tumor lines showing significantly higher

EMT phenotype along with reduced autophagy flux genes and increased cell cycle. The data might suggest

that a proportion of recurring tumor cells in the face of BCG therapy, undergo *in vivo* infection rendering

them less sensitive to cytokine stimulation and chemokine attraction and promoting this stratification where

HLA-E$^{HIGH}$ tumor cells are in significantly closer proximity to CXCL12$^+$ stromal cells and immune cells.

To study the potential interactions between NK and CD8 T cells and HLA-E$^{HIGH}$ tumor cells, we

modeled the interactions using NicheNet, an algorithm that uses a combination of transcriptomic data,

known ligand-receptor interactions, and gene-gene signaling and regulatory relationships to predict

potential interactions between cells of interest (**Fig. 4I**) [33]. The top twelve ranked ligands from KLRC1$^+$

NK, KLRC1$^+$ CD8 T cells, and KLRC1$^+$ T cell-cycle cells predicted to interact with HLA-E$^{BRIGHT}$ tumor

cells were *IFNG, TNF, TNFSF12* (TWEAK), *APOE, TNFSF14* (LIGHT), SEMA4D, *TNFSF10* (TRAIL),

*AREG, GZMB, ALOX5AP, GPI, NUCB2* (**Fig. 4J**). KLRC1-expressing T cell-cycle and CD8 T cells

expressed the highest levels of IFNG, while *AREG* expression was highest in NK cells. Putative receptors

for these ligands, including *IFNGR1, IFNGR2, TNFRSF1A/10B/12A/14/21, SORL1, LDLR, AMFR,*

*ALOX5, EGFR, IGF2R, and PLXNB1* were most highly expressed in the BCG-unresponsive samples,

suggesting the presence of corresponding signaling activity by recurring tumors (**Fig. 4J**). To further

examine genes upregulated in the tumor as a consequence of signals from NK and T cells, a circos plot

revealed the ligand-target weights of the top third of predicted relationships between 'sender' or effector

ligands and 'receiver' or target genes, defining the 'putative effects of the HLA-E/NKG2A axis" in bladder

tumors as well a targeted 'IFNγ-mediated' and 'AREG-mediated' adaptive resistance signatures (**Fig. 4K**).

**Organization of the TME through immune crosstalk with stroma and HLA-E^bright tumors**

388           To spatially resolve the relationships between tumor cells, stroma and immune infiltrates, we

performed spatial transcriptomics (ST-seq) analysis of bladder tumor sections from four treatment-naïve

and four BCG-unresponsive tumors (**Fig. 5**). Clustering analysis revealed 12 unique clusters, where cluster

12 remained unclassified and was removed from subsequent analyses (**Fig. 5A**-**5C**). Clusters were

annotated by their proportional composition by immune, stromal, and/or tumor cells defined using gene

signatures derived from scRNAseq analysis (**Figs. 3** and **4**). Clusters were defined both by the gene content

defining immune, stromal, and/or tumor cells but additionally integrating regional features of the tumor

tissue (**Fig. 5C** and **5D, Extended Figure S4**). We profiled the relative abundance of each spatial cluster

between BCG-naïve (pre-BCG) and BCG-unresponsive (post-BCG) tumors (**Fig. 5E**). We observed clear

differences, where spatial clusters #0, 3, 4, and 6 were substantially enriched in BCG-unresponsive tumors.

Conversely, there was a trend for enrichment of clusters # 1, 2, 5, 7, 8, and 11 in BCG-naïve tumors. Only

spatial cluster # 8 lacked the presence of immune cells, and cluster 10 was defined by neutrophils

exclusively, where they were near regions enriched with DCs, Tregs, T cell-cycle and tumor cells.

401           ST-seq Visium™ spots are 55μm in diameter and require deconvolution to understand which cell

lineages are represented as well as their relative abundances (**Fig. 5F**). We performed neighborhood

analyses to measure proximity of *HLA-E*^HIGH and *HLA-E*^DIM/NEG tumor Visium spots to CD8 T cells and NK

cells (**Fig. 5G**) as well as Tregs (**Fig. 5H**). A summary of the average distances to *HLA-E*^HIGH and *HLA-*

*E*^DIM/NEG tumor Visium spots from BCG-naïve and BCG-unresponsive tumors revealed significantly closer

proximity of *HLA-E*^HIGH tumor spots to NK cells, Tregs, and CD8 T cells. This observation was restricted

to BCG-unresponsive tumors (**Fig. 5I**).

408           To better understand the mechanisms underlying preferential interactions of immune cells with

409     *HLA-E*^HIGH tumor cells, we tested the hypothesis that stroma-produced CXCL12 mediates recruitment of

410     CXCR7+ HLA-E^HIGH tumor cells to regions where macrophages and DCs produce CXCL9, CXCL10, and/or

411     CXCL11 that mediate recruitment of NK cells and T cells (**Fig. 4H**). Spatial relationships were investigated

on the 'nano-environment' level, defined as a Visium spot with a radius of 2.5 adjacent Visium spots, or ~138 microns. A representative BCG unresponsive tumor section with brightfield image of underlying tissue is shown with $HLA\text{-}E^{HIGH}$ tumor/stroma Visium spots shown in blue, and $HLA\text{-}E^{DIM/NEG}$ Visium spots shown in tan represented in **Fig. 6A**. Additional profiling of NK and CD8 T cells and Tregs revealed their distribution being close in proximity to $HLA\text{-}E^{HIGH}$ tumor/stroma Visium spots. BCG-unresponsive tumors revealed significantly higher levels of tumor/stroma spot infiltration with cytotoxic NK and CD8 T cells (7.05 times closer) and regulatory T cells (Tregs, 2.66 times closer) when compared to the BCG-naïve state ($p < 0.0001$ for both). Using the same representative BCG-unresponsive tumor section, a topographical map of HLA-E expressing tumor/stroma spot density is shown overlaid on top of a neighborhood analysis highlighting the intensity of neighboring NK and CD8 T cell infiltration (**Fig. 6B**).

To investigate the biased distribution of NK and T cells near $HLA\text{-}E^{HIGH}$ tumor-stroma spots, we profiled bladder tumor-stroma spots for expression of cytokines and chemokines (**Fig. 4H**). Among the chemokines with detectable gene expression by ST-seq analysis, we observed a strong correlation between CXCL9 (**Fig. 6C-6E, Extended Figure S5**) and HLA-E expression as well as between CXCL12 and HLA-E expression (**Fig. 6F** and **6G**). We next applied our neighborhood analysis to profile tumor expression of known checkpoint ligands, HLA class I expression as well as the IFN-response pathway as a function of proximity to neighboring NK cells and CD8 T cells (**Fig. 6H**). Analysis of 5,319 tumor-stroma spots (defined by >95% tumor- and stroma-specific genes revealed a clear program by tumor/stroma regions increasing expression of inhibitory ligands *CD47*, *CD274* (PD-L1), and *HLA-E* as a greater number of NK cells and CD8 T cells were in contact with tumor-stroma spots (**Fig. 4G**). Additionally, we observed increased tumor-stroma spot expression of the apoptotic genes *PDCD2* and *FAS* and the adhesion molecule *CD58* (LFA-3), known to interact with CD2-expressing NK and T cells. Finally, tumor-stroma spot expression of *IFNGR1* and *IFNGR2* along with *JAK1/JAK2*, *STAT1*, *IRF1* and *IRF9* were increased as a greater number of NK and CD8 T cells were in contact. *STAT3* expression did not change as a function of neighboring NK and CD8 T cells, and *SOCS1* expression was reduced with neighboring NK and CD8 T cells, suggesting that tumor-stroma regions are actively responding to IFN signals. Tumor-stroma expression of these genes was further stratified according to high vs dim/negative expression of *HLA-E* (**Fig. 6I**).

To validate the ST-seq analyses, we used immunofluoresecence and the PhenoCycler/CODEX platform to profile 59 antibodies (**Supplemental Table S2**) in order to better define the interactions between tumor cells and multiple cell lineages. Antibodies targeted markers distinguishing tumor cells (pan-cytokeratin (CK), GATA-3, S100A4) from stromal endothelial cells and fibroblasts (Collagen Type IV, podoplanin, vimentin, PECAM1/CD31), and immune cells (CD45), including monocytes, macrophages, dendritic cells (CD14, CD16, CD68, CD11b, CD11c, HLA-DR), neutrophils (CD66b, CD16), B cells

(CD20, CD138), CD4 and CD8 T cells (CD3, CD4, CD8), Tregs (CD3, CD4, FoxP3), and NK cells

(NKp46, Gzm.B, CD16) (**Fig. 6J** and **6K**).

**Tumor-derived NK and CD8 T cells retain functional competence at time of tumor recurrence.**

To better understand if an elevated expression of cytolytic genes in tumor-derived NKG2A-

expressing NK and CD8 T cell populations correlated with improved functional sensitivity, we used mass

cytometry (CyTOF) to profile the *in vitro* responsiveness of tumor-derived cells from BCG-unresponsive

NMIBC patients at time of tumor recurrence. We tested their ability to respond to HLA-E$^+$ PD-L1$^+$ tumors

in the presence/absence of durvalumab (anti-PD-L1 antibody) and/or monalizumab (anti-NKG2A

antibody), using an artificial K562 acute myeloid leukemia tumor model that allows for uniform HLA-E

and PD-L1 expression via pre-stimulation of tumor lines with exogenous recombinant IFN-γ. Briefly, after

24 hours incubation with recombinant IFN-γ [400ng/ml], stably transduced HLA-E$^+$ K562 cells saw 2.7-

fold and 2.1-fold-increase in expression of HLA-E and PD-L1, respectively, when compared with baseline

expression levels (**Fig. 7A**). Separately, tumor-infiltrating lymphocytes from four BCG-unresponsive

tumors were isolated and then expanded up to 13 days in the presence of low dose IL-2, IL-7, and IL-15 as

well as tetramers targeting CD3 and CD28. Finally, TGF-β, was included in the stimulation cocktail for an

additional three days to promote NKG2A expression by CD8 T cells [24]. Phenograph clustering analysis [34]

of expanded TILs revealed NK cells uniformly expressing NKG2A as well as 10 unique clusters of CD8 T

cells with variegated expression of inhibitory checkpoint receptors (CD161, GITR, LAG3, PD-1, TIGIT,

Tim-3, NKG2A) as well as markers of activation, trafficking, and tissue residence (CD25, CCR4, CCR6,

2B4, CD103, CXCR3) (**Fig. 7B** and **7C**). Four clusters (ID#: 2, 9, 15, 17) expressed elevated levels of

NKG2A, representing ~32.5% of the total CD8 TILs and reflecting the fraction of TILs predicted to react

to NKG2A-blockade. Additionally, ~9% (range, 3.91% - 14.04%) of all expanded TILs were comprised of

NK cells (CD14$^-$ CD19$^-$ CD3$^-$ CD56$^+$) and NK and CD56$^+$ CD8 T cells were strongly enriched for co-

expression of NKG2A. CD56 was then used as a surrogate for NKG2A as use of monalizumab precluded

our ability for detecting NKG2A with antibodies by CyTOF. Further, NKG2A and PD-1 expression by

CD8 T cells was variegated highlighting the dynamic and non-redundant mechanisms for regulating CD8

T cell activation [24].

Following expansion and characterization, we performed 6-hour co-cultures of TILs with K562

cells with and without HLA-E and/or PD-L1 expression (**Fig. 7D-7F**). Additionally, we measured the

effects of PD-L1-blockade or combined PD-L1- and NKG2A-blockade on NK and CD8 T cell

degranulation (CD107a) and expression of IFN-γ. WT K562 cells were used as positive control condition

for measuring NK and NKG2A-expressing CD8 T cell activation. We observed that NK cells and CD56$^+$

CD8 T cells were sensitive to HLA class I$^{-/-}$ K562 cells compared to *ex vivo* activity, and their response was

diminished when K562 cells stably expressed HLA-E or HLA-E and PD-L1. NK cell activation was

restored in response to both HLA-E[+] and HLA-E[+] PD-L1[+] K562 cells when pre-treated with anti-NKG2A

Ab alone or in combination with anti-PD-L1 Ab (CD107a: 2.1-fold increase, p=0.02; IFN-γ: 1.6-fold

increase, p=0.05).

Unlike NK cells, CD56[+] CD8 T cells were more dynamic in their co-expression profiles for

NKG2A and PD-1 staining. Similar to NK cells, untreated CD56[+] CD8 TILs were not reactive to HLA-E[+]

K562 cells with or without co-expression of PD-L1. Activation of CD56[+] CD8 TILs was only significant

in the presence of both anti-NKG2A and anti-PD-L1 Abs (CD107a: 1.75-fold increase, p=0.02; IFN-γ: 1.8-

fold increase, p=0.007) (**Fig. 7F**). Collectively, the data strongly suggest sensitivity of NK and CD8 T cells

to combination immunotherapy strategies following tumor recurrence after failing BCG therapy.

**Discussion:**

Adaptive resistance can be driven by tumor-intrinsic and/or extrinsic mechanisms in overcoming

immune pressures [11, 12, 13, 35]. Evidence from mouse and human studies demonstrates that interferons, while

stimulating a robust anti-tumor response, also upregulate immune-suppressive factors in the setting of

prolonged activation. In melanoma, IFN-γ from CD8 T cells was shown to upregulate tumor PD-L1

expression and mediate infiltration of FOXP3[+] regulatory T cell pathways within the TME [36]. In a broader

meta-analysis across 18 tumor indications, including bladder cancer, inflammatory mediators including

IFN-γ were associated with inhibitory immune checkpoints, including PD-L1/L2 [37, 38]. Despite emerging

evidence positioning pro-tumorigenic roles for IFN-γ, there are well-established anti-tumor functions

mediated through IFN-γ that are critical for anti-PD-L1 immunotherapy [39]. Anti-tumor inflammation,

therefore, exists along a continuum where an equilibrium is necessary for appropriate immunotherapeutic

efficacy[40].

In a recent investigation conducted by Rouanne et al.[26], distinct patterns of disease progression after

BCG therapy and cystectomy were identified in the context of NMIBC. The study revealed that BCG-

unresponsive NMIBC patients characterized by tumors bearing a 'BCG infection' phenotype correlated

with poorer survival. Conversely, BCG-unresponsive patients exhibiting HLA class I proficiency in tumor

cells after BCG therapy (before cystectomy) demonstrated greater abundance of CD8 T cell tumor

infiltrates, upregulation of inflammatory cytokines, and the presence of inhibitory checkpoint molecules.

This latter group experienced significantly improved metastasis-free long-term survival post-cystectomy.

In our own investigation, we delved into the primary resistance mechanisms to BCG therapy and

explored the impact of BCG-induced inflammation on the dynamic interplay between tumor and immune

cell populations. Utilizing both prospective and retrospective NMIBC specimens collected before, during,

and after BCG treatment, our study uncovered a pervasive inflammatory response in both BCG responders

and non-responders alike. Our findings implicated IFN-γ in directly contributing to the upregulation of

HLA-E by approximately 43% of recurring NMIBC tumor cells, and uniquely evident in all BCG-

unresponsive patients. Spatial analyses of tumor cells, stratified by their proximity to infiltrated NK and T

cells, unveiled a program of tumor activation (and potential dysregulation) and selective recruitment to

stromal-rich regions replete with immune effector and regulatory cells that are responding to the same

collection of chemotactic signals.

Distinctive to BCG-unresponsive tumor cells, localized areas or 'nests' exhibiting elevated

expression of HLA-E were enriched for expression of CXCL9 and CXCL12. This activation correlated

with the infiltration of NK cells, CD8 T cells, and Tregs. Notably, even within confined regions of tumor

tissue featuring a mix of HLA-E$^{DIM/NEG}$ and HLA-E$^{HIGH}$ tumor cells, a significant preference was observed

for the proximity of NK and CD8 T cells to HLA-E$^{HIGH}$ tumor cells. These findings hint at an evasion

mechanism wherein HLA-E$^{HIGH}$ tumor cells (responding to stroma-derived CXCL12) appear to play a role

in safeguarding HLA class I negative tumor cells. The expression of the NKG2A receptor also emerged as

a critical factor in categorizing the anti-tumor cytolytic functions of NK and CD8 T cells. However, our

observations revealed that NKG2A$^{+}$ NK and CD8 T cells also expressed several well-established secondary

checkpoint receptors, emphasizing the necessity for a comprehensive understanding of the inhibitory

landscape within patients' tumors to inform effective second-line immunotherapy strategies.

Upon removing NK and CD8 T cells from their suppressive microenvironment, these cells

demonstrated retained functional capacity. Both NK and CD8 T cells, enriched for NKG2A expression,

exhibited highly enhanced cytolytic reactivity to HLA-E$^{+}$ PD-L1$^{+}$ K562 tumors in the presence of

combination NKG2A/PD-L1 blockade. Collectively, these findings portray a landscape of immune

dysregulation prominently characterized by elevated tumor expression of HLA-E and PD-L1 in the context

of BCG-induced chronic activation and inflammation. Importantly, HLA-E$^{HIGH}$ tumor cells similarly

expressed higher levels of CD47, Nectin-2 (CD112), Nectin-4, PVR and at higher levels compared to PD-

L1 expression. Further, HLA-E$^{HIGH}$ tumor cells expressed higher levels of decay accelerating factor (DAF,

*CD55*) and CD59, which block C3 convertase and formation of the membrane attack complex, two critical

steps in driving complement-mediated inflammation and phagocytosis, and apoptosis, respectively. Finally,

the trophoblast-2 (Trop2) antigen was highly expressed on all recurring NMIBC tumor cells. These data

suggest future combination immunotherapy strategies may benefit from additional targets that are strongly

engaged in BCG-unresponsive NMIBC patients. Emphasizing this point, antibody-drug conjugates

targeting Trop2 (sacituzumab govitecan)[41] and Nectin4 (enfortumab vedotin)[42] have demonstrated

promising improvements in patient outcomes for metastatic triple-negative breast cancers and muscle-

invasive bladder cancer, respectively; the latter which, when combined with pembrolizumab (PD-1

blockade), has recently been approved FDA clearance in an all comer population for 1st line treatment of
metastatic urothelial carcinoma [43].

Intravesical administration of BCG is initiated as adjuvant immunotherapy, as complete tumor
resection is the initial diagnostic and therapeutic step. Thus, initiation of standard of care (SOC) BCG
immunotherapy presupposes inflammation without an active tumor to clear [44]. Treatment improvements for
NMIBC and BCG-resistant disease have lagged. This may be, in part, because few studies have attempted
to understand the relationship between the timing of tumor recurrence, reasoning for the recurrence, and
the state of the immune system at the time of recurrence. Poor dosing study designs and lack of
understanding of the mechanisms underlying a therapeutic response to intravesical BCG have led to a
significant gap in knowledge and benefit for patients with NMIBC compared to muscle-invasive or
metastatic disease.

Our findings suggest that all NMIBC patients at the time of tumor recurrence show signs of a
hallmark anti-tumor immune response dominantly driven by IFN-γ. BCG-unresponsive tumors see uniform
increases in chemotactic cytokines and inflammatory pathways that should otherwise function to suppress
tumor growth. Further, increased expression of inhibitory ligands on BCG-unresponsive tumor cells was
observed, suggesting that inflammatory stimuli had been prolonged and triggered feedback mechanisms
responsible for immune evasion. These results are seen on multiple levels, including bulk and single-cell
RNA sequencing, spatial sequencing, multiplex imaging, and mass cytometry. This implies that when
tumors recur, for reasons beyond the scope of this study, they are met by an overactive status of the immune
system ill-equipped to combat them.

Previous analyses profiling urine analytes between the BCG naïve and third dose timepoints
demonstrated that three doses of BCG induced an inflammatory response hundreds of times above baseline
levels of cytokines, including but not limited to IP-10, MIP-1β, IL-8, IL-6, and TNFα [45]. Importantly, the
third dose-response dwarfed the magnitude of the first dose-response, suggesting that repeated exposure
increases the magnitude of inflammation [45]. While these data did not profile out to the time of tumor
recurrence, they lend credence to the theory that all patients experience a ubiquitous and increasingly
powerful immune response to repeated doses of BCG. Importantly, our findings confirm these previous
observations. Further, they demonstrate that inflammation is sustained even throughout the six-week
treatment interruption at time of first cystoscopy and do not equilibrate to baseline levels.

Novel to our study is the spatially-resolved insights into potential consequences of IFN-γ
dysregulation and tumor cell and immune homing patterns. Preferential trafficking of *HLA-E*[HIGH] tumor
cells to immune-rich regions was seen in areas of high expression of *CCL19, CCL21, CXCL9/10/11*, and
CXCL12, and chemokines were detected in the urine early following start of therapy and were sustained
throughout therapy. Further, BCG unresponsive tumor-stroma proximity to cytolytic cells and Tregs was

increased by a median of 7-fold and 2.6-fold, respectively, when comparing *CXCL9/10/11*[HIGH] to

*CXCL9/10/11*[DIM/NEG] tumor nests.

Clinical efficacy in checkpoint blockade is dependent on reinvigorating effector cells expressing

high levels of the targeted checkpoint(s). Congruent with analogous studies, we found that NKG2A[+] NK

and CD8 T cells were enriched in BCG-unresponsive tumors and tumor-derived NK and CD8 T cell anti-

tumor functions were intact at the time of tumor recurrence. Additionally, co-culture experiments with

patient TILs and K562s with and without HLA-E and PD-L1 confirmed the ability of combination anti-

NKG2A plus anti-PD-L1 blockade to enhance the cytolytic function of stimulated TILs from patients at

time of tumor recurrence. This reinforces both the use of NKG2A to reinvigorate cytolytic effector

functions by NK and CD8 T cells, as well as the potential to exploit a critical mechanism of tumor

resistance, whereby chemokines lure in effector cells to take on exhausted phenotypes.

Recent clinical evidence emerging from interim results of the randomly controlled Phase II COAST

trial demonstrated significantly prolonged survival in non-small cell lung cancer (NSCLC) patients treated

with monalizumab combined with durvalumab compared to standard of care durvalumab alone for

treatment of unresectable stage 3 NSCLC [25]. This combination is being further evaluated in a larger planned

Phase III trial (PACIFIC-9, NCT05221840) following these promising early results. Collectively, our data

demonstrate that elevated tumor expression of HLA-E and PD-L1 and chronic activation of NKG2A-

expressing NK and CD8 T cells are hallmark features of resistance to BCG immunotherapy in NMI bladder

tumors and that integrating NKG2A-blockade into combination immunotherapy strategies may prove

particularly effective in settings of BCG-unresponsive, high-grade NMIBC. Based on the data generated

from this study, we recently designed a Phase 2 trial of durvalumab (MEDI4736) and monalizumab in

NMIBC. The Phase 2 ENHANCE (**E**levated **N**KG2A and **H**LA-E **A**mplify **N**K and **C**D8 T-cell **E**ngagers)

(clinicaltrials.gov ID: NCT06503614) will investigate the effects of monalizumab (anti-NKG2A Ab) plus

durvalumab (anti-PD-L1 Ab) for combination checkpoint blockade in high-grade BCG-unresponsive

NMIBC as a bladder-sparing immunotherapy strategy. In this trial, an exploratory objective is to assess the

relationship between baseline tumor HLA-E expression and clinical outcomes to facilitate development of

HLA-E as a potential biomarker for patient selection in future clinical trials intending to target NKG2A.

Our study has notable limitations. Our sample sizes, while being the largest NMIBC spatial

sequencing and single-cell RNA sequencing cohorts available, are small and present a potentially limited

view of the disease. Additionally, as BCG is administered as an adjuvant therapy, we are only afforded

access to bladder tissues after treatment if tumor recurs. Thus, it was not feasible to profile bladder tissues

in the absence of tumor following exposure to BCG. Future studies might benefit from a neoadjuvant

approach, where tumor tissue can be sampled and profiled for potential therapeutic responses to BCG.

Collectively, our analyses suggest that the current guidelines on immunotherapy for BCG-unresponsive NMIBC could be improved via a multi-cell-targeting immunotherapy approach focused on NK cells alongside T cells. Randomized trials from bladder and other tumor indications have shown that PD-1/PD-L1 stratification fails to predict response to immunotherapy with anti-PD-1/PD-L1 antibodies: IMvigor210, JAVELIN bladder 100, and CheckMate-275 (NCT02108652, NCT01772004, NCT02387996) all saw that PD-1/PD-L1 biomarker stratification alone did not effectively stratify response rates. In comparison, recent results using *KLRC1* (NKG2A) expression in the pre-treatment tumor significantly improved predicted anti-PD-L1 response rates in the IMvigor210 cohort, but only in the CD8[HIGH] *PDCD1* (PD-1)[HIGH] group [24]. In fact, where IHC stratification of PD-L1 expression has failed to predict immunotherapeutic responses, *KLRC1* stratification in IMvigor210 showed protective effects were restricted to the PD-L1 IC high group. In conclusion, while our analysis does not exclude the presence of alternative checkpoints, it lends evidence to the hypothesis that combination NK and T cell immunotherapeutic approaches may hold the key to improved outcomes for NMIBC patients.

**Methods**

**Patients and samples**

Patients at Mount Sinai Hospital (MSH) were enrolled in the study following Institutional Review Board (IRB) approval (protocol 10-1180). 10-1180 covers the use of patient tissues in a biorepository and allows for prospective collection of blood, urine, and tissue samples from enrolled patients. Formalin-fixed paraffin-embedded (FFPE) blocks from BCG patients were obtained retrospectively from the biorepository and prospectively for patients receiving treatment. For prospective patients, samples were collected on the day of surgery and throughout BCG immunotherapy. Due to IRB limits on the collection, blood and urine samples were taken at the first dose, third dose, and sixth dose of the induction cycle; at every follow-up cystoscopy; and at the third maintenance cycle dose. Tumor samples were taken at every possible timepoint. BCG naïve was defined as any patient who had yet to receive BCG, regardless of past treatment with other chemotherapies. BCG-unresponsive was defined as any patient with recurrent tumors following at least five of six induction doses of BCG at time of first evaluation.

**Sample processing**

Blood and urine samples from bladder cancer patients were processed to collect PBMCs, serum, and cell-free urine. Blood was spun down (4C, 2000rpm, 10 min) to isolate serum, and PBMCs were collected using Ficoll-Paque isolation. PBMCs were frozen down in a Mr. Frosty at -80 C for 24 hours and

stored at -160 C in 10% DMSO and 90% fetal bovine serum; PBMCs and cell-free urine were stored at -80

C.

Tumor tissues obtained from transurethral resections of bladder tumor (TURBT) were placed into

RPMI medium immediately after removal and transferred to the laboratory for additional processing.

Bladder and lymph nodes obtained from radical cystectomies were sent from the operating room directly

to the pathology suite after completion of the lymph node dissection. The bladder was bivalved; samples of

visible tumor were extracted and placed in RPMI medium, and the tumor was transferred to the laboratory

for additional processing.

Fresh tumor samples underwent a variety of different processing techniques based on the planned

experiment. Tumor used for spatial sequencing was placed in a 10 mm x 10 mm cryomold with optimal

cutting temperature (OCT) media and frozen down on a thermal block immersed in liquid nitrogen. Tumor

tissues were digested using tumor dissociation enzymes (Miltenyi, 130-095-929) and a GentleMACS

machine (program 37C_h_TDK_3) at 37C. Mechanically and enzymatically separated tissues were filtered

through a 70μM cell strainer and assessed on Countess II (ThermoFisher) for viability and cell numbers.

**In situ hybridization and targeted RNA sequencing**

FFPE sections from 40 retrospective NMIBC cases were obtained from the institutional

biorepository and used for targeted RNA sequencing. RNA was extracted from five and ten μm sections.

HTG EdgeSeq lysis buffer was added to lyse and permeabilize the samples. Nuclease protection probes

(NPPs) were added to the lysed samples and hybridized to the target mRNA. A nuclease was added to

digest non-hybridized mRNA and excess NPPs. The nuclease digestion reaction was finalized with a

termination solution followed by heat-mediated denaturation of the enzyme.

Each sample was used as a template for PCR reactions with specially designed primers. Each primer

contains a unique barcode that is used for sample identification and multiplexing. Samples were analyzed

simultaneously on an Illumina sequencing platform to prepare the library. All samples and controls were

quantified in triplicates. No template control (NTC) reactions were made for each master mix used during

the qPCR process to test the absence of a probe or qPCR contamination. Molecular-grade water was used

in place of a test sample in the NTC reactions using the same volume as the template.

Sufficient concentration of sample for library pooling, appropriate dilution for the library pool, and

volume of denaturation reagents to add to the library were determined by HTG library calculator. 2N NaOH

and heat (98C, 4 minutes) were used for library denaturation. The denatured library was loaded into the

well of the NextSeq sequencing cartridge. Sequencing was performed using an Illumina NextSeq sequencer.

The sequencing data on mRNA expression of target genes were imported into HTG EdgeSeq parser
software. HTG biostatistics department performed quality control analyses and normalized the data. Data
were returned from the sequencer as demultiplexed FASTQ files with four files per assay well. The HTG
EdgeSeq parser software aligned the FASTQ files to the probe list and collated the data.

Prior to any analyses, clinical data on all patients was gathered. The following items were collected
for all samples processed by HTG Molecular: patient date of birth; gender; age at BCG induction; BCG
status (unresponsive vs. naïve vs. exposed); date of sample collection; stage at collection; grade at
collection; recurrence date; time to recurrence; time to progression; BCG start date; BCG last exposure;
BCG induction cycle last dose; prior chemotherapy; cystectomy; date of cystectomy; stage at cystectomy.

Following clinical data collection, a gene set enrichment analysis (GSEA) was performed on the
targeted RNA sequencing data. Specifically, we used paired patient samples before and after BCG exposure
in the BCG recurrent patient population only. We used custom gene sets, as well as all Hallmark gene sets
from the Broad Institute's MSigDB, as inputs for the enrichment analysis [27]. Statistical significance was set
at $p < 0.05$, and gene sets found to be significant are listed in **Supplementary Table S1**. All gene sets found
to be statistically significant were evaluated for leading-edge genes, defined as the genes that contribute
most to the enrichment score and associated p-value.

The leading-edge genes from statistically significant gene sets in the GSEA were collated and used
to assess for group differences between the paired HTG patient samples. We performed these analyses
specifically on the BCG-recurrent cohort. Prior to any analyses, a Shapiro-Wilk test, chosen for suitability
in small sample sizes, was used to assess for normality [46]. All samples with $p < 0.05$ were considered not
normally distributed, and a Kruskal-Wallis test was performed to assess for group differences [47]. All other
samples were assessed using an independent T-test. Genes with statistically significant differences between
the BCG naïve and the BCG-unresponsive populations were then visualized on radar plots.

**Protein concentration measurement**
Cell-free urine supernatant and serum samples were randomized in a 96-well plate. Diluted samples
and positive and negative controls were incubated overnight with an incubation mix (incubation solution,
incubation stabilizer, A-probes, and B-probes) at 4°C. Samples were then incubated with an extension mix
(High purity water, PEA solution, PEA enzyme, PCR polymerase) for 5 min and placed on a thermal cycler.
Following the thermal cycler, samples were incubated with a detection mix (detection solution, High purity
water, detection enzyme, PCR polymerase) and transferred to a chip. Primers were loaded onto the chip,
and the chip was profiled using the Fluidigm IFC controller HX with the Fluidigm Biomark Reader. Data

were normalized using extension and interplate controls and a correction factor. The resulting data were
reported in normalized protein expression (NPX) units on a log2 scale.

**IFN-γ co-culture**

Cell lines and CD45- isolated primary tumor cells were incubated in media optimized for high
viability for 72 hours (RPMI-1640 supplemented with 20% fetal bovine serum). Tumor cells were expanded
until they were confluent in 2 T175 flasks. Following expansion, cells were culture in 100 ng / mL of IFN-
γ for a total of 24 hours in a 24 well plate. Following co-culture, cell lines were trypsinized (immortalized),
and primary tumors were gently removed from the solid phase by a cell scraper. Samples with and without
IFN-γ stimulation were measured for HLA-E and PD-L1 protein expression levels.

HLA-E and PD-L1 levels were assessed via FlowCytometry. Cells were stained in 4C FACS buffer
(phosphate-buffered saline (PBS) with 2% heat-inactivated FBS and EDTA 2 mM) for 30 minutes.
Subsequently, cells were washed in PBS, incubated for 20 minutes in a viability dye, washed again with
PBS, and resuspended in 2% paraformaldehyde. The experiment was performed in triplicate, with three
readouts per cell line per experimental condition. FlowCytometry acquisition was performed using an LRS-
Fortessa (BD Biosciences), and data were analyzed using the CytoBank software. When staining for HLA-
E, cells were first stained 20 minutes with HLA-E prior to staining with additional PD-L1. In CytoBank,
several gates were applied to generate the final dataset. A live/dead gate was applied, followed by a gate to
remove doublets and isolate singlets. Lastly, the data was arcsinh transformed prior to analysis.

Once the final dataset had been generated, statistical significance between unstimulated and
stimulated cell lines was assessed. An independent t-test was run with a significance cutoff of $p < 0.05$.

731
**Plasma and cell-free urine supernatant protein concentration analysis**

We used the OLINK Proteomics®- inflammation panel to profile cell-free urine supernatant.
Protein levels were visualized at BCG naïve, third dose, sixth dose, and first-cystoscopy timepoint,
comparing NPX values for all patients between the BCG naïve timepoint and the sixth dose timepoint.
Additionally, urine supernatants from a validation cohort of NMIBC patients treated with BCG were used
from the Aarhus University as previously described [13]. In order to determine the suitable statistical test, a
Shapiro-Wilk's test was used to assess for normality, and a Kruskal-Wallis test was used in every instance
in which one or both samples were not normally distributed. An independent T-test was used in the event
both samples were normally distributed. All statistically significant p values were then used to assess
adjusted p values via the Benjamini-Hochberg correction, with an alpha of 0.05. All statistically significant
genes between the BCG naïve and sixth induction dose time points are shown.

**Multiplex Immunohistochemistry:**

Sections of tumor for immunohistochemical (IHC) staining were taken at a thickness of 3-mm from formalin fixed paraffin-embedded (FFPE) blocks. H&E-stained sections were performed every 5 - 10 slices. The Ventana Discovery Ultra (Roche Diagnostics) machine was used to automatically bake, deparaffinize, and condition the slices. The RUO Discovery Universal (v21.00.0019) was used to perform chromagen IHC on sequential slices. Primary antibodies included (CD3, HLA-E, NKp46, and NKG2A) and were utilized for staining on NMIBC tumors. All slices followed the same protocol, which included a 60 minute incubation at 37°C; secondary antibodies using OmniMap HRP or NP DISCOVERY (Roche Diagnostics); signal detection using Discovery OmniMap. Nuclear counterstaining with Mayer's hematoxylin; and conversion to high-resolution images via the NanoZoomer S10 Digital slide scanner (Hamamatsu).

**Tumor and CD3$^+$ T cell expression and proximity analyses**

Prior to analysis, all slides were reviewed and regions of interest were annotated by a board-certified pathologist (R.B.). Tissue artifacts, including torn, folded, and damaged tissue, were excluded from any analyses. The HALO™ (Indica Labs, Inc.) digital image analysis platform, a semi-automated platform using machine learning to segment and label stained sections, was utilized for quantitative analyses. Halo AI™ and train-by-example classification, segmentation, and random forest classification was used to separate chromogenic stains and generate tabular data for downstream analysis. Slide features of each tumor, including cell lineages (tumor, stroma, and immune) and slide features (such as glass) were characterized. Glass was excluded from all downstream analyese. Classified cell classes were tabulated, and positive staining cells were stratified into expression tertiles (dim, moderate, and bright). Calibration for intensity expression was performed using tonsil tissues from healthy human tonsil. In addition to cell counts, total surface area (mm$^2$) was recorded to facilitate density calculations. Statistical analyses were performed using Python 3.8.1.

**Proximity Ligation Assay**

The NaveniFlex™ Proximity Ligation Assay (PLA) was performed according to the manufacturer's instructions using NaveniFlex Tissue MR ATTO647N (Navinci, Sweden). PLA was performed on sections of tumor taken at a thickness of 3-mm from formalin fixed paraffin-embedded (FFPE) blocks. H&E-stained reviewed with pathologist. Briefly, after deparaffinization, rehydration, and antigen retrieval, slides were blocked with Block NT blocking solution (Navinci, NT.1.100.01) for 60 min at 37 °C in a preheated humidity chamber and then incubated with mouse anti-HLA-E (clone: MEM-E/02, Abcam, 1:200) and rabbit anti-NKG2A (clone: EPR23737-127, Abcam, 1:2000) diluted in Diluent 1 NT solution (Navinci,

NB.1.100.02) overnight at 4°C. As negative controls, two (tonsillectomy) slides were incubated in antibody
diluent with only one primary antibody each. After washing, the slides were incubated with the PLA probes
corresponding to the primary antibodies using anti-mouse Navenibody M1 NT (Navinci, NB.1.100.06) and
anti-rabbit Navenibody R2 NT (NB.1.100.07) in Diluent 2 NT solution (Navinci, Navinci, NF.1.100.03)
for 60 min at 37 °C. Slides were then processed for ligation using reaction 1 reagent containing Buffer 1
NT (Navinci, NB.2.100.17) and Enzyme 1 NT (Navinci, NF.2.100.11) and subsequently reaction 2 reagent
containing Buffer 2 NT (NT.2.100.01) and Enzyme 2 NT (Navinci, NF.2.100.15) and incubated for 30 min
at 37 °C and 90 min at 37 °C, respectively. The slides were washed and incubated with post-block NT
reagent (Navinci, NF.1.100.01) in post-block supplement NT (Navinci, NT.2.100.04) for 30 min at 37 °C,
then processed for detection, counter-stained with DAPI, and mounted with coverslips using Prolong Gold
Antifade reagent (Invitrogen, P36930).

Images were captured at the Microscopy and Advanced Bioimaging Core of the Icahn School of Medicine
at Mount Sinai. A Leica DMi8 (Leica Microsystems, Germany) was equipped with a HC PL APO CS
10x/0.4 (Part Number 506285; Leica Microsystems, Germany) objective lens. A SpectraX fluorescence
illuminator (Lumencor, Oregon, USA) with multiple narrow-band light emitting diodes provided
illumination (LEDs used: 395/25nm for DAPI, 470/24nm for autofluorescence channel, and 640/30nm for
Navinci signal). The microscope and light source were controlled by LAS X software, version 3.7.5.24914
(Leica Microsystems, Germany). For fluorescence excitation, the following illumination settings were used:
a 395nm LED set to 50% (147mW at the SpectraX output port) for DAPI signal, captured at 20 milliseconds
of exposure; a 470nm LED set to 44% (86mW), captured at 70 milliseconds for an autofluorescence
channel; and a 640nm LED set to 100% (231mW), captured at 150 milliseconds for the target signal. A
multi-band pass filter set (Part Number 11525366; Leica Microsystems, Germany) was used to separate
fluorophore emission (Dichroic 415/490/570/660nm; Emission bands: 430/35, 515/40, 595/40,
720/100nm). Images were captured using a Leica DFC9000GT monochrome camera set to 12-bit depth,
2x2 (4-pixel) binning and "Low Noise" Gain mode. Images were captured in montage at 10% overlap,
merged ("Smooth" blending option) and then saved in the proprietary LIF ("Leica Image File") format
before being converted to IMS (Imaris) format for analysis.

Image analysis was performed using Imaris software 10.1.1 (Oxford Instruments, Concord MA). A surface
for the green background channel was created using the surface creation wizard with the following
parameters – Enable Region Of Interest = false, Enable Region Growing = false, Enable Tracking = false,
Enable Classify = false, Enable Shortest Distance = false, Enable Smooth = true, Surface Grain Size = 2.00
$\mu$m, Enable Eliminate Background = false, Active Threshold = true, Enable Automatic Threshold = false,

Manual Threshold Value = 1900, Active Threshold B = false. Masked channels were created by subtracting

the intensities within the green surface from the blue and far-red channels: the mask intensity was set to 0

for inside the green surface while the outside was set to the original channel's value. A new surface was

created using the surface creation wizard for the masked far-red channel using the following parameters –

Enable Region Of Interest = false, Enable Region Growing = false, Enable Tracking = false, Enable Classify

= false, Enable Shortest Distance = false, Enable Smooth = true, Surface Grain Size = 2.00 $\mu$m, Enable

Eliminate Background = false, Active Threshold = true, Enable Automatic Threshold = false, Manual

Threshold Value = 1600, Active Threshold B = false. An area filter was applied to this far-red surface to

remove surfaces whose area was larger than 50um². A DAPI surface was created using surface creation

wizard with the parameters – Enable Region Of Interest = false, Enable Region Growing = true, Enable

Tracking = false, Enable Classify = false, Enable Shortest Distance = true, Enable Smooth = true, Surface

Grain Size = 2.00 $\mu$m, Enable Eliminate Background = true, Diameter Of Largest Sphere = 7.50 $\mu$m, Active

Threshold = true, Enable Automatic Threshold = false, Manual Threshold Value = 10, Active Threshold B

= false, Region Growing Estimated Diameter = 6.00 $\mu$m, Region Growing Morphological Split = false,

Filter Seed Points = "Quality" above 60.0, Filter Surfaces = "Number of Voxels Img=1" between 10.0 and

500, to obtain individual nulcei within the field of views. Finally, a fourth surface was created by applying

the filter – Overlapped Area to Surfaces (Minimum = 0.050 um², Maximum = false) to obtain the DAPI

surfaces that were in contact with the far-red channel. The counts of total number of nuclei and nuclei

overlapping with the masked far-red surfaces were extracted from the statistics tab of Imaris.

**Single-cell RNA sequencing:**

**Data preprocessing**

Single-cell RNA sequencing (scRNA-seq) analysis was performed using Scanpy package in Python

and Seurat package in R. After loading, genes expressed in fewer than three cells were excluded from later

analyses. Cells with < 200 or > 8000 unique genes, as well as cells containing >20% mitochondrial gene

transcripts, were discarded. Cells then go through scrublet to detect doublet population. Subsequent data

normalization was performed by dividing feature counts for each cell by total counts for that cell, scaling

by a factor of 10,000, and natural log transformation. Next, scvi model was set up to correct batch-effects.

Then we performed scaling and principal component analysis (PCA) on the batch-corrected data. Using the

first 50 principal components (PCs), graph-based clustering and UMAP dimensionality reduction was

performed to reveal 16 cell clusters. Cluster-specific marker genes were identified using the FindAllMarkers

function, and marker genes with a natural-log fold change (FC) > 0.25 and expressed in >= 25% of cells

were used to annotate cell cluster identities based on known cell type markers [48, 49, 50]: CD8 T cells (*CD8A*,

*CD3E*), CD8- T cells (*CD3E*), tumor cells (*MUC1*, *KRT7/8/13/17/18*), B cells (*MS4A1*, *CD79A*),

monocytes/macrophages (*CD14*, *LYZ*, *FCGR3A*, *MS4A7*), T regulatory cells (*CD3E*, *FOXP3*), granulocytes

(*C1QA/B/C*, *ITGAM*, *FCGR3A*), NK (*NKG7*, *GNLY*, *KLRD1*, *KLRF1*), fibroblasts (*COL1A1*, *SPARC*),

epithelium (*EMP1*), DC (*FCER1A*, *CST3*, *ITGAX*), plasma cells (*MZB1*, *JCHAIN*, *IGHG3*), and CD8 TRM

cells (*CD3E*, *CD8A*, *KLRD1*). Clusters 22, 24, 25, and 26 containing 224, 142, 141, and 101 cells,

respectively, were ultimately excluded from further analyses due to unclear cell identities. All scRNAseq

analyses were performed using distinct samples without repeated measurements.

**Tumor subclustering analysis**

We performed subclustering analysis on 18,520 bladder cancer (Blca) cells. Preprocessing steps

for the Blca subset included normalization, scaling, and PCA analysis. We then applied Canonical

Correlation Analysis (CCA) integration using the IntegrateLayers() function (dimensional reduction for

correction = pca) in Seurat v5 to correct batch effects [51]. Leiden clustering (resolution = 0.4) was applied to

the shared nearest neighbors (SNN) (dims = 1:10). This resulted in the identification of 9 heterogeneous

tumor subclusters, distinctly separated on the UMAP plot. Each subcluster was profiled by the expression

of tumor marker genes (EPCAM, UPK2) and cytokines (CXCL1, CXCL2, CXCL3, IFNGR1, etc.).

Subcluster B1 was removed from further analysis as it was identified as normal bladder cells with low

EPCAM (tumor-related marker genes) , and subcluster B8 was removed due to high PTPRC expression,

indicating a high presence of immune cells (CD45+). Differentially expressed gene (DEG) analysis (MAST,

R version 1.2.1) [52] and enrichment pathway analysis [53] further characterized the subclusters, highlighting

their distinct biological profiles (Figure 2).

**NicheNet Analysis**

We used the NicheNet R package to infer potential ligand-receptor interactions between NK and T

cells and tumor cells, as well as potential regulatory effects in tumor gene expression as a result of these

interactions [33]. We added known ligand-receptor interactions between ligands IgG1 and IgG3 and the

receptor CD16 (weight = 1) to NicheNet's default ligand-receptor interactions database, and also integrated

the Harmonizome Pathway Commons Protein-Protein Interactions database:

(https://maayanlab.cloud/Harmonizome/dataset/Pathway+Commons+Protein-Protein+Interactions)  with

NicheNet's default signaling network (weight = 1) [54, 55]. We used the modified ligand-receptor interaction

and signaling networks for subsequent analyses. We defined the NK and T cell populations as senders and

tumor cells as receivers and derived a list of 92 receptors expressed in the receiver population by

intersecting the list of receptors from the ligand-receptor interaction network with genes expressed in at

least 10% of tumor cells in the RNA assay. Fifty potential ligands expressed in greater than 10% of the

sender NK and T cells were then derived based on their interaction with the list of 92 receptors. We then

calculated activity scores for each ligand based on its potential to affect the expression of Hallmark

interferon gamma response genes [27,56]. Statistical analyses including Wilcoxon rank-sum tests and Pearson

and Spearman correlations were performed using R v4.0.3.

**Differential gene expression analysis**

Spatial transcriptomic data were generated by Visium Spatial Gene Expression which is a spatial

transcriptomics solution by 10x Genomics. We obtained eight spatial sections consisting of four preBCG

and four postBCG samples. Sequencing data was aligned on GRCh38 and quantified using the Space

Ranger Software Suite (version 1.0, 10x Genomics). We processed the ST data to characterize major cell

types and their spatial patterns using Giotto workflow [57]. Per section, Low quality spots (number of

expressed genes > 100), and lowly expressed genes (expressed in < 10 spots) were removed. The raw counts

were log2(x+1) transformed, followed by adjusting for batch effects and spot wise number of features using

Pearson residuals approach by Lause et al [58]. The spatial network of nearest spots by Delaunay triangulation

was calculated to identify spatially co-expressed genes in neighboring spots via BinSpect-kmeans algorithm

[57], and distinct spatial domains by Hidden Markov Random Field (HMRF) model [57]. Then, different sections

were combined into a pan-section transcriptome by Harmony  to identify pan-section clusters [59]. The pan-

section clusters will be identified by Leiden clustering [57] to detect transcriptionally similar spots shared

across different sections. Subsequently, we identified 12 spatial clusters to profile the spatial transcriptomic

landscape. Differentially expressed genes (DEGs) between KLRC1$^+$ versus KLRC1$^-$ cells were found using

the FindMarkers function from the Seurat R package, where cells with KLRC1 expression greater than 0

in the RNA assay were labeled as KLRC1$^+$.

**Identification of HLA-E high/low tumor subsets in spatial transcriptome**

As HLA-E expressions are not specific to tumors and HLA-E high/low tumor markers have not been

established, we performed a customized cell type presence inference workflow by utilizing Cell Atlas

Reconstruction from Spatially mapped Data (CARD) method [60]. CARD is a computational approach that

reconstructs cell-type-specific expression profiles from spatially mapped transcriptomic data, facilitating

the estimation of cell-type proportions within tissue sections. First, we generated reference single-cell

transcriptomes of HLA-E$^{high}$ and HLA-E$^{low}$ tumor subsets by explicitly labeling tumors with HLA-E

expressions (high: HLA-E > 0, low: HLA-E=0). These tumor labels were utilized as the inputs to calculate

the relative abundances of the tumor subsets on the spatial voxels by CARD.

To further scrutinize the detection of HLA-E high tumors in the spatial transcriptome (ST) data, we

applied additional thresholds to require voxels with i) high HLA-E expressions, and ii) highly abundant

HLA-E high tumor/stromal cells as inferred by CARD [60]. Per section, high HLA-E expression voxels were

identified as those with HLA-E expression values greater than zero (SCT normalized assay) and a z-score

of inferred HLA-E$^{high}$ tumor/stromal cell abundance $> 1.645$ (90% confidence). Conversely, low HLA-E

expression voxels were identified as those with an absence of HLA-E expression (HLA-E expression equal

to zero) and a z-score of inferred HLA-E$^{low}$ tumor/stromal cell abundance $> 1.645$ (90% confidence).

Similarly, we detected high confidence voxels with NK cells, CD8 T-cells and Treg cells by further applying

the abundance z-score $> 1.645$ for further analyses.

**Proximity analysis between different cell types**

We evaluated the proximity between different immune subsets (NK, CD8+ T, and regulatory T (Treg)

cells) and HLA-E$^{high/low}$ tumor cells to observe if distinct cytotoxic immune micro-environments are present,

conditioned on tumoral HLA-E expression status. Upon identifying high confidence voxels with HLA-

E$^{high/low}$ tumor/stromal cells and the immune cell types, we calculated the pairwise Euclidean distances

between the tumor/stromal cells and each immune cell type. To retrieve actual distance between voxels, the

matrix was refined using the computeCellDistance function from the CellChat v2 library in R [61]. For each

voxel with HLA-E$^{high/low}$ tumor/stromal cell presence, its overall distance to an immune cell type was

calculated as the mean distance over the 10 nearest neighboring voxels with the respective immune cell

presence. These distances were summarized across each section by taking the average over all HLA-E$^{high/low}$

tumor/stroma voxels, yielding the overall proximity with each immune cell type per section. These overall

proximity values were compared between pre-BCG and post-BCG groups by Wilcoxon Rank-sum test

(**Figs. 5** and **6**).

**In-depth analysis of cell type enrichment across spatial clusters**

We tested if each pan-section cluster shows enriched presence of distinct cell populations and subsets

as characterized by the scRNA-seq. To this end, for each section, we leveraged the inferred abundances of

different cell types in the spatial voxels by CARD, and tested if a cell type has significantly higher

abundances in each pan-section cluster than the other voxels by Wilcoxon Rank-sum test. Then, the overall

enrichments of each cell type within pre-BCG or post-BCG group were summarized by combining the

Wilcoxon test p-value across the sections via aggregated Cauchy association test (ACAT). The enriched

cell types per pan-section cluster in pre-BCG or post-BCG group were called with ACAT summarized FDR

$< 0.05$ and number of sections with significant enrichments (Wilcoxon FDR $< 0.05$) $> 1$ (Figure 4.3).

**Mass cytometry antibody preparation and staining**

PBMCs from BCG-treated bladder cancer patients at the time of tumor recurrence were isolated
using Ficoll-Paque and resuspended in cell medium (RPMI-1640 medium supplemented with 10% heat-
inactivated FBS, 1% Penicillin, 1% Streptomycin and 1% L-glutamine). PBMCs were conjugated to
antibodies purchased from Fluidigm using the Maxpar X8 and MCP9 labeling kits. Platinum barcodes were
prepared as previously described. All antibodies were titrated prior to conjugation.

Prior to mass cytometry staining, all cells were incubated for 20 minutes at 37C in RPMI cell
medium (described above) and IdU (Fluidigm, t#201127), Rh103 (Fluidigm, #201103A). Following the
incubation, cells were centrifuged, washed with PBS and 0.2% bovine serum albumin (BSA), and incubated
for 3 minutes on ice with an Fc-blocking reagent. Samples were washed again with PBS and 0.2% BSA;
barcoded samples were pooled together and washed; finally, samples were stained with extracellular
antibodies for 30 minutes on ice and in PBS and 0.2% BSA.

When staining four samples, cells were single-barcoded using 194Pt, 195Pt, 196Pt or 198Pt. When
staining five or six samples, cells were stained with combination of barcodes. Cells were again washed
using PBS 0.2% BSA; barcoded samples were pooled together; samples were washed again and stained
with extracellular antibodies for 30 minutes on ice in PBS 0.2% BSA. Cells were co-cultured in the presence
of either WT or HLA-E-transduced K562 cells for a total of 6 hours.

Samples were washed with PBS 0.2% BSA and resuspended in Fixation/Perm buffer (Invitrogen,
#00-5523-00) for 30 minutes on ice. Cells were centrifuged and washed with Maxpar Barcode Perm Buffer
(Fluidigm, #201057) and barcoded using the Cell-ID 20-Plex Pd Barcoding kit (Fluidigm, #201060).
Barcoded samples were washed with permeabilization buffer (Invitrogen, #00-5523-00) and pooled.
Intracellular staining was then performed in permeabilization buffer with Heparin at a concentration of
100U/mL for 30 minutes on ice. Stained cells were washed with permeabilization buffer and resuspended
in PBS with PFA 2.4%, saponin 0.08%, Osmium tetroxide 0.075nM and Ir 0.125uM (Fluidigm, #201192A).
Finally, samples were washed and resuspended in PBS 0.2% BSA and data were acquired within four days,
or frozen in FBS/DMSO 90/10. The antibody panel in Supplemental Table S4 was used to stain bladder
cancer patient samples.

**Mass cytometry sample acquisition and processing:**
Prior to acquisition, samples were washed with cell staining buffer and acquisition solution
(Standard Biotools). Following washing, samples were resuspended in acquisition solution (1 million cells
/ 1 mL) containing a 1:20 dilution of EQ normalization beads. Data were then acquired using the Standard
Biotools Helios mass cytometer with a wide bore injector configuration at an acquisition speed of < 400
cells per second. The output files were normalized and concatenated using Standard Biotools' CyTOF
software, and outputted as FCS files.

The Mount Sinai Human Immune Monitoring Core's (HIMC) pipeline for processing and cleaning

was used to clean the resulting FCS files. Aberrant acquisition time-windows and low DNA intensity events

were stripped out by the sample preprocessing pipeline. Samples were then demultiplexed via the cosine

similarity of the Palladium barcoding channel on a cell-by-cell basis to every possible barcode used in a

batch. Once the cell-barcode labeling has been established, the signal-to-noise (SNR) ratio was calculated

by taking the difference between the highest and second highest similarity scores. Cells with low SNR

ratios were flagged as multiplets and removed. Finally, acquired multiplets are removed based on the

Gaussian parameters residual and offset acquired by the Helios mass cytometer.

**Data processing and analysis:**

Data were uploaded onto Cytobank and processed for downstream analyses. Several gates were

applied: a live dead gate and a doublets gate in sequential order. All data were arc-sinh transformed, and

no batch corrections were performed given all samples were run in a single batch. NK cells and CD8 T cell

subsets were identified via manual gating assignment. Files were downloaded onto Mount Sinai's

supercomputing cluster, Minerva, concatenated into a single object, and clinical data were assigned to each

sample. A phenograph analysis was performed to cluster the cellular data. Wilcoxon tests were run for

statistical significance and the Benjamini-Hochburg correction was applied.

**Tumor Invading Lymphocyte Expansion**

TILs were expanded by seeding single cells tumor dissociate at $10^5$ cells/well in 48-well Costar®

flat-bottom plates (Corning Inc., NY) in complete TIL medium, consisting of RPMI 1640 with L-glutamine

(Corning Inc., NY), 10% human AB serum (MiliporeSigma, MO), 1% nonessential amino acids (Thermo

Fisher Scientific, MA), 1% sodium pyruvate (Thermo Fisher Scientific, MA), 1% penicillin/streptomycin

(Thermo Fisher Scientific, MA), and 3000 IU/ml interleukin-2 (Proleukin®, kindly provided by Clinigen).

Cells were stimulated using 25 μl ImmunoCult™ Human CD3/CD28/CD2 T Cell Activator (STEMCELL

Technologies, Vancouver, BC) and seeded on 1.5 x $10^6$ feeder cells, which were derived from healthy

control PBMCs (New York Blood Center, NY) and irradiated at 50 Gy. TILs were incubated at 37°C, 5%

$CO_2$ and maintained by replacement with fresh complete TIL medium every 2 days. After three weeks,

expanded TIL lines were frozen in 10% DMSO (MiliporeSigma, MO), 90% FBS (MiliporeSigma, MO)

and stored in $LN_2$.

1007

**IFN-γ Stimulation of K562 tumors**

Two separate rounds of stimulation were performed on K562 tumors after reaching steady-state
expansion. First, K562 tumors were co-cultured at a concentration of 0.5 X $10^6$ viable cells / mL with 200
and 400 ng / mL of IFN-γ in R10 media in order to assess the induction of HLA-E and PD-L1 with high
doses of IFN-γ. At every 24-hour interval, cell concentration and viability were assessed using a
hemocytometer, and K562 tumors were split to a concentration of 0.5 X $10^6$ live cells / mL. At 72 hours of
incubation, cells were spun at 1650 RPM, washed with FACS buffer (PBS, 5% FBS, 0.2% EDTA),
resuspended in 5 mL of FACS buffer, and assessed for concentration and viability. 0.5 X $10^6$ live cells were
removed, and spun to remove the FACS buffer. K562 tumors were first incubated with zombie near infrared
(NIR) at a dilution of 1 in 1000 for 20 minutes at room temperature. Following the zombie stain, cells were
washed in 1 mL of FACS buffer, and incubated with PD-L1 (PE; Biolegend cat #124308), and HLA-E (per-
CP CY5.5; Biolegend cat #342609) antibodies in a 1:25 dilution cocktail for 30 minutes on ice in the dark.
Cells were washed again in FACS buffer and suspended in 400μL of fixative buffer (2% paraformaldehyde
in PBS). HLA-E and PD-L1 levels were then assessed using flow cytometry.
1022    
**<u>TIL Co-culture with K562s Expressing Checkpoints</u>**
For co-culture and surface stain of PD-L1 and HLA-E, three separate K562 tumor lines of K562 cells
were generated using 400 ng/ml of IFN-γ for 72 hours, as described by the protocol listed above: WT, E+,
and E+/PD-L1+ lines. Tumor-infiltrating lymphocytes (TILs) from four BCG-unresponsive NMIBC
patients which had been expanded using the protocol listed above were thawed and seeded at a
concentration of 1 X $10^6$ live cells / mL in a stimulatory media consisting of ImmunoCult, CD3/CD28
tetramer (25 uL/mL), IL-2 (10 IU/mL), IL-7 (10 ng/mL), and IL-15 (10 ng/mL). Cells were seeded and
plated in a 96 round-bottom plate. At days three, five, seven, and nine, 50% of the media was aspirated and
replaced with fresh media at twice the initial concentration of cytokines, without CD3/28 tetramer. At day
10, TGF-β (Immunocult, IL-2 (10IU/mL), IL-7 (10ng/mL), IL-15 (10ng/mL), CD3/CD28 tetramer
(25μL/mL), TGF-β (5ng/mL)) was provided to stimulate expression of NKG2A on the TILS.
At day 13, the stimulation was complete, and the TILs were co-cultured with combinations of the K562
cells expressing forms of PD-L1 and HLA-E. In total, 12 experimental co-culture conditions were
conducted for each patients' TIL sample (n=4): two control wells (ex-vivo, and stimulated with a CD3/28
spike); with K562 WT cells, HLA-E+ cells, or HLA-E+ PD-L1+ cells without antibody; TILs in the presence
of HLA-E+ or HLA-E+PD-L1+ K562s with durvalumab alone, or both durvalumab plus monalizumab; and
lastly, TILs and WT K562s in the presence of monalizumab and durvalumab.
TILs were cultured in TIL medium, consisting of RPMI 1640 with L-glutamine (Corning Inc., NY),
10% human AB serum (MiliporeSigma, MO), 1% nonessential amino acids (Thermo Fisher Scientific,
MA), 1% sodium pyruvate (Thermo Fisher Scientific, MA), 1% penicillin, streptomycin (Thermo Fisher

Scientific, MA). Cultures were performed in 96-well u-bottom plates (Corning Inc., NY). TILs were added

to each well at a concentration of at 3 x$10^5$ cells/condition in TIL medium. K562s were resuspended in TIL

medium and added at 5 x $10^4$ singe-cell equivalents/condition. Anti-CD28 (BioLegend, CA) was added to

each well at 1 μg/ml and anti-CD107a-BV785 (BioLegend, CA) was added to each well at 0.3 μg/ml.

Monalizumab was added to the TILs a final concentration of 10 μg/ml, and durvalumab was added to the

K562s for a final concentration of 10 μg/ml. TILs and K562s were cultured separately for 20 minutes at

37°C prior to co-culture. The K562s and TILs were then combined, thoroughly resuspended, and gently

spun for 5 seconds before being returned to the incubator. Plates were incubated at 37°C and 5% $CO_2$, and

after 1 hour of incubation, 0.5X Brefeldin (BioLegend, CA) and 0.5X Monensin (BioLegend, CA) was

added to each tube for a final volume of 500 μl, followed by an additional 15 hours of incubation. At hour

16 plates were spun and supernatant was removed; all samples were washed with 200μLof PBS. Samples

were resuspended in 50μLof zombie near infrared (NIR) at a 1:3000 dilition, and incubated for 20 minutes

in the dark at room temperature. Samples were washed again, spun, supernatant was removed, and

incubated in 50μLof TruStain FCX, diluted 1:200, for 20 minutes in the dark at 4°C. 50μLof the surface

antibody master mix was added at 20 minutes, and returned to the fridge to incubate in the dark at 4°C for

30 more minutes. Plates were washed twice more in FACS buffer, and were suspended in 0.2% FBS FACS

buffer for storage until intracellular staining.

12 hours prior to acquisition by CyTOF, intracellular staining was performed for IFN-γ. 100 μL of

intracellular staining permeabilization wash buffer (BioLegend) was added to each well, the plates were

spun, and supernatant discarded. This was performed 2 additional times with 200 μL of permeabilization

wash buffer. 50 μL of the intracellular staining antibody mix was added to each well, and the plates were

incubated for 20 minutes at 4°C. The plates were washed in 0.2% FBS FACS buffer and resuspended for

storage in 0.2% FBS FACS buffer.

1066

1067 **Data Availability Statement**: data were generated by the authors and have been uploaded to the Gene

1068 Expression Omnibus (GSE276014 and GSE276015) and will be made publicly available upon publication

1069 of this manuscript.

1070

**ACKNOWLEDGMENTS**

We thank Mary Anne O'Donnell (Precision Immunology Institute, Icahn School of Medicine at Mount

Sinai) for critically reviewing the manuscript; Deepta Bhattacharya ´(University of Arizona) for kindly

providing HLA-E$^+$ K562 tumors. We acknowledge the expertise and assistance of the Dean's Flow

Cytometry Center of Research Excellence at Mount Sinai. The A.H. and J.P. labs were supported by funding

from P30CA196521, R01 CA269954-01, R21 CA274148, BCAN No. 961726, and R21AI130760A. The

N.B. lab was supported by funding from the Department of Defense Peer Review Cancer Research program

Translational team Award No. W81XWH1910269 and from the Parker Institute for Cancer Immunotherapy

No.AGR-11611SOW1.

1080

**AUTHOR CONTRIBUTIONS**

D.R., H.Y., Y.A.W., C.B., J.D., B.S., M.G., N.B., J.P.S., and A.H. conceived the project and experiments,

analyzed the data and wrote the manuscript. J.P.S., R.M., P.W., M.G., and R.B. provided access to the

human samples. Additionally, L.D. provided analyses on data derived from human samples at Aarhus

University Hospital. D.R., H.Y., Y.A.W., C.B., A.D., S.B.,W-M.S., J.D., B.S., D-F.R., F.P. E.M., A.T., J-

A.C-F., B.D.H., T.S., S.V.L., S.H., J.K., performed the experiments or analyzed data. A.M.F, M.T., J.Z.,

K.G.B, L.W, R.P.S, S.S, Y-C.W, and Y.A.W. collected the RNA sequencing data. D.G., G.K., R.M.d-R,

B.L., S.K-S. acquired sample data using mass cytometry and Olink proteomics. T.H.T, M.G-B, R.B., E.H-

S, E.M.M, C.B., R.F-G, M.S, H.R, A.N., G.D., and J.J.B-C. performed the imaging experiments. S.S., E.C-

T., D.P., and S.H. provided intellectual input.

1091

1092    **DECLARATION OF INTERESTS**

1100

1101    **LEGENDS TO FIGURES**

1102

1103    **Fig. 1: Chronic activation within the tumor microenvironment of BCG-unresponsive tumors**

1104    **promotes adaptive resistance**.

1105    **A,** Targeted mRNA gene-set enrichment analysis showing statistically significant differences between

1106    BCG-naïve (N=20) and BCG-unresponsive cases (N=20). All gene sets are significant at $p < 0.05$ by

1107    Kruskal-Wallis or independent two-sided t-test.

**B,** Longitudinal protein analysis of urine supernatants across four timepoints in BCG-naïve patients with NMIBC (N=27) with comparisons between BCG-naïve and 6th induction dose timepoints (* p<0.05, ** p<0.001).

**C,** An additional validation cohort of patients with NMIBC receiving BCG therapy at Aarhus University (N=66) showing BCG-naïve vs BCG-exposed timepoints in urine samples (p-values in both cohorts assessed via independent two-sided t-test or Kruskal-Wallis with Benjamini-Hochberg correction for multiple comparisons).

**D,** Differential protein expression analysis of urine before (top row) and after (bottom row) BCG exposure and sampled before (left-side) and after (right-side) transurethral resection of bladder tumors. Number of patients included in analyses: (Top-left) BCG-naïve when tumor present, non-responders, N= 47; responders, N= 60. (Top-right) BCG-naïve after tumor resection, non-responders, N= 5; responders, N= 6. (Bottom-left) BCG-exposed when tumor present, non-responders, N= 49; responders, N= 34. (Bottom-right) BCG-exposed after tumor-resection, non-responders, N= 19; responders, N= 26. P-values assessed with two-sided Wilcoxon rank sum. Proteins listed in orange text indicate unadjusted p-values <0.05. BCG-unresponsive NMIBC was defined as high-grade recurrence within two years after receiving BCG therapy.

**E,** Median fluorescence intensity (MFI) for PD-L1 and HLA-E staining on primary CD45- cells from patients with NMIBC and immortalized bladder tumor lines ex vivo or after 24 hours of incubation with recombinant human (rh) IFN-g stimulation (all comparisons, p<0.05). All stimulation experiments were performed in triplicate.

**Fig: 2: HLA-E tumor expression is highest when near tumor-infiltrating NKG2A+ NK and CD8 T cells**.

**A,** Representative multiplexed immunofluorescence (IF) analysis by PhenoCycler™ (also known as CODEX) of bladder tumor sections from one patient with BCG-unresponsive NMIBC. Representative staining of the entire section is shown for presence of stroma (vimentin, blue) and tumor (S100A4, yellow). Scale bar indicates 2mm.

**B,** Magnified inset from Panel A, highlighting tumor HLA-E expression of and use of digital pathology software to identify NKp46+ NK and CD8 T cells with and without co-expression of NKG2A +/- PD-1. Left image scale bar indicates 400μm and right image scale bar indicates 50μm.

**C,** Magnified inset from Panel B, right image, highlighting individual cells expressing CD8, PD-1, and NKG2A. Scale bars indicate 20μm.

**D,** Representative multiplexed immunohistochemistry (IHC) of bladder tumor from one patient with BCG-

unresponsive NMIBC measuring expression of pan-cytokeratin (pan-CK), HLA-E, CD3, and NKG2A.

Scale bar indicates 100μm.

**E,** Representative digital pathology analyses identifying tumor (red) and adjacent/non-tumor (green) tissue

along with exposed areas of glass (yellow) to be excluded from subsequent analyses. Scale bar indicates

1mm.

**F,** Density map highlighting regions of high HLA-E tumor expression. Blue-red color scale indicates HLA-

E expression intensity. Scale bar indicates 1mm.

**G,** Summary analysis of frequency of tumor cells that are HLA-E-bright (green) and HLA-E-dim/negative

(dim/neg, yellow) in BCG-naïve (N=17) and BCG-unresponsive (unresp., N=24) NMIBC tumors.

**H,** Representative digital pathology analysis on one BCG-naïve (top row) and one BCG-unresponsive

(bottom row) NMIBC tumor section highlighting nuclear expression of DAPI (blue) and identification of

CD3$^+$NKG2A$^+$ CD8 T cells (black), CD3- NKG2A$^+$ NK cells (red), and tumor cells with bright (green) or

dim/negative (yellow) expression of HLA-E.

**I,** Proximity analysis (N=41) measuring cell distance (0-150μm) from HLA-E-bright and HLA-E-

dim/negative tumors  by NKG2A+ NK cells (left side) and NKG2A+ T cells (right side). ****, p<0.00001.

P-values were assessed via independent two-sided t-test.

**J,** Representative proximity ligation assay (PLA) using immunofluorescence to profile interactions between

HLA-E and NKG2A in one BCG-naïve (top row) and one BCG-unresponsive (bottom row) patient with

NMIBC.

**K,** Summary analysis of PLA measuring interactions between HLA-E and NKG2A. P-values were assessed

via independent two-sided t-test.

1161

1162

**Fig. 3: Single-cell RNA sequencing analysis of NKG2A$^+$ NK and CD8 T cells from bladder tumors**

**reveals strong antitumor potential but with co-expression of inhibitory receptors**.

**A,** UMAP visualization of cell lineages from single-cell RNA sequencing analysis of urothelial tumor samples (N=9,

65,324 cells).

**B,** Heatmap summary of genes most differentially expressed on each cell lineage identified.

**C,** Differentially expressed gene (DEG) analysis of bladder tumor-derived NK cells (left), CD8 T cells

(center), and proliferating T cell cycle cells (right) when stratified by high versus dim/negative *KLRC1*

expression.

**D** and **E,** UMAP visualizations of bladder tumor-derived NK cells from unsupervised clustering revealing

(**D**) five clusters or (**E**) expression of *NCAM1*/CD56 for annotation of CD56BRIGHT and CD56DIM NK cells

(N=2,580 cells).

**F,** Pathway analysis of Hallmark gene networks that are significantly differentially expressed (at least

p<0.05) on bladder tumor-derived CD56BRIGHT and CD56DIM NK cells. P-values were determined using a

two-sided t-test.

**G** and **H,** UMAP visualization of bladder tumor-derived NK cells (**G**) clustered according to Groups 1-6

defined by Netskar et al. [30] and (**H**) highlighting expression and distribution of representative tissue

residency genes.

**I,** Heatmap summary showing average expression for genes associated with tissue residency, chemokines,

cytokines, and their receptors on bladder tumor-derived Group 1-6 NK cells (top heatmap) and CD8 T cells

and T cell cycle cells when stratified according to *KLRC1* expression (bottom heatmap).

1183

**Fig. 4: Single-cell RNA sequencing analysis of bladder tumor cells reveals functional differences that**

**segregate the tumor microenvironment**.

**A,** UMAP visualization of bladder tumor cells from unsupervised clustering revealing seven clusters (B1-

B7).

**B,** UMAP visualization of HLA-E expression defined by tertiles, highlighting the top tertile (red dots) as

HLA-E-bright (N=5,461) and bottom tertile (blue dots) as HLA-E-dim/negative (N=9,256) tumor cells.

**C,** Pathway analysis of Hallmark gene networks that are significantly differentially expressed (at least

p<0.05) on HLA-E-bright and HLA-E-dim/negative bladder tumor cells. P-values were determined using

a two-sided t-test.

**D,** Bubble plot showing targeted gene expression across bladder tumor clusters B1-B7 and stratified by

*HLA-E*HIGH or *HLA-E*LOW tumor clusters. Size of bubble indicates percent of cluster or group and color

indicates average gene expression.

**E,** Scatterplot highlighting pseudobulking of bladder tumor cells and expression of *HLA-E* and *ACKR3*

(CXCR7). P-value was determined using a two-sided correlation.

**F,** Stacked bar plot showing the frequency of bladder tumor clusters, B1-B7 that are identified in BCG-

naïve (green) and BCG-unresponsive (grey) NMIBC tumors.

**G,** Heatmap summary of average gene expression of select chemokines, chemokine receptors, amphiregulin

(*AREG*) and *EGFR* across all major clusters identified by scRNAseq of bladder tumors (N=9, 65,324 cells).

**H,** Ligand-receptor interaction analysis of *KLRC1*HIGH cells (NK cells, CD8 T cells, T cell cycle) and *HLA-*

*E*-bright tumor cells (schematic diagram at top) showing the top 12 ranked ligands by KLRC1-expressing

cells and cognate receptors expressed by *HLA-E*[HIGH] tumor cells. Ligand-receptor pairs were selected by

rank-ordering and using cut-off weight of 0.2.

**I,** Circos plot showing top ranked ligands by KLRC1-expressing cells at the bottom and significant genes

expressed by *HLA-E*[HIGH] tumor cells as a result of ligand-receptor interactions highlighting putative effects

of the HLA-E/NKG2A axis (top) or the *IFNG*-mediated (middle) or *AREG*-mediated (bottom) resistance

signatures.

1210

**Fig. 5: Spatial analysis of NMIBC tumors reveals an organization within the TME**.

**A** and **B,** UMAP visualization of spatial transcriptomics sequencing (ST-seq) analysis of BCG-naïve (N=4)

and BCG-unresponsive (N=4) NMIBC tumors identifying 12 ST-clusters and their distributions across

NMIBC specimens.

**C**, Bubble plot showing ST-clusters 1-11 (rows) and their relative composition across immune, stromal,

and tumor subtypes (columns) as defined by scRNAseq analysis. Size of bubble indicates its relative

enrichment and shading of surrounding boxes indicate significance by FDR (corrected p values).

**D,** Representative ST-seq images showing the distribution of ST-clusters 1-11 in one BCG-naïve (left) and

one BCG-unresponsive (right) NMIBC tumor specimens.

**E,** Summary histogram comparing relative enrichment of each ST-cluster in BCG-naïve (grey, N=4) and

BCG-unresponsive (green, N=4) NMIBC tumor specimens.

**F,** Schematic diagram illustrating deconvolution of NK cells and tumor cells across 10x Visium spots.

Cutoff of 20% of spot's composition was used for defining a true signal.

**G** and **H,** Representative ST-seq images from one BCG-naïve (left) and one BCG-unresponsive (right)

NMIBC tumor specimens showing proximity analyses of *HLA-E*[LOW] (black circles) and *HLA-E*[HIGH] (red

circles) tumor cells to (**G**) NK/CD8 T cells (blue spots) or (H) NK cells, CD8 T cells, and Tregs alone or

in combination (colored spots).

**I,** Summary comparisons of proximity of *HLA-E*[HIGH] (red bars) and *HLA-E*[LOW] (grey bars) tumor cells to

NK cells (left), Tregs (middle) and CD8 T cells (right). P-values were determined using two-sided

Wilcoxon test.

1231

**Fig. 6: Recruitment of *HLA-E*[HIGH] tumor cells to stromal regions replete with immune infiltration**

**correlates with BCG-unresponsive NMIBC**.

**A-C,** Representative ST-seq images from one BCG-unresponsive NMIBC tumor highlighting (**A**)

proximity of NK cells, CD8 T cells, and Tregs to tumor and stromal cells according to *HLA-E* expression,

(**B**) overlaid topographical map of *HLA-E* tumor/stromal cell expression with color indicating abundance

of NK and cytolytic CD8 T cells, and (**C**) proximity of NK cells, CD8 T cells, and Tregs to myeloid cells

expressing any combination of *CXCL9*, *CXCL10*, and/or *CXCL11*.

**D,** Correlation analysis of Visium spots, defined as tumor cell with or without stromal cells, for co-

expression of *CXCL9* and *HLA-E* expression in BCG-naïve (N=4, left) and BCG-unresponsive (N=4, right)

NMIBC tumor specimens.

**E,** Correlation analysis of Visium spots, defined as tumor cell with or without stromal cells, for co-

expression of *CXCL12* and *HLA-E* expression in BCG-naïve (N=4, left) and BCG-unresponsive (N=4,

right) NMIBC tumor specimens.

**F,** Meta-analysis of all tumor/stroma-labelled Visium spots showing row-scaled expression of pertinent genes

stratified by number of neighboring infiltrating NK cells and/or CD8 T-cells.

**G,** Z-scored heatmap showing $HLA\text{-}E^{LOW}$, low-cytotoxic infiltrating tumor spots (N = 1,200 Visium spots) vs HLA-

$E^{HIGH}$ and high-NK/CD8 T cell-infiltrated tumor spots (N = 242 Visium spots).

**H,** Representative multiplexed IF analysis by PhenoCycler$^{TM}$ of bladder tumor sections from one patient

with BCG-unresponsive NMIBC highlighting the presence of stroma (vimentin, blue) and tumor (S100A4,

yellow). Scale bar indicates 2mm.

**I,** Magnified inset from Panel H, highlighting tumor S100A4 expression of and use of digital pathology

software to identify NKG2A$^+$ NK and CD8 T cells along FoxP3+ CD4+ Tregs, macrophages (CD68 and

CD163) and dendritic cells (CD11c and/or HLA-DR). Image scale bar indicates 20μm.

1255

**Fig. 7: In vitro combination NKG2A and PD-L1 blockade restores tumor-derived NK and CD8 T**

**cell-mediated antitumor activity.**

**A,** Median fluorescence intensity (MFI) of HLA-E and PD-L1 expression by wild-type (WT) or HLA-E$^+$

K562 tumors cultured overnight in the presence or absence of recombinant human IFN-γ.

**B and C,** Phenograph clustering meta-analysis of tumor-derived and expanded CD56$^+$ CD8 T cells by

CyTOF before co-culture with K562 tumors showing **B,** tSNE analysis of identified CD8 T cell clusters,

and **C,** expression of individual inhibitory and activating receptors by CD8 T cells and their distribution

across clusters.

**D,** Representative CyTOF analysis of CD56 and NKG2A expression on tumor-derived NK (top row) and

CD8 T cells (bottom row) profiled before (left column) and after (right column) expansion with low dose

recombinant human IL-2, IL-7, IL-15 and CD3/CD28 tetramers.

**E,** Representative fluorescence flow cytometric analysis of IFN-g and CD107a expression by CD56$^+$ NK

(top row) and CD8 T cells (bottom row) after 6-hour culture alone or in presence of K562 tumor lines with

or without pre-treatment with anti-PD-L1 antibodies or with anti-PD-L1 + anti-NKG2A antibodies.

**F,** Fold-change differences between the frequencies of NK and CD56$^+$ CD8 T cells when comparing

between experimental conditions (defined by "+"). Individual comparisons are indicated at the bottom. *,

$p < 0.05$; **, $p < 0.001$. P values determined by paired Wilcoxon matched rank test.

1273

1274

1275

1276

**References**

1.  Redelman-Sidi, G., Glickman, M.S. & Bochner, B.H. The mechanism of action of BCG therapy for bladder cancer--a current perspective. *Nat Rev Urol* **11**, 153-162 (2014).

2.  Kamat, A.M. *et al.* Predicting Response to Intravesical Bacillus Calmette-Guerin Immunotherapy: Are We There Yet? A Systematic Review. *Eur Urol* **73**, 738-748 (2018).

3.  Müller, G. *et al.* Association between Development of Metabolic Acidosis and Improvement of Urinary Continence after Ileal Neobladder Creation. *J Urol* **203**, 585-590 (2020).

4.  Hedgepeth, R.C., Gilbert, S.M., He, C., Lee, C.T. & Wood, D.P., Jr. Body image and bladder cancer specific quality of life in patients with ileal conduit and neobladder urinary diversions. *Urology* **76**, 671-675 (2010).

5.  Morales, A. BCG: A throwback from the stone age of vaccines opened the path for bladder cancer immunotherapy. *Can J Urol* **24**, 8788-8793 (2017).

6.  Zlotta, A.R. *et al.* What is the optimal regimen for BCG intravesical therapy? Are six weekly instillations necessary? *Eur Urol* **37**, 470-477 (2000).

7.  Sfakianos, J.P. *et al.* Bacillus Calmette-Guerin (BCG): Its fight against pathogens and cancer. *Urol Oncol* **39**, 121-129 (2021).

8.  Sharma, P., Hu-Lieskovan, S., Wargo, J.A. & Ribas, A. Primary, Adaptive, and Acquired Resistance to Cancer Immunotherapy. *Cell* **168**, 707-723 (2017).

9.  Gocher, A.M., Workman, C.J. & Vignali, D.A.A. Interferon-γ: teammate or opponent in the tumour microenvironment? *Nat Rev Immunol* (2021).

10. Benci, J.L. *et al.* Opposing Functions of Interferon Coordinate Adaptive and Innate Immune Responses to Cancer Immune Checkpoint Blockade. *Cell* **178**, 933-948.e914 (2019).

11. Wang, L. *et al.* Myeloid Cell-associated Resistance to PD-1/PD-L1 Blockade in Urothelial Cancer Revealed Through Bulk and Single-cell RNA Sequencing. *Clin Cancer Res* (2021).

12. Strandgaard, T. *et al.* Field Cancerization Is Associated with Tumor Development, T-cell Exhaustion, and Clinical Outcomes in Bladder Cancer. *Eur Urol* (2023).

13. Strandgaard, T. *et al.* Elevated T-cell Exhaustion and Urinary Tumor DNA Levels Are Associated with Bacillus Calmette-Guérin Failure in Patients with Non–muscle-invasive Bladder Cancer. *European Urology* (2022).

1320   14.   Kates, M. *et al.* Adaptive Immune Resistance to Intravesical BCG in Non-Muscle Invasive
1321         Bladder Cancer: Implications for Prospective BCG-Unresponsive Trials. *Clin Cancer Res* **26**,
1322         882-891 (2020).
1323

1324   15.   Rosenberg, J.E. *et al.* Atezolizumab in patients with locally advanced and metastatic
1325         urothelial carcinoma who have progressed following treatment with platinum-based
1326         chemotherapy: a single-arm, multicentre, phase 2 trial. *Lancet* **387**, 1909-1920 (2016).
1327

1328   16.   Powles, T. *et al.* Efficacy and Safety of Durvalumab in Locally Advanced or Metastatic
1329         Urothelial Carcinoma: Updated Results From a Phase 1/2 Open-label Study. *JAMA Oncol*
1330         **3**, e172411 (2017).
1331

1332   17.   Sharma, P. *et al.* Nivolumab in metastatic urothelial carcinoma after platinum therapy
1333         (CheckMate 275): a multicentre, single-arm, phase 2 trial. *Lancet Oncol* **18**, 312-322
1334         (2017).
1335

1336   18.   Patel, M.R. *et al.* Avelumab in metastatic urothelial carcinoma after platinum failure
1337         (JAVELIN Solid Tumor): pooled results from two expansion cohorts of an open-label, phase
1338         1 trial. *Lancet Oncol* **19**, 51-64 (2018).
1339

1340   19.   Grossman, J.E., Vasudevan, D., Joyce, C.E. & Hildago, M. Is PD-L1 a consistent biomarker
1341         for anti-PD-1 therapy? The model of balstilimab in a virally-driven tumor. *Oncogene* **40**,
1342         1393-1395 (2021).
1343

1344   20.   Kamiya, T., Seow, S.V., Wong, D., Robinson, M. & Campana, D. Blocking expression of
1345         inhibitory receptor NKG2A overcomes tumor resistance to NK cells. *J Clin Invest* **129**, 2094-
1346         2106 (2019).
1347

1348   21.   Pereira, B.I. *et al.* Senescent cells evade immune clearance via HLA-E-mediated NK and
1349         CD8(+) T cell inhibition. *Nat Commun* **10**, 2387 (2019).
1350

1351   22.   van Montfoort, N. *et al.* NKG2A Blockade Potentiates CD8 T Cell Immunity Induced by
1352         Cancer Vaccines. *Cell* **175**, 1744-1755.e1715 (2018).
1353

1354   23.   Andre, P. *et al.* Anti-NKG2A mAb Is a Checkpoint Inhibitor that Promotes Anti-tumor
1355         Immunity by Unleashing Both T and NK Cells. *Cell* **175**, 1731-1743 e1713 (2018).
1356

1357   24.   Salomé, B. *et al.* NKG2A and HLA-E define a novel alternative immune checkpoint axis in
1358         bladder cancer. *Cancer Cell* **40**, 1-17 (2022).
1359

1360   25.   Herbst, R.S. *et al.* COAST: An Open-Label, Phase II, Multidrug Platform Study of
1361         Durvalumab Alone or in Combination With Oleclumab or Monalizumab in Patients With
1362         Unresectable, Stage III Non-Small-Cell Lung Cancer. *J Clin Oncol*, JCO2200227 (2022).
1363

1364  26.  Rouanne, M. *et al.* BCG therapy downregulates HLA-I on malignant cells to subvert
1365       antitumor immune responses in bladder cancer. *J Clin Invest* **132** (2022).
1366

1367  27.  Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set
1368       collection. *Cell Syst* **1**, 417-425 (2015).
1369

1370  28.  Jiang, Y.H. *et al.* Urine cytokines as biomarkers for diagnosing interstitial cystitis/bladder
1371       pain syndrome and mapping its clinical characteristics. *Am J Physiol Renal Physiol* **318**,
1372       F1391-f1399 (2020).
1373

1374  29.  Tyagi, P. *et al.* Urine cytokines suggest an inflammatory response in the overactive bladder:
1375       a pilot study. *Int Urol Nephrol* **42**, 629-635 (2010).
1376

1377  30.  Netskar, H. *et al.* Pan-cancer profiling of tumor-infiltrating natural killer cells through
1378       transcriptional reference mapping. *Nat Immunol* (2024).
1379

1380  31.  Rebuffet, L. *et al.* High-dimensional single-cell analysis of human natural killer cell
1381       heterogeneity. *Nat Immunol* (2024).
1382

1383  32.  Vivier, E. *et al.* Natural killer cell therapies. *Nature* **626**, 727-736 (2024).
1384

1385  33.  Browaeys, R., Saelens, W. & Saeys, Y. NicheNet: modeling intercellular communication by
1386       linking ligands to target genes. *Nat Methods* **17**, 159-162 (2020).
1387

1388  34.  Amir el, A.D. *et al.* viSNE enables visualization of high dimensional single-cell data and
1389       reveals phenotypic heterogeneity of leukemia. *Nat Biotechnol* **31**, 545-552 (2013).
1390

1391  35.  Cristescu, R. *et al.* Pan-tumor genomic biomarkers for PD-1 checkpoint blockade-based
1392       immunotherapy. *Science* **362** (2018).
1393

1394  36.  Spranger, S. *et al.* Up-regulation of PD-L1, IDO, and T(regs) in the melanoma tumor
1395       microenvironment is driven by CD8(+) T cells. *Sci Transl Med* **5**, 200ra116-200ra116
1396       (2013).
1397

1398  37.  Rooney, M.S., Shukla, S.A., Wu, C.J., Getz, G. & Hacohen, N. Molecular and genetic
1399       properties of tumors associated with local immune cytolytic activity. *Cell* **160**, 48-61
1400       (2015).
1401

1402  38.  Ribas, A. Adaptive Immune Resistance: How Cancer Protects from Immune Attack. *Cancer*
1403       *Discov* **5**, 915-919 (2015).
1404

1405  39.  Grasso, C.S. *et al.* Conserved Interferon-γ Signaling Drives Clinical Response to Immune
1406       Checkpoint Blockade Therapy in Melanoma. *Cancer Cell* **38**, 500-515.e503 (2020).
1407

1408 40. Gocher, A.M., Workman, C.J. & Vignali, D.A.A. Interferon-γ: teammate or opponent in the
1409 tumour microenvironment? *Nat Rev Immunol* **22**, 158-172 (2022).

1410

1411 41. Bardia, A. *et al.* Sacituzumab Govitecan in Metastatic Triple-Negative Breast Cancer. *N Engl*
1412 *J Med* **384**, 1529-1541 (2021).

1413

1414 42. Powles, T. *et al.* Enfortumab Vedotin and Pembrolizumab in Untreated Advanced
1415 Urothelial Cancer. *N Engl J Med* **390**, 875-888 (2024).

1416

1417 43. Heath, E.I. & Rosenberg, J.E. The biology and rationale of targeting nectin-4 in urothelial
1418 carcinoma. *Nat Rev Urol* **18**, 93-103 (2021).

1419

1420 44. Chang, S.S. *et al.* Diagnosis and Treatment of Non-Muscle Invasive Bladder Cancer:
1421 AUA/SUO Guideline. *J Urol* **196**, 1021-1029 (2016).

1422

1423 45. Bisiaux, A. *et al.* Molecular analyte profiling of the early events and tissue conditioning
1424 following intravesical bacillus calmette-guerin therapy in patients with superficial bladder
1425 cancer. *J Urol* **181**, 1571-1580 (2009).

1426

1427 46. Shapiro, S.S. & Wilk, M.B. An Analysis of Variance Test for Normality (Complete Samples).
1428 *Biometrika* **52**, 591-611 (1965).

1429

1430 47. Kruskal, W.H. & Wallis, W.A. Use of Ranks in One-Criterion Variance Analysis. *Journal of*
1431 *the American Statistical Association* **47**, 583-621 (1952).

1432

1433 48. Zhang, X. *et al.* CellMarker: a manually curated resource of cell markers in human and
1434 mouse. *Nucleic Acids Res* **47**, D721-d728 (2019).

1435

1436 49. Glass, D.R. *et al.* An Integrated Multi-omic Single-Cell Atlas of Human B Cell Identity.
1437 *Immunity* **53**, 217-232.e215 (2020).

1438

1439 50. Magri, G. *et al.* Human Secretory IgM Emerges from Plasma Cells Clonally Related to Gut
1440 Memory B Cells and Targets Highly Diverse Commensals. *Immunity* **47**, 118-134.e118
1441 (2017).

1442

1443 51. Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-cell
1444 analysis. *Nature biotechnology* **42**, 293-304 (2024).

1445

1446 52. Finak, G. *et al.* MAST: a flexible statistical framework for assessing transcriptional changes
1447 and characterizing heterogeneity in single-cell RNA sequencing data. *Genome biology* **16**,
1448 1-13 (2015).

1449

1450 53. Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**, 1739-
1451 1740 (2011).

1452

54.    Rouillard, A.D. *et al.* The harmonizome: a collection of processed datasets gathered to serve and mine knowledge about genes and proteins. *Database (Oxford)* **2016** (2016).

55.    Galon, J. *et al.* Affinity of the interaction between Fc gamma receptor type III (Fc gammaRIII) and monomeric human IgG subclasses. Role of Fc gammaRIII glycosylation. *Eur J Immunol* **27**, 1928-1932 (1997).

56.    Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**, 1739-1740 (2011).

57.    Dries, R. *et al.* Giotto: a toolbox for integrative analysis and visualization of spatial expression data. *Genome biology* **22**, 1-31 (2021).

58.    Lause, J., Berens, P. & Kobak, D. Analytic Pearson residuals for normalization of single-cell RNA-seq UMI data. *Genome biology* **22**, 1-20 (2021).

59.    Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony. *Nature methods* **16**, 1289-1296 (2019).

60.    Ma, Y. & Zhou, X. Spatially informed cell-type deconvolution for spatial transcriptomics. *Nature biotechnology* **40**, 1349-1359 (2022).

61.    Jin, S., Plikus, M.V. & Nie, Q. CellChat for systematic analysis of cell-cell communication from single-cell and spatially resolved transcriptomics. *BioRxiv*, 2023.2011. 2005.565674 (2023).

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

| | Clinical Cohort Characteristics | | | | |
|---|---|---|---|---|---|
| | HTG | OLINK Urine (Sinai) | Single Cell | Spatial | Cytof |
| n samples, patients | 36, 27 | 42, 26 | 7, 6 | 20, 9 | 16, 16 |
| Female | 4 | 2 | 0 | 0 | 1 |
| BCG Naïve | 19 | 26 | 2 | 4 | 0 |
| BCG Unresponsive | 17 | - | 4 | 5 | 16 |
| Median Age | 64 | 69.5 | 66.9 | 70.6 | 68.5 |
| Median Months to Recurrence | 2.9 | 2.1 | 2.3 | 1.7 | 2.73 |
| Median Months to Progression | 6.7 | - | 14.3 | 23.28 | 4.43 |
| Stage at Collection | | | | | |
| CIS | 4 | - | - | - | - |
| Tis/Ta | 5 | 13 | - | 2 | 8 |
| T1 | 10 | 14 | 4 | 6 | 5 |
| T2 | - | - | - | - | 2 |
| T3 | - | - | 2 | - | 0 |
| T4 | - | - | - | - | 1 |

Table 1.

| Gene Set Names | Defining Genes |
|---|---|
| Interferon Gamma Response Signature | CCL5, CD3D, CXCL10, IL2RB, LAG3, IDO1, IL2RG, STAT1, PRF1, PDCD1, GZMA, GZMB, CD8A, CXCL9, TBX21, CD27, CD4, CD74, IFNG, CCR5, CXCL11, CXCL13, CXCR6, PTPRC |
| B Cell Markers | CD79A, BLNK, CD79B, CCR9, CD19, PNOC, BLK, CD22, HLA-DOB, TCL1A, CR2, SPIB, CD72, TNFRSF17, HLA-DQA1, PAX5, MS4A1, MEF2C, FCRL2 |
| Cytotoxic CD8 T-Cells | GZMK, FLT3LG, KLRK1, KLRD1, PRF1, GZMA, CD3G, CD8B, SIGIRR, GZMB, KLRB1, CD8A, GZMH, PF4, TBX21, CD3E, NKG7, CD27, IFNG, GNLY, EOMES, GZMM, KLRF1, KLRC1, RUNX3 |
| Cytotoxic Receptors and Function | ZNF683, SLAMF6, KLRK1, KLRD1, SLAMF7, NCR1, SH2D1B, GZMA, TYROBP, SH2D1A, GZMB, KLRB1, KIR3DL3, CLEC4C, CLEC4D, KLRC1, GZMH, NKG7, NCR3, GNLY, GZMM, KLRF1 |
| T-Cell Checkpoint Modulation | HLA-DQA2, CD274, TNFRSF14, CD86, CARD11, CD40, AKT1, CD3D, TNFRSF9, KLRK1, LAG3, CSK, TNFRSF13C, CD276, CD80, PDCD1, CD40LG, CD3G, PDCD1LG2, PTPN6, HLA-DQB2, ADORA2A, HLA-DPB1, TNFSF4, TIGIT, HLA-DRB1, CCL19, HAVCR2, CD3E, CD27, CD5, CCL21, CD4, HLA-DQA1, VAV1, CD70, CD24, GRB2, SRC, HLA-DRA, CD28, TNFSF13B, HLA-DPA1, TNFRSF4, CTLA4, ICOS, LYN, BTLA |
| Tumor Inflammation | PRF1, GZMA, GZMK, CCL5, CXCL10, IFNG, CXCL9, STAT1 |
| Th1 Polarization and Response | NFKB1, RELA, IL12B, TBX21, IL2, IFNGR1, IL12RB1, RBPJ, JAK2, TYK2, JAK1, DLL1, IFNG, STAT1, STAT4, IL12A |
| Lymphocyte Trafficking | CD274, FAM129C, CDH2, CXCR4, CD68, PDCD1, PDCD1LG2, PLCG2, CD40LG, HLA-DOB, MAPK11, ITGAL, CD52, HLA-DPB1, HLA-DRB1, PVR, CD4, ITGA5, HLA-DQA1, VAV1, NCAM1, HLA-C, LILRB3, CTLA4, CD83, CD86, LILRB1, HLA-DOA, LILRA3, CXCL12, ROCK2, LILRA5, ITGAX, CD8A, PLCG1, TIGIT, HLA-DMB, ITGA4, MMP9, CD28, CYBB, SELE, ICOS, CLEC4E, ICAM1, HLA-DQA2, SDC1, CEACAM3, CD96, CD80, CD22, CD84, SELL, CD209, PDPN, SIGLEC5, HLA-DRA, SIGLEC1, ITGB2, S1PR4, CD7, CD177, ITGAM, HLA-A, ITK, CD40, ITGB7, CD34, MSN, HLA-F, CD276, PTK2B, RASSF5, CD6, S1PR1, MAPK14, VCAM1, HLA-E, TREM1, HLA-DMA, HLA-DPA1 |
| General T Cell Markers | LCK, CD247, CD2, CD3E, CD3G, CD5, SH2D1A, TNFRSF25, TRAT1, CD96, CD6, LAMP1, CD3D, IL2RB, CD14, ZAP70 |
| Receptor | PDCD1, HAVCR2, TNFRSF14, CD27, CD40, TLR4, KIR2DL1, ICOS, BTLA, TNFRSF9, ADORA2A, LAG3, TNFRSF4, CTLA4, TIGIT, IL2RA, TNFRSF18 |
| Stress Toxicity | RAD17, DDIT3, TLR4, ATR, ADM, BID, CHEK2, HSP90B1, IL1B, RIPK1, CCL2, IL6, HSPA5, CD40LG, CRP, GADD45A, VEGFA, DDB2, HSP90AA1, CASP1, TNFRSF10A, PARP1, PVR, CDKN1A, TP53, TNFRSF10B, NBN, EDN1, FAS, SERPINE1, RAD51, NFAT5, ATG7, CA9, IFNG, ATG5, ATG12, XPC, CALR, GRB2, RAD9A, HMOX1, MMP9, HUS1, EPO, IL1A, CHEK1, ATM, TNF |
| DNA Repair (Hallmark) | TYMS, POLR1D, GTF2H1, ITPA, TAF6, TK2, SDCBP, GPX4, SMAD5, SAC3D1, UMPS, NUDT9, NME3, RPA3, NFX1, DGUOK, GTF2H3, DGCR8, DCTN4, ERCC2, TAF9, POLR2H, NME4, XPC, BOLA2_2B, DAD1, VPS37D, TMED2, NCBP2, RAE1, NUDT21, CCNO, GTF3C5, ERCC4, SUPT5H, ARL6IP1, EIF1B, ZNF707, AK3, VPS28, RAD52, TAF10, ERCC3, FEN1, SUPT4H1, PCNA, POLR2A, POLH, BCAP31, POLB, NT5C, HCLS1, SRSF6, POLR2J, REV3L, USP11, VPS37B, POLR2K, GTF2H5, ADCY6, HPRT1, TP53, PRIM1, RALA, BCAM, POLD1, CLP1, SURF1, POLD4, GMPR2, MPG, PNP, DDB1, POLR2D, RBX1, SF3A3, RFC4, ZWINT, IMPDH2, POLR2C, ADRM1, PDE6G, POLR2G, POLR1C, POLR2F, LIG1, DUT, CMPK2, SEC61A1, RFC3, GUK1, PDE4B, RAD51, GTF2F1, POLA2, STX3, TARBP2, CDA, APRT, ELL, RFC5, POM121, RPA2, POLR3C, NME1, POLR2E, AK1, NPR2, ERCC1, GSDME, TAF12, POLR2I, POLA1, CETN2, POLL, ZNRD1, POLD3, RNMT, TSG101, ELOA, DDB2, UPF3B, SNAPC5, CANT1, ADA, POLR3GL, TAF13, COX17, ERCC8, AAAS, BRF2, POLE4, SSRP1, TAF1C, RFC2, SNAPC4, EDF1, ERCC5, CSTF3, GTF2B, GTF2A2, NELFCD, MRPL40, RRM2B |
| P53 Pathway (Hallmark) | LRMP, IRAK1, DDIT3, AK1, BAK1, ACVR1B, RGS16, ITGB4, IFI30, HBEGF, SDC1, APAF1, MXD1, SOCS1, TXNIP, VWA5A, TRAFD1, BTG2, RAD51C, CEBPA, SPHK1, DGKA, HRAS, BAX, DDB2, GADD45A, CD82, TGFB1, RRAD, ELP1, FBXW7, CYFIP2, TCHH, PTPRE, ADA, PCNA, CDH13, CD81, CASP1, POLH, AEN, SLC3A2, SEC61A1, CDKN1A, LIF, TP53, PHLDA3, FAS, NINJ1, DEF6, FOS, TSPYL2, XPC, PROCR, HDAC3, RAD9A, HMOX1, ERCC5, CCND2, POM121, FOXO3, IL1A, CCND3, TNFSF9, TRAF4, TAP1, APP, HIST1H1C, PPP1R15A, NOTCH1, BTG1 |
| Oxidative Phosphorylation (Hallmark) Peroxisome (Hallmark) | VDAC1, ALAS1, HSD17B10, SDHA, ALDH6A1, ATP5F1, PHB2, BAX, TCIRG1, GPX4, CASP7, NDUFA5, POLR2F, COX17, HTRA2, SURF1, OAT, SLC25A3, DLAT, GPI HSD11B2, ABCD2, ABCB1, ERCC1, ERCC3, HRAS, CDK7, CADM1, FDPS, BCL10, ACSL1, MSH2 |
| Hypoxia | NFKB1, VDAC1, ADORA2B, ATR, NAMPT, ADM, TXNIP, MAP3K1, F3, VEGFA, EGR1, RBPJ, MIF, IGFBP3, NCOA1, HIF1A, PLAU, TP53, EDN1, SERPINE1, FOS, CA9, PER1, NOS3, MMP9, HMOX1, LGALS3, MET, EPO, APEX1, BLM, GPI, PIM1 |
| TGF Beta Signaling | CBL, CREBBP, LTBP1, ID1, TGFB1, SMAD5, MYC, PARP1, TGFBR3, EP300, IFNG, TGFBR1, SMAD2, TGFBR2, THBS1, SMAD3, TGFB3, MAPK3, TNF |

**Extended data Table S1:** Genes defining gene-set enrichment pathways

| Channel | Marker | Clone |
|---|---|---|
| 1 | DAPI | NA |
| 2 | CD45RO | T200/797 |
| 3 | HLA-A | EP1395Y |
| 4 | CD4 | EPR6855 |
| 5 | Podoplanin | 18H5 |
| 6 | CD20 | SP32 |
| 7 | CD68 | RM1031 |
| 8 | CD8 | EPR26538-16 |
| 9 | SMA | EPR5368 |
| 10 | CD11c | FK24 |
| 11 | HLA-E | MEM-E/02 |
| 12 | CD31 | WM59 |
| 13 | HLA-DR | EPR6148 |
| 14 | Granzyme B | EPR20129-217 |
| 15 | Vimentin | LN-6 |
| 16 | CD3e | EPR5361(2) |
| 17 | CD56 | EP2567Y |
| 18 | CD34 | 4H11 |
| 19 | PD-L1 | 28-8 |
| 20 | ICOS | HK53 |
| 21 | CD57 | HNK1 |
| 22 | FOXP3 | 2A11G9 |
| 23 | NKG2D | polyclonal |
| 24 | Pan-Cytokeratin | AE1+AE3 |
| 25 | CD163 | RM1114 |
| 26 | PAX-5 | EPR3730(2) |
| 27 | ASCT2 | CAL33 |
| 28 | PD-1 | CAL20 |
| 29 | CD21 | EP3093 |
| 30 | GLUT1 | EPR3915 |
| 31 | CD11b | M1/70 |
| 32 | Ki67 | EPR3610 |
| 33 | S100A4 | 1F12-1G7 |
| 34 | Collagen IV | HL1351 |
| 35 | CD14 | 1H5D8 |
| 36 | LC3B | EPR21234 |
| 37 | CD45 | EPR27167-58 |
| 38 | IFNG | IFNG/466 |
| 39 | CD79a | EP3618 |
| 40 | IDO1 | EPR20374 |
| 41 | CD44 | IM7.8.1 |
| 42 | NKG2A | ab96319 |
| 43 | GATA3 | EPR16651 |
| 44 | CD66b | polyclonal |
| 45 | CD138 | polyclonal |
| 46 | HIF1A | EPR16897 |
| 47 | MPO | EPR20257 |
| 48 | DNAM-1 | EPR23641-137 |
| 49 | NKp46 | Innate Pharma |
| 50 | CXCR3 | ab71864 |
| 51 | CD38 | EPR22691-219 |
| 52 | CD16 | EPR22409-124 |
| 53 | ATPA5 | polyclonal |
| 54 | CTST | polyclonal |
| 55 | LDHA | EP1563Y |
| 56 | HK1 | EPR10134(B) |
| 57 | MMP9 | EP1254 |
| 58 | PHHH3 | polyclonal |
| 59 | BAX | ab53154 |

**Extended data Table S2:** Antibody panel for CODEX multiplexed immunofluorescence staining

| cell cluster | signature genes |
|---|---|
| normal epithelial cell | C19orf33, SNCG, MIR205HG, KRT19, KRT17, FXYD3, SYT8, SMIM22, ELF3, KRT18, CLDN4 |
| blca | KRT8, CD24, KRT18, TACSTD2, MGST1, FXYD3, ELF3, CLDN4, SMIM22, KLF5, KRT7, KRT19, DHRS2, S100P, GDF15, PSCA, AGR2, SERINC2, S100A14, SDC1, GATA3, KRT20, CLDN7, FOXQ1, CYB5A, AKR1C3, CYP4B1, TMEM54, MDK, RARRES1, RAB25, NQO1, TMEM97, SDC4, VSIG2, RALBP1, EFNA1, CDH1, EHF, JUP, CAPS, IDH1, SPTSSB, HEBP2, SPINT2, CAMK2N1, AL158206.1, MGST2, TSTD1, MAL2, S100A13, ADIRF, HES1, PTPRF, ERBB2, TBX3, PHLDA2, RAB11FIP1, SFN, ID1, ATP1B1, GNL3, CYP4B1, TXNDC17, MARCKSL1, LSR, NR2F6, ABHD11, PLPP2, TNFRSF21, S100A16, SLPI, TSPAN13, TNFAIP2, SPINT1, NEDD4L, STAP2, KRTCAP3, CD9, BEX3, AKR1C2, EPCAM, RAB11A, PPARG, FOXA1, CTTN, GIPC1, ALAS1, AHR, SH3YL1, EMP2, TCEAL4, GGCT, PAFAH1B3, FBP1, NDUFS6, IGFBP3, SNHG18, GPRC5A, AP1S1, CISD3, ECHS1, TIMM13, RBM47, UQCC2, POLR2I, TSPAN6, OCIAD2, GRHL3, ARPC1A, KRT10, CENPX, CRABP2, MRPL14, DCXR, PRDX2 |
| endothelial cell | PCAT19, SPARCL1, RAMP2, VWF, EGFL7, ECSCR, PLVAP, AQP1, TM4SF1, CLDN5, ADGRL4, CLEC14A, CAV1, CRIP2, IGFBP7, EPAS1, RAMP3, EMP1, IFI27, KLF2, SPRY1, EMCN, A2M, ENG, NPDC1, STC1, HYAL2, CAVIN1, TCF4, BCAM, IFITM2, STOM, MMRN2, ESAM, CALCRL, IFITM3, CNN3, CAVIN2, TGFBR2, IGFBP4, CD34, S1PR1, SOCS3, PECAM1, ACKR1, GJA1, CD59, SLC9A3R2, ITGA6, NNMT, PDLIM1, APP, MTUS1, TSPAN7, ADAMTS9, C2CD4B, CDH5, ADAMTS1, APOLD1, MEF2C, LMO2, YBX3, ELK3, TIMP3, CYYR1, CD93, TFPI, THBD, RNASE1, SPTBN1, HLA-DRB5, ITGA5, SOX18, GRASP, CLU, ID3 |
| fibroblast | DCN, CCDC80, IGFBP6, COL1A2, MGP, COL6A2, C1R, PLAC9, FBLN1, CFD, PCOLCE2, SERPING1, LUM, SERPINF1, C1S, MFAP5, MFAP4, SOD3, COL6A1, PI16, OGN, GSN, CLU, SPARCL1, LTBP4, COL6A3, MMP2, EFEMP1, C3, SFRP2, SCARA5, ADH1B, GPX3, SLIT2, CALD1, TIMP2, COL3A1, SPARC, CTSK, FBN1, SPON2, PMP22, PCOLCE, CCN1, PLPP3, OLFML3, EGR1, TFPI, FBLN5, ACKR3, CD34, FBLN2, DPT, RARRES2, APOD, IGFBP5, LRP1, CCN2, ABI3BP, FGFR1, ISLR, GPNMB, AEBP1, CCN5, TIMP1, MXRA8, SOCS3, MFGE8, NNMT, IGFBP7, CEBPD, EMP1, CD248, MYC, C11orf96, AC080038.1, NFIA, FSTL1, TIMP3 |
| pericyte cell | RGS5, PPP1R14A, BGN, TPM2, MYL9, IGFBP7, CALD1, MFGE8, COL4A1, NDUFA4L2, COL4A2, C11orf96, ACTA2, MGP, CAV1, PDGFRB, COX7A1, SOD3, COL18A1, CPE, SPARC, COL3A1, COL6A2, CAVIN3, CSRP2, CRIP2, LHFPL6, COX4I2, WFDC1, COL1A2, A2M, TPM1, PDGFA, TGFB1I1, COL6A1, SPARCL1, MEF2C, FAM162B, HIGD1B, HES4, SELENOM, IGFBP2, CNN3, FN1, FILIP1L, MAP1B, LBH, EHD2, CRYAB, TPPP3, EPS8, IFITM3, PCOLCE, GJA4, CCDC102B, PTP4A3, MYO1B, NR2F2, TINAGL1, PLAC9, RRAS, CYGB, NREP, MYH11, RAMP1, AP002956.1, PLXDC1, RBPMS, SERPING1, ITGA1, LAMA4, PTMS, ACTN1, DSTN |
| schwann cell | NRXN1, CRYAB, GPM6B, S100B, CDH19, SCN7A, MAL, ANGPTL7, LGI4, CHL1, MPZ, GFRA3, PLP1, CADM1, MATN2, SEMA3B, ZEB2-AS1, ANK3, PMP22, AC005747.1, TMEM176B, SPARC, VWA1, PDGFA, CD9, IGFBP7, CLU, SPARCL1, FGL2, RHOB, PRNP, EPB41L2, NDRG2, CNN3, SAMHD1, PDK4, ZEB2 |
| smooth muscle cell | CNN1, ACTG2, PCP4, DES, MYH11, PPP1R14A, TPM2, ACTA2, MYL9, SMTN, FHL1, CSRP1, MYLK, CKB, ECRG4, RAMP1, ACTC1, SORBS1, TPM1, TGFB1I1, SPARCL1, PALLD, TAGLN, PDLIM3, CRYAB, CAV1, SYNPO2, SYNM, SORBS2, TNS1, HSPB6, SLMAP, RBPMS, FERMT2, SELENOM, PSD, NEXN, COX7A1, LIMS2, PTGS1, PDLIM7, CARMN, ACTN1, FN1, AC022868.1, VWA1, CALD1, AF001548.1, MAP1B, COL6A1, LPP, LMOD1, SVIL, FLNA, PGM5, NIBAN1, PKIG, DSTN, FXYD6, CFL2, PITX1, CAVIN1, DMD, PGM5-AS1, FLNC, SLC25A4, ALDH1B1, FBXO32, RBPMS2, HSPB8, LMO3, TNNT2, RBFOX3, DIXDC1, KANK2, TGM2, RRAS, RARRES2, COL4A2, ILK, VCL, CLU, IGFBP7, NFIA, PPP1R12A, COL6A2, NME4, NUPR1, TCEAL4, PLEKHO1, AD000090.1 |
| b cell | IGHM, CD79A, CXCR4, IGKC, CCR7, CD37, VPREB3, ARHGAP24 |
| dc1 | LGALS2, CLEC9A, DNASE1L3, WDFY4, IDO1, XCR1, HLA-DOB, RAB7B, SERPINF2, S100B, NAPSB, BATF3, P2RY6, WFDC21P, C1orf54, SLAMF8, CPVL, CADM1, ASB2, IRF8, SHTN1, PADI2, LYZ, C1orf162, NAAA, HCK, SPI1, SLAMF7, LY86, VMO1, HLA-DQB1, PLEK, ANPEP, FGD2, HLA-DQA1, MPEG1, AIF1, MNDA, CLEC7A, HLA-DRB6, HLA-DPB1, PTPRE, HLA-DPA1, LGALS9, RGS10, HLA-DMB, HLA-DRA, FGL2, HLA-DRB1, HLA-DMA, CD86, CPNE3, SMCO4, SNX3, RAB32, CXCL16, BCL2A1, PPT1, BASP1, LST1, GCA, ITGAX, RNASE6, AP1S2, ADA2, TAP1, RGCC, TGFBI, ITGB2, PARVG, AC004687.1, SELPLG, LSP1, COTL1, SERPINB9, CD83, CTSZ, BID, SEMA4A, GNA15, ASAP1, C20orf27, SAMHD1, UNC119, CCDC88A, PLEKHO1, RGS19, PYCARD, PDLIM7, SERPINB1, ATG3, LIMD2, LCP1, EVI2B, SEPTIN6, TYMP, HLA-DRB5, LSM6, FNBP1, CCND1, PAK1, FAM49B, TNFAIP8, WAS, HCLS1, CD48, NFKBID, TMEM14A, PRCP, PSMB9, SNX10, PPA1, UCP2, DUSP4, DNAJC4, ICAM3, CST7, RNH1, ANXA6, PSME2, NET1, SERPINF1, AC010618.1, GLIPR1, NME4, RNASET2, SGK1, DBNL, MRPL23, VASP, TPP1, FMNL1, ACTN1, CKLF, CFL1, CYB5R3, RHOG, CORO1A, SYNGR2, GPR183, LTB, MZT2A, CAP1, CDK2AP2, GNG5, SERPINB6, GMFG, GPSM3, CRIP1, GABARAP, PLAUR, AP2S1, DUSP2, CTSS, HLA-F, FCGRT, NR4A2, LAPTM5, CD53, RGS2, GTF3C6, BUD23, CD37, MSN, CD52 |
| dc2 | LST1, AIF1, HLA-DQB1, TYROBP, FCER1A, HLA-DQA1, CD1C, FCER1G, HLA-DMB, MS4A6A, C1orf162, SPI1, CLEC10A, GPR183, CD86, NAPSB, JAML, HLA-DPB1, HLA-DMA, HLA-DRB6, LYZ, CD1E, PKIB, LY86, HLA-DRB1, HLA-DPA1, CPVL, RNASE6, MNDA, LGALS2, IGSF6, HCK, HLA-DRA, HLA-DQA2, HLA-DQB2, PLD4, FCGR2B, CALHM6, CXCL16, COTL1, NR4A3, OLR1, CLEC7A, ITGB2, AXL, RGS10, BCL2A1, LGALS9, PAK1, PLAUR, AP1S2, TYMP, CD83, LSP1, ALOX5AP, YWHAH, EVI2B, FAM49B, RGS1, PLSCR1, RGS2, PYCARD, CST7, GPSM3, MACROH2A1, RNASET2, HCLS1, BASP1, CD53, SAMSN1, CORO1A, LAPTM5, CXCR4, GMFG |
| macrophage | AIF1, C1QC, C1QA, C1QB, TYROBP, FCER1G, FCGR2A, MS4A7, MS4A6A, CD14, APOE, SPI1, APOC1, MSR1, FCGR3A, MS4A4A, TGFBI, CSF1R, LST1, CTSZ, HLA-DQA1, SPP1, CD163, CTSB, DAB2, HLA-DPB1, ADAP2, HLA-DMB, TREM2, ITGB2, LAPTM5, LGMN, CD4, CCL3, RNF130, HLA-DQB1, C1orf162, OLR1, |
| mast cell | TPSB2, TPSAB1, CPA3, IL1RL1, RHEX, LTC4S, GATA2, FCER1G, CTSG, HDC, VWA5A, KIT, MS4A2, CLU, ALOX5AP |
| monocyte | AIF1, TYROBP, FCER1G, LST1, LYZ, S100A8, S100A9, SERPINA1, BCL2A1, PLAUR, IL1B, C5AR1, CTSS, G0S2, FCN2, MNDA, SPI1, OLR1, SOD2, TYMP, COTL1, MS4A6A, CSTA, SAMSN1, MS4A7, EREG, CD14, C1orf162, CLEC7A, ITGB2, SRGN, LAPTM5 |
| mreg dc | LAMP3, CCL22, CCL19, NCCRP1, EBI3, CCL17, TVP23A, AOC1, IDO1, CD200, FSCN1, CFP, CCR7, WFDC21P, RASSF4, TMEM176A, CD86, POGLUT1, BIRC3, CD40, IL4I1, TMEM176B, BCL2A1, TBC1D4, RAMP1, NRP2, CD83, MIR155HG, GPR183, LSP1, HLA-DQA1, SAMSN1, IRF4, BASP1, TYROBP, HLA-DQB1, LAD1, IL7R, TXN, KDM2B, HLA-DPB1, MARCKSL1, HLA-DQA2, KYNU, HLA-DRA, NUB1, RFTN1, DUSP5, SYNGR2, ANXA6, RELB, MTHFD2, MGLL, RGS1, LGALS9, HLA-DPA1, TYMP, ERICH1, CSTA, ID2, GRSF1, SPI1, NABP1, CRIP1, KIF2A, HLA-DRB1, PPA1, VOPP1, CFLAR, SOCS1, NFKB1, HMGN3, TNFAIP8, COTL1, CTSH, STK4 |
| neutrophil | G0S2, CSF3R, TREM1, LUCAT1, S100A8, IL1R2, SLC11A1, NAMPT, CXCL8, BCL2A1, S100A9, IVNS1ABP, TPI1, ZFAS1 |
| nk cell | NKG7, GNLY, CCL5, CTSW, CST7, GZMB, TYROBP, PRF1, CD7, GZMA, HCST, KLRD1, CD247, TRDC, PTPRC, KLRC1, CCL4 |
| plasma cell | MZB1, DERL3, CD79A, FKBP11, JCHAIN, IGHG3, TNFRSF17, IGLC1, IGHA1, SEC11C, ANKRD28, SLAMF7, PRDX4, IGHG1, JSRP1, XBP1, IGLC3, IGKC, POU2AF1, CD38, IGHG2, ST6GAL1, TPD52, SPCS3, SDF2L1, ANKRD36BP2, DNAJB9, SPAG4, EAF2, ERLEC1, IGHA2, CRELD2, RNU2-63P, TXNDC15, DTNB, TXNDC5, HERPUD1, KLF13, IGHGP, PRDM1, IGHG4, IGLV3-1, PDE4B, PLPP5, ISG20, GLRX, SSR3, RN7SL2, RN7SL1, NUCB2, MANF, CYTOR, ICAM3, CFLAR, SELENOS |
| plasmacytoid dendritic cell | PLAC8, PLD4, SCT, LILRA4, LRRC26, PTCRA, SPIB, TCL1A, SMPD3, IRF4, NAPSB, LILRB4, DERL3, GZMB, MZB1, IRF7, BCL11A, JCHAIN, MYBL2, IRF8, C12orf75, GPR183, CLIC3, LDLRAD4, SELL, EGLN3, ITM2C, NR4A3, ALOX5AP, THEMIS2, CXCR3, RNASE6, TSPAN13, OPN3, CCDC50, SLC15A4, PPP1R14B, SEC61B, AC004687.1, AREG, FCER1G, DUSP5, SLC7A5, TYROBP, CXCR4, HERPUD1, NR3C1, GNA15, CREM, SERPINF1, ZC3HAV1, N4BP2L1, RGS1, RGS2, TCF4, MS4A6A, HLA-DRB1, PIM2, HLA-DRA, NPC2, LTB, PLP2, HLA-DPB1, ZNF331, CD83, RNASET2, HLA-DMA, SRGN, YPEL5, CTSB, HLA-DPA1, CD37, RPL7, UCP2, NR4A2, HLA-DQB1, LSP1 |
| t cd4 | CD2, CD3D, LTB, IL7R, IL32, CD52, TRAC, TRBC2, CD3E |
| t cd8 | NKG7, CD3D, CCL5, GZMA, TRBC2, CD2, TRAC, CD3G, CD8A, CTSW, LINC01871, HCST, CD3E, CST7, CD52, CD7, IL32, GZMB, PTPRC, CCL4, CORO1A, TRGC2, IL2RG, ALOX5AP, LCK, RGS1, TRBV28, RAC2, KLRD1 |
| t cell cycle | PCLAF, TYMS, CD2, TRBC2, LINC01871, CD3G, NUSAP1, STMN1, CD3D, SMC4, CD7, PTTG1, HMGB2, TOP2A, APOBEC3G, TMPO, CENPM, CENPK, ZWINT, UBE2C, CKS1B, MAD2L1, TIGIT, TK1, LMNB1, SMC2, MKI67, CENPF, SIT1, CDKN3, CLEC2D, CD8A, CDK1, PHF19, PRC1, AURKB, TRGC2, TACC3, KLRC1, CENPU, HELLS, CXCL13, UBE2T, RRM2, ASF1B, H2AC12, LAG3, CTLA4, BIRC5, LINC02195, TUBB, HMGN2, TRAC, RAB27A, PTPN7, GZMB, CORO1A, CENPW, RAC2, CD3E, NKG7, IFNG, AL035456.1, EZH2, GZMA, H2AX, LCK, IL2RB, PCNA, DTYMK, TUBA1B, H2AZ2, CD96, C12orf75, DUT, PSMB9, CD247, CRIP1, MCM7, HMGB1P5, NUDT1, DUT, ANP32E, TNFRSF18, TBC1D10C, SKA2, DEK, PTPN22, DDX39A, PCED1B-AS1, IL2RG, GNG2, CKLF, CALM3, CBX5, SLA, CCL5, LCP1, BUB3, PSMB8, OXNAD1, DNMT1, CCDC167, ANP32B, TUBBP1, PSIP1, IL32, RPA3, HCST, CYTOR, PTPRC, LSP1, GMFG, SEPTIN1, CD52, ANXA6, RANBP1, HOPX, PAXX, CST7, LBH, ICAM3, ITM2A, IFI16, HLA-F, EVL, ACAP1, ALOX5AP, CKS2, LIMD2, FYN, CBX3, NAP1L4, MZT2A, MT2A, IFI27L2, COTL1, CD53, HCLS1, STK17B, ID2, RGS1, SAMSN1, FXYD5, LAPTM5 |
| t reg | LTB, CD3D, TIGIT, IL32, TNFRSF4, TRAC, CD2, TNFRSF18, TRBC2, SPOCK2, CTLA4, CD27, CD3E, FOXP3, CD7, BATF, IL2RG, TRBC1, IL2RA, CARD16, TBC1D4, CORO1A, LCK, RGS1, IL7R, PTPRC, CD3G, ICOS, RAC2, PCED1B-AS1, CYTOR, CD52 |

**Extended data Table S3:** Annotated clusters with list of significantly defining genes for major clusters defined by scRNAseq

| Cluster ID | Cell count: |
|---|---|
| normal epithelial | 3458 |
| HLA-E-bright tumor | 5461 |
| HLA-E-dim/neg tumor | 9256 |
| endothelial cell | 3135 |
| fibroblast | 7630 |
| pericyte cell | 1963 |
| schwann cell | 186 |
| smooth muscle cell | 660 |
| b cell | 3675 |
| dc1 | 80 |
| dc2 | 599 |
| macrophage | 5271 |
| mast cell | 1015 |
| monocyte | 1871 |
| mreg dc | 150 |
| neutrophil | 519 |
| nk cell | 2580 |
| plasma cell | 1301 |
| plasmacytoid dendritic cell | 171 |
| t cd4 | 8548 |
| t cd8 | 5491 |
| t cell cycle | 629 |
| t reg | 1675 |
| | 65,324 |

**Extended data Table S4:** Annotated scRNAseq clusters with numbers of cells

A

B

**Extended data Figure S1:** Cellular sources of IFN-γ in NMIBC by scRNAseq

A

B

C

D

**Extended data Figure S2**

**A** EMT full pathway

**B** Autophagy pathway

**C** Cell cycle pathway

Kruskal-Wallis,
p < 2.2e-16

Kruskal-Wallis,
p < 2.2e-16

Kruskal-Wallis,
p < 2.2e-16

Extended
Data Fig. S3

Extended
Data Fig. S4

| | Clinical Cohort Characteristics | | | | |
|---|---|---|---|---|---|
| | HTG | OLINK Urine (Sinai) | Single Cell | Spatial | Cytof |
| n samples, patients | 36, 27 | 42, 26 | 7, 6 | 20, 9 | 16, 16 |
| Female | 4 | 2 | 0 | 0 | 1 |
| BCG Naïve | 19 | 26 | 2 | 4 | 0 |
| BCG Unresponsive | 17 | - | 4 | 5 | 16 |
| Median Age | 64 | 69.5 | 66.9 | 70.6 | 68.5 |
| Median Months to Recurrence | 2.9 | 2.1 | 2.3 | 1.7 | 2.73 |
| Median Months to Progression | 6.7 | - | 14.3 | 23.28 | 4.43 |
| Stage at Collection | | | | | |
| CIS | 4 | - | - | - | - |
| Tis/Ta | 5 | 13 | - | 2 | 8 |
| T1 | 10 | 14 | 4 | 6 | 5 |
| T2 | - | - | - | - | 2 |
| T3 | - | - | 2 | - | 0 |
| T4 | - | - | - | - | 1 |

Table 1.