1
2
3 **A FUCCI sensor reveals complex cell cycle organization of *Toxoplasma***

4 **endodyogeny.**

7 Mrinalini Batra[1], Clem Marsilia[1], Danya Awshah[1], Lauren M. Hawkins[1], Chengqi Wang[2], Dale

8 Chaput[3], Daria A. Naumova[4] and Elena S. Suvorova[1*]

9
10
[1]Division of Infectious Diseases, Department of Internal Medicine, Morsani College of Medicine,

University of South Florida, Tampa, FL 33612 USA

[2]College of Public Health, University of South Florida, Tampa, FL 33612 USA

[3]Proteomics Core, College of Arts and Sciences, University of South Florida, Tampa, FL 33612 USA

[4]Department of Biology, Krieger School of Arts and Sciences, Johns Hopkins University, Baltimore, MD

21218 USA

17
18
19 Short Title: *Toxoplasma* tachyzoite cell cycle and FUCCI probe

20 Keywords: Apicomplexa, *Toxoplasma gondii*, cell cycle, Cdk-related kinase, FUCCI, PCNA1

21
22

*Address correspondence to:

Elena S. Suvorova

Division of Infectious Diseases and International Medicine

Department of Internal Medicine

Morsani College of Medicine

University of South Florida

3720 Spectrum Blvd, Suite 303

Tampa, FL 33612

email: suvorova@usf.edu

office 813-974-6739

Fax 813-974-0804

In this study, we report the atypical cell cycle organization of the unicellular eukaryotic pathogen

Toxoplasma gondii. The remarkably flexible cell division of *T. gondii* and other apicomplexan parasites

differs considerably from the cell division modes employed by other model eukaryotes. Additionally, there

is a lack of recognizable cell cycle regulators, which have contributed to the difficulties in deciphering the

order of events in the apicomplexan cell cycle. To aid in studies of the cell cycle organization of the *T.*

gondii tachyzoite, we have created the Fluorescent Ubiquitination-based Cell Cycle Indicator (FUCCI)

probes, *Toxo*FUCCI^S and *Toxo*FUCCI^{SC}. We introduced a DNA replication factor TgPCNA1 tagged with

NeonGreen that can be used alone or in conjunction with an mCherry-tagged budding indicator TgIMC3

in the auxin-induced degradation (AID) parental strain. The varied localization and dynamic cell cycle

oscillation have confirmed TgPCNA1 to be a suitable *T. gondii* FUCCI probe. The *Toxo*FUCCI^S analysis

showed that tachyzoite DNA replication starts at or near centromeric regions, has a bell-shaped dynamic

and a significant degree of the cell cycle asynchrony within the vacuoles. Quantitative live and

immunofluorescence microscopy analyses of *Toxo*FUCCI^S and its derivatives co-expressing epitope-

tagged cell cycle markers have revealed an unusual composite cell cycle phase that incorporates

overlapping S, G_2, mitosis and cytokinesis (budding). We identified five intervals of the composite phase

and their approximate duration: S (19%), S/G_2/C (3%), S/M/C (9%), M/C (18%) and C/G_1 (<1%). The

*Toxo*FUCCI^S probe efficiently detected G_2/M and Spindle Assembly Checkpoints, as well as the

SB505124-induced TgMAPK1 dependent block. Altogether, our findings showed an unprecedented

complexity of the cell cycle in apicomplexan parasites.

INTRODUCTION

Cells can produce copies of themselves through a strictly regulated order of events called the cell cycle. These events, or phases, are gated by checkpoints, which are critical for directing cell cycle progression by ensuring specific requirements are met. In G_1 phase, the cell gathers resources in preparation for chromosome replication, which is accomplished during synthesis, or S phase, and any replication errors are subsequently corrected in the dedicated G_2 phase. During G_2, the cell irreversibly blocks chromosomal replication and continues to amass resources for future daughter cells. In mitosis, or M phase, chromosomes are organized and separated into daughter nuclei. The final stage of cell division, cytokinesis, produces two daughter cells. The cyclin-dependent kinases (Cdk) and their corresponding cyclins trigger phase transitions. Loss of control over cell cycle progression can produce severe consequences for an organism, as seen in cancer cells.

While the general processes underlying the cell cycle are largely conserved, cell division is more enigmatic in certain organisms such as unicellular parasites. Our model of choice is *Toxoplasma gondii*, an apicomplexan parasite that infects eukaryotic hosts and is responsible for the disease toxoplasmosis. Most of what we know about parasite replication comes from studies of the tachyzoite, a fast-dividing form of *T. gondii*, which is responsible for the acute phase of infection (6-8). *Toxoplasma* tachyzoites divide within the parasitophorous vacuole by a process called endodyogeny. Two internal buds are formed within the mother cell, when repeated 5 - 6 times, produces 32-64 intravacuolar progeny. We recently characterized the central Crk (CDK-related kinase)-cyclin complexes that are required to advance through multiple points of regulation in the tachyzoite cell cycle (6, 9, 10). This cell cycle process deviates significantly from conventional norms, and our current understanding of specific sequences of events is predominantly based on RNA-seq datasets (1). However, cell cycle regulators are controlled by proteolysis, and there are no tools available that would allow researchers to probe *T. gondii* cell cycle protein regulators spatially and temporally.

Fluorescence Ubiquitination-based Cell Cycle Indicator (FUCCI) is a highly adaptable technology that can be used as a direct readout for the current cell cycle stage of an individual cell at any given time (2-4). The FUCCI approach is based on the universal mechanism that controls cell cycle progression:

ubiquitin-mediated turnover of critical cell cycle regulators. For example, sequential activation of the

major E3 ligase complexes APC/C and SCF^{Skp2} ensures cell cycle-dependent expression of DNA

replication machinery, including the replication factors Geminin (S/G$_2$) and Cdt1 (M/G$_1$). The conventional

FUCCI models use modified Geminin and Cdt1 factors fused to fluorescent proteins of different colors

(3). Consequently, the dividing FUCCI cells change in color of fluorescence depending on the cell cycle

phase transition. The resulting live fluorescence permits real time visualization, isolation, and analysis of

large amounts of cells that can be screened and categorized into subpopulations. Since its invention in

2008, FUCCI has been instrumental in a variety of research questions, such as in evaluating the effects

of drugs on cell replication by visually distinguishing between proliferating and quiescent cells, monitoring

the fates of various stem cell populations, or tracking tumor progression (5, 6). Currently, no FUCCI

analogs have been developed in unicellular organisms, including protozoans. Many checkpoint

regulators were presumed to be lost in the *T. gondii* genome, including the canonical FUCCI probe

Geminin (10, 11). However, since controlled proteolysis has retained its dominant role in regulating the *T.*

gondii cell cycle (9, 10, 12, 13), this suggests that the FUCCI technology could be successfully

implemented in *T. gondii*.

Since apicomplexan parasite division is poorly understood, the FUCCI approach emerged as a

vital tool to decode atypical cell cycle mechanisms and elucidate complex life stage transitions. In this

study we created several *Toxo*FUCCI cell lines that efficiently detected cell cycle blocks induced by

drugs or by knockdown of the endogenous regulators. The *Toxo*FUCCI approach revealed high

complexity of the cell cycle organization and hidden asynchrony of the intravacuolar division in *T. gondii*.

RESULTS

TgPCNA1 role in tachyzoites

The FUCCI technology can provide much-needed clarity to our understanding of the organization of the

T. gondii endodyogenic cell cycle. Modern FUCCI models contain dual markers that are expressed

sequentially in G$_1$ and in S-, G$_2$ phase and mitosis (7). However, the original gene markers used in

FUCCI must be modified for use in apicomplexan parasites, as is the case in plants and flies (4, 8, 9). *T.*

gondii does not encode the replication inhibitor Geminin, while the temporal expression of the CDT1

ortholog TgiRD1 is significantly deviated (10).

Eukaryotic proliferating cell nuclear antigen (PCNA) factors are highly conserved at the protein

level. They form a characteristic ring-like structure comprised of three associated PCNA molecules. This

ring "clamps" around a DNA strand and acts as a hub for replication machinery, notably recruiting the

DNA polymerases δ and ε, and ultimately facilitates efficient DNA replication (11). PCNA proteins from

species as divergent as insects and plants can be substituted for one another *in vitro* and can recruit

endogenous DNA replication machinery (12, 13). *T. gondii* expresses two PCNA factors, though only

TgPCNA1 (TGME49_247460) was found to be essential (14). TgPCNA1 is a conserved protein, which is

predicted to form a ring-shaped homotypic trimer (Fig. S1). The SWISS-Prot modeling revealed structural

conservation of *Toxoplasma* PCNA1 but detected two distinctive regions that are not present in

mammalian PCNA1 (Fig. S1). The coccidian-specific loop and a C-terminal extension, indicating a

species-specific adaptation of TgPCNA1 to a likely altered composition of the *Toxoplasma* replisome.

The function of *Toxoplasma* PCNA1 had not been examined previously. To evaluate whether any

conserved functions were preserved by TgPCNA1, and to determine whether this protein can be used as

a probe for FUCCI, we engineered a transgenic line for the auxin-dependent TgPCNA1 expression and

evaluated the role of TgPCNA1 in tachyzoite division. Immunofluorescence microscopy analyses showed

nuclear TgPCNA1[AID-HA] localization in either a diffused or an aggregated form (Fig. 1A) (14). Co-staining

with Centrin1 and TgIMC1 verified that TgPCNA1[AID-HA] expression patterns change as tachyzoites

progress through the cell cycle. Based on the number of parasites with nuclear TgPCNA1[AID-HA]

aggregates, DNA replication (S-phase) occupies 26.9+/-3.4% of the duration of the cell cycle (Fig. 1A).

Pulldown of either chromatin-bound or chromatin-free TgPCNA1 protein complexes confirmed

that active TgPCNA1 interacts with DNA replication machinery (Fig. 1B and C, Table S2). Only

chromatin-bound TgPCNA1 complexes contained mini-chromosome maintenance (MCM) and

nucleosome assembly factors. Chromatin-bound TgPCNA1 also showed strong associations with DNA

remodeling complexes such as MORC and SAGA (15, 16). Furthermore, TgPCNA1 co-precipitated with

RNA polymerase II and nuclear pore proteins, which verifies that replication and transcription can occur

concurrently in *T. gondii*. Altogether, our TgPCNA1 proteome analyses suggest that DNA replication,

transcription, and epigenetic coding can act cooperatively in *T. gondii*. Specific interactions with

developmental chromatin remodelers indicate that any epigenetic coding that is associated with a

specific developmental stage is likely introduced during S-phase.

Conditional downregulation of TgPCNA1 with auxin treatment verified that TgPCNA1 is essential

for tachyzoite growth. The TgPCNA1^{AID-HA} was efficiently degraded within 30 minutes of auxin treatment

and parasites were unable to form plaques when auxin was present in growth medium for 7 days (Fig.

2A and B). Consistent with its primary role in DNA replication, downregulation of TgPCNA1 was shown to

affect tachyzoites progressing through S-phase. Flow cytometry analyses showed the accumulation of

parasites containing more than 1N and less than 2N DNA content after 4 hours of TgPCNA1 deficiency,

indicating stalled DNA replication (Fig. 2C, red plot, Table S3). Extended TgPCNA1 depletion permitted

slow progression of DNA replication, similar to a slow S-phase phenotype of the fission yeast deficient in

PCNA1 function (Fig. 2C, orange plot) (17). The reduction of 1N DNA peaks indicated that TgPCNA1-

deficient parasites are unable to enter new cell cycle. Our results suggested that TgPCNA1 is required

for an efficient progression of DNA replication.

Our quantitative immunofluorescence microscopy analyses showed increased numbers of

TgPCNA1-deficient parasites with duplicated centrosomes, which suggests they are retained in the post-

G_1 phase (Fig. 2D, Table S3), corroborating our DNA FACScan data. To evaluate how the lack of

TgPCNA1 affects cell cycle progression, we examined parasite morphology by ultra-expansion

microscopy. Co-staining with a mitotic marker TgMORN1 and nuclear stain DAPI revealed that

TgPCNA1-deficient tachyzoites cannot bypass metaphase in mitosis (18-20). The resolution of

TgMORN1-positive structures such as an arrayed centrocone and associated daughter basal complexes

(dBCs) signify a normal progression to anaphase (Fig. 2E), whereas parasites lacking TgPCNA1 had

either an extended or unresolved centrocone and no signs of karyokinesis (nuclear division), indicating

the arrest of mitotic progression prior to anaphase (20). Examining the duplicated centrioles allowed us

to uncover substantial defects in G_2 function, which normally controls reduplication of chromosomes and

centrosomes (10). The TgPCNA1-deficient parasites contained more than 2 centrioles per centrosome and were often separated, which resembles phenotypes that are observed in parasites lacking the G_2 regulator TgiRD1 (Fig. 2F) (10). The centriole reduplication led to the construction of multiple internal buds, sometimes nested within one another in a 'Russian doll' fashion (Fig. 2F, bud-in-the-bud). Multiple daughter cell scaffolds and their associated dBCs were visualized using anti-Tubulin A and anti-TgMORN1 antibodies, respectively (Fig. 2E and F). The fact that the assembly of bud scaffolds took place suggests that that TgPCNA1 does not play a role in parasite budding. Therefore, aside from its role in DNA replication, the downregulation of TgPCNA1 had a major effect on the regulation of centrosome reduplication in G_2 phase. Thus, we concluded that TgPCNA1 expression is required for the normal progression of both S and G_2 phases.

We demonstrated that TgPCNA1 is abundantly expressed in the nucleus. Its maximum expression in S-phase and the factor deficiency phenotype confirm the conserved role for TgPCNA1 as a DNA replication factor. Altogether, this evidence suggests that TgPCNA1 as a suitable candidate in the *Toxoplasma* FUCCI probe set.

Construction of the *Toxoplasma* FUCCI probes.

To temporally segregate *Toxoplasma* FUCCI markers, we brought our focus to S-phase and budding (cytokinesis). FUCCI markers should be abundant and expressed in a large organelle such nucleus, as this is critical for accuracy of detection in live fluorescence. Based on these criteria, we built *T. gondii* RH strains expressing TgPCNA1 fused with two copies of NeonGreen fluorescent protein (TgPCNA1[NG]), either alone or in combination with the alveolar protein TgIMC3, which is tagged with three copies of mCherry fluorescent protein (TgIMC3[Ch]) (21). To preserve their native regulation, the greater genomic loci of TgPCNA1 and TgIMC3 were each integrated in the RH∆*Ku80*∆*HXGPRTosTIR1* parent, which contained the fusion protein coding sequences under the control of their endogenous promoters (Fig. 3C, Fig. S2A and B). Given that TgPCNA1 functions during S-phase, while TgIMC3 is enriched in the cytoskeleton of growing buds in cytokinesis, we have named our transgenic lines *Toxo*FUCCI[S] (expressing TgPCNA1[NG]) and *Toxo*FUCCI[SC] (co-expressing TgPCNA1[NG] and TgIMC3[Ch]). In addition, our

*Toxo*FUCCI parasites are suitable for auxin-induced degradation (AID) conditional expression of the

other protein of interest (22). The introduction of these *Toxo*FUCCI markers did not affect replication

rates in the parental strain, confirming that the new *Toxo*FUCCI probes can be used to reliably represent

the growth characteristics of the type I RH strain (Fig. S2F, Table S4).

TgPCNA1 is a suitable tachyzoite FUCCI marker.

The traditional FUCCI models produce fluorescent color changes in a cell as it transitions through

different cell cycle phases (3). The cell cycle of apicomplexan parasites is uniquely complex, and

features such as internal budding or the ability to produce multiple daughters can further complicate the

task of choosing appropriate FUCCI markers (23). Conveniently, our *Toxo*FUCCI markers not only

display changes in fluorescent expression levels over the course of the cell cycle, but they also

translocate to highlight different cellular compartments, allowing us to incorporate both temporal and

spatial information to identify distinct cell cycle states. Our comprehensive direct and

immunofluorescence microscopy analyses revealed that TgPCNA1[NG] in combination with a DNA dye

DAPI was sufficient to identify major cell cycle stages in endodyogeny (Fig. 3A). In tachyzoites,

TgPCNA1[NG] exists in two major states: sparsely diffused in the nucleus or aggregated in the nucleus at

high abundancy. Co-staining of *Toxo*FUCCI[S] tachyzoites with the centrosome marker Centrin 1

confirmed that diffuse TgPCNA1[NG] expression in the centrally positioned nucleus is characteristic of the

G_1 phase (Fig. 3A, top panel). Early mitosis is detected by the reduction of TgPCNA1[NG] levels in the

nucleus and a shift of the nucleus to the proximal end of the parasite (Fig. 3A, Marker - TgMORN1).

TgPCNA1[NG] was aggregated in bright foci that represent replicative forks, characteristic of tachyzoites in

mid-S phase. We noticed that during budding, the fluorescent signal from partially degraded TgPCNA1[NG]

was accumulated in the apical region (conoid), which can conveniently distinguish parasites in early (G_1,

no conoidal TgPCNA1[NG]) and late (mitosis and budding, conoidal TgPCNA1[NG]) cell cycle phases. Note

that TgPCNA1[AID-HA] did not show this conoidal accumulation in mitotic and budding parasites, making this

conoidal stain specific to NeonGreen fusion to TgPCNA1. The diffuse expression of TgPCNA1[NG] in a U-

shaped or divided nucleus, alongside the conoidal accumulation, is detected in budding parasites (Fig

3A. Marker – TgISP1) (24). Additionally, co-expression of TgPCNA1[NG] and the budding marker

TgIMC3[Ch] in the *Toxo*FUCCI[SC] cell line verified that TgPCNA1[NG] accumulates in the conoidal region of

internal buds (Fig. S2E). Flow cytometry analyses of *Toxo*FUCCI[SC] tachyzoites detected expression of

DNA replication and budding markers (Fig. 3C and D, Fig. S2G). Although the flow cytometry approach

could not provide data that would allow us to distinguish between the aggregated and diffused forms of

TgPCNA[NG], we nevertheless detected double-(NG+/mCh+) and single-positive (NG-/mCh+) populations,

which represents parasites in S-, budding, and G_1 phases.

Tachyzoite DNA replication starts at the centromeric regions.

DNA replication is a highly dynamic process that is executed with two key steps: licensing and firing of

DNA replication origins. In conventional eukaryotes, licensing is carried out in late mitosis and early G_1

by loading the pre-Replication Complex (pre-RC), composed of Cdt1, Cdc6 and MCM2-7 hexamers, on

all origins of DNA replication (25). At the onset of S phase, the sequential loading of the CMG complex

(Cdc45, GINS and MCM2-7) and RCF, PCNA, and two DNA polymerases converts a fraction of pre-RC

into replication forks. Origin firing initiates DNA synthesis, which progresses bidirectionally until

neighboring replication forks meet and fuse, continuing until all chromosomes are replicated (25). In

conventional eukaryotes, PCNA1 serves as a hub for replisome components, such as the δ and ϵ DNA

polymerases, and plays a central role in firing the origins of replication (12, 13).

The dynamics of chromosome replication have yet to be examined in *T. gondii*. Monitoring the

*Toxo*FUCCI[S] marker revealed a bell-shaped progression of DNA replication activity in tachyzoites

(Fig.3A and D). The first replication events take place in a single defined nuclear region, followed by a

massive spread of replication forks across the entire nucleus, which are later reduced to a few TgPCNA1

aggregates as replication comes to an end. In the most studied eukaryotes, the centromeres and

pericentromeric regions, which are critical for sister chromatid adhesion and attachments to

kinetochores, are replicated first (26). To determine if chromosome replication in *T. gondii* also begins at

the centromeres, we introduced the AID-HA epitope to the kinetochore protein TgCenP-C in the

*Toxo*FUCCI[S] probe (Fig. S2C and D, Fig. 3D, red channel). We observed colocalization of the

TgPCNA1^{NG} and TgCenP-C^{AID-HA} factors in the first replication aggregate, suggesting that centromeric

regions are the first DNA regions to be replicated in tachyzoites. By tracking expression of the

*Toxo*FUCCI^{S} probe, we saw that duplicated kinetochores remain closely associated during S-phase and

separate upon the completion of DNA replication (Fig. 3D, late S phase, Table S4).

DNA replication dynamics in tachyzoites

The *Toxo*FUCCI^{S} probe gave us an opportunity to examine the real-time dynamics of DNA replication in

tachyzoites. We measured the perdurance of the aggregated TgPCNA1^{NG} form, corresponding to active

replication forks, and determined that tachyzoites replicate their chromosomes in around 150 minutes

(+/- 10 minutes, Table S5). We found that DNA replication progressed in a bell-shaped dynamic (Fig. 4):

starting with a single aggregate (Fig. 4A and B, first focus), followed by a massive spread of replication

forks across the entire nucleus (Fig. 4C and D) that tapered off to a few TgPCNA1^{NG} aggregates as

replication came to an end (Fig. 4E). We noticed a two-step expansion of replication in early S-phase.

First, the replication focus, which is located near the centromeric region, expanded into a ring-like

structure that was surrounded by sparsely spaced TgPCNA1^{NG} aggregates (Fig. 4A and B, Movie S1). It

is likely that the TgPCNA1^{NG} ring aligned with the intranuclear spindle compartment centrocone, given

the localization of the first aggregate near bundled centromeres (Fig. S4E) (18, 27, 28). During the

second expansion step, the replication ring lost its organization and new replication forks emerged

randomly across the nucleus. The widespread assembly of replication forks marked the peak of DNA

replication. Further examination of the *Toxo*FUCCI^{S} probe during peak time showed rapid and random

movement of the aggregated replication foci (Movie S2). Panel C of Figure 4 depicts the active

rearrangement of TgPCNA1^{NG} aggregates, which were recorded every 30 seconds. As parasites slowly

reached the end of the DNA replication stage, the number of replication aggregates tapered off (Movie

S3. The final replication focus was primarily found in the proximal region of the nucleus (Fig. 4E),

suggesting that tachyzoites maintain some degree of chromatin organization. Interestingly, a time course

of PfPCNA1 displayed similar spatial dynamics of their replication forks: initial aggregation around the

centrocone (the cone of the hemi-spindle), followed by expansion throughout the nucleus, and

completion of DNA replication at the proximal end of the nucleus (29). This suggests that the

organization of the DNA replication process could be conserved across other apicomplexans.

Altogether, the real-time microscopy of the *Toxo*FUCCI[S] probe reveals organized DNA replication

in tachyzoites, which begins at the centromeres, radiates from the centrocone region, and subsides in

the proximal subnuclear compartment.

Asynchrony of the intravacuolar division of tachyzoites

The intravacuolar synchrony of dividing type I RH tachyzoites has been indisputably accepted based on

numerous observations using indirect and real-time microscopy (30, 31). The vast majority of these

studies monitored the budding process by following surface markers such as IMC1, GAP45, and SAG1.

Our *Toxo*FUCCI probe offers an alternative cell cycle marker that has revealed intravacuolar asynchrony

in DNA-replicating parasites. Although the *Toxo*FUCCI probe could not detect asynchrony during mid-S

phase, it allowed us to identify asynchrony at the start and end of DNA replication (Fig. 5A). Quantitative

microscopy showed that, depending on the number of intravacuolar divisions, 7% to 30% of DNA-

replicating vacuoles progressed through S-phase out of synchrony (Fig. 5B, Table S6). This asynchrony

gradually increased during the first three divisions and was easily traceable in the vacuoles of 8

tachyzoites. Figure 5A shows an example of the sequential initiation of DNA replication in *Toxo*FUCCI[S]

parasites that are co-stained with the centromeric marker TgCenP-C (upper panel). Quantifications

across the entire population, which represent a snapshot of all cell cycle phases, showed a similar trend

of increasing asynchrony in the first three intravacuolar divisions (Fig. 5C, Table S6). RH tachyzoites

grow in a characteristic rosette formation, which often splits during the fourth intravacuolar division (from

8 to 16 parasites) (30). Interestingly, we detected a consistent decrease of asynchrony during this stage.

It has been previously shown that a tubular network that connects these parasites plays an important role

in intravacuolar synchronization (30). One plausible explanation is that the rosette split shortens the

communication path between tachyzoites, which results in higher synchronization.

In real-time monitoring of *Toxo*FUCCI[S], we also detected temporal asynchrony in DNA

replication. While the majority of vacuoles displayed only minor discrepancies amongst individual

parasites as they started or ended DNA replication (5-10 minutes apart), a small fraction of vacuoles

contained parasites whose DNA replication processes were offset by 40-45 minutes, which constitutes

25-30% of the total DNA replication time (Fig. S3, Table S6). Moreover, some parasites within a vacuole

would abort DNA replication or had never begun the process, which would likely explain why some

vacuoles contain an odd number of parasites that have been noted in other studies (Fig. 5D, Movie S4)

(32, 33). In conclusion, the *Toxo*FUCCI^S approach revealed the asynchrony of the cell cycle during the

intravacuolar division of *T. gondii* tachyzoites.

Tachyzoite cell cycle has a composite S/G$_2$/M/C phase.

To determine the precise organization of the tachyzoite cell cycle, we evaluated the expression of the

*Toxo*FUCCI^S probe in combination with quantitative immunofluorescence microscopy (Fig. 6). In the

absence of a cell cycle synchronization technique, we quantify and take note of the morphological

changes in specific intracellular structures that indicate the current cell cycle stage (10, 20). In the

experiments described below, we evaluated individual parasites rather than whole vacuoles, given the

substantial degree of cell cycle asynchrony we observed in tachyzoites (Fig. 5).

First, we confirmed that the G$_1$ phase occupies approximately 50% of the cell cycle duration (1,

34), with half of the asynchronous *Toxo*FUCCI^S populations containing tachyzoites with diffused nuclear

TgPCNA^{NG} and a single centrosome (Fig. 6A, Table S4, Marker – TgPCNA^{NG}, Centrin1). We determined

that S-phase runs for about one third of the cell cycle, as 31% of parasites with TgPCNA^{NG} in nuclear

aggregates (replication foci). Co-staining with Centrin1 revealed that 10% of parasites undergoing DNA

replication (occupying 3% of the total cell cycle) had a single centrosome, indicating that centrosome

duplication in *T. gondii* occurs in early S-phase (Fig. 6B and C; red arrow; Table S4). Further

examination of mitotic and budding markers revealed substantial overlaps of cell cycle phases in *T.*

gondii endodyogeny.

S-phase/cytokinesis (budding) overlap: Previous studies suggested that cytoskeletal elements

begin to assemble during S-phase (35). To evaluate this hypothesis, we incorporated an endogenous tag

to TgBCC0, the earliest known marker of cytokinesis in *T. gondii*, and combined this with our

*Toxo*FUCCI^S probe and quantified parasites expressing TgBCC0^{myc} while undergoing DNA replication

(18). Our results confirm a substantial overlap between S-phase and cytokinesis (about 40%), since

TgBCC0 emergence begins as tachyzoites have progressed through 61% of S-phase (Fig. 4AB and C; C

and S; Table S4).

S-phase/cytokinesis/mitosis overlap: Next, we examined the progression of mitosis in the

tachyzoite cell cycle. Conventional mitosis is executed in four major phases: prophase, metaphase,

anaphase, and telophase. Each phase has defined tasks and is phenotypically distinguishable.

Chromatin condenses in prophase, and replicated chromosomes align across the bipolar spindle in

metaphase. Sister chromatid segregation takes place at the transition into anaphase and is regulated by

the Spindle Assembly Checkpoint (SAC) (36). Segregation of two daughter nuclei in closed mitosis, or

restoration of the nuclear membrane around separated sister chromatids in open mitosis, marks

telophase. The lack of chromatin condensation in *T. gondii* prevented the identification of prophase,

while the elements of metaphase and anaphase have been uncovered in several studies using the

TgMORN1 marker (10, 18, 20, 28, 37).

TgMORN1 is a reliable mitotic marker due to its cell cycle-dependent localization in two major

structures: the centrocone and the basal complex (18). Immunofluorescent microscopy studies of mitotic

tachyzoites revealed two patterns of TgMORN1 localization in the perinuclear region: a single centrocone

associated with two daughter basal rings (dBC), or two centrocones associated with one dBC each. The

resolution of the centrocone, caused by spindle break, marks the transition into anaphase (20). Recent

studies dissecting the assembly of the parasite cytoskeleton show that TgMORN1 integrates into the

basal complex after TgBCC0, which according to our quantifications, occupies 3% of the cell cycle (Fig.

6B and C; start of C and M phases; Table S4) (18). Quantitative immunofluorescence microscopy

analyses of *Toxo*FUCCI^S parasites stained with α-TgMORN1 antibody showed that 12% of

asynchronously dividing tachyzoites contain a single centrocone with two dBCs, most of which are

replicating DNA (aggregated TgPCNA1^{NG}). Since cytokinesis (TgBCC0 emergence) is initiated prior to

TgMORN1 integration into the dBC, the tachyzoite S-phase runs concurrently with mitosis and budding

for about 9% of the cell cycle (Fig. 6AB and C; S/M/C interval; Table S4). Considering the vital

importance of the temporal separation of chromosome replication and segregation events, it is likely that tachyzoites containing a single centrocone with two dBCs and undergoing DNA replication were progressing through prophase of mitosis (9% cell cycle). Tachyzoites in a similar stage that completed chromosome replication (diffuse TgPCNA1^{NG} staining and conoidal accumulation) were in metaphase of mitosis (Fig. 6AB and D; Table S4) (20). Centromere segregation became pronounced near the end of DNA replication, which could mark the prophase-to-metaphase-transition (Fig. 4B, red arrow; Table S4).

Mitosis/cytokinesis overlap: Our quantifications revealed that about 12% of tachyzoites with a resolved TgMORN1-positive centrocone contained a single nucleus, which is characteristic of anaphase in mitosis (Fig. 6D, Table S4). Co-staining *Toxo*FUCCI^{S} with the alveolar protein TgIMC1, which localizes to the mother and daughter IMC compartments, and DAPI show that 4% of tachyzoites completed karyokinesis (two nuclei) while retaining detectable mother IMC. This fraction represents mitotic telophase and transition into the G_1 phase of the next cell cycle. (Fig. 6BC and D; Table S4).

S-phase/G_2/cytokinesis overlap: We recently detected a G_2 period of *T. gondii* endodyogeny and showed that the G_2/M checkpoint operated an hour upstream of the SAC in tachyzoites (10). The conventional G_2 period separates S-phase and mitosis, however, our *Toxo*FUCCI^{S} probe detected 30% of tachyzoites replicating DNA that also contained detectable signs of mitosis and budding such as an enlarged centrocone with two dBCs, which implies that *Toxoplasma* G_2 period runs concurrently with S-phase and budding (Table S4). Unfortunately, we cannot determine the onset of G_2 until additional markers of the S/G_2 transition are identified.

Based on the analyses of our *Toxo*FUCCI^{S} model in combination with other cell cycle markers, we concluded that the tachyzoite cell cycle is comprised of G_1 and a composite cell cycle phase. The composite cell cycle phase incorporates five intervals: S, S/G_2/C, S/M/C, M/C, and C/G_1. Four intervals incorporate processes that contain characteristics of their neighboring cell cycle phases, revealing the unprecedented complexity of apicomplexan endodyogeny.

Validation of *Toxo*FUCCI probe using cell cycle arrest models

We introduced our FUCCI markers into the RHΔ*Ku80 TIR1* parental stain, which permits use of auxin-induced degradation models (22). As a result, our *Toxo*FUCCI probes can identify cell cycle blocks resulting from a knockdown of a protein of interest. To validate this function, we created the *Toxo*FUCCI[S] strain, co-expressing the known cell cycle regulators such as TgCrk4[AID-HA] and TgCrk6[AID-HA] (Fig. 7A, C, G; Fig. S2C and D, Fig. S4A) (10, 20, 37). First, we engineered *Toxo*FUCCI[S] strain expressing TgCrk4[AID-HA] and examined G_2/M arrest. We previously showed that a half-division cycle block (4 hours) results in a substantially synchronized population of tachyzoites (10). Staining *Toxo*FUCCI[S]/TgCrk4[AID-HA] parasites with the budding marker TgIMC1 showed the assembly of a single internal bud, signifying G_2/M arrest (Fig. 7B, Fig. S2A). According to our proposed organization of composite cell cycle phases, the G_2/M checkpoint operates at the end of the S/G_2/C interval (Fig. 6). Therefore, TgCrk4-deficient *Toxo*FUCCI[S] parasites should initiate budding and continue DNA replication. Quantitative IFA of *Toxo*FUCCI[S] positive parasites showed that after 4 hours of auxin treatment, 38% tachyzoites actively replicating DNA, and their numbers were increased after 7 hours of auxin treatment, supporting our model (Fig. 7B, D, Table S7). Flow cytometry analyses of DNA content confirmed the enrichment of populations undergoing DNA replication and the emergence of populations with re-replicated DNA, characteristic of a G_2 block (Fig. 7D, 1.8N peak). Likewise, flow cytometry analyses of arrested *Toxo*FUCCI[S]/TgCrk4[AID-HA] populations showed an increase in the number of NeonGreen-expressing parasites (8.4% to 37%) and a distinct population of parasites that re-replicated their DNA (Fig. 7E) (10).

We also created and tested a Spindle Assembly Checkpoint (SAC) *Toxo*FUCCI[S] probe, the *Toxo*FUCCI[S]/TgCrk6[AID-HA] transgenic line (Fig. S2C and D, Fig. S4B; Fig. 7G) (20). The *Toxoplasma* SAC was mapped to the M/C interval (Fig. 6), therefore a SAC block should produce a reduction of parasites replicating their DNA and an increase in parasites undergoing the budding process. Co-staining of *Toxo*FUCCI[S]/TgCrk6[AID-HA] parasites with TgIMC1 or TgMORN1 verified an efficient arrest at the SAC in parasites that were treated with auxin for 4 hours. Tachyzoites that were undergoing metaphase had already completed DNA replication and thus continued budding (Fig. 7F; Fig. S4C). Quantifications of *Toxo*FUCCI[S]/TgCrk6[AID-HA] parasites revealed a drastic reduction in populations replicating their DNA after 7 hours of SAC block (Fig. 7H). This was further validated by FACScan analysis of

*Toxo*FUCCI^S/TgCrk6^{AID-HA} tachyzoites, confirming that auxin treatment resulted in an accumulation of

parasites that completed DNA replication (Fig. 7I). Flow cytometry analysis of SAC-arrested

*Toxo*FUCCI^S/TgCrk6^{AID-HA} parasites revealed a shift toward the composite cell cycle phase (Fig. 7E, NG+

and NG- with 2N DNA content); however, it was not efficient in detecting metaphase block because of

the inability to distinguish between the diffused and aggregated forms of TgPCNA^{NG}.

Overlaying the DNA plots of the asynchronous, G$_2$/M, and SAC arrested populations further

confirmed a temporal segregation of these checkpoints. The G$_2$/M arrested parasites had a pronounced

peak at ~1.8N, confirming that they did not complete DNA replication, while SAC arrested tachyzoites

had a prominent 2N peak, indicating completed DNA replication (Fig. S4D). This observation sheds light

on the mysterious 1.7N-1.8N DNA peak that had been previously observed in asynchronously dividing

tachyzoites (10, 34, 38, 39). Our results suggest that parasites entering mitosis substantially reduce their

rate of DNA replication. According to the *Toxo*FUCCI mapping of the tachyzoite cell cycle, the last 20%

of nuclear DNA is replicated during one-third of S-phase (Fig. 6B). It is a period between G$_2$/M and SAC

that includes mitotic prophase and metaphase, and since we have demonstrated that tachyzoites do not

replicate their DNA during metaphase, it is likely that this slowly progressing DNA replication occurs

during prophase. Such slowed dynamics of DNA replication may provide the time needed to assess DNA

damage and prepare for chromosome segregation.

The effect of SB505124 compound on tachyzoite cell cycle

We had previously demonstrated the critical role of the cell cycle-regulated kinase TgMAPK1 in

controlling the timing of budding (40, 41). Treatment with the TgMAPK1 inhibitor SB505124 effectively

blocked assembly of internal daughter buds in tachyzoites, induced centrosome overamplification, and

led to an acute growth arrest. To determine what cell cycle phase depends on the functioning of

TgMAPK1, we treated our *Toxo*FUCCI^S probe with SB505124 for one division cycle (7 hours). The

results showed that SB505124-dependent TgMAPK1 inhibition affected processes that are carried out

during the S/G$_2$ interval. Specifically, a fraction of the drug treated tachyzoites continued DNA replication

and contained more than 2 centrosomes per cell (Fig. 8A and B, Table S8). Flow cytometry analyses of

DNA content showed a decrease in the non-replicating population (1N, G_1), which was accompanied by

stalled replication (1-2N, S-phase), hold at G_2/M (1.8N), and an increase in the proportion of parasites

with re-replicated DNA (>2N) after 4 hours of treatment (Fig. 8B, red plot). The prolonged treatment with

3µM SB505124 reduced the DNA-replicating population, increased the number of parasites with re-

replicated DNA (>2N), and increased the number of parasites that completed the cell cycle (G_1) (Fig. 8B,

orange plot; Fig. 7C, Table S8). The steady increase of 1.8N and >2N populations, and its similarity to

the TgCrk4-dependent cell cycle block, suggests that the SB505124 compound affects the cell cycle

prior to or at the G_2/M checkpoint.

The *Toxo*FUCCI approach revealed that SB505124 produced different affects across concurrent

cell cycle processes of the composite cell cycle phase. It allowed tachyzoites to postpone budding,

complete mitosis, re-enter S-phase, and start mitosis. Quantification showed an emergence of

binucleated cells that completed mitosis and karyokinesis but did not initiate budding (Fig. 8A and D,

Table S8). In agreement with previously reported defects associated with TgMAPK1 deficiency,

SB505124 treatment delayed parasite budding, leading to an increased occurrence of multiple daughters

(Fig. 8E and F, Table S8) (40, 41). Binucleated cells are extremely rare in tachyzoites dividing normally

(> 0.3%, Fig. 8D), as single nuclei are generally isolated throughout most of the budding process.

Nuclear division takes place in late budding, near the time of the dBC constriction (Fig. 8G, no treatment

panels). The emergence of binucleated cells in SB505124-treated populations suggest that the

tachyzoites had completed mitosis and karyokinesis without budding. Furthermore, the fact that the

binucleated cells contained scaffolds for four daughters suggests that these tachyzoites only initiated

budding after the second round of S-phase and mitosis, which resembles the multinuclear division that

takes place in the cat intestine stages of the parasite development. The *Toxo*FUCCI[S] probe uncovered a

hidden feature of TgMAPK1-deficient tachyzoites: the ability to re-enter S-phase and the loss of

intracellular synchrony that is displayed as a mosaic nuclei phenotype (Fig. 8A, Table S8). Up to 14% of

binucleated cells contained one DNA-replicating (TgPCNA1[NG] positive) and one non-replicating

(TgPCNA1[NG] negative) nucleus (Fig. 8D). Conditional downregulation of TgMAPK1[AID-HA] in the

*Toxo*FUCCI[S] line phenocopied the SB505124 treatment, confirming the specificity of the drug target (Fig.

8A). 50% of the SB505124-treated parasites also lost budding polarity, where internal daughters were

assembled in an antiparallel orientation (Fig. 8F).

In conclusion, our studies of three independent cell cycle regulators validated the *Toxo*FUCCI

probe as an efficient cell cycle sensor. Our results showed that the *Toxo*FUCCI system can facilitate

navigation through the complex and overlapping processes in *T. gondii* endodyogeny. We also showed

that the composite cell cycle phase allows parasites to separate and selectively regulate concurrent

processes such as chromosome replication, chromosome segregation, and cytokinesis (Fig. 8H).

DISCUSSION

In the current study, we determined the cell cycle organization of a single-cell eukaryote, *T. gondii*. We

engineered *Toxo*FUCCI probes and delineated the temporal flow of the endodyogenic cell cycle of *T.*

gondii tachyzoites. We measured the duration of major cell cycle processes and verified the overlaps

between S-phase, G_2 phase, mitosis, and cytokinesis. Our studies found no clear boundaries between

these phases, and thus we refer to the post-G_1 period that consists of five cell cycle intervals as a

composite cell cycle phase (Fig. 9). Real-time microscopy of the *Toxo*FUCCI^S probe revealed an

organized DNA replication process, which suggests the degree of chromatin organization in tachyzoites

is higher than anticipated (42). Additionally, the *Toxo*FUCCI approach uncovered hidden features of the

endodyogenic cell cycle, including the asynchronous progression of the cell cycle within an individual

vacuole.

In the conventional cell cycle, cell cycle phases do not overlap (Fig. 9). Moreover, Gap1 and

Gap2 phases separate the foundational processes of chromosome replication (S-phase) and chromatid

segregation (mitosis) and ensure that there is no overlap, or else a cell risks endangering the proper

inheritance of genetic information (43, 44). *T. gondii* must maintain some features that permit its bypass

of such critical rules. It is possible that *T. gondii* closed mitosis, in combination with its lack of DNA

condensation, eliminates the need for mitotic prophase, which is normally devoted to resolution of the

nuclear membrane and chromosome condensation in organisms that divide by open mitosis. Preparation

for mitosis (G_2/M) can start while parasites complete DNA replication, supported by our observations of

the translocation of replication machinery from centromeres in the apical end to proximal end of the

nucleus. This may provide a temporal window that acts to separate the end of chromosomal replication

and the beginning of chromatid separation. Coincidentally, recent studies in budding yeast have shown

their cell cycle progression occurs in a similar manner, by also starting chromatid segregation before

DNA replication is completed (45). Moreover, budding yeast begin cytokinetic processes at G_1/S and

have a functional G_2/M transition, an overlap similar to that in our findings in *T. gondii* endodyogeny of S-

phase, G_2, mitosis, and cytokinesis; in other words, the existence of a composite cell cycle phase (45,

46). Another organism that potentially utilizes composite cell cycle organization is the apicomplexan

Babesia bovis. scRNAseq analyses of *B. bovis* revealed a loose correlation of the gene expression with

corresponding cell cycle phases, suggesting these cell cycle processes may overlap (47). These

parallels strongly indicate that a composite cell cycle phase may not be such a rare finding, and is likely

an overlooked feature of the eukaryotic cell cycle.

The composite cell cycle phase shortens the duration of cell division, which may benefit *T. gondii*

tachyzoites in the race against the host immune system. Tachyzoites have a limited time to produce

enough progeny for conversion into cysts before the host immune system attacks. Furthermore, it is

tempting to suggest that the composite cell cycle phase permits apicomplexans to manipulate individual

cell cycle processes, and provides a convenient means to switch between binary and multinuclear

division modes. For example, temporal co-repression of G_2 and budding within the S/G_2/C interval would

favor multinuclear divisions such as schizogony and endopolygeny, while restoration of G_2 and budding

will activate a binary division mode. The composite cell cycle encompasses many processes that must

be co-regulated, which may explain why *T. gondii* tachyzoite requires the activity of several Cdk-related

kinases, including TgCrk4, TgCrk6, and possibly TgCrk5, all during the composite phase (Fig. 9) (10, 20,

37, 48). Complex cell cycle regulation is prone to mistakes such as asynchrony of intravacuolar

replication and aborted DNA replication, resulting in in the loss of progeny and the deviation from 2^N

geometric progression of progeny that is expected in binary division. However, apicomplexan parasites

produce large progenies, which likely compensates for mistakes made during cell division.

DNA replication had not been sufficiently examined in apicomplexan parasites. A few critical

observations were recently made in *Plasmodium falciparum*: ori-mapping, measurement of replication

fork rates across the entire genome, and the changing rates of DNA replication during nuclear stages

(49-51). These studies provide great technical details to the DNA replication process in *P. falciparum*,

however, the temporal progression of S-phase in individual nuclei remains unknown. Our *Toxo*FUCCI

studies show that DNA replication in *T. gondii* tachyzoites has defined temporal and spatial dynamics. In

line with previous reports, we established that DNA replication occupied 30% of the tachyzoite division

cycle (52). The estimation of S-phase duration suggested that *T. gondii* tachyzoites duplicate their

chromosomes slower than the red blood stage *P. falciparum* merozoites, which may be due to their

differences in genome size (49, 52). Our estimate based on *Toxo*FUCCI data is that tachyzoites replicate

their 67Mb genome in 150 minutes, at the rate of 530kb per minute. It requires the activity of

approximately 400 replication forks every minute if *T. gondii* retains the replisome rate of 1.2-1.4kb per

minute that has been reported in various eukaryotic cells including *P. falciparum* (49).

Although *T. gondii* follows the canonical rule in beginning its DNA replication at centromeres, the

DNA replication progresses in an unusual bell-shaped dynamic (30). Measurements of the replication

fork rates in *P. falciparum* support the slow initiation of DNA replication at the centromeric regions, as

centromeres are also replicated much slower than the rest of *P. falciparum* genome (49). Interestingly, a

slow replication rate was also associated with *P. falciparum* telomeres, raising the possibility that *T.*

gondii telomeres are also the last DNA loci to be replicated. Alternatively, this period may have been

devoted to regulating entry into mitosis. We showed that tachyzoites in the G_2/M transition have 1.7N-

1.8N DNA content, and thus are continuing to replicate DNA. They complete replication of the remaining

20% of DNA at a pace that is twice as slow while progressing through the last 30% of S-phase and

reaching the SAC. What mechanisms target origins of replication near the centromeres at the onset of S-

phase, and what mechanism is responsible for the reduced replication rate at the end of S-phase? These

important questions will be addressed in future studies.

The task of developing an apicomplexan FUCCI sensor presented several challenges, such as

the absence of conventional markers, and the general difficulty of separating cell cycle phases (10, 49).

Although PCNA1 was tested for a FUCCI probe in mammalian models, it was ultimately rejected due to

insufficient expression levels (3, 50). On the contrary, *Toxoplasma* PCNA1 is a highly abundant nuclear

protein that fits the FUCCI requirements: a transcriptional onset and proteolytic, cell cycle-dependent

offset. Similarly, the GFP-tagged *Plasmodium* PCNA1 had recently been used as a nuclear cycle sensor

to examine multinuclear division of *Plasmodium falciparum* (51). Our results showed that TgPCNA1

alone can identify cell cycle transitions, but nevertheless, we added the budding marker TgIMC3 as an

additional reference for later cell cycle stages (21). The *Toxo*FUCCI approach has great potential to

advance the studies of *T. gondii* biology, not only as a quantitative strategy that permits real-time tracing

of the tachyzoite cell cycle, but can also be adapted to numerous applications in basic and translational

research (4, 6). The *Toxo*FUCCI strains contain auxin-induced degradation modules, allowing

researchers to visualize the effects of targeted protein degradation on cell cycle progression (22). Our

*Toxo*FUCCI strains can be refined by introducing markers of other cell cycle phases or processes. For

example, tachyzoites spend half of their cell division time in the G_1 period, a stage that is "blinded" in the

current *Toxo*FUCCI lines. Our attempts to identify a G_1 protein that would satisfy all the FUCCI

requirements were not successful. Likewise, mitotic markers would enable identification of critical cell

cycle transitions such as the SAC or G_2/M (10, 20).

Unlike mRNA-based technologies, FUCCI provides an unbiased approach to the detection of cell

cycle phases, the major difference being that mRNA-based techniques detect mRNA, the precursor to a

protein, while FUCCI detects expression after translation. While experimental evidence supports the fact

that *T. gondii* structural proteins and virulence factors are expressed "just-in-time", from mRNA to protein

without delay, nearly all major cell cycle transitions operate at the post-translational level (1, 10, 20, 52).

For example, SAC is executed through an order of phosphorylation and dephosphorylation events that

are impossible to detect directly at the mRNA level (20, 53). Similarly, the G_2/M transition in tachyzoites

is primarily regulated by phosphorylation, since we did not detect major changes in protein expression

levels in the checkpoint-arrested populations (10). Although recent single cell mRNA sequencing

(scRNAseq) analyses can precisely detect changes in mRNA expression in *T. gondii*, the cell cycle

phases in these data analyses were approximated, and assigned based on the only available study that

described the relative mRNA expression of known cell cycle markers in a thymidine kinase-dependent

cell cycle block in early S-phase (1, 54). Our *Toxo*FUCCI approach allowed us to reconstruct the

tachyzoite cell cycle, which reveals substantial differences to the currently used cell cycle map (1). While

mRNA-based studies cannot be used to mark all activities and stages of the cell cycle, they are

nonetheless valuable and can be fortified by the information that can be provided from post-translational

systems like FUCCI.

A critical question of apicomplexan cell biology had been posed in 2008: is apicomplexan cell

division closely woven, or loosely stitched (55)? Recent reports, and our current findings, suggest a

relaxed control of the apicomplexan cell cycle, which can overall benefit parasite survival.

MATERIALS AND METHODS

Cell Culture

T. gondii RH*ΔKu80Δhxgprt OsTIR1* strain was maintained in Human foreskin fibroblasts (HFF) (ATCC,

SCRC-1041) cells in DMEM media (Millipore Sigma). Cells were incubated at 37°C and 5% CO_2. Host

cells and parasite strains were tested for mycoplasma contamination using a Mycoplasma detection PCR

kit (MP Biomedicals).

Generation of Transgenic Strains

All transgenic strains and primers created in the current study are described in Table S1.

<u>Endogenous C-terminal tagging</u>. To create conditional expression models of TgPCNA1 and TgCenP-C,

we amplified a fragment of the 3'- end of the gene of interest using PCR from *Toxoplasma* genomic DNA

template. The resulting PCR product was then cloned into pLIC-mAID-3xHA-hxgprt using Gibson

Assembly. The final constructs were linearized within the region amplified from genomic DNA template

with unique endonucleases and transfected into RH*ΔKu80Δhxgprt OsTIR1* parent. A similar approach

was used to build TgCenP-C with 3xmyc epitope tag and TgBCC0 with SM-myc tag. Linearized pLIC-

TgBCC0-SM-myc-CAT construct was transfected into RH Tati*ΔKU80* strain expressing TgPCNA1[NG].

<u>Full ORF gene expression.</u> To create *Toxo*FUCCI^S, we amplified the full ORF of the gene of interest by

PCR from genomic DNA and cloned the amplicons in pLIC-2x Neon^{Green} DHFR. Constructs were

linearized with endonucleases within the gene fragment and electroporated in RH*ΔKu80Δhxgprt*

OsTIR1 parent. The same approach was applied to the construction of LIC IMC3 3x mCherry DHFR.

Dual transgenic strains were developed in a similar fashion and sequentially electroporated into parental

transgenic strains. Constructs that were incorporated in one parent contained differing drug resistance

gene cassettes for selection and screening.

<u>Transfection and selection of transgenic parasites.</u> Freshly lysing parasites were collected, mixed with 50

µg of DNA, and transfected using the Amaxa Nucleofector II device (Lonza) with 100 µl Cytomix buffer

supplemented with 20 mM ATP and 50 mM reduced glutathione. Parasites were then incubated in

regular media for 24 hours at 37°C prior to drug selection. Drug-resistant polyclonal populations were

cloned in 96-well plates by limited dilutions, and clones were verified using immunofluorescence assays

or genomic analyses. We used PCR to verify that tags are incorporated at the appropriate genomic loci

using gene-specific and epitope tag-specific primers where applicable. The expression of tagged proteins

was also confirmed through Western blot and immunofluorescence microscopy analyses.

Plaque Assay

Confluent HFF monolayers were inoculated with freshly egressed parasites grown in presence of either

ethanol or 500 µM Indole-3-Acetic Acid (IAA, auxin) (Millipore Sigma), which prompts degradation of

mAID tagged proteins. One week post infection, infected cells were fixed with 100% methanol (Fisher

Scientific), stained with Crystal Violet, and counted. Three biological replicates of each assay were

performed.

Western Blot

Protein samples derived from isolated parasites or protein extracts were mixed with Laemmli sample

buffer, heated at 95°C for 10 minutes, and sonicated. After separation on SDS-PAGE gels, proteins were

transferred onto nitrocellulose gels and probed with following antibodies: rat α-HA (clone 3F10; Roche

Applied Sciences), rabbit α-mNeonGreen (Cell Signaling Technology), mouse α-GRA7 (kindly provided

by Dr. Peter Bradley, UCLA, CA, USA). Membranes were then incubated with appropriate secondary

antibodies conjugated with horseradish peroxidase (Jackson ImmunoResearch). Signals were detected

using an enhanced chemiluminescent detection kit (Sigma Millipore).

Immunofluorescent Microscopy Analysis

Monolayers infected with tachyzoites were grown under the conditions described above. Cells were

fixed, permeabilized, blocked and treated with antibodies as previously described (10, 20, 37). Primary

antibodies used: rat α-HA (3F10, Roche Applied Sciences), mouse α-Centrin1 (clone 20H5; Millipore

Sigma), rabbit α-Myc (clone 71D10, Cell Signaling), rabbit α-MORN1 (kindly provided by Dr. Marc-Jan

Gubbels, Boston College, MA), mouse monoclonal α-ISP1 (clone 7E8; kindly provided by Dr. Peter

Bradley, UCLA), mouse and rabbit α-IMC1 (kindly provided by Dr. Gary Ward, University of Vermont,

VA), and mouse α-Tubulin A (clone 12G10; DSHB). Alexa-conjugated secondary antibodies of different

emission wavelengths (Thermo Fisher) were used at 1:500 dilution. Nuclei were stained with 4',6-

diamidino-2-phenylindole (DAPI, Sigma). Mounted coverslips were imaged on Zeiss Axiovert Observer

Microscope equipped with a 100✕ objective and an ApoTome slicer. The images were processed in

Zen3.2 and Adobe Photoshop 2024 using linear adjustments when required. All experiments were

performed in triplicate. All quantitative IFA replicates were quantified by different investigators to

minimize the effect of subjective bias.

Ultra-Expansion Microscopy

Samples were prepared as previously described (10). Gel slices were probed with mouse α-Tubulin A

(12G10, DSHB; 1:250) and rabbit α-MORN1 (kindly provided by Dr. Marc-Jan Gubbels, Boston College,

MA; 1:100) and corresponding secondary antibodies for 3 hours at room temperature before a second

round of overnight expansion. Gels were placed on 35 mm glass-bottom dishes coated with poly-D-lysine

(Gibco) and were imaged using Zeiss Axiovert Observer Microscope equipped with a 100✕ objective and

an ApoTome slicer.

FACScan Analysis

Parasite DNA content was measured by Flow Cytometry using Propidium Iodide dye as previously described (10). Infected HFF monolayers were grown overnight at 37°C prior to treatment with 500 μM auxin or 3μM SB505124 (Millipore Sigma). Collected parasites were fixed in 300 μL cold PBS and 700 μL ice cold 100% ethanol and stored at -20°C. To stain DNA, samples were treated with 2 mg/mL propidium iodide (Fisher Scientific) and 10 mg/mL RNase A and incubated for 30 minutes in the dark. DNA content was measured based on the intensity of emission from propidium iodide-stained DNA using a PE-Texas Red-A laser. *T. gondii* size-specific parameters were used to segregate parasite population. The percentages of each cell cycle phase were calculated based on the defined gates for each population.

Quantification of Cell Cycle Phases

All quantifications were performed on all biological triplicates of asynchronously growing populations of *Toxo*FUCCI transgenic lines grown for 18 hours at 37°C.

G_1 phase: We quantified the number of centrosomes in no less than 500 individual parasites with diffuse nuclear TgPCNA^{NG} expression. G1 tachyzoites contain a single centrosome. (G_1-50%)

S phase: We quantified no less than 500 individual parasites to determine the proportion of those expressing aggregated nuclear TgPCNA^{NG}. (S phase- 31%)

Centrosome duplication event: We quantified the number of centrosomes in no less than 200 DNA-replicating parasites (aggregated TgPCNA^{NG}). Centrosome duplication event occurs within 3% of S-phase.

S/C overlap: We quantified no less than 250 DNA-replicating parasites (aggregated TgPCNA^{NG}) and determined the fraction of those that were expressing endogenously tagged TgBCC0^{myc}, constituting 40% of S-phase parasites (31%). (S/C-12%)

S interval: Duration of the S-interval was determined by subtracting the S/C interval: 31%-12%= 19%.

<u>Prophase and Metaphase of mitosis:</u> We quantified no less than 500 individual parasites containing

TgMORN1-positive single centrocone with two dBCs. They constituted 12% of the cell cycle.

<u>S/M/C overlap</u>**:** We quantified no less than 100 DNA replicating parasites (aggregated TgPCNA[NG])

containing TgMORN1-positive single centrocone with two dBCs. They constituted 31% of the S-phase.

(S/M/C-9%)

<u>Metaphase of mitosis:</u> Metaphase duration was determined by subtracting the duration of S/M/C interval

from prophase and metaphase duration, 12%-9% = 3%.

<u>Anaphase of mitosis:</u> We quantified no less than 500 individual parasites with two TgMORN1-positive

centrocones, each associated with a single dBC. (Anaphase-12%).

<u>Telophase of mitosis:</u> We quantified no less than 300 individual parasites containing 2 nuclei.

(Telophase-4%)

<u>S/G$_2$/C interval:</u> To determine the duration of the S/G$_2$/C interval, we summed the duration of the S- and

S/M/C intervals and subtracted the result from the S-phase duration. 31%-(19%+9%) =3%

Proteomic Analysis

Samples were prepared from 2x10^9 parental RHΔ*Ku80*Δ*hxgprt OsTIR1* and tachyzoites expressing

endogenously tagged TgPCNA1[AID-HA] as previously described (10, 20). Chromatin-bound TgPCNA1 was

extracted by treatment with TurboNuclease (Accelogen) following the published protocol (56).

TgPCNA1[HA] complexes were isolated using α-HA magnetic beads (MblBio) and compared to nonspecific

proteins bound from parental parasites. Efficiency of immunoprecipitation was verified by Western blot.

Protein samples were processed for mass spectrometry-based proteomic analysis as previously

described (10, 20). Raw data was processed in MaxQuant (v 1.6.14.0, www.maxquant.org) and queried

against the current UniProt *Toxoplasma gondii Me49* protein sequence database. Proteins were

identified using filtering criteria of 1% protein and peptide false discovery rate. Protein intensity values

were normalized using the MaxQuant LFQ function (57). The computational platform SAINT

(Significance Analysis of Interactome) was used to examine TgPCNA1 interactions (58). The platform

utilizes spectral count data and applies a probabilistic model to gauge the likelihood of authentic

interactions between proteins. We used the SAINT express-spc command in version 3.6.3 with default

parameters, and hits with a cutoff range of 0.5 - 1 were deemed significant.

Phylogenetic Analysis

Protein sequences of PCNA1 orthologs were downloaded from the UniProt database and analyzed in

NG phylogeny (https://ngphylogeny.fr/). The custom workflow included Clustal Omega and BMGE to

align and curate sequences, respectively. Maximum likelihood-based inference with Smart Model

Selection was implemented using the PhyML+SMS algorithm. We applied 1000 bootstraps to test

optimization of the tree. Phylogenetic trees were constructed using Newick utilities (59). Sequences of

the proteins analyzed are included in Table S5.

Data Availability

Mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the

PRIDE (60) partner repository with the dataset identifiers: PXD054719 (TgPCNA1 interactome).

Processed proteomics data are available in Supplementary Table S2.

REFERENCES

1. Behnke MS, Wootton JC, Lehmann MM, Radke JB, Lucas O, Nawas J, et al. Coordinated progression through two subtranscriptomes underlies the tachyzoite cycle of Toxoplasma gondii. PLoS One. 2010;5(8):e12354.

2. Saitou T, Imamura T. Quantitative imaging with Fucci and mathematics to uncover temporal dynamics of cell cycle progression. Dev Growth Differ. 2016;58(1):6-15.

3. Sakaue-Sawano A, Kurokawa H, Morimura T, Hanyu A, Hama H, Osawa H, et al. Visualizing spatiotemporal dynamics of multicellular cell-cycle progression. Cell. 2008;132(3):487-98.

4. Zielke N, Edgar BA. FUCCI sensors: powerful new tools for analysis of cell proliferation. Wiley Interdiscip Rev Dev Biol. 2015;4(5):469-87.

5. Herr P, Bostrom J, Rullman E, Rudd SG, Vesterlund M, Lehtio J, et al. Cell Cycle Profiling Reveals Protein Oscillation, Phosphorylation, and Localization Dynamics. Mol Cell Proteomics. 2020;19(4):608-23.

6. Yano S, Hoffman RM. Real-Time Determination of the Cell-Cycle Position of Individual Cells within Live Tumors Using FUCCI Cell-Cycle Imaging. Cells. 2018;7(10).

7. Ando R, Sakaue-Sawano A, Shoda K, Miyawaki A. Two coral fluorescent proteins of distinct colors for sharp visualization of cell-cycle progression. Cell Struct Funct. 2023;48(2):135-44.

8. Yin K, Ueda M, Takagi H, Kajihara T, Sugamata Aki S, Nobusawa T, et al. A dual-color marker system for in vivo visualization of cell cycle progression in Arabidopsis. Plant J. 2014;80(3):541-52.

9.   Zielke N, Korzelius J, van Straaten M, Bender K, Schuhknecht GFP, Dutta D, et al. Fly-FUCCI: A

versatile tool for studying cell proliferation in complex tissues. Cell Rep. 2014;7(2):588-98.

10.   Hawkins LM, Wang C, Chaput D, Batra M, Marsilia C, Awshah D, et al. The Crk4-Cyc4 complex

regulates G(2)/M transition in Toxoplasma gondii. EMBO J. 2024;43(11):2094-126.

11.   Gonzalez-Magana A, Blanco FJ. Human PCNA Structure, Function and Interactions.

Biomolecules. 2020;10(4).

12.   Boehm EM, Gildenberg MS, Washington MT. The Many Roles of PCNA in Eukaryotic DNA

Replication. Enzymes. 2016;39:231-54.

13.   Strzalka W, Ziemienowicz A. Proliferating cell nuclear antigen (PCNA): a key factor in DNA

replication and cell cycle regulation. Ann Bot. 2011;107(7):1127-40.

14.   Guerini MN, Behnke MS, White MW. Biochemical and genetic analysis of the distinct proliferating

cell nuclear antigens of Toxoplasma gondii. Mol Biochem Parasitol. 2005;142(1):56-65.

15.   Farhat DC, Swale C, Dard C, Cannella D, Ortet P, Barakat M, et al. A MORC-driven

transcriptional switch controls Toxoplasma developmental trajectories and sexual commitment. Nat

Microbiol. 2020;5(4):570-83.

16.   Harris MT, Jeffers V, Martynowicz J, True JD, Mosley AL, Sullivan WJ, Jr. A novel GCN5b lysine

acetyltransferase complex associates with distinct transcription factors in the protozoan parasite

Toxoplasma gondii. Mol Biochem Parasitol. 2019;232:111203.

17.   Alvarez V, Vinas L, Gallego-Sanchez A, Andres S, Sacristan MP, Bueno A. Orderly progression

through S-phase requires dynamic ubiquitylation and deubiquitylation of PCNA. Sci Rep. 2016;6:25513.

18.   Engelberg K, Bechtel T, Michaud C, Weerapana E, Gubbels MJ. Proteomic characterization of

the Toxoplasma gondii cytokinesis machinery portrays an expanded hierarchy of its assembly and

function. Nat Commun. 2022;13(1):4644.

19.   Gubbels MJ, Vaishnava S, Boot N, Dubremetz JF, Striepen B. A MORN-repeat protein is a

dynamic component of the Toxoplasma gondii cell division apparatus. J Cell Sci. 2006;119(Pt 11):2236-

45.

20.   Hawkins LM, Naumov AV, Batra M, Wang C, Chaput D, Suvorova ES. Novel CRK-Cyclin

Complex Controls Spindle Assembly Checkpoint in Toxoplasma Endodyogeny. mBio. 2022:e0356121.

21.   Anderson-White BR, Ivey FD, Cheng K, Szatanek T, Lorestani A, Beckers CJ, et al. A family of

intermediate filament-like proteins is sequentially assembled into the cytoskeleton of Toxoplasma gondii.

Cell Microbiol. 2011;13(1):18-31.

22.   Brown KM, Long S, Sibley LD. Conditional Knockdown of Proteins Using Auxin-inducible Degron

(AID) Fusions in Toxoplasma gondii. Bio Protoc. 2018;8(4).

23.   Francia ME, Striepen B. Cell division in apicomplexan parasites. Nature reviews Microbiology.

2014;12(2):125-36.

24.   Beck JR, Rodriguez-Fernandez IA, de Leon JC, Huynh MH, Carruthers VB, Morrissette NS, et al.

A novel family of Toxoplasma IMC proteins displays a hierarchical organization and functions in

coordinating parasite division. PLoS Pathog. 2010;6(9):e1001094.

25.   Rhind N. DNA replication timing: Biochemical mechanisms and biological significance. Bioessays.

2022;44(11):e2200097.

26.   Ahmad K, Henikoff S. Centromeres are specialized replication domains in heterochromatin. J Cell

Biol. 2001;153(1):101-10.

27.   Brooks CF, Francia ME, Gissot M, Croken MM, Kim K, Striepen B. Toxoplasma gondii sequesters

centromeres to a specific nuclear region throughout the cell cycle. Proc Natl Acad Sci U S A.

2011;108(9):3767-72.

28.   Farrell M, Gubbels MJ. The Toxoplasma gondii kinetochore is required for centrosome

association with the centrocone (spindle pole). Cell Microbiol. 2014;16(1):78-94.

29.   Machado M, Klaus S, Klaschka D, Guizetti J, Ganter M. Plasmodium falciparum CRK4 links early

mitotic events to the onset of S-phase during schizogony. mBio. 2023;14(4):e0077923.

30.   Frenal K, Jacot D, Hammoudi PM, Graindorge A, Maco B, Soldati-Favre D. Myosin-dependent

cell-cell communication controls synchronicity of division in acute and chronic stages of Toxoplasma

gondii. Nat Commun. 2017;8:15710.

31. Ouologuem DT, Roos DS. Dynamics of the Toxoplasma gondii inner membrane complex. J Cell

Sci. 2014;127(Pt 15):3320-30.

32. Dhara A, de Paula Baptista R, Kissinger JC, Snow EC, Sinai AP. Ablation of an Ovarian Tumor

Family Deubiquitinase Exposes the Underlying Regulation Governing the Plasticity of Cell Cycle

Progression in Toxoplasma gondii. MBio. 2017;8(6).

33. Hu K, Roos DS, Angel SO, Murray JM. Variability and heritability of cell division pathways in

Toxoplasma gondii. J Cell Sci. 2004;117(Pt 23):5697-705.

34. Radke JR, Striepen B, Guerini MN, Jerome ME, Roos DS, White MW. Defining the cell cycle for

the tachyzoite stage of Toxoplasma gondii. Mol Biochem Parasitol. 2001;115(2):165-75.

35. Nishi M, Hu K, Murray JM, Roos DS. Organellar dynamics during the cell cycle of Toxoplasma

gondii. J Cell Sci. 2008;121(Pt 9):1559-68.

36. Lara-Gonzalez P, Westhorpe FG, Taylor SS. The spindle assembly checkpoint. Curr Biol.

2012;22(22):R966-80.

37. Alvarez CA, Suvorova ES. Checkpoints of apicomplexan cell division identified in Toxoplasma

gondii. PLoS Pathog. 2017;13(7):e1006483.

38. Chen CT, Gubbels MJ. The Toxoplasma gondii centrosome is the platform for internal daughter

budding as revealed by a Nek1 kinase mutant. J Cell Sci. 2013;126(Pt 15):3344-55.

39. Radke JR, White MW. A cell cycle model for the tachyzoite of Toxoplasma gondii using the

Herpes simplex virus thymidine kinase. Mol Biochem Parasitol. 1998;94(2):237-47.

40. Brown KM, Suvorova E, Farrell A, McLain A, Dittmar A, Wiley GB, et al. Forward genetic

screening identifies a small molecule that blocks Toxoplasma gondii growth by inhibiting both host- and

parasite-encoded kinases. PLoS Pathog. 2014;10(6):e1004180.

41. Suvorova ES, Francia M, Striepen B, White MW. A novel bipartite centrosome coordinates the

apicomplexan cell cycle. PLoS Biol. 2015;13(3):e1002093.

42. Bunnik EM, Venkat A, Shao J, McGovern KE, Batugedara G, Worth D, et al. Comparative 3D

genome organization in apicomplexan parasites. Proc Natl Acad Sci U S A. 2019;116(8):3183-92.

43. Nasmyth K. A prize for proliferation. Cell. 2001;107(6):689-701.

44.   Nurse P. A long twentieth century of the cell cycle and beyond. Cell. 2000;100(1):71-8.

45.   Ivanova T, Maier M, Missarova A, Ziegler-Birling C, Dam M, Gomar-Alba M, et al. Budding yeast

complete DNA synthesis after chromosome segregation begins. Nat Commun. 2020;11(1):2267.

46.   Meitinger F, Palani S. Actomyosin ring driven cytokinesis in budding yeast. Semin Cell Dev Biol.

2016;53:19-27.

47.   Rezvani Y, Keroack CD, Elsworth B, Arriojas A, Gubbels MJ, Duraisingh MT, et al. Comparative

single-cell transcriptional atlases of Babesia species reveal conserved and species-specific expression

profiles. PLoS Biol. 2022;20(9):e3001816.

48.   Naumov A, Kratzer S, Ting LM, Kim K, Suvorova ES, White MW. The Toxoplasma Centrocone

Houses Cell Cycle Regulatory Factors. MBio. 2017;8(4).

49.   White MW, Suvorova ES. Apicomplexa Cell Cycles: Something Old, Borrowed, Lost, and New.

Trends Parasitol. 2018;34(9):759-71.

50.   Kisielewska J, Lu P, Whitaker M. GFP-PCNA as an S-phase marker in embryos during the first

and subsequent cell cycles. Biol Cell. 2005;97(3):221-9.

51.   Klaus S, Binder P, Kim J, Machado M, Funaya C, Schaaf V, et al. Asynchronous nuclear cycles in

multinucleated Plasmodium falciparum facilitate rapid proliferation. Sci Adv. 2022;8(13):eabj5362.

52.   King RW, Deshaies RJ, Peters JM, Kirschner MW. How proteolysis drives the cell cycle. Science.

1996;274(5293):1652-9.

53.   Manic G, Corradi F, Sistigu A, Siteni S, Vitale I. Molecular Regulation of the Spindle Assembly

Checkpoint by Kinases and Phosphatases. Int Rev Cell Mol Biol. 2017;328:105-61.

54.   Xue Y, Theisen TC, Rastogi S, Ferrel A, Quake SR, Boothroyd JC. A single-parasite

transcriptional atlas of Toxoplasma gondii reveals novel control of antigen expression. Elife. 2020;9.

55.   Gubbels MJ, White M, Szatanek T. The cell cycle and Toxoplasma gondii cell division: tightly knit

or loosely stitched? Int J Parasitol. 2008;38(12):1343-58.

56.   Li X, Wang W, Wang J, Malovannaya A, Xi Y, Li W, et al. Proteomic analyses reveal distinct

chromatin-associated and soluble transcription factor complexes. Mol Syst Biol. 2015;11(1):775.

57.   Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M. Accurate proteome-wide label-free

quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. Mol Cell

Proteomics. 2014;13(9):2513-26.

58.   Choi H, Larsen B, Lin ZY, Breitkreutz A, Mellacheruvu D, Fermin D, et al. SAINT: probabilistic

scoring of affinity purification-mass spectrometry data. Nat Methods. 2011;8(1):70-3.

59.   Junier T, Zdobnov EM. The Newick utilities: high-throughput phylogenetic tree processing in the

UNIX shell. Bioinformatics. 2010;26(13):1669-70.

60.   Perez-Riverol Y, Bai J, Bandla C, Garcia-Seisdedos D, Hewapathirana S, Kamatchinathan S, et

al. The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences.

Nucleic Acids Res. 2022;50(D1):D543-D52.

FIGURE LEGENDS

Figure 1. Chromatin-bound *Toxoplasma* PCNA1 interacts with DNA replication machinery.

(A) Fluorescence microscopy analysis of TgPCNA1^{AID-HA} expression in RHΔ*Ku80TIR1* tachyzoites. The

images show co-expression of TgPCNA1^{AID-HA} and centrosome (α-Centrin1/ α-mouse IgG Fluor 568), or

inner membrane complex (α-TgIMC1/ α-rabbit IgG Fluor 568). Cell cycle phases were determined based

on numbers and morphology of the relevant reference structures, as indicated with arrows.

(B) Log2 values of protein spectra detected by mass-spectrometry analysis of TgPCNA1^{AID-HA} complexes

that were chromatin-bound or unbound plotted on graphs. Different color dots represent categories of the

selected TgPCNA1 interactors. Experiments were performed in two biological replicates.

(C) Select set of proteins interacting with chromatin-bound TgPCNA1.

Figure 2. TgPCNA1 controls progression of DNA replication in tachyzoites.

(A) Western Blot analysis of the total lysates of RHΔ*Ku80TIR1* tachyzoites expressing TgPCNA1^{AID-HA}.

Lysates of non-treated parasites, and parasites treated with 500μM IAA for 30 or 90 minutes were

analyzed. Western blots were probed with α-HA (α-rat IgG-HRP) and α-GRA7 (α-mouse IgG-HRP) to

confirm equal loading of total lysates.

(B) Images of stained HFF monolayers infected with RHΔ*Ku80TIR1* TgPCNA1^{AID-HA} tachyzoites grown

with or without 500μM IAA for 7 days. Only TgPCNA1-expressing tachyzoites (-IAA) formed viable

plaques. Experiment was performed in three biological replicates.

(C) FACScan analysis of DNA content obtained from parasites expressing TgPCNA1^{AID-HA} (0, blue plot)

or lacking TgPCNA1^{AID-HA} for 4 hours (+IAA, red plot) and 7 hours (+IAA, orange plot). The results of one

of three independent experiments are shown. Dashed lines indicate the prominent 1.8N DNA content

peak.

(D) Quantification of the G_1 (single centrosome) and S/G_2/M/budding (2 centrosomes) populations. The

TgPCNA1^{AID-HA} -expressing and -deficient (4 and 7 hours with 500μM IAA) parasites were co-stained

with α-Centrin1, α-TgIMC1, and DAPI. 100 random parasite vacuoles were evaluated in three

independent experiments. Mean -/+ SD values are plotted on the graphs.

(E and F) Ultra-expansion microscopy analysis of RHΔ*Ku80TIR1* TgPCNA1^{AID-HA} -expressing (-IAA) and -

deficient (+IAA) tachyzoites. The -IAA images show the 8h mock treated parasites. Centriole (cntl)

counts and localization were evaluated by Centrin1 (α-Centrin/α-mouse IgG Fluor 568) stain. Tubulin A

(α-TubulinA/α-mouse IgG Fluor 568) stains subpellicular MTs (daughter cell scaffold, DCS) and

centrioles (spindle poles). TgMORN1 (α-MORN1/α-rabbit IgG Flour 488) staining shows changes in

morphology of the centrocone and the number of daughter basal complexes (dBC). Nuclei stained with

DAPI (blue). Experiment was performed in three biological replicates.

Figure 3. Characterizing the *Toxo*FUCCI probes.

(A) Fluorescence microscopy analysis of TgPCNA1^{NG} expression in RHΔ*Ku80TIR1* tachyzoites. Images

show immunofluorescence co-expression of TgPCNA1^{NG} and markers for centrosome (α-Centrin1/ α-

mouse IgG Fluor 568), centrocone (α-TgMORN1/ α-rabbit IgG Flour 568), or apical cone (α-TgISP1/ α-

mouse IgG Fluor 568). Guide panel depicts the marker used in grey. Cell cycle phases were determined

based on the numbers and morphology of the relevant reference structures, as indicated with arrows.

(B) Schematics of the tachyzoite cell cycle showing the temporal expression of $Toxo$FUCCI^{SC}

fluorophores.

(C and D) Flow cytometry analysis of $Toxo$FUCCI^{SC} expressing TgPCNA1^{NG} and TgIMC3^{mCh}. The plot

(C) shows distribution of populations differentially expressing TgPCNA1^{NG} (green) and TgIMC3^{mCh} (red).

Cell cycle phases were predicted based on the relative expression of $Toxo$FUCCI^{SC} markers. Data was

analyzed using FlowJo and the results of one of three independent experiments are shown. The plot (D)

shows correlation between the number of cells (events) and expression of the $_{Toxo}$FUCCI^{SC} markers.

(E) Western Blot analysis of total lysates of the parental RH$\Delta$$Ku80TIR1$ and $Toxo$FUCCI^{S} tachyzoites.

Membranes were probed with α-NeonGreen (α-rabbit IgG-HRP) to detect TgPCNA1^{NG} expression and

α-GRA7 (α-mouse IgG-HRP) to confirm equal loading of the total lysates.

(F) Immunofluorescent microscopy analysis of $Toxo$FUCCI^{S} tachyzoites expressing inner kinetochore

marker TgCenP-C^{AID-HA}. Parasites were stained with α-HA (α-rat IgG Flour 568) antibody to detect

TgCenP-C^{AID-HA} in the first replication forks (TgPCNA1^{NG}). Colocalization of the markers is shown in the

enlarged images on the side.

Figure 4. Real-time dynamic of DNA replication in tachyzoites.

(A) Schematics of the DNA replication initiation.

(B) The beginning of DNA replication. Images of the $Toxo$FUCCI^{S} probe were taken every 30 seconds.

The arrow indicates the development of the first replication aggregate.

(C) Expansion of DNA replication forks. Images show changes in the ring-like structure of the replication

aggregates.

(D) The middle of DNA replication. The images show the dynamic rearrangements of the replication foci

that were recorded for 3 minutes.

(E) The end of DNA replication. The series of images depict the offset of DNA replication (last focus).

Note the proximal position of the nucleus and conoidal accumulation of $Toxo$FUCCI^{S} probe in parasites

completing DNA replication.

Figure 5. The intravacuolar asynchrony of the tachyzoite cell cycle.

(A) Images of *Toxo*FUCCI^S parasites expressing the kinetochore marker TgCenP-C^{AID-HA} during early, mid-, and late DNA replication. Representative images of the first three intravacuolar divisions are shown. Stars indicate parasites undergoing DNA replication. The number on the top is an average of the vacuoles containing asynchronous parasites. Schematic on the left shows the periods of the cell cycle represented on the images. The panel of enlarged images on the right depicts different stages of DNA replication in 4 parasites from the same vacuole. Note that replication starts around centromeres (kinetochores) and moves away from centromeres as it progresses to the middle stages.

(B) Quantifications of intravacuolar cell cycle asynchrony in parasites beginning (first replication dot) and completing (last replication dot) DNA replication. 100 random vacuoles of the indicated size containing parasites actively replicating DNA (aggregated TgPCNA1^{NG}) were examined in three independent experiments. Asynchronous vacuoles contained at least one parasite lacking aggregated TgPCNA1^{NG}. Mean -/+ SD values are plotted on the graph.

(C) Quantifications of intravacuolar asynchrony across the entire cell cycle. 100 random vacuoles of the indicated size were examined in three independent experiments. Asynchronous vacuoles contained at least one parasite lacking aggregated TgPCNA1^{NG}. Mean -/+ SD values are plotted on the graph.

(D) Distribution of vacuole size of the *Toxo*FUCCI^S parasites and its parental strain asynchronously grown for 24 hours. The number of parasites per vacuole (shown on the right of the graph) was quantified in a minimum of 300 random vacuoles in three independent experiments. Mean -/+ SD values are plotted on the graph.

Figure 6. Organization of the *T. gondii* tachyzoite cell cycle.

(A) Immunofluorescence microscopy images of TgPCNA1^{NG} (green) and selected cell cycle markers (red) visualized with antibodies against Centrin1 (α-mouse IgG Fluor 568), TgMORN1 (α-rabbit IgG Flour 568), and myc-epitope (α-rabbit IgG Fluor 568). Images are aligned with the corresponding cell cycle intervals shown in panel B. Green double-head arrows show temporal TgPCNA1 expression in replication foci and on the conoid.

(B) Schematic representation of cell cycle phases and intervals inferred from quantitative microscopy analyses of *Toxo*FUCCI^S line and its derivates. The numbers on the top reflect the duration of the intervals in minutes (assuming the division cycle takes 7h) and as a percentage of total time. Dotted lines indicate the putative location of the event. The processes representing conventional cell cycle phases are labeled and shown in green. The red hexagonal signs mark the positions of known checkpoints.

(C) Key events of the composite cell cycle phase. Temporal order was deduced by quantitative microscopy analyses or published studies.

(D) Table summarizing the approaches used to identify mitotic subphases.

Figure 7. The *Toxo*FUCCI approach identifies cell cycle checkpoints.

(A) Schematic representation of cell cycle phases shows the changes in tachyzoite populations conditionally arrested at the indicated checkpoints.

(B and F) Immunofluorescence microscopy images of *Toxo*FUCCI^S +TgCrk4^{AID-HA} (B) and *Toxo*FUCCI^S + TgCrk6^{AID-HA} (F) parasites treated with 500µM auxin for 7 hours.

(C and G) Quantification of the vacuoles of *Toxo*FUCCI^S +TgCrk4^{AID-HA} (C) and *Toxo*FUCCI^S + TgCrk6^{AID-HA} (G) tachyzoites containing aggregated TgPCNA1^{NG}. 100 random vacuoles were examined in three independent experiments. Mean -/+ SD values are plotted on the graph.

(D and H) FACScan analysis of DNA content obtained from *Toxo*FUCCI^S parasites expressing (0, blue plots) or lacking +TgCrk4^{AID-HA} (D) and TgCrk6^{AID-HA} (H) for 4 hours (+IAA, red plot) and 7 hours (+IAA, orange plot). The results of one of three independent experiments are shown. Dashed lines indicate the prominent 1.8N DNA content peak.

(E) Flow cytometry analysis of *Toxo*FUCCI^S +TgCrk4^{AID-HA} (C) and *Toxo*FUCCI^S + TgCrk6^{AID-HA} (G) tachyzoites grown at the indicated conditions. Plots show changes in TgPCNA1^{NG} expression and DNA content (PI). Data was analyzed in FlowJo10.10 and the results of one of three independent experiments are shown.

Figure 8. Mapping TgMAPK1-mediated cell cycle arrest using *Toxo*FUCCI^S model.

(A) Immunofluorescence microscopy images of *Toxo*FUCCI^S tachyzoites with or without 3μM SB505124

treatment for 7 hours, and *Toxo*FUCCI^S + TgMAPK1^{AID-HA} parasites with 500μM auxin treatment for 7

hours. Centrosomes were visualized with antibodies targeting Centrin1 (α-mouse IgG Fluor 568). Green

fluorescence indicates aggregated and conoidal TgPCNA1^{NG}. Blue DAPI stain shows the number of

nuclei per cell.

(B) FACScan analysis of DNA content of *Toxo*FUCCI^S parasites with mock treatment (blue plot) or 4

hours (red plot) and 7 hours (orange plot) treatment with 3μM SB505124. The results of one of three

independent experiments are shown. Dashed lines indicate the prominent 1.8N DNA content peak.

(C) Quantification of *Toxo*FUCCI^S parasites containing aggregated TgPCNA1^{NG} with mock treatment, or

after 4 and 7 hours of 3μM SB505124 treatment. 500 individual parasites were examined in three

independent experiments. Mean -/+ SD values are plotted on the graph.

(D) Quantification of *Toxo*FUCCI^S parasites with two nuclei per cell with mock treatment or after 4 and 7

hours treatment with 3μM SB505124. The graph shows the distribution of non-DNA replicating parasites

undergoing budding (blue bar), parasites with two nuclei containing aggregated TgPCNA1^{NG} (light

green), and parasites with one or two nuclei undergoing DNA replication (dark green). 500 individual

parasites were examined in three independent experiments. Mean -/+ SD values are plotted on the

graph.

(E) Quantification of budding *Toxo*FUCCI^S parasites after mock treatment, or 4 and 7 hours of 3μM

SB505124 treatment. The abnormal budding fraction encompasses parasites containing more or less

than 2 buds per cell and altered bud polarity. 500 individual parasites were examined in three

independent experiments. Mean -/+ SD values are plotted on the graph.

(F) Quantification of budding *Toxo*FUCCI^S parasites showing parallel or anti-parallel bud orientation of

the buds after mock treatment, or 4 and 7 hours of 3μM SB505124 treatment. 500 individual parasites

were examined in three independent experiments. Mean -/+ SD values are plotted on the graph.

(G) Ultra-expansion microscopy analysis of *Toxo*FUCCI^S tachyzoites with mock treatment and 3μM

SB505124 treatment for 7 hours. Tubulin A (α-TubulinA/α-mouse IgG Fluor 568) stain shows mother and

daughter cells' subpellicular MTs (daughter cell scaffold, DCS). TgMORN1 (α-MORN1/α-rabbit IgG Flour

488) stain shows the number of daughter basal complexes (dBC). Nuclei stained with DAPI (nc, blue).

Experiment was performed in three biological replicates.

(H) Schematic representation of the tachyzoite cell cycle. Red lines mark the cell cycle processes that

are affected by SB505124 compound. Flathead line indicates inhibition, and arrowhead line indicates the

promoted process.

1031

Figure 9. Organization of the conventional cell cycle and the endodygenic cycle of _T. gondii_

tachyzoites.

The schematic on the left shows four phases of the conventional cell cycle that do not overlap.

Cytokinesis occupies a fraction of mitotic telophase. On the contrary, four different cell cycle phases run

concurrently within the composite cell cycle phase of _T. gondii_ tachyzoites. Despite drastic differences in

cell cycle organization, _Toxoplasma_ endodyogeny preserved three major cell cycle checkpoints (red stop

symbol).

1039

1040   **Figure S1. _Toxoplasma_ PCNA1 is structurally and evolutionary conserved factor.**

1041   (A) Phylogenetic tree of PCNA1-related proteins.

1042   (B) Folding prediction of TgPCNA1 (AlphaFold2). The structural deviations and predicted

1043   phosphorylation sites (ToxoDB.org) are shown.

1044

Figure S2. Construction of the _Toxo_FUCCI transgenic lines.

(A) Schematics for constructing _Toxo_FUCCI[S] parasites. NeonGreen-tagged TgPCNA1 was introduced as

a second copy under the control of the endogenous promoter. Schematics also indicate the relative

positions of primers used to confirm expression of both TgPCNA1 copies (panel B).

(B and D). PCR analysis of the parental and indicated transgenic lines. Primer combinations used to

detect either native or recombined loci are shown. (B) _Toxo_FUCCI[S] and _Toxo_FUCCI[SC]. (D)

RHΔ_Ku80TIR1_ + TgPCNA1[AID-HA], _Toxo_FUCCI[S] + TgCrk6[AID-HA], _Toxo_FUCCI[S] + TgCenP-C[AID-HA].

(D) Schematics for constructing TgCrk4 and TgCenP-C AID-modified genes. Targeting plasmid included a 3' fragment of *GOI* genomic locus fused with encoded sequence for mini-version of AID (mAID), 3xHA (HA) epitopes, and the drug-selection marker *hxgprt* gene (grey box). Plasmid linearization with a unique endonuclease induced recombination at the *GOI* locus. Schematics also indicate the relative positions of the primers used to confirm GOI knock-in (panel D).

(E) Live fluorescence microscopy analysis of *Toxo*FUCCI[SC] tachyzoites. The top schematic shows temporal expression of *Toxo*FUCCI markers.

(F) The replication rates of parental and transgenic lines. The average number of parasites per vacuole after 24 hours' growth was quantified in three independent experiments. Mean value -/+ SD are plotted on the graph.

(G) Flow cytometry analysis of uninfected and *Toxo*FUCCI[SC]-infected HFF monolayers. The plot shows a fraction of fragmented host cells. Data was analyzed using FlowJo and the results of one of three independent experiments are shown.

Figure S3. The dynamics of TgPCNA1NG during tachyzoite replication.

(A) The end of DNA replication. The series of images depicts the offset of DNA replication in a vacuole of 4 parasites (last focus). Proximal position of the nucleus and conoidal accumulation of *Toxo*FUCCI[S] probe confirm that parasites are ending DNA replication. Note that individual parasites complete DNA replication at different times.

(B) The 3-hour progression of DNA replication. A series of images taken every 5 minutes shows the onset, progression, and ending of DNA replication in a vacuole of 8 parasites (last focus). Note that individual parasites start and complete DNA replication at different times.

Figure S4. Validation of *Toxo*FUCCI lines.

(A and B) Western Blot analyses of the total lysates of the *Toxo*FUCCI[S] tachyzoites expressing TgCrk4[AID-HA] (A) or TgCrk6[AID-HA] (B). Lysates of untreated parasites and parasites treated with 500 µM

IAA for 4 hours were analyzed. Western blots were probed with α-HA (α-rat IgG-HRP) and α-NeonGreen

(α-rabbit IgG-HRP), and with α-GRA7 (α-mouse IgG-HRP) to confirm equal loading of the total lysates.

(C) Quantification of the budding populations of $Toxo$FUCCI^S + TgCrk4^{AID-HA} and $Toxo$FUCCI$^{S\ +}$

TgCrk6^{AID-HA} tachyzoites during cell cycle block (4 hours with 500 μM IAA). 100 random vacuoles of

parasites were examined for α-TgIMC1-positive internal buds in three independent experiments. Mean -

/+ SD values of three independent experiments are plotted on the graph.

(D) FACScan analysis of DNA content of parasites asynchronously growing (blue), and arrested at G2/M

(red: $Toxo$FUCCI^S + TgCrk4^{AID-HA}, 4 hours with 500 μM IAA) or SAC (green: $Toxo$FUCCI$^{S\ +}$ TgCrk6^{AID-HA},

4 hours with 500 μM IAA$)$ checkpoints. Note the differences in DNA content of the checkpoint-arrested

populations. The results of one of three independent experiments are shown.

(E) Immunofluorescent microscopy analysis of $Toxo$FUCCI^S tachyzoites expressing inner kinetochore

marker TgCenP-C^{AID-HA}. Images depict the beginning and end of DNA replication. Parasites were stained

with α-HA (α-rat IgG Flour 568) antibody to detect TgCenP-C^{AID-HA}. The markers' colocalization is shown

in the enlarged image on the side.

(F) Flow cytometry analysis of $Toxo$FUCCI^S tachyzoites grown at the indicated conditions. Plots show no

changes in TgPCNA1^{NG} expression and DNA content (PI). Data was analyzed in FlowJo and the results

of one of three independent experiments are shown.

1095

Table S1. Primers, transgenic strains used in the study, and PCNA1 orthologs.

Table S2. TgPCNA1 proteomes.

Table S3. Growth characteristics of TgPCNA1 AID model.

Table S4. Tachyzoite cell cycle organization.

Table S5. $Toxo$FUCCI real-time microscopy analysis.

Table S6. Intravacuolar asynchrony of $Toxo$FUCCI tachyzoites.

Table S7. Validation of the $Toxo$FUCCI model using checkpoint mutants.

Table S8. Validation of the $Toxo$FUCCI model using drug-induced cell cycle block.

Movie S1. Beginning of DNA replication.

Movie S2. Middle of DNA replication.

Movie S3. End of DNA replication.

Movie S4. Aborted DNA replication.

Figure 1

A TgPCNA1^AID-HA | TgPCNA1^AID-HA Centrin1 DAPI | Cell cycle

aggregated TgPCNA expression

diffused | single cn | *G1*

aggregated | single cn | *S-phase progression*

aggregated | duplicated cn

aggregated | separated cn

TgPCNA1^AID-HA | TgPCNA1^AID-HA TgIMC1 DAPI

diffused | 5μm | *M/C (budding)*

B *Chromatin bound*

TgPCNA1^AID-HA, spectra log2 vs Parent, spectra log2

PCNA1
MCM7
DNA Pol V

Chromatin non-bound

TgPCNA1^AID-HA, spectra log2 vs Parent, spectra log2

PCNA1
MCM7
MCM2

- DNA replication remodeling (red)
- AP2 factors (orange)
- Nuclear pore (green)
- Proteosome (blue)

C *TgPCNA1 interactors on chromatin*

Gene ID TGME49_	Product	SAINT score	Fold Change
DNA replication: MCM complex			
237220	MCM7	1	6.9
214970	MCM2	0.99	5.0
219700	MCM4	0.95	2.9
DNA remodeling: MORC/SAGA complexes			
218960	AP2XII-1	1	250.0
310900	AP2XI-2	1	21.5
305340	MORC	1	18.5
243440	GCN5b	1	10.3
227290	HDAC3	1	6.2

Figure 3

A

	TgPCNA1^NG	Merge Marker / DAPI	Guide	Cell cycle
Centrosome	diffused	Centrin	Single centrosomes	*G₁*
Centromere	aggregated	CenP-C	First replication foci appear near (on) centromeres	*early S phase*
Centrocone	aggregated	MORN1^Cherry	MORN1, cc*, basal complex	*mid-S phase*
Centrocone	aggregated + conoid, conoid	MORN1^Cherry	MORN1, MORN rings, cc*, basal complex	*late S phase mitosis cytokinesis (budding)*
Apical cap	diffused + conoid, conoid	ISP1	mother AC, daughter AC	*cytokinesis (budding)*

DNA replication

5µm

B *Toxo*FUCCI^SC

TgPCNA1^NG
TgIMC3^mCherry

S — M — C — G₁

C

Fluorescence, NeonGreen vs Fluorescence, mCherry

DNA replication — budding

late S + budding
NG(+) mCh(+)
mid-S
early S
NG(-) mCh(-)
NG(-) mCh(+)
G₁

D

events vs Fluorescence intensity

S   M/C   G₁

NG(-) mCh(-)

E

MW kDA	parent	TgPCNA1^NG	
135			
100		▬	α-NG
32	▬	▬	α-GRA7

F *S phase*

kinetochore TgCenP-C^AID-HA
replication foci TgPCNA1^NG
merge

early — 1st focus — non-duplicated
middle — duplicated
late — segregated   non-segregated

2µm

Figure 4

A — *dot* · *ring* · *expanded ring* · *skirt-like* · *disorganized*

centrosomes
centrocone
kinetochores
nucleus

1st (ring) expansion · 2nd (nuclear) expansion

B — *start of replication*

0' · 0.5' · 1' · 1.5' · 2' · 2.5' · 3'

proximal end
apical end
first focus

* ongoing DNA replication

C — *expansion of replication*

0' · 0.5' · 1' · 1.5' · 2' · 2.5' · 3'

proximal end
apical end

ring · ring · lost organization
skirt-like

D — *middle of replication*

0' · 0.5' · 1' · 1.5' · 2' · 2.5' · 3'

apical end
proximal end

E — *end of replication*

0' · 0.5' · 1' · 1.5' · 2' · 2.5' · 3'

conoidal
apical end
last focus
proximal end

4' · 4.5' · 5' · 5.5' · 6' · 6.5' · 7'

5μm

* completed DNA replication

Figure 5

A

1st **7+/-0**   *2nd* **20.3+/- 5.1**   *3rd* **30.3+/- 5.1**   # *division*
   % *asynchrony mean+/-SD*

G₁

kinetochore centromere

early S-phase

1
2
3
4

S

kinetochore

middle S-phase

C   M

conoidal

late S-phase

conoidal

5μm

TgPCNA1^NG   TgCenP-C^AID-HA   DAPI   * *DNA replicating parasite*

B *early and late S-phase*

Intravacuolar asynchrony, %

0.016
0.046

| parasite, # | 2 | 4 | 8 |
| division, # | 1 | 2 | 3 |

C *across cell cycle*

Intravacuolar asynchrony, %

0.26
5.6e-03

| parasite, # | 2 | 4 | 8 | 16 | >32 |
| division, # | 1 | 2 | 3 | 4 | >5 |

D *across population*

Vacuole number, %

vacuole size

parent   FUCCI

Figure 6

Figure 7

A *ToxoFUCCI checkpoint models*

$TgCrk4^{AID}$ $TgCrk6^{AID}$
7h block
4h block

G_2/M SAC

G_1 S G_2 M$_{ana}$ meta   C

B G_2/M IMC1
FUCCI + TgCrk4^{AID-HA}
1 bud
conoid

C *Toxo*FUCCI^S vacuoles
DNA replication, %
0.008
0.03
0   4   7
auxin, h

D
Unit area, %
0h
4h block
7h block
re-replication
PI intensity
1N   1.8N   3.1N

F SAC   MORN1
FUCCI + TgCrk6^{AID-HA}
conoid
1cc
5µm
mBC

G *Toxo*FUCCI^S vacuoles
DNA replication, %
0.0081
0.81
0   4   7
auxin, h

H
Unit area, %
0h
4h block
7h block
2N
PI intensity
1N   1.8N

E
FUCCI + TgCrk6^{AID-HA}
0h, auxin
Fluorescence, NeonGreen
NG(+)   NG(+)
1N DNA   ~2N DNA
62.6%   8.4%
24%   5%
NG(-)   NG(-)
1N DNA   ~2N DNA
PI intensity

FUCCI + TgCrk4^{AID-HA}
7h, auxin
Fluorescence, NeonGreen
NG(+)   NG(+)
1N DNA   ~2N DNA
replication
39.8%   37%
12%   11.2%
NG(-)   NG(-)
1N DNA   ~2N DNA
PI intensity

FUCCI + TgCrk6^{AID-HA}
7h, auxin
Fluorescence, NeonGreen
NG(+)   NG(+)
1N DNA   ~2N DNA
36.8%   23.2%
29%   11%
NG(-)   NG(-)
1N DNA   ~2N DNA
PI intensity

Figure 8

A

ToxoFUCCI^S
no treatment / SB505124

ToxoFUCCI^S + TgMAPK1^AID-HA
IAA

B

C *DNA replication*

D *DNA rereplication*

E *Budding*

F *Budding polarity*

G
no treatment — early budding / late budding complete karyokinesis
SB505124, 7h — complete karyokinesis early & multiple budding / complete karyokinesis early budding

H
G₁ S G₂/M SAC meta M ana SB505124 uncoupling Cytokinesis DNA rereplication

Figure 9

A

RH-TIR1 parent

TgPCNA1

targeting construct

TgPCNA1 2xNG dhfr-ts

B

kb

parent · transgene · gDNA

1+2 · 1+3 · 1+2 · 1+3 · primer set

2.0
1.0

TgPCNA1^NG

wt · wt · 2nd copy

1.0
0.7

TgIMC3^Ch

wt · wt · 2nd copy

C

RH-TIR1 parent

GOI

nuclease DSB · mAID · HA · hxgprt

targeting construct

D

kb

parent · transgene · gDNA

1+2 · 1+3 · 1+2 · 1+3 · primer set

4.0
2.0

TgPCNA1^AID-HA

ToxoFUCCI^S dual transgenic strains

3.0
2.0

TgCrk6

4.0
2.0

TgCenP-C

E

ToxoFUCCI^SC

TgIMC3^Ch
TgPCNA1^NG

G₁ · S phase · budding

1st foci · aggregated · last foci · buds · conoidal · 5μm

F

parasite per vacuole, %

n.s · n.s · n.s

Parent · ToxoFUCCI^S · RH TgPCNA1^AID-HA

24h growth

G

Fluorescence, NeonGreen

non infected host cells
ToxoFUCCI^sc infected host cells

Fluorescence, mCherry

Figure S3

A

end of replication

0' 0.5' 1' 1.5' 2' 2.5' 3' 3.5' 4' 4.5'

5' 5.5' 6' 6.5' 7' 7.5' 8' 8.5' 9' 9.5'

5μm

∗ completed DNA replication

B

0' 5' 10' 15' 20' 25' 30'

35' 40' 45' 50' 55' 60' 65'

70' 75' 80' 85' 90' 95' 100'

105' 110' 115' 120' 125' 130' 135'

140' 145' 150' 155' 160' 165' 170'

175' 180' 185'

5μm

⬭ ongoing DNA replication

Figure S4

A TgCrk4^AID

MW kDA | +IAA |
0 | 4h
135 — | | α-HA
100 — | | α-NG
32 — | | α-GRA7

B TgCrk6^AID

MW kDA | +IAA |
0 | 4h
135 — | | α-HA
100 — | | α-NG
32 — | | α-GRA7

C
Budding, %

- 0h, IAA
- 4h, IAA

TgCrk4^AID: *0.017*
TgCrk6^AID: *0.0016*

D

Counts vs PI intensity

G₂/M TgCrk4
SAC TgCrk6
asynchronous

1N   1.8N   2N

DNA content

E DNA replication

TgPCNA^NG | TgPCNA^NG TgMORN1 DAPI

beggining:
bc, cc
1st replisome / centrocone

end:
last replisome / centrocone+ rings
bc

5μm

F

FUCCI 0h, auxin

Fluorescence, NeonGreen vs PI intensity

NG(+) 1N DNA: 61.5%
NG(+) ~2N DNA: 14.3%
NG(−) 1N DNA: 20.6%
NG(−) ~2N DNA: 3.6%

FUCCI 7h, auxin

NG(+) 1N DNA: 61%
NG(+) ~2N DNA: 14.3%
NG(−) 1N DNA: 20.4%
NG(−) ~2N DNA: 4.12%