β-glucan reprograms alveolar macrophages via neutrophil/IFNγ axis to promote lung injury

Renaud Prével[a], Erwan Pernet[a], Kim A Tran[a], Abderrahmane Sadek[b,c], Mina Sadeghi[a],

Elizabeth Lapshina[a], Leonardo Jurado[a], Arnold S. Kristof[a], Mohieddine Moumni[b], Jérémie

Poschmann[c], Maziar Divangahi[a,d,e,f,*]

[a] Department of Medicine, Meakins-Christie Laboratories, Research Institute McGill

University Health Centre, McGill University, Montreal, Quebec, Canada

[b]Biotechnology and Bioresources Valorization Laboratory, Biology Department, Faculty of

Sciences, Moulay Ismail University of Meknès, Meknès, Morocco

[c]INSERM, Nantes Université, Center for Research in Transplantation and Translational

Immunology (CR2TI), UMR 1064, ITUN, Nantes, France

[d]Department of Microbiology and Immunology, Meakins-Christie Laboratories, Research

Institute McGill University Health Centre, McGill University, Montreal, Quebec, Canada

[e]Department of Pathology, Research Institute McGill University Health Centre, McGill

University, Montreal, Quebec, Canada

[f]McGill International TB Centre, Montreal, Quebec, Canada

*Corresponding author: maziar.divangahi@mcgill.ca

Keywords: acute lung injury; trained immunity; alveolar macrophages; neutrophils

ABSTRACT

Alveolar macrophages (AMs) reside in the lower airways and play a crucial role in lung health and response to sterile inflammation and infections. AMs possess remarkable adaptability to different environmental challenges that can persist through their memory capacity (trained immunity). β-glucan has been characterized as a potent inducer of central trained immunity by reprogramming hematopoietic stem cells (HSCs) in the bone marrow. In the present study, we show that systemic administration of β-glucan induces peripheral trained immunity by reprogramming alveolar macrophages (AMs) in the lungs, in a Dectin1-independent manner. We furthermore demonstrate that AM reprogramming at both the transcriptional and metabolic levels exacerbate lung injury following bacterial (LPS) or viral (polyI:C) challenges via a neutrophil/IFN-γ dependent manner. These findings identify an additional facet of β-glucan in trained immunity involving AM reprogramming and shed light on the potential detrimental effects of trained immunity.

INTRODUCTION

The lungs represent a significant infective niche for numerous infectious agents. However, severe pulmonary disease is increasingly being associated with immune-mediated responses rather than direct damage of invading microbes [1, 2]. Among the various triggers of immunopathology, lipopolysaccharide (LPS), a component of the outer membrane of Gram-negative bacteria, is a potent inducer of systemic inflammation via activation of Toll-like receptor 4 (TLR4)[3]. Excessive inflammation associated with LPS can cause acute lung injury (ALI), a severe inflammatory condition of the lungs characterized by alveolar damage, edema, and impaired gas exchange leading to organ dysfunction, and ultimately, mortality[4, 5]. Thus, the magnitude of inflammatory responses must be regulated to resolve infections while preventing collateral tissue damage. Understanding the mechanisms underlying the regulation of inflammation in ALI is essential for the development of effective therapeutic interventions. Key players in this regulatory network include various immune cells, cytokines, and signalling pathways, which collectively modulate the intensity and duration of inflammatory responses. Dysregulation of these mechanisms can lead to either persistent inflammation or immunosuppression, both of which contribute to the pathogenesis of sepsis [6, 7, 8].

The constant exposure of the lungs to a non-sterile environment leads to the development of a unique immunity by eliminating inhaled foreign particles while minimizing inflammatory-mediated tissue damage. Resident alveolar macrophages (AMs) serve a crucial regulatory function to maintain the delicate balance between inflammation and protecting lung tissue from damage [9]. AMs originate from fetal liver monocytes and act as vigilant sentinels protecting the airways against invading pathogens and pollutants[10] [11]. Thus, AMs play a pivotal role in orchestrating both the initiation and resolution of immune responses within the lung microenvironment.

Adaptations in innate immune cells are diverse with substantial plasticity to various insults that

can be maintained resulting in enhanced (trained immunity) or reduced (tolerance)

inflammatory responses to a second stimuli [12, 13, 14]. Trained immunity is mediated *via* long-

term metabolic reprogramming and epigenetic modifications, which can be induced by various

stimuli such as the attenuated mycobacteria Bacille Calmette-Guérin (BCG) or β-glucan (a

polysaccharidic component of fungi cell wall)[12, 15]. BCG and β-glucan are able to train

monocytes and neutrophils with beneficial impact in cancer or infections *via* the

reprogramming of haematopoietic stem cells (HSC) within the bone marrow[16, 17, 18, 19, 20].

However, there are substantial knowledge gaps in our understanding of trained immunity

within tissue-resident macrophages (TRMs) and its regulatory role in maintaining tissue

homeostasis under stress conditions. For instance, while the initial LPS-stimulation of

macrophages induces a strong inflammatory responses via TLR-4, restimulation with LPS

generates tolerance in these macrophages[21]. Thus, the epigenetic reprogramming of innate

immune cells and their subsequent response depends on two signals: the initial training agent

and the nature of the second stimuli at an inflammatory site. Here we showed that systemic

administration of β-glucan can reprogram AMs in the lungs via neutrophil/type II IFN axis but

independent of Dectin-1 signalling. The unique transcriptomic and metabolic profile of AMs

render them hyperresponsive to both bacterial or viral stimulation causing dysregulated

inflammatory responses with pulmonary damage. The differences in the systemic basal levels

of β-glucan in sepsis patients as well as the pulmonary levels of IFNγ can be major factors in

determining the hyperresponsiveness in sepsis patients and can be potentially targeted for

therapy.

RESULTS

β-glucan-mediated trained immunity aggravates LPS-induced ALI.

To investigate whether systemic administration of β-glucan induces trained immunity within the lung, we sought to assess its impact on Acute Lung Injury (ALI) triggered by LPS treatment. We initially evaluated the consequences of β-glucan-mediated training on ALI by administering LPS seven days after training (Figure 1A). ALI assessment conducted 24 hours post-LPS instillation did not reveal any discernible distinctions attributable to the initial i.p. β-glucan injection. However, ALI was notably exacerbated in mice that had undergone β-glucan training upon subsequent LPS stimulation as demonstrated non-invasively by lung microCT scanners with regards to the increased proportion of poorly- or non-aerated lung segment and to a significant increase in Hounsfield units (Figure 1B). Hounsfield units reflect a coefficient of tissue attenuation and, by extrapolation, tissue composition. Consistent findings were corroborated through two complementary invasive evaluations of ALI. Specifically, the heightened alveolar-capillary permeability following LPS instillation in β-glucan-trained mice was substantiated by elevated lung Evans blue dye concentrations (Figure 1C). Histological examination further unveiled a spectrum of notable alterations following LPS instillation, including cellular infiltration, thickened alveolar walls, and the formation of hyaline membranes, observed in β-glucan-trained mice and not PBS controls (Figure 1D). We additionally monitored weight loss following LPS instillation (Supplementary Figure 1A). ALI-induced inflammation peaks at 24h, reflected in the weight loss of the mice which drops nearly 10% within a day. While there were no differences in weight loss between control and β-glucan-treated mice at this timepoint, we observed that β-glucan mice exhibited a delayed recovery period given they did not return to their original weight by day 4, as opposed to control mice (Supplementary Figure 1B).

The increased cell infiltration 24 hours after LPS instillation in β-glucan-treated mice was due

to a significant recruitment of neutrophils, as assessed in the bronchoalveolar lavage (BAL;

CD11b$^+$Ly6G$^+$). Strikingly, the frequency and total numbers of neutrophils was doubled in the

β-glucan-treated mice, after LPS administration (Figure 1E). Concomitant with the increased

neutrophil recruitment, CXCL1 was significantly increased in β-glucan-treated mice when

compared to control mice after LPS treatment (Figure 1F)[22]. Amplified ALI in β-glucan-

trained mice was also associated with increased proinflammatory cytokines (IL-6 and TNFα)

(Figure 1G-H). As AMs are major producers of CXCL1, IL6 and TNFα, we next assessed

whether β-glucan increased the proportion or number of AMs and found no significant

differences between naïve and β-glucan trained mice (Figure 1I). There was also no changes in

the frequency of CD11b$^+$ AMs at baseline and post-LPS instillation between β-glucan-treated

and control mice (Supplementary Figure 1C). This suggests that β-glucan does not induce the

expansion of monocyte-derived AMs. To further characterize AM phenotype following β-

glucan, we assessed several AM-associated cell surface markers and found no differences in

expression of MHCII, F4/80, CD169 and CD69 (Supplementary Figure 1D). Interestingly, β-

glucan enhanced the frequency of CD80$^+$ AMs, a costimulatory molecule associated with

activation and pro-inflammatory responses (Supplementary Figure 1D)[23]. Thus the increased

cytokine production was not due to an increased number of AMs, but rather intrinsic functional

changes.

Next, we examined whether the effects of β-glucan-induced immune training persisted in ALI

severity. To do so, we administered β-glucan or PBS (control) in mice and after 28 days

challenged them with LPS (Supplementary Figure 1E). Similar to the 7-day timeframe of β-

glucan training, there was increased endothelial permeability of the lungs (Supplementary

Figure 1F), and increased immunopathology (Supplementary Figure 1G) in 28-days β-glucan

trained mice challenged with LPS. This increased ALI in β-glucan treated mice after 28 days

was associated with a higher concentration of CXCL1, IL6 and TNFα (Supplementary Figure

1H) and increased neutrophil infiltration in BAL (Supplementary Figure 1I) following LPS

instillation. The frequency and number of AMs were not changed between groups

(Supplementary Figure 1J). Increased production of cytokines was also observed upon *ex* vivo

stimulation of AMs isolated from treated mice 28-day post-β-glucan (Supplementary Figure

1K, L). Thus, the intrinsic functional changes in AMs suggest that the exacerbation of ALI

induced by β-glucan triggered a long-term reprogramming of immune cells rather than an

additive effect of lingering inflammation from β-glucan injection. To assess whether this

response was specific to bacterial LPS or viral agonists can cause similar responses, we next

challenged β-glucan trained mice with a TLR-3 agonist (poly(I:C)) (Supplementary Figure

2A). Similar to the LPS-model, β-glucan treated mice had heightened poly(I:C)-induced ALI

shown via increased alveolar-capillary permeability, tissue damage (Supplementary Figure 2A-

C). Pro-inflammatory cytokines CXCL1 and IL-6 were also increased while TNFα remains

unchanged following challenge suggesting differing cytokine profiles between LPS and

Poly(I:C)-induced ALI (Supplementary Figure 2D). However, β-glucan-treated mice similarly

had increased neutrophil infiltration following Poly(I:C) instillation (Supplementary Figure

2E). The number of AMs was also unchanged among groups (Supplementary Figure 2F).

Collectively, these results suggest that systemic β-glucan can maintain a long-term

reprogramming in AMs promoting ALI.

β-glucan augmented ALI is mediated via AM

Although there were no differences in the frequency or absolute number of AMs between β-

glucan-treated and control mice, the early heightened response with increased CXCL1

production and the recruitment of neutrophils indicate that the AMs are engaged in the

exacerbation of ALI induced by β-glucan. To further characterize the role of these cells in ALI,

we locally depleted AMs using intranasally administered clodronate liposomes 2 days before

performing the LPS instillation, which is at its peak of AM depletion (Figure 2A). Depletion

of AMs significantly reduced the production of cytokines TNFα, IL-6, and CXCL1 and the

recruitment of neutrophils in BAL, which abolished the β-glucan-induced ALI (Figure 2B-D).

To further confirm that tissue-resident AMs are responsible for the increased β-glucan-induced

ALI, we used *Csf2rb*[-/-] mice (Figure 2E), which do not naturally develop alveolar macrophages

throughout their lifespan but maintain regular levels of bone marrow derived macrophages

(BMDMs) and interstitial macrophages[24]. Similar to the AM depletion, the production of

cytokines and neutrophil recruitment in BAL was similar between β-glucan treated and control

Csf2rb[-/-] mice after LPS administration. Importantly, there was no difference in ALI, as similar

levels of proteins were measured in the BAL of β-glucan treated and control *Csf2rb*[-/-] after the

LPS challenge (Figure 2F-H).

To complement these loss-of-function experiments with a gain-of-function, we adoptively

transferred AMs from adult WT control (PBS) or trained (β-glucan) mice to day 2 *Csf2rb*[-/-]

neonate mice (Figure 2I) and after 6 weeks challenged these mice with LPS. Remarkably, mice

replenished with AMs from β-glucan-trained mice displayed an increased in production of

inflammatory cytokines and the recruitment of neutrophils into the BAL (Figure 2K, L). This

was associated with increased ALI as they had higher protein levels in their BAL compared to

mice who received control AMs (Figure 2J). Collectively, our data demonstrate that the

exacerbation of LPS-induced ALI in β-glucan-trained mice is mediated through AMs, mainly

via functional reprogramming rather than a change in the proportion or number of AMs.

β-glucan reprograms AM

Our findings indicate that β-glucan reprograms AMs causing an increased response to LPS and

subsequently enhanced ALI. To test this hypothesis, we investigated the transcriptional state

of AMs after seven days training with β-glucan *in vivo*, which were then stimulated

with/without LPS *ex vivo* LPS. The rationale for using an *ex vivo* system was to ensure that

AMs are uniformly stimulated with LPS (Figure 3A).

Principal component analysis (PCA) of all expressed genes across the four conditions revealed

that control and β-glucan trained AMs had distinct transcriptional profiles (Figure 3B). More

precisely, the 4 different conditions explained the largest variance as they spread out across the

first principal component (PC), while the variation within replicates was associated to the

second and third PC. Next, we investigated the effect of β-glucan trained AMs by comparing

the differential gene expression between control and β-glucan treated mice (Figure 3C). We

found 88 differentially expressed genes (32 up-regulated and 56 down-regulated, with a false

discovery rate (FDR) adjusted p value < 0.05 and a log2fold-change cut-off of 1). Despite that

only few genes were differentially expressed, gene ontology analysis indicated a strong

enrichment of genes involved in innate immune response and defense response to virus (Figure

3D). Interestingly, LPS response genes were downregulated in β-glucan-treated mice

suggesting that, in response to β-glucan, the LPS response pathway is attenuated or less

activated. Gene set enrichment analysis (GSEA) additionally revealed an increase of genes

involved in IFNα but a significant decrease of genes involved in TNFα signalling (Figure 3E).

Next, we explored how *ex vivo* stimulated AMs responded to LPS using the same differential

gene expression analysis. LPS altered the expression of 525 genes (FDR >0.05 log2FC >1.5)

with the up regulation of 438 genes including numerous pro-inflammatory genes (Figure 3F).

As expected, gene ontology analyses revealed that these genes were involved in cellular

response to LPS, inhibition of viral genome replication, production of interleukin-1β and

cellular response to IL-1 (Figure 3G). Additional analysis of GSEA indicated up-regulation of

genes involved in TNFα signalling (Figure 3H), indicating that LPS and β-glucan both induce

interferon responses but have opposite effects on the TNF response pathway.

To delineate how β-glucan training of AMs may affect the subsequent response to LPS, we

then compared the fold change of gene expression in response to LPS stimulation between β-

glucan trained and untrained AMs. Notably, we observed an upregulation of genes associated

with the LPS response (depicted in blue) in the β-glucan trained group (Figure 3I). This

heightened response included a significant increase in the expression of IL-6, IL-1, TNF, and

other genes (Figure 3I, J). This increased response to LPS by β-glucan-trained AMs was

confirmed at the protein level as they exhibited increased *ex vivo* production of CXCL1 and

TNFα (Figure 3K). Such an increased response to LPS was also obtained in AMs after 28 days

β-glucan training (Supplementary Figure 1G). We also observed similar trends in genes

associated with the defense response to viruses (Supplementary Figure 2G) which is in line

with our findings showing an increased ALI to poly(I:C) (Supplementary Figure 2B-D). Taken

together, β-glucan reprograms AMs to respond robustly to LPS stimulation.

As trained immunity is associated with metabolic rewiring, we next assessed if β-glucan-

induced training modified the AM's metabolic state. Genes involved in both oxidative

phosphorylation and glycolysis were up-regulated during response to LPS in β-glucan-trained

AMs compared to untrained AMs but not at the steady state (Figure 3L). The Seahorse assay

showed while only mitochondrial respiration was increased in β-glucan-trained AMs at the

steady state, both mitochondrial respiration and glycolysis were increased during the response

to LPS in β-glucan-trained AMs, (Figure 3M). Taken together, β-glucan functionally

reprograms AMs by rewiring their transcriptomic and metabolic states.

IFNγ signaling and neutrophils are required for AM reprogramming

Dectin-1 is the receptor that recognizes β-glucan and is involved in mediating the biological

effects of β-glucan-induced training in most cell types[15]. To assess the role of Dectin-1 in β-

glucan increased LPS-induced ALI, we used the same experimental setup using Dectin-1

deficient mice (Supplementary Figure 3A). Surprisingly, the increased neutrophil infiltration

and pro-inflammatory cytokine production in BAL were maintained in β-glucan-treated

Dectin-1[-/-] mice and there was no change in the AM numbers (Supplementary Figure 3B-D).

To determine the effect of Dectin-1 on β-glucan-induced AM function, *Dectin-1*[-/-] mice were

treated with β-glucan and after 7 days AMs were cultured and stimulated *ex vivo* with LPS

(Supplementary Figure 3E). Similar to wild-type AMs, the production of pro-inflammatory

cytokines CXCL1 and TNFα was increased in β-glucan-trained Dectin1-deficient AMs after

LPS stimulation (Supplementary Figure 3F). This indicates that β-glucan-induced AMs

reprogramming is Dectin-1-independent and that AMs are trained via a different signalling

pathway. Since we observed type I IFN gene expression is increased in AMs after β-glucan

treatment (Figure 3E) and type I IFN was previously reported to be involved in AMs training[25],

we next assessed the impact of β-glucan on AMs reprogramming in *IfnaR*[-/-] mice, which lack

type I IFN signalling (Supplementary Figure 3G). Similar to Dectin-1 deficient mice, increased

production of pro-inflammatory cytokines in response to LPS by β-glucan-treated *IfnaR*[-/-] AMs

was maintained, demonstrating that β-glucan-induced AMs reprogramming is type I IFN-

independent (Supplementary Figure 3H).

Considering recent studies suggesting type II IFN can train AMs following BCG [26] or

adenoviral infection [27], using *IfngR*[-/-] mice we next investigated if type II IFN is involved in β-

glucan-induced exacerbation of ALI (Figure 4A). In contrast to *Dectin1*[-/-] *or IfnaR*[-/-] mice, the

increased neutrophils recruitment, inflammatory cytokines production, and ALI were abolished

in β-glucan-trained *IfngR*[-/-] mice (Figure 4B-D). To investigate the role of type II IFN signalling

in β-glucan-induced reprogramming of AMs, we performed *ex vivo* LPS stimulation on β-

glucan trained *IfngR*[-/-] AMs (Figure 4E). The increase of inflammatory cytokines in response

to LPS by β-glucan-trained AMs was abolished (Figure 4F), suggesting that training of AM by

β-glucan is IFNγ-dependent.

Considering the concentration of IFNγ was significantly increased in BAL at day 1 post- β-

glucan administration, which was associated with the increased number of neutrophils (Figure

4G-J), we next assess the direct role of IFNγ in training of AMs by β-glucan. We selectively

depleted IFNγ prior to β-glucan administration *in vivo* (Figure 4K), and found the production

of inflammatory cytokines (CXCL1 and TNFα) was significantly reduced in β-glucan trained

AMs. Similarly, the depletion of neutrophils prior to β-glucan treatment in mice resulted in

significant reduction in inflammatory cytokines in β-glucan trained AMs (Figure 4L). These

findings indicate that β-glucan reprograms AMs via IFNγ- and neutrophils-dependent manner.

DISCUSSION

The evolving field of trained immunity has challenged the boundaries we once drew to discern between the innate and adaptive immune systems. The discovery that the innate immune system can be 'trained' to retain memory-like features has provided novel avenues for prophylactic and therapeutic strategies. The use of an adjuvant like β-glucan has been investigated as an anti-infection agent in acute infections or inflammatory conditions [16, 28] as well as an anti-cancer treatment [29, 30]. While we and others have shown that β-glucan can induce central trained immunity by functionally reprogramming the hematopoietic stem cell compartment in the bone marrow, the present study highlights an additional facet of β-glucan's immunomodulatory effects on residential immune cells. Here we show that a single intraperitoneal (i.p.) injection of β-glucan can induce trained immunity in the lung alveolar macrophages and functionally reprograms AMs at both the transcriptional and metabolic levels. These findings underscore the occurrence of tissue-specific immune training following systemic treatment, demonstrating processes of β-glucan-mediated peripheral trained immunity occurring in parallel with central immunity. We subsequently demonstrate that β-glucan trained AMs can be detrimental after LPS or poly(I:C) administration causing severe ALI. Thus, this study is shedding light on the deleterious effects of trained immunity and its impact on immunopathology, which is incompletely understood.

There are several key features in trained immunity, including shifts in epigenetic and transcriptomic profiles after initial stimulation that returns to steady states. However, this epigenetic and metabolic remodeling result in long-term memory with heightened responses to diverse secondary stimulation [12]. The nature of each stimulus appears to be critical in determining the immunological outcome. While we show here that β-glucan aggravates LPS-induced immunopathology, repeated challenges of LPS has also long-been associated with a dampening of immune responses, termed tolerance [31, 32]. However, ambient exposure of LPS

has also been linked with training leading to enhanced bacterial clearance [25]. Evidently, the

impacts of trained immunity are intricate and context dependent. We postulate two signals are

required for the induction of trained immunity: signal 1, which is essential for reprogramming

innate immune cells centrally (HSCs) or peripherally (e.g. AMs), and signal 2, an

environmental cue critical for activating their functional capacity. In this study, we provide

evidence supporting this conceptual framework.

β-glucan-trained AMs were able to generate a robust response both *in vitro* and *in vivo* to

bacterial (LPS) or viral (poly(I:C)) ligands. The increased response in β-glucan-trained AM

was maintained up to 28 days, which is an indication of long-term reprogramming of AMs.

The reprogramming of trained AMs was supported by significant alterations in both the

transcriptomic and metabolic states. Using adoptive transfer experiments, we then showed that

the functional reprogramming of β-glucan trained AMs persisted even in the absence of the

initial systemic β-glucan administration. This finding highlights the persistence of the intrinsic

training program in AMs.

AMs are originated from yolk sac and fetal liver monocytes, which seeds the lung as soon as

we take our first breath of air [10]. Importantly, AMs are the first immune cell to response to any

particles or pathogens that reach the lower airways of the lungs, thus alterations in their

functional capacity can significantly impact subsequent immune responses. For instance, given

AMs are in direct contact with surfactant (produced by alveolar type 2 epithelial cells) and

invading pathogens, a defect in GM-CSF or TGFβ signalling leads to accumulation of

surfactant (alveolar proteinosis) and increased susceptibility to pulmonary infections[33, 34]. AMs

are able to sense and integrate multiple environmental signals - such as pH[34], temperature[35, 36],

osmolarity[37], metabolites including fatty acids[38, 39], extracellular membrane components and

danger signals [40] – to maintain tissue homeostasis[41]. However, the mechanisms underlying their

ability to adapt to environmental stimuli while maintaining lung tissue homeostasis without

impairing gas exchange is largely unknown[42]. AMs exhibit a remarkable plasticity,

demonstrating a spectrum of functional polarization that ranges from regulatory to pro-

inflammatory and anti-inflammatory states. For instance, AM anti-inflammatory polarization

in a LPS model of sepsis was TNFα dependent, as AMs exposed to TNFα exhibited diminished

phagocytosis, superoxide anion (O2-) and CXCL1 production, with reduced neutrophils

recruitment[32]. Consequently, these mice had a reduction in lung clearance of *P. aeruginosa*

infection. Influenza virus was able to induce similar anti-inflammatory function in AMs with

decreased CXCL1 production and neutrophils recruitment via type I IFN pathway[43], leading to

an increased susceptibility to super bacterial infection. On the other hand, AMs appear to be

more resistant to polarization towards a pro-inflammatory state [44]. A recent study demonstrated

that exposure of lungs to ambient amount of LPS trains AMs in type I IFN dependent, but type

II IFN and T cell independent manner[25]. However, in our model system, systemic

administration of β-glucan trains AMs in a type II IFN dependent, but type I IFN independent

manner. Similarly, in live infection models, we and others have identified IFNγ signaling a key

player in AM training after BCG vaccination[26], influenza infection[45], pneumococcal infection

[46] and intranasal infection with an adenoviral vector [8]. Although we have not identified which

cells produce IFNγ in the β-glucan model, it has been demonstrated by our group and others

that following BCG vaccination, CD4+ T cells are the major source of IFNγ [26], whereas after

pulmonary adenovirus infection, CD8+ T cells predominantly produce IFNγ [27]. Influenza

infection has also been described to induce IFNγ-dependent AM training with NK cells being

the major source [45]. Interestingly, we found that neutrophils were also required for β-glucan-

mediated AMs training. We have been recently shown that β-glucan can reprogram HSCs via

Dectin 1 and type I IFN signaling to promote granulopoiesis and the generation of trained

neutrophils promoting disease tolerance to influenza infection [20]. Although we have shown that

the recruitment of these trained neutrophils and type II IFN signaling were required for training

AMs, the cellular and molecular mechanisms involved in this dialogue is unknown and requires

further investigation. Interestingly, we have recently demonstrated that, in addition to GM-CSF

and TGFβ signalling, neutrophils are critical for AMs self-renewal and maintenance during

early lung development *via* the production of 12-hydroxyeicosatetraenoic acid (12-HETE)[47].

The absence of 12-HETE leads to a significant reduction in the number AMs in adult lungs,

enhanced senescence, and consequently increased susceptibility to IAV or SARS-CoV-2

infection. Thus, there might be a constant bidirectional dialogue between neutrophils and AMs,

with neutrophils providing cues from internal organelles to AMs, and AMs offering signals

from the external environment to neutrophils, which then return to their graveyard in the BM.

Although we have not directly tested the contribution of circulating monocytes in the initial

reprogramming of AMs via β-glucan, the persistence of trained immunity by adoptively

transferred AMs into *Csf2rb*[-/-] mice suggests that the maintenance of the trained AM state was

independent of bone marrow derived monocytes. Additionally, the findings from Theobald *et*

al indicate that Dectin-1 is the receptor responsible for recognizing β-glucan which has been

shown to activate macrophages and induce trained immunity in several models [20, 48, 49, 50]. Here

we found that β-glucan-mediated AM training, when administered systemically, was

independent of Dectin-1. There are two potential explanations for this observation. First, it has

been shown that β-glucan can also activate signaling via other TLRs, particularly TLR2 or

complement receptor 3 (CR3) [51, 52, 53], and second as the β-glucan is a particulate, its

internalization by phagocytes can also initiate signaling [54, 55, 56]. Additionally, our

understanding of how administration of β-glucan in peritoneal cavity leads to HSCs training in

the BM and AMs training in thoracic cavity is extremely limited. Thus, addressing the

molecular mechanisms of β-glucan signaling pathways (e.g., Dectin1-dependent and

independent) in both immune and non-immune cells, as well as its mode of action (e.g., direct

access to an organ versus indirect effects via systemic release of cytokines), is necessary to

delineate the deleterious versus protective effects of β-glucan-mediated trained immunity.

β-glucan is present in the cell wall of all fungi but will vary between different species and

strains. In fact, the differential 1,3 to 1,6 glycosidic branching and molecular weight

significantly impacts the response to the compound with a large variability of scientific findings

contingent on the type of β-glucan used in a study[57, 58, 59]. Fungi make up a portion of the human

microbiome, termed the mycobiome. Several studies have described a high mycotic diversity

between different populations, and even significant variability within an individual overtime

[60, 61, 62]. The variance in gut colonization by fungal species can moreover cause gut dysbiosis

which has been associated with poorer outcomes during SARS-CoV-2 infection, sepsis and

cancer immunotherapy[63, 64 65, 66, 67]. It is tempting to postulate that the levels of the mycobiome

as well as its composition can influence subsequent immune reactions partly due to a distinct

β-glucan makeup. In fact, antibodies to various types of β-glucan was detected in adult sera

with different levels correlating with a person's occupation [68]. Thus, the quantity and quality

of circulating β-glucan in an individual at steady state can remarkably affect the subsequent

immune responses to sterile or microbial inflammation. The concept of detrimental training by

endogenous agents has also been described with heme, which similarly aggravated LPS-

induced inflammation [69]. Basal levels of such immunomodulatory molecules provide a basis

for host response heterogeneity and corresponding excessive inflammation [70]. Understanding

these underlying processes may provide important insights for developing novel therapeutic

approaches against immune-mediated pulmonary disease.

MATERIALS AND METHODS

Mice

C57BL/6, *Csf2rb*$^{-/-}$, *Dectin1*$^{-/-}$, *IfnaR*$^{-/-}$ and *IfngR*$^{-/-}$ mice were purchased from Jackson Laboratories. All animals were housed and inbred at the animal facility of the Research Institute of McGill University under specific pathogen-free conditions with *ad libitum* access to food and water, a temperature of 21 °C (±1 °C), relative humidity of 40–60% (±5%) and light cycle of 12 h on, 12 h off (daily cycle). *Sex as a biological variable*: Mice were randomly allocated to experimental groups, and experiments were performed using both female and male age- and sex-matched mice. Similar findings were reported for both sexes.

β-glucan training

Mice were administered intraperitoneally with 1mg of β-1,3-glucan purified from *Saccharomyces cerevisiae* (Sigma) diluted in 100μL of PBS seven days or twenty-eight days before lung injury or AMs collection for *ex vivo* stimulation.

Acute lung injury models

Mice were administered with 50μg of *Escherichia coli* O55:B55 LPS (Sigma) or poly(I:C) (Invivogen) in PBS (25 μL per mouse, intranasally) to induce TLR-4 or 3-mediated acute lung injury (ALI), respectively.

Lung microCT scan

The trachea was cannulated with a 22-gauge cannula and an intra-thoracic pressure of 25mmH$_2$O was generated using a manometer. Images were acquired right after lung inflation using the nanoScan SPECT + CT (Mediso®) allowing a resolution of 20μm. DICOM software

was used to analyse the microCT scans determining the average lung Hounsfield unit (HU) and

the proportion of non- or poorly-aerated lung (HU -500; +100).

BAL and lung collection

Broncho-alveolar lavage (BAL) samples collected by cannulating the trachea with a 22-gauge

cannula, then washing the lungs with 3× 1 mL of cold, sterile PBS. The total volume of the

recovered fluid after lavage was around 0.7 ml. Samples were centrifuged (1,500 r.p.m.,

10 min). Lung tissues were perfused with 10 mL of PBS, collected and minced before

collagenase type I (3mg, Worthington CLS-1), elastase (3mg, Worthington ESL) and DNase I

(0.4mg, Worthington D) digestion for 30min at 37 °C. Lungs were filtered through a 70 µm

nylon mesh, and red blood cells were lysed.

Endothelial permeability

LPS or poly(I:C)-challenged mice were intravenously injected with 400 µl of Evan's blue dye

(2% in PBS) into the mice. After 1 h, mice were euthanized, and lungs were perfused with

10 ml of PBS. Evan's blue then extracted by overnight incubation in formamide at 56 °C

(lungs) and quantified by spectrophotometry analysis using a standard curve of Evan's blue in

formamide.

Histopathological analysis

Histopathological analysis was performed as previously described[47]. Lungs were inflated and

fixed for 48 h with 10% formalin, and then embedded in paraffin. Sections (5 µm) were cut and

stained with haematoxylin and eosin. Slides were scanned at a resolution of ×40 magnification,

and pictures were taken using a Leica Aperio slide scanner (Leica). Histology samples were

evaluated according to ATS 2011 guidelines regarding "Features and measurements of

experimental acute lung injury in animals" by a blinded observer.

ELISA

CXCL1, TNF-α, IL-6 and IFNγ levels in BAL were assessed by ELISA (R&D Systems).

Protein in BAL

Samples were centrifuged (1,500 r.p.m., 10 min), and total protein content was assessed using

a Pierce BCA Protein assay (ThermoFisher).

Flow cytometry

BAL and total lung cell counts were determined with a haemocytometer, and 1–2 million cells

were used for staining. Cells were initially stained with viability dye e506 (Invitrogen, 20 min,

4 °C) and surface stained with anti-CD16/32 (BD Bioscience) in 0.5% BSA/PBS solution to

block nonspecific AB interaction with Fc receptors (10 min, 4 °C). Cells were then surface-

stained with different combinations of PE-Cy7-conjugated anti-CD11c, BUV786-conjugated

anti-Siglec-F, BUV395-conjugated anti-CD11b, APC-Cy7-conjugated anti-Ly6G, BUV737-

conjugated anti-CD45.2 antibodies (all from BD Biosciences). For IFNγ intra-cellular staining,

cells were fixed and permeabilized using BD CytoFix/CytoPerm (BD Bioscience) before

intracellular staining with PE-conjugated anti-IFNγ antibodies (BD Biosciences). Flow

cytometry was performed using a BD LSR Fortessa X-20 instrument (BD Biosciences) with

FACSDiva software v.8.0.1 (BD Biosciences). Analysis was performed using FlowJo software

v.10.7.1 (Tree Star).

Intravascular staining

In vivo discrimination between pulmonary vasculature and parenchyma was performed as previously described[47]. Adult WT mice were given 2 µg of FITC-conjugated anti-CD45.2 intravenously to label all circulating cells. Three minutes later, mice were euthanized and lungs collected, stained *ex vivo* with BUV395-conjugated anti-CD45.2 antibody to determine the parenchymal (cells only labelled with the *ex vivo* antibody) or vascular localization of the cells (cells labelled with both antibodies).

AM depletion

WT mice were treated with control or clodronate liposomes (70 µl, intranasally; Liposoma BV). The LPS-induced ALI was then performed at day 2 after clodronate instillation.

Adoptive transfer models

AMs from WT mice which received i.p. PBS or β-glucan were collected as described above and resuspended at a density of 5×10^4 cells per 5-7 µL of RPMI1640 medium supplemented with 10% (v/v) FBS, 2 mM l-glutamine, 10 mM HEPES and 100 U ml–1 penicillin–streptomycin. AMs were then transferred by the intranasal route into Day 2 $Csf2rb^{-/-}$ pups. LPS-induced ALI was performed 6 weeks after AM adoptive transfer. BAL and lung tissue were collected and processed as described above for endothelial permeability, flow cytometry, total BAL cytokine and protein content experiments.

Isolation and culture of alveolar macrophages

AMs were collected by BAL of naive mice using cold, sterile PBS (5×1 mL for adult mice). AMs were cultured in the specific media described above. After 1 h of adhesion, AMs were washed with PBS and placed in fresh medium.

Ex vivo stimulation

AMs from WT, *Dectin1[-/-]*, *IfnaR[-/-]* and *IfngR[-/-]* mice which received i.p. PBS or β-glucan were collected as described above and 5×10^4 cells in specific media were distributed per well. AMs were stimulated with 50 ng/ mL of LPS (Sigma) for 4 hours at 37°C.

Library preparation and RNA-seq

Total RNA was collected from BAL AMs from four WT mice per conditions (i.p PBS, no *ex vivo* stimulation / i.p. β-glucan, no *ex vivo* stimulation / i.p PBS, *ex vivo* LPS stimulation / i.p. β-glucan, *ex vivo* LPS stimulation). After RNA quality controls, sequencing libraries were constructed using the Illumina TruSeq protocol. Libraries were sequenced on an Illumina NovaSeq 6000 (paired-end 100 base pair) to an average depth of 51,189,336 reads per sample.

RNA-seq data analysis

RNA-seq reads were aligned to the Mus musculus genome from Ensembl version 99 using STAR (version 2.7.3a) was used[71]. All regions overlapped between referenced exons and alignments were counted using featureCounts (subread-1.6.4)[72]. Low abundance genes were filtered out leaving 12,894 genes for subsequent analysis.

Differential expression analyses were performed using the DESeq2 package (DESeq2 1.40.2)[73]. Gene ontology analysis was realized with the R package TopGO. For gene set enrichment analysis (GSEA) a ranked list of the differentially expressed genes was used with clusterProfiler v4.8.2[74], and the Molecular Signatures Database MSigDB v7.5.1.

Extracellular flux analysis

Seahorse assay of isolated cells was performed as previously described[47]. Real-time OCRs of AMs were measured in XF medium (non-buffered RPMI containing 2 mM l-glutamine, 25 mM

glucose and 1 mM sodium pyruvate) using a Seahorse Xfe 96 Analyzer (Agilent Technologies).

For the mitochondrial stress test, mitochondrial inhibitors oligomycin (1.5 µM), fluorocarbonyl

cyanide phenylhydrazone (FCCP) (1 µM), antimycin A and rotenone (0.5 µM) were used as

per the manufacturer's recommendations. In brief, cells were seeded at a density of 100,000

cells per well and 3 basal measurements were taken. Following this, two consecutive

measurements were taken following each injection of oligomycin, FCCP and antimycin A with

rotenone. All measurements were normalized to cell number using crystal violet dye extraction

assay. Oxygen consumption curves, OCRs and ECARs were generated using Wave Desktop

2.3 (Agilent Technologies).

IFNγ and neutrophils depletion experiments.

Sex- and age-matched adult WT mice received intra-peritoneal 200µg of anti-IFNγ (rat IgG1k,

Biolegend), anti-Ly6G (rat IgG2a,k, Biolegend) or appropriated control isotypes injection at

Day-1, Day 0, Day 2, Day 4 and Day 6 per intraperitoneal injection of β-glucan. AMs collection

for *ex vivo* stimulation was performed at Day 7 after intraperitoneal β-glucan injection.

Ethics statement

All experiments involving animals were approved by the McGill University Animal Care

Committee (permit number 2010–5860) in accordance with the guidelines set out by the

Canadian Council on Animal Care.

Statistical analysis

Data are presented as the median with interquartile range. Statistical analyses were performed

using GraphPad Prism v.9.1.2 software (GraphPad). Statistical differences were determined

using one-way analysis of variance (ANOVA) followed by Dunn's multiple comparisons test,

paired or unpaired two-tailed t-test or two-tailed Mann–Whitney test. Differential gene

expression analysis was carried out using DEseq2 package[73].

Data availability

All data supporting the findings of this study are included in the published article and

supplementary materials. Bulk RNA-seq data have been deposited to the European Nucleotide

Archive and will be made publicly available upon publication. Source data are provided with

this paper.

Acknowledgements

The authors acknowledge technical help from staff at the RI-MUHC histopathology platform

and RI-MUHC Small Animal Imaging Laboratory. M.D. is funded by Canadian Institute of

Health Research (CIHR) Project Grant-168885 and MM1174910, a Fonds de recherche du

Québec–Santé (FRQS) Award, holds the Strauss Chair in Respiratory Diseases and is a fellow

member of the Royal Society of Canada. E.P. and R.P. are fellows supported by a Postdoctoral

Fellowship from the Fonds de Recherche du Québec Santé.

Author contributions

Conceptualization: R.P., E.P. and M.D. Methodology: R.P., E.P., A.S.K., M.S., K.A.T., R.S.,

E.L., L.J., J.P. and M.D. Investigation: R.P., K.A.S., E.P., J.P. and M.D. Funding acquisition:

M.D. Project administration: R.P. and M.D. Supervision: E.P., J.P. and M.D. Writing original

draft: R.P. K.A.T and M.D. Writing, review, and editing: K.A.T, R.P., E.P., K.A.S., J.P. and

M.D.

Competing interests

The authors declare no competing interests.

REFERENCES

1. Xiao, W. *et al.* A genomic storm in critically injured humans. *J Exp Med* **208**, 2581-2590 (2011).

2. Gustine, J.N. & Jones, D. Immunopathology of Hyperinflammation in COVID-19. *Am J Pathol* **191**, 4-17 (2021).

3. Opal, S.M. Endotoxins and other sepsis triggers. *Contrib Nephrol* **167**, 14-24 (2010).

4. Xu, Y. *et al.* Mechanisms of Sepsis-Induced Acute Lung Injury and Advancements of Natural Small Molecules in Its Treatment. *Pharmaceuticals (Basel)* **17** (2024).

5. Gill, S.E., Rohan, M. & Mehta, S. Role of pulmonary microvascular endothelial cell apoptosis in murine sepsis-induced lung injury in vivo. *Respir Res* **16**, 109 (2015).

6. Sun, B. *et al.* Acute lung injury caused by sepsis: how does it happen? *Front Med (Lausanne)* **10**, 1289194 (2023).

7. Kumar, V. Pulmonary Innate Immune Response Determines the Outcome of Inflammation During Pneumonia and Sepsis-Associated Acute Lung Injury. *Front Immunol* **11**, 1722 (2020).

8. Zhou, X. & Liao, Y. Gut-Lung Crosstalk in Sepsis-Induced Acute Lung Injury. *Front Microbiol* **12**, 779620 (2021).

9. Schneider, C. *et al.* Induction of the nuclear receptor PPAR-γ by the cytokine GM-CSF is critical for the differentiation of fetal monocytes into alveolar macrophages. *Nat Immunol* **15**, 1026-1037 (2014).

10. Guilliams, M. *et al.* Alveolar macrophages develop from fetal monocytes that differentiate into long-lived cells in the first week of life via GM-CSF. *J Exp Med* **210**, 1977-1992 (2013).

11. Hussell, T. & Bell, T.J. Alveolar macrophages: plasticity in a tissue-specific context. *Nat Rev Immunol* **14**, 81-93 (2014).

12. Divangahi, M. *et al.* Trained immunity, tolerance, priming and differentiation: distinct immunological processes. *Nat Immunol* **22**, 2-6 (2021).

13. Netea, M.G. *et al.* Defining trained immunity and its role in health and disease. *Nat Rev Immunol* **20**, 375-388 (2020).

14. Netea, M.G. *et al.* Trained immunity: A program of innate immune memory in health and disease. *Science* **352**, aaf1098 (2016).

15. Camilli, G., Tabouret, G. & Quintin, J. The Complexity of Fungal β-Glucan in Health and Disease:
Effects on the Mononuclear Phagocyte System. *Front Immunol* **9**, 673 (2018).

16. Moorlag, S. *et al.* β-Glucan Induces Protective Trained Immunity against Mycobacterium
tuberculosis Infection: A Key Role for IL-1. *Cell Rep* **31**, 107634 (2020).

17. Kaufmann, E. *et al.* BCG Educates Hematopoietic Stem Cells to Generate Protective Innate
Immunity against Tuberculosis. *Cell* **172**, 176-190.e119 (2018).

18. Roquilly, A. *et al.* Alveolar macrophages are epigenetically altered after inflammation, leading
to long-term lung immunoparalysis. *Nat Immunol* **21**, 636-648 (2020).

19. Broquet, A. *et al.* Sepsis-trained macrophages promote antitumoral tissue-resident T cells. *Nat
Immunol* **25**, 802-819 (2024).

20. Khan, N. *et al.* β-Glucan reprograms neutrophils to promote disease tolerance against
influenza A virus. *Nat Immunol* (2025).

21. Foster, S.L., Hargreaves, D.C. & Medzhitov, R. Gene-specific control of inflammation by TLR-
induced chromatin modifications. *Nature* **447**, 972-978 (2007).

22. Wengner, A.M., Pitchford, S.C., Furze, R.C. & Rankin, S.M. The coordinated action of G-CSF
and ELR + CXC chemokines in neutrophil mobilization during acute inflammation. *Blood* **111**,
42-49 (2008).

23. Burastero, S.E. *et al.* Increased expression of the CD80 accessory molecule by alveolar
macrophages in asthmatic subjects and its functional involvement in allergen presentation to
autologous TH2 lymphocytes. *J Allergy Clin Immunol* **103**, 1136-1142 (1999).

24. Robb, L. *et al.* Hematopoietic and lung abnormalities in mice with a null mutation of the
common beta subunit of the receptors for granulocyte-macrophage colony-stimulating factor
and interleukins 3 and 5. *Proc Natl Acad Sci U S A* **92**, 9565-9569 (1995).

25. Zahalka, S. *et al.* Trained immunity of alveolar macrophages requires metabolic rewiring and
type 1 interferon signaling. *Mucosal Immunol* **15**, 896-907 (2022).

26. Tran, K.A. *et al.* BCG immunization induces CX3CR1(hi) effector memory T cells to provide
cross-protection via IFN-γ-mediated trained immunity. *Nat Immunol* **25**, 418-431 (2024).

27. Yao, Y. *et al.* Induction of Autonomous Memory Alveolar Macrophages Requires T Cell Help
and Is Critical to Trained Immunity. *Cell* **175**, 1634-1650.e1617 (2018).

28.   Zervopoulou, E. *et al.* Enhanced medullary and extramedullary granulopoiesis sustain the
652   inflammatory response in lupus nephritis. *Lupus Sci Med* **11** (2024).

653
654   29.   Vuscan, P. *et al.* Potent induction of trained immunity by Saccharomyces cerevisiae β-glucans.
655   *Front Immunol* **15**, 1323333 (2024).

656
657   30.   Kalafati, L. *et al.* Innate Immune Training of Granulopoiesis Promotes Anti-tumor Activity. *Cell*
658   **183**, 771-785.e712 (2020).

659
660   31.   Freudenberg, M.A. & Galanos, C. Induction of tolerance to lipopolysaccharide (LPS)-D-
661   galactosamine lethality by pretreatment with LPS is mediated by macrophages. *Infect Immun*
662   **56**, 1352-1357 (1988).

663
664   32.   Mason, C.M., Dobard, E., Summer, W.R. & Nelson, S. Intraportal lipopolysaccharide
665   suppresses pulmonary antibacterial defense mechanisms. *J Infect Dis* **176**, 1293-1302 (1997).

666
667   33.   Baker, A.D. *et al.* Targeted PPAR{gamma} deficiency in alveolar macrophages disrupts
668   surfactant catabolism. *J Lipid Res* **51**, 1325-1331 (2010).

669
670   34.   Huffman, J.A., Hull, W.M., Dranoff, G., Mulligan, R.C. & Whitsett, J.A. Pulmonary epithelial cell
671   expression of GM-CSF corrects the alveolar proteinosis in GM-CSF-deficient mice. *J Clin Invest*
672   **97**, 649-655 (1996).

673
674   35.   Kashio, M. *et al.* Redox signal-mediated sensitization of transient receptor potential
675   melastatin 2 (TRPM2) to temperature affects macrophage functions. *Proc Natl Acad Sci U S A*
676   **109**, 6745-6750 (2012).

677
678   36.   Link, T.M. *et al.* TRPV2 has a pivotal role in macrophage particle binding and phagocytosis. *Nat*
679   *Immunol* **11**, 232-239 (2010).

680
681   37.   Fang, H.Y. *et al.* Hypoxia-inducible factors 1 and 2 are important transcriptional effectors in
682   primary macrophages experiencing hypoxia. *Blood* **114**, 844-859 (2009).

683
684   38.   Haldar, K. *et al.* The sputum microbiome is distinct between COPD and health, independent
685   of smoking history. *Respir Res* **21**, 183 (2020).

686
687   39.   Okabe, Y. & Medzhitov, R. Tissue-specific signals control reversible program of localization and
688   functional polarization of macrophages. *Cell* **157**, 832-844 (2014).

689
690   40.   Sun, L., Wu, J., Du, F., Chen, X. & Chen, Z.J. Cyclic GMP-AMP synthase is a cytosolic DNA sensor
691   that activates the type I interferon pathway. *Science* **339**, 786-791 (2013).

692
693   41.   Okabe, Y. & Medzhitov, R. Tissue biology perspective on macrophages. *Nat Immunol* **17**, 9-17
694   (2016).

42. Lazarov, T., Juarez-Carreño, S., Cox, N. & Geissmann, F. Physiology and diseases of tissue-
resident macrophages. *Nature* **618**, 698-707 (2023).

43. Shirey, K.A. *et al.* Influenza "Trains" the Host for Enhanced Susceptibility to Secondary
Bacterial Infection. *mBio* **10** (2019).

44. Tomlinson, G.S. *et al.* Adherent human alveolar macrophages exhibit a transient pro-
inflammatory profile that confounds responses to innate immune stimulation. *PLoS One* **7**,
e40348 (2012).

45. Wang, T. *et al.* Influenza-trained mucosal-resident alveolar macrophages confer long-term
antitumor immunity in the lungs. *Nat Immunol* **24**, 423-438 (2023).

46. Arafa, E.I. *et al.* Recruitment and training of alveolar macrophages after pneumococcal
pneumonia. *JCI Insight* **7** (2022).

47. Pernet, E. *et al.* Neonatal imprinting of alveolar macrophages via neutrophil-derived 12-HETE.
Nature **614**, 530-538 (2023).

48. Cheng, S.C. *et al.* mTOR- and HIF-1α-mediated aerobic glycolysis as metabolic basis for trained
immunity. *Science* **345**, 1250684 (2014).

49. Arts, R.J. *et al.* Glutaminolysis and Fumarate Accumulation Integrate Immunometabolic and
Epigenetic Programs in Trained Immunity. *Cell Metab* **24**, 807-819 (2016).

50. Quintin, J. *et al.* Candida albicans infection affords protection against reinfection via functional
reprogramming of monocytes. *Cell Host Microbe* **12**, 223-232 (2012).

51. Yadav, M. & Schorey, J.S. The beta-glucan receptor dectin-1 functions together with TLR2 to
mediate macrophage activation by mycobacteria. *Blood* **108**, 3168-3175 (2006).

52. Thornton, B.P., Větvicka, V., Pitman, M., Goldman, R.C. & Ross, G.D. Analysis of the sugar
specificity and molecular location of the beta-glucan-binding lectin site of complement
receptor type 3 (CD11b/CD18). *J Immunol* **156**, 1235-1246 (1996).

53. Xia, Y. *et al.* The beta-glucan-binding lectin site of mouse CR3 (CD11b/CD18) and its function
in generating a primed state of the receptor that mediates cytotoxic activation in response to
iC3b-opsonized target cells. *J Immunol* **162**, 2281-2290 (1999).

54. McLeish, K.R., Klein, J.B., Coxon, P.Y., Head, K.Z. & Ward, R.A. Bacterial phagocytosis activates
extracellular signal-regulated kinase and p38 mitogen-activated protein kinase cascades in
human neutrophils. *J Leukoc Biol* **64**, 835-844 (1998).

55. Yang, H., He, H. & Dong, Y. CARD9 Syk-dependent and Raf-1 Syk-independent signaling
pathways in target recognition of Candida albicans by Dectin-1. *Eur J Clin Microbiol Infect Dis*
30, 303-305 (2011).

56. Qi, C. *et al.* Differential pathways regulating innate and adaptive antitumor immune responses
by particulate and soluble yeast-derived β-glucans. *Blood* **117**, 6825-6836 (2011).

57. Luo, J. *et al.* Purified β-glucans of Different Molecular Weights Enhance Growth Performance
of LPS-challenged Piglets via Improved Gut Barrier Function and Microbiota. *Animals (Basel)*
9 (2019).

58. Moerings, B.G.J. *et al.* Continuous Exposure to Non-Soluble β-Glucans Induces Trained
Immunity in M-CSF-Differentiated Macrophages. *Front Immunol* **12**, 672796 (2021).

59. Akramiene, D., Kondrotas, A., Didziapetriene, J. & Kevelaitis, E. Effects of beta-glucans on the
immune system. *Medicina (Kaunas)* **43**, 597-606 (2007).

60. Sun, Y. *et al.* Population-Level Configurations of Gut Mycobiome Across 6 Ethnicities in Urban
and Rural China. *Gastroenterology* **160**, 272-286.e211 (2021).

61. Szóstak, N. *et al.* Host Factors Associated with Gut Mycobiome Structure. *mSystems* **8**,
e0098622 (2023).

62. Nash, A.K. *et al.* The gut mycobiome of the Human Microbiome Project healthy cohort.
Microbiome **5**, 153 (2017).

63. Reinold, J. *et al.* The Fungal Gut Microbiome Exhibits Reduced Diversity and Increased Relative
Abundance of Ascomycota in Severe COVID-19 Illness and Distinct Interconnected
Communities in SARS-CoV-2 Positive Patients. *Front Cell Infect Microbiol* **12**, 848650 (2022).

64. Chen, Y. *et al.* Association Between Gut Dysbiosis and Sepsis-Induced Myocardial Dysfunction
in Patients With Sepsis or Septic Shock. *Front Cell Infect Microbiol* **12**, 857035 (2022).

65. Huang, X. *et al.* Multi-kingdom gut microbiota analyses define bacterial-fungal interplay and
microbial markers of pan-cancer immunotherapy across cohorts. *Cell Host Microbe* **31**, 1930-
1943.e1934 (2023).

66. Orieux, A. *et al.* The gut microbiota composition is linked to subsequent occurrence of
ventilator-associated pneumonia in critically ill patients. *Microbiol Spectr* **11**, e0064123
(2023).

67. Prevel, R. *et al.* Gut bacteriobiota and mycobiota are both associated with Day-28 mortality
among critically ill patients. *Crit Care* **26**, 105 (2022).

68. Noss, I. *et al.* IgG to various beta-glucans in a human adult population. *Int Arch Allergy*
Immunol **157**, 98-108 (2012).

785
786 69. Jentho, E. *et al.* Trained innate immunity, long-lasting epigenetic modulation, and skewed
787   myelopoiesis by heme. *Proc Natl Acad Sci U S A* **118** (2021).

788
789 70. Grasselli, G. *et al.* Pathophysiology of COVID-19-associated acute respiratory distress
790   syndrome: a multicentre prospective observational study. *Lancet Respir Med* **8**, 1201-1208
791   (2020).

792
793 71. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

794
795 72. Liao, Y., Smyth, G.K. & Shi, W. featureCounts: an efficient general purpose program for
796   assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).

797
798 73. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
799   RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

800
801 74. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
802   *Innovation (Camb)* **2**, 100141 (2021).

FIGURE LEGENDS

Figure 1. β-glucan-mediated trained immunity increases LPS-induced ALI. A, Schematic of the β-glucan-induced training seven days before lipopolysaccharide (LPS)-induced acute lung injury (ALI) model. Experiments were performed in sex- and age-matched 10-12 weeks old control (i.p. PBS, white bars) and trained (i. p. β-glucan, green bars) WT mice. **B**, Lung micro-CT scan, percentage of poorly- or non-aerated lung and average lung Hounsfield unit. **C**, Alveolar capillary membrane permeability assessed by lung Evans blue dye concentration. **D**, Lung histology after staining with haematoxylin and eosin. **E**, Quantification of BAL neutrophils frequency (left) and absolute count (right) (gated on single live cells, $CD45.2^+$, $CD11c^-$, Siglec-F^-, $CD11b^+$, $Ly6G^+$). **F, G, H** BAL chemokine and pro-inflammatory cytokines concentrations (left to right) (CXCL1: chemokine C-X-C motif ligand 1, IL-6: interleukin-6 and TNF-α: tumor necrosis factor α**). I,** Quantification of BAL alveolar macrophages frequency (left) and absolute count (right) (gated on single live cells, $CD45.2^+$, $CD11c^+$, Siglec-F^+). Data were analysed using one-way ANOVA followed by Dunn's multiple comparisons test. * $p<0.05$, *** $p<0.001$.

Figure 2. Systemic administration of β-glucan enhances ALI via AMs. A, Schematic of the clodronate-mediated alveolar macrophages (AM) depletion experiments, performed in sex- and age-matched 10-12 weeks old control (i.p. PBS, white bars) and trained (i. p. β-glucan, green bars) WT mice. **B**, Alveolar capillary membrane permeability assessed by lung Evans blue dye concentration. **C**, Quantification of BAL neutrophils frequency (left) and absolute count (right) (gated on single live cells, $CD45.2^+$, $CD11c^-$, Siglec-F^-, $CD11b^+$, $Ly6G^+$). **D**, BAL chemokine C-X-C motif ligand 1 (CXCL1) and pro-inflammatory cytokines (IL-6: interleukin-6 and TNF-α: tumor necrosis factor α) concentrations. **E**, Schematic of the β-glucan-induced training and

lipopolysaccharide (LPS)-induced acute lung injury (ALI) model in sex- and age-matched 6 weeks old control (i.p. PBS, white bars) and trained (i. p. β-glucan, green bars) *Csf2rb*[-/-] mice. **F**, BAL total protein concentration. **G**, Quantification of BAL neutrophils frequency (left) and absolute count (right) (gated on single live cells, CD45.2[+], CD11c[-], Siglec-F[-], CD11b[+], Ly6G[+]). **H**, BAL CXCL1, IL-6 and TNF-α concentrations. **I**, Schematic of the adoptive transfer of control (i.p. PBS, white bars) or β-glucan-trained (i. p. β-glucan, green bars) AMs collected from adult WT mice to 2 days old *Csf2rb*[-/-] mice. Lipopolysaccharide (LPS)-induced acute lung injury (ALI) was performed 6 weeks after adoptive transfer. **J**, BAL total protein concentration. **K**, Quantification of BAL neutrophils frequency (left) and absolute count (right) (gated on single live cells, CD45.2[+], CD11c, Siglec-F[-], CD11b[+], Ly6G[+]). **L**, BAL CXCL1, IL-6 and TNF-α concentrations. Data were analysed using one-way ANOVA followed by Dunn's multiple comparisons test. * $p<0.05$, ** $p<0.01$, *** $p<0.001$.

Figure 3. β-glucan reprograms AMs. A, Schematic of control (i.p. PBS) or β-glucan-trained (i. p. β-glucan) AMs collected from adult WT mice *ex vivo* stimulation with LPS (LPS-: unstimulated, LPS+: stimulated in RNAseq analysis). **B**, Discovery plot. **C**, AM differential expression of genes in response to β-glucan training. **D**, Gene ontology in response to β-glucan training. **E**, GSEA in response to β-glucan training. **F**, AM differential expression of genes in response to LPS stimulation. **G**, Gene ontology in response to LPS stimulation. **H**, GSEA in response to LPS stimulation. **I**, AM gene expression in response to LPS in β-glucan-trained AMs. **J**, Examples of genes expression in response to LPS in control *vs* β-glucan-trained AMs. **K**, Chemokine C-X-C motif ligand 1 (CXCL1) and tumor necrosis factor α (TNF-α) concentrations after *ex vivo* LPS stimulation. **L**, GSEA of oxidative phosphorylation (left) and glycolysis (right) pathways according to β-glucan-training in unstimulated (LPS-) and LPS stimulated (LPS+) AMs. **M**, Evaluation of AM metabolism: basal respiration (upper left), ATP

production (upper right), extracellular acidification rate (ECAR, lower left), oxygen

consumption rate (OCR, lower right). Data were analysed using one-way ANOVA followed

by Dunn's multiple comparisons test. ** $p<0.01$, *** $p<0.001$.

Figure 4. IFNγ and neutrophils are required in β-glucan-mediated AM reprogramming.

A, Schematic of the β-glucan-induced training and lipopolysaccharide (LPS)-induced acute

lung injury (ALI) model. Experiments were performed in sex- and age-matched 10-12 weeks

old control (i.p. PBS, white bars) and trained (i. p. β-glucan, green bars) *IfngR$^{-/-}$* mice. **B**,

Alveolar capillary membrane permeability assessed by lung Evans blue dye concentration. **C**,

Quantification of BAL neutrophils proportion (left) and absolute count (right) (gated on single

live cells, CD45.2$^+$, CD11c$^-$, Siglec-F$^-$, CD11b$^+$, Ly6G$^+$). **D**, BAL chemokine C-X-C motif

ligand 1 (CXCL1) concentration (left) and pro-inflammatory cytokines (IL-6: interleukin-6 –

middle- and TNF-α: tumor necrosis factor α –right) concentrations. **E**, Schematic of control

(i.p. PBS, white bars) or β-glucan-trained (i. p. β-glucan, green bars) AMs collected from adult

IfngR$^{-/-}$ mice *ex vivo* stimulation with LPS. **F**, Chemokine C-X-C motif ligand 1 (CXCL1) and

tumor necrosis factor α (TNF-α) concentrations after *ex vivo* LPS stimulation. **G**, Schematic of

the analysis of the effect of i.p. β-glucan injection on interferon-γ (IFNγ) production and

neutrophils expansion before (white bars) and Day 1, Day 3, Day 5 and Day 7 post-injection

(green bars). **H**, BAL IFNγ concentrations. **I**, Quantification of BAL neutrophils proportion

(left) and absolute count (right) (gated on single live cells, CD45.2$^+$, CD11c$^-$, Siglec-F$^-$,

CD11b$^+$, Ly6G$^+$). **J**, Quantification of lung neutrophils proportion (left) and absolute count

(right). **K**, Schematic of control (i.p. PBS, white bars) or β-glucan-trained (i. p. β-glucan, green

bars) AMs *ex vivo* stimulation with 50ng/mL LPS. AMs were collected from control (i.p.

injection of isotypes), neutrophils depleted (i.p. injection of anti-Ly6G antibodies) or IFNγ

antibody-depleted (i.p. injection of anti-IFNγ antibodies) adult WT mice. **L**, CXCL1 and TNF-

α concentrations after *ex vivo* LPS stimulation. Data were analysed using one-way ANOVA

followed by Dunn's multiple comparisons test. * p<0.05, *** p<0.001.

Supplementary Figure 1. Long-term effects of β-glucan-mediated trained immunity on

LPS-induced ALI. A, Schematic of the β-glucan-induced training seven days before

lipopolysaccharide (LPS)-induced acute lung injury (ALI) model for weight loss monitoring.

B, Weight loss curve post-LPS instillation (n=10). **C**, CD11b expression on CD11c$^+$SiglecF$^+$

cells in the bronchoalveolar lavage (BAL) (n=4). **D**, Expression of AM-associated marker (left

to right: MHCII, F4/80, CD169, CD64, CD80) on CD11c$^+$SiglecF$^+$ cells in the bronchoalveolar

lavage (BAL) (n=5). **E**, Schematic of the β-glucan-induced training twenty-eight days before

lipopolysaccharide (LPS)-induced acute lung injury (ALI) model. Experiments were

performed in sex- and age-matched 10-12 weeks old control (i.p. PBS, white bars) and trained

(i. p. β-glucan, green bars) WT mice. **F**, Alveolar capillary membrane permeability assessed

by lung Evans blue dye concentration. **G**, Lung histology after staining with haematoxylin and

eosin. **H**, BAL chemokine C-X-C motif ligand 1 (CXCL1) concentration (left) and pro-

inflammatory cytokines (IL-6: interleukin-6 (middle) and TNF-α: tumor necrosis factor α

(right)). **I**, Quantification of BAL neutrophils frequency (left) and absolute count (right) (gated

on single live cells, CD45.2$^+$, CD11c$^-$, Siglec-F$^-$, CD11b$^+$, Ly6G$^+$). **J,** quantification of BAL

alveolar macrophages frequency (left) and absolute count (right) (gated on single live cells,

CD45.2$^+$, CD11c$^+$, Siglec-F$^+$). **K**, Schematic of control (i.p. PBS, white bars) or 28 days β-

glucan-trained (i. p. β-glucan, green bars) AMs collected from adult WT mice *ex vivo*

stimulation with LPS. **L**, Chemokine C-X-C motif ligand 1 (CXCL1) and tumor necrosis factor

α (TNF-α) concentrations after *ex vivo* LPS stimulation. Data were analysed using unpaired t-

test and one-way ANOVA followed by Dunn's multiple comparisons test. * p<0.05, **

p<0.01, *** p<0.001.

Supplementary Figure 2. β-glucan-mediated trained immunity increases poly(I:C)-induced ALI. A, Schematic of the β-glucan-induced training seven days before poly(I:C)-induced acute lung injury (ALI) model. Experiments were performed in sex- and age-matched 10-12 weeks old control (i.p. PBS, white bars) and trained (i. p. β-glucan, green bars) WT mice. **B**, Alveolar capillary membrane permeability assessed by lung Evans blue dye concentration. **C**, Lung histology after staining with haematoxylin and eosin. **D**, BAL chemokine C-X-C motif ligand 1 (CXCL1) concentration (left) and pro-inflammatory cytokines (IL-6: interleukin-6 (middle) and TNF-α: tumor necrosis factor α (right). **E**, Quantification of BAL neutrophils frequency (left) and absolute count (right) (gated on single live cells, CD45.2$^+$, CD11c$^-$, Siglec-F$^-$, CD11b$^+$, Ly6G$^+$). **F,** Quantification of BAL alveolar macrophages frequency (left) and absolute count (right) (gated on single live cells, CD45.2$^+$, CD11c$^+$, Siglec-F$^+$). **G**, Schematic of control (i.p. PBS, white bars) β-glucan-trained (i. p. β-glucan, green bars) AMs collected from adult WT mice *ex vivo* stimulation with LPS. Differential expression of viral defense genes in response to LPS in β-glucan-trained AMs. Data were analysed using one-way ANOVA followed by Dunn's multiple comparisons test. * $p<0.05$, ** $p<0.01$, *** $p<0.001$.

Supplementary Figure 3. β-glucan-mediated AM reprogramming is independent of Dectin-1 and type I interferon signaling. A, Schematic of the β-glucan-induced training seven days before lipopolysaccharide (LPS)-induced acute lung injury (ALI) model. Experiments were performed in sex- and age-matched 10-12 weeks old control (i.p. PBS, white bars) and trained (i. p. β-glucan, green bars) *Dectin1$^{-/-}$* mice. **B**, BAL chemokine C-X-C motif ligand 1 (CXCL1) concentration (left) and pro-inflammatory cytokines (IL-6: interleukin-6 (middle) and TNF-α: tumor necrosis factor α (right)). **C**, Quantification of BAL neutrophils frequency (left) and absolute count (right) (gated on single live cells, CD45.2$^+$, CD11c$^-$, Siglec-

F⁻, CD11b⁺, Ly6G⁺). **D,** Quantification of BAL alveolar macrophages frequency (left) and

absolute count (right) (gated on single live cells, CD45.2$^+$, CD11c$^+$, Siglec-F$^+$). **E**, Schematic

of control (i.p. PBS, white bars) β-glucan-trained (i. p. β-glucan, green bars) AMs collected

from adult *Dectin1$^{-/-}$* mice *ex vivo* stimulation with LPS. **F**, Chemokine C-X-C motif ligand 1

(CXCL1) and tumor necrosis factor α (TNF-α) concentrations after *ex vivo* LPS stimulation.

G, Schematic of control (i.p. PBS, white bars) β-glucan-trained (i. p. β-glucan, green bars)

AMs collected from adult *IfnaR$^{-/-}$* mice *ex vivo* stimulation with LPS. **H**, Chemokine C-X-C

motif ligand 1 (CXCL1) and tumor necrosis factor α (TNF-α) concentrations after *ex vivo* LPS

stimulation. Data were analysed using one-way ANOVA followed by Dunn's multiple

comparisons test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3