Central histaminergic system reduces response gain and slows visual processing in the retina and lateral geniculate nucleus of awake mice

Authors: Matteo Tripodi[1], Hiroki Asari*[1]

Affiliations:

1. Epigenetics and Neurobiology Unit, EMBL Rome, European Molecular Biology Laboratory, Monterotondo, 00015, Italy

*Corresponding author: Hiroki Asari, E-mail: asari@embl.it

Author contributions: M.T. and H.A. designed the study; M.T. performed experiments; M.T. and H.A. analyzed the results; and M.T. and H.A. wrote the manuscript.

Competing interests: The authors declare no competing financial interests.

Key words: retinal ganglion cells; lateral geniculate nucleus; histaminergic modulation; tuberomammillary nucleus.

Acronyms: AAV, adeno-associated virus; ANOVA, analysis of variance; DLP, digital light processing; DRN, dorsal raphe nucleus; HDC, histidine decarboxylase; GABA, γ-aminobutyric acid; IQR, interquartile range; IR, infrared; LED, light emitting diode; LGN, lateral geniculate nucleus; OT, optic tract; PFA, paraformaldehyde; PSAM/PSEM, pharmacologically selective actuator/effector module; RF, receptive field; RGC, retinal ganglion cell; STA, spike-triggered average; t-SNE, t-distributed Stochastic Neighbor Embedding; UV, ultraviolet.

1   Abstract

Sensory processing is dynamically modulated by an animal's behavior and internal states.

Growing evidence suggests that such modulation starts from early stages, already at the retina,

but the underlying mechanisms remain elusive. Combining pharmacological and

chemogenetic tools with single-unit extracellular recordings in awake head-fixed mice, here

we identified that the visual responses of retinal ganglion cells and the lateral geniculate

nucleus were both made weaker and slower by histaminergic projections from the

tuberomammillary nucleus of the posterior hypothalamus. The observed changes in the visual

responses were, however, not directly linked with histaminergic modulation of pupil dynamics

or locomotion behavior. Instead, our computational modelling analysis suggests that the

temporal response modulations arise from changes in the intrinsic properties of the circuit,

such as gain modulation via the histamine H1 receptors in the retina. Facilitation of visual

processing at low histamine levels may offer an ethological advantage, enabling animals to

respond rapidly to visual threats during periods of reduced activity.

15   Introduction

In the mammalian brain, histaminergic neurons are exclusively located in the

tuberomammillary nucleus (TMN) and surrounding areas of the posterior hypothalamus (Haas

and Panula, 2003). They express histidine decarboxylase (HDC) and innervate widely across

the brain (Lin et al., 2023), including the retina (Gastinger et al., 1999; Repérant et al., 2007;

Greferath et al., 2009; Warwick et al., 2024). These neurons remain silent during sleep, but

increase their firing during wakefulness and reach their maximal activity at the state of high

vigilance, much as other monoaminergic systems do, such as the serotonergic neurons in the

dorsal raphe nucleus (DRN) of the brain stem (Lin et al., 2023; Takahashi et al., 2006; Thakker,

2011). Brain histamine exerts its effects on neurons via three types of receptors: the H1 and

H2 receptors mediate excitation of postsynaptic cells, while the H3 receptors serve as an auto-

receptor to presynaptically inhibit histamine release (Haas and Panula, 2003). The role of the

H4 receptors in the central nervous system remains unknown (Strakhova et al., 2009). The

central histaminergic system has been implicated in regulating an animal's internal state and

a broad range of behavior, including the sleep-wake cycle (John et al., 2004; Mochizuki et al.,

1992), arousal (Parmentier et al., 2002; Scammell et al., 2018), suppression of food intake

(Khouma et al., 2023; Valdés et al., 2010; Wang et al., 2015) and learning (Kamei et al., 1993;

Klapdor et al., 1994; Orsetti et al., 2001; Rubio et al., 2002). Histaminergic neurons, however,

corelease γ-aminobutyric acid (GABA) and interact with other neuromodulatory systems, such

as the cholinergic and serotonergic systems (Khateb et al., 1995; Brown et al., 2002; Lin et

al., 2023). This makes it difficult to dissect the effects of histamine by itself on neural circuit

functions. In particular, histaminergic modulation of sensory processing remains largely

elusive (Lee and Dan, 2012; McCormick et al., 2020).

The retina performs the first stage of visual processing, and conveys visual information

to the brain via spike trains of retinal ganglion cells (RGCs). Growing evidence supports that

the activity of RGCs is correlated with various behavioral measures, such as locomotion and

pupil size (Schröder et al., 2020; Liang et al., 2020; Reggiani et al., 2023; Boissonnet et al.,

2023), pointing to a more dynamic role of the retina than is assumed for a deterministic

processor. The vertebrate retina is part of the central nervous system equipped with both

intrinsic and extrinsic neuromodulation mechanisms, much as other brain regions are (Lee

and Dan, 2012; McCormick et al., 2020). For instance, dopamine is synthesized and released

by a group of retinal amacrine cells under the control of circadian rhythm, and contributes to

adjusting the retinal circuitry for rod or cone vision by modulating gap-junction couplings

(Bloomfield and Völgyi, 2009). Such retinal intrinsic mechanisms of neuromodulation are,

however, much slower than needed to modulate retinal function to follow rapid behavioral

changes (Jacoby et al., 2018; Goel and Mangel, 2021). To this end, the retina seems to exploit

the centrifugal system to receive inputs from a variety of brain regions in a species-specific

manner (Repérant et al., 2007). For example, histaminergic and serotonergic projections from

TMN and DRN, respectively, have been anatomically described in the mammalian retina

(Gastinger et al., 1999; Greferath et al., 2009; Labandeira-Garcia et al., 1990; Warwick et al.,

2024). However, compared to the isthmo-optic pathway in birds (Wilson and Lindstrom, 2011) or the olfacto-retinal circuit in fish (Esposti, et al., 2013), little is known about the function of the mammalian centrifugal system (Vereczki et al., 2024).

Combining in vivo single-unit extracellular recordings from either the lateral geniculate nucleus (LGN) or the optic tract (OT) with pharmacological and chemogenetic tools, here we explored how histamine affects the early visual processing in awake head-fixed mice. During the recording sessions, we simultaneously monitored the subject animal's behavior, such as pupil dynamics and locomotion, to better correlate the effects of histamine on the visual response properties with those on the behavioral measures. We also took a computational approach to clarify the net effect of histamine on the visual responses from functional viewpoints. Because visual processing in the brain fully relies on the signals from the retina, characterization of the retinal operation in vivo is indispensable to better understand how the visual system works.

Results

Histamine leads to weaker and slower visual responses in the mouse lateral geniculate nucleus

To investigate how histamine influences the mouse visual system, we first performed in vivo single-unit extracellular recordings from the lateral geniculate nucleus (LGN) in awake head-fixed mice (Figure 1A,B and Supplementary Figure 1), and compared the visual response properties of the recorded cells before and after perturbing the central histaminergic system (Figures 1 and 2; see Methods for details). Besides directly examining the responses to repeatedly presented visual stimuli, we employed stimulus ensemble statistical techniques ("reverse correlation") to systematically characterize the visual response properties of the recorded cells in the framework of a linear-nonlinear cascade model (Boissonnet et al., 2023). In particular, using full-field white-noise stimuli, we analyzed histaminergic effects on 1) the linear temporal filter, estimated by the spike-triggered average (STA) stimulus – i.e., the mean

stimulus that triggered spiking responses – under a given behavioral state; and 2) the static

nonlinear gain function, i.e., an instantaneous mapping of the STA output to the neural

responses (Figure 2B). We also functionally mapped the receptive field (RF) of the cells by

calculating the linear spatial filter from their responses to random binary "checkerboard" stimuli,

where the contrast of each spatial location was randomly and independently inverted over

time. To manipulate the histaminergic system, we used two different approaches

(Supplementary Figure 2). The first was the chemogenetic activation of the central

histaminergic system (Magnus et al., 2011). We injected Cre-dependent adeno-associated

viruses (AAVs) carrying pharmacologically selective actuator module (PSAM)-5HT3HC

channels into the tuberomammillary nucleus (TMN) of the posterior hypothalamus in histidine

decarboxylase (HDC)-Cre transgenic mice, and then achieved selective activation of the

target neurons by systemic administration of the pharmacologically selective effector molecule

(PSEM; 5mg/kg). The second approach was the use of pharmacological tools in wild-type

mice (Figure 1C; Thakkar, 2011). Specifically, we used chlorphenamine (5 mg/kg; Frisch et

al., 1997; Valle-Baurista et al. 2022), an antagonist of a postsynaptic histamine receptor H1;

and ciproxifan (12 mg/kg; Browman et al., 2004; Muindi et al., 2015), an antagonist of the H3

auto-receptor that presynaptically inhibits histamine release. These drugs effectively down-

and up-regulate the action of histamine in the nervous system, respectively. Thus, resulting

changes in visual processing, if any, were expected to be in opposite directions. While we let

the animals move freely under the head-fixed condition, the reverse correlation analysis was

done on the periods when the animals stayed stationary (<2 cm/s) to minimize the side effect

of locomotion behavior (Supplementary Figure 3; Erisken et al., 2014; Schröder et al., 2020).

Upon chemogenetically activating HDC+ neurons in TMN, we found that LGN neurons

generally showed weaker and slower visual responses. The peak responses to stimulus onset

and offset were both significantly reduced in amplitude and prolonged in latency (Figure 1D-

F). Moreover, the mean firing rate during the white-noise visual stimulation became

significantly lower (median, -4.7 Hz; interquartile range (IQR), 7.9 Hz; p<0.001, Wilcoxon

signed-rank test here and thereafter unless otherwise noted; n=52 from 3 animals); and the

response gain also significantly decreased (median, -33%; IQR, 55%; p<0.001) that we

computed as the ratio of the estimated static nonlinearity functions before and after PSEM

injection (Figure 2C,G). The temporal filter became elongated regardless of their functional

cell types (Supplementary Figure 4), leading to significantly longer peak latencies (median,

+31 ms; IQR, 33 ms; p<0.001) and lower peak frequencies (median, -16%; IQR, 47 %;

p<0.001).

Consistent results were obtained when histamine release was pharmacologically up-

regulated. Specifically, after the application of ciproxifan (H3 antagonist), both the onset and

offset peak responses became weaker and longer (Figure 1G-I). Furthremore, during white-

noise sitmulation, we found significantly reduced mean firing rates (median, -3.0 Hz; IQR, 5.5

Hz; p<0.001; n=126 cells from 4 animals) and response gain (median, -23%; IQR, 65%;

p=0.002); and the linear temporal filters showed significantly longer peak latency (median, +16

ms; IQR, 11 ms; p<0.001) and lower peak frequency (median, -39%; IQR, 20%; p=0.08;

Figure 2D,H). These effects of the pharmacological treatment were less prominent than those

of the chemogenetic manipulation (Figures 1P,Q and 2K-N). None of these treatments

significantly affected the estimated RF size at the population level (Supplementary Figure 5).

In contrast, chlorphenamine (H1 antagonist) led to marginal changes in the visual

response properties of LGN neurons in the opposite direction. The onset/offset stimulus

responses were faster in some cells, but the changes were not statistically significant at the

population level (Figure 1J-L). Likewise, we observed a tendency towards an increased mean

evoked firing rate (median, +1.1 Hz; IQR, 6.8 Hz; p=0.09; n=110 cells from 3 animals) and

response gain (median, +14%; IQR, 94%; p=0.015), and facilitated response kinetics with

significantly higher peak frequency of the linear temporal filter (median, +4%; IQR, 15%;

p=0.02) but no significant changes in its peak latency (median, 0 ms; IQR, 5 ms; p=0.7) or the

RF size (Supplementary Figure 5). Taken together, these results indicate that histamine

reduces the response gain and slows down the response kinetics in the mouse LGN.

Figure 1: Histamine reduced LGN visual responses in awake head-fixed mice. *A,B*: schematic diagram of LGN recordings (A) and a representative histological image showing the electrode location (B; DiI stain, red). *C*: schematic diagram of histamine receptor localizations and their cellular function. Chlorphenamine blocks the postsynaptic receptor H1 (orange) that mediates the action of histamine postsynaptically, while ciproxifan blocks the autoreceptor H3 (green) that inhibits presynaptic histamine release. *D*: visual responses of a representative LGN cell to full-field contrast-inverting stimuli (2 s intervals) before and after chemogenetically activating HDC+ cells in TMN: *top*, spike raster across trials; *bottom*, zoom-in of the spike raster around stimulus onset (-50 ms to 250 ms, red shade on top) and peri-stimulus time histogram, showing a delay of the onset response after the chemogenetic treatment. *E,F*: pairwise comparison of the LGN population responses before and after the chemogenetic treatment (E, peak latency: untreated, 39±17 ms, median ± median absolute deviation; treated, 64±20 ms; p<0.001, Wilcoxon signed-rank test with Bonferroni correction;

F, peak firing rate: untreated, 82±33 Hz; treated, 57±26 Hz; p=0.04, n=26 LGN cells from 3
animals): circle, ON peak; cross, OFF peak; gray, inter-quartile range. *G-I*: corresponding
data before and after ciproxifan administration (G, representative cell's response, with a zoom-
in around stimulus offset; H, peak latency: 37±22 ms versus 50±26 ms, p<0.001; I, peak firing
rate: 95±35 Hz versus 71±31; p<0.001, n=60 LGN cells from 4 animals). *J-L*: corresponding
data before and after chlorphenamine administration (J, representative cell's response; K,
peak latency: 42±20 ms versus 38±16 ms, p=0.3; L, peak firing rate: 81±43 Hz versus 76±33;
p=0.5, n=56 LGN cells from 3 animals). *M-O*: corresponding data before and after saline
administration (M, representative cell's response; N, peak latency: 35±22 ms versus 33±21
ms, p=0.2; O, peak firing rate: 97±48 Hz versus 89±34; p=0.5, n=53 LGN cells from 3 animals).
P,Q: cumulative distributions of modulation indices before and after each treatment (in
corresponding colors): P, peak latencies; Q, peak firing rate; *, p<0.05; **, p<0.01; ***, p<0.001
from the post-hoc test against the saline (control) condition on the average group ranks
(Kruskal-Wallis test).

Supplementary Figure 1: spike waveform and autocorrelation of spike trains of representative cells across different conditions. *A*: average spike waveform (top) and auto-correlogram (bottom) of two representative LGN cells (left and right) before (black) and after (red) chemogenetic activation of TMN HDC+ cells. *B-D*: corresponding data for ciproxifan (B, green), chlorphenamine (C, orange), and saline administration (D, blue), each with two representative LGN cells. *E-I*: corresponding data for RGCs from optic tract recordings (E, chemogenetic activation of HDC+ cells in TMN; F, chemogenetic activation of HDC+ cells outside TMN; G, ciproxifan; H, chlorphenamine; I, saline).

172
Figure 2: Histamine made the visual responses of LGN weaker and slower in awake
head-fixed mice. *A,B*: Schematic diagram of LGN recordings (A) and linear-nonlinear
cascade model (B). Reverse-correlation was used to estimate the linear filter and static
nonlinearity that represent response kinetics and strength, respectively. *C*: Linear filters of two
representative LGN cells before (gray) and after chemogenetic activation of HDC+ cells in
TMN (red). We focused only on those data when the animal stayed stationary (<2 cm/s) to
minimize the side effect of locomotion behavior (Supplementary Figure 3). *D-F*:
Representative LGN linear filters before (gray) and after pharmacological treatment (D, 12
mg/kg ciproxifan, green; E, 5 mg/kg chlorphenamine, orange) or saline injection (F). *G-J*: Static
nonlinearities averaged across LGN population before and after treatment in corresponding
colors (G, chemogenetics, n=52 from 3 animals; H, ciproxifan, n=126 from 4 animals; I,
chlorphenamine, n=110 from 3 animals; J, saline, n=92 from 3 animals): line, median; gray
shade, 25 and 75 percentiles. *K-N*: Cumulative distributions of the modulation index before
and after each treatment (in corresponding color): peak latency (K) and frequency (L) of the
linear temporal filters, mean evoked firing rate during stimulation (M), and static nonlinearity
(i.e., response gain; N): *, p<0.05; **, p<0.01; ***, p<0.001 from the post-hoc test against the
saline control on the average group ranks (Kruskal-Wallis test).

Supplementary Figure 2: Pharmacological and chemogenetic approaches to manipulate the central histaminergic system in the mouse brain. *A,B*: Viral delivery of PSAM 5HT3HC channel to HDC+ cells in the TMN (marked with dotted lines) of the posterior hypothalamus (A) or those in the anterior hypothalamus (B; control). *C-E*: Histological image examples of anterograde tracing of HDC+ cells in TMN. Labelled axons were found in the optic chiasm (C) or the optic nerve (D) after injecting rAAV9::CAG-FLEX-Synaptophysin-GFP or AAV2/1::CAG-FLEX-axon-GFP, respectively, in the TMN of HDC-Cre mice. However, no visible signal was detected in the isolated retinal tissue of all animals examined (n=9; see panel E for example).

Supplementary Figure 3: Locomotion of an animal facilitates the visual responses of LGN and RGCs. *A*: probability distribution of an animal's locomotion speed during recordings before (gray) and after treatment: from left to right, chlorphenamine (CP, orange), ciproxifan (CXN, green), PSAM/PSEM for HDC+ cells in non-TMN (purple) or TMN (red), and saline (blue). *B*: the fraction of running period (>2cm/s) before and after each treatment. No significant change was observed (p=0.07, Kruskal-Wallis test). *C*: the median locomotion speed (during which animals moved at >2 cm/s) before and after each treatment. No significant change was observed (p=0.3). *D*: comparison of the LGN population response properties between stationary and running periods (n=104 from 5 animals with a running period ranging between 20% and 80%): from left to right, peak latency (50±19 ms versus 48±19 ms, median ± median absolute deviation; p<0.001, Wilcoxon signed-rank test), peak frequency (8.2±1.9 Hz versus 8.2±3.0 Hz; p=0.4), mean firing rate (11±6 Hz versus 20±8 Hz; p<0.001). *E*: corresponding data for RGCs (n=44 from 5 animals with a running period ranging between 20% and 80%): from left to right, peak latency (58±19 ms versus 49±18 ms; p<0.001), peak frequency (8.2±2.8 Hz versus 9.2±2.7 Hz; p=0.6), mean firing rate (24±22 Hz versus 23±22 Hz; p=0.5).

Supplementary Figure 4: Functional classification of LGN visual response types and batch effect analysis. *A*: t-SNE embedding of the STAs of LGN cells. Different markers and shadings are used for distinct response categories (shown in B). *B*: Each panel represents one of the six response categories: slow off, off, fast off, fast on, on, and slow on. In each panel, each row represents a cell's STA (color-coded with red and blue hue, indicating positive and negative filter values, respectively); and the overlaid gray line shows the average STAs in each response type. *C-F*: Modulation indices on LGN response characteristics across individual animals: from left to right, peak latency (C), peak frequency (D), mean evoked firing rate (E), and nonlinearity (F). The effects of histamine were largely consistent across animals.

Supplementary Figure 5: Histamine did not affect LGN RF size. *A.B*: Blink (A) and saccade (B) frequencies before and after different treatments: from left to right, chlorphenamine (CP), ciproxifan (CPN), PSAM/PSEM for non-TMN HDC+ cells and TMN HDC+ cells, and saline. None of these changes were statistically significant: p=0.10, Kruskal-Wallis test on blink frequency changes; p=0.15, saccade frequency changes. *C*: Representative time series of the eye position (X- and Y-coordinates of the pupil center extracted from eye-tracking camera images) during white-noise "checkerboard" stimulation (black/red, centered and stable eye position used for reverse correlation; gray, non-centered or non-stable period excluded from the analysis; probability distribution shown on the right), before (left) and after (right) chemogenetic activation of TMN HDC+ cells. *D*: Estimated spatial filter (receptive field; RF) of two example LGN cells before (top) and after (bottom) chemogenetic activation of HDC+ cells in TMN. Note a shift of RF position of all recorded cells due to a shift of the animal's resting

eye position after the treatment (see panel C). *E-H*: Cumulative distribution of the modulation
index of LGN cells before and after treatment (E, receptive field size; F, peak latency; G, peak
frequency; H, mean evoked firing rate): orange, chlorphenamine, n=32 from 3 animals; green,
ciproxifan, N=61 from 4 animals; red, PSAM/PSEM for HDC+ cells in TMN, N=20 cells from 3
animals. Post-hoc test against saline control (N=47 from 3 animals) after Kruskal-Wallis test:
*, p<0.05; **, p<0.01; ***, p<0.001.

Histaminergic modulation starts from the mouse retina

Histaminergic neurons are localized solely in the hypothalamus, most densely in TMN, and

project their axons widely across the brain (Lin et al., 2023), including the mouse retina

(Gastinger et al., 1999; Repérant et al., 2007; Warwick et al., 2024). To test if the observed

histaminergic modulation of the visual processing in LGN originates in the retina, we next

performed in vivo optic tract (OT) recordings to monitor the retinal output responses in awake

head-fixed mice (Figure 3A,B; Boissonnet et al., 2023) before and after manipulating the

central histaminergic system (Figure 3 and Supplementary Figures 6-9).

Consistent with the results in LGN, chemogenetic activation of HDC+ neurons in TMN

led to reduced firing of retinal ganglion cells (RGCs; median, -9.8 Hz; IQR, 13.1 Hz; p=0.01;

n=26 from 4 animals) with lower response gain (median, -28%; IQR, 57 %; p=0.01); and slower

response kinetics with elongated linear temporal filters(Figure 3C,G; peak latency change,

median, 10 ms; IQR, 33 ms; p<0.001; peak frequency change, median, -17%; IQR, 34%,

p=0.003). We also found that the latencies of the peak responses to stimulus onset and offset

were both significantly longer, with a reduced peak firing (Supplementary Figure 6A,F). In

contrast, no change in these response parameters was observed when targeting HDC+

populations in the anterior hypothalamus outside TMN for the PSAM/PSEM activation (firing

rate change, median, 1.7 Hz, IQR, 16.5 Hz, p=0.6; gain change, median, 8%, IQR, 99%,

p=0.5; latency change, median 0 Hz, IQR, 12 Hz, p=0.6; frequency change, median -6%, IQR,

17%, p(U-test)=0.09; N=23 from 3 animals). Neither the onset nor the offset stimulus response

was affected (Supplementary Figure 6B,G). Anterograde tracing of HDC+ cells – via injection

of Cre-dependent AAVs encoding fluorescent marker proteins into the TMN of HDC-Cre

animals – showed labelled axons in the optic chiasm and/or the optic nerve (3 out of 9 animals;

Supplementary Figure 2C,D). Unlike previous reports (Gastinger et al., 1999; Repérant et al.,

2007; Warwick et al., 2024), however, we did not identify any labelled axons in the retinal

tissues (e.g., Supplementary Figure 2E), likely because these projections are sparse and

located far from the cell bodies, making the signals too weak. Nonetheless, our data

collectively support that the histaminergic modulation of the early visual system is mediated

by the centrifugal projections specifically from TMN (Figure 3G-J and Supplementary Figure

2).

The effects of the pharmacological treatments were overall similar between LGN and

RGCs, though certain differences existed. First, pharmacologically blocking the H3 receptors

with ciproxifan led to slower response kinetics in both LGN and RGCs (latency change, median,

10 ms; IQR, 14 ms; $p<0.001$; frequency change, median, -7%; IQR, 25%; $p=0.12$; N=36 from

4 animals), but the firing rates and the response gain remained largely unaffected in RGCs

(Figure 3D,H; firing rate change, median -0.1 Hz; IQR, 12.7 Hz; $p=0.3$; gain change, median,

-7%; IQR, 69%, $p=0.5$). This was also the case with the peak onset/offset stimulus responses

(Supplementary Figure 6C,H). Second, while chlorphenamine (H1 antagonist) drove faster

kinetics with marginally higher firing in both LGN and RGCs, the effects on the temporal filters

were primarily on the peak frequencies in LGN (Figure 2E) but on the peak latencies in RGCs

(Figure 3E; latency change, median, -5 ms; IQR, 8 ms; $p=0.006$; frequency change, median,

+4 Hz; IQR, 25 Hz; $p=0.6$; firing rate change, median, +2 Hz, IQR, 11 Hz; $p=0.09$; gain change,

median, 4%, IQR, 90%, $p=0.3$; N=33 RGCs from 3 animals). Likewise, a shorter peak latency

was observed in the onset/offset stimulus responses of RGCs (Supplementary Figure 6D,I).

Finally, RGC receptive fields did not show significant changes in size, either (Supplementary

Figure 8).

Although diverse effects of histamine across retinal cell types have been reported in

previous ex vivo studies (Gastinger et al 2004; Yu et al., 2009, 2011; Akimov et al., 2010; Vila

et al., 2012; Horio et al., 2018; Warwick et al., 2024), the histaminergic effects observed in this

study were rather uniform across RGC or LGN populations. First, the effects on the peak

responses to the contrast-inverting stimuli did not significantly differ between the stimulus

onset and offset (Figure 1 and Supplementary Figure 2). Second, the effects on the temporal

filter and nonlinearity were not substantially different among the six functional groups that we

identified for RGCs and LGN cells, respectively (Supplementary Figures 4 and 7). Our

datasets are, however, not large enough to identify all the ~40 physiological cell-types reported

thus far (Baden et al., 2016; Rosón et al., 2019); and we could not target a specific cell-type

or tailor visual stimuli for each cell during recordings due to technical limitations. Lastly, we

identified direction-selective (DS) and orientation-selective (OS) cells using the responses to

moving grating stimuli in eight different directions (see Methods for details), but we found no

differences in the DS or OS index values – defined by the normalized population vector length

– before and after each treatment (Supplementary Figure 9).

Taken all together, our data indicate that 1) the mouse early visual processing is

subject to histaminergic modulation controlled by TMN, starting from the retina; 2) histamine

generally suppresses the firing and slows the response kinetics; and 3) response latency and

frequency tuning are likely modulated by different mechanisms.

Figure 3: histamine primarily affected the visual response kinetics of RGCs. *A,B*: Schematic diagram of in vivo optic tract (OT) recordings (A) and a representative histological image showing the electrode location (B; DiI stain, red). *C-F*: Linear filters of two representative RGCs before (gray) and after chemogenetic (A, red) or pharmacological treatment (B, 12mg/kg ciproxifan, green; C, 5 mg/kg chlorphenamine, orange; D, saline, blue). *G-J*: Static nonlinearities averaged across RGC population before and after treatment in corresponding colors (G, chemogenetics, n=26 cells from 4 animals; H, ciproxifan, n=36 from 4 animals; I, chlorphenamine, n=33 from 3 animals; J, saline, n=46 cells): line, median; gray shade, 25 and 75 percentiles. *K-N*: cumulative distributions of the modulation index before and after each treatment (in corresponding color; n=23 from 3 animals for chemogenetic treatment of non-TMN HDC+ cells): peak latency (K) and frequency (L) of the linear temporal filters, mean evoked firing rate during stimulation (M), and static nonlinearity gain function (N); *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$ from the post-hoc test against the saline control condition in the average group ranks (Kruskal-Wallis test).

Supplementary Figure 6: histamine primarily modulated the visual response kinetics of RGCs. *A-E*: Visual responses of a representative RGC to full-field contrast-inverting stimuli (2 s intervals) before and after chemogenetic (A, HDC+ cells in TMN; B, HDC+ cells in non-TMN) or pharmacological treatment (C, ciproxifan; D, chlorphenamine; E, saline); top, spike raster across trials; bottom, zoom-in of the spike raster around stimulus onset or offset (-50 ms to 250 ms, red shade on top) and peri-stimulus time histogram. *F-J*: pairwise comparison of the RGC population responses before and after treatment (top, peak latency; bottom, peak firing rate): *, p<0.05; **, p<0.01, Wilcoxon signed-rank test with Bonferroni correction. F, PSAM/PSEM for TMN HDC+ cells: 57±15 ms versus 66±17 ms peak latency, p=0.005; 122±54 Hz versus 112±66 Hz peak frequency, p=0.08; median ± median absolute deviation, n=16 RGCs from 4 animals. G, PSAM/PSEM for non-TMN HDC+ cells: 42±17 versus 43±14 ms peak latency, p=0.3; 46±30 Hz versus 59±40 Hz peak frequency, p=0.3; n=24 RGCs from 3 animals. H, ciproxifan: 42±31 ms versus 53±31 ms peak latency, p=0.002; 72±80 Hz versus 78±78 Hz peak frequency, p=0.3; n=32 RGCs from 4 animals. I, chlorphenamine: 61±16 ms versus 47±18 ms peak latency, p=0.026; 55±60 Hz versus 102±88 Hz peak frequency, p=0.05; n=22 RGCs from 3 animals. J, saline: 73±30 ms versus 73±31 ms peak latency, p=0.5; 92±50 Hz versus 92±56 Hz peak frequency, p=1; n=37 RGCs from 3 animals. *K,L*: cumulative distributions of the modulation index before and after each treatment (in corresponding colors): K, peak latencies; L, peak firing rate; *, p<0.05 from the post-hoc test against the saline (control) condition on the average group ranks (Kruskal-Wallis test).

Supplementary Figure 7: Functional classification of RGC visual response types and batch effect analysis. *A*: t-SNE embedding of the RGC STAs. Different markers and shadings are used for distinct response categories (shown in B). *B*: Each panel represents one of the six response categories: slow off, off, fast off, fast on, on, and slow on. In each panel, each row represents a cell's STA (color-coded with red and blue hue, indicating positive and negative filter values, respectively); and the overlaid gray line shows the average STAs in each response type. *C-F*: modulation indices on RGC response characteristics across animals: from left to right, peak latency (C), peak frequency (D), mean firing rate (E), and nonlinearity (F). The effects of histamine were generally consistent across animals, and no substantial batch effect was observed.

Supplementary Figure 8: Histamine did not affect RGC RF size. *A*: Representative time
series of the eye position (X- and Y-coordinate of the pupil center extracted from eye-tracking
camera images) during white-noise "checkerboard" stimulus presentation (black/green,
centered and stable eye position used for reverse correlation analysis; gray, non-centered or
non-stable period excluded from the analysis; probability distribution shown on the right), along
with those of firing rate dynamics of two example RGCs and locomotion (from top to bottom),
before (left) and after (right) ciproxifan administration. *B*: Estimated spatial filter (receptive
field) of the two example RGCs (top and middle) before (left) and after (right) ciproxifan
administration. Note a shift of RF position of all recorded cells due to a shift of resting eye
position after the treatment (see panel A). *C-F*: cumulative distribution of the modulation index
of RGCs before and after treatment (E, receptive field size; F, peak latency; G, peak frequency,
H, mean firing rate): orange, chlorphenamine, n=6 cells from 3 animals; green, ciproxifan,
n=25 cells from 4 animals; purple, PSAM/PSEM for HDC+ cells in non-TMN, n=9 cells from 3
animals; red, PSAM/PSEM for HDC+ cells in TMN, n=8 cells from 4 animals. Post-hoc test
against saline control (N=17 from 3 animals) after Kruskal-Wallis test: *, $p<0.05$; **, $p<0.01$.

374
Supplementary Figure 9: Histamine did not affect direction- or orientation-selectivity of
RGCs or LGN cells. *A*: the average firing rate of a representative direction-selective (DS)
RGC in response to moving gratings in eight different directions before (gray) and after
treatment (color-coded): from left to right, PSAM/PSEM for TMN HDC+ cells (red) or non-TMN
cells (purple), ciproxifan (green), chlorphenamine (orange), and saline (blue). *B*: DS indices of
RGC populations before and after treatment. Those with DS index > 0.15 in either condition
were highlighted in the corresponding color (from left to right, n=11, 16, 31, 14, and 30 RGCs,
respectively). No significant change was observed in the DS index values: p>0.2 in all cases
(Wilcoxon signed-rank test). *C*: Corresponding data for the OS indices of RGCs. No significant
change was observed (p>0.7 in all cases). *D,E*: Corresponding population data for the DS and
OS indices of LGN cells (from left to right, n=42, 76, 27, 58 LGN cells; p>0.06 and 0.3 in all
cases, respectively).

Pupil dynamics and locomotion are irrelevant to histaminergic modulation of the early visual processing in mice

387
388

Histamine is a potent neuromodulator implicated in various brain functions and behaviors,

such as sleep-wake cycles and arousal (Haas and Panula, 2003; Thakkar, 2011). Thus, it is

possible that the observed histaminergic effects on the visual response properties in the

mouse early visual system are a result of correlated behavioral modulation. For example, a

larger pupil size leads to stronger responses because more photons reach the retina

(Boissonnet et al., 2023). To address this issue, we examined the relationship between the

histaminergic effects on the RGC/LGN visual responses and simultaneously monitored

behavioral measures (Figure 4 and Supplementary Figure 10).

We first assessed the effects of the pupil size on the RGC/LGN visual responses. In

particular, we performed the reverse correlation analysis using subsampled data with a

constricted or dilated pupil for each recording (below 33 or above 66 percentiles of the pupil

size, respectively; e.g., Figure 4A-C). Under the control condition, we found that the larger the

pupil size, the faster the response kinetics for both RGCs (Figure 4D,E) and LGN cells

(Supplementary Figure 10A,B). We also found that the LGN responses were generally

stronger with a dilated pupil than with a constricted pupil (Supplementary Figure 10C). In

contrast, variations in the mean evoked firing rate of RGCs between constricted and dilated

pupil periods were not uniform across the population, but dependent on their response polarity

(R=0.20, p=0.015, Pearson's correlation with the temporal filter ON-OFF polarity index,

defined by the difference between the peak and valley, normalized by the sum of the two;

n=152 RGCs from 16 animals).

How does this pupil size effect interrelate with the histaminergic effect? As reported

before (Fang et al., 1998), chlorphenamine (H1 antagonist) led to a substantial increase in the

baseline pupil size due to its anticholinergic action (Figure 4K; 67±34%, mean ± standard

deviation of the median pupil size, n=6 animals). We compared the RGC kinetics after blocking

H1 with a constricted pupil (hence relatively slower) to those before the treatment with a dilated

pupil (hence relatively faster), and found no significant difference (Figure 4M). Thus, we cannot

exclude a possible contribution of the baseline pupil dilation to the response facilitation effect

of chlorphenamine. In contrast, chemogenetic activation of TMN HDC+ cells resulted in a

slight increase in the baseline pupil size ($2\pm23\%$, n=6 animals), while ciproxifan (H3

antagonist) treatment led to a slight decrease (Figure 4K; $-7\pm18\%$, n=8 animals), though they

both resulted in slower RGC/LGN response dynamics (Figure 3 and Supplementary Figure 6).

Such discrepancies are likely due to side effects of the manipulations beyond histamine: e.g.,

ciproxifan has modest affinity to adrenergic receptors (Esbenshade et al., 2003) and inhibits

monoamine oxidase A and B (Hagenow et al., 2017), while some HDC+ cells co-release

histamine and GABA (Yu et al., 2015). Here we found that the response kinetics of RGC/LGN

after these manipulations with a dilated pupil (hence relatively faster) were nevertheless

slower than those under the control condition with a constricted pupil (hence relatively slower;

Figure 4L). Therefore, the resulting slow response kinetics after up-regulating histamine

cannot be attributable to the pupil size effects.

We next analyzed the effect of pupil dynamics. In general, changes in pupil size occur

much more slowly than the response dynamics of the early visual system: e.g., pupillary light

reflex has a latency of hundreds of milliseconds and a time-constant of about a second

(Grozdanic et al., 2009; Liu et al., 2017). Nevertheless, when the pupil is dilating, presented

stimuli on the screen should appear brighter on the retina than intended, allowing RGCs to

respond to physically darker stimuli and thereby biasing the response polarity balance.

Conversely, one would expect the opposite effect when the pupil is constricting. We indeed

identified the corresponding changes in the ON/OFF balance of both RGC and LGN responses

(Figure 4G and Supplementary Figure 10D), using the temporal filter ON-OFF polarity index.

An increase of the mean evoked firing rate was also observed in LGN, but not in RGCs, during

pupil dilation (Figure 4J and Supplementary Figure 10G). However, no change in the response

kinetics were observed in both RGCs and LGN between dilating and constricting pupil periods

(Figure 4H,I and Supplementary Figure 10E,F). The frequencies of saccades or eye blinks did

not change, either, after the pharmacological or chemogenetic treatments (Supplementary

Figure 5A,B). We thus conclude that the pupil dynamics should have little to do with the

observed histaminergic effects on the RGC/LGN visual responses.

Locomotion has been reported to profoundly affect information processing throughout

the mouse visual system (Neil and Stryker, 2010; Erisken et al., 2014; Dipoppa et al., 2018;

Savier et al., 2019; Schröder et al., 2020). When we compared the temporal filter

characteristics during running and stationary periods (at a threshold of 2 cm/s), we indeed

found that locomotion facilitated visual responses of RGCs and LGN cells (Supplementary

Figure 3D,E). Importantly, we excluded these running periods from the analysis of the

histaminergic effects described above. Furthermore, the mice remained stationary most of the

time under the head-fixed condition (Supplementary Figure 3A), and neither the chemogenetic

nor the pharmacological treatments resulted in a significant change in either the fraction of

time spent running (Supplementary Figure 3B) or the median running speed (Supplementary

Figure 3C). Thus, locomotion and its effects on RGC/LGN responses are irrelevant to the

observed histaminergic modulation, and likely involve different mechanisms.

Taken together, neither pupil dynamics nor locomotion behavior was causally linked

with the histaminergic modulation of the mouse early visual system. We conclude that the

central histaminergic system modulates the mouse vision, effectively leading to slower and

weaker responses.

Figure 4: histaminergic modulation of pupil dynamics was not causally linked with that of RGC/LGN visual responses. *A*: Representative time series of the pupil size (light yellow, top 33 percentile; dark yellow, bottom 33 percentile) and its time derivative (light blue, positive; dark blue, negative) during randomly flickering full-field stimulus presentation, along with those of the firing rate dynamics of two example RGCs and locomotion (from top to bottom). *B*: Estimated linear filter and nonlinearity for a representative RGC (B, cell 1) using constricted or dilated pupil periods (top) or using constricting or dilating pupil periods (bottom). Note a temporal shift of the linear filter in the former case, while a shift in the filter strength in the latter. *C*: corresponding data for another RGC (cell 2). *D-F*: comparison of the RGC population

response properties between constricted and dilated pupil periods (n=152 RGCs from 16
animals): D, peak latency, 65±19 ms versus 60±17 ms, median ± median absolute deviation,
p<0.001, Wilcoxon signed-rank test; E, peak frequency, 8.2±2.6 Hz versus 8.2±2.7 Hz, p=0.6;
F, mean firing rate, 26±19 Hz versus 28±18 Hz, p=0.6. *G-J*: comparison of the RGC
population response properties between constricting and dilating pupil periods: G, ON-OFF
polarity index of the temporal filters, -0.02±0.33 versus -0.25±0.33, p<0.001; H, peak latency,
62±18 ms versus 63±18 ms, p=0.9; I, peak frequency, 8.3±2.8 Hz versus 8.3±3.0 Hz, p=0.9;
J, mean firing rate, 26±18 Hz versus 28±19 Hz, p=0.7. *K*: probability distributions of relative
pupil size before and after treatment: from left to right, PSAM/PSEM for TMN HDC+ cells (red)
or non-TMN cells (purple), ciproxifan (green), chlorphenamine (orange), and saline (blue). *L*:
comparison of peak latencies before treatment with constricted pupil versus after treatment
with dilated pupil. Histamine nevertheless significantly slowed the RGC population responses
(arrow heads): *, p<0.05; **, p<0.01; ***, p<0.001, Wilcoxon signed rank test. *M*: comparison
of peak latencies before treatment with dilated pupil versus after treatment with constricted
pupil. Effects of H1 blocker could not be distinguished from those of pupil dilation (arrow head,
p=0.5).

486
**Supplementary Figure 10: Histaminergic effects on pupil dynamics are irrelevant to
those on LGN visual responses.** *A-C*: comparison of the LGN population response
properties between constricted and dilated pupil periods (n=350 RGCs from 13 animals): A,
peak latency, 51±20 ms versus 48±19 ms, median ± median absolute deviation, p<0.001,
Wilcoxon signed-rank test; B, peak frequency, 7.9±2.1 Hz versus 8.1±2.4 Hz, p=0.004; E,
mean firing rate, 11±7 versus 15±7 Hz, p<0.001. *D-G*: comparison of the LGN population
response properties between constricting and dilating pupil periods: D, ON-OFF polarity index
of the temporal filters, 0.07±0.22 versus -0.06±0.27, p<0.001; E, peak latency, 49±19 ms
versus 49±19 ms, p=0.8; F, peak frequency, 7.9±2.0 Hz versus 8.1±2.1 Hz, p=0.3; G, mean
firing rate, 11±7 Hz versus 14±7 Hz, p=0.004. H: comparison of peak latencies before
treatment with constricted pupil versus after treatment with dilated pupil: from left to right,
PSAM/PSEM for TMN HDC+ cells (red), ciproxifan (green), chlorphenamine (orange), and
saline (blue): **, p<0.01; ***, p<0.001, Wilcoxon signed rank test. *I*: comparison of peak
latencies before treatment with dilated pupil versus after treatment with constricted pupil.

Response gain modulation is more strongly correlated with peak latency modulation than with peak frequency modulation

What are the mechanisms underlying the histaminergic modulation of the early visual

processing? To address this question, we next took a computational approach to examine the

relationship between different aspects of the visual response properties that we identified to

be affected by histamine (Figure 5). In particular, we simulated the activity of an integrate-and-

fire neuron with Poisson spiking and refractoriness in response to white-noise input stimuli

(Figure 5A), and processed the output spike trains in the framework of the linear-nonlinear

cascade model to characterize the simulated response properties in the same way as the

experimental data (Figure 2B). Here we varied two model parameters, gain and resting

baseline, to examine the effects of their changes on the following four aspects of the simulated

responses: spike counts, the peak latency and frequency of the estimated linear temporal filter,

and the slope of the estimated static nonlinearity (expressed as half-wave rectification). The

rest model parameters were fixed in all simulations, including the underlying linear filter given

by a trigonometric function with logarithmic scale in time, and the refractory period filter given

by an exponential function. The spike threshold was drawn from a random uniform distribution

ranging from 0 to 1 for each time point to approximate the Poisson spiking properties.

When the model neuron had a higher gain or higher resting baseline, subthreshold

signal reached the spike threshold faster and more frequently; hence, total spike counts

became higher (Figure 5B) and the peak latency of the estimated linear temporal filter

appeared shorter (Figure 5C). In contrast, the apparent peak frequency of the simulated

responses was dependent more strongly on the resting baseline than on the gain (Figure 5D).

These changes in the estimated linear temporal filter properties were consistent with the

experimental observation, where the magnitude of the response gain modulation by

histaminergic perturbation was significantly correlated with that of the peak latency modulation

(Figure 5F,H), but much less with that of the peak frequency modulation (Figure 5G,I). This

agreement between the observed and simulated data indicates that, despite the simplicity, our

model captures aspects of neuronal response modulation to a first approximation.

The model analysis supports that the histaminergic effects on RGCs should involve at

least gain and baseline modulations. First, the H1 receptors likely mediate the gain modulation

in RGCs, given that chlorphenamine (H1 antagonist) led to changes in the peak latency but

not the peak frequency, along with changes in the estimated static nonlinearity largely by

scaling (Figure 3). Second, the peak frequency was modulated as well as the peak latency of

RGCs upon chemogenetic activation of HDC+ neurons in TMN, accompanied by a shift of the

estimated static nonlinearity. This implies changes in the baseline resting potential of the cells.

Blocking the H3 receptors, however, led to changes in the RGC response kinetics without

much affecting the mean firing rates. Furthermore, the net effects of histamine perturbations

on LGN responses were more diverse (Figures 1 and 2) and not fully explainable by the simple

model (Figure 5). Thus, we expect that mechanisms besides gain and baseline modulations

are also involved, such as neural circuit effects through retinal amacrine cells for example

(Horio et al., 2018).

Figure 5: Gain modulation accompanies the peak latency modulation but not the peak frequency modulation via histaminergic effects. *A-E*: Using a computational model outlined in A, here we examined the consequences of varying the baseline and gain levels on the response features, such as spike counts (B), peak latency (C) and frequency (D) of the estimated linear filter from the simulated responses, and slope of static nonlinear gain function of the model neuron (E). See Methods for details. *F,G*: Relationship between the magnitudes of the gain modulation and response kinetics modulation by perturbing the histaminergic system for LGN neurons (chemogenetics, red; ciproxifan, green; chlorphenamine, orange; n=300 from 10 animals). Cell types are indicated by different markers (Supplementary Figures 4 and 7). A stronger correlation was found for the peak latencies (F, R = –0.35, p<0.001) than for the peak frequencies (G, Pearson's R = 0.12, p=0.18). Here we combined all data across cell-types and experimental conditions to assess the relationship between the two dependent variables as a result of histamine level manipulations. *H,I*: Corresponding figure panels for RGC responses (n=101 from 11 animals). Significant correlation to the gain modulation was found for the peak latency modulation (R = –0.26, p = 0.03), but not for the peak frequency modulation (R= 0.22, p = 0.11).

Discussion

Here we provide in vivo electrophysiological evidence for histaminergic modulation of the mouse early visual system: the more the histamine released, the slower and weaker the LGN and RGC visual responses in awake head-fixed mice (Figures 1-3). Anterograde tracing of TMN HDC+ neurons confirmed the projection of their axons into the optic chiasm and optic nerve, but not in the retinal tissue (Supplementary Figure 2C-E). Nonetheless, here we used the same mouse lines as previous studies that demonstrated the presence of histaminergic HDC+ axons in the retina (Greferath et al., 2009; Warwick et al., 2024). Moreover, we observed modulation at the level of RGC axons in the optic tract in vivo (Figure 3 and Supplementary Figure 6). It is intriguing that punctate signals of presynaptic markers expressed in TMN HDC+ cells were identified in the optic chiasm (Supplementary Figure 2C), raising a potential role for axonal modulation. However, its contribution should be minimal, if any, because RGC axon potentials should reach the recording site in the optic tract within 1 ms after exiting the eye cup (optic nerve length in mice, <10 mm; conduction velocity, around 10 m/s; Sumitomo et al., 1969). While axonal conduction velocity depends on various factors, including myelination and channel density, its modulation should then occur only on a much shorter time-scale than the observed changes in RGC/LGN response kinetics (up to tens of milliseconds). Taken together, we suggest that the histaminergic modulation likely arises in the retina.

Previous ex vivo studies have reported diverse histaminergic effects in all five major cell types in the retina across species (Vila et al., 2012; Yu et al., 2009, 2011; Horio et al., 2018). Primate RGCs, for instance, either increase or decrease their baseline activity, while their visual responses typically become weaker and slower after histamine application ex vivo (Gastinger et al 2004; Akimov et al., 2010). In contrast, rodent RGCs mostly show an increase in the baseline activity, with various changes in their visual response properties in a histamine dose-dependent manner (Gastinger et al 2004; Warwick et al., 2024). Some of those changes are interrelated: e.g., a broadening of direction-selectivity (DS) tuning can be explained as a

natural consequence of the increased baseline activity because a decrease of the DS index

(DSI) should follow by definition. Indeed, a reduction of DSI was reported at RGC axon

terminals, following an increased baseline activity via arousal or locomotion (Schröder et al.,

2020; Liang et al., 2020). However, the mechanisms underlying the histaminergic modulation

of the retina remain mostly unclear.

We took a computational modelling approach to better understand the net effect of

histamine from functional viewpoints (Figure 5), rather than pursuing the underlying

biophysical mechanisms. For example, we suggest that the H1 receptors mediate gain

modulation in RGCs: chlorphenamine (H1 antagonist) affected the response latency but not

the frequency tuning (Figure 3), a distinctive feature of gain modulation identified by our

computational modelling analysis (Figure 5). This action of histamine does not have to be

directly on RGCs expressing the H1 receptors, but can be mediated indirectly by circuit

mechanisms. Indeed, histamine affects both inward and outward currents of amacrine cells in

the mouse retina (Horio et al., 2018). The effects of histamine can then be more profound than

gain modulation, much as we observed diverse outcomes in RGCs and LGN when different

pharmacological and chemogenetic perturbation tools were employed (see also Warwick et

al., 2024).

Correlation between histamine levels and various behavioral measures has been

widely reported (White and Rumbold, 1988; Haas et al., 2008). For instance, the activity of

central histaminergic neurons generally follows the circadian rhythm, showing high levels of

activity during the active period (i.e., night time for mice as nocturnal animals) and low levels

during the sleep period (Mochizuki et al., 1992; Takahashi et al., 2006). Here we performed

all our experiments during daytime (ZT4-ZT9). Moreover, the mice remained calm under head

fixation, as evidenced by their low spontaneous running behavior (Supplementary Figure 3),

following a week of habituation sessions before recordings. The retinal histamine level of the

subject mice was then expected to be relatively low during the recordings, given that the

mammalian retina contains a comparable level of histamine with other brain areas (around 20-

140 ng/g wet tissue, i.e., about 0.1-1 µM; Nowak, 1993). This will explain why we found

stronger effects by chemogenetically activating HDC+ neurons in TMN or pharmacologically

blocking H3 receptors (mimicking an increase of histamine) than by administering H1

antagonists (mimicking a decrease of histamine). However, regulation of the histamine release

and metabolism is suggested to be both tissue- and specific-specific (Nowak, 1993): e.g., light

stimulation of dark-adapted rabbits at night leads to a substantial reduction of histamine in the

retina and the optic nerve (by about 40-50%), but not in other parts of the central nervous

system. It is a future challenge to determine the exact level of retinal histamine and its

fluctuation under more precisely controlled behavioral contexts.

Our observations on the histaminergic effects have certain discrepancies from those

reported in previous studies. For example, using the mouse retinal explants, Warwick et al.

(2024) showed that histamine increased the baseline activity of RGCs. This effect was dose-

dependent, and saturated at 5 µM histamine with an increased baseline firing by ~4 Hz on

average. We instead found a reduced firing upon H3 antagonist application or chemogenetic

activation of TMN HDC+ cells, while an enhanced firing after blocking H1 receptors (Figure 3

and Supplementary Figure 6). This might be because the baseline activity of RGCs in awake

mice is already high, by ~20 Hz compared to the ex vivo condition (Boissonnet et al., 2023).

Alternatively, the enhancement of the baseline activity in retinal explants could be an artefact

of histamine overdose. An isolated retinal tissue should have no baseline histamine because

the retina itself does not produce histamine (Gastinger et al., 1999; Greferath et al., 2009).

However, bath-application of histamine at the physiological range (0.1-1 µM; Nowak, 1993) or

that of H3 antagonists had no effect on the baseline firing ex vivo (Warwick et al., 2024).

Moreover, chemogenetic activation of HDC+ axon terminals in retinal explants led to a net

reduction of firing, with ~10% of RGCs showing a significant decrease of the baseline firing

while only ~1% showing an increase (Warwick et al., 2022). The retinal environment clearly

differs between ex vivo and in vivo conditions. One should then be cautious as findings from

ex vivo studies may not be directly applicable to the awake, more natural state.

What is the biological relevance of histaminergic modulation of the early visual system,

starting at the retina? We speculate that species-specific ethological factors play a role. For

example, some species may sleep with their eyes open (Rattenborg et al., 2000; Lyamin et

al., 2008, 2021; Pigarev et al., 2011), including mice (Yüzgeç et al., 2018; Karimi Abadchi et

al., 2020; Turner et al., 2020; Senzai and Scanziani, 2022). Furthermore, as nocturnal animals,

mice are less active during daytime. Facilitating visual responses at low histamine level may

then be ethologically beneficial for mice to respond faster to visual threats during daytime,

especially when mice are less active. In contrast, when animals are more active, other

mechanisms, such as arousal, may compensate for the histaminergic effects (Schröder et al.,

2020; Liang et al., 2020). Such arguments also hold for those species without eyelids,

including some amphibians, reptiles, and fish. To our knowledge, we are the first to report

histaminergic effects on LGN in awake mice. Recordings from LGN in cats and guinea pigs

(under the anesthetized condition or brain slices; McCormick and Williamson, 1991; Uhlrich,

et al., 2002) suggest that histamine leads to a slow depolarization of the relay cells, enhancing

the baseline activity but introducing a lag in their visual responses. However, substantial

differences exist in the early visual processing between awake and anesthetized conditions

(Durand et al., 2016; Boissonnet et al., 2023). Further studies are needed to address how

histamine affects visual processing and perception in awake animals in tissue- and species-

specific manners.

Growing evidence supports significant correlations between awake retinal responses

and various behavioral measures, such as pupil dynamics and locomotion (Schröder et al.,

2020; Liang et al., 2020; Boissonnet et al., 2023). Such correlations were also found in our

data sets; however, we demonstrated that an animal's pupil dynamics and locomotion

behavior have little to do with the observed histaminergic effects on RGC/LGN responses

(Figure 4 and Supplementary Figures 3 and 10). This indicates that mechanisms other than

histamine are involved in the retinal modulation by arousal and locomotion. The retina exploits

not just histamine but also many other neuromodulators, synthesized inside (Pourcho, 1996)

or outside the retina (Koves et al., 2016; Vereczki et al., 2024), such as the centrifugal

serotonergic system (Schütte, 1995; Repérant et al., 2000; Gastinger et al., 2006). It is a future

challenge to identify what behavior is controlled by histamine and vice versa, and characterize

the anatomical and physiological features of those HDC+ cells projecting to the retina to clarify

behavioral relevance of the histaminergic modulation of the early visual system.

Materials and Methods

No statistical method was used to predetermine the sample size. The significance level was

0.05 (with multiple comparison correction where appropriate) in all analyses unless noted

otherwise. All experiments were performed under the license 233/2017-PR and 220/2024-PR

from the Italian Ministry of Health, following protocols approved by the Institutional Animal

Care and Use Committee at European Molecular Biology Laboratory. The data analyses were

done in Python and Matlab.

Animals

Animals were housed on a 12h light-dark cycle, with ad libitum access to water and food. All

experiments were performed on female mice. A total of 22 wild-type (C57BL6/J;

RRID:IMSR_JAX:000664) and 12 hemizygous HDC-Cre (Hdctm1.1(icre)Wwis/J;

RRID:IMSR_JAX:021198) mice were used for pharmacological and chemogenetic

manipulations, respectively. Additional 9 hemizygous HDC-Cre mice were used for

anterograde tracing of HDC+ cells in TMN.

Surgical procedures

All surgical procedures were performed in animals from 4 to 19 weeks old (9.2 weeks median

age). Before the surgery, animals were injected with Carprofen (5 mg/kg) and then

anaesthetized with isoflurane (4% for induction, 1% for maintenance in O_2). Throughout the

surgery, the temperature of the animals was kept stable at 37°C using a heating pad

(Supertech Physiological). Ointment (VitA-Pos, Ursapharm) was applied on both eyes to

prevent them from drying during the surgery, and the animals were placed in a stereotaxic

frame (Stoelting 51625). A portion of the scalp was removed to expose the skull and the

periosteum was scraped away with a round scalpel to increase adherence of the dental

cement. For those mice belonging to the chemogenetic manipulation group, the skull was

aligned to level bregma and lambda on the same horizontal plane, and a 100 μm craniotomy

was performed using a motorized drill attached to the stereotaxic arm. Viral solution (250 nL;

AAV9::FLEX-PSAM Y115F,L141F:5HT3 HC; Vector Biolabs, VB773) was injected with an

automatic injector pump at 1 nL/sec, at [-2.6, ±0.8, -5.3] or [-1.7, ±0.5, -5.3] in [Anterior-

Posterior (AP), Medial-Lateral (ML), Dorsal-Ventral (DV)] coordinates for targeting TMN and

non-TMN areas of the hypothalamus, respectively. All animals were then implanted with a

custom-made titanium head plate, cemented on the skull using a mixture of cyanoacrylate

(Loctite 401, Henkel) and dental cement (Paladur, Kulzer). The head plate featured a hole (8

mm diameter), allowing for clear observation of both bregma and lambda. The skull surface

was then glazed with a thick layer of cyanoacrylate to support the skull with mechanical,

atmospheric and biological protection, while still allowing for visual identification of reference

points. After the surgery animals were allowed to recover on a heating pad and then housed

in individual cages. During the following seven days, the mice were administered with

analgesia (Carprofen; 50 mg/mL) diluted in the drinking water.

Anterograde tracing

For anterograde tracing of HDC+ cells, we injected recombinant AAVs carrying Cre-dependent

fluorescent markers (100 nL; rAAV9::CAG-FLEX-Synaptophysin-GFP or AAV2/1::CAG-

FLEX-axon-GFP) into the TMN of HDC-Cre mice (n=9) as described above. At least 10 days

after the injection, the animals were anesthetized (2.5% Avertin, 16 μL/g, intraperitoneal

injection) and perfused with paraformaldehyde (PFA; 4% in phosphate buffer solution). The

brain tissues including the optic chiasms and optic nerves were dissected and post-fixed

overnight in 4% PFA at 4°C. Coronal sections of the brain tissue (thickness, 80 μm) were then

examined under a laser scanning confocal microscope (Leica TCS SP5; Supplementary

Figure 2).

In vivo electrophysiology

After recovery from the surgery, animals were habituated to head fixation by placing them in

the experimental apparatus over the course of one week, twice a day for up to two hours. On

the day of the recording, we first placed the subject animal in the recording set up with its head

fixed, and determined the electrode penetration path to the target area (optic tract, [-1.34,

+1.87, +4.74], [-1.70, +1.87, +4.74], or [-1.82, +2.35, +4.07] in [AP, ML, DV] coordinates; LGN,

[-2.3, +2.3, +2.8]) using the robotic stereotaxic system (StereoDrive, NeuroStar). The animal

was then briefly anesthetized (with isoflurane for about 5 minutes) and a hole was drilled

around the electrode entry point on the skull. After the removal of the anesthesia, an acute

silicone probe (Buzsaki32L, Neuronexus; or P2, Cambridge Neurotech) coated with a

fluorescent dye (DiI stain, Invitrogen, D282) was lowered at 5 µm/s using the robotic arm until

visual responses were found in the target area. All recordings were done during the day time

(between 11am/ZT4 and 4pm/ZT9; 1pm/ZT6 ± 1 hour, mean ± standard deviation; 96% of the

paired recordings were conducted within 2 hours of each other).

After presenting a battery of visual stimuli, we briefly anesthetized the animal with

isoflurane and intraperitoneally injected 0.9% saline solution alone or that containing

chlorphenamine (H1 antagonist, 5 mg/kg; Frisch et al., 1997; Valle-Baurista et al. 2022),

ciproxifan (H3 antagonist, 12 mg/kg; Browman et al., 2004; Muindi et al., 2015), or PSEM (5

mg/kg; Magnus et al., 2011). About 20 minutes after the injection and removal of the

anesthesia, we presented the same battery of visual stimuli to the animal for the second

recording session. Here we did not target the H2 or H4 receptors because blocking the H2

receptors has little effect on retinal physiology ex vivo (Warwick et al., 2024), and the functional

role of the H4 receptors in the central nervous system remains unclear (Strakhova et al., 2009).

In our electrophysiology experiments, we could not keep the same cells long enough to test if

the effects of these manipulations were reversible or not.

After all the recording sessions, the electrode position was verified histologically (e.g.,

Figures 1B, LGN; Figure 3B, optic tract). After retracting the silicon probe, the mice were

anesthetized (2.5% Avertin, 16 μL/g, intraperitoneal injection) and perfused with 4% PFA. The

brain tissue was harvested and post-fixed overnight in 4% PFA at 4°C. Coronal sections of

the brain tissue (thickness, 100-150 μm) were then examined under a fluorescence

microscope (Leica, LMD7000 with N2.1 filter cube) to visualize the trace left by the DiI stain

on the probe. To examine the PSAM expression, staining of the slices was performed

(Supplementary Figure 2A,B): primary staining, 1-hour-incubation with an alpha-

bungarotoxin/biotin-XX conjugate (ThemoFisher, B1196) diluted with PBS with 0.1% Tween

(1/500) at room temperature; secondary staining, streptavidin and alexa-fluorophore, 1/500 in

PBS(T) 0.1% for 2 hours at room temperature.

Visual stimulation

Visual stimuli were presented as described before (Boissonnet et al., 2023). Briefly, visual

stimuli were projected onto a spherical screen (20cm in radius) painted with ultra-violet (UV)

reflective white paint, placed 20 cm from the mouse's left eye. A gamma-corrected digital light

processing device (DLP, Texas Instruments, DLPDLCR3010EVM-LC) was used as a light

source after the green and the red light-emitting diodes (LEDs) were replaced with UV (365nm,

LZ1-00UV00, LED Engine) and infrared (IR; 950nm, SFH 4725S, Osram) LEDs, respectively.

The IR light was used as a synchronization signal and recorded with a photodiode (PDA100A2,

Thorlabs). Stimulation was conducted at 60 frames per second (FPS), and covered 73° in

azimuth and 44° in altitude from the mouse eye position. The maximum light intensity at the

eye position was 31 mW/m^2 (15.4 mW/m^2 for UV LED and 15.9 mW/m^2 for blue LED;

measured with S121C sensor, Thorlabs), leading to mesopic to photopic conditions. We

presented the following stimuli using QDSpy: a randomly flickering full-field stimulus (5

minutes), consisting of a random sequence of black and white frames at 60 FPS; a black-and-

white binary dense noise stimulus (15 minutes), consisting of a 32-by-18 pixels checkerboard,

where each pixel followed an independent random sequence at 60 FPS but the overall

luminance of the frame was kept constant at the mean value; a moving grating stimulus in

eight different directions (spatial frequencies of square waves, 3° or 20°; moving speed, 7.5°/s

or 15°/s); full-field contrast-inverting stimuli in an "OFF-ON-OFF" sequence at maximum

contrast (2 s each), followed by a sinusoid (1.5 Hz) with a linearly increasing amplitude from

0% to 100% contrast over 10 s (10 trials) or a sequence of sinusoids with different temporal

frequencies (1.875, 3.75 and 7.5 Hz, each for 2 s; 15 and 30 Hz for 1 s) at maximum contrast

(10 trials), with an inter-trial interval of 1 s (gray screen).

Electrophysiology data analysis

Here we adapted previously established methods of spiking sorting and optic tract data

analysis (Boissonnet et al., 2023). In brief, we first concatenated the raw data from the same

animal before and after treatment into a single binary file, and used Kilosort 2.0 to sort spikes

with a set of default parameters, except for the spike detection threshold to be 6 during

optimization. Single-units were then identified by clustering in principal component space

using Phy for visualization and manual data curation. Only those units that maintained the

average spike waveforms and autocorrelograms with a minimal refractory period of 1 ms were

kept for subsequent analyses (Supplementary Figure 1). Specifically, we selected RGCs/LGN

cells as those with robust visual responses: i.e., $\text{SNR} = \text{var}[\langle r(t) \rangle]_t / \langle \text{var}[r(t)]_t \rangle > 0.15$, where

$r(t)$ is the response during the "ON-OFF" part of the OFF-ON-OFF stimulus sequence, $\langle \cdot \rangle$

indicates the mean over trials, and $\text{var}[\cdot]_t$ the variance over time t (bin size, 1/60 ms),

respectively; and the estimated temporal filter should have $p < 10^{-10}$ in at least one time bin

within 200 ms from the spike onset (see below for details). Single-units with little or no visual

responses were excluded from the analysis as non-target cells, such as the axons from the

parabigeminal nucleus in the optic tract (Tokuoka et al., 2020; Reinhard et al., 2019). In total,

we obtained 337 RGCs from 20 animals and 557 cells in LGN from 14 mice. These RGCs

typically had a triphasic spike waveform as expected for axonal signals (Barry, 2015).

To characterize the peak latency and firing rate of the responses to full-field contrast-

inverting stimuli (Figure 1 and Supplementary Figure 6), we first identified the cell's response

preference to stimulus polarity using an ON-OFF index defined as $(r_{\mathrm{ON}} - r_{\mathrm{OFF}})/(r_{\mathrm{ON}} + r_{\mathrm{OFF}})$,

where r_{ON} and r_{OFF} are the mean firing rate during the ON and the second OFF periods of the

OFF-ON-OFF stimulus sequence. We then computed the peri-stimulus time histogram (bin

size, 1 ms), smoothed it with a Gaussian filter (kernel width, 6 standard deviation), and

identified the first peak upon stimulus onset or offset for those cells with positive or negative

ON-OFF index values, respectively. We then performed pairwise comparisons on the peak

latencies and firing rates before and after each treatment (e.g., Figure 1F-J). To compare the

changes in the detected peak latency and firing rates across different conditions (e.g., Figure

1K,L), we performed Kruskal-Wallis test on a modulation index defined as

$(a_{\mathrm{after}} - a_{\mathrm{before}})/(a_{\mathrm{after}} + a_{\mathrm{before}})$, where a is either the detected peak latency or peak firing rate

before and after treatment.

For systematically characterizing the visual response properties, we used stimulus

ensemble statistical techniques (reverse correlation methods; 500 ms window; Δt = 1/60 s bin

size) to calculate the linear filter and static nonlinear gain function of the recorded cells in

response to white-noise stimuli as described before (Figure 2B; Boissonnet et al., 2023).

Briefly, we first obtained the linear filter of each cell by calculating a spike-triggered average

(STA) of the stimulus with ±1 being "white" and "black," respectively, under a given behavioral

state (e.g., when the animal stayed stationary below 2 cm/s). As a quality measure, p-value

was computed for each time bin against a null hypothesis that the STA follows a normal

distribution with a mean of zero and a variance of $1/C$, where C is the total number of spikes.

As a measure of the cell's response kinetics, we then estimated the peak latency by fitting a

difference-of-Gaussian curve to the linear filter; and the spectral peak frequency by the Fourier

analysis on the linear filter. The ON-OFF polarity index of a temporal filter was defined as the

difference of its peak and valley, divided by the sum of the two (Figure 4G and Supplementary

Figure 10D). Spatial response properties were examined for STAs from "checkerboard"

stimulation, using the time window where the eye position remained within 1 inter-quartile

range in both horizontal and vertical coordinates for at least 2 seconds (Supplementary

Figures 5C and 8A) to minimize the effects of eye movements while retaining enough data for

RF estimation. In particular, we fitted a two-dimensional Gaussian envelope to the spatial filter

at the peak latency (e.g., Supplementary Figure 5D), and the receptive field (RF) size was

estimated as twice the mean standard deviation (SD) of the long and short axes. Modulation

index was used to characterize the change in a response feature before and after treatment,

such as the peak latency and frequency, the mean firing rate, and the receptive field size

(Figures 2K-N, 3K-N). Here we focused only on those data when the mouse stayed stationary

(<2 cm/s) to minimize the side effects of locomotion behaviour (Erisken et al., 2014; Schröder

et al., 2020; Supplementary Figure 3).

The linear temporal filters from full-field white-noise stimuli were used for broadly

classifying the response types of RGCs and LGN cells, respectively (Supplementary Figures

4A,B and 7A,B). Specifically, we first used t-Distributed Stochastic Neighbor Embedding (t-

SNE) to map the STAs onto two-dimensional space, and then used K-means++ algorithm for

heuristically categorizing the responses into the following six types: fast ON, ON, slow ON,

fast OFF, OFF, and slow OFF. Our data sets were not large enough to physiologically classify

and identify all the cell types reported thus far (Baden et al., 2016; Rosón et al., 2019).

Direction-selectivity (DS) and orientation-selectivity (OS) indices were calculated by projecting

the responses to the moving grating stimuli in eight different directions onto a complex

exponential: $\mathrm{DS/OS\ index} = \left\| \sum_k e^{-i\alpha\omega_k} / \sum_k r_k \right\|$, where ω_k and r_k are the angle of the k-th

direction and the cell's corresponding responses, respectively; and $\alpha = 1$ and 2 for the DS

and OS indices, respectively (Supplementary Figure 9).

Static nonlinear gain function of each cell (bin size, 0.1) was computed as: P(response

| stimulus) = N(stimulus | response) / N(stimulus) / Δt, where N(stimulus | response) and

N(stimulus) are the distributions of spike-triggered stimulus ensembles projected onto the L2-

normalized STA and the entire stimulus ensembles, respectively. The ratio of P(response |

stimulus) before and after the treatment was used as a measure of response gain modulation

for each cell.

Behavioral data analysis

For all experiments, the animal's left eye – i.e., the side presented with visual stimuli – was

recorded using an IR camera (The Imaging Source, DMK23UV024) to monitor its motion and

pupil dynamics at 30-60Hz. Pupil detection was done using a Mask-Region based

Convolutional Neural Network (Mask-RCNN), trained as described previously (Boissonnet et

al., 2023). A two-dimensional ellipse was fit to the pupil, and the large axis diameter was used

as a measure of the pupil size, normalized by the median value before treatment for each

animal (Figure 4A). To examine the pupil size effects on the response dynamics, a threshold

at 33 and 66 percentiles was used to define constricted, neutral, and dilated pupil period,

respectively. To examine the effects of the pupil dynamics, we took the time derivative of the

pupil size dynamics, and set a threshold at ±0.01 point per second to define constricting, stable,

and dilating pupil periods, respectively.

We also monitored the running speed of an animal based on the turning speed of the

custom-made rotary treadmill throughout the recordings. We set a threshold at 2 cm/s to detect

running behavior (Supplementary Figure 3A). For quantification, we measured the fraction of

the running period (Supplementary Figure 3B), and the median running speed when mice

moved at >2 cm/s before and after each treatment (Supplementary Figure 3C)..

Model analysis

To explore how different aspects of neuronal intrinsic properties and output responses are

interrelated between each other, we simulated a model neuron's responses to white-noise

stimuli using different model parameter values (Figure 5A), and characterized these simulated

responses using a linear-nonlinear cascade model as we did for the experimental data.

Specifically, we employed an integrate-and-fire neuron model with Poisson spiking and

refractoriness. A model neuron's discrete linear filter (of length $n_T = 25$) was given by a

trigonometric function with logarithmic scale in time: i.e., $L(i) = \sin[2\pi(10^{t_i} - 1)/(10^{s_t} - 1)]$,

where $s_t = 1.5$ is a temporal spacing parameter and $t_i = i \cdot s_t/(n_T - 1)$ for $i = 0, \dots, n_T - 1$.

The spike-generating function $V(i)$ was then modelled by first projecting the white-noise

Gaussian stimulus $S(i)$ (of length 1,000,000; normalized to have a maximum value of 1) onto

$L(i)$, followed by the addition of baseline B and the multiplication by gain G: i.e., $V(i) = G \cdot$

$\left(\sum_j L(j) \cdot S(i - j) + B\right)$. For each simulation, we chose the value of B from -0.45 to $+0.45$ in

steps of 0.15; and that of G from 1.2^g with $g = -3, \dots, +3$. The spiking response was then

generated by thresholding $V(i)$ (at a random threshold for every time point derived from a

uniform distribution from 0 to 1) and refractoriness (given by an exponential filter: $-\exp[-i/s_h]$

for $i = 0, \dots, 5$ with $s_h = 1$). The output spike trains were then subject to the reverse-correlation

analysis as described above to characterize the apparent linear filter and static nonlinearity of

the model neuron, much as we did for the experimentally recorded RGCs and LGN cells.

Specifically, we focused on the peak latency and frequency of the estimated linear filters and

the slope of the estimated static nonlinear gain function, and examined how these features

depended on the baseline B and the intrinsic gain G of the model neuron to provide a minimal

phenomenological explanation to our observations (Figure 5B-E).

Acknowledgments

This work was supported by research grants from EMBL (H.A.). The EMBL Advanced Light

Microscopy Facility are acknowledged for support in histological image acquisition and

analyses, and EMBL IT Support for provision of computer and data storage servers. We thank

Dmitry Molotkov, Tom Boissonnet, Vanesa Pelcastre, Sahana Rao, Shabnam Chandel, Lily

Knowles for their help in experiments, and all the Asari lab members for many useful

discussions.

References

Akimov, N.P., Marshak, D.W., Frishman, L.J., Glickman, R.D., Yusupov, R.G. (2010). Histamine Reduces Flash Sensitivity of ON Ganglion Cells in the Primate Retina. Invest Ophthalmol Vis Sci. 51(7), 3825–3834.

Baden, T., Berens, P., Franke, K., Rosón, M. R., Bethge, M., Euler, T. (2016). The functional diversity of retinal ganglion cells in the mouse. Nature, 529(7586), 345-350.

Barry, J.M. (2015). Axonal activity in vivo: technical considerations and implications for the exploration of neural circuits in freely moving animals. Front. Neurosci. 9, 153.

Bloomfield, S.A., Völgyi, B. (2009). The diverse functional roles and regulation of neuronal gap junctions in the retina. Nat Rev Neurosci. 10(7), 495–506.

Boissonnet, T., Tripodi, M., Asari, H. (2023). Awake responses suggest inefficient dense coding in the mouse retina. Elife. 12, e78005.

Browman, K.E., Komater, V.A., Curzon, P., Rueter, L.E., Hancock, A.A., Decker, M.W., Fox, G.B. (2004). Enhancement of prepulse inhibition of startle in mice by the H3 receptor antagonists thioperamide and ciproxifan. Behav Brain Res 153(1), 69-76.

Brown, R.E., Sergeeva, O.A., Eriksson, K.S., Haas, H.L. (2002). Convergent Excitation of Dorsal Raphe Serotonin Neurons by Multiple Arousal Systems (Orexin/Hypocretin, Histamine and Noradrenaline). J. Neurosci. 22, 8850–8859.

Dipoppa, M., Ranson, A., Krumin, M., Pachitariu, M., Carandini, M., Harris, K.D. (2018). Vision and Locomotion Shape the Interactions between Neuron Types in Mouse Visual Cortex. Neuron 98(3), 602–615.e8.

Durand, S., Iyer, R., Mizuseki, K., de Vries, S., Mihalas, S., Reid, R. C. (2016). A Comparison of Visual Response Properties in the Lateral Geniculate Nucleus and Primary Visual Cortex of Awake and Anesthetized Mice. J Neurosci, 36(48), 12133–12156.

Erisken, S., Vaiceliunaite, A., Jurjut, O., Fiorini, M., Katzner, S., Busse, L. (2014). Effects
of locomotion extend throughout the mouse early visual system. Curr Biol, 24(24), 2899–
2907.

Esbenshade, T.A., Krueger, K.M., Miller, T.M., Kang, C.H., Denny, L.I., Witte, D.G., Yao,
B.B., Fox, G.B., Faghih, R., Bennani, Y.L., Williams, M., Hancock, A.A. (2003). Two novel
and selective nonimidazole histamine H3 receptor antagonists A-304121 and A-317920: I.
In vitro pharmacological effects. J Pharmacol Exp Ther 305(3), 887–896.

Esposti, F., Johnston, J., Rosa, J. M., Leung, K.-M., Lagnado, L. (2013). Olfactory
Stimulation Selectively Modulates the OFF Pathway in the Retina of Zebrafish. Neuron,
79(1), 97-110.

Fang, S.Y., Druce, H.M., Baraniuk, J.N. (1998). Anticholinergic properties of
brompheniramine, chlorpheniramine, and atropine in human nasal mucosa in vitro. Am J
Rhinol 12(2), 131–133.

Frisch, C., Hasenöhrl, R.U., Huston, J.P. (1997). The histamine H1-antagonist
chlorpheniramine facilitates learning in aged rats. Neurosci Lett 229(2), 89–92.

Gastinger, M.J., O'Brien, J.J., Larsen, J.N.B., Marshak, D.W. (1999). Histamine
Immunoreactive Axons in the Macaque Retina. Invest. Ophthalmol. Vis. Sci. 40, 487–495.

Gastinger, M. J., Tian, N., Horvath, T., Marshak, D. W. (2006). Retinopetal axons in
mammals: Emphasis on histamine and serotonin. Curr. Eye Res. 31, 655–667.

Goel, M., Mangel, S.C. (2021). Dopamine-Mediated Circadian and Light/Dark-Adaptive
Modulation of Chemical and Electrical Synapses in the Outer Retina. Front Cell Neurosci
15, 647541.

Greferath, U., Kambourakis, M., Barth, C., Fletcher, E.L., Murphy, M. (2009).
Characterization of histamine projections and their potential cellular targets in the mouse
retina. Neuroscience 158, 932–944.

Grozdanic, S., Betts, D.M., Allbaugh, R.A., Sakaguchi, D.S., Kwon, Y.H., Kardon, R.H.,
Sonea, I.M. (2003). Characterization of the pupil light reflex, electroretinogram and
tonometric parameters in healthy mouse eyes. Curr Eye Res 26(6), 371–378.

Haas, H., Panula, P. (2003). The role of histamine and the tuberomamillary nucleus in the
nervous system. Nat. Rev. Neurosci. 4, 121–130.

Haas, H.L., Sergeeva, O.A., Selbach, O. (2008). Histamine in the nervous system. Physiol
Rev 88(3), 1183–1241.

Hagenow, S., Stasiak, A., Ramsay, R.R., Stark, H. (2017). Ciproxifan, a histamine H3
receptor antagonist, reversibly inhibits monoamine oxidase A and B. Sci Rep 7, 40541.

Horio, K., Ohkuma, M., Miyachi, E.-I. (2018). The effect of histamine on inward and
outward currents in mouse retinal amacrine cells. Cell. Mol. Neurobiol. 38, 757–767.

Jacoby, J., Nath, A., Jessen, Z.F., Schwartz, G.W. (2018). A Self-Regulating Gap Junction
Network of Amacrine Cells Controls Nitric Oxide Release in the Retina. Neuron 100, 1149-
1162.e5.

John, J., Wu, M.-F., Boehmer, L.N., Siegel, J.M. (2004). Cataplexy-Active Neurons in the
Hypothalamus: Implications for the Role of Histamine in Sleep and Waking Behavior.
Neuron 42, 619–634.

Kamei, C., Okumura, Y., Tasaka, K. (1993). Influence of histamine depletion on learning
and memory recollection in rats. Psychopharmacology (Berl.) 111, 376–382.

Karimi Abadchi, J., Nazari-Ahangarkolaee, M., Gattas, S., Bermudez-Contreras, E.,
Luczak, A., McNaughton, B.L., Mohajerani, M.H. (2020) Spatiotemporal patterns of
neocortical activity around hippocampal sharp-wave ripples. Elife 9, e51972.

Khateb, A., Fort, P., Pegna, A., Jones, B.E., Mühlethaler, M. (1995). Cholinergic nucleus
basalis neurons are excited by histamine in vitro. Neuroscience 69, 495–506.

Khouma, A., Moeini, M.M., Plamondon, J., Richard, D., Caron, A., Michael, N.J. (2023).

Histaminergic regulation of food intake. Front. Endocrinol. 14.

Klapdor, K., Hasenöhrl, R.U., Huston, J.P. (1994). Facilitation of learning in adult and aged

rats following bilateral lesions of the tuberomammillary nucleus region. Behav. Brain Res.

61, 113–116.

Koves, K., Csaki, A., Vereczki, V. (2016). Recent Research on the Centrifugal Visual

System in Mammalian Species. Anat Physiol 6(3), 216.

Labandeira-Garcia, J.L., Guerra-Seijas, M.J., Gonzalez, F., Perez, R., Acuña, C., (1990).

Location of neurons projecting to the retina in mammals. Neurosci. Res. 8, 291–302.

Lee, S.-H., Dan, Y. (2012). Neuromodulation of Brain States. Neuron 76, 209–222.

Liang, L., Fratzl, A., Reggiani, J.D.S., El Mansour, O., Chen, C., Andermann, M.L. (2020).

Retinal Inputs to the Thalamus Are Selectively Gated by Arousal. Curr. Biol. CB 30, 3923–

3934.e9.

Lin, W., Xu, L., Zheng, Y., An, S., Zhao, M., Hu, W., Li, M., Dong, H., Li, A., Li, Y., Gong,

H., Pan, G., Wang, Y., Luo, Q., Chen, Z. (2023) Whole-brain mapping of histaminergic

projections in mouse brain. Proc. Natl. Acad. Sci. U. S. A. 120, e2216231120.

Liu, Y., Rodenkirch, C., Moskowitz, N., Schriver, B., Wang, Q. (2017). Dynamic

Lateralization of Pupil Dilation Evoked by Locus Coeruleus Activation Results from

Sympathetic, Not Parasympathetic, Contributions. Cell Rep, 20(13), 3099–3112.

Lyamin, O.I., Manger, P.R., Ridgway, S.H., Mukhametov, L.M., Siegel, J.M. (2008).

Cetacean sleep: an unusual form of mammalian sleep. Neurosci Biobehav Rev 32(8),

1451–1484.

Lyamin, O.I., Siegel, J.M., Evsigneev, R.V., Nazarenko, E.A., Rozhnov, V.V. (2021).

Characteristics of Sleep-Wakefulness Cycle and Circadian Activity in the Lesser Mouse-

Deer (Tragulus kanchil). Dokl Biol Sci 500(1), 153–158.

Magnus, C.J., Lee, P.H., Bonaventura, J., Zemla, R., Gomez, J.L., Ramirez, M.H., Hu, X.,

Galvan, A., Basu, J., Michaelides, M., Sternson, S.M. (2019). Ultrapotent chemogenetics

for research and potential clinical applications. Science. 364(6436), eaav5282

McCormick, D.A., Nestvogel, D.B., He, B.J., (2020). Neuromodulation of brain state and

behaviour. Ann. Rev. Neurosci. 43(1), 391–415.

McCormick, D.A., Williamson, A. (1991). Modulation of neuronal firing mode in cat and

guinea pig LGNd by histamine: possible cellular mechanisms of histaminergic control of

arousal. J. Neurosci. 11 (10), 3188–3199.

Mochizuki, T., Yamatodani, A., Okakura, K., Horii, A., Inagaki, N., Wada, H. (1992).

Circadian rhythm of histamine release from the hypothalamus of freely moving rats.

Physiol. Behav. 51, 391–394.

Muindi, F., Colas, D., Ikeme, J., Ruby, N.F., Heller, H.C. (2015). Loss of Melanopsin

Photoreception and Antagonism of the Histamine H3 Receptor by Ciproxifan Inhibit Light-

Induced Sleep in Mice. PLoS One 10(6), e0128175.

Niell, C.M., Stryker, M.P. (2010). Modulation of visual responses by behavioral state in

mouse visual cortex. Neuron 65(4), 472–479.

Nowak, J.Z. (1993). Histamine in the retina and some other components of the visual

system. Prog Retin Res 12, 41–74.

Orsetti, M., Ghi, P., Carlo, G.D. (2001). Histamine H3-receptor antagonism improves

memory retention and reverses the cognitive deficit induced by scopolamine in a two-trial

place recognition task. Behav. Brain Res. 124, 235–242.

Parmentier, R., Ohtsu, H., Djebbara-Hannas, Z., Valatx, J.-L., Watanabe, T., Lin, J.-S.

(2002). Anatomical, Physiological, and Pharmacological Characteristics of Histidine

Decarboxylase Knock-Out Mice: Evidence for the Role of Brain Histamine in Behavioral

and Sleep–Wake Control. J. Neurosci. 22, 7695–7711.

Pigarev, I.N., Fedorov, G.O., Levichkina, E.V., Marimon, J.M., Pigareva, M.L., Almirall, H.
(2011). Visually triggered K-complexes: a study in New Zealand rabbits. Exp Brain Res
210(1), 131–142.

Pourcho, R.G. (1996). Neurotransmitters in the retina. Curr Eye Res 15(7), 797–803.

Rattenborg, N.C., Amlaner, C.J., Lima, S.L. (2000). Behavioral, neurophysiological and
evolutionary perspectives on unihemispheric sleep. Neurosci Biobehav Rev 24(8), 817–
842.

Reggiani, J.D.S., Jiang, Q., Barbini, M., Lutas, A., Liang, L., Fernando, J., Deng, F., Wan,
J., Li, Y., Chen, C., Andermann, M.L. (2023). Brainstem serotonin neurons selectively gate
retinal information flow to thalamus. Neuron 111, 711–726.e11.

Reinhard, K., Li, C., Do, W., Burke, E.G., Heynderickx, S., Farrow, K. (2019). A projection
specific logic to sampling visual inputs in mouse superior colliculus. eLife 8, e50697.

Repérant, J., Médina, M., Ward, R., Miceli, D., Kenigfest, N.B., Rio, J.P., Vesselkin, N.P.
(2007). The evolution of the centrifugal visual system of vertebrates. A cladistic analysis
and new hypotheses. Brain Res. Rev. 53, 161–197.

Repérant, J., Araneda, S., Miceli, D., Medina, M., Rio, J. P. (2000). Serotonergic
retinopetal projections from the dorsal raphe nucleus in the mouse demonstrated by
combined [3H] 5-HT retrograde tracing and immunolabeling of endogenous 5-HT. Brain
Res. 878, 213–217.

Rosón, M.R., Bauer, Y., Kotkat, A.H., Berens, P., Euler, T., Busse, L. (2019). Mouse dLGN
Receives Functional Input from a Diverse Population of Retinal Ganglion Cells with Limited
Convergence. Neuron 102(2), 462-476.e8.

Rubio, S., Begega, A., Santin, L.J., Arias, J.L. (2002). Improvement of spatial memory by
(R)-α-methylhistamine, a histamine H3-receptor agonist, on the Morris water-maze in rat.
Behav. Brain Res. 129, 77–82.

Savier, E.L., Chen, H., Cang, J. (2019). Effects of Locomotion on Visual Responses in the

Mouse Superior Colliculus. J Neurosci 39(47), 9360–9368.

Scammell, T.E., Jackson, A.C., Franks, N.P., Wisden, W., Dauvilliers, Y. (2018).

Histamine: neural circuits and new medications. Sleep 42, zsy183.

Schröder, S., Steinmetz, N.A., Krumin, M., Pachitariu, M., Rizzi, M., Lagnado, L., Harris,

K.D., Carandini, M. (2020). Arousal Modulates Retinal Output. Neuron 107, 487–495.e9.

Schütte, M. (1995). Centrifugal innervation of the rat retina. Vis. Neurosci. 12, 1083–1092.

Senzai, Y., Scanziani, M. (2022) A cognitive process occurring during sleep is revealed by

rapid eye movements. Science 377, 999–1004.

Strakhova, M.I., Nikkel, A.L., Manelli, A.M., Hsieh, G.C., Esbenshade, T.A., Brioni, J.D.,

Bitner, R.S. (2009) Localization of histamine H4 receptors in the central nervous system

of human and rat. Brain Res, 1250, 41-48.

Sumitomo, I., Ide, K., Iwama, K., Arikuni, T. (1969). Conduction velocity of optic nerve

fibers innervating lateral geniculate body and superior colliculus in the rat. Exp Neurol

25(3), 378–392.

Takahashi, K., Lin, J.-S., Sakai, K. (2006). Neuronal activity of histaminergic

tuberomammillary neurons during wake-sleep states in the mouse. J. Neurosci. Off. J. Soc.

Neurosci. 26, 10292–10298.

Thakkar, M.M. (2011). Histamine in the regulation of wakefulness. Sleep Med Rev. 15(1),

65-74.

Tokuoka, K., Kasai, M., Kobayashi, K., Isa, T. (2020). Anatomical and electrophysiological

analysis of cholinergic inputs from the parabigeminal nucleus to the superficial superior

colliculus. J Neurophysiol. 124(6), 1968–1985.

Turner, K.L., Gheres, K.W., Proctor, E.A., Drew, P.J. (2020). Neurovascular coupling and

bilateral connectivity during NREM and REM sleep. Elife 9, e62071.

Uhlrich, D. J., Manning, K. A., Xue, J.-T.. (2002). Effects of activation of the histaminergic

tuberomammillary nucleus on visual responses of neurons in the dorsal lateral geniculate

nucleus. J. Neurosci. 22, 1098–1107.

Valdés, J.L., Sánchez, C., Riveros, M.E., Blandina, P., Contreras, M., Farías, P., Torrealba,

F. (2010). The histaminergic tuberomammillary nucleus is critical for motivated arousal.

Eur. J. Neurosci. 31, 2073–2085.

Valle-Bautista, R., Márquez-Valadez, B., Herrera-López, G., Griego, E., Galván, E.J., Díaz,

N.-F.,   Arias-Montaño, J.-A., Molina-Hernández, A. (2022). Long-Term Functional and

Cytoarchitectonic Effects of the Systemic Administration of the Histamine H1 Receptor

Antagonist/Inverse Agonist Chlorpheniramine During Gestation in the Rat Offspring

Primary Motor Cortex. Front Neurosci 15, 740282.

Vereczki, V., Köves, K., Csáki, A. (2024). Current state of knowledge on the centrifugal

visual   system   (including   the   pinealo-to-retinal   connection)   in   mammals   and   its

hypothesized role in circadian rhythms. Explor Neurosci, 3, 51–64.

Vila, A., Satoh, H., Rangel, C., Mills, S.L., Hoshi, H., O'Brien, J., Marshak, D.R., Macleish,

P.R., Marshak, D.W. Histamine receptors of cones and horizontal cells in Old World

monkey retinas. J Comp Neurol. 520(3), 528–543.

Wang, Y.-Q., Li, R., Wu, X., Zhu, F., Takata, Y., Zhang, Z., Zhang, M.-Q., Li, S.-Q., Qu,

W.-M. (2015). Fasting activated histaminergic neurons and enhanced arousal effect of

caffeine in mice. Pharmacol. Biochem. Behav. 133, 164–173.

Warwick, R.A., Riccitelli, S., Heukamp, A.S., Yaakov, H., Ankri, L., Mayzel, J., Gilead, N.,

Parness-Yossifon, R., Rivlin-Etzion, M. (2022). Top-down modulation of the retinal code

via histaminergic neurons of the hypothalamus. bioRxiv, 2022.04.26.489509.

Warwick, R.A., Riccitelli, S., Heukamp, A.S., Yaakov, H., Swain, B.P., Ankri, L., Mayzel,
J., Gilead, N., Parness-Yossifon, R., Di Marco, S., Rivlin-Etzion, M. (2024). Top-down
modulation of the retinal code via histaminergic neurons of the hypothalamus. Sci Adv.
10(35):eadk4062.

White, J.M., Rumbold, G.R. (1988). Behavioural effects of histamine and its antagonists:
a review. Psychopharmacology (Berl) 95(1), 1-14.

Wilson, M., Lindstrom, S.H. (2011). What the bird's brain tells the bird's eye: the function
of descending input to the avian retina. Vis Neurosci, 28(4), 337–350.

Yu, Y., Satoh, H., Wu, S.M., Marshak, D.W. (2009). Histamine enhances voltage-gated
potassium currents of ON bipolar cells in macaque retina. Invest Ophthalmol Vis Sci, 50(2),
959–965.

Yu, Y., Satoh, H., Vila, A., Wu, S.M., Marshak, D.W. (2011). Effects of Histamine on Light
Responses of Amacrine Cells in Tiger Salamander Retina. Neurochem Res, 36, 645–654.

Yu, X., Ye, Z., Houston, C.M., Zecharia, A.Y., Ma, Y., Zhang, Z., Uygun, D.S., Parker, S.,
Vyssotski, A.L., Yustos, R., Franks, N.P., Brickley, S.G., Wisden, W. (2015). Wakefulness
Is Governed by GABA and Histamine Cotransmission. Neuron, 87(1), 164–178.

Yüzgeç, Ö., Prsa, M., Zimmermann, R., Huber, D. (2018). Pupil size coupling to cortical
states protects the stability of deep sleep via parasympathetic modulation. Curr Biol 28,
392–400.e3.