Genetic variation in phenology of wild *Arabidopsis thaliana* plants

Victoria L. DeLeo[1], David L. Des Marais[2], Claire M. Lorts[1], Thomas E. Juenger[3], Jesse R.

Lasky[1,4]

[1]Department of Biology
Pennsylvania State University

[2]Department of Civil and Environmental Engineering
Massachusetts Institute of Technology

[3]Department of Integrative Biology
University of Texas at Austin

[4]Author for correspondence. Email: lasky@psu.edu

16
17
18
19

Short running title: *Phenology of wild Arabidopsis*

Acknowledgements

We thank the herbaria that allowed sampling of seed from specimens for generation of natural
inbred lines used in experiments: Real Jardín Botánico de Madrid, Oslo Natural History
Museum, and Komarov Botanical Institute. We are grateful for assistance with experiments from
P. Patel, C. Yim, J. Kizer, T. Xia, V. Meagher, C. Lorts, K. Turner, and E. Bellis and for helpful
discussions with L. Burghardt. Funding was provided by NIH award R35 GM138300 to JRL.

Author Contribution

JRL conceived of the project with TEJ, DLDM and CML assisted in experiments, VLD
conducted experiments, and analyzed data, VLD and JRL led writing, all authors contributed to
the interpretation and writing of the manuscript.

Data Accessibility Statement

Original data will be included as supplemental tables. Prior studies from which phenology data
was sourced will be listed in a supplemental table.

Conflict of Interest Statement

The authors declare no conflicts of interest.

Abstract

Phenology and the timing of development are often under selection. However, the relative

contributions of genotype, environment, and prior developmental transitions to variance in the

phenology of wild plants is largely unknown. Individual components of phenology (e.g.,

germination) might be loosely related with the timing of maturation due to variation in prior

developmental transitions. Given widespread evidence that genetic variation in life history is

adaptive, we investigated to what degree experimentally measured genetic variation in

Arabidopsis phenology predicts phenology of plants in the wild. As a proxy of phenology, we

obtained collection dates from nature of 227 naturally inbred *Arabidopsis thaliana* accessions

from across Eurasia. We compared this phenology in nature with experimental data on the

descendant inbred lines that we synthesized from two new and 155 published controlled

experiments. We tested whether the genetic variation in flowering and germination timing from

experiments predicted the phenology of the same lines in nature. We found that genetic variation

in phenology from controlled experiments significantly predicts day of collection from wild

individuals, as a proxy for date of flowering, across Eurasia. However, local variation in

collection dates within a region was not explained by genetic variance in phenology in

experiments, suggesting high plasticity across small-scale environmental gradients or complex

interactions between the timing of different developmental transitions. This apparent low

heritability in phenology within regions contrasts with the well-known genetic clines in

phenology, suggesting that strong selection or many generations are required for adaptation and

the emergence of phenological clines.

Introduction

The predictability of trait variation from genetic variation, *i.e.* trait heritability, is a key quantity determining the rate of trait evolution and the ability of populations to adapt to changes in selection (Lush 1937; Gomulkiewicz and Houle 2009). Plant biologists have long relied on common garden experiments to estimate breeding values and heritability for a range of traits (Mazer and LeBuhn 1999). However, natural populations often occur in highly heterogeneous environments to which organismal traits may respond plastically. As a result, heritability in natural populations may be much lower than those in common gardens. Furthermore, even field common gardens might fail to accurately recreate natural selective pressures if the direction and magnitude of selection varies in time (Karrenberg & Widmer 2008). Alternatively, heritability can be estimated using pedigrees or genomic data from wild individuals (Gienapp et al. 2017), but these models rely on assumptions of certain genetic architectures (Mitchell-Olds and Rutledge 1986) and the resulting heritability estimates are not often compared with common garden estimates of trait variation (but see (Weigensberg and Roff 1996)).

Phenology, or the timing of the developmental transitions between an organism's life stages, directly and indirectly influences plant fitness and selection by determining the traits that are expressed at any point throughout the year (Donohue 2005). In seasonally variable environments, traits such as flowering time and dormancy can ameliorate harsh abiotic conditions by timing dormant periods to coincide with unfavorable seasons (e.g. drought escape Ludlow 1989; Lawrence-Paul and Lasky 2024). Flowering, vegetative phase change, and germination timing often show geographic clines, suggesting their importance in adaptation to local environments (Baughman et al. 2019; Martínez-Berdeja et al. 2020; Lawrence-Paul et al. 2023, 2025). Common garden studies often find that genetic variation in individual components of phenology are under selection and have further demonstrated variable strength and direction of selection across field sites (Hall and Willis 2006; Fournier-Level et al. 2013; Ågren et al. 2017; Gamba et al. 2024b). However, the duration of individual developmental transitions (e.g. flowering time) might be loosely related to phenology in nature due to variation in prior developmental transitions (e.g. germination) (Burghardt et al. 2015). While phenology is determined by both genetic factors and plastic responses to environment (Amasino 2004; Andrés and Coupland 2012; Auge et al. 2018), the relative importance of these two sources of

phenotypic variance in the timing of individual developmental transitions determining phenology in nature is unclear, even in model systems.

Much of our understanding of the genetic basis of plant phenology comes from studies in the model *Arabidopsis thaliana* (hereafter Arabidopsis). Common garden trials in both lab and field settings have uncovered genetic loci contributing to much of the observed variation in dormancy and flowering time (Juenger et al., 2005; Brachi et al., 2010; Fournier-Level et al., 2013; Ågren et al., 2017). These traits are also plastic and show genotype by environment interactions (Juenger et al. 2005; Zhou et al. 2005; Wilczek et al. 2009; Lorts and Lasky 2020): germination responds to temperature, photoperiod, moisture, and nutrient availability (Huang et al. 2010, 2018; Penfield and Springthorpe 2012; Footitt et al. 2013; Kenney et al. 2014), while flowering responds to multiple cues such as temperature and daylength (Thomas and Vince-Prue 1997; Lempe et al. 2005; Balasubramanian et al. 2006).

The broad phenological variation described in controlled environment experiments is observed in wild populations as well (Ratcliffe 1976; Simpson and Dean 2002; DeLeo et al. 2020). Arabidopsis can exhibit the life history of a winter annual, germinating in the fall, overwintering as a rosette, and flowering in the spring (Figure 1). However, Arabidopsis can also germinate and flower in a single growing season. These shorter-lived plants tend to be facultative — flowering in spring, summer, or fall — and in some regions this rapid life cycle enables multiple generations within a year. This variation in life histories occurs in many annual plants (Baskin and Baskin 1988) and is made possible, in part, by a range of flowering times and germination traits that can vary independently from each other (Marcer et al. 2018; Martínez-Berdeja et al. 2020). In Arabidopsis, geographic clines in breeding values for phenology, clines in allele frequencies of phenology QTL, and evidence of selection on phenology QTL support the conclusion that phenology is adapted to local environmental conditions in the wild (Caicedo et al. 2004; Stinchcombe et al. 2004; Samis et al. 2012; Fournier-Level et al. 2013; Ågren et al. 2017; Debieu et al. 2018; Exposito-Alonso et al. 2018; Lasky et al. 2018, 2024; Gamba et al. 2024a) (Figure 1). Yet, even within geographic proximity one can find genotypes with substantial genetic differences in flowering time (Jones 1971; Alonso-Blanco et al. 2016; Méndez-Vigo et al. 2022).

Life Cycles Across the Range

(A) Colder Climate (e.g. Scandinavian) Life Cycle

- Late Flowering
- High Germination in Experiment
- Lowest PTU at Flowering/Collection Date

(B) Warm Climate Life Cycle (e.g. Germany)

- Early Flowering
- High Germination in Experiment
- Highest PTU at Flowering/Collection Date

(C) Warmer Climate (e.g. Mediterranean) Life Cycle

- Early Flowering
- Low Germination in Experiment
- Low PTU at Flowering/Collection Date

Genetic Clines, Reaction Norms

(D) Genetic Clines in Flowering Time Across Climates of Origin

(E) Plasticity in Flowering Time Across Growth Environments

(F) Genetic Clines in Germination Across Climates of Origin

Figure 1. *Genotype and environment likely influence phenology of individual Arabidopsis plants across its geographic range the wild. Genetics and environmental cues may determine how long a plant remains in vegetative growth (green) or how long a plant remains dormant as a seed (brown circles), with three representative life cycles shown from Scandinavia to the Mediterranean (A–C), with shaded areas showing accumulation of photothermal units (PTUs) in the growing season (e.g. temperature >4℃). Existing knowledge of clines in flowering time and germination combined with plastic acceleration of flowering in warmer temperatures (D–F).*

There are several mechanisms that may explain why phenology observed in controlled experiments shows poor correlation with phenology observed in nature (Wilczek et al. 2009). While genotype likely influences the phenology of an individual plant in the wild, the dependence of sequential developmental transitions on each other, as well as plasticity in their expression, may limit the translation of genetic values of single stage to natural phenology. First, there are interactions between germination and flowering time, both because of a shared genetic basis and because the timing of early life transitions influences environmental exposures in later life stages (Chiang et al. 2009; Huang et al. 2010; Springthorpe and Penfield 2015; Huo et al. 2016). Secondly, there is stochasticity in individual germination and flowering time (Jimenez-Gomez et al. 2011; Abley et al. 2020). Thirdly, spatial environmental variation is extensive in nature and genotype-environment interactions have major impacts on phenology (Sasaki et al. 2015). Finally, maternal effects on phenological transitions may be strong, as reported extensively for seed dormancy (Boyd et al. 2007; Chiang et al. 2013; Burghardt et al. 2016; Huang et al. 2018). As a result, flowering time loci identified in common garden lab or field experiments might have loose relationships with variation among natural individuals (Chiang et al. 2013); genotypes differing in phenology in one set of conditions may express the same phenology in another set of conditions (Wilczek et al. 2009; Burghardt et al. 2016).

Here, we compare phenological observations of wild individual plants from natural history collections (herbaria and seed stock centers) (Miller-Rushing et al. 2006; MacGillivray et al. 2010; Davis et al. 2015; Ramirez-Parada et al. 2022) with plants in controlled environment experiments. Arabidopsis are naturally inbred (Jones 1971), thus we can compare collection dates of wild individuals with phenology of nearly genetically identical descendants in controlled experiments. This comparison provides a window into how genetic variation shapes phenology in natural populations against the forces of plasticity and GxE. Natural history collections can be used to characterize the climate conditions during development preceding collection (Bontrager et al. 2025b). We estimated phenology as collection date, and using the climate conditions preceding collection, we also estimated phenology as accumulated photothermal units (PTUs) at collection (Burghardt et al. 2015). PTUs allow us to account for weather effects on development and effectively estimate how far into growing seasons plants were collected.

Predictions for Analyses in This Study

(A) Clines, Co-Gradient Plasticity, and Local Genetic
Variation in Flowering Time Cause:

(B) High Germination for Spring Germinants Causes:

(C) Correspondence Between Early Flowering and
Spring Germination Causes:

(D) Germination Correlation with Flowering Time, Clines
in Flowering Time, and Germination Cause:

Figure 2: Existing knowledge of clines in flowering time and germination combined with plastic acceleration of flowering in warmer temperatures (see Figure 1) lead to our predictions (A–D) here) of how flowering time and base germination rates in controlled experiments will correspond to phenology in natural wild plants.

127   We developed three hypotheses for how the phenology of individual plants in nature may

128   relate to genetic variation in phenological traits, illustrated in Figure 2:

129   H1. We predict that Arabidopsis plants collected earlier in a natural growing season

130   will have early flowering and low germination breeding values in controlled

131   environments. Specifically, we predict that genetic clines in flowering time (here,

132   early flowering in warmer climates) and germination rate, combined with plastic

133   acceleration of flowering in warmer conditions (Figure 1), leads to range-wide

134   positive relationships between breeding values for flowering time, germination

135   rate, and collection date (as a proxy for flowering date in the wild; Figure 2).

136   H2. For cumulative photothermal units (PTU) at collection, we predict that spring

137   annuals with high germination rates and early flowering times will have been

138   collected later in growing seasons (high PTU) than winter annuals with later

139   flowering breeding values (Figure 2).

140   H3. Locally, within populations, environmentally induced phenological variation (i.e.,

141   plasticity associated with different timing of spring warming) is likely lower than

142   that seen across the species range. Thus we predict that flowering time breeding

143   values and germination rates would be positively related to collection dates

144   largely due to genetic effects (Figure S1).

Materials and Methods

Natural genotypes

We first conducted an experiment on a set of 101 naturally inbred genotypes ("ecotypes")

selected based on known collection dates either from the herbarium specimen label or recorded

in the Arabidopsis Biological Resource Center https://abrc.osu.edu/ database. 47 of these

ecotypes had not been studied in previous flowering time experiments and 75 had not been

included in previous germination experiments. Seed was ordered from ABRC (94 accessions) or

was germinated from herbarium sheets (7 accessions from Real Jardín Botánico Madrid,

Komarov Botanical Institute, and Oslo Natural History Museum).

Seeds from herbarium specimens were cold-stratified in tap water at pH 7 and placed at

4°C for 7 d. Seeds were then directly sown into damp Fafard germination mix and grown in

Conviron growth chambers under 10/14hr, 18/22°C days/nights. Seven herbarium accessions

germinated using this protocol. In the second round of germination, we surface sterilized seeds

using standard protocols and then cold-stratified, as above. We plated seeds onto

MS+Gamborg's vitamins + agar plates containing 1% sucrose and 10uM GA4 and then placed

them in growth chambers, as above. Two more herbarium accessions flowered using this

protocol; these were transplanted to Fafard Germination mix. To induce flowering for seed

collection, plants were exposed to 30d of 4°C with 10/18hr day/night cycles.

Flowering and germination experiments

Ecotypes were grown in common conditions prior to the flowering time experiment, and

flowering time replicates were descended from a single mother plant. For each ecotype, three

replicates were grown in separate pots. Seeds were stratified at 4°C for 5 days before sowing in

pots. Each pot was thinned to a single individual after the emergence of the second set of true

leaves. Plants were grown at 22°C under 16h days of fluorescent light in a walk-in Conviron

growth chamber (model MTPS144). Day of bolting and day petals appeared were both recorded

as measures of flowering time.

Seeds from each replicate maternal plant in the flowering time experiment were collected

and stored separately in dry conditions until the germination trial. For each treatment, forty seeds

from each parent plant (or as many seeds as were available for replicates with low fecundity)

evenly divided across 2 plates were sown on filter paper in petri dishes and germinated at

23/18°C during day/night with constant 16h daylength in a Conviron growth chamber. In total,

1,752 plates and >50k seeds were assayed.

Seeds were subjected to cold stratification at 4°C in the dark for 3 treatment lengths: 2

weeks, 3 days, and 0 days. Cold stratification can break primary dormancy, however 2 weeks of

chilling can prompt secondary dormancy in the seedbank (Penfield and Springthorpe 2012). The

difference in germination rate between 3 days and 0 days of stratification can therefore indicate

primary dormancy while the difference in germination rate between 3 days and 2 weeks of

stratification may indicate secondary dormancy. As a caveat, lower germination after 2 weeks of

stratification could be due to other, unmeasured negative effects on germination, such as

bacterial infection. We staggered the planting so that all the plates came out of stratification on

the same day. The number of seeds that had germinated in each plate was recorded on days 1, 3,

5, 10, 14, 21, and 28. Seeds were considered germinated if the radicle was visible. Because

maternal plants flowered at different times, germination rates for this experiment may be

influenced by the length of the time between flowering and planting and any after-ripening that

may have occurred. We also tested for these maternal effects among individuals of the same

inbred line (Supplemental Material).

Phenology from published experiments

We searched the literature for experiments on Arabidopsis that measured flowering time or

germination traits across different natural inbred lines (commonly referred to as Arabidopsis

"ecotypes"). The minimum number of ecotypes in any single experiment was 17. Our final set

used data from 38 previous studies (31 included some measure of flowering time, 15 included

germination) that, combined with our new experiments described above, included over 3,000

ecotypes for 86 flowering time experimental conditions and 66 germination experimental

conditions, although all ecotypes were used in only a subset of the trials and only 291 ecotypes

had a reliable date of original collection from the wild.

We used this dataset of phenology measurements from the literature to create an estimate

of genetic variation in flowering time and dormancy among ecotypes. We sought to gain

statistical power by combining data from different experiments. To make phenotypes

comparable, we standardized across treatments and experiments, transforming each flowering

time experiment such that the earliest flowering accession had a value of 0 and the latest

flowering accession had a value of 1.

We averaged standardized experimental flowering times across experiments, keeping

vernalized, non-vernalized, and field experiments separate. Here, we use 'vernalization' to

describe extended cold treatments applied to rosettes. Non-vernalized growing conditions

uncover genetic variation in flowering time due to vernalization requirements that is masked

under vernalized or fall-sown field conditions and may be important for determining phenology

and life history in the wild (Wilczek et al. 2009). However, Arabidopsis plants growing in many

natural settings are expected to experience changes in temperature and photoperiod that would be

more similar to vernalized and field experiments (Li et al. 2010). By keeping the three treatments

separate, we could test whether non-vernalized flowering time or vernalized flowering time was

more predictive of wild phenology. Field flowering time was aggregated across all seasons,

which may lead to less coherent estimates of field genetic flowering time if seasonal differences

lead to meaningful differences in flowering time, and also broken down further into spring (4

treatments over 2 experiments), summer (4 treatments, 1 experiment), and fall plantings (23

treatments, 9 experiments). To control for regional bias, this aggregation was performed both for

all experiments and without experiments that tested only ecotypes collected from a single

country.

Data collection approaches for estimating germination rate and dormancy were more

heterogeneous than for flowering time. Depending on the experiment, dormancy was reported as

the number of days after planting until a set percentage of germination was reached, the

percentage of seed germinated a given number of days after planting, the number of days of

storage until a set percentage of germination, or the germination rate after a given number of

days of storage. Because of the difference in metrics across experiments, dormancy values were

standardized by rank within each experiment with zero indicating low to no dormancy and one

indicating high dormancy. Measures that reported a percentage germinated were ranked in the

opposite direction from measures that reported number of days until germination or the number

of days of storage before 50% germination. We found that standardized rank-based metrics that

increase with dormancy (such as days to 50% germination) were correlated with rank based on

metrics that decrease with dormancy (such as percentage of seeds germinated after a set number

of days, Pearson's $r = 0.485$).

These experiments captured variation in primary dormancy, or recalcitrance to germinate

immediately after harvest. Secondary dormancy, or dormancy induced when a seed experiences

conditions unfavorable to germination, has been studied in experiments that measured an

increase in dormancy during storage. Primary and secondary dormancy could lead to different

phenological and ecological outcomes (Martinez-Berdeja et al. 2020), so we averaged across

experiments that measured primary or secondary dormancy separately to estimate each of these

two traits. Secondary dormancy was not included in the Generalized Additive Models described

below. Maternal conditions may be an important source of phenological plasticity; thus we also

used our dormancy experiment to examine how variation in flowering time among maternal

plants influences germination (Supplemental Material).

Wild phenology

The date of collection (for ecotypes maintained by ABRC) or collection date (for ecotypes grown

from a known herbarium record) was used as an estimate of reproductive phenology of

individuals in the wild (Primack et al. 2004; Davis et al. 2015). We excluded records from
regions where Arabidopsis has been recently introduced, like North America and Japan.
Accessions from outside the native range or collected after the 320[th] day of the year (which we
deemed likely errors based on location, sometimes due to intentional plantings, and were greater
than 2 standard deviations from the mean) were removed. Our geographic limits also excluded
island accessions to the south of the Mediterranean, e.g. Cape Verde Island.

We hypothesized that genetic differences in wild phenology may be more apparent if we
account for spatiotemporal environmental fluctuations causing plasticity. Therefore, we
calculated photothermal units (PTUs) for each accession from the day of collection using
monthly climate time series data from CRU (Harris et al. 2014), beginning with January 1 of
each year, following the methods of (DeLeo et al. 2020) and (Burghardt et al. 2015). PTUs
integrate the temperature and light experienced by a plant at a given location through the
growing season and therefore may capture environmental cues relevant to phenology and better
describe genetic variation in development (Wilczek et al. 2009; Brachi et al. 2010). We only
sought to use PTUs to estimate how far into growing seasons plants were collected; we did not
attempt to estimate PTUs since germination since germination timing was unknown. The models
described below use the square root of PTU because the resulting distribution was closer to
normal than the log transformation.

Statistical comparison of collection date in the wild versus phenology in experiments
H1: Range-wide comparison of phenology breeding values vs. collection date
We tested if genetic variation in normalized phenology measured on naturally inbred lines
explained variation in the wild phenology of the parent of the line (Figure 2), using linear
regression between the normalized phenology breeding values (flowering time and dormancy
rank) and collection day. Vernalized, non-vernalized, and field flowering times were modelled
separately, because the genetic variation in phenology uncovered by each treatment could relate
to wild phenology in different ways. Under the hypothesis that vernalization and field conditions
better recreate environments that a plant would experience at their geographic origin, these
genetic flowering time measures should be more positively related to wild flowering time.
Likewise, non-vernalized flowering times may better recreate the original temporal niches of
summer annuals and thus be positively related to flowering time in these ecotypes. However, it is

also possible that long non-vernalized flowering times indicate obligate winter annuals. In these

plants, non-vernalized flowering times would be negatively related to wild phenology since later

flowering times in non-vernalized experiments would indicate plants that overwinter and are

collected early the next year (Figure 2).

We also investigated how range-wide differences in wild phenology related to regional

differences in breeding values using a larger sample size, taken from a broader collection of

ecotypes than just those with documented collection dates. This comparison included stock

center ecotypes with experimental phenology data and known location-of-origin, but no recorded

date of wild collection, and herbarium specimens for which we had not germinated seed and

grown plants to measure traits in experiments. To do so, we first estimated wild flowering time

for each stock center ecotype in experiments using herbarium records near the ecotype collection

location. These wild flowering times were estimated from a previously published generalized

additive model (GAM) with spatially-varying intercepts of herbaria collection dates from 2,655

Eurasian Arabidopsis records used in (DeLeo et al. 2020), which included year of collection as a

nuisance variable, as collection date has changed over time across the range of Arabidopsis

(DeLeo et al. 2020). Values of the smooth intercept surface were extracted at the coordinates of

stock center ecotypes having an experimentally measured flowering time. These estimated wild

flowering times were regressed against breeding values for flowering times under vernalized,

non-vernalized, and field experiments and rank primary dormancy. In addition, for these same

data we performed Spearman rank correlation between estimated wild flowering times and

phenology breeding values.

H2: Range-wide comparison of phenology breeding values vs. PTU

To account for plasticity in phenology due to differences among locations in the timing and

progression of growing seasons, we also tested the relationship between phenology breeding

values and the estimated PTUs at the time of collection from the wild. We conducted the same

analyses across the range of Arabidopsis as described in H1, except that the PTU at collection

was used instead of collection date. Specifically, we conducted linear regression between

normalized breeding values with PTU at collection and we also conducted linear regression

between breeding values and the PTUs at collection of nearby herbarium specimens.

H3: Within population variation

Next, we tested for the effects of genetic variation in flowering time and dormancy on

phenological variation *within* populations in the wild. By first accounting for geographic

variation in mean flowering time, we aimed to isolate local within-population variation (Figure

S1). This analysis, in essence, asks whether genetic variation in individual phenological stages

explains natural phenological variation within populations. Generalized Additive Models

(GAMs) allow for model parameters to vary smoothly across space and thus can capture spatially

varying patterns, e.g. in local average collection date (Yee and Mackenzie 2002; Yee and

Mitchell 2006). We built a GAM for the dependent variable of collection date (Y) at location j of

individual i which included covariates of experimental flowering time (β_1), dormancy (β_2),

elevation (β_3) and a spatially varying intercept (μ_j):

Equation 1.

$Y_{ij} = \mu_j + \beta_1 \text{Flowering Time}_{ij} + \beta_2 \text{Primary Dormancy}_{ij} + \beta_3 \text{Elevation}_j + \varepsilon_{ij}$

The smooth spatially varying intercept term μ_j estimates spatial variation in mean

collection date (e.g., due to environmental gradients). The upper limit on degrees of freedom for

the spatially varying intercept was increased to 45 following the recommendations of Wood

(2006). In models that allowed for higher degrees of freedom, the effective degrees of freedom

did not meaningfully increase. Elevation was included because of its known importance to

flowering times and spring onset (Vidigal et al. 2016; Gamba et al. 2023) and the high resolution

of elevation data compared to smooth variation in GAM parameter surfaces. We fit GAMs using

the 'gam' function in the 'mgcv' package in R (Wood 2006) using restricted maximum

likelihood (REML), although Generalized Cross Validation returned similar estimates. Model

fitting allowed for penalization of smooth terms to 0 so that uninformative covariates could be

removed from the equation. Residuals were plotted using the 'gam.check' function in 'mgcv'

(Wood 2006), and one ecotype (Nok-10) was removed from our analyses that was an extreme

outlier based on its residuals. Spatial variation in β coefficients for flowering time and dormancy

covariates (Eq 1) were not significant, so a model using a constant coefficient was used in our

analyses. Field, vernalized, and non-vernalized flowering times were tested in separate versions

of the model and tested different hypotheses with regards to wild phenology. To compare among

the three measures of flowering time, a version of Equation 1 was fit on a subset of ecotypes that

had all three flowering time measures, and Akaike's Information Criterion (Akaike 1974) was

compared.

In the wild, both flowering time and dormancy contribute to phenology. Thus, we also

tested whether including interactions between flowering time and primary dormancy (β_4)

improved the model:

Equation 2.

$Y_{ij} = \mu_j + \beta_1$Flowering Time$_{ij} + \beta_2$Primary Dormancy$_{ij} + \beta_3$Elevation$_j + \beta_4$Flowering Time$_{ij}$ *

Primary Dormancy$_{ij} + \varepsilon_{ij}$

Finally, given the importance of plasticity in response to temperature in the timing of

germination and flowering time, PTU might better capture genetic variation in phenology in the

wild (Figure 2 "Predictions"). Therefore, we also tested the models above using PTU in place of

collection day to correct for climate.

Results

Genetic correlation among phenology traits in controlled experiments

Across all compiled experiments (both previously published and our new experiments) mean

flowering times of ecotypes measured in vernalized and non-vernalized experiments were

strongly positively correlated (Pearson's $r = 0.90$, $p < 0.001$). Although fall-sown field trials

might be expected to expose plants to cold seasonal temperatures that give vernalization cues,

standardized flowering time breeding values in both non-vernalized and vernalized experiments

were similarly correlated to published field experiment values ($r_{\text{non-vernalized}} = 0.77$, $r_{\text{vernalized}} =$

0.72, $p < 0.001$ for both). Primary and secondary dormancy were negatively correlated, but not

significantly so ($r = -0.15$, $p = 0.11$). Despite known interactions between flowering time and

dormancy due to seed maturation environment and pleiotropy of causal loci, only non-vernalized

flowering time was significantly correlated with primary dormancy ($r = -0.17$, $p = 0.02$, Figure 3;

for a heatplot of all phenotypes, see Figure S2). These correlations were qualitatively unchanged

when using an alternate standardization method across experiments (Figure S3).

Figure 3. Standardized phenology traits combined across multiple published controlled experiments combined with our new experiments. Standardized flowering times were correlated across all three types of treatments (vernalized, non-vernalized, field), but they were weakly correlated with dormancy.

H1: Comparing wild phenology with breeding values in experiments across the range

Experimental breeding values for phenology traits were modest predictors of collection day in simple linear models, with the only statistically significant predictors being non-vernalized flowering time (estimated slope = 27.7 days/standardized flowering time, $r^2 = 0.09$, p < 0.001) and primary dormancy (-17.1 days/standardized rank dormancy, $r^2 = 0.02$, p = 0.02). Flowering time breeding values under vernalized or field conditions did not significantly predict day of collection with a linear model (p > 0.05) (Figure 4). Flowering time and germination measured within individual experiments did not predict wild collection day better than our averaged

breeding values, suggesting some power was gained by combining individual experiments into

standardized values (Figure S7).

Figure 4. Date of collection (left
panels) and PTUs (photothermal
units) at collection (right panels)
of wild plants compared to
standardized breeding values for
phenological traits from
experiments. Collection day of
wild plants was positively related
to non-vernalized flowering time
in experiments. Collection day
was negatively related to primary
dormancy. The values shown for
field flowering time experiments
were calculated from fall-sown
experiments only.

Flowering time and dormancy breeding values predict regional variation in collection date of

nearby herbarium specimens

Our model of collection date, above, was limited to the 227 ecotypes with both a known
collection date and experimental phenology data. To expand our predictions across more of the
Arabidopsis range, we tested whether genetic variation in flowering time in experiments predicts
local mean collection dates of herbarium specimens across the landscape. First, we found that
local mean collection dates across a landscape from >2500 herbarium collections were positively
correlated with actual collection days of individual stock center ecotypes (which were not used in
fitting the model, $\rho = 0.51$, $p < 0.001$ Figure S8), validating these local predictions of phenology
in the wild based on herbarium collections.

Figure 5. Genetic variation in flowering time experiments predicts collection dates of nearby

herbarium specimens. A) There is geographic variation in collection dates as shown by

estimated mean date from GAMs with spatially varying means, with earlier collections (light

green) around the Mediterranean (DeLeo et al. 2020). *B) Local mean collection dates were*

positively correlated with genetic variation in vernalized flowering time experiments (p < 0.01),

and all ecotypes from locations with earlier mean collection dates in the wild (largely

Mediterranean) had rapid flowering breeding values. Primary dormancy breeding values were

negatively correlated with local mean collection dates (p < 0.001).

As with the collection dates of the original maternal sources of stock center lines, breeding values for non-vernalized flowering times significantly predicted local mean herbarium collection dates ($r^2 = 0.10$, $p < 0.001$). For comparison, latitude was a better predictor of mean herbarium collection date ($r^2 = 0.37$, $p < 0.001$). These patterns were similar for vernalized, non-vernalized, and field measurements, although not significant for field trials when fall plantings were grouped with spring and summer plantings ($r^2 = 0.07$, $p = 0.002$, $r^2 = 0.10$, $p < 0.001$, and $r^2 = 0.02$, $p = 0.13$, respectively, Figure 5). These results show that range-wide genetic variation in phenology measured in controlled environments corresponds to actual phenology of plants in nature. However, because plants in nature from different regions experience different environments, these patterns cannot determine the degree to which genetic variation causes the variation in natural phenology as opposed to genetic variation being correlated with environmental variation that causes phenological plasticity in nature (i.e. co-gradient variation, Figure 1).

H2: Range-wide comparison of phenology breeding values vs. Photothermal Units

Because environmental differences among locations can influence phenology (Fournier-Level et al. 2013), we also calculated PTU at collection and compared to these values to traits from experiments. However, PTU at collection was not significantly predicted by any phenology trait breeding values (Figure 4). Thus, while ecotypes with later flowering time breeding values tend to be collected later in the year, these later flowering ecotypes are not collected at higher PTU, i.e. later in the local growing season. The fact that phenology breeding values are correlated with

date, but not PTU at collection, is consistent with the hypothesis that geographic climate

variation maintains clines in breeding values due to local adaptation and clines in collection date

partly due to plasticity (co-gradient variation), generating a breeding value-collection date

correlation. These correlations were qualitatively unchanged when using an alternate

standardization across experiments and when excluding herbarium specimens (Figure S4–S6).

H3: Variation in phenology breeding values does not predict local, within-region variation in

wild collection dates

We next examined whether genetic values could explain variation in natural phenology *within*

regions (Figure S1). We used GAMs that included spatially varying intercepts and an elevation

covariate to account for the among population variation in environmental effects on collection

dates, leaving local variation to be potentially explained by genetic variation. However, we found

that normalized flowering time (slope of non-vernalized flowering time = -8.2, p = 0.098) and

dormancy (slope of dormancy = 2.2, p = 0.51) were not significantly related to day of collection

in this GAM (Equation 1), although elevation effects were significant (slope of elevation =

0.012, p = 0.049, Figure 6, Figure S9, Table S1). In comparing models using vernalized, non-

vernalized, and field experiments, we found that standardized non-vernalized flowering times led

to lower AIC compared to other flowering time measures but higher AIC than a model without

flowering time at all, although the difference was slight (difference < 2 for an AIC of 530). We

used non-vernalized flowering times in the final models, in part because there were more

ecotypes with non-vernalized flowering times (184 for non-vernalized vs 113 for vernalized or

78 for field).

Figure 6. Top: Spatial variation in date of collection for 227 ecotypes with known collection

dates. A) Color represents the intercept value for collection day at a location after fitting a

linear (i.e. non-spatially varying) relationship with elevation, experimental primary dormancy

and non-vernalized flowering time. Much of the variation in collection date was explained by

the spatially varying intercept rather than being explained by experimental flowering time. B)

Residuals of a GAM predicting collection date using only intercept and elevation parameters,

compared to non-vernalized flowering time, primary dormancy, or fitted collection dates.

Although we found that flowering time, dormancy, and the interaction between the two

were not significantly related to collection day, including phenology breeding values did lower

the AIC and slightly increase the deviance explained by the model. This suggests that dormancy

and flowering time could be related to variation in phenology within regions. However, most of

the variation was described by the spatially varying intercept –a model with only a spatially

varying intercept had a deviance explained of 80.1% – which could account for geographic

genetic clines in phenology and plastic responses to geographic climate gradients (Figure S8).

As with the linear models, we also tested whether genetic variation in phenology in

experiments could explain local variation PTU of collection date in nature. Again, we found that

flowering time and dormancy were not significantly related to wild PTU at collection in GAMs

with spatially varying intercepts. Elevation was negatively related to PTU, indicating that higher

elevation plants were collected earlier in growing seasons, reflecting a potential winter annual

strategy (Figure 1, slope: -0.018 $\sqrt{PTU}$/m, $p < 0.01$). Additionally, elevation was positively

correlated with date of collection (slope: 0.012 days/m, $p < 0.05$).

Discussion

Plant phenology comprises multiple traits that contribute to fitness. In Arabidopsis, flowering

time and germination can vary independently to create a landscape of possible life histories

across environments (Debieu et al. 2013; Marcer et al. 2018; Martínez-Berdeja et al. 2020). This

variation in phenology is likely partly maintained by selection, given that the traits vary

geographically in association with climate (Fournier-Level et al. 2011; Vidigal et al. 2016;

Exposito-Alonso 2020) and biotic pressures (Lyons et al. 2015; Davila Olivas et al. 2017), that

QTL show evidence of local adaptation (Gamba et al. 2023; Lasky et al. 2024), and that the

timing of individual phenological transitions is often correlated with fitness in experiments

(Korves et al. 2007; Stock et al. 2015). Standing genetic variation is necessary for the timing of

developmental transitions to evolve in response to selection by local environments, but little is

known about the heritability of phenology in natural, wild individuals. Furthermore, because

multiple developmental transitions influence phenology, the effects of variation in the timing of

one stage might be compensated by (or, in under-parameterized experiments, confounded by)

variation in the timing of other stages. Given widespread evidence that genetic variation in life

history is adaptive, we investigated to what degree experimentally measured genetic variation in

Arabidopsis phenology predicts phenology of plants in the wild, based on collection dates of

natural history records.

The influence of genotype and environment on flowering time and dormancy have been

well studied experimentally in Arabidopsis. Yet, common garden and controlled environment

experiments must be designed thoughtfully to highlight genetic differences between ecotypes

that are relevant to selection in natural environments (Karrenberg and Widmer 2008). While

experimental design has increasingly recognized the importance of field conditions to acquire a

measure of phenology that is more representative of nature (Wilczek et al. 2009; Anderson et al.

2012; Brachi et al. 2013; Poorter et al. 2016), there has been little comparison of controlled

experiments, field or otherwise, to phenology in wild, naturally cycling individual plants. We

found that flowering time and dormancy breeding values are related to date of collection and

capture variation in phenology primarily *among* populations in the wild, but explain little

variation *within* regions/populations.

Genetic variation in flowering time and dormancy predicts variation in wild phenology among

populations across the species range

Across the species range, both primary dormancy and non-vernalized flowering time

significantly predicted collection dates. These phenological transitions are not independent of

each other. As seen in individual experiments (Martínez-Berdeja et al. 2020), we found that the

average breeding values for primary dormancy and non-vernalized flowering time were

negatively correlated (Figure 3). Furthermore, dormancy is strongly affected by environmental

conditions during seed maturation (Penfield and Springthorpe 2012; Huang et al. 2015;

Burghardt et al. 2016). Thus, interaction between flowering time and dormancy in the wild could

stem from both genetic correlations and environmentally induced interactions. Despite our

expectation for interactions between germination and flowering traits, models of collection day

that included flowering time by dormancy interactions had a higher AIC and did not explain

more deviance than a model with both traits separately. Including interactions between flowering

time and dormancy in our models did not help to explain collection date in the wild.

We found that predicted wild collection date was positively related to flowering time

breeding values when we extended our analysis to predict wild flowering times from a smoothed

surface of collection dates fit to herbarium records. In our linear models, average flowering time

breeding value was more closely related to this estimated day of collection than was the actual

date of collection of ecotypes in experiments. However, a large portion of phenological variation

within populations remained unexplained even with phenological breeding values. We attempted

to avoid records that were unusually young or old by only using specimens that had both flowers

and fruits; moreover, both our herbarium and seed collections spanned nearly the entire year

(Julian days 5–350 for herbarium records, 43–346 for seed collections). Still, herbarium records

are known to skew slightly towards earlier in seasons (though removing herbarium specimens

from our analyses did not change results) (Daru et al. 2018), while seed collections must be

collected long enough after the initiation of flowering to allow for the development of some

mature seeds. A single observation during reproduction, common for natural history collection

vouchers, cannot resolve the uncertainty around when plants begin their vegetative growth in the

wild, making it difficult to describe phenology of the full life cycle. With better information and

in climates where growing conditions are clearly defined, it may be possible to estimate

germination dates and full life cycle phenology from herbarium specimens (Bontrager et al.

2025a). Nevertheless, natural history collections could perhaps provide large numbers of

observations useful for fine tuning models of landscape phenology.

Spring onset differs among locations, and so a calendar date at a higher latitude or

elevation may be functionally earlier in the season than a lower latitude or elevation. PTUs

represent how much of a growing season has passed by a given date by recording temperature

and daylight above a threshold. Thus, plants collected at lower PTUs may be early flowering

(developmentally) despite a later collection date. Higher PTU may indicate plants growing as

summer or fall annuals that germinated later in the year. In that case, ecotypes with higher PTUs

would likely be fast cycling, earlier flowering plants (Figure 2). While we did find a negative

correlation between flowering time breeding values in field and vernalized experiments versus

PTU, we see the opposite in non-vernalized experiments. This last observation is somewhat

surprising, since we expected plants that flower later in the absence of vernalization to be more

likely to grow as winter annuals in the wild and flower early in the spring (Figure 1). Higher

PTUs in these presumptive winter annuals may suggest that plants in these locations regulate

their life histories to flower later in the year than expected by temperature and daylight alone.

There is some evidence that later flowering ecotypes could in fact be more flexible in their

flowering time relative to germination than early flowering ecotypes (Miryeganeh et al. 2018),

which could hide the signal of early spring flowering we expected in late flowering Arabidopsis.

Within regions, breeding values do not predict phenology in the wild

Because Arabidopsis exhibits substantial local within-population variation in phenology (Jones

1971; Brachi et al. 2013; Alonso-Blanco et al. 2016; DeLeo et al. 2020), we asked how genetic

variation in phenology was related to phenological variation within populations. This within-

population phenological variation can have fitness consequences. Variation contributes to

population persistence in variable environments, as when different germination behavior

provides bet hedging in the seedbank (Cohen 1967; Gremer and Venable 2014).

We found that neither dormancy nor flowering time breeding values from experiments

were significantly related to collection date when we accounted for geographic variation in local

mean collections dates in a GAM. However, flowering time and dormancy did improve GAMs

for collection date over a model that included only geographic location and elevation, suggesting

that genetic variation in flowering time does explain a minor fraction of the phenological

variation within regions. Non-vernalized flowering time was negatively, non-significantly related

to local variation in collection date, perhaps because in many locations, genotypes flowering

later in experiments without vernalization are more likely to behave as winter annuals, flowering

early in the spring before spring annuals.

In our models, there was very little difference between non-vernalized, vernalized, and

field-measured flowering times in predicting collection dates, despite the expectation that field

experiments recreate conditions similar to nature. However, data from field experimental trials

were available for a smaller set of ecotypes than non-vernalized and vernalized indoor trials, and

we lacked Iberian ecotypes with field experimental data and date of collection, limiting the

usefulness of this measure across the range. Our model provided weak evidence that the breeding

values for flowering time and dormancy explained within-population variation in wild collection

day. However, our set of ecotypes with known collection day numbered 258, potentially still too

few to detect strong statistical significance given the noise arising from plasticity in responses to

local environmental gradients.

Conclusion

We demonstrated that experimental measures of genetic variation in the timing of individual

developmental transitions in Arabidopsis are significantly, but modestly, correlated with

phenological variation in wild plants. On one hand, the existence of a relationship suggests there

is some relevance of phenotypes measured in experiments to those exhibited in nature. However,

the weakness of the relationships (especially locally, within regions) suggests the existence of

either a) complex interactions between developmental stages to determine phenology, (b) low

heritability in nature, or (c) extensive rank changing genotype-environment interactions across

microsites. Genetic differences in Arabidopsis phenology across the species range are often cited

as evidence of the adaptive importance of phenology (Stinchcombe et al. 2004; Fournier-Level et

al. 2011; Samis et al. 2012; Brachi et al. 2013; Exposito-Alonso 2020). Incorporating

information on between-population and within-population phenology diversity helps to clarify

how selection may be acting on phenology across the landscape and how influential plasticity is

in the phenology of this annual herb. Yet, genetic flowering times among populations across the

species range are associated with phenological variation in natural history collections, due to

some combination of genotypic and environmental effects (and their possible co-gradient

variation) (Jones et al. 2024). While phenological plasticity is often of large magnitude, it may

not be enough to maintain fitness under environmental change (Zettlemoyer et al. 2024).

Ultimately, controlled experiments on many plants have suggested phenology is under selection

in nature, but understanding more subtle environmental differences and stochasticity may help to

clarify the evolution of phenology and translate genetic values into reliable predictions with and

between populations.

References

Abley, K., P. Formosa-Jordan, H. Tavares, E. Chan, O. Leyser, and J. C. W. Locke. 2020. An ABA-GA
bistable switch can account for natural variation in the variability of Arabidopsis seed germination
time. bioRxiv, doi: https://doi.org/10.1101/2020.06.05.135681.

Ågren, J., C. G. Oakley, S. Lundemo, and D. W. Schemske. 2017. Adaptive divergence in flowering time
among natural populations of Arabidopsis thaliana: Estimates of selection and QTL mapping.
Evolution (N Y) 71:550–564.

Akaike, H. 1974. A New Look at the Statistical Model Identification. IEEE Trans Automat Contr 19:716–
723.

Alonso-Blanco, C., J. Andrade, C. Becker, F. Bemm, J. Bergelson, K. M. M. Borgwardt, J. Cao, E. Chae, T.
M. M. Dezwaan, W. Ding, J. R. R. Ecker, M. Exposito-Alonso, A. Farlow, J. Fitz, X. Gan, D. G. G.
Grimm, A. M. M. Hancock, S. R. R. Henz, S. Holm, M. Horton, M. Jarsulic, R. A. A. Kerstetter, A.
Korte, P. Korte, C. Lanz, C. R. Lee, D. Meng, T. P. P. Michael, R. Mott, N. W. W. Muliyati, T. Nägele,
M. Nagler, V. Nizhynska, M. Nordborg, P. Y. Y. Novikova, F. X. Picó, A. Platzer, F. A. A. Rabanal, A.
Rodriguez, B. A. A. Rowan, P. A. A. Salomé, K. J. J. Schmid, R. J. J. Schmitz, Ü. Seren, F. G. G.
Sperone, M. Sudkamp, H. Svardal, M. M. M. Tanzer, D. Todd, S. L. L. Volchenboum, C. Wang, G.
Wang, X. Wang, W. Weckwerth, D. Weigel, and X. Zhou. 2016. 1,135 genomes reveal the global
pattern of polymorphism in *Arabidopsis thaliana*. Cell 166:481–491.

Alpert, P., and E. L. Simms. 2002. The relative advantages of plasticity and fixity in different
environments: When is it good for a plant to adjust? Evol Ecol 16:285–297.

Amasino, R. 2004. Vernalization, competence, and the epigenetic memory of winter. Plant Cell, doi:
10.1105/tpc.104.161070.

Andrés, F., and G. Coupland. 2012. The genetic basis of flowering responses to seasonal cues.

Auge, G. A., L. K. Blair, A. Karediya, and K. Donohue. 2018. The autonomous flowering-time pathway
pleiotropically regulates seed germination in Arabidopsis thaliana. Ann Bot 121:183–191.

Balasubramanian, S., S. Sureshkumar, J. Lempe, and D. Weigel. 2006. Potent induction of *Arabidopsis*
thaliana flowering by elevated growth temperature. PLoS Genet 2:0980–0989.

Baskin, C. C., and J. M. Baskin. 1988. Germination ecophysiology of herbaceous plant species in a
temperate region. Am J Bot 75:286–305.

Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using lme4. J
Stat Softw 67:1–48.

Baythavong, B. S. 2011. Linking the spatial scale of environmental variation and the evolution of
phenotypic plasticity: Selection favors adaptive plasticity in fine-grained environments. American
Naturalist 178:75–87.

Boyd, E. W., L. A. Dorn, C. Weinig, and J. Schmitt. 2007. Maternal effects and germination timing
mediate the expression of winter and spring annual life histories in *Arabidopsis thaliana*. Int J Plant
Sci 168:205–214.

Brachi, B., N. Faure, M. Horton, E. Flahauw, A. Vazquez, M. Nordborg, J. Bergelson, J. Cuguen, and F.
Roux. 2010. Linkage and association mapping of *Arabidopsis thaliana* flowering time in nature.
PLoS Genet, doi: 10.1371/journal.pgen.1000940.

Brachi, B., R. Villoutreix, N. Faure, N. Hautekèete, Y. Piquot, M. Pauwels, D. Roby, J. Cuguen, J. Bergelson,
and F. Roux. 2013. Investigation of the geographical scale of adaptive phenological variation and its
underlying genetics in *Arabidopsis thaliana*. Mol Ecol 22:4222–4240.

Burghardt, L. T., B. R. Edwards, and K. Donohue. 2016. Multiple paths to similar germination behavior in
Arabidopsis thaliana. New Phytologist 209:1301–1312.

Burghardt, L. T., C. J. E. Metcalf, A. M. Wilczek, J. Schmitt, and K. Donohue. 2015. Modeling the influence
of genetic and environmental variation on the expression of plant life cycles across landscapes. Am
Nat 185:212–227. University of Chicago Press Chicago, IL.

Caicedo, A. L., J. R. Stinchcombe, K. M. Olsen, J. Schmitt, and M. D. Purugganan. 2004. Epistatic
interaction between Arabidopsis FRI and FLC flowering time genes generates a latitudinal cline in a
life history trait. Proc Natl Acad Sci U S A 101:15670–15675.

Chiang, G. C. K., D. Barua, E. Dittmar, E. M. Kramer, R. R. de Casas, and K. Donohue. 2013. Pleiotropy in
the wild: The dormancy gene dog1 exerts cascading control on life cycles. Evolution (NY) 67:883–
893.

Chiang, G. C. K., D. Barua, E. M. Kramera, R. M. Amasino, and K. Donohue. 2009. Major flowering time
gene, FLOWERING LOCUS C, regulates seed germination in Arabidopsis thaliana. Proc Natl Acad Sci
USA 106:11661–11666.

Cohen, D. 1967. Optimizing reproduction in a randomly varying environment when a correlation may
exist between the conditions at the time a choice has to be made and the subsequent outcome. J
Theor Biol 16:1–14.

Daru, B. H., D. S. Park, R. B. Primack, C. G. Willis, D. S. Barrington, T. J. S. Whitfeld, T. G. Seidler, P. W.
Sweeney, D. R. Foster, A. M. Ellison, and C. C. Davis. 2018. Widespread sampling biases in herbaria
revealed from large-scale digitization. New Phytologist 217:939–955.

Davila Olivas, N. H., E. Frago, M. P. M. Thoen, K. J. Kloth, F. F. M. Becker, J. J. A. van Loon, G. Gort, J. J. B.
Keurentjes, J. van Heerwaarden, and M. Dicke. 2017. Natural variation in life history strategy of
Arabidopsis thaliana determines stress responses to drought and insects of different feeding guilds.
Mol Ecol 26:2959–2977.

Davis, C. C., C. G. Willis, B. Connolly, C. Kelly, and A. M. Ellison. 2015. Herbarium records are reliable
sources of phenological change driven by climate and provide novel insights into species'
phenological cueing mechanisms. Am J Bot 102:1599–1609.

Debieu, M., C. Tang, B. Stich, T. Sikosek, S. Effgen, E. Josephs, J. Schmitt, M. Nordborg, M. Koornneef,
and J. de Meaux. 2013. Co-Variation between Seed Dormancy, Growth Rate and Flowering Time
Changes with Latitude in *Arabidopsis thaliana*. PLoS One, doi: 10.1371/journal.pone.0061075.

DeLeo, V. L., D. N. L. Menge, E. M. Hanks, T. E. Juenger, and J. R. Lasky. 2020. Effects of two centuries of
global environmental variation on phenology and physiology of *Arabidopsis thaliana*. Glob Chang
Biol 26:523–538.

Donohue, K. 2005. Niche construction through phenological plasticity: Life history dynamics and
ecological consequences.

Exposito-Alonso, M. 2020. Seasonal timing adaptation across the geographic range of Arabidopsis
thaliana. Proceedings of the National Academy of Sciences 117:9665–9667.

Exposito-Alonso, M., F. Vasseur, W. Ding, G. Wang, H. A. Burbano, and D. Weigel. 2018. Genomic basis
and evolutionary potential for extreme drought adaptation in *Arabidopsis thaliana*. Nat Ecol Evol,
doi: 10.1038/s41559-017-0423-0.

Footitt, S., Z. Huang, H. A. Clay, A. Mead, and W. E. Finch-Savage. 2013. Temperature, light and nitrate
sensing coordinate Arabidopsis seed dormancy cycling, resulting in winter and summer annual
phenotypes. Plant Journal 74:1003–1015.

Fournier-Level, A., A. Korte, M. D. Cooper, M. Nordborg, J. Schmitt, and A. M. Wilczek. 2011. A map of
local adaptation in *Arabidopsis thaliana*. Science (1979), doi: 10.1126/science.1209271.

Fournier-Level, A., A. M. Wilczek, M. D. Cooper, J. L. Roe, J. Anderson, D. Eaton, B. T. Moyers, R. H.
Petipas, R. N. Schaeffer, B. Pieper, M. Reymond, M. Koornneef, S. M. Welch, D. L. Remington, and J.
Schmitt. 2013. Paths to selection on life history loci in different natural environments across the
native range of *Arabidopsis thaliana*. Mol Ecol 22:3552–3566.

Franco, M., and J. Silvertown. 1996. Life history variation in plants: An exploration of the fast-slow
continuum hypothesis. Philosophical Transactions of the Royal Society B: Biological Sciences
351:1341–1348.

Gamba, D., C. Lorts, A. Haile, S. Sahay, L. Lopez, T. Xia, E. Kulesza, D. Elango, J. Kerby, M. Yifru, C. E.
Bulafu, T. Wondimu, K. Glowacka, and J. R. Lasky. 2023. The genomics and physiology of abiotic
stressors associated with global elevation gradients in *Arabidopsis thaliana*. bioRxiv preprint.

Gremer, J. R., and D. L. Venable. 2014. Bet hedging in desert winter annual plants: Optimal germination
strategies in a variable environment. Ecol Lett 17:380–387.

Harris, I., P. D. Jones, T. J. Osborn, and D. H. Lister. 2014. Updated high-resolution grids of monthly
climatic observations–the CRU TS3. 10 Dataset. International journal of climatology 34:623–642.
Wiley Online Library.

Hu, J., L. Lei, and J. De Meaux. 2017. Temporal fitness fluctuations in experimental *Arabidopsis thaliana*
populations. PLoS One 12.

Huang, X., J. Schmitt, L. Dorn, C. Griffith, S. Effgen, S. Takao, M. Koornneef, and K. Donohue. 2010. The
earliest stages of adaptation in an experimental plant population: Strong selection on QTLS for
seed dormancy. Mol Ecol 19:1335–1351.

Huang, Z., S. Footitt, A. Tang, and W. E. Finch-Savage. 2018. Predicted global warming scenarios impact
on the mother plant to alter seed dormancy and germination behaviour in Arabidopsis. Plant Cell
Environ 41:187–197.

Huang, Z., H. Ölçer-Footitt, S. Footitt, and W. E. Finch-Savage. 2015. Seed dormancy is a dynamic state:
Variable responses to pre- and post-shedding environmental signals in seeds of contrasting
Arabidopsis ecotypes. Seed Sci Res 25:159–169.

Huo, H., S. Wei, and K. J. Bradford. 2016. DELAY of GERMINATION1 (DOG1) regulates both Seed
dormancy and flowering time through microRNA pathways. Proc Natl Acad Sci U S A 113:E2199–
E2206.

Jimenez-Gomez, J. M., J. A. Corwin, B. Joseph, J. N. Maloof, and D. J. Kliebenstein. 2011. Genomic
analysis of QTLs and genes altering natural variation in stochastic noise. PLoS Genet 7.

Johnston, I. G., and G. W. Bassel. 2018. Identification of a bet-hedging network motif generating noise in
hormone concentrations and germination propensity in Arabidopsis. J R Soc Interface 15.

Jones, C. V, C. E. Regan, J. A. Firth, E. F. Cole, and B. C. Sheldon. 2024. Environmental similarity between
relatives reduces heritability of reproductive timing in wild great tits. bioRxiv 2024.03.11.584392.

Juenger, T. E., S. Sen, K. A. Stowe, and E. L. Simms. 2005. Epistasis and genotype-environment
interaction for quantitative trait loci affecting flowering time in *Arabidopsis thaliana*. Pp. 87–105 *in*
Genetica.

Jump, A. S., R. Marchant, and J. Peñuelas. 2009. Environmental change and the option value of genetic
diversity.

Karrenberg, S., and A. Widmer. 2008. Ecologically relevant genetic variation from a non-Arabidopsis
perspective.

Kenney, A. M., J. K. McKay, J. H. Richards, and T. E. Juenger. 2014. Direct and indirect selection on
flowering time, water-use efficiency (WUE, δ13C), and WUE plasticity to drought in *Arabidopsis*
thaliana. Ecol Evol 4:4505–4521.

Korves, T. M., K. J. Schmid, A. L. Caicedo, C. Mays, J. R. Stinchcombe, M. D. Purugganan, and J. Schmitt.
2007. Fitness Effects Associated with the Major Flowering Time Gene FRIGIDA in *Arabidopsis*
thaliana in the Field . Am Nat, doi: 10.1086/513111.

Lasky, J. R., B. R. Forester, and M. Reimherr. 2018. Coherent synthesis of genomic associations with
phenotypes and home environments. Molecular Ecology Resources 18:91–106.

Lasky, J. R., M. Takou, D. Gamba, and T. H. Keitt. 2024. Estimating scale-specific and localized spatial
patterns in allele frequency. Genetics 227:3, doi: 10.1093/genetics/iyae082.

Lawrence-Paul, E. H., and J. R. Lasky. 2024. Ontogenetic changes in ecophysiology are an understudied
yet important component of plant adaptation. Am J Bot e16294.

Lempe, J., S. Balasubramanian, S. Sureshkumar, A. Singh, M. Schmid, and D. Weigel. 2005. Diversity of
flowering responses in wild *Arabidopsis thaliana* strains. PLoS Genet, doi:
10.1371/journal.pgen.0010006.

Li, Y., Y. Huang, J. Bergelson, M. Nordborg, and J. O. Borevitz. 2010. Association mapping of local
climate-sensitive quantitative trait loci in *Arabidopsis thaliana*. Proc Natl Acad Sci USA 107:21199–
21204.

Lorts, C. M., and J. R. Lasky. 2020. Competition × drought interactions change phenotypic plasticity and
the direction of selection on Arabidopsis traits. New Phytologist n/a. John Wiley & Sons, Ltd.

Ludlow, M. M. 1989. Strategies of response to water stress. Pp. 269–281 *in* K. H. Kreeb, H. Richter, and
T. M. Hinckley, eds. Structural and functional responses to environmental stresses: water shortage.
SPB Academic Publishers, Berlin.

Lyons, R., A. Rusu, J. Stiller, J. Powell, J. M. Manners, and K. Kazan. 2015. Investigating the association
between flowering time and defense in the *Arabidopsis thaliana-Fusarium oxysporum* interaction.
PLoS One, doi: 10.1371/journal.pone.0127699.

MacGillivray, F., I. L. Hudson, and A. J. Lowe. 2010. Herbarium collections and photographic images:
Alternative data sources for phenological research. Pp. 425–461 *in* Phenological Research:
Methods for Environmental and Climate Change Analysis. Springer Netherlands, Dordrecht.

Marcer, A., D. S. Vidigal, P. M. A. James, M. J. Fortin, B. Méndez-Vigo, H. W. M. Hilhorst, L. Bentsink, C.
Alonso-Blanco, and F. X. Picó. 2018. Temperature fine-tunes Mediterranean *Arabidopsis thaliana*
life-cycle phenology geographically. Plant Biol 20:148–156.

Martínez-Berdeja, A., M. C. Stitzer, M. A. Taylor, M. Okada, E. Ezcurra, D. E. Runcie, and J. Schmitt. 2020.
Functional variants of DOG1 control seed chilling responses and variation in seasonal life-history
strategies in *Arabidopsis thaliana*. Proc Natl Acad Sci USA 117:2526–2534.

Méndez-Vigo, B., A. R. Castilla, R. Gómez, A. Marcer, C. Alonso-Blanco, and F. X. Picó. 2022.
Spatiotemporal dynamics of genetic variation at the quantitative and molecular levels within a
natural *Arabidopsis thaliana* population. Journal of Ecology 110.

Méndez-Vigo, B., N. H. Gomaa, C. Alonso-Blanco, and F. Xavier Picó. 2013. Among- and within-
population variation in flowering time of Iberian *Arabidopsis thaliana* estimated in field and
glasshouse conditions. New Phytologist 197:1332–1343.

Miller-Rushing, A. J., R. B. Primack, D. Primack, and S. Mukunda. 2006. Photographs and herbarium
specimens as tools to document phenological changes in response to global warming. Am J Bot
93:1667–1674.

Miryeganeh, M., M. Yamaguchi, and H. Kudoh. 2018. Synchronisation of Arabidopsis flowering time and
whole-plant senescence in seasonal environments. Sci Rep 8.

Onofri, A., P. Benincasa, M. B. Mesgaran, and C. Ritz. 2018. Hydrothermal-time-to-event models for seed
germination. European Journal of Agronomy 101:129–139.

Penfield, S., and V. Springthorpe. 2012. Understanding chilling responses in Arabidopsis seeds and their
contribution to life history. Philosophical Transactions of the Royal Society B: Biological Sciences
367:291–297.

Poorter, H., F. Fiorani, R. Pieruschka, T. Wojciechowski, W. H. van der Putten, M. Kleyer, U. Schurr, and J.
Postma. 2016. Pampered inside, pestered outside? Differences and similarities between plants
growing in controlled conditions and in the field.

Postma, F. M., and J. Ågren. 2018. Among-year variation in selection during early life stages and the
genetic basis of fitness in *Arabidopsis thaliana*. Mol Ecol 27:2498–2511. Blackwell Publishing Ltd.

Primack, D., C. Imbres, R. B. Primack, A. J. Miller-Rushing, and P. Del Tredici. 2004. Herbarium specimens
demonstrate earlier flowering times in response to warming in Boston. Am J Bot 91:1260–1264.

Ranal, M. A., and D. G. De Santana. 2006. How and why to measure the germination process?

Ratcliffe, D. 1976. Germination characteristics and their inter- and intra-population variability in
Arabidopsis. Arabidopsis Information Service 34–45.

Reich, P. B. 2014. The world-wide "fast-slow" plant economics spectrum: a traits manifesto. Journal of
Ecology 102:275–301.

Ritz, C., F. Baty, and J. C. Streibig. 2015. Dose-Response Analysis Using R. PLoS One 10:e0146021.

Salguero-Gómez, R., O. R. Jones, E. Jongejans, S. P. Blomberg, D. J. Hodgson, C. Mbeau-Ache, P. A.
Zuidema, H. De Kroon, and Y. M. Buckley. 2016. Fast-slow continuum and reproductive strategies
structure plant life-history variation worldwide. Proc Natl Acad Sci U S A 113:230–235.

Samis, K. E., C. J. Murren, O. Bossdorf, K. Donohue, C. B. Fenster, R. L. Malmberg, M. D. Purugganan, and
J. R. Stinchcombe. 2012. Longitudinal trends in climate drive flowering time clines in North
American *Arabidopsis thaliana*. Ecol Evol 2:1162–1180.

Simpson, G. G., and C. Dean. 2002. Arabidopsis, the Rosetta stone of flowering time?

Springthorpe, V., and S. Penfield. 2015. Flowering time and seed dormancy control use external
coincidence to generate life history strategy. Elife 2015.

Stearns, S. C. 1989. Trade-Offs in Life-History Evolution. Funct Ecol 3:259. JSTOR.

Stinchcombe, J. R., C. Weinig, M. Ungerer, K. M. Olsen, C. Mays, S. S. Halldorsdottir, M. D. Purugganan,
and J. Schmitt. 2004. A latitudinal cline in flowering time in *Arabidopsis thaliana* modulated by the
flowering time gene FRIGIDA. Proceedings of the National Academy of Sciences, doi:
10.1073/pnas.0306401101.

Stock, A. J., B. V. McGoey, and J. R. Stinchcombe. 2015. Water availability as an agent of selection in
introduced populations of *Arabidopsis thaliana*: Impacts on flowering time evolution. PeerJ
2015:1–16.

Thomas, B., and D. Vince-Prue. 1997. Photoperiodism in plants. 2nd ed. Academic Press, San Diego, CA,
USA.

Vidigal, D. S., A. C. S. S. Marques, L. A. J. Willems, G. Buijs, B. Méndez-Vigo, H. W. M. Hilhorst, L.
Bentsink, F. X. Picó, and C. Alonso-Blanco. 2016. Altitudinal and climatic associations of seed
dormancy and flowering traits evidence adaptation of annual life cycle timing in *Arabidopsis*
thaliana. Plant Cell Environ 39:1737–1748.

Walker, M. D., R. C. Ingersoll, and P. J. Webber. 1995. Effects of interannual climate variation on
phenology and growth of two alpine forbs. Ecology 76:1067–1083.

Wilczek, A. M., L. T. Burghardt, A. R. Cobb, M. D. Cooper, S. M. Welch, and J. Schmitt. 2010. Genetic and
physiological bases for phenological responses to current and predicted climates. Philosophical
Transactions of the Royal Society B: Biological Sciences 365:3129–3147.

Wilczek, A. M., J. L. Roe, M. C. Knapp, M. D. Cooper, C. Lopez-Gallego, L. J. Martin, C. D. Muir, S. Sim, A.
Walker, J. Anderson, J. F. Egan, B. T. Moyers, R. Petipas, A. Giakountis, E. Charbit, G. Coupland, S.
M. Welch, and J. Schmitt. 2009. Effects of genetic perturbation on seasonal life history plasticity.
Science (1979) 323:930–934.

Wood, S. N. 2006. Generalized additive models: an introduction with R. 1st ed. Chapman and Hall/CRC.

Yee, T. W., and M. Mackenzie. 2002. Vector generalized additive models in plant ecology. Ecol Modell,
doi: 10.1016/S0304-3800(02)00192-8.

Yee, T. W., and N. D. Mitchell. 2006. Generalized additive models in plant ecology. Journal of Vegetation
Science, doi: 10.2307/3236170.

Zettlemoyer, M., R. Conner, M. Seaver, E. Waddle, and M. DeMarche. 2024. A Long-Lived Alpine
Perennial Advances Flowering under Warmer Conditions but Not Enough to Maintain Reproductive
Success. Am Nat 203.

Zhou, D., T. Wang, and I. Valentine. 2005. Phenotypic plasticity of life-history characters in response to
different germination timing in two annual weeds. Canadian Journal of Botany 83:28–36.

 Supplemental Material

Plasticity due to maternal effects

Maternal conditions may be an important source of phenological plasticity; thus we used our

dormancy experiment to examine how variation in flowering time among maternal plants

influences germination. Variation in phenology may not be adaptive in regions where there is

harsh seasonality or environmental conditions are less variable year to year (Cohen 1967).

Similarly, observations of phenology in the wild could be temporally variable because of year-to-

year environmental differences (Walker et al. 1995; Hu et al. 2017; Postma and Ågren 2018) or

genetically diverse individuals germinating from the seedbank (Ratcliffe 1976). Among replicate

maternal individuals in the same controlled experiment, variation in phenology is likely due to

responses to very subtle environmental differences and some developmental stochasticity.

For each ecotype and for each maternal replicate, we used R packages 'drc' (Ritz et al.

2015) and 'drcSeedGerm' (Onofri et al. 2018) to fit a logistic function to germination counts to

estimate three parameters: maximum germination proportion, time to 50% germination, and

slope of the germination curve. A Generalized Linear Mixed Model was fit using the lmer

package in R (Bates et al. 2015) to estimate the influence of relative flowering time of maternal

replicates of each ecotype on germination traits. Because plasticity or responsiveness to

environmental cues may confer greater fitness in some environments and not others (Alpert and

Simms 2002; Baythavong 2011), germination traits and variation were regressed against

maternal flowering time and location of origin. We performed hierarchical clustering of a

distance matrix of phenotypic variation among ecotypes to group ecotypes that reacted similarly

to different stratification treatments.

We found that some ecotypes had more consistent phenology across maternal

lines. For example, DAM1 maternal replicates clustered together in germination across cold

treatments, whereas Yeg-7 replicates did not (Figure S10A). Cold stratification altered

expression of variance among maternal sources. In Yeg-7, for example, maternal lines P vs A

had similar total germination following 3 days or 2 weeks of stratification but different

germination for 0 days stratification. Thus, stratification (or lack of stratification) could

contribute to diverse phenology for similar genetic backgrounds under natural conditions. The

first and second principal components in an analysis of flowering and germination traits

explained 24.55% and 17.85% of the variation respectively, but again maternal replicates of

ecotypes did not always cluster in PC space (Figure S10B). The variance in phenology for an

ecotype was not related to collection date or geographic origin (latitude and longitude, Figure

S10C), suggesting that clinal variation in maternal plasticity effects do not confound our earlier

large-scale analyses comparing breeding values with phenology in nature. Nevertheless, these

effects may further obscure genetic contributions to phenology in nature.

We expected to see more variation among ecotypes from regions with favorable growing

conditions year-round. However, the lack of a geographic association between variance in

germination traits and flowering time among replicate individuals of an ecotype may indicate

geographically consistent selection on the canalization of phenology.

Figure S1. An illustration of the importance of accounting for local mean flowering dates when
testing for correlations between breeding value and observed flowering dates in the presence of
co-gradient variation (correlated plasticity and genetic clines). In both A and B there is co-
gradient variation, but in A the breeding values predict variation in actual natural phenology
(flowering date) while in B the breeding values do not predict local phenology independent of
plasticity and environmental differences among sites.

Figure S2. 42 ecotypes have measures of vernalized, non-vernalized, and field flowering times as

well as primary and secondary dormancy traits (standardized trait units are shown). The latest

flowering accessions tend to be late flowering across all treatments, but primary and secondary

dormancy vary independently from flowering time.

Figure S3. Standardized phenology traits combined across multiple published controlled
experiments combined with our new experiments, using an alternate standardization (compare
with Figure 3). Here, each experiment's values were standardized to mean zero unit standard
deviation.

Figure S4. Date of collection (left panels) and PTUs (photothermal units) at collection (right

panels) of wild plants compared to standardized breeding values for phenological traits from

experiments, here using an alternate standardization (compare to Figure 4). Here, each

experiment's values were standardized to mean zero unit standard deviation. The values shown

for field flowering time experiments were calculated from fall-sown experiments only.

Figure S5. Date of collection (left panels) and PTUs (photothermal units) at collection (right panels) of wild plants compared to standardized breeding values for phenological traits from experiments, here using an alternate standardization and excluding herbarium specimens (compare to Figure 4). Here, each experiment's values were standardized to mean zero unit standard deviation. The values shown for field flowering time experiments were calculated from fall-sown experiments only.

Figure S6. Date of collection (left panels) and PTUs (photothermal units) at collection (right

panels) of wild plants compared to standardized breeding values for phenological traits from

experiments, excluding herbarium specimens (compare to Figure 4). The values shown for field

flowering time experiments were calculated from fall-sown experiments only.

Figure S7. Flowering times and germination rates within experiments do not predict wild

phenology better than average values across multiple experiments. Left, non-vernalized

flowering times at 16°C (using the same criteria to remove collection day and geographical

outliers as in the full set of accessions) Alonso-Blanco et al. (2016) did not significantly predict

collection day (p > 0.2, n = 173). Right, germination rates after 4 days of stratification at 4°C

Martínez-Berdeja et al. (2020) did not significantly predict collection day (p > 0.1, n = 84).

Figure S8: Predicted collection day vs. actual collection day. A) Difference between collection

day and predicted collection day in ecotype accessions is designated by the size of the black

circles. Accessions from Turkey were collected earlier than expected in our model, while

expectations for Western Europe were closer to observed values. Herbarium records on which

the model was built are shown as gray circles. B) Density plots of predicted collection days

(black) and actual collection days (gray). Collections tended to be later than model predictions.

C) Relationship between non-vernalized flowering time and collection day. Both predicted

(black) and observed (gray) collection days are positively related to non-vernalized flowering

times, but the observed collection days have greater range and a lower slope. Model-predicted

flowering times underestimate the variation in collection days, which might be due to actual

collection days picking up plants on the tails of the flowering distribution.

Figure S9. Spatially varying intercept of models of within-population phenological variation.

The spatially varying intercept absorbed most of the phenological variation in collection date and

photothermal units. Red, dashed lines represent areas where collection day was earlier while

green, dotted lines represent areas where collection day was later. Collections are represented as

circles. The model attributes a great deal of variation to geographic location, despite including

phenological breeding values. Elevation was more important in predicting photothermal units

than collection date.

Figure S10: Variation in experimental phenology among maternal lines. A) Early flowering lines

tended to have lower germination under 0 days stratification relative to 3 days or 2 weeks

stratification, and later-flowering lines tended to have longer times to 50% germination.

However, there was variation across traits. Replicate maternal lines of the same ecotype did not

always cluster. B) Maternal lines of the same ecotype did not always cluster together in PCA of

phenology traits. Lines belonging to the same ecotype are represented with the same color,

although colors are not unique across ecotypes. PC1 was positively associated with time to 50%

germination at 0 days and 3 days stratification and flowering time and negatively associated

with time to 50% germination at 2 weeks stratification and max germination under all

treatments. C. Flowering time, collection day, or geographic origin did not explain variance

among maternal lines.

998 Table S1: Model statistics for other variables in models of within-population phenological

999 variation. P-values for the estimates are given in parentheses. Phenology breeding values did not

1000 significantly predict within-population variation in phenology.

Model	Non-Vernalized Flowering Time	Primary Dormancy	FT*Dormancy	Elevation	Adj R^2
Collection Date ~ μ + FT + Elevation	-3.25 (0.35)	NA	NA	**0.0148 (0.006)**	0.883
PTU$^{1/2}$ ~ μ + FT + Elevation	-1.21 (0.75)	NA	NA	**-0.0147 (0.006)**	0.876
Collection Date ~ μ + Dormancy + Elevation	NA	1.98 (0.56)	NA	**0.0126 (0.04)**	0.876
PTU$^{1/2}$ ~ μ + Dormancy + Elevation	NA	2.56 (0.48)	NA	**-0.0181 (0.009)**	0.858
Collection Date ~ μ + FT + Dormancy + Elevation	-8.19 (0.10)	2.19 (0.51)	NA	**0.0120 (0.05)**	0.881
PTU$^{1/2}$ ~ μ + FT*Dormancy + Elevation	-5.41 (0.56)	5.39 (0.43)	-6.86 (0.66)	**-0.0183 (0.009)**	0.864

1001

1002

1003

1004