Does genetic variation in controlled experiments predict phenology of wild plants?

Victoria L. DeLeo[1], David L. Des Marais[2], Claire M. Lorts[1], Thomas E. Juenger[3], Jesse R.

Lasky[1,4,5]

[1]Department of Biology
Pennsylvania State University
USA

[2]Department of Civil and Environmental Engineering
Massachusetts Institute of Technology
USA

[3]Department of Integrative Biology
University of Texas at Austin
USA

[4]PAC Herbarium
Pennsylvania State University
USA

[5]Author for correspondence. Email: lasky@psu.edu

Short running title: *Phenology of wild Arabidopsis*

Acknowledgements

We thank the herbaria that allowed sampling of seed from specimens for generation of natural
inbred lines used in experiments: Real Jardín Botánico de Madrid, Oslo Natural History
Museum, and Komarov Botanical Institute. We are grateful for assistance with experiments from
P. Patel, C. Yim, J. Kizer, T. Xia, V. Meagher, K. Turner, and E. Bellis and for helpful
discussions with L. Burghardt. We thank reviewers for their constructive comments. Funding
was provided by NIH award R35 GM138300 to JRL.

Author Contribution

JRL conceived of the project with TEJ, DLD assisted in experiments, VLD conducted
experiments, and analyzed data, VLD and JRL led writing, all authors contributed to the
interpretation and writing of the manuscript.

Data Accessibility Statement

Original data will be included as supplemental tables. Prior studies from which phenology data
was sourced will be listed in a supplemental table.

Conflict of Interest Statement

The authors declare no conflicts of interest.

Abstract

Phenology and the timing of development are often under selection. However, the relative contributions of genotype, environment, and prior developmental transitions to variance in the phenology of wild plants is largely unknown. Individual components of phenology (e.g., germination) might be loosely related with the timing of maturation due to variation in prior developmental transitions. Given widespread evidence that genetic variation in life history is adaptive, we investigated to what degree experimentally measured genetic variation in Arabidopsis phenology predicts phenology of plants in the wild. As a proxy of phenology, we obtained collection dates from nature of 227 naturally inbred *Arabidopsis thaliana* accessions from across Eurasia. We compared this phenology in nature with experimental data on the descendant inbred lines that we synthesized from two new and 155 published controlled experiments. We tested whether the genetic variation in flowering and germination timing from experiments predicted the phenology of the same lines in nature. We found that genetic variation in phenology from controlled experiments significantly predicts day of collection from wild individuals, as a proxy for date of flowering, across Eurasia. However, local variation in collection dates within a region was not explained by genetic variance in phenology in experiments, suggesting high plasticity across small-scale environmental gradients or complex interactions between the timing of different developmental transitions. While experiments have shown phenology is under selection, understanding the subtle environmental and stochastic effects on phenology may help to clarify the heritability and evolution of phenological traits in nature.

Introduction

The predictability of trait variation from genetic variation, *i.e.* trait heritability, is a key quantity determining the rate of trait evolution and the ability of populations to adapt to changes in selection (Lush 1937; Gomulkiewicz and Houle 2009). Plant biologists have long relied on common garden experiments to estimate breeding values and heritability for a range of traits (Mazer and LeBuhn 1999). However, natural populations often occur in highly heterogeneous environments to which organismal traits may respond plastically. As a result, heritability in natural populations may be much lower than those in common gardens. Furthermore, even field common gardens might fail to accurately recreate natural selective pressures if the direction and magnitude of selection varies in time (Karrenberg & Widmer 2008). As an alternative to common gardens, heritability can be estimated using pedigrees or genomic data from wild individuals (Gienapp et al. 2017). However, these models rely on assumptions of certain genetic architectures (Mitchell-Olds and Rutledge 1986) and the resulting heritability estimates are not often compared with common garden estimates of trait variation (but see (Weigensberg and Roff 1996)).

Phenology, or the timing of the developmental transitions between an organism's life stages, directly and indirectly influences plant fitness and selection by determining the traits that are expressed at any point throughout the year (Donohue 2005). In seasonally variable environments, traits such as flowering time and dormancy can ameliorate harsh abiotic conditions by timing dormant periods to coincide with unfavorable seasons (e.g. drought escape Ludlow 1989; Lawrence-Paul and Lasky 2024). Flowering, vegetative phase change, and germination timing often show geographic clines, suggesting their importance in adaptation to local environments (Baughman et al. 2019; Martínez-Berdeja et al. 2020; Lawrence-Paul et al. 2023, 2025). Common garden studies often find that genetic variation in individual components of phenology are under selection and have further demonstrated variable strength and direction of selection across field sites (Hall and Willis 2006; Fournier-Level et al. 2013; Ågren et al. 2017; Gamba et al. 2024b). However, the duration of individual developmental transitions (e.g. flowering time) might be loosely related to phenology in nature due to variation in prior developmental transitions (e.g. germination) (Burghardt et al. 2015). While phenology is determined by both genetic factors and plastic responses to environment (Amasino 2004; Andrés

and Coupland 2012; Auge et al. 2018), the relative importance of these two sources of

phenotypic variance in the timing of individual developmental transitions determining phenology

in nature is unclear, even in model systems.

Much of our understanding of the genetic basis of plant phenology comes from studies in

the model *Arabidopsis thaliana* (hereafter Arabidopsis). Common garden trials in both lab and

field settings have uncovered genetic loci contributing to much of the observed variation in

dormancy and flowering time (Juenger et al. 2005; Atwell et al. 2010; Brachi et al. 2010;

Fournier-Level et al. 2013; Ågren et al. 2017). Flowering time and seed dormancy are

determined by complex, overlapping gene networks, leading to pleiotropy involving these traits

(Simpson and Dean 2002; Wilczek et al. 2010; Auge et al. 2019). These traits are also plastic and

show genotype by environment interactions (Juenger et al. 2005; Zhou et al. 2005; Wilczek et

al. 2009; Lorts and Lasky 2020): germination responds to temperature, photoperiod, moisture,

and nutrient availability (Huang et al. 2010, 2018; Penfield and Springthorpe 2012; Footitt et al.

2013; Kenney et al. 2014), while flowering responds to multiple cues such as temperature and

daylength (Thomas and Vince-Prue 1997; Lempe et al. 2005; Balasubramanian et al. 2006).

The broad phenological variation described in controlled environment experiments is

observed in wild populations as well (Ratcliffe 1976; Simpson and Dean 2002; DeLeo et al.

2020). Arabidopsis can exhibit the life history of a winter annual, germinating in the fall,

overwintering as a rosette, and flowering in the spring (Figure 1). However, Arabidopsis can also

germinate and flower in a single growing season. These shorter-lived plants tend to be facultative

— flowering in spring, summer, or fall — and in some regions this rapid life cycle enables

multiple generations within a year. This variation in life histories occurs in many annual plants

(Baskin and Baskin 1988) and is made possible, in part, by a range of flowering times and

germination traits that can vary independently from each other (Marcer et al. 2018; Martínez-

Berdeja et al. 2020). In Arabidopsis, geographic clines in genetic values for phenology, clines in

allele frequencies of phenology QTL, and evidence of selection on phenology QTL support the

conclusion that phenology is adapted to local environmental conditions in the wild (Caicedo et

al. 2004; Stinchcombe et al. 2004; Samis et al. 2012; Fournier-Level et al. 2013; Ågren et al.

2017; Debieu et al. 2018; Exposito-Alonso et al. 2018; Lasky et al. 2018, 2024; Gamba et al.

2024a) (Figure 1). Yet, even within geographic proximity one can find genotypes with

substantial genetic differences in flowering time (Jones 1971; Alonso-Blanco et al. 2016;

Méndez-Vigo et al. 2022).

Life Cycles Across the Range   **Genetic Clines, Reaction Norms**

(A) Colder Climate (e.g. Scandinavian) Life Cycle

- Late Flowering
- High Germination in Experiment
- Lowest PTU at Flowering/Collection Date

PTU at Flowering/Collection Date

4°C

Jan.   Dec.
        Time of Year

Temperature

(D) Genetic Clines in Flowering Time Across Climates of Origin

Flowering Time

Genetic Cline

Cold   Warm
        Climate of Origin

(B) Warm Climate Life Cycle (e.g. Germany)

- Early Flowering
- High Germination in Experiment
- Highest PTU at Flowering/Collection Date

PTU at Flowering/Collection Date

4°C

Jan.   Dec.
        Time of Year

Temperature

(E) Plasticity in Flowering Time Across Growth Environments

Flowering Time

Plasticity

Cold   Warm
        Growth Environment

(C) Warmer Climate (e.g. Mediterranean) Life Cycle

- Early Flowering
- Low Germination in Experiment
- Low PTU at Flowering/Collection Date

PTU at Flowering/Collection Date

4°C

Jan.   Dec.
        Time of Year

Temperature

(F) Genetic Clines in Germination Across Climates of Origin

Base Germination (1/Primary Dormancy)

Genetic Cline

Cold   Warm
        Climate of Origin

127   ***Figure 1.*** *Genotype and environment likely influence phenology of individual Arabidopsis plants*

128   *across its geographic range in the wild. Genetics and environmental cues may determine how*

129   *long a plant remains in vegetative growth (green) or how long a plant remains dormant as a*

130   *seed (brown circles), with three representative life cycles shown from Scandinavia to the*

131   *Mediterranean (A–C), with shaded areas showing accumulation of photothermal units (PTUs) in*

There are several mechanisms that may explain why phenology observed in controlled experiments shows poor correlation with phenology observed in nature (Wilczek et al. 2009). While genotype likely influences the phenology of an individual plant in the wild, the dependence of sequential developmental transitions on each other, as well as plasticity in their expression, may limit the translation of genetic values of single stage to natural phenology. First, there are interactions between germination and flowering time, both because of a shared genetic basis and because the timing of early life transitions influences environmental exposures in later life stages (Chiang et al. 2009; Huang et al. 2010; Springthorpe and Penfield 2015; Huo et al. 2016). Secondly, there is stochasticity in individual germination and flowering time (Jimenez-Gomez et al. 2011; Abley et al. 2020). Thirdly, spatial environmental variation is extensive in nature and genotype-environment interactions have major impacts on phenology (Sasaki et al. 2015). Finally, maternal effects on phenological transitions may be strong, as reported extensively for seed dormancy (Boyd et al. 2007; Chiang et al. 2013; Burghardt et al. 2016; Huang et al. 2018). As a result, flowering time loci identified in common garden lab or field experiments might have loose relationships with variation among natural individuals (Chiang et al. 2013); genotypes differing in phenology in one set of conditions may express the same phenology in another set of conditions (Wilczek et al. 2009; Burghardt et al. 2016).

Here, we compare phenological observations of wild individual plants from natural history collections (herbaria and seed stock centers) with plants in controlled environment experiments. Collection dates of natural history accessions have been demonstrated as accurate estimates of phenology (Miller-Rushing et al. 2006; MacGillivray et al. 2010; Davis et al. 2015; Ramirez-Parada et al. 2022). Across Eurasian Arabidopsis, we showed previously that herbarium collection dates were accelerated by approximately one week for each degree increase in spring temperatures and that there was a latitudinal cline in mean collection dates from early March to late June (DeLeo et al. 2020). Arabidopsis are naturally inbred (Jones 1971), thus we can compare collection dates of wild individuals with phenology of nearly genetically identical descendants in controlled experiments. This comparison provides a window into how genetic variation shapes phenology in natural populations against the forces of plasticity and GxE.

Natural history collections can be used to characterize the climate conditions during development

preceding collection (Bontrager et al. 2025b). We estimated phenology as collection date, and

using the climate conditions preceding collection, we also estimated phenology as accumulated

photothermal units (PTUs) at collection (Burghardt et al. 2015). PTUs allow us to account for

weather effects on development and in essence allow estimation of how far into growing seasons

plants were collected.

Predictions for Analyses in This Study

(A)   Clines, Co-Gradient Plasticity, and Local
Genetic Variation in Flowering Time Cause:

(C)   Correspondence Between Early Flowering
and Spring Germination Causes:

(D)   Germination Correlation with Flowering Time,
Clines in Flowering Time, and Germination Cause:

(B)   High Germination for Spring Germinants Causes:

Figure 2: Existing knowledge of clines in flowering time and germination combined with plastic acceleration of flowering in warmer temperatures (see Figure 1) lead to our predictions (A–D here) of how flowering time and base germination rates in controlled experiments will correspond to phenology in natural wild plants.

We developed three hypotheses for how the phenology of individual plants in nature may relate to genetic variation in phenological traits, illustrated in Figures 2-3:

H1. We predict that Arabidopsis plants collected earlier in a natural growing season will have early flowering and low germination genetic values in controlled environments. We hypothesize that this pattern is due to co-gradient variation in flowering time along temperature gradients (i.e. when genetic clines and plastic responses to environment are positively correlated) (Levins 1968). Specifically, genetic clines in flowering time (here, early flowering in warmer climates) and germination rate, combined with plastic acceleration of flowering in warmer conditions (Figure 1), leads to range-wide positive relationships between genetic values for flowering time (Figure 2A), germination rate (Figure 2B), and collection date (as a proxy for flowering date in the wild).

H2. For cumulative photothermal units (PTU) at collection, we predict that spring annuals with high germination rates and early flowering times will have been collected later in growing seasons (high PTU) than winter annuals with later flowering genetic values (Figure 2C, 2D).

H3. Within local regions, environmentally induced phenological variation (i.e., plasticity associated with different timing of spring warming) is likely lower than that seen across the species range. Thus we predict that flowering time genetic values and germination rates would be positively related to local within-region variation in collection dates largely due to genetic effects (Figure 3).

Partially Supported by Existing Literature:

Legend:

- 🔴 Early flowering genotypes in experiments
- 🟣 Intermediate flowering genotypes in experiments
- 🔵 Late flowering genotypes in experiments
- ◇ Local mean date

(A) Genetic cline in flowering time.
Colors change between environments.

Cogradient plastic response to environment, e.g. purple genotype flowers earlier in warm environments.

Untested Competing Hypotheses:

(B) Genetic values predict local variation in flowering date. Flowering date has high h² within natural populations. Residuals from local mean date are correlated with genetic variation in experiments.

The purple genotype has a **late flowering residual from the local mean**, and flowers later than the orange genotype in experiments.

(C) Genetic values _do not_ predict local variation in flowering date. Flowering date has low h² within natural populations. Residuals from local mean date are not correlated with genetic variation in experiments.

The purple genotype has an **early flowering residual from the local mean**, and flowers earlier than the orange genotype in experiments.

Figure 3. An illustration of hypotheses for local genetic variation in phenology within regions and the relationship with phenology of wild plants. In the presence of co-gradient variation (correlated plasticity and genetic clines, panel A) it is important to account for local mean collection dates (diamonds) when testing for correlations between genetic values for phenology versus observed collection dates). In both (B) and (C)there is co-gradient variation, but in (B) the genetic values for phenology traits (colors) predict local variation in natural phenology (residuals from local mean collection date) while in (C) the genetic values (colors) do not predict local phenology (residuals from local mean collection date).

Materials and Methods

Natural genotypes

We first conducted an experiment on a set of 101 naturally inbred genotypes ("ecotypes")

selected based on known collection dates either from the herbarium specimen label or recorded

in the Arabidopsis Biological Resource Center https://abrc.osu.edu/ database. 47 of these

ecotypes had not been studied in previous flowering time experiments and 75 had not been

included in previous germination experiments. Seed was ordered from ABRC (94 accessions) or

was germinated from herbarium sheets (7 accessions from Real Jardín Botánico Madrid,

Komarov Botanical Institute, and Oslo Natural History Museum).

Seeds from herbarium specimens were cold-stratified in tap water at pH 7 and placed at

4°C for 7 d. Seeds were then directly sown into damp Fafard germination mix and grown in

Conviron growth chambers under 14/10hr, 22/18°C days/nights. Four herbarium accessions

germinated using this protocol. In a second round of germination, we surface sterilized seeds

using standard protocols and then cold-stratified, as above. We plated these sterilized seeds onto

MS+Gamborg's vitamins + agar plates containing 1% sucrose and 10uM GA4 and then placed

them in growth chambers, as above. Three additional herbarium accessions flowered using this

protocol; these were transplanted to Fafard Germination mix. To induce flowering for seed

collection, plants from both rounds of germination were exposed to 30d of 4°C with 10/18hr

day/night cycles.

Flowering and germination experiments

Ecotypes were grown in common conditions prior to the flowering time experiment, and

flowering time replicates were descended from a single mother plant. For each ecotype, three

replicates were grown in separate pots. Seeds were stratified at 4°C for 5 days before sowing in

pots. Each pot was thinned to a single individual after the emergence of the second set of true

leaves. Plants were grown at 22°C under 16h days of fluorescent light in a walk-in Conviron

growth chamber (model MTPS144). Day of bolting and day petals appeared were both recorded

as measures of flowering time.

Seeds from each replicate maternal plant in the flowering time experiment were collected

and stored separately in dry conditions until the germination trial. Non-stratified seeds were kept

in dry boxes at room temperature. For each treatment, forty seeds from each parent plant (or as

many seeds as were available for replicates with low fecundity) evenly divided across 2 plates

were sown on filter paper in petri dishes and germinated at 23/18°C during day/night with

constant 16h daylength in a Conviron growth chamber. In total, 1,752 plates and >50k seeds

were assayed.

Seeds were subjected to cold stratification at 4°C in the dark for 3 treatment lengths: 2

weeks, 3 days, and 0 days. Cold stratification can break primary dormancy, however 2 weeks of

chilling can prompt secondary dormancy in the seedbank (Penfield and Springthorpe 2012). The

difference in germination rate between 3 days and 0 days of stratification can therefore indicate

primary dormancy while the difference in germination rate between 3 days and 2 weeks of

stratification may indicate secondary dormancy. As a caveat, lower germination after 2 weeks of

stratification could be due to other, unmeasured negative effects on germination, such as

bacterial infection. We staggered the planting so that all the plates came out of stratification on

the same day. The number of seeds that had germinated in each plate was recorded on days 1, 3,

5, 10, 14, 21, and 28. Seeds were considered germinated if the radicle was visible. Because

maternal plants flowered at different times (and seeds were harvested when maternal plants

senesced), germination rates for this experiment may be influenced by the length of the time

between flowering and planting and any after-ripening that may have occurred. We also tested

for these maternal effects among individuals of the same inbred line (Supplemental Material).

Phenology from published experiments

We searched the literature for experiments on Arabidopsis that measured flowering time or

germination traits across different natural inbred lines (commonly referred to as Arabidopsis

"ecotypes"). The minimum number of ecotypes in any single experiment was 17. Our final set

used data from 38 previous studies (31 included some measure of flowering time, 15 included

germination, Table S1) that, combined with our new experiments described above, included over

3,000 ecotypes for 86 flowering time experimental conditions and 66 germination experimental

conditions, although all ecotypes were used in only a subset of the trials and only 291 ecotypes

had a reliable date of original collection from the wild.

We used this dataset of phenology measurements from the literature to create an estimate

of genetic variation in flowering time and dormancy among ecotypes. We sought to gain

statistical power by combining data from different experiments. To make phenotypes

comparable, we standardized across treatments and experiments, transforming each flowering

time experiment such that the earliest flowering accession had a value of 0 and the latest

flowering accession had a value of 1.

We averaged standardized experimental flowering times across experiments, keeping

vernalized, non-vernalized, and field experiments separate. Here, we use 'vernalization' to

describe extended cold treatments applied to rosettes. Non-vernalized growing conditions

uncover genetic variation in flowering time due to vernalization requirements that is masked

under vernalized or fall-sown field conditions and may be important for determining phenology

and life history in the wild (Wilczek et al. 2009). However, Arabidopsis plants growing in many

natural settings are expected to experience changes in temperature and photoperiod that would be

more similar to vernalized and field experiments (Li et al. 2010). By keeping the three treatments

separate, we could test whether non-vernalized flowering time or vernalized flowering time was

more predictive of wild phenology. Field flowering time was handled in two distinct manners:

first, aggregated across all seasons, which may lead to less coherent estimates of field genetic

flowering time if seasonal differences lead to meaningful differences in flowering time, and

second, broken down further into spring (4 treatments over 2 experiments), summer (4

treatments, 1 experiment), and fall plantings (23 treatments, 9 experiments). To control for

regional bias, this aggregation was performed both for all experiments and without experiments

that tested only ecotypes collected from a single country.

Data collection approaches for estimating germination rate and dormancy were more

heterogeneous than for flowering time. Depending on the experiment, dormancy was reported as

the number of days after planting until a set percentage of germination was reached, the

percentage of seed germinated a given number of days after planting, the number of days of

storage until a set percentage of germination, or the germination rate after a given number of

days of storage. Because of the difference in metrics across experiments, dormancy values were

standardized by rank within each experiment with zero indicating low to no dormancy and one

indicating high dormancy. Measures that reported a percentage germinated were ranked in the

opposite direction from measures that reported number of days until germination or the number

of days of storage before 50% germination. We found that standardized rank-based metrics that

increase with dormancy (such as days to 50% germination) were correlated with rank based on

metrics that decrease with dormancy (such as percentage of seeds germinated after a set number

of days, Pearson's $r = 0.485$).

300   These experiments captured variation in primary dormancy, or recalcitrance to germinate

301 immediately after harvest. Secondary dormancy, or dormancy induced when a seed experiences

302 conditions unfavorable to germination, has been studied in experiments that measured an

303 increase in dormancy during storage. Primary and secondary dormancy could lead to different

304 phenological and ecological outcomes (Martinez-Berdeja et al. 2020), so we averaged across

305 experiments that measured primary or secondary dormancy separately to estimate each of these

306 two traits. Secondary dormancy was not included in the Generalized Additive Models described

307 below. Maternal conditions may be an important source of phenological plasticity; thus we also

308 used our dormancy experiment to examine how variation in flowering time among maternal

309 plants influences germination (Supplemental Material).

311 *Wild phenology*

312 The date of collection (for ecotypes maintained by ABRC) or collection date (for ecotypes grown

313 from a known herbarium record) was used as an estimate of reproductive phenology of

314 individuals in the wild (Primack et al. 2004; Davis et al. 2015). We excluded records from

315 regions where Arabidopsis has been recently introduced, like North America and Japan.

316 Accessions from outside the native range or collected after the 320[th] day of the year (which we

317 deemed likely errors based on location, sometimes due to intentional plantings, and were greater

318 than 2 standard deviations from the mean) were removed. Our geographic limits also excluded

319 island accessions to the south of the Mediterranean, e.g. Cape Verde Island.

320   We hypothesized that genetic differences in wild phenology may be more apparent if we

321 account for spatiotemporal environmental fluctuations causing plasticity. Therefore, we

322 calculated photothermal units (PTUs) for each accession from the day of collection using

323 monthly climate time series data from CRU (Harris et al. 2014), beginning with January 1 of

324 each year, following the methods of (DeLeo et al. 2020) and (Burghardt et al. 2015). PTUs

325 integrate the temperature and light experienced by a plant at a given location through the

326 growing season and therefore may capture environmental cues relevant to phenology and better

327 describe genetic variation in development (Wilczek et al. 2009; Brachi et al. 2010). We only

328 sought to use PTUs to estimate how far into growing seasons plants were collected; we did not

329 attempt to estimate PTUs since germination since germination timing was unknown. The models

described below use the square root of PTU because the resulting distribution was closer to

normal than the log transformation.

Statistical comparison of collection date in the wild versus phenology in experiments

H1: Range-wide comparison of phenology genetic values vs. collection date

We tested if genetic variation in normalized phenology measured on naturally inbred lines

explained variation in the wild phenology of the parent of the line (Figure 2), using linear

regression between the normalized phenology genetic values (flowering time and dormancy

rank) and collection day. Vernalized, non-vernalized, and field flowering times were modelled

separately, because the genetic variation in phenology uncovered by each treatment could relate

to wild phenology in different ways. Under the hypothesis that vernalization and field conditions

better recreate environments that a plant would experience at their geographic origin, these

genetic flowering time measures should be more positively related to wild flowering time.

Likewise, non-vernalized flowering times may better recreate the original temporal niches of

summer annuals and thus be positively related to flowering time in these ecotypes. However, it is

also possible that long non-vernalized flowering times indicate obligate winter annuals. In these

plants, non-vernalized flowering times would be negatively related to wild phenology since later

flowering times in non-vernalized experiments would indicate plants that overwinter and are

collected early the next year (Figure 2).

We also investigated how range-wide differences in wild phenology related to regional

differences in genetic trait values using a larger sample size, taken from a broader collection of

ecotypes than just those with documented collection dates. This comparison included stock

center ecotypes with experimental phenology data and known location-of-origin, but no recorded

date of wild collection, and herbarium specimens for which we had not germinated seed and

grown plants to measure traits in experiments. To do so, we first estimated wild flowering time

for each stock center ecotype in experiments using herbarium records near the ecotype collection

location. These wild flowering times were estimated from a previously published generalized

additive model (GAM) with spatially-varying intercepts of herbaria collection dates from 2,655

Eurasian Arabidopsis records used in (DeLeo et al. 2020), which included year of collection as a

nuisance variable, as collection date has changed over time across the range of Arabidopsis

(DeLeo et al. 2020). Values of the smooth intercept surface were extracted at the coordinates of

stock center ecotypes having an experimentally measured flowering time. These estimated wild

flowering times were regressed against genetic values for flowering times under vernalized, non-

vernalized, and field experiments and rank primary dormancy. In addition, for these same data

we performed Spearman rank correlation between estimated wild flowering times and phenology

genetic values.

H2: Range-wide comparison of phenology genetic values vs. PTU

To account for plasticity in phenology due to differences among locations in the timing and

progression of growing seasons, we also tested the relationship between phenology genetic

values and the estimated PTUs at the time of collection from the wild. We conducted the same

analyses across the range of Arabidopsis as described in H1, except that the PTU at collection

was used instead of collection date. Specifically, we conducted linear regression between

normalized genetic values with PTU at collection and we also conducted linear regression

between genetic values and the PTUs at collection of nearby herbarium specimens.

H3: Local within region variation

Next, we tested for the effects of genetic variation in flowering time and dormancy on

phenological variation *within* local regions in the wild. By using a model term to first account for

geographic, *among* region, variation in mean flowering time, we aimed to isolate the remaining

local, *within*-region variation (Figure 3). This analysis, in essence, asks whether genetic variation

in individual phenological stages explains natural phenological variation within local regions.

Generalized Additive Models (GAMs) allow for model parameters to vary smoothly across space

and thus can capture spatially varying patterns, e.g. in local average collection date (Yee and

Mackenzie 2002; Yee and Mitchell 2006). We built a GAM for the dependent variable of

collection date (*Y*) at location *j* of individual *i* which included covariates of experimental

flowering time (β_1), dormancy (β_2), elevation (β_3) and a spatially varying intercept (μ_j):

Equation 1.

$Y_{ij} = \mu_j + \beta_1\text{Flowering Time}_{ij} + \beta_2\text{Primary Dormancy}_{ij} + \beta_3\text{Elevation}_j + \varepsilon_{ij}$

The smooth spatially varying intercept term μ_j estimates spatial variation in mean

collection date (e.g., due to environmental gradients). The upper limit on degrees of freedom for

the spatially varying intercept was increased to 45 following the recommendations of Wood

(2006). In models that allowed for higher degrees of freedom, the effective degrees of freedom

did not meaningfully increase. Elevation was included because of its known importance to

flowering times and spring onset (Vidigal et al. 2016; Gamba et al. 2023) and the high resolution

of elevation data compared to smooth variation in GAM parameter surfaces. We fit GAMs using

the 'gam' function in the 'mgcv' package in R (Wood 2006) using restricted maximum

likelihood (REML), although Generalized Cross Validation returned similar estimates. Model

fitting allowed for penalization of smooth terms to 0 so that uninformative covariates could be

removed from the equation. Residuals were plotted using the 'gam.check' function in 'mgcv'

(Wood 2006), and one ecotype (Nok-10) was removed from our analyses that was an extreme

outlier based on its residuals. Spatial variation in β coefficients for flowering time and dormancy

covariates (Eq 1) were not significant, so a model using a constant coefficient was used in our

analyses. Field, vernalized, and non-vernalized flowering times were tested in separate versions

of the model and tested different hypotheses with regards to wild phenology. To compare among

the three measures of flowering time, a version of Equation 1 was fit on a subset of ecotypes that

had all three flowering time measures, and Akaike's Information Criterion (Akaike 1974) was

compared.

In the wild, both flowering time and dormancy contribute to phenology. Thus, we also

tested whether including interactions between flowering time and primary dormancy (β_4)

improved the model:

Equation 2.

$Y_{ij} = \mu_j + \beta_1 \text{Flowering Time}_{ij} + \beta_2 \text{Primary Dormancy}_{ij} + \beta_3 \text{Elevation}_j + \beta_4 \text{Flowering Time}_{ij} *$

$\text{Primary Dormancy}_{ij} + \varepsilon_{ij}$

Finally, given the importance of plasticity in response to temperature in the timing of

germination and flowering time, PTU might better capture genetic variation in phenology in the

wild (Figure 2). Therefore, we also tested the models above using PTU in place of collection day

to correct for climate.

Results

Genetic correlation among phenology traits in controlled experiments

Across all compiled experiments (both previously published and our new experiments) mean flowering times of ecotypes measured in vernalized and non-vernalized experiments were strongly positively correlated (Pearson's $r = 0.90$, p < 0.001). Although fall-sown field trials might be expected to expose plants to cold seasonal temperatures that give vernalization cues, standardized flowering time genetic values in both non-vernalized and vernalized experiments were similarly correlated to published field experiment values ($r_{non-vernalized} = 0.77$, $r_{vernalized} = 0.72$, p < 0.001 for both). Primary and secondary dormancy were negatively correlated, but not significantly so ($r = -0.15$, p $= 0.11$). Despite known interactions between flowering time and dormancy due to seed maturation environment and pleiotropy of causal loci, only non-vernalized flowering time was significantly correlated with primary dormancy ($r = -0.17$, p $= 0.02$, Figure 4; for a heatplot of all phenotypes, see Figure S1). These correlations were qualitatively unchanged when using an alternate standardization method across experiments (Figure S2).

Figure 4. Standardized phenology traits combined across multiple published controlled

experiments combined with our new experiments. Standardized flowering times were correlated

across all three types of treatments (vernalized, non-vernalized, field), but they were weakly

correlated with dormancy. See Methods for details on standardization method.

H1: Comparing wild phenology with genetic values in experiments across the range

Experimental genetic values for phenology traits were modest predictors of collection day in

simple linear models, with the only statistically significant predictors being non-vernalized

flowering time (estimated slope = 27.7 days/standardized flowering time, $r^2 = 0.09$, $p < 0.001$)

and primary dormancy (-17.1 days/standardized rank dormancy, $r^2 = 0.02$, $p = 0.02$). Flowering

time genetic values under vernalized or field conditions did not significantly predict day of

collection with a linear model ($p > 0.05$) (Figure 5). Flowering time and germination measured

within individual experiments did not predict wild collection day better than our averaged

genetic values, suggesting some power was gained by combining individual experiments into

standardized values (Figure S3). These correlations were qualitatively unchanged when using an

alternate standardization across experiments and when excluding herbarium specimens (Figures

S4–S6).

Figure 5. Date of collection (left panels) and PTUs (photothermal units) at collection (right panels) of wild plants (y-axes) compared to standardized genetic values for phenological traits from experiments (x-axes). Collection day of wild plants was positively related to non-vernalized flowering time in experiments. Collection day was negatively related to primary dormancy. The values shown for field flowering time experiments were calculated from fall-sown experiments only.

Flowering time and dormancy genetic values predict regional variation in collection date of

nearby herbarium specimens

Our model of collection date, above, was limited to the 227 ecotypes with both a known
collection date and experimental phenology data. To expand our predictions across more of the
Arabidopsis range, we tested whether genetic variation in flowering time in experiments predicts
local mean collection dates of herbarium specimens across the landscape. First, we found that
local mean collection dates across a landscape from >2500 herbarium collections were positively
correlated with actual collection days of individual stock center ecotypes (which were not used in
fitting the model, $\rho = 0.51$, $p < 0.001$ Figure S7), validating these local predictions of phenology
in the wild based on herbarium collections.

Figure 6. Genetic variation in flowering time experiments predicts collection dates of nearby

herbarium specimens. A) There is geographic variation in collection dates as shown by

estimated mean date from GAMs with spatially varying means, with earlier collections (light

green) around the Mediterranean (DeLeo et al. 2020). *B) Local mean collection dates were*

positively correlated with genetic variation in vernalized flowering time experiments (p < 0.01),

and all ecotypes from locations with earlier mean collection dates in the wild (largely

Mediterranean) had rapid flowering genetic values. Primary dormancy genetic values were

negatively correlated with local mean collection dates (p < 0.001).

As with the collection dates of the original maternal sources of stock center lines, genetic

values for non-vernalized flowering times significantly but modestly predicted local mean

herbarium collection dates ($r^2 = 0.10$, $p < 0.001$). For comparison, latitude was a better predictor

of mean herbarium collection date ($r^2 = 0.37$, $p < 0.001$). These patterns were similar for

vernalized ($r^2 = 0.07$, $p = 0.002$, Figure 6), non-vernalized ($r^2 = 0.10$, $p < 0.001$), and field

measurements (all seasons grouped, $r^2 = 0.02$, $p = 0.13$, although not significant in this last case).

These results show that range-wide genetic variation in phenology measured in controlled

environments likely corresponds to actual phenology of plants in nature. However, because

plants in nature from different regions experience different environments, these patterns cannot

determine the degree to which genetic variation causes the variation in natural phenology as

opposed to genetic variation being correlated with environmental variation that causes

phenological plasticity in nature (i.e. co-gradient variation, Figure 1).

H2: Range-wide comparison of phenology genetic values vs. Photothermal Units

Because environmental differences among locations can influence phenology (Fournier-Level et

al. 2013), we also calculated PTU at collection and compared to these values to traits from

experiments. However, PTU at collection was not significantly predicted by any phenology trait

genetic values (Figure 5). Thus, while ecotypes with later flowering time genetic values tend to

be collected later in the year, these later flowering ecotypes are not collected at higher PTU, i.e.

later in the local growing season. The fact that phenology genetic values are correlated with date,

but not PTU at collection, is consistent with the hypothesis that geographic climate variation

maintains genetic clines in flowering time and germination, while plastic acceleration of

flowering in warmer conditions (Figure 1) also contributes to the breeding value-collection date

correlation (co-gradient variation).

H3: Variation in phenology genetic values does not predict local, within-region variation in wild

collection dates

We next examined whether genetic values could explain variation in natural phenology *within*

regions (Figure 3). We used GAMs that included spatially varying intercepts and an elevation

covariate to account for the among region variation in environmental effects on collection dates, leaving local variation to be potentially explained by genetic variation. However, we found that normalized flowering time (slope of non-vernalized flowering time = -8.2, p = 0.098) and dormancy (slope of dormancy = 2.2, p = 0.51) were not significantly related to day of collection in this GAM (Equation 1), although elevation effects were significant (slope of elevation = 0.012, p = 0.049, Figure 7, Table S2). In comparing models using vernalized, non-vernalized, and field experiments, we found that standardized non-vernalized flowering times led to lower AIC compared to other flowering time measures but higher AIC than a model without flowering time at all, although the difference was slight (difference < 2 for an AIC of 530). We used non-vernalized flowering times in the final models, in part because there were more ecotypes with non-vernalized flowering times (184 for non-vernalized vs 113 for vernalized or 78 for field).

Figure 7. Top: Spatial variation in date of collection for 227 ecotypes with known collection dates. A) Color represents the intercept value for collection day at a location after fitting a linear (i.e. non-spatially varying) relationship with elevation, experimental primary dormancy

and non-vernalized flowering time. Much of the variation in collection date was explained by

the spatially varying intercept rather than being explained by experimental flowering time. B)

Residuals of a GAM predicting collection date using only intercept and elevation parameters,

compared to non-vernalized flowering time, primary dormancy, or fitted collection dates.

Although we found that flowering time, dormancy, and the interaction between the two were not significantly related to collection day, including phenology genetic values did lower the AIC and slightly increase the deviance explained by the model. This suggests that dormancy and flowering time could be related to variation in phenology within regions. However, most of the variation was described by the spatially varying intercept –a model with only a spatially varying intercept had a deviance explained of 80.1% – which could account for geographic genetic clines in phenology and plastic responses to geographic climate gradients.

As with the linear models, we also tested whether genetic variation in phenology in experiments could explain local variation in PTU of collection date in nature. Again, we found that flowering time and dormancy were not significantly related to wild PTU at collection in GAMs with spatially varying intercepts. Elevation was negatively related to PTU, indicating that higher elevation plants were collected earlier in growing seasons, reflecting a potential winter annual strategy (Figure 1, slope: -0.018 $\sqrt{PTU}$/m, $p < 0.01$). Additionally, elevation was positively correlated with date of collection (slope: 0.012 days/m, $p < 0.05$).

Discussion

Plant phenology comprises multiple traits that contribute to fitness. In Arabidopsis, flowering time and germination can vary independently to create a landscape of possible life histories across environments (Debieu et al. 2013; Marcer et al. 2018; Martínez-Berdeja et al. 2020). This variation in phenology is likely partly maintained by selection, given that the traits vary geographically in association with climate (Fournier-Level et al. 2011; Vidigal et al. 2016; Exposito-Alonso 2020) and biotic pressures (Lyons et al. 2015; Davila Olivas et al. 2017), that QTL show evidence of local adaptation (Gamba et al. 2023; Lasky et al. 2024), and that the timing of individual phenological transitions is often correlated with fitness in experiments

(Korves et al. 2007; Stock et al. 2015). Standing genetic variation is necessary for the timing of

developmental transitions to evolve in response to selection by local environments, but little is

known about the heritability of phenology in natural, wild individuals. Furthermore, because

multiple developmental transitions influence phenology, the effects of variation in the timing of

one stage might be compensated by (or, in under-parameterized experiments, confounded by)

variation in the timing of other stages. Given widespread evidence that genetic variation in life

history is adaptive, we investigated to what degree experimentally measured genetic variation in

Arabidopsis phenology predicts phenology of plants in the wild, based on collection dates of

natural history records.

The influence of genotype and environment on flowering time and dormancy have been

well studied experimentally in Arabidopsis. Yet, common garden and controlled environment

experiments must be designed thoughtfully to highlight genetic differences between ecotypes

that are relevant to selection in natural environments (Karrenberg and Widmer 2008). While

experimental design has increasingly recognized the importance of field conditions to acquire a

measure of phenology that is more representative of nature (Wilczek et al. 2009; Anderson et al.

2012; Brachi et al. 2013; Poorter et al. 2016), there has been little comparison of controlled

experiments, field or otherwise, to phenology in wild, naturally cycling individual plants. We

found that flowering time and dormancy genetic values are related to date of collection and

capture variation in phenology primarily *among* populations in the wild, but explain little

variation *within* regions/populations.

Genetic variation in flowering time and dormancy predicts variation in wild phenology among
regions across the species range

Across the species range, both primary dormancy and non-vernalized flowering time

significantly predicted collection dates, though much variation remained unexplained. These

phenological transitions are not independent of each other. As seen in individual experiments

(Martínez-Berdeja et al. 2020), we found that the average genetic values for primary dormancy

and non-vernalized flowering time were negatively correlated (Figure 4). Furthermore, dormancy

is strongly affected by environmental conditions during seed maturation (Penfield and

Springthorpe 2012; Huang et al. 2015; Burghardt et al. 2016). Thus, interaction between

flowering time and dormancy in the wild could stem from both genetic correlations and

environmentally induced interactions. Despite our expectation for interactions between

germination and flowering traits, models of collection day that included flowering time by

dormancy interactions had a higher AIC and did not explain more deviance than a model with

both traits separately. Including interactions between flowering time and dormancy in our

models did not help to explain collection date in the wild.

We found that predicted wild collection date was positively related to flowering time

genetic values when we extended our analysis to predict wild flowering times from a smoothed

surface of collection dates fit to herbarium records. In our linear models, average flowering time

breeding value was more closely related to this estimated day of collection than was the actual

date of collection of ecotypes in experiments. However, a large portion of phenological variation

within local regions remained unexplained even with phenological genetic values. We attempted

to avoid records that were unusually young or old by only using specimens that had both flowers

and fruits; moreover, both our herbarium and seed collections spanned nearly the entire year

(Julian days 5–350 for herbarium records, 43–346 for seed collections). Still, herbarium records

are known to skew slightly towards earlier in seasons (though removing herbarium specimens

from our analyses did not change results) (Daru et al. 2018), while seed collections must be

collected long enough after the initiation of flowering to allow for the development of some

mature seeds. A single observation during reproduction, common for natural history collection

vouchers, cannot resolve the uncertainty around when plants begin their vegetative growth in the

wild, making it difficult to describe phenology of the full life cycle. With better information and

in climates where growing conditions are clearly defined, it may be possible to estimate

germination dates and full life cycle phenology from herbarium specimens (Bontrager et al.

2025a). Nevertheless, we found the expected strong latitudinal cline in collection dates of

Arabidopsis, going from March to late June across the study region, and we previously found

that collection date accelerates by ~one week for every 1ºC spring warming (DeLeo et al. 2020),

suggesting collection dates do reflect natural phenology.

Spring onset differs among locations, and so a calendar date at a higher latitude or

elevation may be functionally earlier in the season than a lower latitude or elevation. PTUs

represent how much of a growing season has passed by a given date by recording temperature

and daylight above a threshold. Thus, plants collected at lower PTUs may be early flowering (developmentally) despite a later collection date. Higher PTU may indicate plants growing as summer or fall annuals that germinated later in the year. In that case, ecotypes with higher PTUs would likely be fast cycling, earlier flowering plants (Figure 2). While we did find a negative correlation between flowering time genetic values in field and vernalized experiments versus PTU, we see the opposite in non-vernalized experiments. This last observation is somewhat surprising, since we expected plants that flower later in the absence of vernalization to be more likely to grow as winter annuals in the wild and flower early in the spring (Figure 1). Higher PTUs in these presumptive winter annuals may suggest that plants in these locations regulate their life histories to flower later in the year than expected by temperature and daylight alone. There is some evidence that later flowering ecotypes could in fact be more flexible in their flowering time relative to germination than early flowering ecotypes (Miryeganeh et al. 2018), which could hide the signal of early spring flowering we expected in late flowering Arabidopsis.

Within regions, genetic phenology values do not predict phenology in the wild

Because Arabidopsis exhibits substantial local within-population variation in phenology (Jones 1971; Brachi et al. 2013; Alonso-Blanco et al. 2016; DeLeo et al. 2020; Arana and Picó 2025), we asked how genetic variation in phenology was related to phenological variation within local regions. This within-region phenological variation can have fitness consequences. Variation contributes to population persistence in variable environments, as when different germination behavior provides bet hedging in the seedbank (Cohen 1967; Gremer and Venable 2014).

We found that neither dormancy nor flowering time genetic values from experiments were significantly related to collection date when we accounted for geographic variation in local mean collections dates in a GAM. However, flowering time and dormancy did improve GAMs for collection date over a model that included only geographic location and elevation, suggesting that genetic variation in the timing of germination and flowering time does explain a minor fraction of the phenological variation within regions. Non-vernalized flowering time was negatively, non-significantly related to local variation in collection date, perhaps because in many locations, genotypes flowering later in experiments without vernalization are more likely to behave as winter annuals, flowering early in the spring before spring annuals.

In our models, there was very little difference between non-vernalized, vernalized, and

field-measured flowering times in predicting collection dates, despite the expectation that field

experiments recreate conditions similar to nature. However, data from field experimental trials

were available for a smaller set of ecotypes than non-vernalized and vernalized indoor trials, and

we lacked Iberian ecotypes with field experimental data and date of collection, limiting the

usefulness of this measure across the range. Our model provided weak evidence that the genetic

values for flowering time and dormancy explained within-region variation in wild collection day.

However, our set of ecotypes with known collection day and experimental indoor flowering time

numbered 258, potentially still too few to detect strong statistical significance given the noise

arising from plasticity in responses to local environmental gradients.

Conclusion

We demonstrated that experimental measures of genetic variation in the timing of individual

developmental transitions in Arabidopsis are significantly, but modestly, correlated with

phenological variation in wild plants. On one hand, the existence of a relationship suggests there

is some relevance of phenotypes measured in experiments to those exhibited in nature. However,

the weakness of the relationships (especially locally, within regions) suggests the existence of

either a) complex interactions between developmental stages to determine phenology, (b) low

heritability in nature, or (c) extensive rank changing genotype-environment interactions across

microsites. Genetic differences in Arabidopsis phenology across the species range are often cited

as evidence of the adaptive importance of phenology (Stinchcombe et al. 2004; Fournier-Level et

al. 2011; Samis et al. 2012; Brachi et al. 2013; Exposito-Alonso 2020). Incorporating

information on between-population and within-population phenology diversity helps to clarify

how selection may be acting on phenology across the landscape and how influential plasticity is

in the phenology of this annual herb. Yet, genetic flowering times among populations across the

species range are associated with phenological variation in natural history collections, due to

some combination of genotypic and environmental effects (and their possible co-gradient

variation) (Jones et al. 2024). While phenological plasticity is often of large magnitude, it may

not be enough to maintain fitness under environmental change (Zettlemoyer et al. 2024).

Ultimately, controlled experiments on many plants have suggested phenology is under selection

in nature, but understanding more subtle environmental differences and stochasticity may help to

clarify the evolution of phenology and translate genetic values into reliable predictions with and

between populations.

References

Abley, K., P. Formosa-Jordan, H. Tavares, E. Chan, O. Leyser, and J. C. W. Locke. 2020. An ABA-GA
bistable switch can account for natural variation in the variability of Arabidopsis seed germination
time. bioRxiv, doi: https://doi.org/10.1101/2020.06.05.135681.

Ågren, J., C. G. Oakley, S. Lundemo, and D. W. Schemske. 2017. Adaptive divergence in flowering time
among natural populations of Arabidopsis thaliana: Estimates of selection and QTL mapping.
Evolution (N Y) 71:550–564.

Akaike, H. 1974. A New Look at the Statistical Model Identification. IEEE Trans Automat Contr 19:716–
723.

Alonso-Blanco, C., J. Andrade, C. Becker, F. Bemm, J. Bergelson, K. M. M. Borgwardt, J. Cao, E. Chae, T.
M. M. Dezwaan, W. Ding, J. R. R. Ecker, M. Exposito-Alonso, A. Farlow, J. Fitz, X. Gan, D. G. G.
Grimm, A. M. M. Hancock, S. R. R. Henz, S. Holm, M. Horton, M. Jarsulic, R. A. A. Kerstetter, A.
Korte, P. Korte, C. Lanz, C. R. Lee, D. Meng, T. P. P. Michael, R. Mott, N. W. W. Muliyati, T. Nägele,
M. Nagler, V. Nizhynska, M. Nordborg, P. Y. Y. Novikova, F. X. Picó, A. Platzer, F. A. A. Rabanal, A.
Rodriguez, B. A. A. Rowan, P. A. A. Salomé, K. J. J. Schmid, R. J. J. Schmitz, Ü. Seren, F. G. G.
Sperone, M. Sudkamp, H. Svardal, M. M. M. Tanzer, D. Todd, S. L. L. Volchenboum, C. Wang, G.
Wang, X. Wang, W. Weckwerth, D. Weigel, and X. Zhou. 2016. 1,135 genomes reveal the global
pattern of polymorphism in *Arabidopsis thaliana*. Cell 166:481–491.

Alpert, P., and E. L. Simms. 2002. The relative advantages of plasticity and fixity in different
environments: When is it good for a plant to adjust? Evol Ecol 16:285–297.

Amasino, R. 2004. Vernalization, competence, and the epigenetic memory of winter. Plant Cell, doi:
10.1105/tpc.104.161070.

Andrés, F., and G. Coupland. 2012. The genetic basis of flowering responses to seasonal cues.

Auge, G. A., L. K. Blair, A. Karediya, and K. Donohue. 2018. The autonomous flowering-time pathway
pleiotropically regulates seed germination in Arabidopsis thaliana. Ann Bot 121:183–191.

Balasubramanian, S., S. Sureshkumar, J. Lempe, and D. Weigel. 2006. Potent induction of *Arabidopsis*
thaliana flowering by elevated growth temperature. PLoS Genet 2:0980–0989.

Baskin, C. C., and J. M. Baskin. 1988. Germination ecophysiology of herbaceous plant species in a
temperate region. Am J Bot 75:286–305.

Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using lme4. J
Stat Softw 67:1–48.

Baythavong, B. S. 2011. Linking the spatial scale of environmental variation and the evolution of
phenotypic plasticity: Selection favors adaptive plasticity in fine-grained environments. American
Naturalist 178:75–87.

Boyd, E. W., L. A. Dorn, C. Weinig, and J. Schmitt. 2007. Maternal effects and germination timing
mediate the expression of winter and spring annual life histories in *Arabidopsis thaliana*. Int J Plant
Sci 168:205–214.

Brachi, B., N. Faure, M. Horton, E. Flahauw, A. Vazquez, M. Nordborg, J. Bergelson, J. Cuguen, and F.
Roux. 2010. Linkage and association mapping of *Arabidopsis thaliana* flowering time in nature.
PLoS Genet, doi: 10.1371/journal.pgen.1000940.

Brachi, B., R. Villoutreix, N. Faure, N. Hautekèete, Y. Piquot, M. Pauwels, D. Roby, J. Cuguen, J. Bergelson,
and F. Roux. 2013. Investigation of the geographical scale of adaptive phenological variation and its
underlying genetics in *Arabidopsis thaliana*. Mol Ecol 22:4222–4240.

Burghardt, L. T., B. R. Edwards, and K. Donohue. 2016. Multiple paths to similar germination behavior in
Arabidopsis thaliana. New Phytologist 209:1301–1312.

Burghardt, L. T., C. J. E. Metcalf, A. M. Wilczek, J. Schmitt, and K. Donohue. 2015. Modeling the influence
of genetic and environmental variation on the expression of plant life cycles across landscapes. Am
Nat 185:212–227. University of Chicago Press Chicago, IL.

Caicedo, A. L., J. R. Stinchcombe, K. M. Olsen, J. Schmitt, and M. D. Purugganan. 2004. Epistatic
interaction between Arabidopsis FRI and FLC flowering time genes generates a latitudinal cline in a
life history trait. Proc Natl Acad Sci U S A 101:15670–15675.

Chiang, G. C. K., D. Barua, E. Dittmar, E. M. Kramer, R. R. de Casas, and K. Donohue. 2013. Pleiotropy in
the wild: The dormancy gene dog1 exerts cascading control on life cycles. Evolution (NY) 67:883–
893.

Chiang, G. C. K., D. Barua, E. M. Kramera, R. M. Amasino, and K. Donohue. 2009. Major flowering time
gene, FLOWERING LOCUS C, regulates seed germination in Arabidopsis thaliana. Proc Natl Acad Sci
USA 106:11661–11666.

Cohen, D. 1967. Optimizing reproduction in a randomly varying environment when a correlation may
exist between the conditions at the time a choice has to be made and the subsequent outcome. J
Theor Biol 16:1–14.

Daru, B. H., D. S. Park, R. B. Primack, C. G. Willis, D. S. Barrington, T. J. S. Whitfeld, T. G. Seidler, P. W.
Sweeney, D. R. Foster, A. M. Ellison, and C. C. Davis. 2018. Widespread sampling biases in herbaria
revealed from large-scale digitization. New Phytologist 217:939–955.

Davila Olivas, N. H., E. Frago, M. P. M. Thoen, K. J. Kloth, F. F. M. Becker, J. J. A. van Loon, G. Gort, J. J. B.
Keurentjes, J. van Heerwaarden, and M. Dicke. 2017. Natural variation in life history strategy of
Arabidopsis thaliana determines stress responses to drought and insects of different feeding guilds.
Mol Ecol 26:2959–2977.

Davis, C. C., C. G. Willis, B. Connolly, C. Kelly, and A. M. Ellison. 2015. Herbarium records are reliable
sources of phenological change driven by climate and provide novel insights into species'
phenological cueing mechanisms. Am J Bot 102:1599–1609.

Debieu, M., C. Tang, B. Stich, T. Sikosek, S. Effgen, E. Josephs, J. Schmitt, M. Nordborg, M. Koornneef,
and J. de Meaux. 2013. Co-Variation between Seed Dormancy, Growth Rate and Flowering Time
Changes with Latitude in *Arabidopsis thaliana*. PLoS One, doi: 10.1371/journal.pone.0061075.

DeLeo, V. L., D. N. L. Menge, E. M. Hanks, T. E. Juenger, and J. R. Lasky. 2020. Effects of two centuries of global environmental variation on phenology and physiology of *Arabidopsis thaliana*. Glob Chang Biol 26:523–538.

Donohue, K. 2005. Niche construction through phenological plasticity: Life history dynamics and ecological consequences.

Exposito-Alonso, M. 2020. Seasonal timing adaptation across the geographic range of Arabidopsis thaliana. Proceedings of the National Academy of Sciences 117:9665–9667.

Exposito-Alonso, M., F. Vasseur, W. Ding, G. Wang, H. A. Burbano, and D. Weigel. 2018. Genomic basis and evolutionary potential for extreme drought adaptation in *Arabidopsis thaliana*. Nat Ecol Evol, doi: 10.1038/s41559-017-0423-0.

Footitt, S., Z. Huang, H. A. Clay, A. Mead, and W. E. Finch-Savage. 2013. Temperature, light and nitrate sensing coordinate Arabidopsis seed dormancy cycling, resulting in winter and summer annual phenotypes. Plant Journal 74:1003–1015.

Fournier-Level, A., A. Korte, M. D. Cooper, M. Nordborg, J. Schmitt, and A. M. Wilczek. 2011. A map of local adaptation in *Arabidopsis thaliana*. Science (1979), doi: 10.1126/science.1209271.

Fournier-Level, A., A. M. Wilczek, M. D. Cooper, J. L. Roe, J. Anderson, D. Eaton, B. T. Moyers, R. H. Petipas, R. N. Schaeffer, B. Pieper, M. Reymond, M. Koornneef, S. M. Welch, D. L. Remington, and J. Schmitt. 2013. Paths to selection on life history loci in different natural environments across the native range of *Arabidopsis thaliana*. Mol Ecol 22:3552–3566.

Franco, M., and J. Silvertown. 1996. Life history variation in plants: An exploration of the fast-slow continuum hypothesis. Philosophical Transactions of the Royal Society B: Biological Sciences 351:1341–1348.

Gamba, D., C. Lorts, A. Haile, S. Sahay, L. Lopez, T. Xia, E. Kulesza, D. Elango, J. Kerby, M. Yifru, C. E. Bulafu, T. Wondimu, K. Glowacka, and J. R. Lasky. 2023. The genomics and physiology of abiotic stressors associated with global elevation gradients in *Arabidopsis thaliana*. bioRxiv preprint.

Gremer, J. R., and D. L. Venable. 2014. Bet hedging in desert winter annual plants: Optimal germination strategies in a variable environment. Ecol Lett 17:380–387.

Harris, I., P. D. Jones, T. J. Osborn, and D. H. Lister. 2014. Updated high-resolution grids of monthly climatic observations–the CRU TS3. 10 Dataset. International journal of climatology 34:623–642. Wiley Online Library.

Hu, J., L. Lei, and J. De Meaux. 2017. Temporal fitness fluctuations in experimental *Arabidopsis thaliana* populations. PLoS One 12.

Huang, X., J. Schmitt, L. Dorn, C. Griffith, S. Effgen, S. Takao, M. Koornneef, and K. Donohue. 2010. The earliest stages of adaptation in an experimental plant population: Strong selection on QTLS for seed dormancy. Mol Ecol 19:1335–1351.

Huang, Z., S. Footitt, A. Tang, and W. E. Finch-Savage. 2018. Predicted global warming scenarios impact
on the mother plant to alter seed dormancy and germination behaviour in Arabidopsis. Plant Cell
Environ 41:187–197.

Huang, Z., H. Ölçer-Footitt, S. Footitt, and W. E. Finch-Savage. 2015. Seed dormancy is a dynamic state:
Variable responses to pre- and post-shedding environmental signals in seeds of contrasting
Arabidopsis ecotypes. Seed Sci Res 25:159–169.

Huo, H., S. Wei, and K. J. Bradford. 2016. DELAY of GERMINATION1 (DOG1) regulates both Seed
dormancy and flowering time through microRNA pathways. Proc Natl Acad Sci U S A 113:E2199–
E2206.

Jimenez-Gomez, J. M., J. A. Corwin, B. Joseph, J. N. Maloof, and D. J. Kliebenstein. 2011. Genomic
analysis of QTLs and genes altering natural variation in stochastic noise. PLoS Genet 7.

Johnston, I. G., and G. W. Bassel. 2018. Identification of a bet-hedging network motif generating noise in
hormone concentrations and germination propensity in Arabidopsis. J R Soc Interface 15.

Jones, C. V, C. E. Regan, J. A. Firth, E. F. Cole, and B. C. Sheldon. 2024. Environmental similarity between
relatives reduces heritability of reproductive timing in wild great tits. bioRxiv 2024.03.11.584392.

Juenger, T. E., S. Sen, K. A. Stowe, and E. L. Simms. 2005. Epistasis and genotype-environment
interaction for quantitative trait loci affecting flowering time in *Arabidopsis thaliana*. Pp. 87–105 *in*
Genetica.

Jump, A. S., R. Marchant, and J. Peñuelas. 2009. Environmental change and the option value of genetic
diversity.

Karrenberg, S., and A. Widmer. 2008. Ecologically relevant genetic variation from a non-Arabidopsis
perspective.

Kenney, A. M., J. K. McKay, J. H. Richards, and T. E. Juenger. 2014. Direct and indirect selection on
flowering time, water-use efficiency (WUE, $\delta13C$), and WUE plasticity to drought in *Arabidopsis*
thaliana. Ecol Evol 4:4505–4521.

Korves, T. M., K. J. Schmid, A. L. Caicedo, C. Mays, J. R. Stinchcombe, M. D. Purugganan, and J. Schmitt.
2007. Fitness Effects Associated with the Major Flowering Time Gene FRIGIDA in *Arabidopsis*
thaliana in the Field . Am Nat, doi: 10.1086/513111.

Lasky, J. R., B. R. Forester, and M. Reimherr. 2018. Coherent synthesis of genomic associations with
phenotypes and home environments. Molecular Ecology Resources 18:91–106.

Lasky, J. R., M. Takou, D. Gamba, and T. H. Keitt. 2024. Estimating scale-specific and localized spatial
patterns in allele frequency. Genetics 227:3, doi: 10.1093/genetics/iyae082.

Lawrence-Paul, E. H., and J. R. Lasky. 2024. Ontogenetic changes in ecophysiology are an understudied
yet important component of plant adaptation. Am J Bot e16294.

Lempe, J., S. Balasubramanian, S. Sureshkumar, A. Singh, M. Schmid, and D. Weigel. 2005. Diversity of
flowering responses in wild *Arabidopsis thaliana* strains. PLoS Genet, doi:
10.1371/journal.pgen.0010006.

Li, Y., Y. Huang, J. Bergelson, M. Nordborg, and J. O. Borevitz. 2010. Association mapping of local
climate-sensitive quantitative trait loci in *Arabidopsis thaliana*. Proc Natl Acad Sci USA 107:21199–
21204.

Lorts, C. M., and J. R. Lasky. 2020. Competition × drought interactions change phenotypic plasticity and
the direction of selection on Arabidopsis traits. New Phytologist n/a. John Wiley & Sons, Ltd.

Ludlow, M. M. 1989. Strategies of response to water stress. Pp. 269–281 *in* K. H. Kreeb, H. Richter, and
T. M. Hinckley, eds. Structural and functional responses to environmental stresses: water shortage.
SPB Academic Publishers, Berlin.

Lyons, R., A. Rusu, J. Stiller, J. Powell, J. M. Manners, and K. Kazan. 2015. Investigating the association
between flowering time and defense in the *Arabidopsis thaliana-Fusarium oxysporum* interaction.
PLoS One, doi: 10.1371/journal.pone.0127699.

MacGillivray, F., I. L. Hudson, and A. J. Lowe. 2010. Herbarium collections and photographic images:
Alternative data sources for phenological research. Pp. 425–461 *in* Phenological Research:
Methods for Environmental and Climate Change Analysis. Springer Netherlands, Dordrecht.

Marcer, A., D. S. Vidigal, P. M. A. James, M. J. Fortin, B. Méndez-Vigo, H. W. M. Hilhorst, L. Bentsink, C.
Alonso-Blanco, and F. X. Picó. 2018. Temperature fine-tunes Mediterranean *Arabidopsis thaliana*
life-cycle phenology geographically. Plant Biol 20:148–156.

Martínez-Berdeja, A., M. C. Stitzer, M. A. Taylor, M. Okada, E. Ezcurra, D. E. Runcie, and J. Schmitt. 2020.
Functional variants of DOG1 control seed chilling responses and variation in seasonal life-history
strategies in *Arabidopsis thaliana*. Proc Natl Acad Sci USA 117:2526–2534.

Méndez-Vigo, B., A. R. Castilla, R. Gómez, A. Marcer, C. Alonso-Blanco, and F. X. Picó. 2022.
Spatiotemporal dynamics of genetic variation at the quantitative and molecular levels within a
natural *Arabidopsis thaliana* population. Journal of Ecology 110.

Méndez-Vigo, B., N. H. Gomaa, C. Alonso-Blanco, and F. Xavier Picó. 2013. Among- and within-
population variation in flowering time of Iberian *Arabidopsis thaliana* estimated in field and
glasshouse conditions. New Phytologist 197:1332–1343.

Miller-Rushing, A. J., R. B. Primack, D. Primack, and S. Mukunda. 2006. Photographs and herbarium
specimens as tools to document phenological changes in response to global warming. Am J Bot
93:1667–1674.

Miryeganeh, M., M. Yamaguchi, and H. Kudoh. 2018. Synchronisation of Arabidopsis flowering time and
whole-plant senescence in seasonal environments. Sci Rep 8.

Onofri, A., P. Benincasa, M. B. Mesgaran, and C. Ritz. 2018. Hydrothermal-time-to-event models for seed
germination. European Journal of Agronomy 101:129–139.

Penfield, S., and V. Springthorpe. 2012. Understanding chilling responses in Arabidopsis seeds and their
contribution to life history. Philosophical Transactions of the Royal Society B: Biological Sciences
367:291–297.

Poorter, H., F. Fiorani, R. Pieruschka, T. Wojciechowski, W. H. van der Putten, M. Kleyer, U. Schurr, and J.
Postma. 2016. Pampered inside, pestered outside? Differences and similarities between plants
growing in controlled conditions and in the field.

Postma, F. M., and J. Ågren. 2018. Among-year variation in selection during early life stages and the
genetic basis of fitness in *Arabidopsis thaliana*. Mol Ecol 27:2498–2511. Blackwell Publishing Ltd.

Primack, D., C. Imbres, R. B. Primack, A. J. Miller-Rushing, and P. Del Tredici. 2004. Herbarium specimens
demonstrate earlier flowering times in response to warming in Boston. Am J Bot 91:1260–1264.

Ranal, M. A., and D. G. De Santana. 2006. How and why to measure the germination process?

Ratcliffe, D. 1976. Germination characteristics and their inter- and intra-population variability in
Arabidopsis. Arabidopsis Information Service 34–45.

Reich, P. B. 2014. The world-wide "fast-slow" plant economics spectrum: a traits manifesto. Journal of
Ecology 102:275–301.

Ritz, C., F. Baty, and J. C. Streibig. 2015. Dose-Response Analysis Using R. PLoS One 10:e0146021.

Salguero-Gómez, R., O. R. Jones, E. Jongejans, S. P. Blomberg, D. J. Hodgson, C. Mbeau-Ache, P. A.
Zuidema, H. De Kroon, and Y. M. Buckley. 2016. Fast-slow continuum and reproductive strategies
structure plant life-history variation worldwide. Proc Natl Acad Sci U S A 113:230–235.

Samis, K. E., C. J. Murren, O. Bossdorf, K. Donohue, C. B. Fenster, R. L. Malmberg, M. D. Purugganan, and
J. R. Stinchcombe. 2012. Longitudinal trends in climate drive flowering time clines in North
American *Arabidopsis thaliana*. Ecol Evol 2:1162–1180.

Simpson, G. G., and C. Dean. 2002. Arabidopsis, the Rosetta stone of flowering time?

Springthorpe, V., and S. Penfield. 2015. Flowering time and seed dormancy control use external
coincidence to generate life history strategy. Elife 2015.

Stearns, S. C. 1989. Trade-Offs in Life-History Evolution. Funct Ecol 3:259. JSTOR.

Stinchcombe, J. R., C. Weinig, M. Ungerer, K. M. Olsen, C. Mays, S. S. Halldorsdottir, M. D. Purugganan,
and J. Schmitt. 2004. A latitudinal cline in flowering time in *Arabidopsis thaliana* modulated by the
flowering time gene FRIGIDA. Proceedings of the National Academy of Sciences, doi:
10.1073/pnas.0306401101.

Stock, A. J., B. V. McGoey, and J. R. Stinchcombe. 2015. Water availability as an agent of selection in
introduced populations of *Arabidopsis thaliana*: Impacts on flowering time evolution. PeerJ
2015:1–16.

Thomas, B., and D. Vince-Prue. 1997. Photoperiodism in plants. 2nd ed. Academic Press, San Diego, CA,
USA.

Vidigal, D. S., A. C. S. S. Marques, L. A. J. Willems, G. Buijs, B. Méndez-Vigo, H. W. M. Hilhorst, L.
Bentsink, F. X. Picó, and C. Alonso-Blanco. 2016. Altitudinal and climatic associations of seed
dormancy and flowering traits evidence adaptation of annual life cycle timing in *Arabidopsis*
thaliana. Plant Cell Environ 39:1737–1748.

Walker, M. D., R. C. Ingersoll, and P. J. Webber. 1995. Effects of interannual climate variation on
phenology and growth of two alpine forbs. Ecology 76:1067–1083.

Wilczek, A. M., L. T. Burghardt, A. R. Cobb, M. D. Cooper, S. M. Welch, and J. Schmitt. 2010. Genetic and
physiological bases for phenological responses to current and predicted climates. Philosophical
Transactions of the Royal Society B: Biological Sciences 365:3129–3147.

Wilczek, A. M., J. L. Roe, M. C. Knapp, M. D. Cooper, C. Lopez-Gallego, L. J. Martin, C. D. Muir, S. Sim, A.
Walker, J. Anderson, J. F. Egan, B. T. Moyers, R. Petipas, A. Giakountis, E. Charbit, G. Coupland, S.
M. Welch, and J. Schmitt. 2009. Effects of genetic perturbation on seasonal life history plasticity.
Science (1979) 323:930–934.

Wood, S. N. 2006. Generalized additive models: an introduction with R. 1st ed. Chapman and Hall/CRC.

Yee, T. W., and M. Mackenzie. 2002. Vector generalized additive models in plant ecology. Ecol Modell,
doi: 10.1016/S0304-3800(02)00192-8.

Yee, T. W., and N. D. Mitchell. 2006. Generalized additive models in plant ecology. Journal of Vegetation
Science, doi: 10.2307/3236170.

Zettlemoyer, M., R. Conner, M. Seaver, E. Waddle, and M. DeMarche. 2024. A Long-Lived Alpine
Perennial Advances Flowering under Warmer Conditions but Not Enough to Maintain Reproductive
Success. Am Nat 203.

Zhou, D., T. Wang, and I. Valentine. 2005. Phenotypic plasticity of life-history characters in response to
different germination timing in two annual weeds. Canadian Journal of Botany 83:28–36.

Supplemental Material

Plasticity due to maternal effects

Maternal conditions may be an important source of phenological plasticity; thus we used our dormancy experiment to examine how variation in flowering time among maternal plants influences germination. Variation in phenology may not be adaptive in regions where there is harsh seasonality or environmental conditions are less variable year to year (Cohen 1967). Similarly, observations of phenology in the wild could be temporally variable because of year-to-year environmental differences (Walker et al. 1995; Hu et al. 2017; Postma and Ågren 2018) or genetically diverse individuals germinating from the seedbank (Ratcliffe 1976). Among replicate maternal individuals in the same controlled experiment, variation in phenology is likely due to responses to very subtle environmental differences and some developmental stochasticity.

For each ecotype and for each maternal replicate, we used R packages 'drc' (Ritz et al. 2015) and 'drcSeedGerm' (Onofri et al. 2018) to fit a logistic function to germination counts to estimate three parameters: maximum germination proportion, time to 50% germination, and slope of the germination curve. A Generalized Linear Mixed Model was fit using the lmer package in R (Bates et al. 2015) to estimate the influence of relative flowering time of maternal replicates of each ecotype on germination traits. Because plasticity or responsiveness to environmental cues may confer greater fitness in some environments and not others (Alpert and Simms 2002; Baythavong 2011), germination traits and variation were regressed against maternal flowering time and location of origin. We performed hierarchical clustering of a distance matrix of phenotypic variation among ecotypes to group ecotypes that reacted similarly to different stratification treatments.

We found that some ecotypes had more consistent phenology across maternal lines. For example, DAM1 maternal replicates clustered together in germination across cold treatments, whereas Yeg-7 replicates did not (Figure S8A). Cold stratification altered expression of variance among maternal sources. In Yeg-7, for example, maternal lines P vs A had similar total germination following 3 days or 2 weeks of stratification but different germination for 0 days stratification. Thus, stratification (or lack of stratification) could contribute to diverse phenology for similar genetic backgrounds under natural conditions. The first and second principal components in an analysis of flowering and germination traits explained 24.55% and 17.85% of

the variation respectively, but again maternal replicates of ecotypes did not always cluster in PC

space (Figure S8B). The variance in phenology for an ecotype was not related to collection date

or geographic origin (latitude and longitude, Figure S8C), suggesting that clinal variation in

maternal plasticity effects do not confound our earlier large-scale analyses comparing trait

genetic values with phenology in nature. Nevertheless, these effects may further obscure genetic

contributions to phenology in nature.

We expected to see more variation among ecotypes from regions with favorable growing

conditions year-round. However, the lack of a geographic association between variance in

germination traits and flowering time among replicate individuals of an ecotype may indicate

geographically consistent selection on the canalization of phenology.

Figure S1. 42 ecotypes have measures of vernalized, non-vernalized, and field flowering times as
well as primary and secondary dormancy traits (standardized trait units are shown). The latest
flowering accessions tend to be late flowering across all treatments, but primary and secondary
dormancy vary independently from flowering time.

Figure S2. Standardized phenology traits combined across multiple published controlled
experiments combined with our new experiments, using an alternate standardization (compare
with Figure 3). Here, each experiment's values were standardized to mean zero, unit standard
deviation.

Figure S3. Flowering times and germination rates within experiments do not predict wild

phenology better than average values across multiple experiments. Left, non-vernalized

flowering times at 16°C (using the same criteria to remove collection day and geographical

outliers as in the full set of accessions) Alonso-Blanco et al. (2016) did not significantly predict

collection day (p > 0.2, n = 173). Right, germination rates after 4 days of stratification at 4°C

Martínez-Berdeja et al. (2020) did not significantly predict collection day (p > 0.1, n = 84).

996 Figure S4. Date of collection (left panels) and PTUs (photothermal units) at collection (right

997 panels) of wild plants compared to standardized genetic values for phenological traits from

998 experiments, here using an alternate standardization (compare to Figure 4). Here, each

999 experiment's values were standardized to mean zero unit standard deviation. The values shown

1000 for field flowering time experiments were calculated from fall-sown experiments only.

1001

Figure S5. Date of collection (left panels) and PTUs (photothermal units) at collection (right panels) of wild plants compared to standardized genetic values for phenological traits from experiments, here using an alternate standardization and excluding herbarium specimens (compare to Figure 4). Here, each experiment's values were standardized to mean zero unit standard deviation. The values shown for field flowering time experiments were calculated from fall-sown experiments only.

1009 Figure S6. Date of collection (left panels) and PTUs (photothermal units) at collection (right

1010 panels) of wild plants compared to standardized genetic values for phenological traits from

1011 experiments, excluding herbarium specimens (compare to Figure 4). The values shown for field

1012 flowering time experiments were calculated from fall-sown experiments only.

1013

Figure S8: Variation in experimental phenology among maternal lines. A) Early flowering lines
tended to have lower germination under 0 days stratification relative to 3 days or 2 weeks
stratification, and later-flowering lines tended to have longer times to 50% germination.
However, there was variation across traits. Replicate maternal lines of the same ecotype did not
always cluster. B) Maternal lines of the same ecotype did not always cluster together in PCA of
phenology traits. Lines belonging to the same ecotype are represented with the same color,
although colors are not unique across ecotypes. PC1 was positively associated with time to 50%
germination at 0 days and 3 days stratification and flowering time and negatively associated with
time to 50% germination at 2 weeks stratification and max germination under all treatments. C.
Flowering time, collection day, or geographic origin did not explain variance among maternal
lines.

1025

1026

1027

1028

1029

1030   Table S1 [see attached supplemental Excel file]. Experiments included in our study and

1031   phenotypes represented.

1032

Table S2: Model statistics for other variables in models of within-region phenological variation. P-values for the estimates are given in parentheses. Phenology genetic values did not significantly predict within-region variation in phenology.

Model	Non-Vernalized Flowering Time	Primary Dormancy	FT*Dormancy	Elevation	Adj R^2
Collection Date $\sim \mu + FT +$ Elevation	-3.25 (0.35)	NA	NA	**0.0148 (0.006)**	0.883
$PTU^{1/2} \sim \mu + FT +$ Elevation	-1.21 (0.75)	NA	NA	**-0.0147 (0.006)**	0.876
Collection Date $\sim \mu +$ Dormancy $+$ Elevation	NA	1.98 (0.56)	NA	**0.0126 (0.04)**	0.876
$PTU^{1/2} \sim \mu +$ Dormancy $+$ Elevation	NA	2.56 (0.48)	NA	**-0.0181 (0.009)**	0.858
Collection Date $\sim \mu + FT +$ Dormancy $+$ Elevation	-8.19 (0.10)	2.19 (0.51)	NA	**0.0120 (0.05)**	0.881
$PTU^{1/2} \sim \mu +$ FT*Dormancy $+$ Elevation	-5.41 (0.56)	5.39 (0.43)	-6.86 (0.66)	**-0.0183 (0.009)**	0.864