Structure-guided secretome analysis of gall-forming microbes offers insights into effector diversity and evolution

Soham Mukhopadhyay[1-3]*, Muhammad Asim Javed[1-3], Jiaxu Wu[1-3] and Edel Pérez-López[1-3]*

[1]Départment de Phytologie, Faculté des sciences de l'agriculture et de l'alimentation, Université Laval, Quebec City, Quebec, Canada.

[2]Centre de recherche et d'innovation sur les végétaux (CRIV), Université Laval, Quebec City, Quebec, Canada.

[3]L'Institute EDS, Université Laval, Quebec City, Québec City, Québec, Canada

*Corresponding: somuk3@ulaval.ca, edel.perez-lopez@fsaa.ulaval.ca

ABSTRACT

Phytopathogens secrete effector molecules to manipulate host immunity and metabolism. Recent advances in structural genomics have identified fungal effector families whose members adopt similar folds despite sequence divergence, highlighting their importance in virulence and immune evasion. To extend the scope of comparative structure-guided analysis to more evolutionarily distant phytopathogens with similar lifestyles, we used AlphaFold2 to predict the 3D structures of the secretome from selected plasmodiophorid, oomycete, and fungal gall-forming pathogens. Clustering protein folds based on structural homology revealed species-specific expansions and a low abundance of known orphan effector families. We identified novel sequence-unrelated but structurally similar (SUSS) effector clusters, rich in conserved motifs such as 'CCG' and 'RAYH'. We demonstrate that these motifs likely play a central role in maintaining the overall fold. We also identified a SUSS cluster adopting a nucleoside hydrolase-like fold conserved among various gall-forming microbes. Notably, ankyrin proteins were significantly expanded in gall-forming plasmodiophorids, with most being highly expressed during clubroot disease, suggesting a role in pathogenicity. Altogether, this study advances our understanding of secretome landscapes in gall-forming microbes and provides a valuable resource for broadening structural phylogenomic studies across diverse phytopathogens.

KEYWORDS: Effector biology, AlphaFold, Plasmodiophorid, Gall-forming pathogens, SUSS

INTRODUCTION

The evolutionary arms race between host plants and their microbial pathogens is a fascinating example of adaptation and counteradaptation. Central to this battlefield are the effectors — secreted proteins from pathogens that manipulate host cellular processes to facilitate infection and colonization [1]. Effectors can also be recognized by plant receptor and trigger immunity and defense responses in the host [2]. Due to their central role in pathogen-host interactions, effectors must continually evolve to evade detection by the host's immune system [3]. This arms race drives rapid changes in effector sequences, often resulting in high mutation rates and diversification [4]. Additionally, effectors should maintain structural integrity for functionality while altering surface residues to avoid immune recognition [5]. Recent studies have defined these groups of effectors as Sequence-Unrelated Structurally Similar (SUSS), which despite lacking sequence similarity, share significant structural homology [6,7]. For example, the MAX effector family, which includes proteins from various fungal pathogens like *Magnaporthe oryzae* and *Pyrenophora tritici-repentis*, exhibits a conserved β-sandwich fold[8]. Other examples include the RXLR-WY effector families in oomycetes [9,10], the LARS effectors in *Cladosporium fulvum* and *Leptosphaeria maculans* [11], the RALPH effectors in *Blumeria graminis* [12], and the FOLD effectors in *Fusarium oxysporum f. sp. lycopersici* (D. S. Yu et al., 2024) demonstrating structural conservation. FOLD effectors have also recently been found in the secretome of unrelated pathogenic and symbiotic fungi, pointing to the fold's relevance in plant colonization and expansion in different evolutive groups [14]. A recent study that classified orphan effector candidates (OECs) from Ascomycota into 62 main structural groups, proposes that such structural conservation can be explained by changes in thermodynamic frustration at surface residues, which increase the robustness of the protein structure while altering potential interaction sites [5].

Some of the discussed studies have been fueled by the emergence of machine-learning tools like AlphaFold [15], which has revolutionized the field of protein modeling and enabled the computational prediction of pathogen effector structures. The utility of AlphaFold in plant-microbe interactions has been further demonstrated by its ability to predict the structure of the highly conserved AvrE-family of effector proteins [16]. These proteins are crucial in the pathogenesis of various phytopathogenic bacteria but are challenging to study due to their large size, toxicity to plant and animal cells, and low sequence similarity to known proteins [17]. The structure prediction revealed β-barrel structures similar to bacterial porins, allowing to modulate host cell functions by facilitating the movement of small molecules across plasma membrane. AlphaFold Multimer [18], an

extension of AlphaFold designed to predict in-silico protein-protein interactions, has been recently

used to identify 15 effector candidates capable of targeting the active sites of chitinases and

proteases [19], expanding the applicability of these tools to advance the tailored functional

characterization of effector proteins.

Recent advances in structure-guided secretome analysis have mostly focused on fungal

pathogens due to their significant economic impact and the availability of robust genomic and

transcriptomic resources. However, in recent years, there is a growing interest in protist pathogens,

which are quickly becoming a threat to agriculture and environment [20]. For example, the gall-

forming *Plasmodiophora brassicae*, a protist belonging to the class Plasmodiophorid, can cause

significant yield loss in canola fields [21,22]. Moreover, these obligate biotrophic protists are

impossible to culture in axenic media and therefore difficult to transform [23]. Although the effector

repertory for some of these protists has been predicted, the majority of their secretome remains

uncharacterized due to the absence of known protein domains [20]. To gain more insights into the

secretome composition and effector biology of these understudied pathogens, we conducted

structural similarity-based clustering of the predicted effectors of selected plasmodiophorid,

oomycete, and fungal gall-farming pathogens. Here we examined (*i*) if the primary secretome

families present in each pathogen share common folds with known fungal effector families; (*ii*) if

they share common folds that could be associated to their biotrophic lifestyle and gall- forming

pathogenicity strategy; and (*iii*) if some of the known effectors from these pathogens are part of

SUSS effector families. By comparing the secretome of gall-forming pathogens from distant

lineages, we provide a comprehensive overview of the uniqueness and commonality of the

secretome landscape and offers more insights into the protist effector families by bringing them

into the structural genomics era.

RESULTS

Secretome prediction and structural modeling of gall forming pathogens

Based on genome availability, phylogenetic distance, and economic significance, six gall-

forming pathogens were selected for secretome analysis- two plasmodiophorids, *Plasmodiophora*

brassicae and *Spongospora subterranea*, oomycete *Albugo candida*, and three fungi from different

lineages, *Taphrina deformans, Ustilago maydis*, and *Synchytrium endobioticum* (**Fig. 1a**). To gain

better understanding of plasmodiophorids, for which structural data is scarce, we also included

Polymyxa betae, a non-gall-forming plasmodiophorid vector of the beet necrotic yellow vein virus,

causing Rhizomania disease [24] (**Fig. 1a**). To identify the putative secretome of these plant

pathogens, we first employed SignalP [25] to predict sequences with N-terminal signal peptide.

Sequences carrying a signal peptide were subjected to DeepTMHMM [26] search to remove

sequences with transmembrane domains. Mature protein sequences greater than 1000 amino acid

length were also removed (**Fig. 1b**). The *Ustilago maydis* secretome and corresponding structures

were obtained from a recent study [7] using similar filtering steps. This resulted in a total of 4197

proteins from seven gall-forming or related plant pathogens (**Table S1**). Next, the structures of

3575 proteins, excluding the *U. maydis* secretome which was already available, were modeled

using AlphaFold2. InterproScan [27] was performed on all 4197 proteins against pfam, Gene3D and

SUPERFMAILY databases, identifying 41.59% of the proteins analyzed carrying a known protein

domain (**Table S1**). Next, 2349 structures with pLDDT > 65 were selected for further analysis

(**Table S1**). Although pLDDT > 70 is recommended as the threshold for structures with reliable

confidence by AlphaFold developers, we selected a score of 65 or higher to include *AvrSen1*

(pLDDT 67), the only known *S. endobioticum* avirulent gene [28] (**Fig. 1c**). This resulted in the

inclusion of 298 candidate effectors which would have otherwise been excluded (**Table S1**).

IUPred3 [29] was also used to score the proteins as disordered or not based on whether 50% of the

sequence position was predicted to be disordered (Table **S1**). Most of the proteins (97%) with

pLDDT > 70 were not disordered, while 26% of the secretome with pLDDT < 70 was disordered,

thus showing the limitation of Alpahfold2 in modelling such effectors (**Table S1**).

Structure-based clustering reveals species-specific folds and low homology with known

fungal effector families

Structural similarity among the predicted structures were assessed using TMAlign [30] by scoring

all structures against each other. To study the similarity with known effector families, we also

included 19 crystal structures from the DELD, FOLD, LARS, MAX, KP6, RALPH, NTF2-like,

ToxA, C2-like, and Zn-binding effector families, as previously utilized in a recent study [14] (**Table**

S2). Additionally, we also included 62 structural families of orphan candidate effectors (OCE)

recently identified in the Ascomycota lineage [5] (**Table S2**). Comparisons with TMScore greater

than 0.5 were considered positive for structural similarity. Markov clustering [31,32] with an inflation

value of two was applied to cluster the secretome based on structural similarity score. This resulted

in 254 structural clusters with at least two members (**Fig. 2**, **Table S3**).

Ankyrin repeat containing proteins were the largest cluster for plasmodiophorid *P. brassicae*

(n=42) and *S. subterranea* (n=39) (**Fig. 2**). The largest cluster detected on *A. candida* secretome is

formed by 'CCG' [33] class of effectors (n=48), while *U. maydis* largest cluster was composed of the

Tin2-like proteins [34] (n=31). For *S. synchytrium*, the largest cluster (n=64) contains AvrSen1

virulence factor (**Fig. 2a**). *P. betae* and *T. deformans* do not carry large (n > 30) effector clusters

which could be due to *P. betae's* vector-like nature and *T. deformans*'s reduced genome size

compared to other fungi [35]. *T. deformans*'s largest cluster is primarily composed of various

Glycoside hydrolases while *P. betae*'s largest cluster consists of orphan helical proteins (**Table**

S3). A 20-member Kinase family (cluster 13) was noted in *P. brassicae* which was absent in *A.*

candida and 3 other fungi. Chitin deacetylases, which have been reported to convert chitin to less

immunogenic chitosan [36] was expanded (n=11) in *P. brassicae* (**Table S3**). Chitin deacetylases

were found in 5 out of 7 pathogens tested, except in oomycete *A. candida* and ascomycete *T.*

deformans, which despite being a fungus contains very little chitin in the cell wall [37] (**Table S3**).

In *S. subterranea* and *P. brassicae*, a unique TauD/TfdA protein cluster was also identified,

typically found in bacteria for taurine utilization as a sulfur source [38]. *P. betae* carries a 6 member

G-domain protein cluster, with similarity to it's host *Beta vulgari*'s GTPases (**Fig. S1**, **Table S3**).

Among the well characterized fungal effector families, only three KP6 fold [39] and one RALPH like

fold were found in *U. maydis* (**Fig. S1**, **Table S3**). *T. deformans* carry 2 ToxA homologs (**Fig. S1**,

Table S3). .

Effector folds conserved across kingdoms

Six protein folds—Hydrolases (clusters 2, 8, 9), Carboxypeptidases (cluster 12), Aspartyl

proteases (cluster 3), Lectins (cluster 38), SCP domain (cluster 24), and an orphan group (cluster

5)—contained proteins from all the pathogens investigated in this study, indicating deep

evolutionary conservation of these folds across kingdoms (**Table S3**). In fact, 20 out of 62

Ascomycete orphan effector groups had at least one structural homolog in plasmodiophorids and

oomycetes tested in this study, although a complete evolutionary connection would require the

comparison of a much larger number of pathogens (**Table S3**). We didn't find any specific fold (n

> 1 per species) conserved only among the gall-forming pathogens studied, indicating that this

virulence strategy can be achieved by different mechanisms without necessarily converging onto

common effector folds.

Structural search identifies a nucleoside hydrolase-like fold conserved in some gall-forming pathogens

Effectors are notorious for carrying unknown domains, making it difficult to predict the putative function of promising candidates [41]. We searched for uncharacterized *P. brassicae* candidate effectors within the same cluster that were also overexpressed during infection. Two candidate effectors, PBTT_09143 and PBTT_07479, which are the first and fifth most expressed proteins at 16 days post inoculation (dpi) during clubroot disease, belong to the cluster 21 grouping with PBTT_0412, none of which carries a known domain (**Fig. 3a-c**). Interestingly, the cluster also includes ten members from *A. candida* secretome, with eight carrying a predicted nucleoside hydrolase domain (**Table S4**). We subjected the three *P. brassicae* candidates to a FoldSeek-mediated [42] structural search against the PDB100 and AFD-proteome databases. Nucleoside hydrolases always emerged as the top hit (E-value < 10^-5, Prob ~1, TMscore >=0.5) (**Table S4**). Next, searching AFDB cluster webtool [43], which allows for the identification of structural homologs across the known protein space, AFDB clusters A0A0G4IP88 and A0A024FV66 emerged as the top hits. The members of these clusters are often gall-forming, carry predicted signal peptide and belong to various biotrophs *like Melanopsichium pennsylvanicum, Ustilago maydis, Albugo candida, Sporisorium scitamineum, Testicularia cyperi, Colletotrichum orbiculare,* among others (**Fig 3b, d**). Some of the members also do not carry identifiable domains (**Table S4**) and show limited sequence similarity among themselves (**Fig 3b**, **Table S4**). Interestingly, the cluster also includes bacterial Type III effector HopQ1 from *Pseudomonas syringae* (**Table S4**). Unlike *P. brassicae* effectors, HopQ1 is predicted to carry a nucleoside hydrolase domain, although the domain has been reported to be unable to bind standard nucleosides [44]. HopQ1 has been reported to be associated with 14-3-3 plant proteins to promote virulence [44]. Thus, it's possible that the nucleoside hydrolase-like fold present in various gall-forming fungal, protist, and oomycete pathogens might have also neo-functionalized and be involved in new molecular strategies during the infection.

A fungal effector family shows structural homology despite extreme sequence diverge

The Mig1 protein in *Ustilago maydis* is a maize-induced effector that plays a crucial role in the biotrophic interaction between the fungus and its host [45]. It is specifically induced during the biotrophic phase, contributing to the fungus's pathogenicity. Cluster 30, specifically found in *Ustilago maydis*, contains this effector (**Table S5**). Upon examining the sequence and structural

similarities between the members of the cluster, we discovered instances of structural homology

(TMScore > 0.5) despite pairwise sequence identity being as low as 0.6% and no higher than 30%

(**Fig. 4a-d**, **Table S5**). When aligning the protein sequences of 13 members, four cysteine residues

were found to be strongly conserved (**Fig. 4e**). These four residues form two disulfide bridges

(**Fig. 4f**), likely playing a crucial role in maintaining the overall fold despite significant sequence

dissimilarity. All 13 members of the cluster were expressed upon infection (**Table S6**). A variable

expression was observed for the *Mig1*-like genes located in the same genomic region, including

members with completely opposite patterns of induction (**Table S6**), suggesting a possible

regulatory role or the acquisition of new functions.

Identification of new SUSS effector families enriched in known motifs

The fungi *S. endobioticum* and *A. candida* encode large effector clusters which included

previously identified avirulent factors AvrSen1 and CCG28/31/33/70, respectively (**Table S7**) [46]. It

was previously shown that the N-terminal region of the CCG28 ,33 and 70 share structural

homology [46]. To examine if these clusters represent SUSS families, we carried out the sequence-

based clustering of the 4197 proteins from the seven pathogens investigated here. We performed

BlastP search in all-vs-all mode and kept only those results with E-value lower than 10^-04 and

bidirectional coverage of 50%. Markov clustering revealed 642 sequence-based clusters with at

least two members (**Table S7**). We searched for sequence clusters having members from the same

structural clusters. This revealed the presence of 12 sequence-related clusters associated with

AvrSen1 structural cluster and 11 sequence-related clusters associated with CCG structural cluster

(**Fig. 5a**, **Table S7**). It was previously reported that CCG containing effectors share limited

similarity around the CCG motif and can be grouped in several clades based on sequence

similarity [33]. Thus, AvrSen1 and CCG represent novel SUSS effector families whose similarities

can't be delineated by sequence search alone. Integration of sequence and structure data increased

the member count of AvrSen1 and CCG clusters to 124 and 50 from previous 64 and 48,

respectively (**Table S8**).

It has been reported that *S. endobioticum* secretome is enriched in RAYH [47].While it is

understood that the CCG class of effectors derived its name due to the presence of the CCG motif,

it was not immediately clear if Avrsen1 cluster was also the source of the conserved RAYH motif.

To verify that, we subjected the mature protein sequences of the two pathogen secretome to motif

search using MEME [48]. Here we identified a 16 amino acid long RAYH motif present in 118 *S.*

endobioticum proteins and 15 amino acid long 'CCG' motif present in 74 *A. candida* proteins (E <

0.1, combined match p < 0.001) (**Fig. 5b**, **Fig. S2, S3**, **Table S8**). Of the 118 sequences proteins

carrying the RAYH motif, 79 were members of the AvrSen1 cluster in *S. endobioticum* (**Table S8**).

Thus, the expansion of SUSS effectors in *A. candida* and *S. synchytrium* has resulted in the

enrichment of common motifs, something that has recently been observed for the Y/F/WxC motif

in *Blumeria graminis* RNA-like effector cluster [7].

Selection pressure analysis on *A. candida* CCG members shows that both cysteine residues in

the 'CCG' motifs and two additional cysteine residues within 50 amino acids of the motif are often

under purifying selection. (**Fig. S4, S5**). Visualizing these four cysteines on the predicted structure

shows that they form disulfide bridges and probably play a crucial role in overall maintenance of

the fold (**Fig. 5c**). The CCG motif seems to be a crucial part of a module consisting of two parallel

alpha-helices joined to a beta-sheet and CCG effectors are often composed of several of these

modules (**Fig. 5c**). The RAYH motif was also found to be part of the core structure of most

AvrSen1-like effectors, forming a long alpha-helix (**Fig 5d**). Apart from the RAYH motif region,

several surrounding hydrophobic residues were also strongly conserved, likely playing a role in

maintaining the structure (**Fig. S6**). Examining the sequence-related subclusters of the AvrSen1-

like family we found that the effectors are evolving by keeping the core alpha-helix bundle fixed

while diversifying the peripheral stretches (**Fig 5d**).

Ankyrin repeat containing proteins are a common feature of gall forming plasmodiophorids

The largest structural cluster in *P. brassicae* and *S. subterranea* consists of ankyrin repeat-

containing proteins (hereafter ANK proteins) (**Fig. 6a**). The presence of only five members in the

non-gall-forming plasmodiophorid *P. betae*, in contrast to over 40 members in the gall-forming *P.*

brassicae and *S. subterranea*, highlights the importance of this domain in their pathogenicity

strategies (**Table S9**). Interestingly, InterProScan identified 32 additional *P. brassicae* proteins

with ANK domains, while only two *S. subterranea* ANK proteins could be identified outside of the

cluster (**Table S9**). We found that *P. brassicae* secretome is richer in repeat proteins compared to *S.*

subterranean ANK proteins repertory (**Fig 6a**), with 19 additional leucine-rich repeats (LRRs), ten

tetratricopeptide repeats, and three MORN repeats (**Table S9**). Although 40 out of 74 ankyrins are

over expressed (TPM >10) at 16- and 26 dpi in *Arabidopsis thaliana*, only five LRRs are induced

(**Table S9**). Notably, 17 ankyrin-repeat proteins also carry the SKP1/BTB/POZ domain, which is

often involved in ubiquitination [49] (**Fig. 6b**, **Table S9**).

A MEME motif scan with the members of the ankyrin cluster identified a 33 amino acid long

motif in *P. brassicae* and a 32 amino acid length motif in *S. spongospora* (**Fig 6c**, **Table S9**).

Aligning the MEME profile of the two identified ankyrin motifs shows a strong conservation of

two Leucine and one Alanine residues (**Fig. 6d**) and upon visualization, those residues form the

hydrophobic pocket between the two alpha helices of the ankyrin repeat, stabilizing the structure

(**Fig 6d**). The rest of the non-conserved residues were found to be highly polymorphic (**Fig. 6d**).

Finally, to have an idea of the possible role of ANK proteins in plasmodiophorid virulence, we

selected 70 ankyrin domain-containing proteins from *P. brassicae* and screened them against 20

key immune-related genes in *Arabidopsis* using AlphaFold-Multimer (**Table S10**). Protein-protein

interactions were considered significant if the inter-chain Predicted Aligned Error (PAE) value was

below 10, and the iPTM+pTM score was 70 or higher. Among the identified interactions, MPK3,

MAPK4, MAPK6 SnRK1, NPR1, XCP1, CNGC4, and BAK1 were targeted by a total of ten

ankyrin domain proteins (**Fig. 6e**, **Table S11**). This dataset should serve as a valuable starting

point for further understanding the role of ANKs in the virulence mechanisms of

plasmodiophorids.

DISCUSSION

This study identified the primary protein folds in gall-forming pathogens secretome supporting

the idea that pathogen secretome are often dominated by expansion of specific folds which have

been adopted and diversified over the course of evolution. Characterizing these primary effector

folds in understudied plasmodiophorids like *P. brassicae* and *S. subterranea* would offer valuable

insights into their virulence strategies, which remain largely enigmatic [20,50]. Here we found that the

ankyrin proteins, which are significantly expanded in gall-forming plasmodiophorids but less so in

the related species *P. betae*, may be key to their ability to manipulate host immune responses and

promote gall formation (**Fig. 2**, **Fig. 6**). This finding aligns with previous research indicating the

importance of repeat-containing proteins in the virulence of plant pathogens [51]. Example of that

are *Phytophthora* spp. effectors containing tandem repeats of the "(L)WY" motif, whose

modularity and elaborate mimicry of a host phosphatase helps to promote infection [52]. ANK motifs

are well-known for mediating protein-protein interactions [53], and have been identified as type IV

effectors in the intracellular human pathogens *Legionella pneumophila* and *Coxiella burnetii* [54].

Thus, we can hypothesize that the highly polymorphic surface residues of plasmodiophorid ANK motifs, coupled with the variable frequency of occurrence in member proteins, results in different binding surfaces for the host target proteins.

This study also identified conserved protein folds across multiple kingdoms, particularly the hydrolase, carboxypeptidase, and aspartyl protease folds (**Table S3**). These folds appear to play a fundamental role in the virulence strategies of these pathogens, likely due to their ability to perform essential biochemical functions that facilitate infection [55–57]. The conservation of these folds across diverse species highlights their evolutionary significance and suggests that they may represent mechanisms of host manipulation that have been retained through speciation. We also identified a nucleoside hydrolase like fold in evolutionary distant gall-forming pathogens which has homology to bacterial effector HopQ1 (**Fig. 3**). Nucleoside hydrolases are involved in the purine salvage pathway in various pathogens [58], but similar to HopQ1's mode of action, these gall-forming biotrophs might also be targeting 14-3-3 proteins, which are implicated in hormonal signaling [59]. Although HopQ1 is widely conserved within the *Pseudomonas* species complex, our FoldSeek-mediated search identified hits in *Pseudomonas savastanoi* isolates, some of which are known to form galls on woody plants [60,61]. It remains to be seen if some of the HopQ1 homologs have been specifically adapted in these bacteria to support a particular lifestyle.

Here we also provided further evidence for the divergent evolution model of effector evolution, which describes how members of the same effector family can exhibit extreme sequence dissimilarity over a long period while retaining the core fold intact [7]. We show that this evolution mechanism occurs in both fungi and oomycetes and often involves the conservation of cysteine or hydrophobic residues to maintain the original fold (**Fig. 4**, **Fig. 5**). Given that sequence dissimilarity between homologs can be extreme, as exemplified by the Mig1 family in *U. maydis*, it would become common practice among researchers to incorporate structural knowledge into sequence searches to accurately gauge the diversity of the effector families (**Fig. 4**).

Overall, our study underscores the power of structural genomics and machine learning tools like AlphaFold2 in uncovering the complexities of pathogen effector repertoires. The findings presented here open new avenues for research into the evolution of virulence strategies in phytopathogens and highlight the potential for these insights to inform the development of novel approaches to plant disease management. As we continue to expand our understanding of effector biology, particularly in under-studied pathogens, it will be crucial to integrate these structural

insights with functional studies to fully elucidate the roles of these proteins in host-pathogen
interactions.

MATERIALS AND METHODS

Secretome prediction

The proteome of *P. brassicae* was derived from our recent study generating the first complete
genome of the clubroot pathogen [62]. We are thankful to Prof. Anne Legrève for providing the
updated annotation of the *P. betae* genome and proteome previously published by them [63,64]. The
rest of the proteome for *S. subterranea* [65], *A. candida* [66], *S. endobioticum* [47], *U. maydis* [67] and *T.*
deformans [35] were downloaded from Uniprot database. SignalP 6 was utilized to identify
sequences with predicted signal peptides, which were subsequently removed. DeepTMHMM was
run through pybiolib package. InterProScan 5.61-93.0 was used to confirm the presence of known
domains using the Pfam, Gene3D, and SUPERFAMILY databases.

Structure prediction

A total of 3615 mature protein sequences were selected to be modeled using AlphaFold 2, but 40
predictions repeatedly failed at the MSA construction step, resulting in 3575 structures. To
expedite the process, ParaFold 2.0 [68] was used, which employs AlphaFold 2.3.1 internally but
distributes the CPU and GPU tasks to facilitate parallelization. 'Valeria' compute cluster
(https://valeria.science/accueil) of Université Laval was used for the structure prediction. The full
database was used to construct the MSA (multiple sequence alignment), and models were
predicted in 'monomer' mode, resulting in five PDB structures sorted by pLDDT scores. The
Rank_0 PDB was used for subsequent studies. Finally, 2000 models with pLDDT scores over 65
were selected for downstream analysis.

Similarity search, clustering, and network plots

TM-Align was used to perform an all-versus-all structural comparison of 2000 models, and
those comparisons with a normalized TM-score above 0.5 were considered significant. All-versus-
all sequence comparison was performed using BlastP [69] with an E-value < 10^-4 and bidirectional
coverage of at least 50%. Structure and sequence similarity data, represented by three columns
with the first two as target and source IDs, and the third one being TM-score/E-value, were
clustered using the Markov clustering with an inflation value of two. For sequence clustering, E-

values were loaded following the recommendation of the MCL workflow [mcxload -abc seq.abc --
stream-mirror --stream-neg-log10 -stream-tf 'ceil(200)']. Custom Python scripts were written to
find the sequence-related subclusters belonging to the same structural cluster and to count the
occurrence of cluster members (https://github.com/Edelab). Plots were generated using Chiplot
(https://www.chiplot.online/) and ggplot2 [70].
354

Sequence Alignment

356 Pairwise alignment of protein sequences was done by EMBOSS Needle [71]. Clustal Omega was
357 used to generate multiple protein sequence alignment [72]. Kalign 3.4.0 [73] was used to generate
358 alignment of extremely divergent Mig1 cluster.
359

Selection pressure analysis and structure visualization

Coding sequences (CDS) were obtained from the Ensembl Fungi/ENA database. Multiple
nucleotide sequences were aligned using the Kc-Align codon-aware aligner [74]. Positions with more
than 50% gaps were removed using the Clipkit [75] online tool in 'gappy' mode. The trimmed
alignments were manually analyzed using Geneious (http://www.geneious.com/) for correct codon
alignment. The resulting alignment was uploaded to the Datamonkey server
(http://www.datamonkey.org/), which hosts the HyPhy package [76]. All the branches were used as
input for FEL [77] to identify sites under purifying selection (p < 0.01). The ESPript 3.0 web server
was used to generate multiple sequence alignments [78]. Multiple structures were aligned using
mTm-Align [79]. PyMOL 3.0.4 [80] was used to visualize the PDB files and color conserved sites or
disulfide bridges.
371

Expression analysis

Datasets for the RNA-Seq reads obtained at 16- and 26 dpi during P. brassicae infection were
downloaded from the EBI server (accession number PRJEB12261). The reads from the infected
samples were mapped to the *A. thaliana* genome TAIR10 (Genbank accession number
GCF_000001735.4) using HiSAT2 [81] to remove host contaminant sequences. To use Salmon [82], an
index file was created by concatenating the *P. brassicae* genome and CDS sequences. The
remaining RNA-Seq reads were quasi-mapped to the index using Salmon 1.10.0 to generate
normalized transcripts per million (TPM) counts for all 10,521 genes. TPM values from three

replicates were averaged. Pre-processed gene expression data for *U. maydis* was publicly available

[83].

Motif scanning

384   MEME 5.5.5 was used to scan the list of amino acid sequences in '-anr' mode (E < 0.1) to

385   discover motifs of any length and frequency. MAST 5.5.5 was used to protein sequences with

386   MEME motifs. Sequence profiles of *P. brassicae* and *S. subterranea* ANK motifs were aligned

387   using Tomtom 5.5.5 [84].

Structural homology search

390   FoldSeek (https://search.foldseek.com/search) was used to search for structural homology

391   against Uniprot50, Swiss-Prot, the AlphaFold proteome, and PDB. The AFDB cluster database

392   (https://cluster.foldseek.com/) was searched to find cluster members in the AlphaFold database.

In-Silico protein-protein interaction prediction

ANK proteins were screened for interaction against a list of *A. thaliana* immune genes using

AlphaPulldown v1.0.4 [85]. It utilizes AlphaFold Multimer but separates the CPU and GPU jobs and

reuses the MSA to reduce compute time. The full AlphaFold 2.3.0 database was used for MSA

creation. The resulting models from the AlphaPulldown run were parsed with the supplied

singularity image alpha-analysis_jax_0.4.sif to produce the final iPTM+pTM score table.

ChimeraX 1.8 [86] was used to visualize the predicted aligned error for residue pairs under 4

Ångströms at the interface between the two chains produced by AlphaFold-Multimer.

ACKNOWLEDGMENTS

We are grateful to the bioinformatics support personnel and infrastructure at IBIS, Université

Laval, for their constant assistance throughout this project, and to Prof. Sylvain Raffaele for his

help in selecting the representative accessions of 62 orphan effector family clusters in

Ascomycota. This work was funded by the Canola Agronomic Research Program, Grant ID

2021.4, Western Grain Research Foundation, Canola Council of Canada, Alberta Canola and

Manitoba Canola Growers Association; and Discovery Program, Grant ID RGPIN-2021-02518,

Natural Sciences and Engineering Research Council of Canada.

COMPETING INTERESTS

None declared.

AUTHOR CONTRIBUTIONS

S.M., M.A.J. and E.P.L. conceived the research; S.M. conducted the research; S.M., and E.P.L. wrote the manuscript and prepared figures; S.M., M.A.J., J.W and E.P.L. edited the manuscript; E.P.L. supervised the research.

ORCID

Soham Mukhopadhyay: 0000-0002-5279-0396

Muhammad Asim Javed: 0000-0003-1658-3565

Jiaxu Wu : 0000-0002-9165-2076

Edel Pérez-López: 0000-0003-2841-9391

DATA AVAILABILITY

The datasets used in this study can be downloaded from Zenodo: https://doi.org/10.5281/zenodo.11152389 , and the scripts from GitHub: https://github.com/Edelab

REFERENCES

1. Toruño, T. Y., Stergiopoulos, I. & Coaker, G. Plant-Pathogen Effectors: Cellular Probes Interfering with Plant Defenses in Spatial and Temporal Manners. *Annu Rev Phytopathol* **54**, 419–441 (2016).

2. Chen, J., Zhang, X., Rathjen, J. P. & Dodds, P. N. Direct recognition of pathogen effectors by plant NLR immune receptors and downstream signalling. *Essays Biochem* **66**, 471–483 (2022).

3. Martel, A. *et al.* The ETS-ETI cycle: evolutionary processes and metapopulation dynamics driving the diversification of pathogen effectors and host immune factors. *Curr Opin Plant Biol* **62**, 102011 (2021).

4. Liu, L. *et al.* Arms race: diverse effector proteins with conserved motifs. *Plant Signal Behav* (2019).

5.   Derbyshire, M. C. & Raffaele, S. Surface frustration re-patterning underlies the structural
landscape and evolvability of fungal orphan candidate effectors. *Nat Commun* **14**, 5244
(2023).

6.   Seong, K. & Krasileva, K. V. Computational Structural Genomics Unravels Common Folds
and Novel Families in the Secretome of Fungal Phytopathogen Magnaporthe oryzae.
Molecular Plant-Microbe Interactions® **34**, 1267–1280 (2021).

7.   Seong, K. & Krasileva, K. V. Prediction of effector protein structures from fungal
phytopathogens enables evolutionary analyses. *Nat Microbiol* **8**, 174–187 (2023).

8.   Guillen, K. de *et al.* Structure Analysis Uncovers a Highly Diverse but Structurally
Conserved Effector Family in Phytopathogenic Fungi. *PLoS Pathog* **11**, e1005228 (2015).

9.   Combier, M., Evangelisti, E., Piron, M.-C., Schornack, S. & Mestre, P. Candidate effector
proteins from the oomycetes Plasmopara viticola and Phytophthora parasitica share similar
predicted structures and induce cell death in Nicotiana species. *PLoS One* **17**, e0278778
(2022).

10.   Win, J. *et al.* Sequence Divergent RXLR Effectors Share a Structural Fold Conserved across
Plant Pathogenic Oomycete Species. *PLoS Pathog* **8**, e1002400 (2012).

11.   Lazar, N. *et al.* A new family of structurally conserved fungal effectors displays epistatic
interactions with plant resistance proteins. *PLoS Pathog* **18**, e1010664 (2022).

12.   Cao, Y. *et al.* Structural polymorphisms within a common powdery mildew effector scaffold
as a driver of coevolution with cereal immune receptors. *Proceedings of the National
Academy of Sciences* **120**, e2307604120 (2023).

13.   Yu, D. S. *et al.* The structural repertoire of Fusarium oxysporum f. sp. lycopersici effectors
revealed by experimental and computational studies. *Elife* **12**, (2024).

14.   Teulet, A. *et al.* A pathogen effector FOLD diversified in symbiotic fungi. *New Phytologist*
239, 1127–1139 (2023).

15.   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583–589 (2021).

16.   Nomura, K. *et al.* Bacterial pathogens deliver water- and solute-permeable channels to plant
cells. *Nature* **621**, 586–591 (2023).

17.   Xin, X.-F. *et al.* Pseudomonas syringae Effector Avirulence Protein E Localizes to the Host
Plasma Membrane and Down-Regulates the Expression of the NONRACE-SPECIFIC
DISEASE RESISTANCE1/HARPIN-INDUCED1-LIKE13 Gene Required for Antibacterial
Immunity in Arabidopsis. *Plant Physiol* **169**, 793–802 (2015).

18.   Evans, R. *et al.* Protein complex prediction with AlphaFold-Multimer. Preprint at
477   https://doi.org/10.1101/2021.10.04.463034 (2022).

478   19.   Homma, F., Huang, J. & van der Hoorn, R. A. L. AlphaFold-Multimer predicts cross-
479   kingdom interactions at the plant-pathogen interface. *Nat Commun* **14**, 6040 (2023).

480   20.   Mukhopadhyay, S., Garvetto, A., Neuhauser, S. & Pérez-López, E. Decoding the Arsenal:
481   Protist Effectors and Their Impact on Photosynthetic Hosts. *Molecular Plant-Microbe*
482   *Interactions®* **37**, 498–506 (2024).

483   21.   Javed, M. A. *et al.* The clubroot pathogen Plasmodiophora brassicae: A profile update. *Mol*
484   *Plant Pathol* **24**, 89–106 (2023).

485   22.   Ochoa, J. C. *et al.* Natural variation in Arabidopsis responses to Plasmodiophora brassicae
486   reveals an essential role for Resistance to Plasmodiophora brasssicae 1 (RPB1). *The Plant*
487   *Journal* **116**, 1421–1440 (2023).

488   23.   González-García, M. & Pérez-López, E. Looking for a Cultured Surrogate for Effectome
489   Studies of the Clubroot Pathogen. *Front Microbiol* **12**, (2021).

490   24.   Decroës, A. *et al.* Rhizomania: Hide and Seek of Polymyxa betae and the Beet Necrotic
491   Yellow Vein Virus with Beta vulgaris. *Molecular Plant-Microbe Interactions®* **35**, 989–
492   1005 (2022).

493   25.   Teufel, F. *et al.* SignalP 6.0 predicts all five types of signal peptides using protein language
494   models. *Nat Biotechnol* **40**, 1023–1025 (2022).

495   26.   Hallgren, J. *et al.* DeepTMHMM predicts alpha and beta transmembrane proteins using
496   deep neural networks. Preprint at https://doi.org/10.1101/2022.04.08.487609 (2022).

497   27.   Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics*
498   **30**, 1236–1240 (2014).

499   28.   van de Vossenberg, B. T. L. H. *et al.* The Synchytrium endobioticum AvrSen1 Triggers a
500   Hypersensitive Response in Sen1 Potatoes While Natural Variants Evade Detection.
501   *Molecular Plant-Microbe Interactions®* **32**, 1536–1546 (2019).

502   29.   Erdős, G., Pajkos, M. & Dosztányi, Z. IUPred3: prediction of protein disorder enhanced
503   with unambiguous experimental annotation and visualization of evolutionary conservation.
504   *Nucleic Acids Res* **49**, W297–W303 (2021).

505   30.   Zhang, Y. & Skolnick, J. TM-align: a protein structure alignment algorithm based on the
506   TM-score. *Nucleic Acids Res* **33**, 2302–2309 (2005).

507   31.   Enright, A. J., Van Dongen, S. & Ouzounis, C. A. An efficient algorithm for large-scale
508   detection of protein families. *Nucleic Acids Res* **30**, 1575–1584 (2002).

32.   Van Dongen, S. Graph Clustering Via a Discrete Uncoupling Process. *SIAM Journal on*
Matrix Analysis and Applications **30**, 121–141 (2008).

33.   Furzer, O. J. *et al.* An Improved Assembly of the Albugo candida Ac2V Genome Reveals
the Expansion of the "CCG" Class of Effectors. *Molecular Plant-Microbe Interactions* **35**,
39–48 (2022).

34.   Tanaka, S. *et al.* Neofunctionalization of the secreted Tin2 effector in the fungal pathogen
Ustilago maydis. *Nat Microbiol* **4**, 251–257 (2019).

35.   Cissé, O. H. *et al.* Genome Sequencing of the Plant Pathogen Taphrina deformans, the
Causal Agent of Peach Leaf Curl. *mBio* **4**, 10.1128/mbio.00055–13 (2013).

36.   Gao, F. *et al.* Deacetylation of chitin oligomers increases virulence in soil-borne fungal
pathogens. *Nature Plants 2019 5:11* **5**, 1167–1176 (2019).

37.   Petit, M. & Schneider, A. Chemical analysis of the wall of the yeast form of Taphrina
deformans. *Arch Microbiol* **135**, 141–146 (1983).

38.   Eichhorn, E., Ploeg, J. R. van der, Kertesz, M. A. & Leisinger, T. Characterization of α-
Ketoglutarate-dependent Taurine Dioxygenase from Escherichia coli *. *Journal of*
Biological Chemistry **272**, 23031–23036 (1997).

39.   Thynne, E. *et al.* An array of Zymoseptoria tritici effectors suppress plant immune
responses. Preprint at https://doi.org/10.1101/2024.03.12.584321 (2024).

40.   Akum, F. N., Steinbrenner, J., Biedenkopf, D., Imani, J. & Kogel, K.-H. The Piriformospora
indica effector PIIN_08944 promotes the mutualistic Sebacinalean symbiosis. *Front Plant*
Sci **6**, (2015).

41.   Lovelace, A. H. *et al.* Effector Identification in Plant Pathogens. *Phytopathology* **113**, 637–
650 (2023).

42.   van Kempen, M. *et al.* Fast and accurate protein structure search with Foldseek. *Nat*
Biotechnol **42**, 243–246 (2024).

43.   Barrio-Hernandez, I. *et al.* Clustering predicted structures at the scale of the known protein
universe. *Nature* **622**, 637–645 (2023).

44.   Li, W., Yadeta, K. A., Elmore, J. M. & Coaker, G. The Pseudomonas syringae Effector
HopQ1 Promotes Bacterial Virulence and Interacts with Tomato 14-3-3 Proteins in a
Phosphorylation-Dependent Manner1[C][W][OA]. *Plant Physiol* **161**, 2062–2074 (2013).

45.   Basse, C. W., Stumpferl, S. & Kahmann, R. Characterization of a Ustilago maydisGene
Specifically Induced during the Biotrophic Phase: Evidence for Negative as Well as Positive
Regulation. *Mol Cell Biol* **20**, 329–339 (2000).

46.   Redkar, A. *et al.* The Arabidopsis WRR4A and WRR4B paralogous NLR proteins both
543   confer recognition of multiple Albugo candida effectors. *New Phytol* **237**, 532–547 (2023).

544   47.   van de Vossenberg, B. T. L. H. *et al.* Comparative genomics of chytrid fungi reveal insights
545   into the obligate biotrophic and pathogenic lifestyle of Synchytrium endobioticum. *Sci Rep*
546   **9**, 8672 (2019).

547   48.   Bailey, T. L. & Gribskov, M. Combining evidence using p-values: application to sequence
548   homology searches. *Bioinformatics* **14**, 48–54 (1998).

549   49.   Geyer, R., Wee, S., Anderson, S., Yates, J. & Wolf, D. A. BTB/POZ Domain Proteins Are
550   Putative Substrate Adaptors for Cullin 3 Ubiquitin Ligases. *Mol Cell* **12**, 783–790 (2003).

551   50.   Pérez-López, E. *et al.* Identification of Plasmodiophora brassicae effectors — A challenging
552   goal. *Virulence* **9**, 1344–1353 (2018).

553   51.   Mesarich, C. H., Bowen, J. K., Hamiaux, C. & Templeton, M. D. Repeat-containing protein
554   effectors of plant-associated organisms. *Front Plant Sci* **6**, 160681 (2015).

555   52.   Li, H. *et al.* Pathogen protein modularity enables elaborate mimicry of a host phosphatase.
556   *Cell* **186**, 3196-3207.e17 (2023).

557   53.   Li, J., Mahajan, A. & Tsai, M. D. Ankyrin repeat: A unique motif mediating protein-protein
558   interactions. *Biochemistry* **45**, 15168–15178 (2006).

559   54.   Pan, X., Lührmann, A., Satoh, A., Laskowski-Arce, M. A. & Roy, C. R. Ankyrin repeat
560   proteins comprise a diverse family of bacterial type IV effectors. *Science (1979)* **320**, 1651–
561   1654 (2008).

562   55.   Bradley, E. L. *et al.* Secreted Glycoside Hydrolase Proteins as Effectors and Invasion
563   Patterns of Plant-Associated Fungi and Oomycetes. *Front Plant Sci* **13**, 853106 (2022).

564   56.   Liu, K. *et al.* Effector Protein Serine Carboxypeptidase FgSCP Is Essential for Full
565   Virulence in Fusarium graminearum and Is Involved in Modulating Plant Immune
566   Responses. *https://doi.org/10.1094/PHYTO-02-24-0068-R* (2024) doi:10.1094/PHYTO-02-
567   24-0068-R.

568   57.   Wang, C. *et al.* The secreted FolAsp aspartic protease facilitates the virulence of Fusarium
569   oxysporum f. sp. lycopersici. *Front Microbiol* **14**, 1103418 (2023).

570   58.   Hofer, A. Targeting the nucleotide metabolism of Trypanosoma brucei and other
571   trypanosomatids. *FEMS Microbiol Rev* **47**, 1–20 (2023).

572   59.   Camoni, L., Visconti, S., Aducci, P. & Marra, M. 14-3-3 Proteins in Plant Hormone
573   Signaling: Doing Several Things at Once. *Front Plant Sci* **9**, (2018).

574   60.   Harmon, C. L. *et al.* Bacterial gall of Loropetalum chinense caused by Pseudomonas
575   amygdali pv. loropetali pv. nov. *Plant Dis* **102**, 799–806 (2018).

61. Ramos, C., Matas, I. M., Bardaji, L., Aragón, I. M. & Murillo, J. Pseudomonas savastanoi
pv. savastanoi: some like it knot. *Mol Plant Pathol* **13**, 998–1009 (2012).

62. Javed, M. A., Mukhopadhyay, S., Normandeau, E., Brochu, A.-S. & Pérez-López, E.
Telomere-to-telomere Genome Assembly of the Clubroot Pathogen Plasmodiophora
Brassicae. *Genome Biol Evol* **16**, evae122 (2024).

63. Decroës, A., Calusinska, M., Delfosse, P., Bragard, C. & Legrève, A. First Draft Genome
Sequence of a Polymyxa Genus Member, Polymyxa betae, the Protist Vector of
Rhizomania. *Microbiol Resour Announc* **8**, 10.1128/mra.01509–18 (2019).

64. Decroës, A. *et al.* Metagenomics approach for Polymyxa betae genome assembly enables
comparative analysis towards deciphering the intracellular parasitic lifestyle of the
plasmodiophorids. *Genomics* **114**, 9–22 (2022).

65. Ciaghi, S., Neuhauser, S. & Schwelm, A. Draft Genome Resource for the Potato Powdery
Scab Pathogen Spongospora subterranea. *Molecular Plant-Microbe Interactions®* **31**,
1227–1229 (2018).

66. McMullan, M. *et al.* Evidence for suppression of immunity as a driver for genomic
introgressions and host range expansion in races of Albugo candida, a generalist parasite.
Elife **4**, e04550 (2015).

67. Kämper, J. *et al.* Insights from the genome of the biotrophic fungal plant pathogen Ustilago
maydis. *Nature* **444**, 97–101 (2006).

68. Zhong, B. *et al.* ParaFold: Paralleling AlphaFold for Large-Scale Predictions. Preprint at
https://doi.org/10.48550/arXiv.2111.06340 (2021).

69. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421
(2009).

70. Wickham, H. ggplot2. (2016) doi:10.1007/978-3-319-24277-4.

71. Rice, P., Longden, I. & Bleasby, A. EMBOSS: The European Molecular Biology Open
Software Suite. *Trends in Genetics* **16**, 276–277 (2000).

72. Sievers, F. & Higgins, D. G. Clustal Omega for making accurate alignments of many
protein sequences. *Protein Science* **27**, 135–145 (2018).

73. Lassmann, T. Kalign 3: multiple sequence alignment of large datasets. *Bioinformatics* **36**,
1928–1929 (2020).

74. Nicholas Keener. davebx/kc-align: Codon-aware aligner. https://github.com/davebx/kc-align
(2020).

75. Steenwyk, J. L., Buida, T. J., Li, Y., Shen, X. X. & Rokas, A. ClipKIT: A multiple sequence
alignment trimming software for accurate phylogenomic inference. *PLoS Biol* **18**, e3001007
(2020).

76. Kosakovsky Pond, S. L., Frost, S. D. W. & Muse, S. V. HyPhy: hypothesis testing using
phylogenies. *Bioinformatics* **21**, 676–679 (2005).

77. Kosakovsky Pond, S. L. & Frost, S. D. W. Not So Different After All: A Comparison of
Methods for Detecting Amino Acid Sites Under Selection. *Mol Biol Evol* **22**, 1208–1222
(2005).

78. Robert, X. & Gouet, P. Deciphering key features in protein structures with the new
ENDscript server. *Nucleic Acids Res* **42**, W320–W324 (2014).

79. Dong, R., Peng, Z., Zhang, Y. & Yang, J. mTM-align: an algorithm for fast and accurate
multiple protein structure alignment. *Bioinformatics* **34**, 1719–1725 (2018).

80. Schrödinger LLC. *The PyMOL Molecular Graphics System, Version~1.8.* (2015).

81. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–915
(2019).

82. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and
bias-aware quantification of transcript expression. *Nat Methods* **14**, 417–419 (2017).

83. Lanver, D. *et al.* The Biotrophic Development of Ustilago maydis Studied by RNA-Seq
Analysis. *Plant Cell* **30**, 300–323 (2018).

84. Gupta, S., Stamatoyannopoulos, J. A., Bailey, T. L. & Noble, W. S. Quantifying similarity
between motifs. *Genome Biol* **8**, R24 (2007).

85. Yu, D., Chojnowski, G., Rosenthal, M. & Kosinski, J. AlphaPulldown—a python package
for protein–protein interaction screens using AlphaFold-Multimer. *Bioinformatics* **39**,
(2023).

86. Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis. *Protein
Science* **32**, e4792 (2023).

FIGURE LEGENDS

Fig. 1. Description of the pathogens included in the study, workflow overview and statistics

of structure prediction. A, Cladogram of the pathogens used in the study. The schematics

represent the disease symptoms on their respective hosts, with the white areas representing galls.

The secretome count indicates the number of proteins per species predicted to be secreted, and the

functional annotation shows the percentage of the secretome predicted to contain a known protein
domain in the Pfam, SUPERFAMILY, or Gene3D databases. **B**, Flowchart of the workflow used to
predict the secretome and the corresponding 3D structures. **C**, Raincloud plot showing the median
and density distribution of pLDDT scores of the predicted structures in each pathogen.

Fig. 2. Visualization of dominant protein folds present in each pathogen. A, Network plot of
structurally similar secretome clusters with at least 15 members. Not all 255 clusters are shown to
reduce complexity. Each node represents a single protein, and an edge between two nodes
represents structural similarity (TMScore > 0.5). **B**, Representative structure of the dominant fold
in each pathogen. Since Ankyrin repeats are common in both *P. brassicae* and *S. subterranea*, they
are represented only once.

Fig. 3. Sequence and structural similarity among HopQ1 homologs in *U. maydis*, *A. candida*
and *P. brassicae*. A, Network plot showing the structural similarity between the members of
cluster 21. Edges denote structural similarity (TMScore > 0.5). **B**, Pairwise sequence identity
between selected HopQ1 structural homologs from plasmodiophorids, oomycetes, and fungi,
illustrating sequence dissimilarity between some proteins despite structural homology. **C**, Gene
expression values (log2 TPM) of two highly induced *P. brassicae* genes at 16 and 26 dpi. **D**, 3D
structure of the mature protein sequences, assuming a HopQ1-like fold.

Fig. 4. Sequence and structural similarity among Mig1 homologs in *Ustilago maydis*. A,
Similarity matrix showing the pairwise sequence identity (%) between Mig1 cluster members. **B**,
Similarity matrix showing the pairwise structural homology scores (TMScore) between Mig1
cluster members. **C**, Superimposition of two Mig1 homologues, illustrating structural similarity
despite extreme sequence divergence. **D**, Differential gene expression patterns of two Mig1
tandem duplicates. **E**, Multiple alignment of protein sequences, highlighting the conservation of
cysteine residues (marked in yellow). **F**, Visualization of the conserved cysteine residues forming
disulfide bridges.

Fig. 5. SUSS effector families are enriched in common motifs. A, Network plots demonstrating
that the two primary effector families in *A. candida* and *S. endobioticum* can only be grouped
together when structural data is incorporated into the sequence-based clustering. The plots also

indicate which sequence-based clusters contain the known effectors from these groups. **B**, 'RAYH'

and 'CCG' motif patterns identified by MEME scan. **C**, Disulfide bridges in the 'CCG' motif,

likely playing a pivotal role in structural maintenance, are highlighted in the virulence factor

CCG30. A zoomed-in view of the 'CCG module' shows the four conserved cysteine residues

forming disulfide bridge. **D**, The 'RAYH' motif, occupying the central position in the core alpha-

helix bundle, is highlighted in six sequence-based subclusters within the AvrSen1-like cluster in *S.*

endobioticum.

Fig. 6. Diversity, structural features and host immune targets of ankyrin repeats in

Plasmodiophorids. A, Frequency of repeat-containing proteins in *P. brassicae* and *S. subterranea*.

B, Network plot showing structural homology within *P. brassicae* Ankyrin repeats, also

highlighting the Ankyrin repeats with SKP1/BTB/POZ superfamily domains. **C**, Alignment of

Ankyrin motifs from *P. brassicae* and *S. subterranea*. **D**, Visualization of conserved hydrophobic

residues in a single Ankyrin repeat module. **E**, Number of Ankyrin repeat proteins predicted to

target *Arabidopsis* immune proteins. **F**, AlphaFold Multimer predicted complex of MPK3 and

PBTT_00818, highlighting the predicted aligned errors of surface contacts under 4 Ångströms.

SUPPLEMENTARY INFORMATION

Table S1. Detailed information about the pathogens investigated in this study and their secretome

composition with pLDDT scores, IUPred3 results, and InterproScan identity.

Table S2. Known/orphan effector families searched in gall-forming secretome investigated in this

study.

Table S3. Structural clusters generated in this study; ID of the top 26 clusters detected; and

frequency of clusters, orphan effectors and known fungal effectors per species studied here.

Table S4. Detailed information of cluster 2, FoldSeek search results, AFDB cluster webtool, and

sequence identity matrix with HopQ1.

Table S5. Sequence identity matrix of *Ustilago maydis* Mig1 protein with other members of

cluster 30.

Table S6. Expression of *Ustilago maydis mig1* and members of the cluster 30 in axenic, 1, 2, 4, 6, 8, and 12 dpi.

Table S7. Detailed information of the new SUSS effector families identified in this study. Structure based and Sequence based clustering of CCG effectors. Structure based and Sequence based clustering of AvrSen1-like effectors. MEME output of CCG and RAYH motif scan. Selection of one representative member (highest pLDDT) from each sequence-based cluster for selection pressure analysis in CCGs. Selection of one representative member (highest pLDDT) from each sequence-based cluster to examine RAYH motif conservation in AvrSen1-like cluster.

Table S8. Sequence clusters generated in this study.

Table S9. Detailed information of proteins containing repeats. Ankyrin repeat containing proteins, leucine-rich repeats, and Ankyrin repeat containing proteins with SPK1 domains in *P. brassicae* and *S. subterranean*. Expression of *P. brassicae* ankyrin repeat containing proteins and leucine-rich repeats at 16- and 26 dpi in *A. thaliana*. *P. brassicae* and *S. subterranean* ankyrin repeat containing proteins MEME scan.

Table S10. List of *A. thaliana* and *P. brassicae* proteins used to study modeled interactions with AlphaFold-Multimer.

Table S11. Output of the AlphaFold-Multimer predicting interaction between *A. thaliana* immunity hubs and *P. brassicae* Ankyrin repeat containing proteins.

Fig. S1. The structures of *Beta vulgari*'s GTPases, KP6 fold, RALPH-like fold and ToxA fold share structural similarity with candidate effectors found in P. *betae*, *U. maydis* and *T. deformans*,, respectively. The number of clusters where proteins share structural similarities with these folds is provided.

Fig. S2. Location of CCG motifs identified by MEME scan.

Fig. S3. Location of RAYH motifs identified by MEME scan.

Fig. S4. Analysis evidencing purifying selection in the CCG effector family. Plots have been

downloaded from the Datamonkey server, which hosts the FEL package. The indicated codons are

numbered according to the codon-aligned nucleotide alignment file, where positions with 50%

gaps have been trimmed.

Fig. S5. Multiple alignment of representative members of CCG sequence-based clusters.

Conserved residues (>70% similar by biochemical properties in ESPript3.0) are highlighted.

Fig. S6. Multiple alignment of representative members of AvrSen1-like sequence-based clusters.

Conserved residues (>70% similar by biochemical properties in ESPript3.0) are highlighted.

(a)

Plasmodiophorids | Oomycota | Ascomycota | Basidiomycota | Chytridiomycota

Pathogen	*Plasmodiophora brassicae*	*Polymyxa betae*	*Spongospora subterranea*	*Albugo candida*	*Taphrina deformans*	*Ustilago maydis*	*Synchytrium endobioticum*
Primary host	Brassicaceae	Sugar beet	Potato	Brassicaceae	Peach	Maize	Potato
Secretome count	806	714	576	594	199	582	726
Functional annotation	63 / 37	35 / 65	31 / 69	36 / 64	64 / 36	51 / 49	21 / 79

% annotated
% not annotated

(b)

Entire proteome of the pathogen of interest

↓

SignalP — — → Prediction of sequences with signal peptide

↓

DeepTMHMM — — → Removal of sequences with transmembrane domains

↓

Removal of sequences longer than 1000 AA

↓

Secretome List → InterProScan against the Pfam, GENE3D and SUPERFAMILY database

↓

AlphaFold2

↓

Selection of structures with pLDDT > 65

(c)

P. brassicae n = 800
S. subterranea n = 574
P. betae n = 687
A. candida n = 594
T. deformans n = 195
U. maydis n = 582
S. endobioticum n = 724

pLDDT score

40   60   80   100

- *P. betae*
- *P. brassicae*
- *S. subterranea*
- *U. maydis*
- *A. candida*
- *T. deformans*
- *S. endobioticum*

1 PBTT_04511 Ankyrin repeats

4 A0A507CY85 AvrSen1-like

2 R4XM90 CAZYmes

11 UMAG_05302 Tin-2-like

6 A0A024FTM3 CCG family

5 JG262 Helical

(a)

Cluster 21
- *U. maydis*
- *A. candida*
- *P. brassicae*

(b)

P. syringae HopQ1
P. brassicae PBTT_07479
A. candida A0A024FXU1
C. orbiculare N4USF2
P. brassicae PBTT_09143
S. scitamineum A0A0F7S5Z6
U. maydis A0A0D1E2J4
M. pennsylvanicum A0A077RCL5
T. cyperi A0A317XP18

P. syringae HopQ1
P. brassicae PBTT_07479
A. candida A0A024FXU1
C. orbiculare N4USF2
P. brassicae PBTT_09143
S. scitamineum A0A0F7S5Z6
U. maydis A0A0D1E2J4
M. pennsylvanicum A0A077RCL5
T. cyperi A0A317XP18

Sequence Identity (%)

(c)

PBTT_09143
PBTT_07479

TPM (log2)

Days post inoculation
16   26

(d)

P. brassicae PBTT_09143

P. syringae HopQ1

A. candida A0A024FXU1

U. maydis A0A0D1E2J4

C. orbiculare N4USF2

M. pennsylvanicum A0A077RCL5

S. scitamineum A0A0F7S5Z6

T. cyperi A0A317XP18

(a)

(b)

(c)
UMAG_12257
UMAG_12215
Seq Identity 0.6%
TM Score 0.5

(d)
UMAG_12257
UMAG_12258

(e)

	50	60	70
Mig1	....LPRYDLYHE..HC	EADAG.PSGSDPKWP	C	FTMW
UMAG_01300	...PSPRFSWNA..LC	GSHAH.SS...IKHI	C	FDSL
UMAG_00558	.VSPGNSYDE..YC	KGSGT.KF..RSSRL	C	VNVK
UMAG_01298	...ISHDVAYHL..LC	DQGTSLPLSPSVPRA	C	FETR
UMAG_01301	....EQPP.NWTVGYY	C	SDLSS.SS.....RV	C	FTTD
UMAG_10418	...........GWLV..YC	GGQAG.DIPPGYIYA	C	FESK
UMAG_01299	FDKSAYGQAEWSR..YC	SSVS.KSADLLAHAC	FTIH
UMAG_01297	....ERNSHMYHS..LC	HPHSGLAQNDHRSKA	C	FDTR
UMAG_12258	....QQN..EYDA..YC	SAHAS.PA...VGAI	C	FDTS
UMAG_01302	.VSPGNSWAD..YC	QPGGS.KF..DTHHL	C	VSSS
UMAG_12257	......TWSL..YC	NG.GG.TVPRDTSYT	C	FESD
UMAG_12215	...IEITPLYVDTII	Y	GVRTS.RSQPEIHFE	L	VGHG
UMAG_12216	.GPNSDSKW....Y	C	FAHNG.ND..RLETV	S	TKSG

	140	150	160
Mig1	DTG	C	YKITLK.HKEGFHRR....IWVSDEAGKD	R	DID
UMAG_01300	DQ.	C	IAYDVQ.DVSGN.D........LKNGLV	C	PGQ
UMAG_00558	DSI	C	ASVQIW.TNSAVST.......VIGDQQV	C	YSD
UMAG_01298	DQS	C	VDVVVY.LLPERES.......TVRESW	C	PGQ
UMAG_01301	SQR	C	ADGSVA.LSSGR.L........IDTTTF	C	NGT
UMAG_10418	G..	C	VNLIFF.SSKGPEH.......ISNLIK..T	C	PVA
UMAG_01299	NV.	S	VRAPIR.VSRGRTKEGQEPALAPFMEGTY	C	AGS
UMAG_01297	KLG	C	ARVKIY.HLPTWTL.......EVDQVW	C	AGH
UMAG_12258	YP.	C	VQYTIS.STKGRKD.......SRVGHF	C	PGD
UMAG_01302	DPM	C	GWVVIQ.KRGGDGS.......NLFQQTY	C	KGD
UMAG_12257	G..	C	LS...Y.SINKRND.......FSDLISFLH	C	PVN
UMAG_12215	STK	C	FDIMLLRDVSKKYR......MTIGKTVDG	P	DFD
UMAG_12216	DTG	C	YKIHLSGREQGRIH.......VTTTDENY	R	DID

(f)
UMAG_01300
Disulfide bridge
CYS-15
CYS-27
CYS-88
CYS-107

(a)

Seq cluster 43 with
CCG 28/33/71

A. candida
CCG cluster

Sequence
similar cluster
Singleton
Structural
similarity

Seq cluster 18
with AvrSen1

Seq cluster 201
with CCG 30

S. endobioticum
Avrsen1-like cluster

(b)

RAYH motif

CCG motif

(c)

Disulfide
bridges

A. candida CCG30
(A0A024FTM3)

Cys 73   Cys 19

Cys 67

Cys 22

(d) RAYH motif

A0A507BZL2
seq cluster 2

A0A507C8C0
seq cluster 40

A0A507CJ21
seq cluster 96

A0A507CLL0
seq cluster 98

A0A507CHN9
seq cluster 306

A0A507D0R9
seq cluster 69