Original Manuscript

Title Gene Age Gap Estimate (GAGE) for major depressive disorder: a penalized biological age

model using gene expression

[1]Yijie (Jamie) Li, [2]Rayus Kuplicki, [3]Bart N. Ford, [1]Elizabeth Kresock, [2]Leandra Figueroa-Hall,

[2,4]Jonathan Savitz, [1,5,*]Brett A. McKinney

[1]Tandy School of Computer Science, The University of Tulsa, Tulsa, OK, USA

[2]Laureate Institute for Brain Research, Tulsa OK, USA
10
11   [3]Department of Pharmacology and Physiology, Oklahoma State University Center for Health
12   Sciences, Tulsa, OK, USA.
13
14   [4]Oxley College of Health and Natural Sciences, The University of Tulsa, Tulsa OK, USA.
15
16   [5]Department of Mathematics, The University of Tulsa, Tulsa, OK, USA

17   [*]corresponding, brett-mckinney@utulsa.edu

19
20

Abstract

Recent associations between Major Depressive Disorder (MDD) and measures of premature

aging suggest accelerated biological aging as a potential biomarker for MDD susceptibility or

MDD as a risk factor for age-related diseases. Residuals or "gaps" between the predicted

biological age and chronological age have been used for statistical inference, such as testing

whether an increased age gap is associated with a given disease state. Recently, a gene

expression-based model of biological age showed a higher age gap for individuals with MDD

compared to healthy controls (HC). In the current study, we propose an approach that simplifies

gene selection using a least absolute shrinkage and selection operator (LASSO) penalty to

construct an expression-based Gene Age Gap Estimate (GAGE) model. We train a LASSO gene

age model on an RNA-Seq study of 78 unmedicated individuals with MDD and 79 HC, resulting

in a model with 21 genes. The L-GAGE shows higher biological aging in MDD participants than

HC, but the elevation is not statistically significant. However, when we dichotomize

chronological age, the interaction between MDD status and age has a significant association with

L-GAGE. This effect remains statistically significant even after adjusting for chronological age

and sex. Using the 21 age genes, we find a statistically significant elevated biological age in

MDD in an independent microarray gene expression dataset. We find functional enrichment of

infectious disease and SARS-COV pathways using a broader feature selection of age related

genes.

1. Introduction

Major depressive disorder (MDD) has been hypothesized to show characteristics of premature aging (Ford and Savitz (2022)). Biological aging can be measured in multiple dimensions such as telomere length, immunosenescence, brain volume, and gene expression. These measures of biological aging are correlated with chronological age, but environmental and genetic factors can increase or decrease an individual's biological age relative to their chronological age and influence their risk for age related diseases. For example, MDD has been associated with markers of cellular and immune aging including shortened leukocyte telomere length (Darrow et al., 2016; Ridout et al., 2016), elevated indicators of oxidative stress (Ait Tayeb et al., 2023), and elevated circulating inflammatory cytokines (Raison et al., 2006). Epigenetic clocks predicting biological age based on the accumulation of methylated CpG sites have found higher biological age in MDD participants compared with healthy controls (Protsenko et al., 2021). Brain age models constructed from T1-weighted magnetic resonance image (MRI) data from 2,188 healthy controls predicted a gap of +1.08 years (SE 0.22) between predicted and chronological age across 2,675 depressed participants (Han et al., 2021).

A recent RNA-Seq MDD study found that gene expression-based biological aging was elevated in MDD participants compared to Healthy Controls (HC) (Cole et al., 2021). The PBMC samples included four groups: 44 healthy controls and a mixture of MDD participants: 94 treatment-resistant, 47 treatment-responsive, and 46 untreated (Cole et al., 2021). They selected age genes iteratively by varying the P-value threshold for the t-test between upper and lower chronological age quartiles. For a given iteration, a biological age was computed for each subject based on the signed z-score of the age-related genes, and the P-value threshold was chosen to optimize the

correlation between biological and chronological age of the participants (Spearman Correlation

Coefficient (SCC) = 0.72, p < 0.01). A linear model of biological age was fit to chronological

age and association with MDD was computed by comparing the number of MDD and HC

participants above and below the regression line.

In the current study, we describe a different gene age approach that simplifies the gene selection

procedure by modeling age from RNA-Seq gene expression using a multivariate LASSO

penalized regression rather than an iterative univariate test. The LASSO approach has the

potential to capture more variation because it is multivariate, plus it automates gene selection by

cross-validation of the penalty and it reduces the amount of correlation in the selected features.

We use age as a quantitative variable during the LASSO feature selection in linear regression, as

opposed to using age quartiles, which is another way for the model to include more variation

when estimating age. After training the gene age model, we dichotomize chronological age when

using it as a covariate for the association of the gene age gap with MDD.

The current study is outlined as follows. We describe the LASSO biological age model trained

on an existing RNAseq dataset consisting of 157 individuals (78 with MDD and 79 healthy

controls) (Li et al., 2022). The residual is an estimate of the gap between an individual's

chronological age and their biological gene age, which we refer to as the LASSO Gene Age Gap

Estimate (L-GAGE). We describe the use of L-GAGE for testing elevated biological aging in

MDD. We find L-GAGE is elevated in MDD participants compared to HC, but the elevation is

not statistically significant. However, when we dichotomize chronological age into older and

younger, the interaction between MDD status and age is significantly associated with the L-

GAGE residual. We use the top L-GAGE genes to train a gene age model in an independent

public dataset for MDD, and the residual shows a statistically significant increase in MDD

compared to HC. Finally, we use machine learning feature selection to explore biological

pathways that are significantly enriched for the gene sets identified as being associated with

aging.

2. Materials and Methods

2.1. Gene Expression Data

To train our biological age models, we use an extant RNA-Seq dataset (Le et al., 2020). The

study was approved by the Western Institutional Review Board and conducted according to

the principles expressed in the Declaration of Helsinki. The data consists of 78 MDD and 79 HC

participants (91 females and 66 males). Individuals with current symptoms of depression met

DSM-IV-TR criteria for MDD based on the Structural Clinical Interview for DSM-IV-TR Axis I

Disorders and an unstructured psychiatric interview. HC individuals had no personal or

immediate family history of major psychiatric disorders. MDD participants were unmedicated

for at least 3 weeks prior to study entry. Exclusion criteria included major medical or

neurological illness, psychosis, traumatic brain injury, and a history of drug/alcohol abuse within

1 year. There is a higher female/male ratio for MDD (51/27) than HC (40/39), compatible with

trends in the general population. The age distribution is slightly skewed towards younger

individuals with age range from 18 to 55 (Fig. 1). The 8,923 genes in the RNA-Seq gene

expression data are normalized by counts per million reads, which we then quantile normalize

and log2 transform to stabilize variance. We removed genes with a low coefficient of variation

(standard deviation divided by absolute mean). We chose a threshold of 0.045 to obtain 5,587

genes.

To test the generalizability of the gene age model, we use an independent microarray study of

MDD from the gene expression omnibus (GEO) with accession number GSE98793 (Leday et al.,

2018). This data skews older than the discovery data (ages ~30–70 years) and includes MDD

with anxiety. We exclude anxiety, resulting in 64 participants with MDD and 64 HC.

Gene Age Gap Estimate (GAGE)

We use LASSO for gene selection and modeling biological age, and then we use the residual of

this model, which we call LASSO Gene-Age Gap Estimate (L-GAGE), for association testing

with MDD. For the LASSO biological aging model, we build a full penalized regression model

with all gene expression variables and with chronological age as the outcome variable. We

include both MDD and HC samples in the age model, which was also the approach in Ref. (Cole

et al., 2021). Our biological age model is based on the non-zero coefficient genes from the

lambda-1se LASSO penalty (the largest λ for which the average cross-validation (CV) error is

within one standard error of the minimum CV error). We compute the gap/residuals of the

LASSO model between predicted biological age and chronological age (i.e., the L-GAGE score).

Our goal is to use L-GAGE to test for increased biological age in MDD participants (Fig. 2).

For the replication microarray dataset, we retrain the gene-age regression coefficients for the top

L-GAGE genes because we could not match two of the gene symbols that are part of the

multivariate L-GAGE model from the RNA-Seq data. We use Ridge rather than LASSO because

we first perform feature selection in the discovery data and LASSO may force too many genes to

zero. We test Ridge residual (R-GAGE) for association with MDD in the microarray data.

2.2. Relationship between gene age gap, chronological age, MDD and sex

It is important to consider adjustments for chronological age in biological age models because of

regression to the mean as discussed for brain age models (Le et al., 2018), but sex is also an

important covariate for MDD. To further explore covariate effects, we add MDD x Age and

MDD x Sex interactions for L-GAGE associations with MDD. We use the OLS model

$$LGAGE = \beta_0 + \beta_1 MDD + \beta_2 Z + \beta_3 (MDD * Z) + \varepsilon, \qquad \text{(Eq. 1)}$$

where Z represents the adjustment or interaction variable (Age or Sex). We focus on the effect

of β_3, which represents how much the average L-GAGE of the MDD group changes for the Z=1

condition.

We consider two cases when age is used as a covariate with interactions (Z in Eq. 1): as

continuous and as dichotomous with a threshold. To verify our choice of age threshold, we use a

threshold regression model in the "chngpt" package in R (Fong et al., 2017). We use this

approach to check for possible nonlinear relationship between MDD and age and whether the

effect of chronological age on MDD increases at some threshold point. The mean function of the

threshold model is:

$$\eta = \alpha_1 + \alpha_2 z + \beta_1 I(x > e), \qquad \text{(Eq. 2)}$$

where x stands for chronological age, e is the age threshold and z are additional predictors. "I" is

a step indicator function. The threshold is optimized using the exact criterion function with a

logistic-based smooth function.

2.3. Feature selection, Gene-Age pathway Enrichment, and interpretable classifier

We use LASSO to create the gene-based residual age model, L-GAGE, but LASSO feature

selection also results in a set of age-related genes. As a secondary analysis, we use LASSO and

other feature selection methods to identify important age-related genes for pathway enrichment

to understand the biological mechanisms of the age models. We use univariate linear regression,

random forest (RF) regression, and nearest-neighbor projected distance regression (NPDR) (Le

et al., 2020) as feature selection methods. RF has the ability to find more complex models than

LASSO and linear regression, but RF has limited ability to detect interactions (McKinney et al.,

2009), whereas NPDR has the ability to detect interaction effects (Le et al., 2020). For univariate

feature selection, we use a linear model of individual genes with age, and we use a P-value

threshold of 0.05 (uncorrected for improved pathway overlap). We use the standard NPDR with

an adjusted P-value threshold of 0.05 FDR, and we use the LASSO penalized NPDR. For NPDR,

we use the imbalanced k-nearest-neighbor value (k=47) that approximates the 0.5 standard

deviation of the hyper-radius (Le et al., 2020). We use permutation variable importance with RF.

We use the Reactome Pathway database in MSigDB (Subramanian et al., 2005) for biological

pathway enrichment of age related genes.

For additional interpretation of the gene-age prediction of MDD along with consideration for

other covariates, we train a decision tree to predict MDD based on L-GAGE, chronological age,

and sex. Decision trees have high variance, but they are useful for interpreting the relationships

between covariates.

3.   Results and Discussion

3.1. Association Testing of Gene Age L-GAGE with MDD.

We test for association of the LASSO Gene Age Gap Estimate (L-GAGE) score with MDD

status. L-GAGE is the residual from a LASSO gene expression model of chronological age. The

LASSO model uses the cross-validation tuned lambda-1se value ($\lambda = 1.636048$), which is the

largest λ at which the mean-squared error (MSE) is within one standard error of the minimum

MSE. The residuals are constant, and heteroscedasticity is not present based on the Non-constant

Variance Score Test. The penalty results in a multivariate linear model of age with 22 genes and

a Spearman Correlation Coefficient (SCC) with chronological age of 0.77 (Fig. 2). Counting the

number of HC or MDD above or below the regression line (Fig. 2), we find that the biological

age is greater in MDD participants than HC (HC – 45 (56.96 %) below, 34 (43.037%) above,

MDD 35 (44.87%) below, 43 (55.128%) above). The P-value of the Chi-squared test of GAGE

sign (above or below the line) for MDD is not significant (0.1753). The greater L-GAGE in

MDD versus HC can be seen in L-GAGE density (Fig. 3A). The L-GAGE distribution for males

and females is very similar (Fig. 3B). While L-GAGE is greater in MDD than HC participants,

we do not find a statistically significant replication of the effect found in Ref. (Cole et al., 2021).

However, we do see a comparable effect size to what was previously found. Using the same

genes as their model also did not yield a statistically significant MDD association.

3.2 Testing MDD-Age interaction for L-GAGE association model.

We test for the effect of L-GAGE on MDD by introducing an MDD-Age interaction term (Eq. 1).

Dichotomizing age at threshold 40, MDD alone is not significant, but we find a statistically

significant effect of the interaction between MDD and Age 40 on L-GAGE (Table 1 and Fig. 4).

For individuals younger than 40, L-GAGE shows very little difference between MDD and HC,

but for older individuals, there is greater biological aging (L-GAGE) for the MDD versus HC

group (Table 1 and Fig. 4). Age alone is also statistically significant (Table 1). These age effects

remain significant when we add sex as a covariate (Table 1B), but sex is not significant (Table

1B and Table 2).

The MDD-Age interaction and the MDD term (Eq. 1) do not have a significant effect on L-

GAGE when age is treated as a continuous variable (MDD P-value = 0.364, Age P-value = 0.316,

MDD*Age P-value = 0.197). Also, there is no direct statistical association between MDD and

age and between MDD and sex (Two Sample T-test of MDD and Chronological age: P-value =

0.167; Chi-squared-test of MDD and sex: P-value = 0.08716). To further support our choice of

age threshold, we use a threshold regression (Eq. 2). The change point for age in relation to

MDD is estimated to be 39 years (Fig. 5). Combined with the third quartile being age 41, the

threshold regression suggests that age 40 is a suitable cutoff point for dividing the participants

into two age groups.

Additional support for the age-40 threshold can be seen in the decision tree for predicting MDD

(Fig. 6), where age with threshold 39.5 is the second important split variable, following L-GAGE.

The decision tree also suggests interaction effects, where the effect of L-GAGE on MDD is

conditioned on chronological age. If L-GAGE (node 1, Fig. 6) is below a threshold, participants

tend to be HC. If the L-GAGE is below the threshold and chronological age is above 39.5 (i.e.,

an interaction), participants tend to be MDD. However, for chronological age less than 39.5.

(node 3, Fig. 6), the prediction of MDD is considerably more complex. We note that this

decision tree was trained on the full dataset to maximize power, but it is instructional for

interpretation.

3.3 Characterizing Age-Associated Genes

The LASSO regression used in L-GAGE selected 21 age genes with non-zero coefficients (Table

3). In order to perform pathway enrichment for age related genes, we expand the number of

genes and feature selection methods to include linear regression, RF, and nearest-neighbor

projected distance regression (NDPR) (Le et al., 2020). Using the Reactome database, we find

enrichment for Infectious Disease, Adaptive Immune System, and SARS-CoV-2 Infection

pathways for top genes with P-value<0.05 from linear regression (Table 5) and NPDR (Table 6).

SARS-CoV-2 can cause neurological complications, and a recent study showed that

differentially expressed genes for COVID infection overlap with many gene associations for

neuropsychiatric disorders including depression (Quincozes-Santos et al., 2021). We also found

across all feature selection methods (including LASSO), the four common age genes are NAA20

(N-alpha-acetyltransferase 20), CCNE1 (Cyclin E1), and SESTD1 (SET domain containing

protein 1A), and TAF9 (TATA-box-binding protein associated factor 9). These genes will be

discussed further.

3.4 Association Test of L-GAGE in independent MDD study

To evaluate the generalizability of the 21 genes from the L-GAGE model, we train a Ridge

regression model of age using a microarray study by (Leday et al., 2018). Out of the 21 genes in

our L-GAGE model, 19 were found in the microarray dataset by matching gene symbols (genes

ANTXRL and UBE2F-SCLY were not present). Using the 19 genes in the microarray data with

64 MDD and 64 HC, the Ridge model of chronological age (lambda penalty 1.991) has

SCC=0.465 (Fig. 7).   In the scatter plot of Ridge gene-age versus chronological age (Fig. 7)

41(23) MDD are above(below) the regression line and 23(41) HC are above(below) the

regression line. The elevated Ridge Gene Age Gap Estimate (R-GAGE residual) in MDD

compared to HC is statistically significant (Fig. 8, t-test P-value = 0.000186). Also note there is

no association between MDD/HC and chronological age (t-test P-value=0.9756).

Conclusion

We presented a procedure for creating an expression-based biological age model using LASSO

penalized regression, and we explored the association between the residual (the LASSO-based

Gene Age Gap Estimate or L-GAGE) and MDD while adjusting for chronological age and sex.

We found increased biological aging based on L-GAGE in MDD versus HC participants with an

effect size similar to a previous study (Cole et al., 2021), but the difference was not statistically

significant (see discussion below about replication). We found a statistically significant MDD-

Age interaction for L-GAGE when age is dichotomized with threshold 40 years. We used

multiple statistical criteria to verify this threshold based on the age distribution. Using a higher

threshold results in a very sparse older group. The MDD-Age interaction could indicate an effect

of lifetime number of MDD episodes on biological aging that is not detectable until middle-age.

The interaction effect remained significant when adjusting for chronological age and sex, and we

emphasize the importance of including age as a covariate in biological age association tests to

avoid confounding due to regression to the mean (Le et al., 2018).

Elevated gene aging found previously in MDD participants was also observed in our data based

on L-GAGE but was not statistically significant. However, we found evidence for the

generalizability of the L-GAGE model by using the 21 LASSO-selected genes to test for

elevated gene age in an independent microarray gene expression dataset (recall the discovery

data used RNA-Seq). A limitation of this replication is the retraining of the model coefficients,

which was necessary because only 19 of the 21 genes could be reliably mapped to microarray

probes. Also, distribution differences between platforms changes the scale of the original

regression coefficients. And so, while we did retrain the coefficients for microarray data, we

restricted ourselves to the same genes selected in the discovery cohort, and we found a

statistically significant elevation of gene age in MDD compared to HC. The age distributions are

also somewhat different, with the discovery set skewing younger and the replication set skewing

older. An interesting future work would be to integrate these data and other cohorts to get a

broader age range with a gene age model with more generalizability.

We note a few of the top age-associated genes, such as CCNE1, NAA20, SESTD1, and TAF9,

that have been associated with aging, senescence, and infectious disease. In a study of Lung

Adenocarcinoma, CCNE1 gene expression was found to be correlated with patients' age (Ullah

et al., 2022), and NAA20 and SETD1A are involved in senescence, which is related to aging and

age-related diseases. It was shown that depletion of NAA20 in non-transformed mammal cells

led to senescence (Elurbide et al., 2023), and in another study knockdown of SETD1A triggered

cellular senescence. (Tajima et al., 2019). TAF9 cross-reactivity was shown to be associated with

immunity to CMV (human cytomegalovirus) in the context of autoimmune disease (Chen et al.,

2021).

Pathway enrichment of a larger set of age genes, beyond the 21 L-GAGE genes, resulted in the

detection of Infectious Disease, Adaptive Immunity, and SARS-CoV Infection pathways. As

noted in (Cole et al., 2021), evaluating PBMC transcription can increase the risk for false

positive immune pathways. Future work will involve pathway analysis based on other biological

specimens and testing of the gene age model for MDD in RNA-seq data from postmortem brain

areas. For example, the Stanley Medical Research Institute data includes frontal and cingulate

cortex and hippocampus as well as several subcortical areas. Brain region-specific gene age

effects on MDD could provide valuable insights into the etiology of MDD.

This study contributes a new approach to estimating biological aging and contributes to the

evidence for the role of aging and inflammation in depression. Future studies are needed with

broader age ranges, more uniform age distributions, large sample sizes, and utilization of MDD

age-of-onset and number of depressive episodes. Future gene age models may help identify

individuals that need different treatment or management for depression due to an increase in their

relative biological age.

Research data for this article

Data and code for this research are available at https://github.com/insilico/GeneAgeMDD.

Funding

BAM and JS received support from the National Institute of Mental Health (R01MH098099).

Reference

Ait Tayeb, A.E.K., Poinsignon, V., Chappell, K., Bouligand, J., Becquemont, L., Verstuyft, C., 2023. Major Depressive Disorder and Oxidative Stress: A Review of Peripheral and Genetic Biomarkers According to Clinical Characteristics and Disease Stages. Antioxidants 12(4), 942. doi:10.3390/antiox12040942

Chen, Y.F., Hsieh, A.H., Wang, L.C., Yu, K.H., Kuo, C.F., 2021. Cytomegalovirus-Associated Autoantibody against TAF9 Protein in Patients with Systemic Lupus Erythematosus. J Clin Med 10(16). DOI: 10.3390/jcm10163722

Cole, J.J., McColl, A., Shaw, R., Lynall, M.E., Cowen, P.J., de Boer, P., Drevets, W.C., Harrison, N., Pariante, C., Pointon, L., consortium, N., Goodyear, C., Bullmore, E., Cavanagh, J., 2021. No evidence for differential gene expression in major depressive disorder PBMCs, but robust evidence of elevated biological ageing. Transl Psychiatry 11(1), 404. DOI: 10.1038/s41398-021-01506-4

Darrow, S.M., Verhoeven, J.E., Révész, D., Lindqvist, D., Penninx, B.W., Delucchi, K.L., Wolkowitz, O.M., Mathews, C.A., 2016. The Association Between Psychiatric Disorders and Telomere Length: A Meta-Analysis Involving 14,827 Persons. Psychosom Med 78(7), 776-787. DOI: 10.1097/psy.0000000000000356

Elurbide, J., Carte, B., Guedes, J., Aldabe, R., 2023. NatB Catalytic Subunit Depletion Disrupts DNA Replication Initiation Leading to Senescence in MEFs. Int J Mol Sci 24(10). DOI: 10.3390/ijms24108724

Fong, Y., Huang, Y., Gilbert, P.B., Permar, S.R., 2017. chngpt: threshold regression model estimation and inference. BMC Bioinformatics 18(1), 454. DOI: 10.1186/s12859-017-1863-x

Ford, B.N., Savitz, J., 2022. Depression, aging, and immunity: implications for COVID-19 vaccine immunogenicity. Immunity & Ageing 19(1), 32. DOI: 10.1186/s12979-022-00288-7

Han, L.K.M., Dinga, R., Hahn, T., Ching, C.R.K., Eyler, L.T., Aftanas, L., Aghajani, M., Aleman, A., Baune, B.T., Berger, K., Brak, I., Filho, G.B., Carballedo, A., Connolly, C.G., Couvy-Duchesne, B., Cullen, K.R., Dannlowski, U., Davey, C.G., Dima, D., Duran, F.L.S., Enneking, V., Filimonova, E., Frenzel, S., Frodl, T., Fu, C.H.Y., Godlewska, B.R., Gotlib, I.H., Grabe, H.J., Groenewold, N.A., Grotegerd, D., Gruber, O., Hall, G.B., Harrison, B.J., Hatton, S.N., Hermesdorf, M., Hickie, I.B., Ho, T.C., Hosten, N., Jansen, A., Kähler, C., Kircher, T.,

Klimes-Dougan, B., Krämer, B., Krug, A., Lagopoulos, J., Leenings, R., MacMaster, F.P., MacQueen, G., McIntosh, A., McLellan, Q., McMahon, K.L., Medland, S.E., Mueller, B.A., Mwangi, B., Osipov, E., Portella, M.J., Pozzi, E., Reneman, L., Repple, J., Rosa, P.G.P., Sacchet, M.D., Sämann, P.G., Schnell, K., Schrantee, A., Simulionyte, E., Soares, J.C., Sommer, J., Stein, D.J., Steinsträter, O., Strike, L.T., Thomopoulos, S.I., van Tol, M.J., Veer, I.M., Vermeiren, R., Walter, H., van der Wee, N.J.A., van der Werff, S.J.A., Whalley, H., Winter, N.R., Wittfeld, K., Wright, M.J., Wu, M.J., Völzke, H., Yang, T.T., Zannias, V., de Zubicaray, G.I., Zunta-Soares, G.B., Abé, C., Alda, M., Andreassen, O.A., Bøen, E., Bonnin, C.M., Canales-Rodriguez, E.J., Cannon, D., Caseras, X., Chaim-Avancini, T.M., Elvsåshagen, T., Favre, P., Foley, S.F., Fullerton, J.M., Goikolea, J.M., Haarman, B.C.M., Hajek, T., Henry, C., Houenou, J., Howells, F.M., Ingvar, M., Kuplicki, R., Lafer, B., Landén, M., Machado-Vieira, R., Malt, U.F., McDonald, C., Mitchell, P.B., Nabulsi, L., Otaduy, M.C.G., Overs, B.J., Polosan, M., Pomarol-Clotet, E., Radua, J., Rive, M.M., Roberts, G., Ruhe, H.G., Salvador, R., Sarró, S., Satterthwaite, T.D., Savitz, J., Schene, A.H., Schofield, P.R., Serpa, M.H., Sim, K., Soeiro-de-Souza, M.G., Sutherland, A.N., Temmingh, H.S., Timmons, G.M., Uhlmann, A., Vieta, E., Wolf, D.H., Zanetti, M.V., Jahanshad, N., Thompson, P.M., Veltman, D.J., Penninx, B., Marquand, A.F., Cole, J.H., Schmaal, L., 2021. Brain aging in major depressive disorder: results from the ENIGMA major depressive disorder working group. Mol Psychiatry 26(9), 5124-5139. DOI: 10.1038/s41380-020-0754-0

Le, T.T., Dawkins, B.A., McKinney, B.A., 2020. Nearest-neighbor Projected-Distance Regression (NPDR) for detecting network interactions with adjustments for multiple tests and confounding. Bioinformatics 36(9), 2770-2777. DOI: 10.1093/bioinformatics/btaa024

Le, T.T., Kuplicki, R.T., McKinney, B.A., Yeh, H.W., Thompson, W.K., Paulus, M.P., Tulsa, I., 2018. A Nonlinear Simulation Framework Supports Adjusting for Age When Analyzing BrainAGE. Front Aging Neurosci 10, 317. DOI: 10.3389/fnagi.2018.00317

Leday, G.G.R., Vértes, P.E., Richardson, S., Greene, J.R., Regan, T., Khan, S., Henderson, R., Freeman, T.C., Pariante, C.M., Harrison, N.A., Perry, V.H., Drevets, W.C., Wittenberg, G.M., Bullmore, E.T., 2018. Replicable and Coupled Changes in Innate and Adaptive Immune Gene Expression in Two Case-Control Studies of Blood Microarrays in Major Depressive Disorder. Biol Psychiatry 83(1), 70-80. DOI: 10.1016/j.biopsych.2017.01.021

Li, Y.J., Kresock, E., Kuplicki, R., Savitz, J., McKinney, B.A., 2022. Differential expression of MDGA1 in major depressive disorder. Brain Behav Immun Health 26, 100534. DOI: 10.1016/j.bbih.2022.100534

McKinney, B.A., Crowe, J.E., Guo, J., Tian, D., 2009. Capturing the spectrum of interaction effects in genetic association studies by simulated evaporative cooling network analysis. PLoS Genet 5(3), e1000432. DOI: 10.1371/journal.pgen.1000432

Protsenko, E., Yang, R., Nier, B., Reus, V., Hammamieh, R., Rampersaud, R., Wu, G.W.Y., Hough, C.M., Epel, E., Prather, A.A., Jett, M., Gautam, A., Mellon, S.H., Wolkowitz, O.M., 2021. "GrimAge," an epigenetic predictor of mortality, is accelerated in major depressive disorder. Translational Psychiatry 11(1), 193. DOI: 10.1038/s41398-021-01302-0

Quincozes-Santos, A., Rosa, R.L., Tureta, E.F., Bobermin, L.D., Berger, M., Guimaraes, J.A.,
Santi, L., Beys-da-Silva, W.O., 2021. COVID-19 impacts the expression of molecular markers
associated with neuropsychiatric disorders. Brain Behav Immun Health 11, 100196. DOI:
10.1016/j.bbih.2020.100196

Raison, C.L., Capuron, L., Miller, A.H., 2006. Cytokines sing the blues: inflammation and the
pathogenesis of depression. Trends Immunol 27(1), 24-31. DOI: 10.1016/j.it.2005.11.006

Ridout, K.K., Ridout, S.J., Price, L.H., Sen, S., Tyrka, A.R., 2016. Depression and telomere
length: A meta-analysis. J Affect Disord 191, 237-247. DOI: 10.1016/j.jad.2015.11.052

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., Mesirov, J.P., 2005. Gene set
enrichment analysis: A knowledge-based approach for interpreting genome-wide expression
profiles. Proceedings of the National Academy of Sciences 102(43), 15545-15550. DOI:
doi:10.1073/pnas.0506580102

Tajima, K., Matsuda, S., Yae, T., Drapkin, B.J., Morris, R., Boukhali, M., Niederhoffer, K.,
Comaills, V., Dubash, T., Nieman, L., Guo, H., Magnus, N.K.C., Dyson, N., Shioda, T., Haas,
W., Haber, D.A., Maheswaran, S., 2019. SETD1A protects from senescence through regulation
of the mitotic gene expression program. Nature Communications 10(1), 2854. DOI:
10.1038/s41467-019-10786-w

Ullah, M.A., Farzana, M., Islam, M.S., Moni, R., Zohora, U.S., Rahman, M.S., 2022.
Identification of the prognostic and therapeutic values of cyclin E1 (CCNE1) gene expression in
Lung Adenocarcinoma and Lung Squamous Cell Carcinoma: A database mining approach.
Heliyon 8(9), e10367. DOI: 10.1016/j.heliyon.2022.e10367

457 **Figures**
458

459
460
461 **Figure 1. Histogram of chronological ages with a bin size of 1:** Bars are separated by Healthy
462 Control (HC, red) and major depressive disorder (MDD, blue). There are more younger
463 participants in the dataset with the same age, especially from age 20~28. For example, there are

464   15 participants that are 24 years of age. Chronological age is not associated with MDD versus
465   HC (T-test P-value 0.167).
466

467
468   **Figure 2. Scatter plot with regression line of biological age and chronological age:**
469   Biological age model is based on LASSO regression and the residual is later used for LASSO
470   Gene Age Gap Estimate (L-GAGE). The points are colored by MDD (blue) and HC (red)**.** The
471   points are shaped by Female (circle) and Male (triangle)**.** Spearman Correlation Coefficient (SCC
472   = 0.77, slope P-value < 0.01)**.**
473
474
475
476

477
478
479 **Figure 3. Density plots of the LASSO based Gene Age Gap Estimate (L-GAGE) separated**
480 **by MDD (A) and sex (B)**. A positive gene-age residual (x-axis) indicates a sample above the
481 gene age regression line and negative below. **A**. Biological age relative to chronological age (L-
482 GAGE) is greater in MDD participants than in HC. **B**. The L-GAGE difference between males
483 and females is less pronounced.

484
485
486
487

488
489
Figure 4. MDD x Age interaction for L-GAGE with age 40 threshold. A. The average L-
GAGE for people older than 40 with MDD is higher than the L-GAGE value for people younger
than 40 with MDD (blue line), whereas in the HC group the average L-GAGE is lower for
people older than 40 than for people younger than 40 (red line). **B.** For individuals younger than
40, L-GAGE shows very little difference between MDD and HC. For older individuals, there is
greater biological aging (L-GAGE) for the MDD versus HC group. The L-GAGE association
with MDD is still significant when adjusted by age and sex.

497
498

Figure 5. Effect of Chronological Age on MDD Determined by Threshold Regression Model. A. Threshold regression (Eq. 2) shows the nonlinear relationship between MDD and chronological age. The prediction indicates an increase in MDD up to the age of 39, which is identified as the change point by the model. **B.** The likelihood analysis of the threshold regression model also indicates that age 39 is the optimal threshold, having the highest model likelihood.

510
Figure 6. Gene age decision tree for MDD with covariates. For added interpretation, we train a decision tree on all samples to predict MDD. The model identifies the gene age residual L-GAGE as the most important predictor, with chronological age being the second most significant factor. In the first split, if the gene age gap is low, L-GAGE < -2.251 (Node 1), there is high probability for a subject to be HC (Node 2). If the gene age gap is higher, L-GAGE ≥ -2.251, the model becomes more complex and initially depends on chronological age with split 39.5 years (Node 3). If L-GAGE is high and Age ≥ 39.5, then there is a high probability a subject is MDD (Node 15). When Age < 39.5, the model again becomes dependent on L-GAGE, and at a certain split, females exhibit a higher probability of MDD compared to males (Nodes 8 and 9).

Figure 7. Gene age versus chronological age in replication microarray data (GSE98793).
Gene age regression line (blue line, Spearman correlation coefficient (SCC) = 0.46, P-value <
0.01) is based on Ridge regression in the replication microarray data using only the 19 genes
from the original L-GAGE model (21 genes less 2 missing). The points are colored by MDD
(blue) and HC (red) and shaped by Female (circle) and Male (triangle). For MDD, 41(23) are
above(below) the regression line and for HC, 23(41)are above(below) the regression line. The
Ridge Gene Age Gap Estimate (R-GAGE) is higher for MDD than HC participants (see Fig. 8, t-
test P-value = 0.0001856).

556
557

558
**Figure 8. Density plot of the Ridge based Gene Age Gap Estimate (R-GAGE) in replication
microarray data (GSE98793).** The Ridge age model (Fig. 7) is trained on the replication
microarray data using19 genes from the original L-GAGE model. A positive R-GAGE residual
(x-axis) indicates a sample above the gene age regression line and negative below (Fig. 7).
Biological age relative to chronological age (R-GAGE) is higher in MDD (blue) participants
than in HC (red) (T-test P-value = 0.0001856).

579 **Tables**
580

581 **Table 1**

582 *LASSO Gene-age-gap estimate (L-GAGE) association with MDD and dichotomized age*

583 *interaction.*

584 **A**

	Estimate	Std. Error	t value	Pr(>|t|)
(Intercept)	0.1225	0.3495	0.35	0.7265
MDD	-0.1286	0.5203	-0.247	0.8052
Age40	-1.606	0.7535	-2.131	0.0347*
MDD*Age40	2.2764	0.9984	2.28	0.024*
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1				

586 **B**

	Estimate	Std. Error	t value	Pr(>|t|)
(Intercept)	0.2644	0.4172	0.634	0.5273
MDD	-0.1870	0.5296	-0.353	0.7246
SexMale	-0.2838	0.4534	-0.626	0.5324
Age40	-1.6144	0.7551	-2.138	0.0341*
MDD*Age40	2.3274	1.0038	2.319	0.0217*
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1				

587
588 *Note.* **A.** Based on the ordinary least squares model (Eq. 1 with Z=Age), where chronological age

589 is dichotomized with threshold is Age>=40 and Age<40, the MDD x Age interaction is

590 significant. Biological age (L-GAGE)) is similar for MDD and HC when Age<40, but when the

591 chronological age is higher than 40, biological age is significantly greater in MDD individuals

592 than HC. **B**. The MDD x Age interaction remains significant when Sex is added as a covariate.

602 **Table 2**

603 *Gene-age-gap regression with MDD-sex interaction with Female and Male.*

	Estimate	Std. Error	t value	Pr(>|t|)
(Intercept)	-0.24481	0.44221	-0.554	0.581
MDD	0.64637	0.5907	1.094	0.276
Male	0.04395	0.62938	0.07	0.944
MDD* Male	-0.55121	0.91608	-0.602	0.548
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1				

604 *Note.* Based on the ordinary least squares model (Eq. 1 with Z=Male/Female instead of age), L-

605 GAGE score of MDD in males is slightly lower than the L-GAGE score of MDD in females, but

606 the interaction term MDD*Male is not statistically significant.

608 **Table 3**

609 *Age associated genes selected by LASSO.*

Down Regulated with Increasing Age		Up Regulated with Increasing Age	
Gene	Coefficient	Gene	Coefficient
NAA20	-6.7070152	CCNE1	14.2689027
ZNF347	-2.9514771	SESTD1	8.8624231
PRMT6	-2.4559818	ZNF334	2.4209761
WDR13	-1.7979357	ANTXRL	2.0255277
DDX19B	-1.3737037	DTD2	1.8502139
TAF9	-1.2672137	CYTH3	1.5349361
ADSS	-1.1724134	DYRK1A	1.2905045
TGFBR3	-1.0316785	HTATSF1	1.078388
SMYD5	-0.8454683	SFXN4	0.7870119
CISD1	-0.6212633	UBE2F-SCLY	0.2252943
TGIF2-C20orf24	-0.5057642		

611 *Note.* Multivariate coefficients are shown that survived LASSO penalty. Negative coefficients

612 (left columns) indicate higher expression of the gene tends to occur with younger age. Positive

613 coefficients (right columns) indicate higher expression of the gene tends to occur in older

614 individuals. These genes are used in the gene age model and the L-GAGE residual.

620 **Table 4**

621 *Age associated genes selected by linear regression with adjusted P-value 0.05 FDR.*

Down Regulated with Increasing Age				Up Regulated with Increasing Age			
Gene	Coefficient	P-value	Adjusted P-value	Gene	Coefficient	P-value	Adjusted P-value
NAA20	-16.1918	8.86E-08	0.0005	CCNE1	42.8022	5.59E-07	0.0013
CIART	-22.7969	6.87E-07	0.0013	SESTD1	12.2045	1.19E-05	0.0111
TAF9	-21.2804	2.85E-06	0.0040	ITGB1BP1	10.7847	2.46E-05	0.0197
MLXIPL	-20.0949	4.55E-06	0.0051	ANTXRL	13.8739	4.23E-05	0.0295
TGFBR3	-17.7019	7.91E-05	0.0491				

622 *Note.* Negative coefficients (left columns) indicate higher expression of the gene tends to occur

623 with younger age. Positive coefficients (right columns) indicate that higher expression of the

624 gene tends to occur in older individuals. These genes are shown for comparison but not used in

625 the gene age model.

645   **Table 5**

646   *MSigDB Reactome results of the age genes selected by linear regression.*

Gene Set Name	Genes in Gene Set (K)	Description	Genes in Overlap (k)	k/K	p-value
REACTOME_RNA_POLYMERASE_II_TRANSCRIPTION	1393	RNA Polymerase II Transcription	46	0.0330	3.84E-11
REACTOME_POST_TRANSLATIONAL_PROTEIN_MODIFICATION	1442	Post-translational protein modification	44	0.0305	1.21E-09
REACTOME_METABOLISM_OF_RNA	714	Metabolism of RNA	29	0.0406	2.05E-09
REACTOME_TRANSCRIPTIONAL_REGULATION_BY_TP53	363	Transcriptional Regulation by TP53	20	0.0551	4.74E-09
REACTOME_INFECTIOUS_DISEASE	1019	Infectious disease	33	0.0324	3.95E-08
REACTOME_MEMBRANE_TRAFFICKING	629	Membrane Trafficking	23	0.0366	6.11E-07
REACTOME_METABOLISM_OF_LIPIDS	742	Metabolism of lipids	25	0.0337	8.86E-07
REACTOME_SUMOYLATION	187	SUMOylation	12	0.0642	1.19E-06
REACTOME_SARS_COV_INFECTIONS	471	SARS-CoV Infections	19	0.0403	1.4E-06
REACTOME_VESICLE_MEDIATED_TRANSPORT	724	Vesicle-mediated transport	23	0.0318	6.34E-06

648   *Note.* We collect the 464 age associated genes with P-value lower than 0.05 (not adjusted for

649   better pathway detection) and query MSigDB Reactome database for pathway enrichment.

650   Notably, these age associated genes are enriched for infectious disease and SARS-CoV

651   Infections pathways.

661   **Table 6**
662   *MSigDB Reactome results of the 145 age genes selected by nearest-neighbor projected distance*
663   *regression (NPDR) with LASSO penalty.*
664

Gene Set Name	Genes in Gene Set (K)	Description	Genes in Overlap (k)	k/K	p-value
REACTOME_NEF_ME DIATES_DOWN_MOD ULATION_OF_CELL_S URFACE_RECEPTORS _BY_RECRUITING_TH EM_TO_CLATHRIN_A DAPTERS	21	Nef-mediates down modulation of cell surface receptors by recruiting them to clathrin adapters	4	0.1905	7.44E-07
REACTOME_NEF_ME DIATED_CD4_DOWN_ REGULATION	9	Nef Mediated CD4 Down-regulation	3	0.3333	3.22E-06
REACTOME_RNA_PO LYMERASE_II_TRANS CRIPTION	1393	RNA Polymerase II Transcription	16	0.0115	2.38E-05
REACTOME_LDL_CLE ARANCE	19	LDL clearance	3	0.1579	3.62E-05
REACTOME_TRANSC RIPTIONAL_REGULA TION_BY_TP53	363	Transcriptional Regulation by TP53	8	0.022	3.78E-05
REACTOME_MHC_CL ASS_II_ANTIGEN_PRE SENTATION	126	MHC class II antigen presentation	5	0.0397	7.56E-05
REACTOME_ADAPTIV E_IMMUNE_SYSTEM	829	Adaptive Immune System	11	0.0133	1.38E-04
REACTOME_TRAFFIC KING_OF_AMPA_REC EPTORS	31	Trafficking of AMPA receptors	3	0.0968	1.63E-04
REACTOME_TP53_RE GULATES_METABOLI C_GENES	87	TP53 Regulates Metabolic Genes	4	0.046	2.32E-04

665