Short-term memory errors are strongly associated with a drift in

neural activity in the posterior parietal cortex

Joon Ho Choi[1,¶,*], Sungwon Bae[1,2,¶], Jiho Park[3], Minsu Yoo[1,3], Chul-Hoon Kim[2], L. Ian

Schmitt[4], Ji-Woong Choi[3], and Jong-Cheol Rah[1, 5,*]

[1]Sensory and Motor Systems Neuroscience Group, Korea Brain Research Institute, 61

Cheomdan-ro, Dong-Gu, Daegu, 41062, Korea [2]Department of Pharmacology, College of

Medicine, Yonsei University, 50 Yonsei-ro, Seodaemun-Gu, Seoul, 03722, Korea [3]Department

of Electrical Engineering and Computer Science, Daegu Gyeongbuk Institute of Science and

Technology (DGIST), 333 Techno Jungang-daero, Dalseong-gun, Daegu, 42988, Korea

[4]Center for Brain Science, RIKEN, Wako, Saitama, Japan. [5]Department of Brain Sciences,

DGIST, Daegu, 42988, Korea

[¶] These authors contributed equally

*Corresponding authors:

Joon Ho Choi

Sensory and Motor Systems Neuroscience Group, Korea Brain Research Institute, 61

Cheomdan-ro, Dong-Gu, Daegu 41062, Korea

joonchoi@kbri.re.kr

Jong-Cheol Rah

Sensory and Motor Systems Neuroscience Group, Korea Brain Research Institute, 61

Cheomdan-ro, Dong-Gu, Daegu 41062, Korea

jcrah@kbri.re.kr

Abstract

Understanding the neural mechanisms behind short-term memory (STM) errors is crucial for

unraveling cognitive processes and addressing related deficits in neuropsychiatric disorders. This

study investigates whether STM errors result from misrepresentation of sensory information or a

decay in these representations over time. Utilizing 2-photon calcium imaging in the posterior

parietal cortex (PPC) of mice engaged in a delayed match-to-sample task, we identified a subset

of PPC neurons exhibiting both directional and temporal selectivity. Contrary to the idea that

STM errors primarily stem from mis-encoding during the sample phase, our findings indicate

that these errors are more closely associated with a drift in neural activity during the delay

period. This drift results in a gradual shift away from the correct representation, ultimately

leading to incorrect behavioral responses. These results emphasize the importance of maintaining

stable neural representations in the PPC for accurate STM. Our findings also suggest that

targeting PPC activity stabilization during delay periods could be a potential therapeutic strategy

for mitigating cognitive impairments in disorders like schizophrenia.

Keywords:

Short-term memory, posterior parietal cortex, 2-photon calcium imaging, error trials, neural

activity

Introduction

Short-term memory (STM) is a capacity to retain immediately relevant information over a brief

period, typically on the order of seconds to minutes. While STM plays a pivotal role in a

multitude of cognitive functions, representations in this system are fragile and prone to errors in

information retention leading to sub-optimal behavior. Despite their behavioral impact, there is

currently no circuit-level understanding of how STM errors occur due to a lack of systematic

investigation. Gaining insight into the circuit mechanisms underlying STM errors is crucial for

comprehending information retention and its relevance to various psychiatric disorders,

including schizophrenia and attention deficit hyperactivity disorder, which frequently manifest

deficits in STM[1–3] This study aims to examine the circuit underpinnings of STM errors. To this

end, we considered two potential mechanisms which may underlie error generation: (1)

Inadequately encoding of information in the neural activity, and (2) Disruption of encoded

information during the delay period.

Along with the prefrontal cortex (PFC), the posterior parietal cortex (PPC) is a critical brain

region responsible for maintaining information during STM[4–6]. We chose to elucidate the activity

of the PPC for several reasons. As a major multimodal association cortex, the PPC is ideally

positioned to integrate multimodal sensory information and relay this accumulated information

to prefrontal areas[6,7]. Thus, if inadequate evidence accumulation or misperception were

represented in both the PFC and PPC during error trials, the PPC is more likely the source of the

erroneous activity that leads to the representation error. Furthermore, in both primates and

rodents, a subset of PPC neurons exhibits direction-predictive firing rate changes in a graded

manner, reflecting the clarity of evidence during a decision-making task such as the random dot

motion discrimination task or Poisson click task[8,9]. Meanwhile, neurons in the prefrontal motor

area demonstrate more categorized activity changes, suggesting that the cumulative evidence

information in the PPC transformed into a discrete decision in the PFC. Therefore, we

hypothesized that the gradual degradation of encoded information could be more readily

observed in the PPC compared with the PFC. By focusing on the PPC, we aimed to capture these

subtle variations in sensory encoding and retention.

Despite extensive evidence implicating the PPC as a substrate for STM, the nature of

information maintenance in this circuit remains incompletely understood. In a previous study,

mice were trained to navigate through a virtual T-maze by choosing a left or right turn according

to visual cues presented while PPC activity was measured. Since the cue was presented only in

the first half of each trial, the mice had to rely on their STM to perform this task. This study

revealed that the PPC exhibits a complex population encoding characterized by direction- and

phase-selective sequences of activity across multiple neurons[6]. However, given that PPC activity

shows navigational attributes, the PPC activity during the task cannot be interpreted as simply

reflecting visual STM[10–12]. Indeed, Krumin and colleagues have illustrated that sequential PPC

activity can be accurately predicted by the combination of heading direction and spatial location

within the virtual corridor during a comparable behavioral task[13]. Task-relevant increases or

suppressions in PPC activity have also been observed in mice performing a visually guided

delayed go/no-go task[6,14]. Notably, 'go' signals exhibited selectivity and were represented by a

substantial number of neurons, while 'no-go' signals were represented relatively weakly[6,14],

raising questions about whether the activities genuinely represented the retained visual signal

itself, or rather pertained to the suppression and release of impulsive movement[15].

To more directly investigate how retained sensory information during the delay period may be

encoded in PPC activity, and how it may be disrupted in errors, we recorded PPC activity while

mice performed a delayed match-to-sample task that did not require navigation. A large fraction

of the imaged neurons in PPC showed activity that reflected memory-guided decision-making,

and the activity pattern of neurons in this population were remarkably similar to those identified

in previous STM studies[6] consistent with a common mechanism of encoding for short-term

representations across tasks. Comparison of PPC activity between correct and incorrect trials in

this task revealed that STM errors were associated with weak initial encoding of the signal

followed by spontaneous drift towards incorrect representation during the delay. These findings

elucidate the circuit mechanisms underpinning errors in STM, providing insight into the nature

of this function and opening future avenues to study how it might be disrupted in

neuropsychiatric disorders.

Results

A sequence of choice-dependent neuronal activity in PPC

Our aim is to determine the extent to which PPC activity plays a pivotal role in maintaining

sensory information within the context of a STM-dependent perceptual decision-making task,

with an exclusion of spatial location and head-direction involvement. To achieve this, we trained

seven transgenic mice expressing GCaMP6f (C57BL/6J-Tg(Thy1-GCaMP6f)GP5.5Dkim/J) to

perform STM tasks. In brief, the mice were head-restrained, and gratings were presented on a

screen in front of them, moving either to the left or right direction in a pseudo-random manner.

Following around 2-second delay in which visual stimulation was omitted, a bifurcated lick port

was presented within the reach of the mice. Correct responses, licking the corresponding lick

port, were rewarded with approximately 5 μl of tap water, while incorrect responses, licking the

wrong lick port, led to an extended inter-trial interval of 2 seconds (time-out) (see Fig. 1A). After

thorough training, the mice successfully learned this task, achieving a discrimination rate

exceeding 70%. Importantly, mice did not exhibit a bias towards either direction after training

(Fig. 1B-C).

During task performance, we recorded the activities of PPC neurons by analyzing the

fluorescence changes in the GCaMP6f-expressing layer (L) 2/3 neurons with two-photon laser

scanning microscopy. The identification of the PPC region was based on stereotaxic coordinates

derived from mesoscopic connectivity analysis, which revealed distinct antero- and retrograde

connections compared to neighboring areas[16]. Active neurons were identified through *post hoc*

analysis using the CaImAn software[17]. While various types of responses were observed from the

PPC neurons, a large proportion of neurons consistently exhibited selectivity based on direction,

task phase, or both across trials (Fig. 1D). In addition to the remarkable response consistency

across trials, the time-wise correlation of mean response vectors, computed every 0.5s (15 image

frames) across all recorded neurons, demonstrated similarity within behavioral phases (Fig. 1E,

see Methods).

To objectively categorize active neurons based on their response patterns, we employed a

modified generalized linear model (GLM), as detailed in the Methods section[18]. The time-

correlation of the activity vectors from the neurons sorted by our GLM was nearly identical to

those from all the recorded neurons (Supplementary Fig. 1). This suggests that the phase-

dependent feature of the activity is well preserved in the GLM-sorted neurons. Our analysis

revealed that nearly half of the active neurons in the PPC satisfied the criteria for phase or

stimulus direction selectivity ($p < 0.01$), amounting to 2,664 out of 4,569 neurons

(Supplementary Fig. 2). Approximately 60% of these neurons (1,609/2,664 neurons, 873 for the

right signal and 736 for the left signal) exhibited directional selectivity, the remaining neurons

responded similarly to both directions (Supplementary Fig. 3A-C). Out of the 1,609 directionally

selective neurons, approximately 41% displayed phase selectivity, with 661 neurons exhibiting

phase selectivity during the sample phase, 315 during the delay phase. Consistent with previous

studies, PPC neurons rarely displayed sustained activity across multiple phases of the task [6] (Fig.

1F and Supplementary Fig. 4) with less than 8% (202/2664) of neurons demonstrating phase

selectivity during both the sample and delay phases.

When the neurons were sorted based on the onset of the peak responses, it became evident that a

directional-selective sequence of neuronal activity emerged with a remarkable similarity to the

findings of Harvey and colleagues [6] (Fig. 2A). This result underscores the replicability of the

directional-selective sequence of neural activity in the PPC within a behavioral paradigm that

excluded spatial navigation. To assess predictive capabilities of the GLM-selected neurons, the

direction selectivity of each neuron was verified through receiver operating characteristics

(ROC) analysis (Fig. 2C-D). It is noteworthy that while ROC analysis identifies neurons with

significant differences in condition-dependent $\Delta F/F$ amplitude, modified GLM analysis detects

neurons based on condition-dependent response selectivity. We introduced a preference metric,

preference index (PI), which was proportional to the hyperbolic area of the ROC curve,

calculated as (AUC-0.5)*2. The preference index of the neurons with temporal and directional

selectivity was significantly higher compared to the shuffled responses, particularly for the

delay-preferring neurons (Fig. 2C, D). It was noteworthy that over 30% of the phase- and

direction-selective neurons exhibited a preference index above 0.9, indicating a near-complete

separation in their responses to the two visual stimuli (Fig. 2B-C). Because PI was calculated

based on the averaged responses[19], we also sought to quantify the consistency of the responses

across trials. To this end, we constructed a reliability index (RI) proportional to the hyperbolic

area of the area under ROC curve from the $\Delta F/F$ of all trials (see methods). Similar to the PI, a

significantly high RI was observed in the direction-selective neurons compared to the shuffled

control or non-directional neurons (Supplementary Fig. 3). Interestingly, direction-selective

delay neurons showed a greater RI than sample neurons, suggesting refinement of the encoded

information following stimulus-offset (Supplementary Fig. 3D-I).

Phase-selective activities were further validated across the population by using principal

components analysis (PCA) to captures the temporal activity pattern of each neuron throughout

the entire behavioral session (Fig. 2E). The GLM-identified phase-selective neurons (dots with

corresponding colors) form distinct populations in the principal component space, demonstrating

that our modified GLM algorithm accurately identifies neurons with phase-specific firing

patterns.

To determine if the activity patterns we observed were organized anatomically, we further

assessed the relative co-localization of PPC neurons with similar response properties.

Specifically, we compared the distance within groups of neurons with shared functional

characteristics to the distance across groups (Supplementary Fig. 5). Consistent with previous

results for STM activity in navigation tasks[6], we found that the distances to functionally similar

neurons were not significantly different from the distances to neurons with distinct firing

patterns, refuting the hypothesis that activity patterns were specifically localized.

In summary, our results support the view that activity sequences in the PPC can encode short-

term memory of sensory input, even in a delayed match-to-sample task that does not require

encoding of heading direction or spatial location.

Functional characteristics of directional neurons in error trials

Having established that the PPC neurons exhibit direction- and phase- selectivity without the

involvement of spatial navigation, we next sought to discern the mechanisms by which PPC

activity may be disrupted on error trials. Specifically, we aimed to determine whether

inadequately encoded sensory inputs during the sample phase, poor retention of the encoded

sensory information during the delay phase, or both, could reflect disrupted STM in error trials.

The activity of selective PPC neurons identified in correct trials showed markedly lower phase-

or direction-selectivity on error trials (Fig. 3A). At the level of individual neurons, we often

observed that neurons with high PI exhibited opposite directional selectivity, on error trials (Fig.

3B). It is important to note that not all the directionally selective neurons responded in the

opposite direction, suggesting a mixed population response but one in which there is some

degree of mis-encoding of the STM. There was a significant reduction in the PI during incorrect

trials, particularly among the neurons that had a high PI in correct trials (Fig. 3C). In fact, the

magnitude of reduction in PI was greater for the neurons that had a high PI in correct trials (Fig.

3D and Supplementary Fig. 6). The median ΔPI for the neurons with an initial PI between 0.9

and 1 was approximately -1, suggesting that nearly half of the highly reliable neurons lost

response-selectivity or responded to the opposite direction during error trials. To assess how

activity changes were related to the consistency of responses on individual trials, we also

examined the trial accuracy-dependent difference in RI value for each phase- and direction-

selective neuron as well as non-direction selective neurons (Fig. 4A-I and Supplementary Fig. 6).

This analysis revealed that a greater proportion of sample-phase neurons hold directional

information regardless of trial accuracy, while delay neurons exhibited greater activation in the

opposite direction during error trials. Notably, highly reliable directional delay neurons showed

the greatest reduction in the PI and RI in error trials (Fig. 4 B-I). Together, these results indicate

that directional delay neurons with high reliability are likely to show a response pattern opposite

to their responses on correct trials.

A reversal of the response firing for neurons with high PI would predict that the overall

difference between preferred and non-preferred directions would be diminished. Indeed, we

found a significant reduction in the difference between the ΔF/F of preferred and opposite

directional cues for error trials (Fig. 4J). This reduced difference was more pronounced for the

delay-selective neurons. We observed a reduction in ΔF/F in the preferred direction and an

increase in ΔF/F in the opposite direction, rendering the two responses during the delay period

indistinguishable (Fig. 4K). This suggests that the response selectivity of delay neurons was

more affected than that of sample neurons in error trials.

Activity trajectory distances of the direction-selective PPC neurons during the task

The observation that a fraction of the direction-selective neurons showed activity patterns

corresponding to their response to the opposite directional signal during error trials is potentially

consistent with either mis-encoding or disrupted retention of STM information. To more directly

compare the relative contributions of these two possible mechanisms, we next examined the

temporal changes in the population-wide activity of directional neurons. Specifically, we sought

to determine if the STM associated population dynamics initially resembled the opposite

direction during the sample phase, if it gradually became more like the opposite direction as the

trials proceeded, or both.

To this end, we first captured the dynamics of direction-selective neurons in a reduced dimension

as a trajectory through activity space using PCA. The activity trajectories from the correct

preferred direction responses of direction-selective neurons (specifically, the left trial responses

of left-preferring neurons and the right trial responses of right-preferring neurons) were

designated as preferred directional correct responses or PP. Conversely, the trajectories from the

correct opposite-directional responses (i.e., the right trial responses of left-preferring neurons and

the left trial responses of right-preferring neurons) were dubbed opposite directional responses or

OO. (Fig. 5 and Supplementary Fig. 8). We next calculated the distance between these

trajectories as a metric of the activity difference across conditions (Fig. 5A). The trajectory

distance between PP and OO (d1) remained stably high throughout the sample period and

increased further during the delay period (Fig. 5C, d1). As a control, we also calculated the

distance between randomly selected preferred trials and the remaining preferred trials to assess

the magnitude of variability in activity (Fig. 5C, d0).

To assess the changes in the activity of direction-selective neurons during error trials, we

compared the trajectory distances from error trials to both correct trials, PP and OO. During error

trials where preferred directional inputs resulted in an opposite direction response (PO), such as

left-preferred neuron activity in left stimuli-right response and right-preferred neuron activity in

right stimuli-left response errors, the trajectory distance to the correct OO activity (Fig. 5D, d2)

increased and more or less saturated around the first 0.5 seconds of the sample period. However,

d2 in the sample phase saturated at a lower distance than d1, suggesting a relatively poor activity

distinction by stimulus direction (two-tailed unpaired t-test; n=100, $p < 0.001$ Supplementary

Fig. 8A). The d2 then decreased from the late sample phase through the delay period (gray area).

Conversely, the trajectory distance between PO and PP (Fig. 5D, d3) decreased during the

sample period yet remained significantly greater than d0 (two-tailed unpaired t-test; n=100, $p <$

0.001, Supplementary Fig. 8B). The d3 then increased dramatically during the delay phase.

Together, these results suggest that during the sample phase of PO errors, PPC neurons fired

similarly to PP but with relatively poor distinction by stimulus direction. Over the late sample

phase to the delay phase, the activity increasingly resembled the opposite directional firing.

Similarly, the activity in error trials where opposite stimuli resulted in preferred directional

responses (OP) showed decreasing distance to OO during the sample phase and drifted

significantly during the delay phase (Fig. 5E, d4). The activity distance between OP and PP (d5)

increased during the sample phase and showed a slight decrease at the end of the delay phase.

However, because the error activity trajectories are not necessarily located within the same

activity plane as the PP and OO trajectories, a closer distance to PP does not guarantee a distant

relationship to OO. To evaluate the relative distances in a multi-dimensional space, we devised a

trajectory similarity index (TSI). The TSI measures the differences in distances to PP and OO,

normalized by the sum of those distances, so that a higher absolute TSI indicates greater

similarity. The polarity of the TSI value indicates whether the activity is more similar to PP

(positive) or to OO (negative) (illustration in Supplementary Fig. 8C, see Methods for details).

In line with the analysis of distances to the PP or OO activity (d2-d5), the TSI during the sample

phase showed weak yet significant directional activity according to the given stimuli. However,

as the delay progressed, the activity drifted toward the opposite direction (Fig. 5F).

Population activity of PPC neurons during the task

The activity trajectories in the principal component dimension were computed using the

combined activities of direction-selective neurons across sessions. To understand how these

dynamics might relate to the overall population activity, we next employed measures that assess

activity across all recorded neurons. Specifically, we applied linear discriminant analysis (LDA)

to compare population activity during error trials with that observed in correct trials, where the

stimulus direction was either the same or opposite. This approach provided a measure of the

signal-to-noise ratio (SNR) of the encoded signal across the population, where the signal denotes

the distance between the two responses in the activity space, and the noise reflects the variance

within the responses (see Methods). We measured the SNR changes of error trials within a

moving window of 50 frames in the hyperplane that best discriminates between correct left- and

right-trials (Fig. 6). Consistent with the trajectory distances of direction-selective neurons, the

SNR between correct and same-stimuli error trials (correct left trials (LL) vs. Left stimulus-Right

(LR) response and correct right trials (RR) vs. Right stimulus-Left response (RL)) initially

decreased, reaching saturation around the middle of the sample period (approximately 30

frames). As the frames from the delay period were included (after the 70th frame), the SNR

increased (Fig. 6A, left). On the other hand, the SNR between correct and incorrect trials with

the opposite stimuli (same response error, LL vs. RL and RR vs. LR) exhibited an opposite

pattern: an initial increase and saturation over 30 frames during the sample period, followed by a

subsequent decrease during the delay period (Fig. 6A, right). The patterns of same-stimuli and

same-response errors aligned with the TSI changes in OP and PO errors, respectively (Fig. 5F).

While the grouping of 50 frames across trials ensures the separation between groups, this

analysis obscures the SNR changes during the delay period. To better visualize the SNR changes

during the delay period, we replotted the measured SNR for categorizing every 30 frames of all

trials within the delay period (Fig. 6B). While the same signal errors became more differentiated

during the delay period, a decreasing SNR was observed in the same response errors. Across all

frames, during same-stimuli errors, PPC activities were relatively similar in the sample phase but

diverged more significantly during the delay phase. In contrast, in same-response errors, the

activity differences driven by stimulus direction in the sample phase diminished during the delay

phase (Fig. 6C). These results highlight the role of activity drift toward the opposite direction in

contributing to the erroneous STM.

While our analysis using PCA and LDA revealed that the activity difference in error trials

compared to correct trials, the observed activity difference may not directly reflect a

representation of the directional content of sensory inputs. To address this issue, we employed a

dimensionality reduction approach to identify population activity modes that distinguish features

of interest, particularly the signal corresponding to the direction of motion in the visual stimulus

(coding direction or CD). To directly compare signal direction-dependent activity modes, we

computed projections along a vector in the activity space of all recorded PPC neurons that

maximally differentiate sensory direction in the sample (CD_{sample}) and delay phase (CD_{delay}) (Fig.

6D). To compare the separation of the direction signals, we projected the error trial activities

onto the vector. In comparison to correct trials, the directional sensory signals were weaker and

less distinguishable between directions, particularly during the delay phase of error trials (Fig.

6D-E).

Consequently, we aimed to assess how accurately correct or error trials could be predicted. We

trained a Support Vector Machine (SVM) model using the activities of PPC neurons during

correct trials. Subsequently, we evaluated the accuracy of decoding sensory direction from the

trial-by-trial population activity of PPC neurons using the SVM decoders. We labeled PPC

activity from randomly chosen 70% of correct trials in a session as belonging to either the left or

right direction, and the remaining trials were used to test prediction accuracy (Fig. 6F). For

correct trials, the prediction accuracy of the classifier significantly increased above the shuffled

control level (grey line) at the onset of the sample phase and remained elevated until the end of

the delay period. When assessing error trials using the same hyperplane generated from correct

trials, the prediction accuracy showed a slight temporal delay but eventually reached a level

comparable to that of correct trials. This result suggests that despite a significant reduction in the

distinction between stimulus directions, the population activity of PPC neurons still encodes the

correct direction. However, during the delay period of error trials, the prediction accuracy

dropped rapidly, falling below chance level by the end, indicating a tendency toward choosing

the opposite direction. Taken together, our results demonstrate that population activity in PPC,

resembles the responses to the opposite directional signal in error trials especially during the

delay phase, and thus that STM memory errors are associated with a reversal in the stimulus

content encoding within PPC.

Discussion

Distinguished sensory information maintenance from navigational and movement signals in PPC

activity during STM

In the current study, we observed propagating neural activity that tracked the sensory

information being maintained during STM task performance. While similar observations have

been made previously in STM-based virtual T-maze task[6] a subsequent study have raised

questions about the interpretation of these findings, pointing out that the choice-dependent

activity sequence could be predicted by heading direction and the animal's position in the virtual

corridor and so may reflect a navigational signal[13]. This view is consistent with earlier studies

reporting PPC neurons exhibited activity changes due to locomotor actions, including turning

and forward motion, when these motions were associated with spatial references[11,12,20,21]. Thus,

distinguishing between the activity related to sensory information maintenance and that

influenced by navigational context is crucial for a better understanding of the PPC function in

STM. Sensory cue-dependent go/no-go tasks also have limitations due to possible alternative

interpretations, such as preparatory activity of impulsive licking behavior and their

suppression[6,14,19,22]. In our study, we re-examined PPC activity during an STM task that required

retaining visual information and making choices with an equivalent directional licking behavior

without a spatial component. Despite this, we found remarkably similar activity in PPC neurons.

We observed direction-selective activity sequences that predicted sensory direction in the PPC

not only in the absence of navigation but also without ongoing stimulus during the delay period.

Moreover, the disruptions of activity patterns associated with error trials were consistent with a

drift towards an erroneous representation of stimulus direction, further supporting the view that

the observed activity reflects an STM signal.

Difference in population activity in error trials

A sizeable proportion of the PPC neurons exhibited significant directional preferences, with over

30% showing exceptionally high selectivity (PI> 0.9, Fig. 2A-C). The majority of the direction

selective neurons were tuned to either of the phases simultaneously, rather than exhibiting

sustained activity throughout both periods (Fig. 2A and D). Highly reliable directional firing or

high PI could be ensured by the wiring motif observed in the PPC, where co-selective neurons

have strong and more frequent synaptic connections while inhibiting anti-selective neurons[23].

In the PCA-driven activity subspace, which captured over 70 % of the selective neurons' activity

(Supplementary Fig. 8), trajectories driven by the two-directional visual stimuli were

differentiated immediately after the stimuli onset and continued to diverge during the delay

period (Fig. 5C and 6B). It was rather unexpected that the activity trajectories were further

differentiated during the delay period in the absence of sensory stimuli. This phenomenon was

also consistent with the significantly greater direction-selectivity of the delay neurons compared

to sample neurons (Fig. 2D). These findings support the hypothesis that the primary function of

the PPC is to maintain the differentiated sensory inputs collected during the sample period, rather

than to detect and encode this signal. This latter function may instead depend on early processing

within sensory cortical areas[6,14,24]. The continuous increase in activity differences during the

delay phase may reflect the processing time required to optimize the acquired visual information

and minimize activity variability, occurring in parallel with the time needed for premotor

neurons to prepare and execute invariant motor responses[25,26]. These possibilities could be

distinguished through a systematic comparison of PPC activities with varying stimulus length

and complexity.

Our various, complementary multivariate population analyses consistently demonstrated a

retention of differences in population activity in response to different stimuli in correct trials

(Fig. 5C, and 6F). Although the underlying population activity is spread across multiple neurons,

this encoding pattern is nonetheless reminiscent of the increased (or decreased) and sustained

firing rate in PPC neurons of primates[8,27] or rats performing tasks that require STM[9]. Hence, it is

tempting to propose that the population activity difference in mouse PPC is analogous to the

quality of evidence. In this scenario, differences in PPC activity between cue directions that

develop during the sample period would serve as the basis for stimulus encoding, while the

maintenance of these differences over the delay period would correspond to retention of this

information. Consistent with this idea, we observed weaker directional differentiation and poor

maintenance of differences in PPC activities during error trials. These observations suggest that

STM errors are associated with weaker encoding of sensory information and a progressive

weakening of this information throughout the STM task (Fig. 5-6). However, it is noteworthy

that despite the 'misunderstanding' of some neurons and the resulting reduced activity distinction

during the sample phase of error trials, population-level activity in the PPC still sufficiently

encodes the correct direction (Fig. 6F).

Increasing opposite directional activity of PPC neurons in error trials

While the origin of the reduced distance in activity space on error trials remains unclear, changes

in the arousal state of the animal may account for this poor encoding. The level of arousal varies

continuously and neuronal responses in sensory cortices as well as high-order cortical areas

including PPC are critically modulated by attentional states[6,28–33]. In error trials with insufficient

arousal, the activity of PPC neurons in response to the preferred visual stimuli would decrease,

leading to relatively poor separation in the activity space (Fig. 3D and 4G-H). However, neurons

with negative PI values, which exhibit greater firing to the opposite directional stimuli than the

preferred direction, cannot be explained by this modulation. Instead, we believe that PPC activity

encodes how the stimulus was 'understood' rather than the stimulus identity per se, particularly

for delay neurons and to a lesser degree for directional sample neurons (Fig. 3 and 4). Supporting

this idea, neurons in the PPC of macaques showed firing changes that could predict behavioral

choices even with 0% coherence in a random dot motion task[8].

Recent anatomical analysis has demonstrated that the probability and strength of the L2/3 PPC

neurons are dependent on the similarity of directional selectivity[23]. Moreover, PPC subnetworks

with opposite selectivity appear to suppress one another through mutual inhibitory

connections[23,34,35]. Based on these findings and our results, we propose that direction-selective

delay neurons form stronger subnetworks than sample neurons. In error trials, although the

opposite-directional PPC activity during the sample phase does not initially alter the perceived

direction of the stimuli, it may be amplified through tightly interconnected delay neurons,

ultimately leading to an inversion in the encoded direction and resulting in an error. This idea is

consistent with our findings, which illustrate a significantly greater PI and RI in delay neurons

compared to sample neurons (Fig. 2D and 4), as well as the drifting of population activity toward

the opposite direction as the delay progresses (Fig. 5F and 6).

Limitation of the study

Due to the limited number of task-relevant neurons recorded in a single trial, compounded by the

intrinsic noise of cortical activities, we based much of our analyses on averaged activity across

numerous trials of the STM task. As a result, it was not plausible to reliably track the onset and

amplification of inappropriate directional firing in the current study, both the origin of erroneous

activity within the subnetwork and the nature of its amplification remains unresolved. Further,

even though the PPC activities were acquired while mice were engaged in head fixation to

perform the task, unintended movement or arousal state may occur that could modulate PPC

activity [33,36]. It would be an intriguing avenue for future study to access trial variability with

variable attentional conditions with the aid of high-density electrophysiological recordings[37–41].

Conclusion

In conclusion, our study sheds light on the neural mechanisms underlying sensory encoding and

retention in STM tasks, emphasizing the role of the PPC in these processes. By elucidating the

relationship between PPC activity patterns and task performance, we provide valuable insights

into the cognitive functions of the PPC and with major implications for understanding STM

errors. Since reduced STM performance is one of the key symptoms in schizophrenia[42–44] as well

as an endophenotype of other neuropsychiatric disorders[45] insight into the nature of errors occur

in STM also hold major translational relevance. Thus, our findings open new avenues for

research aimed at developing novel therapeutic approaches for these disorders. Future research

exploring trial variability and attentional conditions using advanced recording techniques will

further enhance our understanding of PPC function in STM as well as how it may be disrupted in

disease.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used ChatGPT-OpenAI in order to correct

grammatical errors. After using this tool, the authors reviewed and edited the content as needed

and take full responsibility for the content of the publication.

Acknowledgements

Korea Brain Research Institute Research Programs, grant 23-BR-01-01, 23-BR-03-01, 24-BR-03-08, and 23-BR-04-04 (JCR). National Research Foundation of Korea, funded by the Ministry of Science and Information and Communication Technology, grant NRF-2017M3A9G8084463 and NRF-2022R1A2C1004216 (JCR).

Author contributions

J.C.R. and J.H.C. planned the study. J.H.C. instrumented and performed the experiments. S.B. and J.P. analyzed the data under the supervision of J.C.R., J.H.C., J.W.C., and L.I.S. The manuscript was prepared by S.B. and J.C.R. with input from all the authors.

Competing interest

The authors declare no competing interests.

Supplementary information

Supplemental Figure S1: Time-correlation of activity dynamics along the coding direction by the

PPC neurons in the STM task.

Supplemental Figure S2: Logic diagram of directional and non-directional phase-selective

neurons identified by the modified General Linear Model (GLM).

Supplemental Figure S3: Reliability index of directional and non-directional neurons.

Supplemental Figure S4: Supplementary Figure 4. Phase selectivity of GLM-defined phase-

selective and non-selective neurons.

Supplemental Figure S5: Anatomical distances and functional similarities.

Supplemental Figure S6: Reliability index of direction- and phase-selective neurons in error

trials.

Supplemental Figure S7: Variance explained by the number of principal components.

Supplemental Figure S8: Discriminability between the trajectory distances and schematic of the

trajectory similarity index (TSI)

Supplemental Figure S9: Framework of the proposed method for automatic event-related ROI

detection.

Main Figure

Figure 1 Ca²⁺ responses of PPC neurons during short-term memory task. A. The schematic

illustrates the behavioral task designed to measure short-term memory in head-restrained mice.

During the sample phase, black-and-white phase-reversing gratings moved either leftward or

rightward (or downward or rightward) for 4 seconds on a screen in front of the mouse. After a 2-

second delay, two laterally positioned lick ports were moved to a location within reach of the

mouse's tongue. Correct licking of the corresponding lick port was rewarded with water, while

incorrect licking resulted in a time-out as a form of punishment. **B.** Four possible behavioral

outcomes from the task. **C.** Performance in right and left trials from thoroughly trained mice. No

significant difference was observed (two-tailed paired t-test; n=11, $p < 0.05$). Error bars

represent the standard error of the mean (SEM) across sessions. n.s: not significant. **D.** Examples

of neuronal activity recorded during behavioral tasks. Each row illustrates the color-coded

normalized **Δ**F/F. **E.** The color-coded autocorrelation diagram of the mean response vectors, *w*,

derived from coding direction (CD) of correct trials. *w* was calculated over moving 0.5s time

windows. Data from 11 sessions, 6 mice, and 4756 total neurons S: sample phase, D: delay

phase, and R: response phase. **F.** The proportion of neurons displaying task-dependent activity

defined by a General Linear Model (GLM). (Left) The fraction of neurons with single-phase

selectivity, and neurons with activity in two or more phases out of all recorded neurons. (Middle)

Percentages of neurons with and without directional selectivity out of single-phase selective

neurons. (Right) Percentages of neurons with corresponding phase selectivity out of single-phase

selective neurons.

Figure 2 Identification and characterization of the direction- and phase-selective neurons.

A. Color-coded trial average ΔF/F of right (upper) or left (lower) directional phase selective

neurons. ΔF/F was normalized to the maximum ΔF/F and aligned based on the time-to-peak of

each neuron. **B.** Example traces (left) and histogram of trial-averaged ΔF/F from highly

directional phase-selective PPC neurons. The thickness of the traces represents the standard

error. Blue and red in the histogram illustrate left and right trials, respectively. **C.** (Left)

Histogram of preference indices (PI) of GLM-defined directional neurons (light blue) and those

calculated with shuffled trials (gray). (Right) Same as left but shown as cumulative distribution

function (CDF). **D.** PI histogram of directional neurons during the sample phase (upper) and

delay phase (lower). **E.** Principal components analysis (PCA) capturing the temporal activity

pattern of each neuron. GLM-defined sample (black), delay (blue), or reward (red) phase

selectivity is indicated by colors. Variance explained by the principal components.

Figure 3. Responses of the direction- and phase-selective neurons in error trials. A. Same as

Fig. 2A but in error trials. **B.** The same example neurons shown at the first and third row in Fig.

2B (left) and their mean traces in error trials. **C.** (Left) Histogram of preference indices (PI) of

GLM-defined directional neurons in correct (light blue), error (red open), and shuffled (gray)

trials. (Right) Cumulative distribution function (CDF) of the left histogram. **D.** A box plot

illustrating the difference of PIs (ΔPI) between error and correct trials for each neuron, as a

function of PIs in correct trials.

Figure 4. Comparison of responses between delay phase-selective neurons and sample phase-selective neurons during error trials. A. Preference indices (PI) histogram of GLM-defined directional single-phase selective neurons in correct (light blue), error (red open), and shuffled (gray) trials. **B.** PI histogram of directional sample-phase selective neurons in correct (black), error (red open), and shuffled (gray) trials. **C.** PI histogram of directional delay-phase

selective neurons in correct (blue), error (red open), and shuffled (gray) trials. **D.** (Left)

Reliability indices (RI) histogram of GLM-defined directional single-phase selective neurons in

correct (light blue), error (red open), and shuffled (gray) trials. (Right) Cumulative distribution

function (CDF) of the left histogram. **E.** (Left) RI histogram of directional sample-phase

selective neurons in correct (black), error (red open), and shuffled (gray) trials. (Right)

Cumulative distribution function (CDF) of the left histogram. **F.** (Left) RI histogram of

directional delay-phase selective neurons in correct (blue), error (red open), and shuffled (gray)

trials. (Right) Cumulative distribution function (CDF) of the left histogram. **G.** (Left) Scatter plot

comparing reliability index in PPC correct and incorrect. A reliability index value close to 1 for

'Correct' indicates a significant response in the preferred direction, while a reliability index value

close to -1 for 'Incorrect' suggests a response closer to the opposite direction during error trials.

(Right) Scatter plot comparing the reliability index (Correct) with the difference between

reliability index (Incorrect) and reliability index (Correct) in PPC. Δ reliability index values

closer to -1 indicate that the response during incorrect trials was closer to the opposite direction

compared to correct trials. **H.** (Left) Scatter plot comparing the reliability index of GLM-defined

direction-selective sample neurons in PPC during correct and incorrect trials. (Right) Scatter plot

comparing the reliability index (Correct) of GLM-defined direction-selective sample neurons

with the difference between the reliability indices for incorrect and correct trials. **I.** (Left) Scatter

plot comparing the reliability index of GLM-defined direction-selective delay neurons in PPC

during correct and incorrect trials. (Right) Scatter plot comparing the reliability index (Correct)

of GLM-defined direction-selective delay neurons with the difference between the reliability

indices for incorrect and correct trials. **J.** (Upper) Mean normalized ΔF/F of direction-selective

sample neurons in response to the preferred (red) and the opposite (blue) directional signal in

correct (left) and incorrect (right) trials. (Lower) Same as upper plots but for direction-selective

delay neurons. **K.** (Upper) Δ mean normalized $\Delta F/F$ of direction-selective sample neurons in

correct trials (black) or incorrect trials (gray), calculated by subtracting the value of the opposite

directional signal from the preferred directional signal. (Lower) (Upper) Δ mean normalized

$\Delta F/F$ of direction-selective delay neurons in correct trials (black) or incorrect trials (gray),

calculated by subtracting the value of the opposite directional signal from the preferred

directional signal.

Figure 5. Differences in population activity of direction selective neurons in correct and

error trials. A. Schematic of assessing population activity difference using the dimensionality

reduction approach. **B.** Activity trajectories that capture the population activity of directional

neurons in the preferred (dark colors) or opposite (light colors) directional stimulation in correct

(blue) and incorrect (red) trials. Although the first three principal components are illustrated for

visualization purposes, trajectories were projected into a five-dimensional subspace. Eigenvector

was generated using 50% of all sorted neurons, and the remaining 50% were projected onto this

eigenvector. The faint lines in each graph represent values from 100 random iterations **C.**

(Upper) A schematic of correct preferred and correct anti-preferred, or opposite trial (e.g. for

left-preferring neurons, correct left trials vs. correct right trials). (Lower) Temporal changes of

the Euclidean distance between the activity trajectories of correct preferred direction responses

and correct opposite direction responses (PP-OO) (d1). d0 illustrates trial-by-trial activity

variation, showing the distance between randomly selected preferred direction responses and the

rest of the preferred direction responses in correct trials. **D.** d2: The distance between the activity

trajectories of correct opposite direction responses and incorrect preferred direction responses.

(PO-OO). d3: The inter-trajectory distance of correct preferred direction responses and incorrect

preferred direction responses (PP-PO). **E.** d4: Inter-trajectory distance between the correct

opposite direction response and incorrect opposite responses (OP-OO). d5: Inter-trajectory

distance between the correct preferred direction response and incorrect opposite direction

responses (PP-OP). **F.** Temporal changes in the trajectory similarity index for the incorrect

preferred responses (orange) or the incorrect opposite responses (blue) (see also Supplementary

Fig. 8).

Figure 6. Population dynamics of PPC neurons in correct and incorrect trials. A. Time-

dependent changes in the normalized signal-to-noise ratio (SNR) in distinction between

responses in activity space using the linear discrimination task (LDA). SNR was calculated over

a sliding window of 50 frames, advancing by one frame at a time. The vertical dashed line

represents the initiation point where the window commences its entry into the delay phase. (Left)

Mean normalized SNR between clusters of frames representing the ensemble of neural activity in

correct trials and same stimulation error trials (i.e. LL vs LR and RR vs RL) as a function of

time. The black horizontal line (paired t-test; frame 8 ~ 107, $p < 0.05$) and the bold black line

(paired t-test; frame 11 ~ 65, $p < 0.01$) indicate time points significantly different from the delay

phase. (Right) Same as left but between correct and same choice error trials (i.e. LL vs RL and

RR vs LR). The black line (paired t-test; frame 3 ~ 112, $p < 0.05$) and the bold black line (paired

t-test; frame 6 ~ 105, $p < 0.01$) indicate time points significantly different from the delay phase.

Line thickness indicates mean ± SEM across sessions. **B.** Mean normalized SNR of LDA during

the delay phase. SNR was calculated over a sliding window of 30 frames during the delay phase.

(Gray) LDA between correct trials and same stimulation error trials. (Blue) LDA between correct

and same choice error trials as a function of time. The thickness of the lines represents standard

error. **C.** Comparison of normalized SNR of LDA between correct and same stimulus error (left)

or same choice error (right) calculated from the same number of frames in the sample phase

(frames 66 ~ 115) and delay phase (frames 121 ~ 170). Statistical analyses were conducted using

paired t-test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$). Error bars indicate mean ± standard error

across sessions. **D.** Projections of activity along CD_{sample} and CD_{delay} in each direction (red: right,

blue: left) and response accuracy (dark: correct, light: incorrect). Line thickness indicates mean ±

standard error across sessions. **E.** Residual separation defined as (Right incorrect - Left incorrect)

/ (Right correct - Left correct). (paired t-test, $p < 0.05$). **F.** Accuracy of decoded neural activity

prediction by support vector machine (SVM) plotted as a function of time. Line thickness

indicates mean ± standard error across sessions.

Methods

Animals

All animal experiments were conducted with the approval of the Animal Experiment Ethics

Committee (approval no. IACUC-17-00031) at the Korea Brain Research Institute (KBRI). All

experiments were performed using male C57BL/6J-Tg(Thy1-GCaMP6f)GP5.5Dkim/J transgenic

Mice (n = 6).

Cranial window and head plate preparation

Mice aged between 8 to 12 weeks underwent cranial window and head plate implantation

surgery. To ease pain and reduce inflammation, they received subcutaneous injections of

ketoprofen (5 mg/kg) prior to surgery. For deep anesthesia, a mixture of 1–2% isoflurane or a

ketamine/xylazine solution (100 mg/kg and 10 mg/kg) was used. After confirming anesthesia by

the absence of pedal withdrawal reflex, we covered both eyes with petroleum jelly and removed

hair from the incision site. The scalp was cleaned and sterilized with ethanol and betadine and

then removed. Using a scalpel, we stripped away the periosteum and roughened the skull's

surface with a drill burr (Miniature Carbide Burr Drill Bits Set, Stoelting) to ensure the dental

cement would adhere properly. The skull was cleaned with Ringer solution, and a craniotomy

approximately 3 mm in diameter was performed over the posterior parietal cortex of the right

hemisphere. The craniotomy was then sealed with a double-layered glass coverslip (round 3/5

mm, Harvard Apparatus combined with NOA71, Norland) and secured with surgical adhesive

(Vetbond, 3M). The dura mater remained unharmed. A tailor-made commercially pure titanium

head plate (rectangular shape, 19 mm x 12 mm x 1 mm) with a 4.5 mm central opening was

affixed on top of the window using dental cement (Super-Bond C&B kits, Sun Medical,

darkened with black pigment). After the surgery, mice were placed in a warmed chamber for a

minimum of four hours for recovery and post-operative monitoring. Consequently, the animals

were housed in individual cages.

Behavior task

After full recovery from surgery, the mice were provided with limited water (1 ml/day) while

their body weights were monitored. If a mouse's body weight dropped below 75% of its initial

weight, water restriction and training were stopped. The mice were trained to perform a delayed

match-to-sample task while their heads were fixed in place using a holder (modified MAG-1,

Narishige Scientific Instrument, Tokyo, Japan). The visual signal to discriminate was high-

contrast sine-squared wave gratings, moving either left/right or down/right, with specific spatial

(3 cm/cycle) and temporal (2 Hz) frequencies. The signal was provided by a monitor placed

directly in front of the mice at a distance of 25cm. Custom written software running in Raspberry

Pi 3 (RPi3) was used for both stimuli generation and behavioral training. Left/right or down/right

were trained in a separate group of mice and all the behavioral and imaging data acquired from

left and down trials were pooled and analyzed as the same group.

The position of the drinking spouts was adjusted to fit each individual mouse. The lick detection

circuit and reward solenoid were controlled by Arduino Mega2560 (rev 3, Arduino) connected to

RPi3. During training, mice earned the majority of their daily water intake (1 mL) through

training. Each visual stimulus lasted 4 seconds, followed by a 2-second pause. After this pause,

the spouts moved quickly to a position accessible by the tongue for 1 second. Successful licks

(licks on the left-hand side spout in response to leftward or downward moving signals, or licks

on the right-hand side spout in response to rightward moving signals) during this window were

rewarded with approximately 5 µl of tap water. Incorrect licks resulted in no reward and were

punished with a 2-second time-out, after which the trial continued. After a reward period or upon

an incorrect lick, the spout withdrew quickly, staying out of reach until the commencement of

the next trial, which was set to begin 4 or more seconds later. Training progressed through the

following stages. First, the mice learned to lick a non-moving lick port located at the center with

either of visual cues. Then they learned to match the direction and rewarding spout out of the two

non-moving spouts. We then gradually increased the delay by introducing moving lick ports.

Once mice consistently achieved high performance (>70% correct rate) with 2s delay, behavioral

performance and PPC activities were recorded for analyses.

In vivo two-photon calcium imaging

We conducted two-photon imaging utilizing HyperScope (Scientifica) and ScanImage (MBF

Bioscience) with a Ti:Sapphire laser (Chameleon Ultra II, Coherent). The imaging setup targeted

the GCaMP6f calcium indicator excited at 910 nm through an 16x water immersion lens

(N16XLWD-PF, Nikon). The objective lens has an attached light barrier (custom made) isolating

the imaging system from visual stimuli. We captured 512×512 pixel images(700umX700um

ROI) at 30Hz in the PPC layer 2/3. The average imaging laser power was less than 100mW at

the objective lens. The imaging and behavior system was synchronized by frame trigger signal

output counted by Mega 2560.

Image preprocessing and cell selection

The processing of calcium imaging data utilized the CaImAn software framework. This included

adjusting for motion in segments of 128 by 128 pixels, ensuring each segment overlapped by 48

pixels to maintain continuity. For identifying neuronal sources, we applied the 'greedy_roi'

technique, sizing the expected neurons at 4 by 4 pixels for detailed imaging and 3 by 3 pixels for

imaging the entire brain. Only components with a signal-to-noise ratio of at least 2.0 and 90%

pixel correlation were considered reliable. We normalized the fluorescence signal decay rates,

3.0 seconds for indicators localized in the nucleus and 1.0 second for those in the cytosol, using

z-scoring for subsequent analyses. To ensure coherence between behavioral actions and neural

activity, we aligned and adjusted all time series data to a uniform sampling rate of 30 Hz,

synchronized with the camera and microscope's frame capture times

Automated event-related neuron detection in calcium imaging data

We used generalized linear models (GLM) to regress recorded calcium signals against a time

series of task events. We propose a novel method for automatically searching event-related

neurons in calcium imaging data (Supplementary Fig. 9). Our method searches the ROI

pixelwise using event timing information. By utilizing a statistical method to determine the

relationship between the calcium trace of each pixel and the event timing, the searched ROI

pixels are clustered. This process enables the automatic detection of event-related ROIs without

requiring any additional detection procedures, while also enabling the detection of components

with low SNR or fine size as ROI.

The proposed method determines the event relevance of calcium traces to detect event-related

neurons in calcium imaging data. Rather than relying on the morphological shape of the neuron

for ROI detection, the proposed method determines the event relevance of each pixel's calcium

trace independently. If ROI detection is performed by reflecting the neuron's morphological

shape, the detection performance may decrease with low SNR or fine-size components.

The proposed method utilizes a generalized linear model (GLM) to detect ROIs by assessing the

relationship between the calcium trace of each pixel and the calcium signal model that is strongly

associated with the event. To create a calcium signal model that is strongly associated with

events, a temporal component model of calcium dynamics and event timing is necessary.

Event timing refers to the timing of a task or stimulus. In an event-related neuron, there is a

calcium activity after at the event timing. Non-event timing refers to other times where there are

just activations that are not related to events, including spontaneous activation occurs. Neurons

strongly associated with events show high activation at event timing while low activation at non-

event timing. Therefore, we define a spike train $\tilde{s}(t)$ of the calcium signal most strongly

associated with the event:

$$\tilde{s}(t) = \sum_{i=1}^{k} \beta_i \delta(t - t_i) \tag{1}$$

where $\tilde{s}(t)$ is composed of the product of the β (amplitude of the spike) and the impulse train.

An event-related neuron is expected to activate at least once after an event. Denoting the i^{th} event

timing by t_i, $\tilde{s}(t)$ becomes a spike train of event-related neurons with an amplitude of β_i at event

timings.

The temporal component of calcium dynamics is determined by the activity of a calcium

indicator. When the calcium indicator is combined with calcium, luminous intensity changes.

The fluorescence trace of the calcium indicator was modeled in the form of a double exponential

and is expressed as:

$$c(t) = e^{\frac{-t}{\tau_d}} - e^{\frac{-t}{\tau_r}} \tag{2}$$

800   where τ_r and τ_d are the time constants for the increase and decrease of fluorescence,

801   respectively[1].

802   We express the modeled event-related calcium trace $\tilde{y}(t)$ of the neuron by the fluorescence trace

803   model of the calcium indicator $c(t)$ with the convolution of the spike train $\tilde{s}(t)$.

$$\tilde{y}(t) = \int_{-\infty}^{\infty} \tilde{s}(t - \tau)c(\tau)\, d\tau \tag{3}$$

We defined the event-related calcium trace model $\tilde{y}(t)$. The ROI was detected by determining

the relationship between this model and the pixel's calcium trace $y(t)$ using the least square

method and 1-sampled t-test. Through the least square method, it is possible to find the estimator

of the regression coefficient β (the amplitude of the spike in equation (1)) that minimizes the

residual of the pixel's calcium trace $y(t)$ and the event-related calcium trace model $\tilde{y}(t)$ of the

neuron. The β estimator matrix $\hat{\beta}$ up to the kth event was expressed as:

$$\hat{\beta} = [\beta_1\ \beta_2\ \beta_3\ \cdots\ \beta_k] = \|y(t) - \tilde{y}(t)\|^2 \tag{4}$$

Notice that $\tilde{y}(t)$ has a weight $c(t)$ for event timing. By $c(t)$, as the event timing and activation

timing are closer, the value of the regression coefficient β increases. It was determined whether a

neuron is event-related through the distribution of β. As the mean of $\hat{\beta}$ is close to 0 or the

variance of $\hat{\beta}$ is larger, the activation timing of neurons has little relation to the event timing.

Therefore, we tested the null hypothesis that β is extracted from a normal distribution with a zero

mean and unknown variance, using a 1-sampled *t*-test. At this time, the test statistic is as shown:

$$test\ statistic = \frac{\bar{\beta}}{s/\sqrt{k}} \tag{5}$$

where $\underline{\beta}$ is the mean of the $\hat{\beta}$, s is the standard deviation of the $\hat{\beta}$, and k is the size of $\hat{\beta}$, also

known as the trial number of events. The proposed method uses a *p*-value of 0.05 to determine a

pixel with a statistically significant difference from a pixel of an event-related neuron.

When the event association determination is completed for all pixels of the calcium image, the

proposed method finally defines the event-related ROI by clustering the pixels associated with

the event. The proposed method clusters pixels to ROIs automatically, where the clustering is

performed based on the Moore-Neighbor tracking algorithm and determines connectivity

between pixels [2]. Finally, the proposed method provides ROI clusters corresponding to event-

related neurons (Supplementary Fig. 9).

Selectivity

Through our uniquely developed Generalized Linear Model (GLM), we pinpointed neurons that

were activated in specific phases across different behavioral patterns. To confirm these

observations, we employed a preference index, calculated by conducting a receiver operating

characteristic (ROC) analysis. This analysis measures the theoretical observer's capability to

distinguish between types of trials using the neural responses from individual trials. The

preference index itself is determined from the area under the ROC curve (auROC), using the

formula $2 \times (auROC - 0.5)$, producing a range from -1 to 1. We applied this index to the

segments representing each behavioral phase, using the average neural activity from trials to

derive specific values. Notably, the preference index (PI) compares trial-averaged values

between right and left conditions. Additionally, the reliability index (RI) is calculated by

comparing all individual trials between right and left, without averaging. Furthermore, the phase

selective index (PSI) is derived by comparing the trial-averaged values from the phase of interest

with those from non-related phases. This method allowed us to quantitatively assess the degree

to which individual neurons preferentially respond to different trial types, providing a clearer

understanding of the role each sorted neuron plays in the observed behavioral patterns.

PCA

We used Principal component analysis (PCA) on the neurons sorted through the Generalized

Linear Model (GLM) to infer a shared space, an intrinsic manifold shared by neuronal

population activity. We calculated the variance explained by principal components of correct

trials. We observed five principal components explained similar level of variance of our data

both in right-preferring neurons, left preferring neurons, and directional neurons.(right-preferring

neurons: 72.4%, left-preferring neurons: 69.0%, directional neuron: 70.4%). A specific

procedure for conducting PCA on PPC neuron data is as follows:

We organized the optical data from an experiment into an $X = M \times N$ matrix, where M is the

number of frames per trial, and N is the number of sorted neurons in a session $\left(N = N_{sample} + \right.$

$\left. N_{delay} + N_{reward}\right)$. Directional neurons, comprising both right-preferring and left-preferring

neurons, were grouped together. Therefore, we proceeded with Principal Component Analysis

(PCA) after aligning each neuron according to its preferred direction. We applied z-score

normalization to each column of the matrix to ensure comparability across neurons. Z-score

normalization adjusts the data to have a mean of 0 and a standard deviation of 1, enhancing the

robustness of PCA against variations in scale among different neurons.

$$Z_{correct} = \frac{(X - \mu)}{\sigma} \tag{6}$$

865   Where μ is the mean and σ is the standard deviation of neuron activity.

866   We calculated the covariance matrix, C, from the normalization data Z, to understand how the

867   variables relate to one another.

$$C_{correct} = \frac{1}{n-1} Z_{correct}^T Z_{correct} \tag{7}$$

Where n is the number of neurons.

We computed the eigenvalues and eigenvectors of the covariance matrix C. The eigenvectors

represent the directions of maximum variance in the data, and the eigenvalues represent the

magnitude of these variances. Neuronal activity data were projected onto the principal

components to transform the data into a new space defined by these components. To compute

relative changes in incorrect trials compared to correct trials, we projected the incorrect trials

onto the eigenvector induced by correct trials and quantified the variations through the distances

between trajectories.

$$Z_{pca} = Z_{incorrect} \times V_{correct} \tag{8}$$

Where $V_{correct}$ represents eigenvectors and Z_{pca} represents the data transformed into the

principal component space.

Coding direction

This procedure closely follows the methodology outlined in [3,4]. The population dynamics of n

neurons captured concurrently throughout a session were examined (Fig. 6). The population

activity of n neurons created a trajectory in the n - dimensional activity space during each

session, where each dimension corresponds to a single neuron activity. In the activity space, we

found directions (modes) that maximally separated the neuronal pathways for various trial

conditions and specific phase. In trials with lick-correct left and lick-correct right instructions,

the trial-averaged activity differences of n neurons were averaged within a 1.7s (121 ~ 170

frame) window after the visual cue offset. This resulted in the delay mode (CD_{delay}), which was

calculated as follows:

$$\Delta\underline{r} = \underline{r}_{correct\ right} - \underline{r}_{correct\ left} \qquad (9)$$

893   The resulting vector was normalized by its l^2 norm, resulting in a vector of weights (one weight

894   per neuron i) as follows:

$$m = \frac{\Delta\bar{r}}{\sqrt{\sum_i^N |\Delta\bar{r_i}|^2}} \qquad (10)$$

896   Projections of the neural activity along the mode over time were calculated as follows:

$$P^k = (X^k)^T m \tag{11}$$

Where X^k is the $N \times T$ matrix of averaged activity of neurons over time for session k.

To maintain consistency in the analysis despite the varying number of neurons recorded in each

trial, we normalized each mode by its l^2 norm, as indicated in equation (10). This normalization

ensured that the projected neural activity, P^k remained unaffected by the quantity of neurons

simultaneously recorded. In instances where a specific neuron was not showed a specific activity

during a particular trial, we assigned a weight of zero to that neuron in both equations (10) and

(11) for the purpose of calculating the projected activity for that trial. This approach guaranteed

that the analysis could accurately reflect the contribution of active neurons in each trial, without

skewing results due to differences in neuron sampling across trials.

The sample mode (CD_{sample}) and the reward mode (CD_{reward}) were defined to capture distinct

aspects of neural activity corresponding to different phases of the trials, specifically focusing on

the behavioral outcomes of licking correctly to the right or left.

CD_{sample} : This mode was determined by constructing an $n \times 1$ vector representing the

differences in the average activity of n neurons during trials that resulted in correct licks to the

right or left. These differences were averaged over a 1.7-second window (from frame 56 to frame

115) coinciding with the beginning of the stimulus presentation. By focusing on this early

window, CD_{sample} captures the neural activity patterns triggered by the initial stimulus,

reflecting how neurons differentiate between trial types in response to the stimulus.

CD_{reward} : Similarly, this mode was defined by an $n \times 1$ vector, but it focused on the average

activity differences of n neurons during trials with correct licks to the right or left, within a 1-

second window (from frame 171 to frame 200) at the conclusion of the delay phase. The

CD_{reward} specifically targets the neural response patterns associated with the anticipation or

receipt of a reward, highlighting how neuronal activity aligns with behavioral outcomes

following the stimulus-response interval.

In addition to the selective modes that focused on differentiating neural activity based on specific

trial outcomes, we identified a nonselective ramping mode (D_{reward}). This mode was designed

to capture general changes in neural activity towards the end of the delay phase, irrespective of

the trial outcome (whether the lick was correct to the right or left). To compute this ramping

mode, we adopted a methodology similar to that used for the reward mode. However, instead of

distinguishing between trial outcomes, we aggregated neural responses from both correct-right

and correct-left licks. We then calculated the difference in trial-averaged activity during the last

second of the delay phase (from frame 141 to frame 170) compared to the first second of the

reward phase (from frame 171 to frame 200).

$$\Delta \underline{r} = \underline{r}_{correct\ reward} - \underline{r}_{correct\ last\ delay} \qquad (12)$$

To facilitate a direct comparison between the sample and delay phases while considering their

distinct neural dynamics, we employed the Gram-Schmidt process for orthogonalization of these

phases in a specific sequence. This process was applied to the modes in two different orders:

first, starting with the CD_{reward}, then the D_{reward}, followed by the CD_{sample}; and second,

beginning with the CD_{reward}, then the D_{reward}, and finally, the CD_{delay}. Orthogonalization

using the Gram-Schmidt process ensures that each mode represents a unique aspect of neural

activity, independent of the others, thus allowing for a clearer analysis of the distinct

contributions of sample and delay phases to overall neural dynamics.

Furthermore, to assess the neural activity's alignment with behavioral outcomes - specifically,

the differences between correct and incorrect responses - we projected the trial-averaged activity

from incorrect trials onto the modes defined by correct trials across sessions. This projection

allowed us to compare the neural activity associated with correct and incorrect trials within the

same mode, providing insights into how specific patterns of neural dynamics correlate with

behavioral performance. This approach helps in understanding the neural underpinnings of

decision-making and error processing by quantifying the extent to which neural activity during

incorrect trials deviates from the established patterns of correct trials.

After projecting the neural activity data along the direction defined by CD (which represents the

difference in neural activity between conditions), we normalized the resulting projections. This

normalization involved adjusting the projections based on the average value obtained from

correct rightward trials during the specific phase associated with each mode, for every session.

This step ensures that the comparison across different trials and sessions is on a consistent scale,

facilitating a more accurate analysis of the neural activity patterns.

To further investigate the differences in neural activity patterns between incorrect and correct

trials, we introduced the concept of "residual separation." This metric quantifies the deviation in

the neural trajectories of incorrect trials from those of correct trials. The residual separation

effectively captures the extent to which the neural representation of incorrect trials diverges from

the expected pattern observed in correct trials, providing insights into the neural mechanisms

underlying error processing and decision-making accuracy. This analysis allows us to better

understand how the brain's neural activity aligns with behavioral outcomes and how it adapts or

fails in cases of incorrect responses.

$$Residual\ separation = \frac{mean(Traj_{incorrect\ right}) - mean(Traj_{incorrect\ left})}{mean(Traj_{correct\ right}) - mean(Traj_{correct\ left})} \quad (13)$$

Mean ($Traj$) represents averaged trajectory at the time corresponding to each mode.

Linear discriminant analysis

This procedure closely follows the methodology outlined in[5]

In examining a neural network that processes stimulus-tuned input under noisy conditions, with

the objective of accurately reporting the stimulus identity, we delve into the nature of the input

signals. Assuming that for a particular stimulus, the input time series can be modeled by a

multivariate normal distribution characterized by temporally uncorrelated, stimulus-independent

noise, our analysis reveals a statistically optimal strategy for distinguishing between two

different stimuli. This strategy encompasses executing a linear projection alongside temporal

filtering of the input time series.

By applying this approach, we calculate the ideal projection weights and filtering mechanism,

alongside the signal-to-noise ratio (SNR) achievable through any given projection and filtering

combination. It's important to highlight that the selection of the input dimensionality and

statistical properties was guided primarily by the aims of enhancing visualization and deepening

conceptual understanding. However, our analytical framework is versatile, accommodating

inputs from an arbitrary number of neurons, each with unique stimulus tuning and noise

covariance patterns. This flexibility underscores the broad applicability of our findings,

extending beyond specific configurations to encompass a wide range of neural input scenarios.

For the task of distinguishing between two types of stimulations S_i , where $i = 1$ corresponds to

a correct right decision and $i = 2$ to a correct left decision, we consider the dynamics of

feedforward input at each time step t. The input, $u(s_i, t)$, follows a multivariate normal

distribution, characterized by a stimulus-dependent mean and a stimulus-independent covariance.

Specifically, the mean for the stimulation S_1 is denoted as $g(s_1) = [1, 2]$, and for S_2 it is

$g(s_2) = [2, 1]$. The covariance matrix, independent of the stimulus, is given by $\Sigma_\eta = [2, 1; 1, 2]$.

To analyze the inputs, we employ a projection onto a linear discriminant defined by $W_{LD} =$

$\Sigma_\eta^{-1}(g(s_2) - g(s_1))$. This approach effectively differentiates the inputs based on their stimulus-

dependent means while considering the uniform noise structure. The projected time series,

$d_{W_{LD}}(s, t) = w_{LD}^T u(s, t)$ then undergoes cumulative summation over time to yield $D_{W_{LD}}(s, t) =$

$\sum_{t'=1}^{t} d_{W_{LD}}(s, t)$.

This methodology leverages the linear discriminant W_{LD} to filter and aggregate the signal in a

way that enhances the separation between the two stimuli. By summing the projected signals

cumulatively, the process amplifies the distinction between the neural responses to S_1 and S_2,

facilitating the identification of the stimulus based on the accumulated evidence over time.

To elucidate the processing of instantaneous input $u(s, t)$ onto a vector w, we derive expressions

for the signal difference $(\Delta\mu_{input}(w))$, noise $(\sigma_{input}(w))$, and signal-to-noise ratio (SNR)

associated with the projection $d_w(s, t) = w^T u(s, t)$. The analytical expressions are as follows:

1000

$$\Delta\boldsymbol{\mu_{input}}(\boldsymbol{w}) \equiv \langle \boldsymbol{d_w(s_2, t) - d_w(s_1, t)} \rangle = \boldsymbol{w^T}\big(\boldsymbol{g(s_2) - g(s_1)}\big) \tag{14}$$

1001

1002   Signal difference ($\Delta\mu_{input}(w)$) is the difference in mean projections between the two stimuli. It

1003   quantifies how distinct the mean responses to the two stimuli are when projected onto w.

1004

$$\sigma_{input}(w) \equiv \sqrt{0.5 \sum_{i=1,2} \langle (d_w(s_i,t) - \langle d_w(s_i,t) \rangle)^2 \rangle} = \sqrt{w^T \sum_{\eta} w} \tag{15}$$

1005

1006   Noise ($\sigma_{input}(w)$) is the standard deviation of the projection. It measures the variability of the

1007   projection, irrespective of the stimulus.

1008

$$SNR_{input}(w) = \Delta\mu_{input}(w)/\sigma_{input}(w) \tag{16}$$

1009

1010   Signal to noise ratio (SNR) assesses the clarity of the signal relative to the noise for the chosen

1011   projection vector w.

1012   Following temporal integration, to calculate the cumulative effect over time, we consider

1013 $D_w(s,t) = \sum_{t'=1}^{t} d_w(s,t')$ and derive the temporally integrated quantities:

1014

$$\Delta\mu_{input}(w,t) \equiv \langle D_w(s_2,t) - D_w(s_1,t) \rangle = \Delta\mu_{input}(w)t \tag{17}$$

1015

1016   Integrated signal difference ($\Delta\mu_{input}(w,t)$) grows linearly with time, reflecting how the

1017   separation between stimuli accumulates over time.

1018

$$\sigma_{input}(\text{w}, t) \equiv \sqrt{0.5 \sum_{i=1,2} \langle \left(D_w(s_i, t) - \langle D_w(s_i, t) \rangle \right)^2 \rangle} = \sigma_{input}(\text{w})\sqrt{t} \qquad (18)$$

1019   The standard deviation of the integrated signal scales with the square root of time. Integrated

1020   noise ($\sigma_{input}(w, t)$) indicates how noise accumulates over time.

1021

$$\boldsymbol{SNR_{input}(w, t) \equiv \Delta\mu_{input}(w, t)/\sigma_{input}(w, t) = SNR_{input}(w, t)\sqrt{t}} \qquad (19)$$

1022

Integrated SNR ($SNR_{input}(w, t)$) demonstrates how the clarity of the signal improves with

temporal integration, as the signal difference accumulates faster than noise over time.

The trials were organized into four categories based on the mouse's response to directional cues,

enabling a precise analysis of behavioral patterns:

(1) Trials where the mouse licked right in response to a right-directed cue. (2) Trials where the

mouse licked left in response to a left-directed cue. (3) Trials where the mouse licked left in

response to a right-directed cue. (4) Trials where the mouse licked right in response to a left-

directed cue.

The linear discriminant (w_{LD}) was determined based on the differentiation between the two types

of stimulus inputs in correctly responded trials.

To assess the SNR within the time-series data meticulously, we employed a specific windowing

technique. A window spanning 50 frames was defined, and this window was progressively

shifted by one frame forward to scrutinize the SNR variations across the different groups over

time. The SNR within each window was computed as the ratio of the mean signal activity to the

mean noise activity, enabling a quantitative measurement of how clearly the signal of each group

stood out against the background noise.

To standardize comparisons across different sessions and ensure a uniform basis for analysis, the

SNR values observed were normalized against the maximum SNR value observed in the

respective session. This step was critical for drawing direct comparisons across sessions,

effectively illuminating the signal quality differences between them. The utilization of Matlab

for all data analyses provided a robust framework for dissecting and understanding the dynamic

processes underlying the behavioral responses.

1045

1046

1047   **Data availability**

1048

1049   The data and custom MATLAB analysis code supporting the findings of this study are available

1050   on Figshare at 10.6084/m9.figshare.26778301

1051

1052

1053

1054

1055

1056

1057

1058

1059

Reference

1. Lee, J., and Park, S. (2005). Working memory impairments in schizophrenia: A meta-

analysis. J Abnorm Psychol *114*, 599–611.

2. Reichenberg, A., Caspi, A., Harrington, H., Houts, R., Keefe, R.S.E., Murray, R.M.,

Poulton, R., and Moffitt, T.E. (2010). Static and dynamic cognitive deficits in childhood

preceding adult schizophrenia: A 30-year study. American Journal of Psychiatry *167*, 160–

169.

3. Forbes, N.F., Carrick, L.A., McIntosh, A.M., and Lawrie, S.M. (2009). Working memory in

schizophrenia: a meta-analysis. Psychol Med *39*, 889–905.

4. Chafee, M. V, and Goldman-Rakic, P.S. (1998). Matching Patterns of Activity in Primate

Prefrontal Area 8a and Parietal Area 7ip Neurons During a Spatial Working Memory Task. J

Neurophysiol *79*, 2919–2940.

5. Chafee, M. V, and Goldman-Rakic, P.S. (2000). Inactivation of Parietal and Prefrontal

Cortex Reveals Interdependence of Neural Activity During Memory-Guided Saccades. J

Neurophysiol *83*, 1550–1566.

6. Harvey, C.D., Coen, P., and Tank, D.W. (2012). Choice-specific sequences in parietal cortex

during a virtual-navigation decision task. Nature *484*, 62–68.

1083

1084   7.   Zingg, B., Hintiryan, H., Gou, L., Song, M.Y., Bay, M., Bienkowski, M.S., Foster, N.N.,

1085   Yamashita, S., Bowman, I., Toga, A.W., et al. (2014). Neural networks of the mouse

1086   neocortex. Cell *156*, 1096–1111.

1087

1088   8.   Shadlen, M.N., and Newsome, W.T. (2001). Neural basis of a perceptual decision in the

1089   parietal cortex (area LIP) of the rhesus monkey. J Neurophysiol *86*, 1916–1936.

1090

1091   9.   Hanks, T.D., Kopec, C.D., Brunton, B.W., Duan, C.A., Erlich, J.C., and Brody, C.D. (2015).

1092   Distinct relationships of parietal and prefrontal cortices to evidence accumulation. Nature

1093   *520*, 220–223.

1094

1095   10.   Save, E., and Poucet, B. (2009). Role of the parietal cortex in long-term representation of

1096   spatial information in the rat. Neurobiol Learn Mem *91*, 172–178.

1097

1098   11.   Nitz, D.A. (2006). Tracking Route Progression in the Posterior Parietal Cortex. Neuron *49*,

1099   747–756.

1100

1101   12.   Mcnaughton, B.L., Mizumori, S.J.Y., Barnes, C.A., Leonard, B.J., Marquis, M., and Green,

1102   E.J. (1994). Cortical Representation of Motion during Unrestrained Spatial Navigation in the

1103   Rat. Cerebral Cortex *4*, 27–39.

1104

1105   13. Krumin, M., Lee, J.J., Harris, K.D., and Carandini, M. (2018). Decision and navigation in

1106   mouse parietal cortex. Elife *7*.

1107

1108   14. Goard, M.J., Pho, G.N., Woodson, J., and Sur, M. (2016). Distinct roles of visual, parietal,

1109   and frontal motor cortices in memory-guided sensorimotor decisions. Elife *5*.

1110

1111   15. Shenoy, P., and Yu, A.J. (2012). Strategic Impatience in Go/NoGo versus Forced-Choice

1112   Decision-Making. NeurIPS.

1113

1114   16. Oh, S.W., Son, S.J., Morris, J.A., Choi, J.H., Lee, C., and Rah, J.C. (2021). Comprehensive

1115   Analysis of Long-Range Connectivity from and to the Posterior Parietal Cortex of the

1116   Mouse. Cereb Cortex *31*, 356–378.

1117

1118   17. Kleinfeld, D., Giovannucci, A., Friedrich, J., Gunn, P., ré mie Kalfon, J., Brown, B.L., Ann

1119   Koay, S., Taxidis, J., Najafi, F., Gauthier, J.L., et al. (2019). CaImAn an open source tool for

1120   scalable calcium imaging data analysis. Elife.

1121

1122   18. Nelder, J.A., and Wedderburn, R.W.M. (1972). Generalized Linear Models.

1123

1124   19. Pho, G.N., Goard, M.J., Woodson, J., Crawford, B., and Sur, M. (2018). Task-dependent

1125   representations of stimulus and choice in mouse parietal cortex. Nature Communications

1126   2018 9:1 *9*, 1–16.

1127

1128   20. Chen, L.L., Lin, L.H., Green, E.J., Barnes, C.A., and McNaughton, B.L. (1994). Head-

1129   direction cells in the rat posterior cortex. I. Anatomical distribution and behavioral

1130   modulation. Exp Brain Res *101*, 8–23.

1131

1132   21. Nitz, D.A. (2012). Spaces within spaces: rat parietal cortex neurons register position across

1133   three reference frames. Nature Neuroscience 2012 15:10 *15*, 1365–1367.

1134

1135   22. Li, B., Nguyen, T.P., Ma, C., and Dan, Y. (2020). Inhibition of impulsive action by

1136   projection-defined prefrontal pyramidal neurons. Proc Natl Acad Sci U S A *117*, 17278–

1137   17287.

1138

1139   23. Kuan, A.T., Bondanelli, G., Driscoll, L.N., Han, J., Kim, M., Hildebrand, D.G.C., Graham,

1140   B.J., Wilson, D.E., Thomas, L.A., Panzeri, S., et al. (2024). Synaptic wiring motifs in

1141   posterior parietal cortex support decision-making. Nature 2024 627:8003 *627*, 367–373..

1142

1143   24. Cui, H., and Andersen, R.A. (2007). Posterior parietal cortex encodes autonomously selected

1144   motor plans. Neuron *56*, 552–559.

1145

1146   25. Crammond, D.J., and Kalaska, J.F. (2000). Prior information in motor and premotor cortex:

1147   activity during the delay period and effect on pre-movement activity. J Neurophysiol *84*,

1148   986–1005.

1149

1150 26. Churchland, M.M., Yu, B.M., Ryu, S.I., Santhanam, G., and Shenoy, K. V. (2006). Neural

1151   Variability in Premotor Cortex Provides a Signature of Motor Preparation. Journal of

1152   Neuroscience *26*, 3697–3712.

1153

1154 27. Pesaran, B., Pezaris, J.S., Sahani, M., Mitra, P.P., and Andersen, R.A. (2002). Temporal

1155   structure in neuronal activity during working memory in macaque parietal cortex. Nature

1156   Neuroscience 2002 5:8 *5*, 805–811.

1157

1158 28. Gottlieb, J.P., Kusunoki, M., and Goldberg, M.E. (1998). The representation of visual

1159   salience in monkey parietal cortex. Nature 1998 391:6666 *391*, 481–484.

1160

1161 29. Hulsey, D., Zumwalt, K., Mazzucato, L., McCormick, D.A., and Jaramillo, S. (2024).

1162   Decision-making dynamics are predicted by arousal and uninstructed movements. Cell Rep

1163   *43*, 113709.

1164

1165 30. Reynolds, J.H., and Chelazzi, L. (2004). Attentional modulation of visual processing. Annu

1166   Rev Neurosci *27*, 611–647.

1167

1168 31. Ito, M., and Gilbert, C.D. (1999). Attention Modulates Contextual Influences in the Primary

1169   Visual Cortex of Alert Monkeys. Neuron *22*, 593–604.

1170

1171 32. McAdams, C.J., and Maunsell, J.H.R. (2000). Attention to both space and feature modulates

1172   neuronal responses in macaque area V4. J Neurophysiol *83*, 1751–1755.

1173

1174   33. Herrington, T.M., and Assad, J.A. (2010). Temporal sequence of attentional modulation in

1175   the lateral intraparietal area and middle temporal area during rapid covert shifts of attention.

1176   J Neurosci *30*, 3287–3296.

1177

1178   34. Cossell, L., Iacaruso, M.F., Muir, D.R., Houlton, R., Sader, E.N., Ko, H., Hofer, S.B., and

1179   Mrsic-Flogel, T.D. (2015). Functional organization of excitatory synaptic strength in

1180   primary visual cortex. Nature 2015 518:7539 *518*, 399–403.

1181

1182   35. Wilson, D.E., Scholl, B., and Fitzpatrick, D. (2018). Differential tuning of excitation and

1183   inhibition shapes direction selectivity in ferret visual cortex. Nature *560*, 97–101.

1184

1185   36. Herrington, T.M., Masse, N.Y., Hachmeh, K.J., Smith, J.E.T., Assad, J.A., and Cook, E.P.

1186   (2009). The effect of microsaccades on the correlation between neural activity and behavior

1187   in middle temporal, ventral intraparietal, and lateral intraparietal areas. J Neurosci *29*, 5793–

1188   5805.

1189

1190   37. Chen, S., Liu, Y., Wang, Z.A., Colonell, J., Liu, L.D., Hou, H., Tien, N.W., Wang, T.,

1191   Harris, T., Druckmann, S., et al. (2024). Brain-wide neural activity underlying memory-

1192   guided movement. Cell *187*, 676-691.e16.

1193

1194   38. Steinmetz, N.A., Zatka-Haas, P., Carandini, M., and Harris, K.D. (2019). Distributed coding

1195   of choice, action and engagement across the mouse brain. Nature 2019 576:7786 *576*, 266–

1196   273.

1197

1198   39. Steinmetz, N.A., Aydin, C., Lebedeva, A., Okun, M., Pachitariu, M., Bauza, M., Beau, M.,

1199   Bhagat, J., Böhm, C., Broux, M., et al. (2021). Neuropixels 2.0: A miniaturized high-density

1200   probe for stable, long-term brain recordings. Science (1979) *372*.

1201

1202   40. Chen, H., Chen, S., and Deng, X. (2019). A Universal Nonparametric Event Detection

1203   Framework for Neuropixels Data. bioRxiv, 650671.

1204

1205   41. Stine, G.M., Trautmann, E.M., Jeurissen, D., and Shadlen, M.N. (2023). A neural

1206   mechanism for terminating decisions. Neuron *111*, 2601-2613.e5.

1207

1208   42. Aleman, A., Hijman, R., De Haan, E.H.F., and Kahn, R.S. (1999). Memory impairment in

1209   schizophrenia: A meta-analysis. American Journal of Psychiatry *156*, 1358–1366.

1210

1211   43. Cellard, C., Tremblay, S., Lehoux, C., and Roy, M.A. (2007). Processing spatial–temporal

1212   information in recent-onset schizophrenia: The study of short-term memory and its

1213   susceptibility to distraction. Brain Cogn *64*, 201–207.

1214

1215   44. Guo, J.Y., Ragland, J.D., and Carter, C.S. (2018). Memory and cognition in schizophrenia.

1216   Molecular Psychiatry 2018 24:5 *24*, 633–642.

1217

1218 45. Moses, M., Tiego, J., Demontis, D., Bragi Walters, G., Stefansson, H., Stefansson, K.,

1219   Børglum, A.D., Arnatkeviciute, A., and Bellgrove, M.A. (2022). Working memory and

1220   reaction time variability mediate the relationship between polygenic risk and ADHD traits in

1221   a general population sample. Molecular Psychiatry 2022 27:12 *27*, 5028–5037.

1222

1223 46. Pnevmatikakis, E.A., and Giovannucci, A. (2017). NoRMCorre: An online algorithm for

1224   piecewise rigid motion correction of calcium imaging data. J Neurosci Methods *291*, 83–94.

1225

1226 47. Digital Image Processing Using Matlab (Gonzalez) (2011).

1227

1228 48. Li, N., Daie, K., Svoboda, K., and Druckmann, S. (2016). Robust neuronal dynamics in

1229   premotor cortex during motor planning. Nature *532*, 459–464.

1230

1231 49. Finkelstein, A., Fontolan, L., Economo, M.N., Li, N., Romani, S., and Svoboda, K. (2021).

1232   Attractor dynamics gate cortical information flow during decision-making. Nat Neurosci *24*,

1233   843–850.

1234

1235 50. Chadwick, A., Khan, A.G., Poort, J., Blot, A., Hofer, S.B., Mrsic-Flogel, T.D., and Sahani,

1236   M. (2023). Learning shapes cortical dynamics to enhance integration of relevant sensory

1237   input. Neuron *111*, 106-120.e10.

1238

1239