1 **Sharp waves, bursts, and coherence: Activity in a songbird vocal**
2 **circuit is influenced by behavioral state**
3
4 Abbreviated title: Pallial-striatal dynamics and behavioral state
5

6 Author Names and affiliations: Corinna Lorenz[1,2]*, Anindita Das[2]*, Eduarda Gervini Zampieri
7 Centeno[3,5], Hamed Yeganegi[4,6], Robin Duvoisin[1], Roman Ursu[3], Aude Retailleau[3], Nicolas
8 Giret[2], Arthur Leblois[3], Richard H. R. Hahnloser[1], and Janie M. Ondracek[4†]
9
[1] Institute of Neuroinformatics, University of Zurich and ETH Zurich, 8057, Zurich,
Switzerland
[2] Institut des Neurosciences Paris-Saclay, CNRS, 151 route de la rotonde, 91400 Saclay,
France
[3] Bordeaux Neurocampus, Institut des Maladies Neurodégénératives, 146 rue Léo-Saignat -
33076 Bordeaux Cedex
[4] Technical University of Munich, TUM School of Life Sciences, Chair of Zoology, Liesel-
Beckmann-Straße 4, 85354 Freising, Germany
[5] Amsterdam UMC location Vrije Universiteit Amsterdam, Anatomy and Neurosciences, De
Boelelaan 1117, Amsterdam, The Netherlands
[6] Graduate School of Systemic Neurosciences, Ludwig-Maximilians-University Munich,
Großhaderner Str. 2, 82152, Planegg, Germany
(*) Authors contributed equally
(†) Corresponding author
24
25 Corresponding author email address: janie.ondracek@tum.de

26
27 Conflict of interest statement: The authors declare no competing financial interests.
28

Acknowledgements: This research was supported by the Agence Nationale de la Recherche
(SleepinBrainDyn) and the CNRS (International Emerging Action, project BirdSleep) to AD;
the EUR NEURO (EUR/7410-5) and Bordeaux Neurocampus - GPR BRAIN 2030
(GP/7200R-12) to EGZC; the Deutsche Forschungsgemeinschaft (ON 151/1-1) and the
Daimler und Benz Stiftung (PN 32-09/18) to JMO; the BayFrance program (FK23 2019) to
CL, HY, NG, and JMO. The authors would further like to thank and acknowledge Melanie
Dumont for excellent animal caretaking and Manon Rolland for help with experiments.

38
39

Abstract

Similar to motor skill learning in mammals, vocal learning in songbirds requires a set of interconnected brain areas that make up an analogous basal ganglia-thalamocortical circuit known as the anterior forebrain pathway (AFP). Although neural activity in the AFP has been extensively investigated during awake singing, very little is known about its neural activity patterns during other behavioral states. Here, we used chronically implanted Neuropixels probes to investigate spontaneous neural activity in the AFP during natural sleep and awake periods in male zebra finches. We found that during sleep, neuron populations in the pallial region LMAN (lateral magnocellular nucleus of the nidopallium) spontaneously exhibited synchronized bursts that were characterized by a negative sharp deflection in the local field potential (LFP) and a transient increase in gamma power. LMAN population bursts occurred primarily during non-rapid eye movement (NREM) sleep and were highly reminiscent of sharp-wave ripple (SWR) activity observed in rodent hippocampus. We also examined the functional connectivity within the AFP by calculating the pairwise LFP coherence. As expected, delta and theta band coherence within LMAN and Area X was higher during sleep compared to awake periods. Contrary to our expectations, we did not observe strong coherence between LMAN and Area X during sleep, suggesting that the input from LMAN into Area X is spatially restricted. Overall, these results provide the first description of spontaneous neural dynamics within the AFP across behavioral states.

Significance statement

Although cortical and basal ganglia circuits are known to be required for learning in both mammals and birds, little is known about the ongoing spontaneous activity patterns within these circuits, or how they are modulated by behavioral state. Here we prove the first description of cortical-basal ganglia network activity during sleep and awake periods in birds.

64 Within the pallial area LMAN, we observed population-wide bursting events that were highly

65 reminiscent of hippocampal sharp-wave ripple (SWR) activity, suggesting that large-scale

66 population events have diverse functions across vertebrates.

68 Introduction

During vocal learning, a juvenile bird transitions from acoustically simple, highly variable

"subsongs" to complex and stereotypical adult songs through a process of motor learning

(Kollmorgen et al., 2020; Konishi, 1965, 1970; McCasland & Konishi, 1981). Critically

involved in this learning process is a set of interconnected brain areas that make up a basal

ganglia-thalamocortical circuit (Fig. 1A) known as the AFP (Andalman & Fee, 2009; Bottjer

et al., 1984; Brainard & Doupe, 2000; Farries & Perkel, 2002; Scharff & Nottebohm, 1991)

Within the AFP, the pallial structure LMAN exerts a direct influence on vocal output through

its projection to premotor area RA (robust nucleus of the arcopallium), presumably

modulating downstream neural activity through bursts of spikes (Kao et al., 2008; Kojima et

al., 2013; Palmer et al., 2021). Through this influence, LMAN drives behavioral variability

that can be used for learning (Tumer & Brainard, 2007). Area X, part of the avian striatum

(Person et al., 2008; Pfenning et al., 2014), is thought to receive an efference copy of activity

in LMAN (Andalman & Fee, 2009; Vates et al., 1997; Vates & Nottebohm, 1995), thus

placing Area X in a position to evaluate LMAN exploratory activity in the context of the

ongoing song via a dopaminergic reward signal (Duffy et al., 2022; Kasdin et al., 2025; Qi et

al., 2025).

While the AFP shares many similarities with the mammalian corticostriatal circuit, including

cell types, neural connectivity, and gene expression patterns (Bolhuis et al., 2010; Gale &

Perkel, 2010; Murugan et al., 2013), it is anatomically distinct from the mammalian circuit.

For example, the avian pallium (including LMAN) displays a widely different anatomical

organization compared to mammalian cortex and is remarkable due to its lack of cellular

layers. Similarly, the songbird Area X is an aggregate of neural populations that are typically

segregated across various nuclei in mammalian basal ganglia (BG; (Person et al., 2008)).

Despite these differences, the overall function of the circuit seems to be similar across birds

and mammals, and many studies have shown that corticostriatal circuits are required for

learning in both birds (Ali et al., 2013; Andalman & Fee, 2009; Bottjer et al., 1984; Olveczky

et al., 2005; Scharff & Nottebohm, 1991; Tumer & Brainard, 2007; Zai et al., 2020) and

mammals (Kawai et al., 2015; Lemke et al., 2021; Santos et al., 2015).

Though the anatomical similarities and differences between these circuits have been well

characterized (Gale & Perkel, 2010), it remains unclear how these circuits compare in terms

of neural activity. Spontaneous network activity that occurs during non-task-oriented

behaviors, such as sleep or non-active awake periods, might be one way to broadly compare

circuit dynamics across species. While aspects of the mammalian corticostriatal circuit have

been well-studied in terms of neural activity across behavioral states (D. Liu et al., 2020) and

sleep (Mizrahi-Kliger et al., 2018, 2020), including its role in the control of sleep and

wakefulness (Lazarus et al., 2013) comparatively little is known about ongoing activity within

the AFP. With the exception of one study that characterized the presence of high gamma

oscillations in Area X during sleep (S. Y. Yanagihara & Hessler, 2012), spontaneous neural

activity patterns in the AFP - and how they are modulated by behavioral state - remain

largely unexplored.

We used a new approach (Lorenz, 2024) to chronically implant Neuropixels1.0 probes (Jun

et al., 2017) in order to investigate neural activity in the pallial-striatal network of the AFP

during periods of natural sleep and awake behavior in seven male zebra finches. By

characterizing spontaneous neural spiking activity in relation to ongoing LFP oscillations in

this avian pallial-striatal circuit, we gain insight into the different regimes of neural activity

that are present within this circuit across different behavioral states, and which helps inform

further studies of the analogous corticostriatal network in mammals.

Materials and Methods

Subjects

Seven male zebra finches (*Taeniopygia guttata*) were used in this study; data were pooled

across two laboratories in France (the Neuroscience Paris-Saclay Institute and the Bordeaux

Neurocampus). The age of the animals ranged from 76 days post-hatch (dph) to 232 dph at

the time of recording. Where it was possible in the figures, we have identified the 3 youngest

sub-adult animals (ZF-1, ZF-5, and ZF-7) with pink shading of their names. All animals were

bred and housed in the animal facilities of the laboratories. During the experiment, animals

were housed individually with water and food *ad libitum* under a 14:10 hr light/dark cycle.

Details about the individual animals can be found in Table 1.

All experimental procedures were approved by the French Ministry of Research and the

ethical committee "Paris-Sud et Centre" (CEEA No. 59, project 2017-25) and "Poissons

Oiseaux Nouvelle Aquitaine" (CEEA No. 73, project S73) and performed under the license

2015-25 and APAFIS#13413-2018020713145795.

Surgery and experiments

Chronic Neuropixels implantation surgery

The day of the surgery, food and water sources were removed from the cage approximately

30 minutes before the start of the surgery. Animals were anesthetized with isoflurane (0.6-

1.5% inhalation) and placed in the stereotactic apparatus once the toe flexion reflex was no

longer observed. Feathers were removed from the scalp and the area was disinfected with

ethanol and betadine. 0.05 microliters of diluted lidocaine (0.5mg/kg) was administered

subcutaneously and a local anesthetic (0.5 g Emla® creme) was applied to the exposed

skin. The scalp was resected and the exposed skull was prepared for the implant by drilling

several small holes in the outer bone layer. A small craniotomy contralateral to the final

implant was made to place a silver wire in contact with the dura that served as the electrical

ground.

The main craniotomy was made on the right hemisphere at approximately 1.7 mm lateral to

the superior sagittal sinus and 4.5 mm anterior to lambda, the confluence of sinuses, with

the flat part of the anterior skull rotated to a 50-degree angle. The positions of LMAN and

Area X were mapped with high-impedance single-electrodes in order to identify the brain

areas based on the firing rate properties thereof.

A Neuropixels probe version 1.0 (Jun et al., 2017) was implanted 4 to 5 mm deep to

penetrate the estimated center of LMAN and Area X. The custom-made holder that carried

the Neuropixels probe was fixed to the skull with dental cement and the craniotomy was

sealed with a two-part silicone gel (Dow Dowsil™, PN 3-4680) before it was covered with a

custom-made casing. In order to fix the Neuropixels headstage to the bird's head in a natural

position, the bird's head was rotated to a 35-degree angle in the stereotaxic during the

fixation. Finally, the ground wire was connected and open holes along the implant were

sealed with fast drying two-component silicone gel (Kwik-Sil).

The weight of the implant (~1.9 g) was immediately offset by a weight-relieving system. The

headstage was connected to an external counterweight using a nylon thread that was routed

through simple guide points positioned above and to the side of the setup, forming an

inverted U-shape. This allowed the counterweight to remain outside the recording area while

still providing vertical support and offsetting the implant's weight. The counterweight was

manually adjusted to ensure that the animals could move comfortably and were not

burdened by the implant.

Histology

Post-hoc control of the electrode location in the brain was performed through histological

examination of brain tissue. At the end of the experiment, birds were given a lethal injection

(i.p) of Exagon (pentobarbital sodium, 400 mg/mL), and in some cases perfused

intracardially (0.01 M Phosphate-Buffered Saline (PBS) followed by 4% paraformaldehyde

(PFA)). The brain was removed and post-fixed in 4% PFA for at least 24 hr. For 2 birds

recorded in Bordeaux, the brains were subsequently cryoprotected in 30% sucrose and cut

using a freezing microtome; for 3 birds recorded in Paris, brains were cut fresh at room

temperature using a vibratome. In all cases, the right hemisphere was cut in the sagittal

plane at a thickness of 50-80-microns. Images of the individual brain slices were taken using

bright-field and epiflurescence microscopy and analyzed using ImageJ software (Rasband

WS, NIH, Bethesda, Maryland, USA). The electrode tract, as indicated by

colored/fluorescent dye (DiI fluorescent dye; $DiIC_{18}(3)$), was clearly visible on consecutive

brain slices (see Fig. 1C; tissue from ZF2 and prepared on a vibratome).

2 birds died during the Covid-19 laboratory lockdown, and we were not able to retrieve the

brains in time to perform histology (Table 1). For these animals, the channel assignment was

performed on the basis of the electrophysiology, as described in detail below.

Electrode channel mapping

We defined the boundaries of LMAN and Area X based on darkness of the tissue in regular

light or the fluorescent background, as these two brain areas showed a clear contrast with

surrounding tissue due to higher myelination, cell density, and size (Nixdorf-Bergweiler et al.,

2007). The reconstruction of electrode channels to be assigned to LMAN or X was

performed based on the location of the electrode tip on the histological pictures, the

coordinates of the channels on the electrode and the above-mentioned boundaries of LMAN

and X. The limits of LMAN were also confirmed based on spontaneous activity, including

more bursts of large spikes in LMAN as compared to surrounding tissue.

For the 2 birds with missing histology, we relied heavily on these electrophysiological

properties, including the spike amplitude, rate, and response mapping that was performed

during the surgical probe implantation. Generally, LMAN channels were associated with

larger spike amplitudes, whereas Area X channels had higher spike rates.

Experimental design

Animals were housed individually in sound-attenuated recording chambers and neural

recordings were performed in sessions (one to 24 hours of continuous neural recording).

Electrophysiology

Extracellular activity was recorded with the chronically implanted Neuropixels 1.0 probe

which splits the signal on chip and samples the spike signal (bandpass filtered at 0.3 and 10

kHz) at 30 kHz and the LFP signal (bandpass filtered at 0.5 and 500 Hz) at 2500 Hz. Audio

signals were simultaneously recorded using a wall-attached microphone, pre-amplified and

digitized at 10, 20 or 32 kHz depending on the bird.

Sleep Recordings

Sleep-related activity was recorded either continuously throughout the night (n=3 birds;

Table 1) or at the beginning or end of the night (n=4). In the case of continuous recordings, a

one-hour segment was selected by visually inspecting the microphone data for a

consolidated quiet period indicative of sleep. In 5 out of 7 birds, sleep-related activity was

recorded with the light off during the normal dark period; however, in 2 birds (see Table 1),

the dark period was initiated earlier than normal, and recording sessions started in the late

afternoon (with light off).

Awake Recordings

For 4 out of the 7 birds (ZF-1, ZF-2, ZF-3 and ZF-5), we also examined brain activity that

occurred during normal daytime hours in the light. The behavior of the birds during this time

was characterized by singing, active movement, and periods of rest. Because it was possible

that birds were napping during periods of daytime rest, we classified the awake data into two

behavioral states on the basis of sounds detected from the microphone. "Light on active"

periods corresponded to microphone data which contained singing or sounds related to

movements. (These periods were distinct from movement-related voltage artifacts which

were detected as described below in section "Sound detection and movement artifact

identification"). "Light on non-active" periods corresponded to microphone data where no

sounds were detected. These could be periods where the bird was awake but not moving or

in a state of quiet daytime rest (see section "Sound detection and movement artifact

identification" and Table 2 for more information).

Note that by segmenting activity solely based on microphone-detected sounds, we may

neglect neural activity changes occurring in Area X and LMAN prior to song initiation

(Hessler & Doupe, 1999a; Woolley et al., 2014),. Such premotor activity could therefore be

misclassified as occurring during "Light on non-active." However, this misclassification is

likely uncommon, as zebra finches are generally behaviorally active in the periods

surrounding song bouts (Zann, 1996).

Statistical analysis

Data preprocessing and spike clustering

For neural spiking analysis, spike sorting was performed in four of the seven birds on all

active channels using Kilosort 1.0 or Kilosort 2.5 (Pachitariu et al., 2016; see Table 1 for

more information) and manually curated using phy (https://github.com/kwikteam/phy) in four

birds (ZF-1, ZF-2, ZF-3 and ZF-5) or automatically curated using SpikeInterface (script

available in our data repository) in one bird (ZF-4). Regardless of the sorting process, a

user-based quality check of the final dataset ensured that the quality of sorted units was

satisfactory and homogeneous.

Sound detection and movement artifact identification

We identified periods of putative awakening during the light-off period by analyzing the

microphone recordings. We calculated the root mean square (RMS) of the microphone

signal and manually defined a threshold that was just above the noise level (Fig. 3A, Fig.

S1). Each threshold crossing identified a putative movement-related sound, and 1-s bins

during which sound was detected were excluded from further analysis.

During the light-off recordings, it is possible that the bird could awaken briefly but not move,

which would not be identified as a putative awakening by our sound detection approach.

Although we are not able to quantify the extent to which such events may contribute to our

dataset, other papers which used video recordings to confirm sleep states have reported

discarding between 2% (Yeganegi & Ondracek, 2025) and 3% (Yeganegi & Ondracek, 2023)

of sleep EEG data due to movement-related artifacts. Across our birds, the percentage of

data discarded using the sound detection method ranged from 0.12% to 6.02% (see Fig. S1

for exact values). Therefore, although our approach using sound detections to identify

putative awakenings cannot identify quiet awakenings in the dark, we estimate that this

would affect at most an additional 3% of the data. We evaluate and discuss the potential

inclusion of quiet awakenings in our sleep analysis in the Discussion and in Text S1.

For the analysis of awake neural activity, we similarly leveraged the microphone recording to

distinguish between silent periods and periods of activity. Periods where sounds were

detected were identified as "Light on active" periods, and silent periods were identified as

"Light on non-active."

The awake data often exhibited movement-related artifacts in the LFP which we did not

observe in the putative sleep recordings. These daytime movement artifacts were

characterized by synchronous fluctuations across all channels. To address this, we applied a

modified version of the Artifact Subspace Reconstruction (ASR, (Chang et al., 2020; Kothe &

Makeig, 2013) algorithm to identify and exclude artifact-contaminated periods. Briefly, we

performed an eigenvalue decomposition of the data covariance matrix. Components

associated with large eigenvalues indicative of high-variance noise, were flagged as

potential artifacts. These eigenvalues were identified using a threshold defined as a multiple

q of the median eigenvalue. To detect specific times of contamination, we reconstructed the

LFPs using only the artifact-related components and computed the summed signal across all

channels. Time bins where this signal exceeded a threshold—defined as the mean plus k

times the standard deviation—were marked as contaminated and excluded from further

analysis. Both k (ranging between 3 and 12) and q (ranging between 10 and 30) were

manually tuned for each bird to conservatively exclude movement artifacts without

discarding valid neural activity.

Sleep staging

The sleep staging was performed using an unsupervised k-means clustering algorithm

previously developed in songbirds to classify sleep data on the basis of multiple EEG

features (Canavan & Margoliash, 2020; Low et al., 2008). This algorithm classifies sleep into

one of 3 states: SWS, REM sleep, and intermediate sleep (IS) which has been compared to

the mammalian NREM sleep stage N2 and often contains a mixture of low-amplitude delta

and higher-frequency elements such as theta (Canavan & Margoliash, 2020). Our use of this

method diverged from its original implementation in the following ways: 1) we used LFP

instead of EEG to compute the sleep stages, 2) our exclusion of awake periods was based

on sound detection from the microphone (as described above), and did not include manual

scoring of video data to confirm that animals in the dark were sleeping prior to sleep state

classification, as was done in (Canavan & Margoliash, 2020). The LMAN LFP channel used

in the analysis was a centrally located LMAN channel along the dorsoventral axis of

recording. Spectral parameters were computed from the multi-taper spectral analysis

(Prerau et al., 2017) of the filtered (0.5 - 60 Hz) and down sampled (250 Hz) LFP using 3 s

bins (1 s overlap). A small number of bins were not classified as REM, SWS or IS using the

two-step clustering algorithm (Canavan & Margoliash, 2020) and were labeled 'Not

classified/Nc'. Nc classifications arise when a bin is classified as SWS in the first clustering

round and then as REM in the second clustering round.

Our use of a centrally located LMAN LFP channel to classify sleep stages (instead of a

superficially located EEG electrode) means that our sleep staging results may diverge

slightly from values reported elsewhere (Canavan & Margoliash, 2020; Low et al., 2008;

Yeganegi & Ondracek, 2023, 2025). However, LFP signals were filtered in the same way

that EEG signals were filtered (i.e., between 0.5 and 60 Hz), which would eliminate high-

frequency characteristics that might be present in an LFP recording but not in an EEG

recording.

Another concern was that the LFP recording centered in LMAN might not reflect "global"

sleeping activity. Indeed, recent work has shown that sleeping brain dynamics are not as

global as previously thought, and different sleep states might occur in different brain regions,

as was also reported for songbirds (Yeganegi & Ondracek, 2025). To investigate this in our

own data (Fig. S2), we examined the differences in sleep classifications in a superficial LFP

located in the hyperpallium (comparable to a superficially implanted EEG electrode) and

compared it to a centrally located LMAN channel (Fig. S2A). We found that almost 90% of

the sleep scoring matched between the LMAN channel and the superficial LFP channel (Fig.

S2B), suggesting that the use of an LMAN channel to assess sleep state is a fair

assessment of the ongoing sleep states that would have been recorded using a superficial

EEG electrode. Of the 11% of mis-matched sleep states, the largest category involved a

mismatch between SWS and IS, such that sleep states that were identified as SWS in LMAN

were identified as IS in the hyperpallium (Fig. S2C).

LFP Coherence Analysis

Data inclusion

For the coherence analysis we only used bins identified as either REM or SWS. Time points

at which movements were detected, as described in the previous section (Sound detection

and movement artifact identification) were excluded for the coherence analysis.

LFP pre-processing

First, we extracted the LFP data from *.bin* files utilizing SpikeInterface

(https://spikeinterface.readthedocs.io/en/latest/overview.html) with phase shift correction

using *spikeinterface.preprocessing.phase_shift*. Next, we removed artifacts from the LFP

signal by establishing a threshold based on 80% of the maximum absolute trace value and

interpolating the signal around the artifact time point with a margin of 20 samples before and

after. Next, we z-scored the trace using *elephant.signal_processing.zscore.*

Coherence Analysis – Sleep recordings

Pairwise LFP coherence was computed from the z-scored LFP traces. We included

every 4[th] LFP channel (corresponding to a dorso-ventral linear distance of approximately 40

microns between them) spanning Area X and LMAN to reduce the computation time while

still retaining enough spatial resolution to obtain a clear estimate of the spatial variation. We

performed pairwise LFP coherence per identified state. For sleep recordings, this involved

computing coherence for the SWS and REM states respectively. Specifically, we computed

the magnitude squared coherence $C_{1,2}$ of LFP signals L1 and L2 as follows:

$$C_{1,2} = {|P_{1,2}|^2}\Big/{P_{1,1}\, P_{2,2}}$$

where $P_{1,1}$ and $P_{2,2}$ are power spectral density estimates of L1 and L2 using Welch's

method, and $|P_{1,2}|$ is the absolute value of the cross spectral density estimate. We used the

scipy.signal.coherence (https://docs.scipy.org/doc/).

The accuracy of coherence magnitude estimates depends on 1) the length of the segments

used for computation that is usually determined by the lowest frequency of interest (FOI) and

2) the total number of such segments obtained (equal to the total length of the signal being

analyzed divided by the segment length). In our case, these parameters were constrained by

our lowest FOI which was 1 Hz and the total duration of REM or SWS states in our

recordings per bird. We selected a segment length of 3 sec for our analysis, since 3 sec

segments were sufficient to give an accurate coherence estimate for our lowest FOI (1 Hz)

and our total length of recordings corresponding to SWS or REM were large enough to

obtain sufficient number of 3-sec segments.

After accounting for removal of movement artifacts and the analysis of only the REM and

SWS segments, the total percentage of data analyzed in the coherence analysis per bird

were: ZF1: 78.2%, ZF2: 78.4%, ZF3: 65.9%, ZF4: 76.2%, ZF5: 85.8%, ZF6: 77.4%, ZF7:

71.4%. Because the number of SWS segments were more numerous that the number of

REM segments, for each channel comparison, we initially performed the coherence

computation using a randomly selected number of SWS segments equal to the total number

of REM segments. We then averaged across all SWS segments after ascertaining that the

standard error of mean for the averaged coherence for all the SWS segments was low (<

0.05 for all channel comparisons for all birds) indicating low variation between randomly

selected SWS segments.

In order to evaluate statistical significance in the coherency analysis between SWS and

REM states, we first selected a minimum number of channels to compare that was the same

for all birds for each brain area (LMAN = 7; equal to ~240 microns, Area X = 26; equal to

~1000 microns). This was necessary as the probe implantation was variable across animals

and resulted in different number of total channel counts in the brain areas of different birds.

Then we identified the most central channel in each brain area and included neighboring

channels along the probe length. After pooling these channel comparisons across birds for

SWS and REM states, we identified significant differences across brain states in the

coherence analysis using a Wilcoxon Signed Rank test.

Coherence Analysis – Awake recordings

A similar procedure was followed for analysis of the awake daytime recordings (obtained from 4 out of the 7 birds). Here our goal was to determine whether behavioral state affected the coherence between brain areas. That is, we compared sleep-related activity to periods of Light on active and Light on non-active states. Only 3 out of the 4 birds (ZF-1, ZF-2, and ZF-4) had sufficiently long periods of both Light on active states to allow us to compare the pair-wise coherence analysis (see Table S1 for more information). Coherence values were pooled across birds as described above. To compare the awake and putative coherence results for these 3 birds, we performed statistical testing using a Friedman Test followed by the Nemeyni posthoc tests to account for multiple comparisons between non-independent samples (same channels and 4 different behavioral states).

Analysis of spiking variability

To assess how spike activity varied over time within LMAN and Area X during putative sleep, we computed the coefficient of variation (CV) in consecutive, non-overlapping segments of 10 seconds. Within each segment, we counted the total number of spikes (ns, from all units either in LMAN or in Area X) in sub-windows of 20 milliseconds. The CV was then computed as the standard deviation of spike counts divided by the mean spike count across all sub-windows within that segment (CV =σ_{ns}/μ_{ns}). Because we were interested in the relationship between the temporal variability of spike activity within a population and how it related to putative sleep, we computed the δ/γ ratios for the LFP signal of the central electrode in LMAN in the same 10-second segment. We then used Pearson's correlation coefficient (Pearson, 1895) and type-II linear regression (Ricker, 1973) to assess the relationship between δ/γ ratios and the CV of spike activation. The average firing rate of neurons in LMAN and Area X was calculated by determining the total spike count per second within the same windows where CV was computed. We performed this analysis for sleep recordings in

five birds (ZF-1 to ZF-5), excluding segments that were contaminated by potential movement

(see *Sound detection and movement artifact identification*).

Population synchronized burst detection and spectral analysis of associated

LFP

For each neural unit (single or multiunit), bursts were identified when the Instantaneous

Firing Rate (IFR) exceeded 100 Hz. The fraction of simultaneously bursting LMAN units was

determined in a sliding 50 ms window with 10 ms step size. The resulting trace was

smoothed using a Gaussian kernel with 15 ms standard deviation, and candidate events of

synchronized bursts were classified as peaks exceeding the 99^{th} percentile.

We performed this analysis for sleep-related recordings of five birds (ZF-1 to ZF-5). Events

that occurred during the 1-sec segments that were flagged as putative awakenings or Nc

sleep states were excluded. Similarly, in daytime recordings, events that overlapped with

LFP movement artifacts were also excluded (see *Sound detection and movement artifact*

identification).

To create control events, we used random time points in the recording. For assessing the

LFP around synchronized bursts and control time points, we selected the central LMAN LFP

and compared raw traces that were offset-corrected by median subtraction. Interested in a

high-resolution description of the non-stationary spectro-temporal content of the associated

LFP, we applied a complex wavelet transform using complex Morlet (Gabor) wavelets (*cwt,*

MATLAB, The MathWorks, Natick, MA). Scalograms were computed over a 1-sec window

centered on each event. Vice versa, to assess the oscillatory context of these events, we

calculated the power estimate in the δ- (1-4 Hz) and γ-band (30 – 90 Hz) in the 3-sec

window around the event, the same bin size used for sleep staging, using Thomson's

multitaper power spectral density method (*pmtm*, MATLAB, The MathWorks, Natick, MA).

For the high γ-band (90 – 145 Hz), we used a narrower window of 300 ms centered to the

event.

Data and software availability

All research data as well as scripts to replicate the results are available in the corresponding

repository: https://gin.g-node.org/E4-Leblois/npx_sleep_2024

Spectral analysis and visualizations in Fig. 3B-D and S4 were computed with an adapted

code of the following repository https://github.com/cortex-lab/spikes.

Results

Distinct regimes of activity are present during avian sleep

We investigated neural activity in the pallium and basal ganglia of male zebra finches (n = 7)

chronically implanted with Neuropixels 1.0 probes (Fig. 1B). In each animal, the probe was

implanted such that both the pallial structure LMAN and the striatal structure Area X within

the avian BG (Person et al., 2008) were targeted (Fig. 1C).

We observed that during putative sleep, slow wave delta activity (1-4 Hz) in the LFP was

widespread throughout the pallium and BG, spanning more than 4 mm (Fig. 1D, black

arrowheads). Neural spiking was synchronized with the slow wave dynamics and was

observable as sparsely bursting patterns of activity in both pallial and BG neurons (Fig. 1D-

Spiking Activity). These slow wave dynamics and periods of synchronized bursting were

entirely absent from awake data (Fig. 1 E).

We used two complementary approaches to further analyze putative sleep states: 1) we

analyzed sleep as a continuous variable using the LFP δ/γ power ratio, and 2) we

categorized sleep into discrete stages using an automated avian sleep clustering algorithm

(Canavan & Margoliash, 2020; Low et al., 2008). Whereas the former approach

acknowledges the continuous nature of sleeping brain dynamics, the latter approach allowed

us to numerically characterize certain aspects of avian sleep and compare our results with

other avian sleep studies.

To analyze the continuous nature of sleeping brain dynamics, we calculated the δ/γ ratio as

the ratio of delta activity (δ, 1-4 Hz) to gamma activity (γ, 30-90 Hz). This method is

analogous to metrics that have been used with rodents (Csicsvari et al., 1998; O'Keefe,

1976), but accounts for the dominant frequency bands present during avian sleep (Low et

al., 2008; Yeganegi et al., 2019; Yeganegi & Ondracek, 2023).

We calculated the δ/γ ratio for a large range of pallial and BG sites (Fig. 2A). Whereas δ/γ

ratios calculated for LFPs within the BG were mostly flat (Fig. 2A, blue lines), δ/γ ratios

calculated for pallial LFPs were large and dynamic (Fig. 2A, pink lines, also black line).

Within the pallium, we observed that LPF traces with low δ/γ values had REM sleep-like

characteristics (i.e., low amplitude LFP signals), while LFP traces with high δ/γ had SWS-like

characteristics (i.e., large amplitude LFP signals; Fig. 2B). Median δ/γ values (Fig. 2C) were

calculated using a centrally located pallial LMAN LFP for each bird individually and showed

some variability across birds that matched the inter-individual variability reported previously

using similar metrics (Yeganegi & Ondracek, 2023). δ/γ values calculated using a LMAN

LFP were generally higher than δ/γ values reported using a hyperpallial EEG electrode

(Yeganegi & Ondracek, 2023). We used the centrally located LMAN channel quantified in

Fig. 2C for all subsequent analysis using the δ/γ ratio.

In addition to the δ/γ ratio, we used a clustering-based sleep scoring method (Canavan &

Margoliash, 2020; Low et al., 2008) to classify sleep stages into 3 states (see Methods for

more details). For this analysis, we used the same centrally located LMAN LFP that we used

in the δ/γ analysis. We observed LFP segments that were classified as SWS were indeed

characterized by high δ/γ values, and LFP segments that were classified as REM sleep were

characterized by low δ/γ values (Fig. 2D, E) with IS states falling in between the two

distributions (Fig. 2F). 3-D plots showed that sleep states were well-segregated on the basis

of the spectral features used in the clustering-based approach (Fig. 2G, H). See Fig. S3 for

sleep classification distributions for all birds.

Different spectral features characterize sleep and awake states

Were the activity patterns that we observed during sleep different than those observed

during wakefulness? In the next analysis, we examined the spectral features of the LFP

signals recorded in LMAN and the surrounding pallium as well as Area X and the

surrounding BG during awake and putative sleep phases (Fig. 3A, B). In contrast to the

sleep-related activity which was recorded in the dark, the awake recordings took place in the

light during the normal day-time hours and captured awake behavior (e.g., singing, moving,

resting; Fig. S1).

We observed in LMAN (Fig. 3C, dark gray shading) and the surrounding pallial channels

(Fig. 3C, light gray shading) that the power was higher for a broad range of low frequencies

during sleep compared to awake states (Fig. 3C, 1-4 Hz, 4-12 Hz, 12-30 Hz, black

arrowheads, thin line compared to thick line). In contrast, for LFPs recorded within Area X

(Fig. 3C, dark green shading) and the surrounding BG (Fig. 3C, light green shading) only the

power in the theta range (Fig. 3C, 4-12 Hz) was occasionally higher during putative sleep

compared to awake states (Fig. 3C, black arrowhead, thin line compared to thick line; see

also Fig. S4 for data from other birds).

The power in the high-frequency gamma range (Fig. 3C, 90-145 Hz) was higher for LMAN

LFPs during awake states compared to sleep states in all birds (Fig. 3C, black arrowhead,

see also Fig. S4).

Across depths, we saw that putative sleep states were associated with higher power in low

frequency bands compared to awake states (Fig. 3D), which is in line with results from brain

areas including RA (Hahnloser et al., 2006) and Area X (S. Y. Yanagihara & Hessler, 2012).

Importantly, the LFP associated with the LMAN depth (Fig. 3D, black line; 2.6 mm) was

strongly modulated by sleep. We saw that LFPs at the deepest part of the probe followed the

common 1/f power spectrum but generally carried less power (Fig. 3D, light green line: 0.1

mm from probe tip).

Within-area coherence is high during SWS in LMAN and Area X

Given the clear spectral differences that we observed between LMAN and Area X during

sleep and awake states, we focused our next analysis on investigating the functional

connectivity between these brain areas. We characterized the LFP coherence 1) within

LMAN, 2) within Area X, and 3) across LMAN and Area X for 3-s SWS and REM sleep LFP

segments and for different canonical frequency ranges that have been shown to play an

important role in behavior (Buzsáki & Draguhn, 2004) for all birds (Fig. 4A, B).

We observed the average pairwise LFP coherence across birds to be significantly modulated

by sleep states (Fig. 4B, Fig. 4C). Whereas coherence in the low frequency ranges was

spatially widespread during SWS (Fig. S5), gamma band coherence was limited to spatially

neighboring channels with the coherence values dropping to 50% of its peak value within

150 microns. This is consistent with the idea that gamma oscillations reflect local network

dynamics (Buzsáki & Wang, 2012; Cardin, 2016; Cardin et al., 2009).

Since the pairwise LFP coherence varied as a function of the distance between the electrode

sites and the total number of electrodes within brain regions varied slightly across birds due

to variations in implant angle or exact implant co-ordinates, we compared the same number

of LMAN and Area X electrodes sites across birds to compare the median coherences

between sleep states. The sites were selected such that they spanned the same distance

from the identified centers of these brain areas for all 7 birds (see Methods for more details).

We found that SWS and REM sleep coherence values were significantly different for all

frequency bands analyzed and across all brain areas (Fig. 4D; Table 2). During SWS within

LMAN, we observed very high delta band coherence, ranging from 0.86 - 0.99 (min - max,

median = 0.96; Table 2). During REM sleep within LMAN, the delta band coherence was still

high (median = 0.87), but the range of values was larger (0.63 - 0.99; min - max), indicating

a decrease in coherence between some LMAN pairs during REM sleep.

Within Area X, median coherence values were lower for all frequency ranges and sleep

states compared to within LMAN coherence values (Fig. 4B, C; Table 2). Median coherence

values within Area X were lower during REM sleep compared to SWS for all frequency

ranges (Fig. 4C, Fig. 4B green dots). This finding is consistent with primate literature

demonstrating significantly lower coherence within basal ganglia compared to cortical areas

during SWS (Mizrahi-Kliger et al., 2018).

LMAN sends direct projections to Area X, which in turn sends feedback projections to LMAN

through thalamic area DLM (Fig. 1A; (Farries et al., 2005)). If the closed-loop structure of the

circuit plays an important role in modulating sleep dynamics, we may expect strong cross-

area coherence.

While median coherence values were low across frequency bands for LMAN-Area X LFP

pairs compared to within-LMAN or within-Area X LFP pairs (Table 2), a few individual LMAN-

Area X LFP pairs showed high coherence values > 0.5, especially for the low frequencies (1-

12 Hz; Fig. 4B, blue dots). This finding suggests that the input from LMAN to Area X may be

spatially restricted, as only a few pairs shared similar oscillatory drive during sleep.

We did not find a strong effect of sleep state on the across-area LMAN-Area X coherence.

Although significant differences were found across all frequency bands (Fig. 4C, right), the

median differences between SWS and REM sleep were quite small (Table 2).

LMAN-Area X dynamics are modulated by behavioral state

We were curious as to how the LFP coherence changed as an effect of behavioral state, and

so we compared the sleep-related LFP coherence with coherence analyzed for daytime

"Light on active" and "Light on non-active" periods for a small group of birds (n=3 birds; Fig.

4D). The sleep-related coherence analysis for this smaller group of birds was consistent with

the results of the larger group of seven birds (see Table S2): coherence values were highest

within LMAN compared to within Area X and across LMAN and Area X pairs for all frequency

bands, and coherence values were higher during SWS compared to REM sleep across

frequency bands, with the exception of high gamma within LMAN. In this case, REM

coherence was higher compared to SWS coherence within LMAN; however, this difference

was not significant for the smaller group of three birds (Fig. 4D, LMAN-LMAN comparison;

High gamma, n.s.).

When we compared sleep-related coherence to daytime coherence (i.e., Light on active and

Light on non-active), we saw that the delta and theta band coherence values were

significantly lower for daytime periods, suggesting that periods of putative sleep in the dark

(i.e., SWS and REM states) were distinct from periods of active and non-active behaviors

recorded during the daytime.

Interestingly, when we analyzed the coherence trends within the daytime periods, we

observed that periods of daytime active behavior were associated with higher delta band

coherence values compared to non-active periods of daytime. This was true for within

LMAN, within Area X, and across LMAN-Area X LFP pairs, suggesting that oscillatory

activity in the AFP differs during non active daytime periods compared to nighttime sleep

Phases of high δ/γ activity during sleep are associated with

increased spiking variability for neurons in LMAN but not Area X

Using our Neuropixels recordings, we leveraged the high-pass filtered electrode signals to

investigate the spiking activity in LMAN and Area X during sleep in n = 5 birds (Table 1).

Spikes were identified and assigned using state-of-the-art automatic detection and sorting

algorithms, either in a fully automated way or followed by meticulous manual curation to

ensure accuracy and reliability (see Methods). The results of any spiking analysis did not

differ qualitatively across these two preprocessing approaches.

During sleep, we observed increased but sparse population-wide spiking activity which was

locked to negative LFP deflections and interleaved with periods of neural silence, followed

by sudden switches to periods of random spiking and lower LFP amplitudes (see Fig. 1D).

To quantify the different patterns of spiking activity that we observed during sleep, we

calculated the coefficient of variation (CV, standard deviation divided by the mean) for

identified LMAN and Area X neuron populations in 10 s bins.

The CV can be thought of as a measure of collective variability in coincident spike trains: CV

values for the population rate will approach zero when all neurons are statistically

independent (Kobak et al., 2019). This is because random irregularities stemming from

individual neurons will average out as the population size increases, ultimately leading to a

stable population rate. However, when neurons are synchronized and activity oscillates, the

variability from one moment to another in the population rate is high.

We found that median CVs ranged from 0.27 in Area X to 2.14 in LMAN (Fig. 5A; see Table

4 for individual values), and CV values were significantly higher for LMAN neurons

compared to Area X neurons in all birds (median [min, max] LMAN CV over all birds = 1.44

[0.73, 2.14]); median [min, max] Area X CV for all birds was = 0.61 [0.27, 1.04]).

We observed that the CV values fluctuated closely with δ/γ values, such that high CV values

were associated with high δ/γ values (Fig. 5B). This trend was especially obvious for LMAN

neurons (Fig. 5C, D). Indeed, regression fits between CV values for LMAN neurons and δ/γ

values returned higher slopes (Fig. 5C, D) and both variables were highly correlated for all

birds (Table 5). This was not the case for Area X neurons: estimated regression slopes were

lower, and only 2/5 birds showed significant positive correlations between CV values and δ/γ

values (ZF-1 and ZF3), whereas 1 bird showed a significant negative correlation (ZF-4,

Table 5; Fig. 5C, filled symbols).

High CV values for LMAN neurons compared to lower CV values for Area X neurons (Fig.

5A) suggest that the spike trains for LMAN neurons were more irregular than for Area X

neurons. Indeed, when we examined spike rasters corresponding to high CV values for

LMAN populations (Fig. 5D, E), spiking patterns were conspicuously irregular - albeit largely

synchronized as bursts of spikes across the population of LMAN neurons. In contrast, CV

values were significantly lower for Area X neurons (Fig. 5A), capturing their regular spiking

patterns and high firing rates (Fig. 5D, F).

It is important to note that the δ/γ trace used in this analysis was calculated from a centrally

located LMAN LFP (see Methods for details). Therefore, strong correlations between CV

values and δ/γ values in LMAN may reflect the tight correspondence between spiking activity

and LFP oscillations in LMAN. The correspondence between spiking activity in Area X and

the LMAN LFP was much more heterogeneous, suggesting that oscillatory activity in LMAN

did not directly influence Area X spiking activity during sleep as it did in LMAN.

Synchronized population bursting in LMAN during sleep is

associated with a negative deflection in the LFP

Our findings suggested that within LMAN, populations of neurons are co-activated at specific

time points during putative sleep. In order to explore the synchronized bursting of LMAN

neurons during putative sleep - and how they are related to the LFP (Fig. 6A) - we identified

and thresholded bursting events in the population of LMAN neurons (Fig. 6B). Then, we

used these synchronized bursts to identify activity patterns in the LFP, which we called

"burst-aligned LFP" (Fig. 6C). Individual burst-aligned LFP segments generally had higher

power in frequency ranges < 10 Hz (example in Fig. 6D), and the average burst-aligned LFP

showed a prominent low-frequency power burst centered on zero-time lag (Fig. 6E). Indeed,

when we compared the bursting-aligned LFP segments (Fig. 6C, lower blue line) to

randomly selected LFP segments (Fig. 6C, lower black line), the bursting-aligned LFP

segments showed a notable increase in power for frequencies lower than 10 Hz. This

phenomenon was present across birds: every bird examined had a notable negative

deflection that corresponded to the population synchronized bursting (Fig. 6F).

Having established that synchronized bursting in LMAN during putative sleep is consistently

associated with prominent low-frequency deflections in the LFP, we next asked whether

similar events occur during wakefulness. Specifically, in the four birds for which we recorded

neural activity during daytime periods, we examined whether synchronized LMAN bursting

events also occur during active and resting periods, and if so, whether the associated burst-

aligned LFP exhibits features similar to those observed during putative sleep.

During light on non-active periods, burst-aligned LFPs in ZF-2 and ZF-3 resembled those

seen during putative sleep, exhibiting a negative deflection in the LFP at the time of the

synchronized bursting and elevated low-frequency power in the scalogram, although with

reduced overall magnitude (Fig. 6G; Fig. S6). In contrast, ZF-1 exhibited no detected events

during rest, and ZF-5 showed only 11 events, none of which were associated with a clear

LFP deflection.

In active light on periods, we observed numerous synchronized bursting events across all

birds except ZF-3, which was generally inactive during this recording (Fig. S1). The detected

events coincided with behaviors such as singing, calling, or general movement, consistent

with the increased bursting activity characteristic during vocal production (Hessler & Doupe,

1999b). However, the associated LFPs remained flat across all birds, showing no consistent

deflection time-locked to the bursts (Fig. 6G, long dashed line; Fig. S6). This absence of

coordinated low-frequency structure in the LFP distinguishes these active-state bursting

events from those observed during putative sleep, where synchronized bursts are reliably

coupled with prominent negative LFP deflections (Fig. 6F).

Consistent with these observations, Fig. 6H summarizes the average LFP amplitude at the

time of synchronized bursts across states, showing the strongest negative deflections during

putative sleep, weaker deflections during light on rest, and values near zero during active

wakefulness. This graded pattern supports a state-dependent modulation of LFP responses

to synchronized bursting in LMAN.

679 Synchronized population bursts in LMAN occur predominantly

680 during NREM sleep

We were curious as to whether these synchronized bursting events in LMAN occurred during

a specific sleep regime, i.e. either during NREM sleep (i.e., SWS and IS sleep) or during

REM sleep (Fig. 7A). With the exception of 1 bird (Fig 7A, ZF-4: upward triangle), the

synchronized bursting events occurred predominantly during NREM sleep.

We also examined the δ/γ values surrounding the synchronized bursting events, with the

expectation that these events might occur during LFP segments with higher δ/γ values (i.e.,

during SWS). Interestingly, we found that the bursting events were not consistently

associated with lower δ/γ values compared to control events across birds (Fig. 7B; see

Methods for details). Because the δ/γ is a ratio, low δ/γ values can result from a decrease in

δ power or an increase in γ power. We examined the δ and γ power separately for bursting

events compared to control events. We found that while the power in the δ range was indeed

elevated during bursting events compared to control events for all but one bird, ZF-4, (Fig.

7C), a larger increase in γ power during the bursting event (Fig. 7D) compared to control

segments led to the overall inconsistent δ/γ values we observed during bursting events

compared to controls (Fig. 7B).

When visualized as the difference between the burst and control segments (Fig. 7E), this

burst-associated increase in power extended from delta up through high gamma and was

present for all birds analyzed (Fig. 7F).

702 Discussion

In order to explore the ongoing spontaneous activity patterns of the AFP during sleep and

awake behavior in songbirds, we investigated the activity patterns of neurons in LMAN and

Area X (Fig. 1A, C) using chronically implanted Neuropixels probes (Fig. 1B).

Coherence between field potentials has been used in a multitude of studies as a mesoscopic

measure of functional connectivity between brain areas (Schneider et al., 2021) and is

believed to reflect the neuronal connectivity (Buzsáki & Schomburg, 2015) and oscillatory

synchronization (Bastos et al., 2015) in these brain areas.

Importantly, we observed strong significant differences between sleep-related coherence

(i.e., SWS and REM) and awake-related coherence (i.e., light on non-active and light on

active) across brain regions and passbands (Fig. 4D, see exact values in Table 3). This

finding demonstrates that the birds were in a different behavioral state in the dark compared

to recordings in the light, and suggests that our sleep recordings were not overly

contaminated with undetected quiet awakenings. If this had been the case, we hypothesize

that the sleep-related coherence values would more strongly resemble the light on non-

active coherence values, which they do not (see Text S1 for further discussion).

Contrary to our expectations, we did not observe strong across-area coherence between

LMAN and Area X on average, although individual LMAN-Area X pairs did show strong

coherence > 0.5. In contrast, studies of motor skill learning in rodents found an increase in

cross-area coherence between motor cortex and striatum during sleep (Koralek et al., 2013;

Lemke et al., 2021). The fact that we did not observe strong cross-area coherence may stem

from the fact that the birds in this study were not actively engaged in a learning paradigm.

A previous study demonstrated the presence of phasic LFP gamma oscillations in Area X

during sleep, with singing-modulated Area X units phase-locked to gamma oscillations (S. Y.

Yanagihara & Hessler, 2012). Our pairwise LFP coherence results, albeit a mesoscale

measure of synchronization across brain areas, could be a reflection of this phase-locking

and the concerted activity of anatomically connected LMAN units and Area X units during

awake periods. Verification of this hypothesis would require further data collection in awake

and singing birds to investigate the specific neural activity of different neuronal subtypes and

their relation to the ongoing LFP gamma oscillations.

During putative sleep, we observed that populations of LMAN neurons spontaneously

reactivated in bursting events. These events occurred predominantly during NREM sleep

and were associated with a transient increase in gamma band power. Across animals, the

population bursting events were preceded by a large negative deflection of the LFP, pointing

to the possibility that they were triggered by the ongoing large-scale global oscillatory activity

in LMAN.

The combination of large negative LFP deflection and population bursting activity in LMAN is

highly reminiscent of SWR activity observed during SWS and quiet rest in rodent

hippocampus (Buzsáki, 1986, 2015; O'Keefe, 1976).

The sharp wave and ripple LFP patterns that characterize mammalian SWRs are thought to

arise from the specific anatomy of the hippocampus. The sharp wave reflects massive

excitation of CA1 neurons by the pyramidal cells in CA3. This excitation synchronizes the

interneuron network and generates a ripple in the pyramidal layer (Buzsáki et al., 1983;

Girardeau & Zugaro, 2011; A. A. Liu et al., 2022). In rodents, the ripple oscillation is

characterized by a frequency in the range of 110-200 Hz, overlapping with typical gamma

band activity (Buzsáki, 2015; A. A. Liu et al., 2022).

Our observation of SWR-like activity in LMAN suggests that sharp waves coinciding with

gamma oscillations may not be unique to mammalian hippocampus. However, the function

of this non-hippocampal SWR-like activity remains unclear.

Hippocampal SWR activity has been typically considered a "biomarker" that accompanies

spatial navigation learning and episodic memory (Buzsáki, 2015); these learning paradigms

are fundamentally different from the auditory and motor learning that characterizes vocal

learning in songbirds. Furthermore, the birds used in our study sang mature crystallized

songs characterized by stable motif structure, suggesting that the developmental learning

period had already ended. Nevertheless, LMAN has been shown to have an important role in

the ongoing maintenance of adult song via N-methyl-D-aspartate (NMDA) receptor-

dependent transmission from LMAN to the pallial premotor nucleus RA (Brainard & Doupe,

2000; Solis et al., 2000; Warren et al., 2011). Bursting in LMAN influences RA and thereby

affects the spectral variability of song (Kao et al., 2005; Moorman et al., 2021; Tian et al.,

2023; Woolley & Kao, 2015). From these observations a potential role of avian SWR activity

emerges in the consolidation of experience in a broader sense (Tachibana et al., 2022).

It remains to be determined whether the LMAN synchronized bursts that we observed during

sleep represent true replay events related to, for example, daytime singing. "Replay"-like

reactivations have also been reported during sleep in RA (Dave & Margoliash, 2000;

Elmaleh et al., 2021, 2023). Given the widespread LFP dynamics that we observed across

LMAN, it would be important to investigate how synchronized bursting in LMAN and RA are

temporally related, and specifically whether the widespread LFP oscillations in LMAN

influence RA bursting during sleep.

Our investigation of the spontaneous network activity of the avian AFP has revealed two

major insights that help inform our understanding of how the AFP compares to the

mammalian corticostriatal circuit.

Similar to what has been reported in primates, we observed an overall desynchronization of

activity in BG compared to pallial areas during SWS sleep in songbirds. Specifically, pairwise

LFP coherence values within Area X were lower compared to LMAN for low frequency

ranges during SWS (Fig. S5: delta, theta, beta). This finding is consistent with primate

literature demonstrating significantly lower coherence within BG compared to cortical areas

during SWS (Mizrahi-Kliger et al., 2018).

In contrast, the presence of SWR-like activity in LMAN is unexpected and suggests that

multiple pallial neural architectures may give rise to SWR-like activity patterns.

During embryonic development in mammals, parts of the ventral pallium give rise to the

amygdalar complex (Garcia-Calero & Puelles, 2021). In birds and reptiles, however, the

ventral pallium gives rise to the sauropsid brain area known as the dorsal ventricular ridge

(DVR; (Puelles et al., 2000, 2017)). This large pallial region dorsal to the BG is unique to

reptiles and birds. In birds, the DVR has functionally been compared to the mammalian

neocortex due to its role in avian cognition (Güntürkün, 2005): high-level cognition in corvids

(Nieder et al., 2020), vocal learning in songbirds (Roberts et al., 2017; S. Yanagihara &

Yazaki-Sugiyama, 2016), and visual imprinting in chicken (Horn, 2004) are all behaviors that

critically involve the DVR. In birds, the DVR is subdivided into the mesopallium, nidopallium,

entopallium, and arcopallium (Briscoe & Ragsdale, 2018); LMAN is part of the nidopallium.

Importantly, similar SWR-like LFP patterns have also been recorded during sleep in the

reptilian DVR (Albeck et al., 2022; Libourel et al., 2018; Shein-Idelson et al., 2016). The fact

that SWR-like activity patterns arise from both reptilian and avian DVR during sleep

suggests that some version of the neural circuit that gives rise to mammalian hippocampal

SWRs may also be present in the avian DVR – despite the absence of clear anatomical

lamination in the DVR (although see (Stacho et al., 2020)). As highlighted above, however,

the function of this DVR-specific, SWR-like activity is still a mystery. Further work is required

to determine whether the SWR-like activity observed in the avian DVR is an

"epiphenomenon" related to the ongoing dynamics of the sleeping brain, or whether it has a

true function in learning and memory.

818   References

Albeck, N., Udi, D. I., Eyal, R., Shvartsman, A., & Shein-Idelson, M. (2022). Temperature-

robust rapid eye movement and slow wave sleep in the lizard Laudakia vulgaris.

Communications Biology, *5*(1), Article 1. https://doi.org/10.1038/s42003-022-04261-4

Ali, F., Otchy, T. M., Pehlevan, C., Fantana, A. L., Burak, Y., & Ölveczky, B. P. (2013). The

Basal Ganglia Is Necessary for Learning Spectral, but Not Temporal, Features of

Birdsong. *Neuron*, *80*(2), 494–506. https://doi.org/10.1016/j.neuron.2013.07.049

Andalman, A. S., & Fee, M. S. (2009). A basal ganglia-forebrain circuit in the songbird

biases motor output to avoid vocal errors. *Proceedings of the National Academy of*

Sciences of the United States of America, *106*(30), 12518–12523.

https://doi.org/10.1073/pnas.0903214106

Bastos, A. M., Vezoli, J., & Fries, P. (2015). Communication through coherence with inter-

areal delays. *Current Opinion in Neurobiology*, *31*, 173–180.

https://doi.org/10.1016/j.conb.2014.11.001

Bolhuis, J. J., Okanoya, K., & Scharff, C. (2010). Twitter evolution: Converging mechanisms

in birdsong and human speech. *Nature Reviews Neuroscience*, *11*(11), 747–759.

https://doi.org/10.1038/nrn2931

Bottjer, S. W., Miesner, E. A., & Arnold, A. P. (1984). Forebrain lesions disrupt development

but not maintenance of song in passerine birds. *Science (New York, N.Y.)*,

224(4651), 901–903. https://doi.org/10.1126/science.6719123

Brainard, M. S., & Doupe, A. J. (2000). Interruption of a basal ganglia-forebrain circuit

prevents plasticity of learned vocalizations. *Nature*, *404*(6779), 762–766.

https://doi.org/10.1038/35008083

Briscoe, S. D., & Ragsdale, C. W. (2018). Homology, neocortex, and the evolution of

developmental mechanisms. *Science*, *362*(6411), 190–193.

https://doi.org/10.1126/science.aau3711

Buzsáki, G. (1986). Hippocampal sharp waves: Their origin and significance. *Brain*

Research, *398*(2), 242–252. https://doi.org/10.1016/0006-8993(86)91483-6

Buzsáki, G. (2015). Hippocampal sharp wave-ripple: A cognitive biomarker for episodic

memory and planning. *Hippocampus*, *25*(10), 1073–1188.

https://doi.org/10.1002/hipo.22488

Buzsáki, G., & Draguhn, A. (2004). Neuronal Oscillations in Cortical Networks. *Science*,

304(5679), 1926–1929. https://doi.org/10.1126/science.1099745

Buzsáki, G., Leung, L. W. S., & Vanderwolf, C. H. (1983). Cellular bases of hippocampal

EEG in the behaving rat. *Brain Research*, *287*(2), 139–171.

Buzsáki, G., & Schomburg, E. W. (2015). What does gamma coherence tell us about inter-

regional neural communication? *Nature Neuroscience*, *18*(4), 484–489.

https://doi.org/10.1038/nn.3952

Buzsáki, G., & Wang, X.-J. (2012). Mechanisms of Gamma Oscillations. *Annual Review of*

Neuroscience, *35*, 203–225. https://doi.org/10.1146/annurev-neuro-062111-150444

Canavan, S. V., & Margoliash, D. (2020). Budgerigars have complex sleep structure similar

to that of mammals. *PLOS Biology*, *18*(11), e3000929.

https://doi.org/10.1371/journal.pbio.3000929

Cardin, J. A. (2016). Snapshots of the Brain in Action: Local Circuit Operations through the

Lens of γ Oscillations. *Journal of Neuroscience*, *36*(41), 10496–10504.

https://doi.org/10.1523/JNEUROSCI.1021-16.2016

Cardin, J. A., Carlén, M., Meletis, K., Knoblich, U., Zhang, F., Deisseroth, K., Tsai, L.-H., &

Moore, C. I. (2009). Driving fast-spiking cells induces gamma rhythm and controls

sensory responses. *Nature*, *459*(7247), 663–667.

https://doi.org/10.1038/nature08002

Chang, C.-Y., Hsu, S.-H., Pion-Tonachini, L., & Jung, T.-P. (2020). Evaluation of Artifact

Subspace Reconstruction for Automatic Artifact Components Removal in Multi-

Channel EEG Recordings. *IEEE Transactions on Biomedical Engineering*, *67*(4),

1114–1121. https://doi.org/10.1109/TBME.2019.2930186

Csicsvari, J., Hirase, H., Czurko, A., & Buzsáki, G. (1998). Reliability and state dependence

of pyramidal cell-interneuron synapses in the hippocampus: An ensemble approach

in the behaving rat. *Neuron*, *21*(1), 179–189. https://doi.org/10.1016/s0896-

6273(00)80525-5

Dave, A. S., & Margoliash, D. (2000). Song replay during sleep and computational rules for

sensorimotor vocal learning. *Science (New York, N.Y.)*, *290*(5492), 812–816.

https://doi.org/10.1126/science.290.5492.812

Duffy, A., Latimer, K. W., Goldberg, J. H., Fairhall, A. L., & Gadagkar, V. (2022). Dopamine

neurons evaluate natural fluctuations in performance quality. *Cell Reports*, *38*(13),

110574. https://doi.org/10.1016/j.celrep.2022.110574

Elmaleh, M., Kranz, D., Asensio, A. C., Moll, F. W., & Long, M. A. (2021). Sleep replay

reveals premotor circuit structure for a skilled behavior. *Neuron*, *109*(23), 3851-

3861.e4. https://doi.org/10.1016/j.neuron.2021.09.021

Elmaleh, M., Yang, Z., Ackert-Smith, L. A., & Long, M. A. (2023). Uncoordinated sleep replay

across hemispheres in the zebra finch. *Current Biology*, *33*(21), 4704-4712.e3.

https://doi.org/10.1016/j.cub.2023.09.005

Farries, M. A., Ding, L., & Perkel, D. J. (2005). Evidence for "direct" and "indirect" pathways

through the song system basal ganglia. *The Journal of Comparative Neurology*,

484(1), 93–104. https://doi.org/10.1002/cne.20464

Farries, M. A., & Perkel, D. J. (2002). A Telencephalic Nucleus Essential for Song Learning

Contains Neurons with Physiological Characteristics of Both Striatum and Globus

Pallidus. *Journal of Neuroscience*, *22*(9), 3776–3787.

https://doi.org/10.1523/jneurosci.22-09-03776.2002

Gale, S. D., & Perkel, D. J. (2010). Anatomy of a songbird basal ganglia circuit essential for

vocal learning and plasticity. *Journal of Chemical Neuroanatomy*, *39*(2), 124–131.

https://doi.org/10.1016/j.jchemneu.2009.07.003

Garcia-Calero, E., & Puelles, L. (2021). Development of the mouse anterior amygdalar radial

unit marked by Lhx9-expression. *Brain Structure & Function*, *226*(2), 575–600.

https://doi.org/10.1007/s00429-020-02201-8

Girardeau, G., & Zugaro, M. (2011). Hippocampal ripples and memory consolidation. *Current*

Opinion in Neurobiology, *21*(3), 452–459. https://doi.org/10.1016/j.conb.2011.02.005

Güntürkün, O. (2005). The avian 'prefrontal cortex' and cognition. *Current Opinion in*

Neurobiology, *15*(6), 686–693. https://doi.org/10.1016/j.conb.2005.10.003

Hahnloser, R. H. R., Kozhevnikov, A. A., & Fee, M. S. (2006). Sleep-related neural activity in

a premotor and a basal-ganglia pathway of the songbird. *Journal of Neurophysiology*,

96(2), 794–812. https://doi.org/10.1152/jn.01064.2005

Hessler, N. A., & Doupe, A. J. (1999a). Singing-related neural activity in a dorsal forebrain-

basal ganglia circuit of adult zebra finches. *The Journal of Neuroscience : The*

Official Journal of the Society for Neuroscience, *19*(23), 10461–10481.

Hessler, N. A., & Doupe, A. J. (1999b). Social context modulates singing-related neural

activity in the songbird forebrain. *Nature Neuroscience*, *2*(3), 209–211.

https://doi.org/10.1038/6306

Horn, G. (2004). Pathways of the past: The imprint of memory. *Nature Reviews*

Neuroscience, *5*(2), 108–120. https://doi.org/10.1038/nrn1324

Jun, J. J., Steinmetz, N. A., Siegle, J. H., Denman, D. J., Bauza, M., Barbarits, B., Lee, A. K.,

Anastassiou, C. A., Andrei, A., Aydın, Ç., Barbic, M., Blanche, T. J., Bonin, V., Couto,

J., Dutta, B., Gratiy, S. L., Gutnisky, D. A., Häusser, M., Karsh, B., … Harris, T. D.

(2017). Fully integrated silicon probes for high-density recording of neural activity.

Nature, *551*(7679), 232–236. https://doi.org/10.1038/nature24636

Kao, M. H., Doupe, A. J., & Brainard, M. S. (2005). Contributions of an avian basal ganglia–

forebrain circuit to real-time modulation of song. *Nature*, *433*(7026), 638–643.

https://doi.org/10.1038/nature03127

Kao, M. H., Wright, B. D., & Doupe, A. J. (2008). Neurons in a forebrain nucleus required for

vocal plasticity rapidly switch between precise firing and variable bursting depending

on social context. *The Journal of Neuroscience : The Official Journal of the Society*

for Neuroscience, *28*(49), 13232–13247. https://doi.org/10.1523/JNEUROSCI.2250-

08.2008

Kasdin, J., Duffy, A., Nadler, N., Raha, A., Fairhall, A. L., Stachenfeld, K. L., & Gadagkar, V.

(2025). Natural behaviour is learned through dopamine-mediated reinforcement.

Nature, *641*(8063), 699–706. https://doi.org/10.1038/s41586-025-08729-1

Kawai, R., Markman, T., Poddar, R., Ko, R., Fantana, A. L., Dhawale, A. K., Kampff, A. R., &

Ölveczky, B. P. (2015). Motor Cortex Is Required for Learning but Not for Executing a

Motor Skill. *Neuron*, *86*(3), 800–812. https://doi.org/10.1016/j.neuron.2015.03.024

Kobak, D., Pardo-Vazquez, J. L., Valente, M., Machens, C. K., & Renart, A. (2019). State-

dependent geometry of population activity in rat auditory cortex. *eLife*, *8*.

https://doi.org/10.7554/eLife.44526

Kojima, S., Kao, M. H., & Doupe, A. J. (2013). Task-related "cortical" bursting depends

critically on basal ganglia input and is linked to vocal plasticity. *Proceedings of the*

National Academy of Sciences, *110*(12), 4756–4761.

https://doi.org/10.1073/pnas.1216308110

Kollmorgen, S., Hahnloser, R. H. R., & Mante, V. (2020). Nearest neighbours reveal fast and

slow components of motor learning. *Nature*, *577*(7791), 526–530.

https://doi.org/10.1038/s41586-019-1892-x

Konishi, M. (1965). The role of auditory feedback in the control of vocalization in the white-

crowned sparrow. *Zeitschrift Für Tierpsychologie*, *22*(7), 770–783.

Konishi, M. (1970). Comparative neurophysiological studies of hearing and vocalizations in

songbirds. *Zeitschrift Für Vergleichende Physiologie*, *66*(3), 257–272.

https://doi.org/10.1007/BF00297829

Koralek, A. C., Costa, R. M., & Carmena, J. M. (2013). Temporally Precise Cell-Specific

Coherence Develops in Corticostriatal Networks during Learning. *Neuron*, *79*(5),

865–872. https://doi.org/10.1016/j.neuron.2013.06.047

Kothe, C. A., & Makeig, S. (2013). BCILAB: A platform for brain-computer interface

development. *Journal of Neural Engineering*, *10*(5), 056014.

https://doi.org/10.1088/1741-2560/10/5/056014

Lazarus, M., Chen, J. F., Urade, Y., & Huang, Z. L. (2013). Role of the basal ganglia in the

control of sleep and wakefulness. *Current Opinion in Neurobiology*, *23*(5), 780–785.

https://doi.org/10.1016/j.conb.2013.02.001

Lemke, S. M., Ramanathan, D. S., Darevksy, D., Egert, D., Berke, J. D., & Ganguly, K.

(2021). Coupling between motor cortex and striatum increases during sleep over

long-term skill learning. *eLife*, *10*, 1–25. https://doi.org/10.7554/eLife.64303

Libourel, P.-A., Barrillot, B., Arthaud, S., Massot, B., Morel, A. L., Beuf, O., Herrel, A., &

Luppi, P. H. (2018). Partial homologies between sleep states in lizards, mammals,

and birds suggest a complex evolution of sleep states in amniotes. *PLoS Biology*,

16(10), e2005982. https://doi.org/10.1371/journal.pbio.2005982

Liu, A. A., Henin, S., Abbaspoor, S., Bragin, A., Buffalo, E. A., Farrell, J. S., Foster, D. J.,

Frank, L. M., Gedankien, T., Gotman, J., Guidera, J. A., Hoffman, K. L., Jacobs, J.,

Kahana, M. J., Li, L., Liao, Z., Lin, J. J., Losonczy, A., Malach, R., … Buzsáki, G.

(2022). A consensus statement on detection of hippocampal sharp wave ripples and

differentiation from other fast oscillations. *Nature Communications*, *13*(1), Article 1.

https://doi.org/10.1038/s41467-022-33536-x

Liu, D., Li, W., Ma, C., Zheng, W., Yao, Y., Tso, C. F., Zhong, P., Chen, X., Song, J. H.,

Choi, W., Paik, S.-B., Han, H., & Dan, Y. (2020). A common hub for sleep and motor

control in the substantia nigra. *Science (New York, N.Y.)*, *367*(6476), 440–445.

https://doi.org/10.1126/science.aaz0956

Lorenz, C. (2024). *Song-related neuron populations in the zebra finch forebrain chronically*

recorded with a custom-designed Neuropixels device [Doctoral Thesis, ETH Zurich].

https://doi.org/10.3929/ethz-b-000690541

Low, P., Shank, S. S., Sejnowski, T. J., & Margoliash, D. (2008). Mammalian-like features of

sleep structure in zebra finches. *Proc Natl Acad Sci U S A*, *105*(26), 9081–9086.

https://doi.org/0703452105 [pii]10.1073/pnas.0703452105

McCasland, J. S., & Konishi, M. (1981). Interaction between auditory and motor activities in

an avian song control nucleus. *Proceedings of the National Academy of Sciences of*

the United States of America, *78*(12), 7815–7819.

Mizrahi-Kliger, A. D., Kaplan, A., Israel, Z., & Bergman, H. (2018). Desynchronization of slow

oscillations in the basal ganglia during natural sleep. *Proceedings of the National*

Academy of Sciences of the United States of America, *115*(18), E4274–E4283.

https://doi.org/10.1073/pnas.1720795115

Mizrahi-Kliger, A. D., Kaplan, A., Israel, Z., Deffains, M., & Bergman, H. (2020). Basal

ganglia beta oscillations during sleep underlie Parkinsonian insomnia. *Proceedings of*

the National Academy of Sciences, *117*(29), 17359–17368.

https://doi.org/10.1073/pnas.2001560117

Moorman, S., Ahn, J.-R., & Kao, M. H. (2021). Plasticity of stereotyped birdsong driven by

chronic manipulation of cortical-basal ganglia activity. *Current Biology*, *31*(12), 2619-

2632.e4. https://doi.org/10.1016/j.cub.2021.04.030

Murugan, M., Harward, S., Scharff, C., & Mooney, R. (2013). Diminished FoxP2 Levels

Affect Dopaminergic Modulation of Corticostriatal Signaling Important to Song

Variability. *Neuron*, *80*(6), 1464–1476. https://doi.org/10.1016/j.neuron.2013.09.021

Nieder, A., Wagener, L., & Rinnert, P. (2020). A neural correlate of sensory consciousness

in a corvid bird. *Science*, *369*(6511), 1–5. https://doi.org/10.1126/science.abb1447

Nixdorf-Bergweiler, B. E., Bischof, H.-J., Nixdorf-Bergweiler, B. E., & Bischof, H.-J. (2007). *A*

Stereotaxic Atlas Of The Brain Of The Zebra Finch, Taeniopygia Guttata. National

Center for Biotechnology Information (US).

O'Keefe, J. (1976). Place units in the hippocampus of the freely moving rat. *Experimental*

Neurology, *51*(1), 78–109.

Olveczky, B. P., Andalman, A. S., & Fee, M. S. (2005). Vocal experimentation in the juvenile

songbird requires a basal ganglia circuit. *PLoS Biology*, *3*(5), e153.

https://doi.org/10.1371/journal.pbio.0030153

Palmer, S. E., Wright, B. D., Doupe, A. J., & Kao, M. H. (2021). Variable but not random:

Temporal pattern coding in a songbird brain area necessary for song modification.

Journal of Neurophysiology, *125*(2), 540–555. https://doi.org/10.1152/jn.00034.2019

Person, A. L., Gale, S. D., Farries, M. A., & Perkel, D. J. (2008). Organization of the songbird

basal ganglia, including area X. *Journal of Comparative Neurology*, *508*(5), 840–866.

https://doi.org/10.1002/cne.21699

Pfenning, A. R., Hara, E., Whitney, O., Rivas, M. V., Wang, R., Roulhac, P. L., Howard, J. T.,

Wirthlin, M., Lovell, P. V., Ganapathy, G., Mouncastle, J., Moseley, M. A., Thompson,

J. W., Soderblom, E. J., Iriki, A., Kato, M., Gilbert, M. T. P., Zhang, G., Bakken, T., …

Jarvis, E. D. (2014). Convergent transcriptional specializations in the brains of

humans and song-learning birds. *Science*, *346*(6215).

https://doi.org/10.1126/science.1256846

Prerau, M. J., Brown, R. E., Bianchi, M. T., Ellenbogen, J. M., & Purdon, P. L. (2017). Sleep

Neurophysiological Dynamics Through the Lens of Multitaper Spectral Analysis.

Physiology (Bethesda, Md.), *32*(1), 60–92.

https://doi.org/10.1152/physiol.00062.2015

Puelles, L., Kuwana, E., Puelles, E., Bulfone, A., Shimamura, K., Keleher, J., Smiga, S., &

Rubenstein, J. L. (2000). Pallial and subpallial derivatives in the embryonic chick and

mouse telencephalon, traced by the expression of the genes Dlx-2, Emx-1, Nkx-2.1,

Pax-6, and Tbr-1. *The Journal of Comparative Neurology*, *424*(3), 409–438.

https://doi.org/10.1002/1096-9861(20000828)424:3<409::aid-cne3>3.0.co;2-7

Puelles, L., Sandoval, J. E., Ayad, A., del Corral, R., Alonso, A., Ferran, J. L., & Martínez-de-

la-Torre, M. (2017). The Pallium in Reptiles and Birds in the Light of the Updated

Tetrapartite Pallium Model. In *Evolution of Nervous Systems* (Vol. 2).

https://doi.org/10.1016/b978-0-12-804042-3.00014-2

Qi, J., Schreiner, D. C., Martinez, M., Pearson, J., & Mooney, R. (2025). Dual

neuromodulatory dynamics underlie birdsong learning. *Nature*, *641*(8063), 690–698.

https://doi.org/10.1038/s41586-025-08694-9

Roberts, T. F., Hisey, E., Tanaka, M., Kearney, M. G., Chattree, G., Yang, C. F., Shah, N.

M., & Mooney, R. (2017). Identification of a motor-to-auditory pathway important for

vocal learning. *Nature Neuroscience*, *20*(7), 978–986.

https://doi.org/10.1038/nn.4563

Santos, F. J., Oliveira, R. F., Jin, X., & Costa, R. M. (2015). Corticostriatal dynamics encode

the refinement of specific behavioral variability during skill learning. *eLife*, *4*, e09423.

https://doi.org/10.7554/eLife.09423

Scharff, C., & Nottebohm, F. N. (1991). A comparative study of the behavioral deficits

following lesions of various parts of the zebra finch song system: Implications for

vocal learning. *The Journal of Neuroscience : The Official Journal of the Society for*

Neuroscience, *11*(9), 2896–2913.

Schneider, M., Broggini, A. C., Dann, B., Tzanou, A., Uran, C., Sheshadri, S., Scherberger,

H., & Vinck, M. (2021). A mechanism for inter-areal coherence through

communication based on connectivity and oscillatory power. *Neuron*, *109*(24), 4050-

4067.e12. https://doi.org/10.1016/j.neuron.2021.09.037

Shein-Idelson, M., Ondracek, J. M., Liaw, H., Reiter, S., & Laurent, G. (2016). Slow waves,

sharp waves, ripples, and REM in sleeping dragons. *Science (New York, N.Y.)*,

352(6285), 590–595. https://doi.org/10.1126/science.aaf3621

Solis, M. M., Brainard, M. S., Hessler, N. A., & Doupe, A. J. (2000). Song selectivity and

sensorimotor signals in vocal learning and production. *Proceedings of the National*

Academy of Sciences of the United States of America, *97*(22), 11836–11842.

https://doi.org/10.1073/pnas.97.22.11836

Stacho, M., Herold, C., Rook, N., Wagner, H., Axer, M., Amunts, K., & Gütürkün, O. (2020).

A cortex-like canonical circuit in the avian forebrain. *Science*, *369*(6511).

https://doi.org/10.1126/science.abc5534

Tachibana, R. O., Lee, D., Kai, K., & Kojima, S. (2022). Performance-Dependent

Consolidation of Learned Vocal Changes in Adult Songbirds. *Journal of*

Neuroscience, *42*(10), 1974–1986. https://doi.org/10.1523/JNEUROSCI.1942-

21.2021

Tian, L. Y., Warren, T. L., Mehaffey, W. H., & Brainard, M. S. (2023). Dynamic top-down

biasing implements rapid adaptive changes to individual movements. *eLife*, *12*,

e83223. https://doi.org/10.7554/eLife.83223

Tumer, E. C., & Brainard, M. S. (2007). Performance variability enables adaptive plasticity of

"crystallized" adult birdsong. *Nature*, *450*(7173), 1240–1244.

https://doi.org/10.1038/nature06390

Vates, G. E., & Nottebohm, F. (1995). Feedback circuitry within a song-learning pathway.

Proceedings of the National Academy of Sciences of the United States of America,

92(11), 5139–5143.

Vates, G. E., Vicario, D. S., & Nottebohm, F. (1997). Reafferent thalamo- "cortical" loops in

the song system of oscine songbirds. *The Journal of Comparative Neurology*, *380*(2),

275–290.

Warren, T. L., Tumer, E. C., Charlesworth, J. D., & Brainard, M. S. (2011). Mechanisms and

time course of vocal learning and consolidation in the adult songbird. *Journal of

Neurophysiology*, *106*(4), 1806–1821. https://doi.org/10.1152/jn.00311.2011

Woolley, S. C., & Kao, M. H. (2015). Variability in action: Contributions of a songbird cortical-

basal ganglia circuit to vocal motor learning and control. *Neuroscience*, *296*, 39–47.

https://doi.org/10.1016/j.neuroscience.2014.10.010

Woolley, S. C., Rajan, R., Joshua, M., & Doupe, A. J. (2014). Emergence of context-

dependent variability across a basal ganglia network. *Neuron*, *82*(1), 208–223.

https://doi.org/10.1016/j.neuron.2014.01.039

Yanagihara, S. Y., & Hessler, N. A. (2012). Phasic basal ganglia activity associated with

high-gamma oscillation during sleep in a songbird. *Journal of Neurophysiology*,

107(1), 424–432. https://doi.org/10.1152/jn.00790.2011

Yanagihara, S., & Yazaki-Sugiyama, Y. (2016). Auditory experience-dependent cortical

circuit shaping for memory formation in bird song learning. *Nature Communications*,

7(May 2015), 11946. https://doi.org/10.1038/ncomms11946

Yeganegi, H., Luksch, H., & Ondracek, J. M. (2019). Hippocampal-like network dynamics

underlie avian sharp wave-ripples. *bioRxiv*, 825075. https://doi.org/10.1101/825075

Yeganegi, H., & Ondracek, J. M. (2023). Multi-channel recordings reveal age-related

differences in the sleep of juvenile and adult zebra finches. *Scientific Reports*, *13*(1),

Article 1. https://doi.org/10.1038/s41598-023-35160-1

Yeganegi, H., & Ondracek, J. M. (2025). Local sleep in songbirds: Different simultaneous

sleep states across the avian pallium. *Journal of Sleep Research*, *34*(3), e14344.

https://doi.org/10.1111/jsr.14344

Zai, A. T., Cavé-Lopez, S., Rolland, M., Giret, N., & Hahnloser, R. H. R. (2020). Sensory

substitution reveals a manipulation bias. *Nature Communications*, *11*(1), 5940.

https://doi.org/10.1038/s41467-020-19686-w

Zann, R. A. (1996). *The Zebra Finch: A Synthesis of Field and Laboratory Studies*. Oxford

University Press.

1106

1107

Figures and Table Legends

1108

1109

1110

Figure 1: Pallial-basal ganglia dynamics during sleep and awake behavior in a

songbird. (A) Schematic depicts the basal ganglia-thalamocortical circuit (blue arrows).

Black arrows indicate other important connections of the song system. HVC, proper name;

RA, robust nucleus of the arcopallium; LMAN, lateral magnocellular nucleus of the

nidopallium; Area X, proper name; DLM, medial dorsolateral nucleus of the thalamus. **(B)**

Image of a male zebra finch implanted with a Neuropixels1.0 probe. **(C)** Histological brain

slice depicting the Neuropixels electrode track (red DiI fluorescence). Area X is indicated

with a dotted black line. St, striatum; d, dorsal; a, anterior; v, ventral; p, posterior; for other

abbreviations, see (A). Histology from ZF-2, cut at room temperature on a vibratome. **(D)** 7 s

segment of the bandpass-filtered (0.5-250Hz) local field potentials (LFP) recorded during

putative sleep in subadult ZF-1 and corresponding microphone signal (top, arbitrary units

(AU)). Black arrowheads (LFP, top) indicate large amplitude negative LFP deflections.

Recording sites were located in the pallium (gray shading) and basal ganglia (green

shading) and are organized according to depth. Dark gray box (LMAN, right side) indicates

LFPs that correspond to LMAN; dark green box (Area X, right side) indicates LFPs that

correspond to Area X. Below, corresponding spiking activity for identified LMAN or Area X

neurons. Firing rate is indicated with the color scale, and neurons are sorted according to

firing rate. Note how neural activity transitions from sparse and synchronous firing aligned to

the LFP troughs to asynchronous firing at 912-914 s. **(E)** 7 s segment of the bandpass-

filtered LFP recorded during active daytime behavior in ZF-1. Figure conventions same as in

(D). Note the calling and singing behavior recorded on the microphone signal (top) and the

corresponding activity in LMAN and Area X. Note also the absence of large amplitude

negative deflections in the LFP as observed during putative sleep.

1134

Figure 2: Identification of sleep stages and representative neural dynamics. (A) Ratio

of δ/γ power for consecutive depths along the Neuropixels shank. Each trace is separated by

200 um. Black trace indicates the δ/γ trace calculated for a centrally located pallial (LMAN)

LFP. Numbers and arrows indicate low (1), middle (2), and high (3) δ/γ values. δ/γ values

calculated for a 3-s LFP window, as depicted in (B). Data from sub-adult ZF-1 (pink

shading). **(B)** 3-second segments of local field potential (LFP) activity from pallial and BG

sites for the low, middle, and high δ/γ values indicated in (A). Each trace is separated by 80

um. Note the large amplitude slow wave dynamics present for high δ/γ values indicated in

(3). Linear color scale for visualization purposes only. Data from sub-adult ZF-1. **(C)** Box

plots indicate average δ/γ values calculated in 3-s windows from a centrally located LMAN

channel for the duration of the sleep phase per bird. The left and right edges of the box

represent the 25th and 75th percentile, respectively, and the middle dot represents the

median. Whiskers extend to the most extreme datapoints not considered outliers, and

datapoints beyond this range are indicated as dots. Sub-adult animals are indicated with

pink shading of their names. **(D)** δ/γ values for an entire recording duration for an example

bird (sub-adult ZF-5) plotted with the identified sleep stages as colored bars below. SWS,

slow wave sleep; IS, intermediate sleep; REM, rapid eye-movement sleep; Nc, not

categorized. Dotted black box indicated with the black arrow is enlarged in (E). **(E)** 50 s trace

of raw LFP from the LMAN channel used for sleep staging (see Methods for details). Color-

coded bars indicate classified sleep stages. Note how SWS segments correspond to a large

amplitude LFP signal and REM segments correspond to a low amplitude LFP signal. **(F)**

Histogram of δ/γ values grouped according to identified sleep states together with the

proportions of sleep stages depicted in a pie chart for the bird in (D). Note how SWS

segments correspond to high δ/γ values and REM segments correspond to low δ/γ values.

(G, H) Plots show that sleep state clusters are segregated in 3-D space according to the

features used for classification, e.g., mean LFP magnitude, the delta (δ) and gamma (γ)

power, δ/γ values. Each dot represents a 1-s snippet of sleeping data (resolution of the

automated sleep staging procedure). Same bird as in (D-F).

1163

Figure 3: Sleep-related modulation of low-frequency power is higher in pallium compared to the striatum (A) Comparison of microphone signals recorded during the light on (awake) period and the light off (sleep) period. Top traces (black) indicate the raw microphone signal; scale is the same across recordings (arbitrary units, AU). Bottom traces (gray) indicate the root mean square (RMS) of the microphone signal. Black dotted line indicates the manually set threshold, and threshold crossings indicate sound detections (red lines). The overall percent of data for which sounds were detected is stated (top). Recording from sub-adult ZF-1. Microphone data from other birds can be found in Fig. S1. **(B)** The color-coded power spectrum density (PSD) of the LFP signal for each electrode is displayed, organized by depth (dorsal to ventral), for both the awake state (left) and sleep state (right) for sub-adult ZF-1. Warm color indicates high power; cool colors indicate low power. The electrode depths corresponding to LMAN (gray box, dashed line) and Area X (green box, dashed line) are indicated. The analysis is based on the average PSD derived from 10 randomly selected 10-sec windows. **(C)** Power distributions across electrode sites provide a detailed comparison between awake (thin lines) and sleep (thick lines) for different frequency bands. LMAN exhibits higher gamma band power (90-145 Hz) during awake states compared to sleep states (LMAN; black arrowheads). In contrast LMAN exhibits higher

power in low frequency bands (1-30 Hz) during sleep compared to awake states (LMAN;

black arrowheads). A difference in sleep and awake states in Area X is largely absent, with

the exception of the theta band power (4-12 Hz), which showed a slight increase in two birds

during sleep (Area X; black arrowhead). **(D)** The marginal distribution of power across

depths is displayed for awake and sleep states for each bird. Note that the distances in the

legend refer to the distance from the probe tip, (i.e., 3.6 mm from the probe tip). The black

line indicates depth associated with LMAN (i.e., 2.6 mm); note how it is strongly modulated

by sleep. Gray lines indicate other depths associated with the pallium. The dark green line

indicates the depth associated with Area X; light green lines indicate other depths associated

with the BG. Dotted line indicates 0 dB. Spectral data for other birds are depicted in Fig. S4.

1192

1193 **Figure 4: LFP coherence in the pallial-striatal circuit is modulated by behavioral**

1194 **states. (A)** Heat maps represent a color-coded representation of the average pairwise

1195 coherence calculated for all passbands for all SWS segments (top) and REM sleep

1196 segments (bottom). Warm colors indicate high coherence values, and cool colors indicate

1197 low coherence values. The diagonal white line indicates symmetry. The white triangles along

1198 the line of symmetry indicate the pallial electrode sites that were included in the LMAN-

1199 LMAN coherency analysis (corresponding to the gray LMAN shading; small black triangle,

1200 right side) or the BG electrode sites that were included in the Area X-Area X coherence

1201 analysis (green Area X shading; large dark green triangle, right side). The white rectangle

1202 (lower right quadrant) indicates the sites that were included in the LMAN-Area X coherency

1203 analysis (blue rectangle, right side). Data from sub-adult ZF-5. **(B)** Scatter plots (top row)

1204 depict the average REM sleep versus SWS coherence value for each electrode pair plotted

1205 for different frequency bands in each panel for sub-adult ZF-5. LMAN-LMAN comparisons

1206 (gray dots); Area X- Area X comparisons (green dots) and LMAN-Area X comparisons (blue

1207 dots). Black lines at 0.5 for visualization purposes only. Bottom row depicts the mean

1208 coherence values for all 7 birds (symbols); colors indicate the coherence comparisons (e.g.,

1209 LMAN-LMAN in dark gray). Subadult birds are indicated with pink shading of their names.

1210 **(C)** Comparison between REM and SWS coherences for different passbands. Box plots

1211 indicate the median coherency values across different frequency bands for within-LMAN

1212 comparisons (left panel), within-Area X comparisons (middle panel) and LMAN-Area X

1213 comparisons (right panels). Blue boxes indicate SWS, red boxes indicate REM sleep. The

1214 box bottom and top edges represent the 25th and 75th percentile, respectively, and the

1215 middle line represents the median. Whiskers extend to the most extreme datapoints not

1216 considered outliers, and datapoints beyond this range are indicated as circles. Asterisks

1217 indicate significance (Wilcoxon sign rank test; ***$p \leq 0.001$, exact p values are stated in Table

1218 2). **(D)** Comparison between awake and sleep coherences for different passbands for 3

1219 birds. SWS (blue box); REM sleep (red box); Light on non-active (gray box); Light on active

1220 (white box). Delta and theta band coherences were significantly lower during the awake

1221 states compared to the sleep states for all channel comparisons. Figure conventions same

1222 as in (C). Asterisks indicate significance (Friedman test for multiple comparisons;

1223 ***$p \leq 0.001$, exact p values are stated in Table 3). The non-significant comparisons from the

 post-hoc Nemeyni test are marked in the figure (n.s). All other pairwise comparisons were

1225   significant with the post-hoc test.

1226

Figure 5: Spiking variability is significantly higher for LMAN neurons compared to

Area X neurons. (A) Bar plots summarize CV values for 7 finches (black bars, LMAN

population; green bar, Area X population). CV values were higher for LMAN populations

compared to Area X populations. Sub-adult birds are indicated by pink shading of their

names. **(B)** δ/γ values (top, blue line) were highly correlated with CV values (bottom) for both

LMAN and Area X populations. δ/γ values and CV values computed in 10 s windows. **(C)**

The slope of regression fits between CV and δ/γ was consistently higher in LMAN compared

to Area X across birds. Filled symbols indicate significant correlations (p < 0.01, Table 5).

(D) Scatter plot indicates the CV versus δ/γ values for ZF-1. LMAN, black dots; Area X,

green dots. Black and green arrows indicate the data points that correspond to the 10 s

snippet of data plotted in (D) and (E). The marginal distributions for the CV values are

plotted (right) for LMAN (black line) and Area X neurons (green line). **(E)** Spiking raster plot

for 33 LMAN neurons for an example time window with low CV value (left plot) and a high

CV value (right box). Each tick corresponds to a spike and neurons are sorted by their

average spiking activity from high (top) to low (bottom). Corresponding 10 s LFP trace is

plotted at the bottom of the raster plot. Vertical bar is 500 uV. Data correspond to black

arrows in (D). **(F)** Spiking raster plot for 27 Area X neurons for a low CV value (left plot) and

a high CV value (right box). Figure convention same as in (E). Data correspond to green

arrows in (D). Vertical bar is 500 uV.

1246

Figure 6: Synchronized bursting in LMAN is associated with a negative deflection in

the LFP (A) Two examples of synchronized bursting activity in LMAN from ZF-2. Top panel

depicts high-pass filtered (HPF) electrode signals from eight electrodes in LMAN. Middle

panel depicts the detection of synchronized bursting (black line) based on the smoothed

fraction of bursting units in a 50 ms window exceeding the 99.5[th] percentile threshold.

Bottom panel depicts the raw LFP (blue, same electrode as in top panel). Area indicated by

gray shading is expanded in (B). **(B)** Zoom-in on the highlighted gray window in (A). Top

panel depicts the instantaneous firing rate (IFR) of all LMAN units (n = 32), color-coded and

clipped at 100 Hz. The panel below depicts the occurrence of bursts in black for each LMAN

unit. Lower panels same as in (A). **(C)** Upper panel depicts 195 bursting-aligned LFP

segments detected from channel 236, stacked and color-coded. Events are sorted by the

total fraction of bursting units. Lower panel indicates the average bursting-aligned LFP (blue)

and the same statistics for randomly selected LFP segments (black, gray shading is +/- SD).

(D) Two raw LFP snippets (top panel) and their corresponding scalograms (bottom panel).

The left example shows a bursting aligned LFP. The right depicts random control (right). **(E)**

The average bursting aligned-LFP scalogram (left) and its difference to the average random

control scalogram (right). Note how the power density for the synchronized burst is highly

elevated at low frequencies. Sb, synchronized burst. **(F)** LFPs show consistent negative dip

during a synchronized burst. Each line indicates the bursting-aligned LFP normalized to the

random control (z-norm ± SEM) for each bird. **(G)** The average burst-aligned LFP ± SEM

across different states for ZF-2. Note that the LFP dip remains visible during light on non-

active periods (short-dashed line), though it is reduced in magnitude, and becomes further

diminished during light on active periods (long-dashed line). **(H)** Comparison of LFP dips

across all birds with available awake data shows that the strongest deflections occur during

light-off periods. No bursting events were detected for ZF-1 during light on active periods.

See Fig. S6 for more details.

1273

Figure 7: Synchronized bursts occur during NREM sleep and are associated with

higher γ activity. (A) Bar plots indicate median number of bursts associated with each

sleep state normalized by the overall sleep state duration in seconds across birds (see Fig.

2F, G). With the exception of ZF-4, most birds showed more bursts detected during NREM

states compared to REM sleep states. NREM states included both SWS and IS classified

states. Symbols indicate median data for each bird. Sub-adults are indicated with pink

shading of their names. **(B)** Bar plots indicate median δ/γ value calculated for 500 ms

segments of LFP that were centered on a synchronized burst detection (Bursts, grey bar)

compared to the mean value of 500 ms control segments of LFP that were temporally shifted

away from the burst detection by a random offset (Control, white bar; see Methods for

details). Symbols indicate median data for each bird. Note how δ/γ values are on average

higher for the control LFP segments. **(C)** Same convention as in (B) for δ values. Note how δ

values are higher for the burst LFP segments compared to the control LFP segments. **(D)**

Same convention as in (B) and (C) for γ values. Note the higher γ values during the burst

LFP segments compared to the control LFP segments. **(E)** Spectrograms display the

average peri-burst time window surrounding a burst event (top), a control event (middle),

and the difference in power between the burst event and the control event (bottom) for ZF-2.

Note the increase in power surrounding the burst event that extends up into the high γ

range. **(F)** Panels display the normalized PSD for different frequency ranges: δ (1-4 Hz, top),

low γ (30-90 Hz, middle), and high γ (90-145 Hz, bottom plot). Normalized PSD indicates the

difference between the burst events and the control events divided by the standard deviation

of the control events. Different line colors indicate different birds. Note how data for all birds

fall above the zero (dotted line) indicating higher values during the burst event compared to

control events.

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

ZF	Animal ID	Age (dph)	Rec location	Start of recording	Data available	# LMAN units	# Area X units	Histology	Spike-sorting***
1	g4r4	84	Paris	03:58 pm*	LFP+SU	30	26	Vibratome	KS 1.0 +Phy
2	r15v15	125	Paris	8:42 pm	LFP+SU	32	6	Vibratome	KS 1.0 + Phy
3	j8v8	124	Paris	05:18 am	LFP+SU	40	19	Missing**	KS 1.0 + Phy
4	Redred	177	Bordeaux	05:30 am	LFP+SU	28	7	Cryostat	KS 2.5 + SI
5	r5n5	85	Paris	5:21 pm*	LFP+SU	6	3	Vibratome	KS 1.0 + Phy
6	black/cyan	232	Bordeaux	04:17 am	LFP	-	-	Cryostat	-
7	no_label	76	Bordeaux	00:28 am	LFP	-	-	Missing**	-

1310

1311 **Table 1: Information related to experimental animals**

1312 Age refers to the age at the recording time in days post hatch (dph).

1313

1314 For all birds recorded in Bordeaux, data were recorded continuously throughout the night,
1315 and a 1-hour time period was selected for the analysis in this paper. The times indicated in
1316 the "Start of Recording" column indicates the start of the 1-hour recording time for those
1317 birds.

1318

1319 * For some Paris birds, the dark period (light off) time was initiated earlier than normal and
1320 recording sessions started in the late afternoon (in the dark).

1321

1322 LFP, local field potential
1323 SU, spiking unit (including single and multi-unit activity from high-pass filtered voltage traces)

1324

1325 ** These animals died during the Covid laboratory lock-down in 2020 and it was not possible
1326 to retrieve the brains in time to perform histology. Electrode reconstruction was then based
1327 on response mapping of the brain areas that was performed during the probe implantation,
1328 as well as by visualizing the sorted spiking data for LMAN and Area X, which was
1329 distinguishable on the basis of spike amplitude and rate.

1330

***Different spikesorting pipelines were setup in Paris and Bordeaux and largely reflected the
different use of Matlab versus Python. When the data were first acquired in Paris, Kilosort
1.0 (KS 1.0) was the only version available; the data were sorted with Kilosort 1.0 and
curated with Phy. Later, as more data were acquired in Bordeaux, the analysis pipelines
there incorporated the newer version of Kilosort 2.5 (KS 2.5) and reflected E.G.C.Z.'s work
with the Spikeinterface (SI) development team.

Brain Area	Frequency range	Median coherence across channels during SWS (Min, Max values)	Median coherence across channels during REM (Min, Max values)	Wilcoxon signed rank test	
				z value	*p* value
LMAN (N=147)	Delta	0.965 (0.861, 0.996)	0.872 (0.633, 0.989)	10.52	7.15E-26
	Theta	0.968 (0.876, 0.995)	0.893 (0.754, 0.988)	10.52	7.15E-26
	Beta	0.821 (0.521, 0.969)	0.680 (0.311, 0.963)	10.52	7.15E-26
	Low gamma	0.500 (0.119, 0.854)	0.484 (0.106, 0.862)	6.244	4.26E-10
	High gamma	0.397 (0.074, 0.769)	0.451 (0.063, 0.806)	-5.67	1.43E-8
Area X (N=2275)	Delta	0.606 (0.043, 0.946)	0.409 (0.006, 0.888)	39.34	0.00E+00
	Theta	0.671 (0.131, 0.977)	0.561 (0.101, 0.939)	41.00	0.00E+00
	Beta	0.394 (0.062, 0.928)	0.304 (0.046, 0.912)	41.28	0.00E+00
	Low gamma	0.189 (0.024, 0.770)	0.178 (0.023, 0.703)	31.30	5.44E-215
	High gamma	0.146 (0.021, 0.691)	0.139 (0.020, 0.648)	24.77	1.69E-135
LMAN – Area X (N=1274)	Delta	0.169 (0.003, 0.820)	0.163 (0.011, 0.782)	10.50	9.05E-26
	Theta	0.244 (0.011, 0.818)	0.234 (0.008, 0.749)	4.79	1.70E-6
	Beta	0.107 (0.013, 0.464)	0.117 (0.015, 0.480)	-8.37	5.71E-17
	Low gamma	0.069 (0.019, 0.153)	0.077 (0.020, 0.172)	-14.35	1.06E-46
	High gamma	0.055 (0.014, 0.171)	0.040 (0.010, 0.139)	30.84	8.16E0-209

1337

1338 **Table 2: Average coherence values for all birds (n = 7) used for analysis in Figure 4 C.**
1339 Frequency ranges: Delta (1 – 4 Hz), Theta (4 – 12 Hz), Beta (12 – 30 Hz), Low gamma (30 –
1340 90 Hz), High gamma (90 – 140 Hz). N corresponds to the number of LFP pairs.
1341

Brain Area	Frequency range	Median coherence across channels during SWS (Min, Max values)	Median coherence across channels during REM (Min, Max values)	Median coherence across channels Light on non-active (Min, Max values)	Median coherence across channels Light on active (Min, Max values)	Friedman test (k=4, df=3) Q	p	Post-hoc Nemenyi test (k=3) SWS vs REM	SWS vs light on non-active	SWS vs light on active	REM vs light on non active	REM vs light on active	light on active vs non active
LMAN (N=63)	Delta	0.975 (0.908, 0.995)	0.882 (0.633, 0.975)	0.322 (0.176, 0.883)	0.362 (0.209, 0.886)	171.02	1.66E-37	.001	.001	.001	.001	.001	.001
	Theta	0.977 (0.922, 0.995)	0.898 (0.769, 0.975)	0.224 (0.039, 0.826)	0.166 (0.032, 0.818)	171.48	6.11E-37	.001	.001	.001	.001	.001	.001
	Beta	0.836 (0.656, 0.956)	0.622 (0.399, 0.867)	0.831 (0.422, 0.974)	0.689 (0.232, 0.963)	90.77	1.50E-19	.001	.001	.001	.001	.0019	.001
	Low gamma	0.529 (0.227, 0.836)	0.466 (0.149, 0.806)	0.149 (0.024, 0.457)	0.122 (0.016, 0.544)	143.02	8.44E-31	.001	.001	.001	.001	.001	.001
	High gamma	0.402 (0.124, 0.753)	0.446 (0.092, 0.792)	0.298 (0.084, 0.807)	0.306 (0.089, 0.798)	7.51	0.0572	No post-hoc test					
Area X (N=975)	Delta	0.678 (0.141, 0.948)	0.469 (0.112, 0.899)	0.266 (0.103, 0.915)	0.311 (0.138, 0.914)	1449.20	0.00E-00	.001	.001	.001	.001	.001	.001
	Theta	0.719 (0.246, 0.977)	0.626 (0.226, 0.928)	0.207 (0.029, 0.905)	0.179 (0.013, 0.896)	2028.41	0.00E-00	.001	.001	.001	.001	.001	.001
	Beta	0.379 (0.067, 0.878)	0.277 (0.050, 0.767)	0.618 (0.076, 0.976)	0.331 (0.066, 0.972)	1489.37	0.00E-00	.001	.001	.001	.001	.001	.001
	Low gamma	0.157 (0.024, 0.680)	0.145 (0.023, 0.669)	0.158 (0.017, 0.882)	0.114 (0.011, 0.879)	541.32	5.29E-117	.001	.001	.001	.001	.001	.001
	High gamma	0.123 (0.021, 0.611)	0.114 (0.019, 0.593)	0.172 (0.022, 0.823)	0.154 (0.021, 0.827)	688.60	6.21E-149	.001	.001	.001	.001	.001	.001
LMAN – Area X (N=546)	Delta	0.309 (0.087, 0.819)	0.310 (0.079, 0.782)	0.272 (0.119, 0.891)	0.311 (0.153, 0.894)	238.38	2.13E-51	.001	.001	.001	.001	.001	.001
	Theta	0.341 (0.128, 0.818)	0.368 (0.091, 0.749)	0.133 (0.033, 0.916)	0.105 (0.017, 0.858)	367.37	2.58E-79	.001	.001	.001	.001	.001	.001
	Beta	0.154 (0.035, 0.438)	0.163 (0.030, 0.394)	0.307 (0.043, 0.975)	0.196 (0.056, 0.969)	873.38	5.27E-189	.001	.001	.001	.001	.001	.001
	Low gamma	0.056 (0.020, 0.146)	0.068 (0.020, 0.130)	0.153 (0.021, 0.894)	0.116 (0.015, 0.893)	651.25	7.82E-141	.001	.001	.001	.001	.001	.001
	High gamma	0.052 (0.019, 0.100)	0.036 (0.009, 0.079)	0.056 (0.015, 0.819)	0.053 (0.017, 0.817)	496.50	2.73E-107	.001	.900	.001	.001	.001	.001

Table 3: Average coherence values for all birds (n = 3) used for analysis in Figure 4D.
Frequency ranges: Delta (1 – 4 Hz), Theta (4 – 12 Hz), Beta (12 – 30 Hz), Low gamma (30 – 90 Hz), High gamma (90 – 140 Hz). N corresponds to the number of LFP pairs.

Bird	N	LMAN CV median (min, max)	Area X CV median (min, max)	Wilcoxon ranksum test z Value	p Value
ZF-1	532	0.73 (0.33-2.68)	0.28 (0.14-0.37)	28.18	1.08E-174
ZF-2	315	2.14 (0.32-5.16)	0.62 (0.39-0.99)	18.13	1.74E-73
ZF-3	990	0.80 (0.39 -6.42)	0.27 (0.16-0.54)	38.12	6.71E-318
ZF-4	318	1.44 (0.44-5.44)	0.67 (0.43-0.98)	15.79	3.94E-56
ZF-5	400	1.62 (0.88-5.08)	1.04 (0.86-1.22)	18.36	3.00E-75
Population median	5	1.44	0.62		

Table 4: Median CV values were significantly higher for LMAN neurons compared to Area X neurons.

Median values and range. Population statistics are computed across median CVs. p value is the result of a Wilcoxon rank sum test.

1354

Bird		LMAN: CV versus δ/γ			Area X: CV versus δ/γ	
	N	R	p		R	p
ZF-1	532	0.67	7.58E-71		0.73	2.45E-88
ZF-2	315	0.43	2.08E-15		0.05	0.42
ZF-3	990	0.32	1.17E-25		0.24	2.06E-14
ZF-4	318	0.16	0.0034		-0.28	3.65E-07
ZF-5	400	0.71	5.40E-63		-0.09	0.085

1355

1356 **Table 5: LMAN CV values were significantly and positively correlated with δ/γ values**

1357 **for 5/5 zebra finches.** Correlations were more heterogeneous for Area X. R, Pearson

1358 correlation coefficient.