Cholinergic signaling differentially regulates song premotor circuits to stabilize songs in songbirds

Ning Xu,[1,3] Yutao Zhang,[1,3] Yalun Sun,[1,3] Xueqing Song,[1] Yangyang Cao,[1] Xinqi Yang,[1] Songhua Wang,[1,*] and Wei Meng[1,2,*]

[1] Jiangxi Province Key Laboratory of Organic Functional Molecules; Institute of Organic Chemistry, Jiangxi Science and Technology Normal University, Nanchang, China

[2] Lead contact

[*] Correspondence: meng7883@163.com (W. M.), wangsonghua18@126.com (S. W.)

Tel: +86-138-7062-0922

Postal address: 605 Fenglin Rd, Nanchang, Jiangxi Province 330013 China

[3] These authors contributed equally.

Running Title: Cholinergic Signaling Differentially Regulates Birdsong

ABSTRACT

Cholinergic modulation plays an important role in motor skill learning, including vocal learning. In songbirds, song premotor nucleus RA simultaneously receives inputs from song nuclei HVC and LMAN, and then its projection neurons (RAPNs) generate song motor control output. Using electrophysiological and pharmacological methods, we found that cholinergic signaling can enhance song stability by reducing HVC-RAPN excitatory synaptic transmission in adult male zebra finches, mediated by mAChRs. Although nAChRs are not effective overall, cholinergic signaling can also decrease LMAN-RAPN excitatory synaptic transmission induced by electrical stimulation via nAChRs, suggesting the potential role of cholinergic regulation in song behavior through LMAN-RA pathway. On the contrary, in adult female zebra finches, only LMAN-RAPN synaptic transmission was reduced by cholinergic signaling via mAChRs. The role of differential cholinergic regulation of song premotor circuits in songbirds' singing provides insights into the neural processes of motor skill learning.

Key words: cholinergic modulation, mAChRs, nAChRs, songbirds, song control pathways, synaptic transmission, song behavior, sex differences

INTRODUCTION

The learning and long-term retention of motor skills play a fundamental role throughout an individual's lifespan, such as handwriting, cycling, piano playing, dancing, and effective speech communication.[1] These are typically acquired gradually through multiple learning sessions until performance reaches stability,[2] within which cholinergic signaling assumes a pivotal role.[3, 4] However, the process and mechanism of cholinergic regulation on motor skill learning remain unclear.

Similar to human speech, songbird singing is a rare vocal learning behavior in animals and also a form of complex motor skill learning.[5] Juvenile songbirds optimize the matching degree between their own singing and the instructional song template through repetitive practice and self-correction relying on auditory feedback, and their songs progressively stabilize during the course of attaining adulthood.[6] Two well-defined neural pathways in songbird brain, namely vocal motor pathway (VMP) responsible for vocalization and anterior forebrain pathway (AFP) responsible for juvenile song learning and adult song plasticity, coordinately control singing.[7, 8] VMP is composed of the song premotor nuclei HVC (proper name) and the robust nucleus of the arcopallium (RA), as well as brainstem motor nucleus; AFP represents a neural circuit, where its output nucleus the lateral magnocellular nucleus of the anterior nidopallium (LMAN) projects to RA of VMP, while its input nucleus area X receives the afferent from HVC of VMP.[9]

As an analogous structure of human laryngeal motor cortex, RA serves as the convergence nucleus within song premotor circuits, and concurrently receives glutamatergic projections from its upstream motor nucleus HVC and LMAN of AFP (Figure 1A).[10, 11] Meanwhile, RA receives cholinergic projections from the ventral pallidum (VP) in basal forebrain.[12, 13] During the critical period of song learning, an elevation of acetylcholine (ACh) concentration within RA was observed in male zebra finches.[14] The combined infusion of muscarinic acetylcholine receptor (mAChR) and nicotinic acetylcholine receptor (nAChR) antagonists into RA leads to abnormal song development during the critical period.[15] RA consists of two types of neurons, including RA interneurons and RA projection neurons (RAPNs) that encode the acoustic properties of song syllables and input the information to brainstem motor nucleus.[16, 17, 18] Our previous work showed that cholinergic signaling primarily modulates RAPNs' electrophysiological activities in adult male zebra finches via

mAChRs, but not nAChRs.[19] These studies indicate that RA is a crucial target of

cholinergic modulation. Nonetheless, the essential mechanism by which cholinergic

signaling governs RA to impact song production remains unknown. To clarify this

issue, it is indispensable to understand the mechanism through which cholinergic

signaling regulates the song premotor circuits with RA as the core.

In this study, we initially employed in vitro patch-clamp whole-cell recording

combined with pharmacological approaches to separately examine the roles of two

cholinergic receptors, mAChRs and nAChRs, in cholinergic modulation of RAPNs'

excitatory synaptic afferents in adult male zebra finches. Given that the song premotor

signals input to RA are respectively derived from HVC and LMAN, we conducted a

further investigation into the influences of different cholinergic receptors on HVC-

RAPN and LMAN-RAPN excitatory synaptic transmission. Subsequently, we used in

vivo targeted pharmacological manipulation combined with song analysis to detect

the effects of different cholinergic receptors within RA on birdsongs. Moreover,

considering that zebra finches manifest typical sexual dimorphism, which only males

sing and females never do,[10] we also explored the cholinergic modulation of the song

premotor circuits similar to that of males in adult female zebra finches. Overall, our

findings delineate a gender-differentiated mechanism by which cholinergic signaling

differentially regulates song premotor circuits via mAChRs and nAChRs, thereby

potentiating song behavior. Finally, we put forward a rational hypothesis concerning

cholinergic modulation of song learning process.

RESULTS

Cholinergic signaling reduces RAPNs' excitatory synaptic afferents in adult male

and female zebra finches via mAChRs rather than nAChRs

To investigate the cholinergic modulation of RAPNs' excitatory synaptic

afferents in adult zebra finches, we first respectively recorded the influences of

cholinergic receptor agonists and antagonists on RAPNs' mEPSCs. The results

showed that the non-selective cholinergic receptor agonist carbachol (CAR)

significantly decreased the frequence and amplitude of RAPNs' mEPSCs in both

males (Pre: 7.14 ± 0.88 Hz, 10.59 ± 0.98 pA; CAR: 5.07 ± 0.63 Hz ($p = 0.0008$, $t =$

5.117, DOF = 8), 8.50 ± 1.11 pA ($p = 0.0006$, $t = 5.427$, DOF = 8); n = 9, Figure 1B)

and females (Pre: 6.09 ± 0.32 Hz, 10.84 ± 0.62 pA; CAR: 4.82 ± 0.34 Hz ($p = 0.0023$,

t = 4.366, DOF = 8), 8.54 ± 0.24 pA (p = 0.0041, t = 3.958, DOF = 8); n = 9, Figure 2B), indicating that CAR can reduce RAPNs' excitatory synaptic afferents in adult male and female zebra finches through both pre- and postsynaptic mechanisms.

The mAChR agonist Oxotremorine M iodide (OM) mimicked the effects of CAR, which also significantly reduced the frequency and amplitude of RAPNs' mEPSCs in both males (Pre: 6.31 ± 0.54 Hz, 13.14 ± 0.70 pA; OM: 4.59 ± 0.72 Hz (p = 0.0007, t = 5.694, DOF = 7), 11.98 ± 0.61 pA (p = 0.0023, t = 4.652, DOF = 7); n = 8, Figure 1C) and females(Pre: 5.20 ± 0.59 Hz, 11.52 ± 0.56 pA; OM: 3.66 ± 0.29 Hz (p = 0.0194, t = 3.019, DOF = 7), 10.21 ± 0.26 pA (p = 0.0451, t = 2.435, DOF = 7); n = 8, Figure 2C). However, the nAChR agonist DMPP had no effect on the frequency and amplitude of RAPNs' mEPSCs in both males(Pre: 5.28 ± 0.40 Hz, 14.45 ± 1.16 pA; DMPP: 5.22 ± 0.31 Hz (p = 0.8747, t = 0.163, DOF = 8), 14.65 ± 1.09 pA (p = 0.3649, t = -0.961, DOF=8); n = 9, Figure 1D).and females (Pre: 3.85 ± 0.28 Hz, 12.86 ± 0.54 pA; DMPP: 3.65 ± 0.23 Hz (p = 0.4310, t = 0.829, DOF = 8), 12.68 ± 0.47 pA (p = 0.7110, t = 0.384, DOF = 8); n = 9, Figure 2D). If CAR and the mAChR antagonist atropine (ATR) were simultaneously added, the effect of CAR was blocked by ATR, and no significant changes were observed in the frequency and amplitude of RAPNs' mEPSCs in both males (Pre: 7.01 ± 0.76 Hz, 9.43 ± 0.57 pA; CAR+ATR: 6.90 ± 0.73 Hz (p = 0.2611, t = 1.209, DOF = 8), 9.21 ± 0.36 pA (p = 0.6275, t = 0.504, DOF = 8); n = 9, Figure 1E) and females (Pre: 5.07 ± 0.41 Hz, 9.82 ± 0.85 pA; CAR+ATR: 4.93 ± 0.45 Hz (p = 0.6909, t = 0.417, DOF = 6), 9.65 ± 0.97 pA (p = 0.4046, t = 0.896, DOF = 6); n = 7, Figure 2E). When CAR and the nAChR antagonist mecamylamine (MEC) were co-administered, the effect of CAR was not blocked by MEC, and the frequency and amplitude of RAPNs' mEPSCs remained significantly decreased in both males (Pre: 8.16 ± 0.38 Hz, 12.12 ± 1.89 pA; CAR + MEC: 7.00 ± 0.51 Hz (p = 0.0072, t = 3.742, DOF = 7), 9.56 ± 1.67 pA (p = 0.0008, t = 5.661, DOF = 7); n = 8, Figure 1F) and females (Pre: 4.47 ± 0.58 Hz, 9.55 ± 0.60 pA; CAR + MEC: 3.02 ± 0.73 Hz (p = 0.0087, t = 3.597, DOF = 7), 8.21 ± 0.58 pA (p = 0.0463, t = 2.416, DOF = 7); n = 8, Figure 2F). This finding suggests that cholinergic signaling reduces RAPNs' excitatory synaptic afferents in adult male and female zebra finches via mAChRs but not nAChRs.

Figure 1 Cholinergic modulation of RAPNs' mEPSCs in adult male zebra finches

(A) Experiment schematic. In vitro whole- cell patch clamp recording of RAPNs' mEPSCs in adult male zebra finches.

(B) Top, example traces of adult male RAPNs' mEPSCs in Pre (gray) and CAR (blue); Bottom, quantification of mEPSCs' frequency (left) and amplitude (right) in Pre and CAR, n = 9.

(C-F) Similar to (B). OM (C), n = 8; DMPP (D), n = 9; CAR+ATR (E), n = 7; CAR+MEC (F), n = 8. ** $p \leq 0.01$, *** $p \leq 0.001$, paired t-test. Plots represent mean ± SEM.

Figure 2 Cholinergic modulation of RAPNs' mEPSCs in adult female zebra finches

(A) Experiment schematic. In vitro whole-cell patch clamp recording of RAPNs' mEPSCs in adult female zebra finches.

(B) Top, example traces of adult female RAPNs' mEPSCs in Pre (black) and CAR (red); Bottom, quantification of mEPSCs' frequency (left) and amplitude (right) in Pre and CAR, n = 9.

(C-F) Similar to (B). OM (C), n = 8; DMPP (D), n = 9; CAR+ATR (E), n = 9; CAR+MEC (F), n = 8. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, paired *t*-test. Plots represent mean ± SEM.

Cholinergic signaling reduces HVC-RAPN excitatory synaptic transmission in adult male zebra finches via mAChRs rather than nAChRs

HVC-RA pathway is a crucial excitatory synaptic afferent of RA. Our further results showed that CAR significantly reduced the amplitude of RAPNs' evoked EPSCs (eEPSCs) induced by stimulating HVC-RA projection fibers in adult males (Pre: 120.05 ± 9.19 pA, CAR: 71.53 ± 6.56 pA, n = 10, $p = 0.00004$, $t = 7.350$, DOF = 9, Figure 3B), indicating that CAR can prominently decrease HVC-RAPN excitatory synaptic transmission. Meanwhile, the effect of CAR was accompanied by a significant increase in paired-pulse facilitation ratio (PPR) (Pre: 0.92 ± 0.08, CAR: 1.28 ± 0.07, n = 7, $p = 0.0007$, $t = -6.266$, DOF = 6, Figure 3B), suggesting that

presynaptic mechanism was involved in the role of CAR on HVC-RAPN excitatory
synaptic transmission.

OM mimicked the effect of CAR, which likewise significantly reduced HVC-
RAPN eEPSCs' amplitude (Pre: 137.07 ± 14.06 pA, OM: 73.43 ± 9.23 pA, n = 8, $p =$
0.0002, $t = 6.678$, DOF = 7, Figure 3C) and caused a significant elevation in PPR (Pre:
0.97 ± 0.07, OM: 1.16 ± 0.07, n = 7, $p = 0.0379$, $t = -2.652$, DOF = 6, Figure 3C).
However, DMPP had no effect on HVC-RAPN eEPSCs' amplitude (Pre: $122.73 \pm$
14.78 pA, DMPP: 120.19 ± 11.75 pA, n = 11, $p = 0.7904$, $t = 0.273$, DOF = 10, Figure
3D). If CAR and ATR were concurrently administered, the inhibitory action of CAR
on HVC-RAPN eEPSCs' amplitude was obstructed by ATR (Pre: 119.45 ± 8.56 pA,
CAR+ATR: 117.50 ± 15.11 pA, $n = 8$, $p = 0.8948$, $t = 0.137$, DOF = 7, Figure 5E).
Nevertheless, when CAR and MEC were concurrently administered, MEC failed to
neutralize the effect of CAR (Pre: 130.82 ± 5.30 pA, CAR+MEC: 93.80 ± 8.79 pA, n
$= 11$, $p = 0.0002$, $t = 5.606$, DOF = 10, Figure 5F). These results suggest that
cholinergic signaling reduces HVC-RAPN excitatory synaptic transmission in adult
male zebra finches via mAChRs but not nAChRs.
213
214

Figure 3 Cholinergic modulation of HVC-RAPN eEPSCs in adult male zebra finches

(A) Experiment schematic. In vitro whole-cell patch clamp recording of HVC-RAPN eEPSCs in adult male zebra finches.

(B) Top, example traces of adult male HVC-RAPN eEPSCs in Pre (black) and CAR (red). Quantification of eEPSCs' amplitude in Pre and CAR, n = 10; Bottom, example traces of eEPSCs in paired pulse stimulation. Quantification of paired-pulse ratio in Pre and CAR, n = 7.

(C) Similar to (B). OM: Top, n = 8; Bottom, n = 7.

(D-F) Example traces of adult male HVC-RAPN eEPSCs in Pre (black), DMPP (D, red), CAR+ATR (E, red) and CAR+MEC (F, red). Quantification of eEPSCs' amplitude in Pre, DMPP (D, n = 11), CAR+ATR (E, n = 8), and CAR+MEC (F, n = 11). * $p \leq 0.05$, *** $p \leq 0.001$, paired *t*-test. Plots represent mean ± SEM.

Cholinergic signaling reduces LMAN-RAPN excitatory synaptic transmission in adult male zebra finches via nAChRs rather than mAChRs

LMAN-RA pathway is another important excitatory synaptic afferent of RA. CAR significantly decreased the amplitude of RAPNs' eEPSCs elicited by stimulating LMAN-RA projection fibers in adult males (Pre: 126.48 ± 9.88 pA, CAR: 98.30 ± 10.11 pA, n = 9, $p = 0.0001$, $t = 6.619$, DOF = 8, Figure 4B), indicating that CAR can markedly reduce LMAN-RAPN excitatory synaptic transmission. During the CAR effect, PPR was significantly increased (Pre: 0.92 ± 0.05, CAR: 1.14 ± 0.07, n = 8, $p =$

0.0191, t = -3.031, DOF = 7, Figure 4B), suggesting that presynaptic mechanism was involved in the role of CAR on LMAN-RAPN excitatory synaptic transmission.

OM exerted no influence on LMAN-RAPN eEPSCs' amplitude (Pre: 131.67 ± 12.60 pA, OM: 122.81 ± 12.13 pA, n = 8, p = 0.2537, t = 1.243, DOF = 7, Figure 4C). However, DMPP simulated the effect of CAR, significantly reducing LMAN-RAPN eEPSCs' amplitude (Pre: 146.85 ± 10.60 pA, DMPP: 102.52 ± 9.45 pA, n = 8, p = 0.0008, t = 5.537, DOF = 7, Figure 4D), and simultaneously causing a significant increase in PPR (Pre: 0.87 ± 0.05, DMPP: 1.14 ± 0.07, n = 8, p = 0.0020, t = -4.747, DOF = 7, Figure 4D). When CAR and ATR were applied together, the reduction effect of CAR on LMAN-RAPN eEPSCs' amplitude was not blocked by ATR (Pre: 110.24 ± 9.16 pA, CAR+ATR: 79.38 ± 7.17 pA, n = 10, p = 0.0031, t = 3.992, DOF = 9, Figure 4E). If CAR and MEC were applied simultaneously, the effect of CAR was blocked by MEC (Pre: 124.01 ± 10.32 pA, CAR+MEC: 119.83 ± 11.34 pA, n = 8, p = 0.6553, t = 0.466, DOF = 7, Figure 4F). These results demonstrate that cholinergic signaling reduces LMAN-RAPN excitatory synaptic transmission in adult male zebra finches via nAChRs but not mAChRs. Namely, receptor mechanisms of cholinergic modulation on the excitatory synaptic transmission of the two song premotor pathways, LMAN-RA and HVC-RA, in males are different.

Figure 4 Cholinergic modulation of LMAN-RAPN eEPSCs in adult male zebra finches

(A) Experiment schematic. In vitro whole-cell patch clamp recording of LMAN-RAPN eEPSCs in adult male zebra finches.

(B) Top, example traces of adult male LMAN-RAPN eEPSCs in Pre (gray) and CAR (blue). Quantification of eEPSCs' amplitude in Pre and CAR, n = 9; Bottom, example traces of eEPSCs in paired pulse stimulation. Quantification of paired-pulse ratio in Pre and CAR, n = 8.

(C) Example traces of adult male LMAN-RAPN eEPSCs in Pre (gray) and OM (blue). Quantification of eEPSCs' amplitude in Pre and OM, n = 8.

(D) Similar to (B). DMPP: Top, n = 8; Bottom, n = 8.

(E-F) Similar to (C). CAR+ATR (E), n = 10; CAR+MEC (F), n = 8. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, paired t-test. Plots represent mean ± SEM.

Cholinergic signaling has no effect on HVC-RAPN excitatory synaptic transmission in adult female zebra finches

In contrast to the findings in males, CAR exerted no impact on HVC-RAPN eEPSCs' amplitude in adult females (Pre: 122.97 ± 8.65 pA, CAR: 115.66 ± 11.75 pA, n = 10, $p = 0.4391$, $t = 0.809$, DOF = 9, Figure 5B). Additionally, neither DMPP (Pre:

277　126.04 ± 11.88 pA, DMPP: 121.63 ± 10.59 pA, n = 8, p = 0.5647, t = 0.604, DOF = 7,

278　Figure 5C) nor OM (Pre: 126.26 ± 14.72 pA, OM: 117.62 ± 13.01 pA, n = 9, p =

279　0.1521, t = 1.583, DOF = 8, Figure 5D) had any effect on HVC-RAPN eEPSCs'

280　amplitude. This postulates that HVC-RAPN excitatory synaptic transmission in adult

281　female zebra finches is not regulated by cholinergic signaling.

Figure 5 Cholinergic modulation of HVC-RAPN and LMAN-RAPN eEPSCs in
adult female zebra finches
(A) Experiment schematic. In vitro whole-cell patch clamp recording of HVC-RAPN
and LMAN-RAPN eEPSCs in adult female zebra finches.
(B) Example traces of adult female HVC-RAPN eEPSCs in Pre (black) and CAR
(red). Quantification of eEPSCs' amplitude in Pre and CAR, n = 10.
(C-D) Similar to (B). DMPP (C), n = 8; OM (D), n = 9.
(E) Top, example traces of adult female LMAN-RAPN eEPSCs in Pre (gray) and

CAR (blue). Quantification of eEPSCs' amplitude in Pre and CAR, n = 10; Bottom,
example traces of eEPSCs in paired pulse stimulation. Quantification of paired-pulse
ratio in Pre and CAR, n = 7.
(F) Similar to (E). OM: Top, n = 8; Bottom, n = 7.
(G) Example traces of adult female LMAN-RAPN eEPSCs in Pre (gray) and DMPP
(blue). Quantification of eEPSCs' amplitude in Pre and DMPP, n = 8.
(H-I) Similar to (G). CAR+ATR (H), n = 10; CAR+MEC (I), n = 9. * $p \leq 0.05$, ** $p \leq$
0.01, *** $p \leq 0.001$, paired t-test. Plots represent mean ± SEM.
300

301 Cholinergic signaling reduces LMAN-RAPN excitatory synaptic transmission in

302 adult female zebra finches via mAChRs rather than nAChRs

Similar to males, CAR significantly reduced LMAN-RAPN eEPSCs' amplitude

in adult females (Pre: 122.74 ± 8.71 pA, CAR: 79.81 ± 6.61 pA, n = 10, $p = 0.0006$, t

= 5.110, DOF = 9, Figure 5E), and led to a significant increase in PPR (Pre: 1.02 ±

0.07, CAR: 1.37 ± 0.08, n = 7, $p = 0.0071$, $t = -2.782$, DOF = 6, Figure 5E),

suggesting that there are also presynaptic mechanisms involved in the CAR

modulation of LMAN-RAPN excitatory synaptic transmission in females.

OM mimicked the reduction effect of CAR on LMAN-RAPN eEPSCs'

amplitude in females (Pre: 130.25 ± 8.59 pA, OM: 92.08 ± 7.44 pA, n = 8, $p = 0.0001$,

$t = 7.614$, DOF = 7, Figure 5F), accompanied by a significant increase in PPR (Pre:

1.02 ± 0.06, OM: 1.36 ± 0.10, $n = 7$, $p = 0.0318$, $t = -3.997$, DOF = 6, Figure 5F).

Whereas, DMPP had no influence on LMAN-RAPN eEPSCs' amplitude in females

(Pre: 141.88 ± 6.44 pA, DMPP: 149.49 ± 10.03 pA, n = 8, $p = 0.3943$, $t = -0.907$,

DOF= 7, Figure 5G). Simultaneously applying CAR and ATR, the effect of CAR was

blocked by ATR (Pre: 122.27 ± 8.41 pA, CAR+ATR: 118.89 ± 10.46 pA, n = 10, $p =$

0.7415, $t = 0.340$, DOF= 9, Figure 5H). While simultaneously applying CAR and

MEC, the effect of CAR was not blocked by MEC (Pre: 132.91 ± 12.19 pA,

CAR+MEC: 115.44 ± 10.91 pA, n = 9, $p = 0.0361$, $t = 2.513$, DOF= 8, Figure 5I).

These results indicate that, contrary to the receptor mechanism of cholinergic

modulation on LMAN-RAPN excitatory synaptic transmission in males, cholinergic

signaling reduces LMAN-RAPN excitatory synaptic transmission in adult female

zebra finches via mAChRs but not nAChRs.

325 Cholinergic signaling within RA enhances song stability of adult male zebra

326 finches via mAChRs rather than nAChRs

To further validate the impact of cholinergic signaling within RA on adult song

behavior and its receptor mechanisms, the results of behavioral experiments combined

with an in vivo pharmacological method (Figure 6A and B) showed that CAR

microinjection onto RA significantly elevated the similarity to own song of adult

males ((Pre: 87.64 ± 0.49%, CAR: 92.78 ± 0.95%, n = 4, p = 0.0264, t = -4.087, DOF

= 3, Figure 6C and D) and reduced the song entropy (Pre: -2.78 ± 0.05, CAR: -3.05 ±

0.09, n = 4, p = 0.0098, t = 5.875, DOF = 3, Figure 6C and E). OM microinjection

onto RA exhibited a comparable effect to CAR, which also increased song similarity

(Pre: 85.98 ± 3.01%, OM: 92.61 ± 1.70%, n = 4, p = 0.0282, t = -3.987, DOF = 3,

Figure 6C and D) and decreased song entropy (Pre: -2.89 ± 0.18, OM: -3.05 ± 0.17, n

= 4, p = 0.0017, t = 10.591, DOF = 3, Figure 6C and E), whereas DMPP

microinjection had no effect on song similarity (Pre: 93.00 ± 1.67%, DMPP: 93.06 ±

1.46%, n = 4, p = 0.8414, t = -0.218, DOF = 3, Figure 6C and D) and entropy (Pre: -

2.91 ± 0.18, DMPP: -2.91 ± 0.11, n = 4, p = 0.9968, t = -0.004, DOF = 3,Figure 6C

and E). For comparison, the results of phosphate buffered saline (PBS) microinjection

showed that there was no obvious change in song similarity (Pre: 92.02 ± 1.72%, PBS:

92.68 ± 2.25%, n = 4, p = 0.3002, t = -1.249, DOF = 3, Figure 6C and D) and entropy

(Pre: -2.22 ± 0.09, PBS: -2.25 ± 0.07, n = 4, p = 0.2270, t = 1.515, DOF = 3, Figure

6C and E), thereby ruling out the influence brought by the microinjection operation

per se. In addition, other song acoustic characteristics were not notably influenced by

the diverse drug microinjections (Figure S1). These results manifest that cholinergic

signaling within RA enhances song stability of adult male zebra finches via mAChRs

but not nAChRs.

Figure 6 Cholinergic signaling within RA enhances song stability of adult male zebra finches via mAChRs rather than nAChRs

(A) Experiment schematic. In vivo pharmacological microinjection onto bilaterally RA and song recording.

(B) Left, a free-moving adult male zebra finch in a recording studio with cannulas bilaterally implanted over RA; Right, cannula tracks in a coronal section.

(C) Example spectrograms in Pre and CAR, OM, DMPP or PBS. 'iabcde' indicates the syllables constituting a song motif.

(D) Similarity to own songs in Pre and CAR (black, n = 4), Pre and OM (red, n = 4), Pre and DMPP (gray, n = 4), Pre and PBS (blue, n = 4).

(E) Song entropy in Pre and CAR (black, n = 4), Pre and OM (red, n = 4), Pre and DMPP (gray, n = 4), Pre and PBS (blue, n = 4). ** $p \leq 0.01$, paired *t*-test. Plots represent mean ± SEM.

DISCUSSION

To illuminate cholinergic modulation of songbirds' song behavior, it is indispensable to elucidate the cholinergic regulatory mechanism at the neural pathway level. We discover a mechanism by which cholinergic signaling separately regulates song premotor circuits via different receptors, thereby enhancing song stability of adult males. Moreover, a similar mechanism exists in the same circuits of adult females. Furthermore, we postulate a hypothesis concerning cholinergic regulatory process of song learning in juveniles and song maintenance in adults. From a broader perspective, our findings provide significant insights into cholinergic regulation of motor skill learning.

Songbirds are an excellent animal model for revealing cholinergic modulation of motor skill learning

Central cholinergic system exists in the brains of fish,[20, 21, 22] amphibians,[23, 24] reptiles,[23, 24] mammals,[25, 26, 27, 28] and birds,[29, 30, 31, 32] which mainly participates in various higher neural activities such as motor skill acquisition,[3, 33] learning and memory,[34, 35] sleep and wakefulness,[36, 37] and attention maintenance.[38, 39] ACh exerts different effects by binding to two different types of receptors, mAChRs (G-protein coupled receptors) and nAChRs (ligand-gated ion channel receptors). Both receptors are expressed in all layers of the vertebrate cerebral cortex,[20, 28, 40] with different expression patterns among different layers and cell types.[41] For example, it has been reported that nAChRs is mainly expressed on interneurons in the prefrontal cortical layer 2/3 of rats, while only a small portion of pyramidal neurons express nAChRs.[42] It was found that there are regional and interlayer differences in the expression of the two main cortical mAChR subtypes (M1 and M2) in the prefrontal cortex of rhesus monkeys.[43]

The construction of avian subcortical cholinergic system is analogous to that of mammals. Research has confirmed that the cholinergic system of pigeon in basal forebrain, epithalamus, isthmus, and hindbrain closely resembles that of reptiles and mammals.[29] Further research indicates that the basal telencephalic cholinergic system in pigeons and budgerigars is anatomically very similar to the cholinergic basal nuclear system in mammals, particularly in terms of their projection patterns in

cerebral cortex or pallium.[32, 44] Although the neurons in avian pallium are not

hierarchically organized like those in mammalian cerebral cortex, cholinergic receptor

distribution in avian brain also shows regional specificity.[30, 31]

In songbirds, both male and female zebra finches have cholinergic projection

fibers in song nuclei and auditory nuclei.[45, 46] Studies have shown that HVC and RA

are rich in acetylcholinesterase, suggesting a strong cholinergic dominance over these

two nuclei.[45, 46, 47] The cholinergic innervation of HVC and RA originates from VP,

which is analogous to the mammalian basal forebrain cholinergic system, specifically

the nucleus basalis of Meynert.[12]

Evidence suggests that mAChRs are expressed within song nuclei,[48, 49] and

nAChRs are also present in multiple song nuclei.[50] Asogwa et al. cloned four out of

the five mammalian mAChR subunits (Chrm2-5) in male zebra finches. They found

that, at each developmental stage, the expression of excitatory subunits (chrm3 and

chrm5) consistently exhibits lower levels compared to inhibitory subunits (chrm2 and

chrm4) within most song nuclei, including RA.[51] Recently, all 15 types of nAChR

subunits in male zebra finch brain were cloned. It was confirmed that most nAChR

subunits (except for ChrnA1, A6, A9, and A10) are expressed in song related

pathways during the critical period of song learning, while only 6 nAChR subunits

(ChrnA2-5, A7, and B2) are expressed in adulthood.[52] This suggests that the

expression of most nAChR subunits undergoes continuous changes throughout song

formation process in various song nuclei, and the expression types of nAChR subunits

in adulthood are much fewer than those during song learning.

Central cholinergic system plays a crucial role in the regulation of motor skill

learning.[53, 54, 55] However, the precise mechanisms through which this system governs

such learning remain elusive, and elucidating its role poses significant challenges.

Songbird's song learning is a complex form of sensory-motor learning, which is an

extremely rare vocal learning behavior similar to human language learning.[5]

Pharmacological and behavioral research showed that subcutaneous administration of

mid (0.18mg/kg) and high (0.54mg/kg) dose of the nAChR agonist nicotine can elicit

sensitized responses of song production and locomotor activity in adult male zebra

finches.[56] Continuous nicotine administration for 7 days can alter the tempo and

rhythm of adult crystallized songs, and this effect may persist for an extended period

(over 2 months) after discontinuation of nicotine administration.[57] These findings

indicate that central cholinergic system likely plays a significant role in the
vocalization and song learning processes of songbirds. Therefore, songbirds could
serve as an excellent research model for in-depth exploration of cholinergic regulation
on complex motor control and learning mechanisms, including human speech.

A mechanism by which cholinergic signaling separately regulates song premotor
circuits of male songbirds through different receptors thereby promoting song
stability

The electrophysiological activities of HVC and RA are associated with the neural
encoding of song motor. The motor control signals generated by HVC require
processing through RA in order to ultimately produce appropriate song motor
encoding for coordinated activation of syringeal and respiratory muscles.[58, 59] Shea
and Margoliash found that cholinergic signaling attenuates the auditory responses of
HVC and RA to bird's own song in adult male zebra finches.[60] Furthermore, distinct
HVC projection neurons (HVC-to-RA and HVC-to-area X projection neurons) exhibit
differential responses to cholinergic signaling, and this diversity of response may
facilitate the coordination of cholinergic regulation between HVC-RA motor
command transmission and the reception of HVC motor signals by AFP.[61] Jaffe and
Brainard found that dialyzing CAR into HVC increases HVC neuron activities,
resulting in an elevation of song pitch, amplitude, tempo and stereotypy to levels
approaching those of direct courtship songs in adult male Bengalese finches, which
can be weakened by blocking mAChRs.[62]

An early study revealed a notable elevation in ACh levels within HVC, LMAN
and RA of zebra finches during the critical period for song acquisition, followed by a
gradual decline as the birds approach adulthood.[14] ACh transient increase within RA
during critical period is associated with heightened acetyltransferase (ChAT)
activity.[63] Additionally, biochemical research has shown that during the critical period
of song learning in zebra finches, CAR significantly increases the phosphoinositide
turnover within the synaptoneurosomes of RA neurons.[64] Through intracellular
electrophysiological recordings in brain slices, Salgado-Commissariat et al.
discovered that nicotine enhances RA neuron excitability in adult male zebra finches,
without distinguishing between the types of RA neurons (projection neurons or
interneurons). Furthermore, by activating different nAChR subtypes, they

demonstrated that tetanic stimulation induces bidirectional synaptic plasticity (long-
term potentiation and long-term depression) in LMAN-RA pathway.[65]

RAPNs are homologous to layer 5 pyramidal neurons in mammalian motor
cortex, and their activities govern song production.[66, 67] Our previous results obtained
from brain slice whole-cell current-clamp recordings indicate that cholinergic
signaling can induce a hyperpolarization of membrane potential, an increase in
afterhyperpolarization potential and membrane conductance, and a reduction in action
potential firing in RAPNs of adult male zebra finches via mAChRs rather than
nAChRs.[19, 68] This suggests that under physiological conditions, cholinergic signaling
may influence RAPN activities by altering their intrinsic membrane properties,
thereby regulating song behavior. In present study, the results of brain slice whole-cell
voltage-clamp recordings showed that cholinergic signaling significantly reduces
RAPNs' mEPSCs in adult male zebra finches through mAChRs rather than nAChRs,
indicating that RAPNs' excitatory synaptic afferents in adulthood is regulated by
cholinergic signaling, and mAChR mediated action remains predominant. Cholinergic
regulation on RAPNs' intrinsic membrane properties and excitatory synaptic afferents
is predominantly mediated by the inhibitory effect of mAChRs to reduce excitability,
which is consistent with the circumstance that the inhibitory mAChR subunits are
expressed significantly more than the constitutively low-expressed excitatory mAChR
subunits within RA at all development stages.[51]

In mammals, similar to our results, ACh local application inhibits the excitability
of layer 5 pyramidal neurons in rat prefrontal cortex, somatosensory cortex, and visual
cortex.[69] Another study has confirmed that ACh induces membrane potential
hyperpolarization and enhances membrane conductance of layer 5 pyramidal neurons
in rat neocortex via mAChRs, thereby exerting an inhibitory action.[70] Additionally, it
has been reported that CAR reduces glutamatergic excitatory synaptic transmission of
layer 5 pyramidal neurons in rat sensory cortex, and this effect is blocked by mAChR
antagonists.[71] In fact, the impacts of cholinergic signaling on neurons in diverse
cortical areas and different layers are not homogeneous. A study has found that CAR
increases the firing of rat neocortical neurons through different mAChR subunits, but
reduces their synaptic transmission.[72] Generally, ACh can exert an excitatory action
on neurons at the deepest layer (layer 6) of mouse primary cortex, including primary
somatosensory cortex, primary motor cortex and combined medial prefrontal cortex.

It has been disclosed through pharmacological methods that nAChRs and mAChRs

jointly contribute to mediating the ACh's action on the neurons in layer 6 of primary

motor cortex. However, the neurons in layer 6 of combined medial prefrontal cortex

display a stronger response to ACh, and the effect is mainly mediated by nAChRs.

The response of layer 6 neurons in primary somatosensory cortex to ACh is relatively

weak, and the receptor mechanism of neurons therein affected by ACh is

inconsistent.[26] By comparing the aforementioned studies on mammals with our

experimental results, it can be noted that the cholinergic regulation of song premotor

brain regions in songbirds exhibits both similarities to that of cerebral cortex in

mammals and interspecific dissimilarities or its functional specificity.

It is worth noting that the pharmacological research on layer 6A pyramidal cell

types of rat somatosensory cortex in recent years has revealed that low concentration

(30 μM) ACh can induce hyperpolarization of corticocortical pyramidal cells via M4

and M1 mAChRs, while depolarizing corticothalamic pyramidal cells. However,

under the influence of high concentration (1 mM) ACh, corticothalamic pyramidal

cells can be depolarized via α4β2 nAChRs, while the membrane potential of

corticocortical pyramidal cells is unaffected. Nevertheless, mEPSCs' frequency of

corticocortical pyramidal cells is reduced, whereas that of corticothalamic pyramidal

cells is increased. Additionally, glutamate synaptic transmission in corticocortical

pyramidal cells is inhibited via M4 mAChRs, but in the corticothalamic pyramidal

cells, it is enhanced via α4β2 nAChRs. This study indicates a potential mechanism

through which ACh might separately regulate diverse projection neurons in the same

cortical region via both nAChRs and mAChRs, thus achieving the coordination

among synaptic transmission functions.[41] However, within the song premotor nucleus

RA of songbirds, there is merely one type of projection neurons (projecting to

brainstem motor nucleus). The integration of song premotor signals from HVC-RA

and LMAN-RA pathways within RAPNs is the critical determinant in generating the

encoded output for controlling song production.[73, 74] The more crucial mechanism

underlying cholinergic modulation of song behavior might lie in how to achieve the

discrete regulation of HVC-RA pathway and LMAN-RA pathway, concomitant with

the reciprocal coordination and collaboration in the regulation between the signal

transmissions of the two pathways. Our present results revealed that cholinergic

signaling markedly decreases glutamatergic HVC-RAPN eEPSCs' amplitude via

mAChRs in adult male zebra finches. On the contrary, the reduction of glutamatergic

LMAN-RAPN eEPSCs' amplitude caused by cholinergic signaling was mediated by

nAChRs in adult males. Meanwhile, cholinergic regulation of both HVC-RAPN and

LMAN-RAPN synaptic transmission is accompanied by alterations in PPR (an

electrophysiological indicator suggesting presynaptic mechanisms[75]). Our discoveries

demonstrate a mechanism by which cholinergic signaling achieves the distinct

regulation of song premotor circuits through two receptors, nAChRs and mAChRs.

Recently, it has also been found in adult mice that cholinergic signaling can

differentially regulate the glutamatergic synaptic transmission of the projections from

prelimbic cortex and thalamus to Basolateral Amygdala pyramidal neurons,

respectively, through different receptors.[76] The disclosure of the differential regulatory

mechanism of cholinergic receptors further underpins the crucial role of cholinergic

regulation adapted to specific functions in relevant brain regions.

Most tellingly, our further behavioral research combined with in vivo targeted

pharmacological manipulation indicated that CAR microinjection onto RA

significantly enhances song stability of adult male zebra finches. Furthermore,

microinjection of mAChRs agonist has the same effect as CAR, whereas

microinjection of nAChRs agonist has no effect. The result is consistent with the

finding of Jaffe and Brainard in adult male Bengalese finches that dialysis infusion of

CAR into HVC affects singing, both increasing song stability.[62] Concurrently, the

result indicates that cholinergic signaling within RA governs song behavior in

adulthood predominantly via mAChRs rather than nAChRs. This receptor mechanism

is consistent with that of cholinergic signaling regulating RAPNs' mEPSCs and

intrinsic membrane properties in adult males, all through mAChRs but not nAChRs.[19]

Taken together, given that RAPNs are marked by the electrophysiological

characteristic of high excitability,[77, 78] cholinergic signaling can diminish the

glutamatergic excitatory synaptic transmission of song premotor pathways in adult

males via mAChRs, thus appropriately lowering the high excitability of RAPNs and

potentially enhancing their information integration capacity by improving signal-to-

noise ratio through this mechanism, ultimately promoting song stability. We recorded

that RAPNs' excitability in 26-day-old fledglings (at subsong stage) is conspicuously

higher than that in adults at song crystallization stage (data not shown), which is in

accordance with our anticipation that the decrease in RAPNs' excitability is associated

with the enhancement of song stability. Moreover, evidence indicates that apart from cholinergic signaling, RA is also controlled by a multitude of other neurotransmitters, such as monoamine neurotransmitters.[79, 80, 81, 82, 83] Cholinergic signaling could merely be one constituent in the modulation of the chemical "cocktail". Meanwhile, RAPNs' inhibitory synaptic inputs from the GABAergic interneuron network would exert an important balancing effect.[84]

Another investigation has validated that cholinergic signaling within RA also has a substantial impact on song learning during developmental stage. Puzerey et al. employed in vivo reverse microdialysis technique to administer nAChR and mAChR antagonists in juvenile zebra finch RA for the purpose of chronically blocking cholinergic signaling in RA during song learning. After several weeks, it was discovered that singing quantity declines, and song learning process becomes disordered, such as excessive increase in song variability, abnormal acoustic features, and a decrease in similarity to tutor song.[15] Furthermore, throughout the process of development and song acquisition, the synaptic density and amount of HVC-RA projection fibers increase conspicuously, yet the synaptic density and amount of LMAN-RA projection fibers diminish markedly.[11, 85] Although LMAN-RA pathway assumes a pivotal role in song learning[86, 87] and provides motor correction signals for HVC-RA pathway,[59, 88, 89] and the lesion of LMAN in juvenile zebra finches can disrupt song acquisition, the lesion of LMAN in adulthood has no impact on the already stable songs.[90, 91] Consequently, based on the findings of predecessors and ours, it is reasonable to believe that during the song learning period of juvenile zebra finches, the mutual coordination and collaboration between the cholinergic regulation of LMAN-RA pathway mediated by nAChRs and that of HVC-RA pathway mediated by mAChRs exert a significantly facilitating effect on the process of song learning. However, during the process of song gradually stabilizing, with the weakening of LMAN-RA pathway and the strengthening of HVC-RA pathway, the cholinergic modulation of LMAN-RA pathway mediated by nAChRs also weakens accordingly. Therefore, the reinforcing effect of cholinergic signaling on the stability of adult birdsongs is mainly regulated by HVC-RA pathway mediated via mAChRs. Herein, we propose a mechanism hypothesis that cholinergic signaling influences song production by regulating song premotor circuits (Figure 7A), and we are currently undertaking a study on cholinergic modulation of song premotor circuits during song

learning period of juvenile zebra finches. The finding of Asogwa et al. that the

expression types of nAChR subunits in adulthood within song related pathways,

including LMAN-RA pathway, are much fewer than those in critical period of song

learning offers support for our hypothesis.[52]

Although cholinergic modulation of LMAN-RA pathway mediated by nAChRs

turns quiescent or latent in adulthood, our experimental observations still imply that in

the context of electrical stimulation of LMAN-RA projection fibers, cholinergic

signaling can give rise to a decline in LMAN-RAPN excitatory synaptic transmission

in adult males via nAChRs, suggesting that this pathway is likely to undergo adaptive

modifications under the elicitation of certain physiological conditions (such as

experience-dependent song plasticity[92]) or pathological conditions. For example, in

the event that the auditory feedback of adult songbirds undergoes modifications or is

absent, LMAN-RA pathway will facilitate variations in the signal output of HVC-RA

pathway and induce changes in vocalization.[93] It was reported that by activating

different nAChR subtypes, long-term synaptic plasticity could be induced in the

LMAN-RA pathway of adult male zebra finches through tetanic stimulus,[65]

suggesting the potential cholinergic regulation of LMAN-RA pathway in adulthood.

614
Figure 7 Mechanism hypotheses on the cholinergic modulation of song premotor
circuits in male and female zebra finches
(A) A hypothesis regarding the mechanism through which cholinergic signaling
influences song production by regulating song premotor circuits in male zebra finches.
(B) A hypothesis regarding the mechanism through which cholinergic signaling
impacts the perception and/or selection of male songs by regulating song premotor
circuits in adult female zebra finches. Details in "DISCUSSION".
622

Cholinergic modulation of LMAN-RA pathway mediated by mAChRs may be a potential mechanism for perception and selection of birdsongs in females

Zebra finches are a species among passerines that exhibits distinct sexual dimorphism in both appearance and song behavior. Adult male birds attract females through singing, whereas females do not sing.[94] Correspondingly, most of song control nuclei in males are much larger than those in females.[95] Studies have demonstrated that the neurons within HVC and RA of females are smaller and less numerous than those of males, so the volumes of the two nuclei in females are considerably smaller than those in males.[96] Such a discrepancy is associated with sex hormones[97] and may even be linked to sex-specific transcriptomes.[8] It has been reported that RAPNs' excitability of adult female zebra finches is significantly lower than that of males.[98] Our previous studies have demonstrated that both excitatory and inhibitory synaptic transmission of RAPNs have sexual dimorphism. In adult female zebra finches, both the frequency and amplitude of RAPNs' spontaneous EPSCs (sEPSCs) and mEPSCs are notably lower than those of males.[99] Moreover, both the frequency and amplitude of spontaneous inhibitory postsynaptic currents (sIPSCs) and mIPSCs of RAPNs in females are also conspicuously lower than those in males.[84] A study has shown that neuropeptides expression in the neurons of forebrain song control nuclei in female zebra finches is significantly lower than that in males.[100] However, the in-depth research regarding the effect of neurotransmitters on the song related nuclei of female songbirds is rather limited.

Previous immunohistochemical results demonstrated that the size and density of ChAT-immunoreactive somata projected from VP to HVC and RA in males are significantly higher than those in females,[101] and the density of ChAT immunoreactive positive-labeled fibers around area X in males is also conspicuously greater than that in females,[47] indicating sex differences in the cholinergic modulation of song control system. Our present study showed that, similar to males, cholinergic signaling can significantly reduce RAPNs' mEPSCs in adult female zebra finches via mAChRs rather than nAChRs, indicating that RAPNs' excitatory synaptic afferents in adult females is regulated by cholinergic signaling, and the role of mAChRs predominates. To our knowledge, this should be the first time to provide direct evidence for cholinergic modulation of female song control nuclei.

Studies have indicated that the development and formation of HVC-RA song

premotor pathway in female zebra finches is incomplete.[102, 103, 104] Moreover, it has been reported that sex differences in the myelination of zebra finch HVC-RA tract are significant, and the degree of myelination in females is lower than that in males and remains unchanged with development.[105] Additionally, the quantity of RAPNs in females is markedly less than that in males.[106] These pieces of evidence seemingly offer a rational explanation for the fact that females do not sing. Nevertheless, it has still been proved that female zebra finches have HVC-RA projection fibers like males.[107] Meanwhile, a study of Wang et al. demonstrated that electrical stimulation of HVC-RA projection fibers in adult female zebra finches could result in the corresponding neural activity in RA, indicating the presence of HVC-RA synaptic transmission in females.[108] In accordance with this finding, our present results indicate that RAPNs' eEPSCs can be recorded by electrically stimulating HVC-RA projection fibers in adult female zebra finches, reconfirming the presence of HVC-RA synaptic transmission in adult females. However, we found that, in contrast to males, cholinergic signaling has no effect on HVC-RAPN eEPSCs in females, suggesting that HVC-RAPN excitatory synaptic transmission of females is not regulated by cholinergic signaling. This result indicates that cholinergic modulation of HVC-RA song premotor pathway in adult zebra finches exhibits a typical sex dimorphism.

Among all song nuclei, only the LMAN of females is approximately the same in volume as that of males.[109] Consistent with males, the LMAN of female zebra finches likewise forms neural connections projected to RA in early developmental stage.[103] Moreover, a study has found that in adulthood, the number of synapses in the LMAN of females is significantly higher than that of males.[110] However, it has also been found that LMAN neurons projecting to RA in adult female zebra finches are much less than that in adult males, which might be caused by the massive loss of LMAN-to-RA projection neurons in females during the process of sexual differentiation and song learning.[111] Our further examination revealed that cholinergic signaling significantly reduces the amplitude of RAPNs' eEPSCs evoked by electrically stimulating the LMAN-RA projection fibers of adult female zebra finches. Moreover, this process is accompanied by the alteration of PPR, indicating that the presynaptic mechanism participated in cholinergic regulation on LMAN-RAPN excitatory synaptic transmission in females. However, unlike the cholinergic regulation of LMAN-RAPN synaptic transmission in males mediated by nAChRs, that in females is

mediated by mAChRs rather than nAChRs, which is in line with the receptor

mechanism of cholinergic signaling regulating RAPNs' mEPSCs in females. Like

zebra finches, in another songbird cowbirds (Molothrus ater), only males sing.

Although females do not sing, they can distinguish males by perceiving birdsongs.

Research has shown that the volume of female LMAN and the quantity of neurons

therein are positively correlated with mating choices, indicating that non-singing

females may perceive and choose male songs by LMAN.[112] Hence, female zebra

finches, comparable to their male counterparts, might establish memories of song

templates (songs heard during juvenile period, such as father's song[113, 114]) in the early

life stages and gradually develop song perception, among which LMAN may assume

a pivotal role.[110] Based on this clue, the cholinergic modulation of LMAN-RA

pathway in females might be implicated in the procedure of their perception and

selection of male songs.

Truthfully, the perception of auditory events in female zebra finches is not

necessarily less competent than that in males.[115, 116, 117] Previous research has verified

that in female canaries, HVC plays a crucial role in the perception of conspecific

song.[118] Further research indicates that the HVC-RA pathway of female zebra finches

can mediate predictive call timing in the same way as that of males, and the adaptive

timing plasticity of females is stronger than that of males.[115] However, considering the

substantial sex dimorphism in the structure and function of HVC-RA pathway, the

mechanisms of song perception in females and males may not be identical. Drawing

on the interaction and relationship between HVC-RA and LMAN-RA pathways of

males, and based on the findings of predecessors and ours, we put forward a

mechanism hypothesis that cholinergic signaling impacts the perception and/or

selection of male songs in adult female zebra finches by regulating song premotor

circuits (Figure 7B). Although cholinergic signaling exerts no influence on HVC-

RAPN excitatory synaptic transmission of adult females, it can reduce LMAN-RAPN

excitatory synaptic transmission via mAChRs, weaken the dependence on the auditory

feedback of LMAN-RA pathway, and thereby relatively strengthen the song

perception (including conspecific songs, and predictive call timing) and/or preference

selection of female HVC-RA pathway for males.

Limitations of the study

While the roles of different cholinergic receptors were unambiguously identified by pharmacological means, the impact of inhibitory interneuron network within RA on RAPNs could not be precluded in in vivo experiments. The cholinergic modulation of RAPNs' inhibitory synaptic afferents and whether it enhances or attenuates the effect of cholinergic signaling on birdsongs require further investigation. Our hypothesis regarding the cholinergic receptor mechanism in the song learning process of juveniles also remains to be verified, and we are currently attempting to undertake this work. Additionally, the hypothesis concerning the significance of cholinergic modulation on song premotor circuits in females still requires elaborately designed behavioral experiments for validation.

SUPPLEMENTAL INFORMATION

Figure S1 Other song acoustic characteristics are not influenced by cholinergic signaling within RA.

ACKNOWLEDGMENTS

This research was funded by the National Natural Science Foundation of China (32160123, 32360222), the Key Project of Natural Science Foundation of Jiangxi Province in China (20212ACB205002), and Jiangxi Province Key Laboratory of Organic Functional Molecules (2024SSY05141).

AUTHOR CONTRIBUTIONS

Investigation, Ning Xu, Yutao Zhang, and Yalun Sun; methodology, Wei Meng and Songhua Wang; data curation, Ning Xu, Yutao Zhang, Yalun Sun, and Xueqing Song; formal analysis, Wei Meng, Ning Xu, YangYang Cao, and Xinqi Yang; writing – original draft, Wei Meng and Songhua Wang; writing – review & editing, Wei Meng and Ning Xu; conceptualization, Wei Meng; resources, supervision, and project administration, Wei Meng; funding acquisition, Wei Meng and Songhua Wang.

DECLARATION OF INTERESTS

The authors declare no competing interests.

INCLUSION AND DIVERSITY

We support inclusive, diverse, and equitable conduct of research.

REFERENCES

1. Wood AN. New roles for dopamine in motor skill acquisition: lessons from primates, rodents, and songbirds. *Journal of neurophysiology* **125**, 2361-2374 (2021).

2. Dayan E, Cohen LG. Neuroplasticity subserving motor skill learning. *Neuron* **72**, 443-454 (2011).

3. Conner JM, Culberson A, Packowski C, Chiba AA, Tuszynski MH. Lesions of the Basal forebrain cholinergic system impair task acquisition and abolish cortical plasticity associated with motor skill learning. *Neuron* **38**, 819-829 (2003).

4. Bariselli S, Mateo Y, Reuveni N, Lovinger DM. Gestational ethanol exposure impairs motor skills in female mice through dysregulated striatal dopamine and acetylcholine function. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* **48**, 1808-1820 (2023).

5. Zhang Y, Zhou L, Zuo J, Wang S, Meng W. Analogies of human speech and bird song: From vocal learning behavior to its neural basis. *Frontiers in psychology* **14**, 1100969 (2023).

6. Veit L, Tian LY, Monroy Hernandez CJ, Brainard MS. Songbirds can learn flexible contextual control over syllable sequencing. *eLife* **10**, (2021).

7. Brainard MS, Doupe AJ. What songbirds teach us about learning. *Nature* **417**, 351-358 (2002).

8. Friedrich SR, Nevue AA, Andrade ALP, Velho TAF, Mello CV. Emergence of sex-specific transcriptomes in a sexually dimorphic brain nucleus. *Cell reports* **40**, 111152 (2022).

9. Nottebohm F, Stokes TM, Leonard CM. Central control of song in the canary, Serinus canarius. *The Journal of comparative neurology* **165**, 457-486 (1976).

10. Vicario DS. Organization of the zebra finch song control system: II. Functional organization of outputs from nucleus Robustus archistriatalis.

The Journal of comparative neurology **309**, 486-494 (1991).

11.   McDonald KS, Kirn JR. Anatomical plasticity in the adult zebra finch
song system. *The Journal of comparative neurology* **520**, 3673-3686
(2012).

12.   Li R, Sakaguchi H. Cholinergic innervation of the song control nuclei by
the ventral paleostriatum in the zebra finch: a double-labeling study
with retrograde fluorescent tracers and choline acetyltransferase
immunohistochemistry. *Brain research* **763**, 239-246 (1997).

13.   Reiner A*, et al.* Revised nomenclature for avian telencephalon and
some related brainstem nuclei. *The Journal of comparative neurology*
473, 377-414 (2004).

14.   Sakaguchi H, Saito N. The acetylcholine and catecholamine contents in
song control nuclei of zebra finch during song ontogeny. *Brain research*
Developmental brain research **47**, 313-317 (1989).

15.   Puzerey PA, Maher K, Prasad N, Goldberg JH. Vocal learning in
songbirds requires cholinergic signaling in a motor cortex-like nucleus.
Journal of neurophysiology **120**, 1796-1806 (2018).

16.   Yu AC, Margoliash D. Temporal hierarchical control of singing in birds.
Science (New York, NY) **273**, 1871-1875 (1996).

17.   Leonardo A, Fee MS. Ensemble coding of vocal control in birdsong.
The Journal of neuroscience : the official journal of the Society for
Neuroscience **25**, 652-661 (2005).

18.   Hahnloser RH, Kozhevnikov AA, Fee MS. An ultra-sparse code
underlies the generation of neural sequences in a songbird. *Nature* **419**,
65-70 (2002).

19.   Meng W, Wang S, Yao L, Zhang N, Li D. Muscarinic Receptors Are
Responsible for the Cholinergic Modulation of Projection Neurons in
the Song Production Brain Nucleus RA of Zebra Finches. *Frontiers in*
cellular neuroscience **11**, 51 (2017).

20.   Pérez SE, Yáñez J, Marín O, Anadón R, González A, Rodríguez-
Moldes I. Distribution of choline acetyltransferase (ChAT)
immunoreactivity in the brain of the adult trout and tract-tracing
observations on the connections of the nuclei of the isthmus. *The*
Journal of comparative neurology **428**, 450-474 (2000).

21.   Clemente D*, et al.* Cholinergic elements in the zebrafish central
844   nervous system: Histochemical and immunohistochemical analysis.
845   *The Journal of comparative neurology* **474**, 75-107 (2004).

847   22.   López JM, Domínguez L, Morona R, Northcutt RG, González A.
848   Organization of the cholinergic systems in the brain of two lungfishes,
849   Protopterus dolloi and Neoceratodus forsteri. *Brain structure & function*
850   **217**, 549-576 (2012).

852   23.   Hoogland PV, Vermeulen-VanderZee E. Distribution of choline
853   acetyltransferase immunoreactivity in the telencephalon of the lizard
854   Gekko gecko. *Brain, behavior and evolution* **36**, 378-390 (1990).

856   24.   Powers AS, Reiner A. The distribution of cholinergic neurons in the
857   central nervous system of turtles. *Brain, behavior and evolution* **41**,
858   326-345 (1993).

860   25.   Pediconi MF, Roccamo de Fernández AM, Barrantes FJ. Asymmetric
861   distribution and down-regulation of the muscarinic acetylcholine
862   receptor in rat cerebral cortex. *Neurochemical research* **18**, 565-572
863   (1993).

865   26.   Tian MK, Bailey CD, Lambe EK. Cholinergic excitation in mouse
866   primary vs. associative cortex: region-specific magnitude and receptor
867   balance. *The European journal of neuroscience* **40**, 2608-2618 (2014).

869   27.   Ghimire M, Cai R, Ling L, Hackett TA, Caspary DM. Nicotinic Receptor
870   Subunit Distribution in Auditory Cortex: Impact of Aging on Receptor
871   Number and Function. *The Journal of neuroscience : the official journal*
872   *of the Society for Neuroscience* **40**, 5724-5739 (2020).

874   28.   Benoy A, Bin Ibrahim MZ, Behnisch T, Sajikumar S. Metaplastic
875   Reinforcement of Long-Term Potentiation in Hippocampal Area CA2 by
876   Cholinergic Receptor Activation. *The Journal of neuroscience : the*
877   *official journal of the Society for Neuroscience* **41**, 9082-9098 (2021).

879   29.   Medina L, Reiner A. Distribution of choline acetyltransferase
880   immunoreactivity in the pigeon brain. *The Journal of comparative*
881   *neurology* **342**, 497-537 (1994).

883   30.   Dietl MM, Cortés R, Palacios JM. Neurotransmitter receptors in the
884   avian brain. II. Muscarinic cholinergic receptors. *Brain research* **439**,
885   360-365 (1988).

31.   Lohmann TH, Torrão AS, Britto LR, Lindstrom J, Hamassaki-Britto DE.
A comparative non-radioactive in situ hybridization and
immunohistochemical study of the distribution of alpha7 and alpha8
subunits of the nicotinic acetylcholine receptors in visual areas of the
chick brain. *Brain research* **852**, 463-469 (2000).

32.   Semba K. Phylogenetic and ontogenetic aspects of the basal forebrain
cholinergic neurons and their innervation of the cerebral cortex.
Progress in brain research **145**, 3-43 (2004).

33.   Ztaou S*, et al.* Involvement of Striatal Cholinergic Interneurons and M1
and M4 Muscarinic Receptors in Motor Symptoms of Parkinson's
Disease. *The Journal of neuroscience : the official journal of the*
Society for Neuroscience **36**, 9161-9172 (2016).

34.   Power AE, Vazdarjanova A, McGaugh JL. Muscarinic cholinergic
influences in memory consolidation. *Neurobiology of learning and*
memory **80**, 178-193 (2003).

35.   Hasselmo ME. The role of acetylcholine in learning and memory.
Current opinion in neurobiology **16**, 710-715 (2006).

36.   Greco MA, McCarley RW, Shiromani PJ. Choline acetyltransferase
expression during periods of behavioral activity and across natural
sleep-wake states in the basal forebrain. *Neuroscience* **93**, 1369-1374
(1999).

37.   Nikonova EV*, et al.* Transcriptional Profiling of Cholinergic Neurons
From Basal Forebrain Identifies Changes in Expression of Genes
Between Sleep and Wake. *Sleep* **40**, (2017).

38.   Sarter M, Lustig C, Blakely RD, Koshy Cherian A. Cholinergic genetics
of visual attention: Human and mouse choline transporter capacity
variants influence distractibility. *Journal of physiology, Paris* **110**, 10-18
(2016).

39.   Orciani C, Hall H, Pentz R, Foret MK, Do Carmo S, Cuello AC. Long-
term nucleus basalis cholinergic depletion induces attentional deficits
and impacts cortical neurons and BDNF levels without affecting the
NGF synthesis. *Journal of neurochemistry* **163**, 149-167 (2022).

40.   Williams VM*, et al.* Distribution of cholinergic neurons in the brains of a
lar gibbon and a chimpanzee. *Anatomical record (Hoboken, NJ : 2007)*

930   **305**, 1516-1535 (2022).

932   41.   Yang D, Günter R, Qi G, Radnikow G, Feldmeyer D. Muscarinic and
933     Nicotinic Modulation of Neocortical Layer 6A Synaptic Microcircuits Is
934     Cooperative and Cell-Specific. *Cerebral cortex (New York, NY : 1991)*
935     **30**, 3528-3542 (2020).

937   42.   Poorthuis RB, Bloem B, Schak B, Wester J, de Kock CP, Mansvelder
938     HD. Layer-specific modulation of the prefrontal cortex by nicotinic
939     acetylcholine receptors. *Cerebral cortex (New York, NY : 1991)* **23**,
940     148-161 (2013).

942   43.   Tsolias A, Medalla M. Muscarinic Acetylcholine Receptor Localization
943     on Distinct Excitatory and Inhibitory Neurons Within the ACC and LPFC
944     of the Rhesus Monkey. *Frontiers in neural circuits* **15**, 795325 (2021).

946   44.   Roberts TF, Hall WS, Brauth SE. Organization of the avian basal
947     forebrain: chemical anatomy in the parrot (Melopsittacus undulatus).
948     *The Journal of comparative neurology* **454**, 383-408 (2002).

950   45.   Ryan SM, Arnold AP. Evidence for cholinergic participation in the
951     control of bird song; acetylcholinesterase distribution and muscarinic
952     receptor autoradiography in the zebra finch brain. *The Journal of*
953     *comparative neurology* **202**, 211-219 (1981).

955   46.   Sadananda M. Acetylcholinesterase in central vocal control nuclei of
956     the zebra finch (Taeniopygia guttata). *Journal of biosciences* **29**, 189-
957     200 (2004).

959   47.   Zuschratter W, Scheich H. Distribution of choline acetyltransferase and
960     acetylcholinesterase in the vocal motor system of zebra finches. *Brain*
961     *research* **513**, 193-201 (1990).

963   48.   Lovell PV, Huizinga NA, Friedrich SR, Wirthlin M, Mello CV. The
964     constitutive differential transcriptome of a brain circuit for vocal learning.
965     *BMC genomics* **19**, 231 (2018).

967   49.   Lovell PV, Clayton DF, Replogle KL, Mello CV. Birdsong
968     "transcriptomics": neurochemical specializations of the oscine song
969     system. *PloS one* **3**, e3440 (2008).

971   50.   Watson JT, Adkins-Regan E, Whiting P, Lindstrom JM, Podleski TR.
972     Autoradiographic localization of nicotinic acetylcholine receptors in the
973     brain of the zebra finch (Poephila guttata). *The Journal of comparative*

neurology **274**, 255-264 (1988).

51.   Asogwa NC, Mori C, Sánchez-Valpuesta M, Hayase S, Wada K. Inter-
        and intra-specific differences in muscarinic acetylcholine receptor
        expression in the neural pathways for vocal learning in songbirds. *The
        Journal of comparative neurology* **526**, 2856-2869 (2018).

52.   Asogwa NC*, et al.* Nicotinic acetylcholine receptors in a songbird brain.
        The Journal of comparative neurology **530**, 1966-1991 (2022).

53.   Conner JM, Kulczycki M, Tuszynski MH. Unique contributions of
        distinct cholinergic projections to motor cortical plasticity and learning.
        Cerebral cortex (New York, NY : 1991) **20**, 2739-2748 (2010).

54.   Li Y, Hollis E. Basal Forebrain Cholinergic Neurons Selectively Drive
        Coordinated Motor Learning in Mice. *The Journal of neuroscience : the
        official journal of the Society for Neuroscience* **41**, 10148-10160 (2021).

55.   Voegtle A*, et al.* Cholinergic modulation of motor sequence learning.
        The European journal of neuroscience, (2024).

56.   Cappendijk SL*, et al.* In vivo nicotine exposure in the zebra finch: a
        promising innovative animal model to use in neurodegenerative
        disorders related research. *Pharmacology, biochemistry, and behavior*
        96, 152-159 (2010).

57.   Perry WM, Cappendijk SL. Effects of nicotine administration on spectral
        and temporal features of crystallized song in the adult male zebra finch.
        *Nicotine & tobacco research : official journal of the Society for
        Research on Nicotine and Tobacco* **16**, 1409-1416 (2014).

58.   Chi Z, Margoliash D. Temporal precision and temporal drift in brain and
        behavior of zebra finch song. *Neuron* **32**, 899-910 (2001).

59.   Tian LY, Warren TL, Mehaffey WH, Brainard MS. Dynamic top-down
        biasing implements rapid adaptive changes to individual movements.
        eLife **12**, (2023).

60.   Shea SD, Margoliash D. Basal forebrain cholinergic modulation of
        auditory activity in the zebra finch song system. *Neuron* **40**, 1213-1226
        (2003).

61.   Shea SD, Koch H, Baleckaitis D, Ramirez JM, Margoliash D. Neuron-
        specific cholinergic modulation of a forebrain song control nucleus.

1018   *Journal of neurophysiology* **103**, 733-745 (2010).

1019

1020   62.   Jaffe PI, Brainard MS. Acetylcholine acts on songbird premotor circuitry
1021   to invigorate vocal output. *eLife* **9**, (2020).

1022

1023   63.   Sakaguchi H, Saito N. Developmental change of cholinergic activity in
1024   the forebrain of the zebra finch during song learning. *Brain research*
1025   *Developmental brain research* **62**, 223-228 (1991).

1026

1027   64.   Sakaguchi H. Developmental changes in carbachol-stimulated
1028   phosphoinositide turnover in synaptoneurosomes of the robust nucleus
1029   of the archistriatum in the zebra finch. *Neuroreport* **6**, 1901-1904 (1995).

1030

1031   65.   Salgado-Commissariat D, Rosenfield DB, Helekar SA. Nicotine-
1032   mediated plasticity in robust nucleus of the archistriatum of the adult
1033   zebra finch. *Brain research* **1018**, 97-105 (2004).

1034

1035   66.   Pfenning AR*, et al.* Convergent transcriptional specializations in the
1036   brains of humans and song-learning birds. *Science (New York, NY)* **346**,
1037   1256846 (2014).

1038

1039   67.   Jarvis ED. Evolution of vocal learning and spoken language. *Science*
1040   *(New York, NY)* **366**, 50-54 (2019).

1041

1042   68.   Meng W, Wang SH, Li DF. Carbachol-Induced Reduction in the Activity
1043   of Adult Male Zebra Finch RA Projection Neurons. *Neural plasticity*
1044   **2016**, 7246827 (2016).

1045

1046   69.   Gulledge AT, Park SB, Kawaguchi Y, Stuart GJ. Heterogeneity of phasic
1047   cholinergic signaling in neocortical neurons. *Journal of neurophysiology*
1048   **97**, 2215-2229 (2007).

1049

1050   70.   Gulledge AT, Stuart GJ. Cholinergic inhibition of neocortical pyramidal
1051   neurons. *The Journal of neuroscience : the official journal of the*
1052   *Society for Neuroscience* **25**, 10308-10320 (2005).

1053

1054   71.   Levy RB, Reyes AD, Aoki C. Nicotinic and muscarinic reduction of
1055   unitary excitatory postsynaptic potentials in sensory cortex; dual
1056   intracellular recording in vitro. *Journal of neurophysiology* **95**, 2155-
1057   2166 (2006).

1058

1059   72.   Gigout S, Jones GA, Wierschke S, Davies CH, Watson JM, Deisz RA.
1060   Distinct muscarinic acetylcholine receptor subtypes mediate pre- and
1061   postsynaptic effects in rat neocortex. *BMC neuroscience* **13**, 42 (2012).

1062

73. McCasland JS. Neuronal control of bird song production. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **7**, 23-39 (1987).

74. Bottjer SW, Brady JD, Cribbs B. Connections of a motor cortical region in zebra finches: relation to pathways for vocal learning. *The Journal of comparative neurology* **420**, 244-260 (2000).

75. Glasgow SD, McPhedrain R, Madranges JF, Kennedy TE, Ruthazer ES. Approaches and Limitations in the Investigation of Synaptic Transmission and Plasticity. *Frontiers in synaptic neuroscience* **11**, 20 (2019).

76. Tryon SC, Bratsch-Prince JX, Warren JW, Jones GC, McDonald AJ, Mott DD. Differential Regulation of Prelimbic and Thalamic Transmission to the Basolateral Amygdala by Acetylcholine Receptors. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **43**, 722-735 (2023).

77. Spiro JE, Dalva MB, Mooney R. Long-range inhibition within the zebra finch song nucleus RA can coordinate the firing of multiple projection neurons. *Journal of neurophysiology* **81**, 3007-3020 (1999).

78. Zemel BM*, et al.* Motor cortex analogue neurons in songbirds utilize Kv3 channels to generate ultranarrow spikes. *eLife* **12**, (2023).

79. Wood WE, Lovell PV, Mello CV, Perkel DJ. Serotonin, via HTR2 receptors, excites neurons in a cortical-like premotor nucleus necessary for song learning and production. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **31**, 13808-13815 (2011).

80. Wood WE, Roseberry TK, Perkel DJ. HTR2 receptors in a songbird premotor cortical-like area modulate spectral characteristics of zebra finch song. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **33**, 2908-2915 (2013).

81. Wang S*, et al.* Dopamine Modulates Excitatory Synaptic Transmission by Activating Presynaptic D1-like Dopamine Receptors in the RA Projection Neurons of Zebra Finches. *Frontiers in cellular neuroscience* **14**, 126 (2020).

82. Solis MM, Perkel DJ. Noradrenergic modulation of activity in a vocal

control nucleus in vitro. *Journal of neurophysiology* **95**, 2265-2276 (2006).

83.   Sizemore M, Perkel DJ. Noradrenergic and GABA B receptor activation differentially modulate inputs to the premotor nucleus RA in zebra finches. *Journal of neurophysiology* **100**, 8-18 (2008).

84.   Wang S*, et al.* Sexual dimorphism of inhibitory synaptic transmission in RA projection neurons of songbirds. *Neuroscience letters* **709**, 134377 (2019).

85.   Herrmann K, Arnold AP. The development of afferent projections to the robust archistriatal nucleus in male zebra finches: a quantitative electron microscopic study. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **11**, 2063-2074 (1991).

86.   Scharff C, Nottebohm F. A comparative study of the behavioral deficits following lesions of various parts of the zebra finch song system: implications for vocal learning. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **11**, 2896-2913 (1991).

87.   Warren TL, Tumer EC, Charlesworth JD, Brainard MS. Mechanisms and time course of vocal learning and consolidation in the adult songbird. *Journal of neurophysiology* **106**, 1806-1821 (2011).

88.   Kao MH, Doupe AJ, Brainard MS. Contributions of an avian basal ganglia-forebrain circuit to real-time modulation of song. *Nature* **433**, 638-643 (2005).

89.   Andalman AS, Fee MS. A basal ganglia-forebrain circuit in the songbird biases motor output to avoid vocal errors. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 12518-12523 (2009).

90.   Aronov D, Andalman AS, Fee MS. A specialized forebrain circuit for vocal babbling in the juvenile songbird. *Science (New York, NY)* **320**, 630-634 (2008).

91.   Bottjer SW, Miesner EA, Arnold AP. Forebrain Lesions Disrupt Development But Not Maintenance of Song in Passerine Birds. *Science (New York, NY)* **224**, 901-903 (1984).

92.   Brainard MS. Contributions of the anterior forebrain pathway to vocal plasticity. *Annals of the New York Academy of Sciences* **1016**, 377-394

93. Nordeen KW, Nordeen EJ. Deafening-induced vocal deterioration in adult songbirds is reversed by disrupting a basal ganglia-forebrain circuit. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **30**, 7392-7400 (2010).

94. Chen Z, Ye R, Goldman SA. Testosterone modulation of angiogenesis and neurogenesis in the adult songbird brain. *Neuroscience* **239**, 139-148 (2013).

95. Nottebohm F, Arnold AP. Sexual dimorphism in vocal control areas of the songbird brain. *Science (New York, NY)* **194**, 211-213 (1976).

96. Brenowitz EA. Testosterone and brain-derived neurotrophic factor interactions in the avian song control system. *Neuroscience* **239**, 115-123 (2013).

97. Wade J, Arnold AP. Sexual differentiation of the zebra finch song system. *Annals of the New York Academy of Sciences* **1016**, 540-559 (2004).

98. Liu XL, Hou GQ, Liao SQ, Li DF. Sexual dimorphism of the electrophysiological properties of the projection neurons in the robust nucleus of the arcopallium in adult zebra finches. *Neuroscience bulletin* **26**, 147-152 (2010).

99. Wang S, Meng W, Liu S, Liao C, Huang Q, Li D. Sex differences of excitatory synaptic transmission in RA projection neurons of adult zebra finches. *Neuroscience letters* **582**, 75-80 (2014).

100. Bottjer SW, Roselinsky H, Tran NB. Sex differences in neuropeptide staining of song-control nuclei in zebra finch brains. *Brain, behavior and evolution* **50**, 284-303 (1997).

101. Sakaguchi H, Li R, Taniguchi I. Sex differences in the ventral paleostriatum of the zebra finch: origin of the cholinergic innervation of the song control nuclei. *Neuroreport* **11**, 2727-2731 (2000).

102. Konishi M, Akutagawa E. Neuronal growth, atrophy and death in a sexually dimorphic song nucleus in the zebra finch brain. *Nature* **315**, 145-147 (1985).

103. Mooney R, Rao M. Waiting periods versus early innervation: the

development of axonal connections in the zebra finch song system. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **14**, 6532-6543 (1994).

104. Holloway CC, Clayton DF. Estrogen synthesis in the male brain triggers development of the avian song control pathway in vitro. *Nature neuroscience* **4**, 170-175 (2001).

105. Diez A, Wang S, Carfagnini N, MacDougall-Shackleton SA. Sex differences in myelination of the zebra finch vocal control system emerge relatively late in development. *Developmental neurobiology* **82**, 581-595 (2022).

106. Johnson F, Sellix M. Reorganization of a telencephalic motor region during sexual differentiation and vocal learning in zebra finches. *Brain research Developmental brain research* **121**, 253-263 (2000).

107. Shaughnessy DW, Hyson RL, Bertram R, Wu W, Johnson F. Female zebra finches do not sing yet share neural pathways necessary for singing in males. *The Journal of comparative neurology* **527**, 843-855 (2019).

108. Wang J, Sakaguchi H, Sokabe M. Sex differences in the vocal motor pathway of the zebra finch revealed by real-time optical imaging technique. *Neuroreport* **10**, 2487-2491 (1999).

109. Nixdorf-Bergweiler BE. Divergent and parallel development in volume sizes of telencephalic song nuclei in male and female zebra finches. *The Journal of comparative neurology* **375**, 445-456 (1996).

110. Nixdorf-Bergweiler BE. Lateral magnocellular nucleus of the anterior neostriatum (LMAN) in the zebra finch: neuronal connectivity and the emergence of sex differences in cell morphology. *Microscopy research and technique* **54**, 335-353 (2001).

111. Nordeen EJ, Grace A, Burek MJ, Nordeen KW. Sex-dependent loss of projection neurons involved in avian song learning. *Journal of neurobiology* **23**, 671-679 (1992).

112. Hamilton KS, King AP, Sengelaub DR, West MJ. A brain of her own: a neural correlate of song assessment in a female songbird. *Neurobiology of learning and memory* **68**, 325-332 (1997).

113. Riebel K, Smallegange IM, Terpstra NJ, Bolhuis JJ. Sexual equality in

zebra finch song preference: evidence for a dissociation between song recognition and production learning. *Proceedings Biological sciences* **269**, 729-733 (2002).

114.   Terpstra NJ, Bolhuis JJ, Riebel K, van der Burg JM, den Boer-Visser AM. Localized brain activation specific to auditory memory in a female songbird. *The Journal of comparative neurology* **494**, 784-791 (2006).

115.   Benichov JI, Benezra SE, Vallentin D, Globerson E, Long MA, Tchernichovski O. The Forebrain Song System Mediates Predictive Call Timing in Female and Male Zebra Finches. *Current biology : CB* **26**, 309-318 (2016).

116.   Kriengwatana B, Spierings MJ, ten Cate C. Auditory discrimination learning in zebra finches: effects of sex, early life conditions and stimulus characteristics. *Animal Behaviour* **116**, 99-112 (2016).

117.   Rouse AA, Patel AD, Wainapel S, Kao MH. Within-species differences in vocal production learning in a songbird are associated with differences in flexible rhythm pattern perception. *bioRxiv*, 2022.2007.2013.499954 (2023).

118.   Brenowitz EA. Altered perception of species-specific song by female birds after lesions of a forebrain nucleus. *Science (New York, NY)* **251**, 303-305 (1991).

119.   Zhang Y*, et al.* BDNF enhances electrophysiological activity and excitatory synaptic transmission of RA projection neurons in adult male zebra finches. *Brain research* **1801**, 148208 (2023).

120.   Mehaffey WH, Doupe AJ. Naturalistic stimulation drives opposing heterosynaptic plasticity at two inputs to songbird cortex. *Nature neuroscience* **18**, 1272-1280 (2015).

121.   Wang S, Liao C, Li F, Liu S, Meng W, Li D. Castration modulates singing patterns and electrophysiological properties of RA projection neurons in adult male zebra finches. *PeerJ* **2**, e352 (2014).

122.   Sizemore M, Perkel DJ. Premotor synaptic plasticity limited to the critical period for song learning. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 17492-17497 (2011).

123.   Meitzen J, Moore IT, Lent K, Brenowitz EA, Perkel DJ. Steroid hormones act transsynaptically within the forebrain to regulate neuronal

1282   phenotype and song stereotypy. *The Journal of neuroscience : the*

1283   *official journal of the Society for Neuroscience* **27**, 12045-12057 (2007).

1284 **Methods**

1285 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, peptides, and recombinant proteins		
Carbachol, CAR	MCE	Cat.# HY-B1208
Atropine sulfate, ATR	MCE	Cat.# HY-B1205
Mecamylamine,MEC	MCE	Cat.# HY-B1395
1,1-dimethyl-4-phenylpiperazinium iodid,DMPP	MCE	Cat.# HY-W009190
Oxotremorine M iodide,OM	MCE	Cat.# HY-101372A
Tetrodotoxin, TTX	Sigma	CAS: 4368-28-9
Picrotoxin, PTX	Sigma	CAS: 124-87-8
Software and algorithms		
Sound Analysis Pro 2011	Wenchan Zhao et al. ,2019	soundanalysispro.com
Cool Edit Pro 2.0	Adobe Systems	N/A
MiniAnalysis 6.0	Synaptosoft	N/A
Clampfit 10.7	Molecular Devices	N/A
Other		
3.5mm Cannula-Single	RWD	Cat.# 62004
4.0mm Injector-Single	RWD	Cat.# 62204

1286

1287 **RESOURCE AVAILABILITY**

1288 **Lead contact**

1289 Further information and requests for data should be directed to and will be fulfilled by

1290 the lead contact, Wei Meng (meng7883@163.com).

1291

1292 **Materials availability**

1293 This study did not generate new unique reagents.

1294

Data and code availability

The original data generated during the current study are available from the lead

contact upon requests.

This paper does not report original code.

Any additional information required to reanalyze the data reported in this work is

available from the lead contact upon requests.

1301

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Ethics statement

All of the work described in this study was approved by the Institutional Animal Care and Use Committee of Jiangxi Science and Technology Normal University (3601020137931).

Animals

The adult male and female zebra finches (Taeniopygia guttata) were obtained from a reputable supplier. These birds, all of which were older than 120 days, had previously been raised in a spacious aviary under a light/dark cycle of 14 hours of light and 10 hours of darkness, at a temperature of 24 □. Data were collected from a total of 81 adult males and 65 females.

Slice preparation

Following deep anesthesia, the brain was carefully extracted. Subsequently, the fresh brain slices were sectioned using a vibrating microtome (7000 smz; Campden Instruments, UK) with a thickness of 250 μm in the coronal plane, and immediately transferred to ice-cold slice solution with an initial pH of 7.3–7.4, osmolarity of 330–340 mOsm, bubbled with 95% O_2 and 5% CO_2 carbogen. The slice solution was composed of 62.5mM NaCl, 5mM KCl, 28 mM $NaHCO_3$, 248 mM sucrose, 1.3mM $MgSO_4 \cdot 7H_2O$, 10mM glucose, and 1.26mM $NaH_2PO_4 \cdot H_2O$. The slices were transferred to a holding chamber containing oxygenated artificial cerebrospinal fluid (ACSF) at 35°C. The ACSF was composed of 125mM NaCl, 2.5mM KCl, 1.2mM $MgSO_4 \square 7H_2O$, 1.27mM $NaH_2PO_4 \square H_2O$, 25mM $NaHCO_3$, 25mM glucose, and 2.0mM $CaCl_2$, pH 7.3-7.4. The slices were incubated for at least 0.5 h and equilibrated to room temperature prior to electrophysiological recording.

Patch-clamp electrophysiology

The visualization of RA in brain slices was achieved using infrared differential interference contrast microscopy (IR-DIC) through a microscope (BX51WIF, Olympus, Tokyo, Japan). Whole-cell patch clamp recordings were conducted at a temperature of approximately 24 °C. Slices were perfused with carbogen-bubbled ACSF at a rate of 1-2 ml/min, while neurons were visualized under a 40× water

immersion lens. Patch pipettes were fabricated from standard borosilicate capillary glass (BF150-117-10, Sutter Instruments, CA, USA) using a Flaming/Brown micropipette puller (P-1000, Sutter Instruments, CA, USA). All recording pipettes had an open-tip resistance ranging from 3.0 to 6.0 MΩ in the bath solution. The patch pipettes were filled with a solution containing the following concentrations: 120 mM KMeSO4, 5 mM NaCl, 10 mM HEPES, 2 mM EGTA, 5 mM QX-314, 2 mM ATP, and 0.3 mM GTP (pH adjusted to range 7.3–7.4; osmolarity set at 340 mOsm). RAPNs exhibit regular spontaneous firing of action potentials and demonstrate slow time-dependent inward rectification of hyperpolarizing current pulses. RA interneurons do not display spontaneously firing action potentials, lack time-dependent inward rectification of hyperpolarizing current pulses, and exhibit irregular spiking in response to depolarizing current injection.[77, 119]

To record the miniature excitatory postsynaptic currents (mEPSCs) of RAPNs, a concentration of 1 µM TTX was applied to ASCF in order to block spontaneous events driven by intrinsic Na+ channel-mediated action potentials. During eEPSCs recordings of RAPNs, a bath application of 150 µM picrotoxin (PTX) was employed to block GABAA receptor-mediated inhibitory synaptic currents. Reported holding potentials have been corrected for corresponding liquid junction potential. The signals underwent sampling at 10 kHz and filtering at 2 kHz using the MultiClamp 700B amplifier (Molecular Devices, CA, USA). EPSCs were evoked through a concentric needle electrode placed in the afferent trace of HVC (as shown in Figure 3A, 5A) or LMAN (as shown in Figure 4A, 5A).[120] The paired-pulse ratio (PPR), which quantifies the relative amplitude of the second pulse compared to the first pulse of evoked excitatory postsynaptic currents, was assessed at an inter-pulse interval of 50 ms. Alterations in PPR reflect involvement of presynaptic mechanisms.

Drug Application

CAR (30 µM); OM (an agonist of mAChRs, 10 µM); ATR (an antagonist of mAChRs, 10 µM); DMPP (an agonist of nAChRs, 10 µM); MEC (an antagonist of nAChRs, 10µM). The effects of these drugs on RAPNs were assessed through bath perfusion.

Song recording

Song recordings were conducted in a recording studio measuring 2.1 × 1.2 × 1 m,

equipped with a TAKSTAR directional microphone (Guangdong Victory Electronics

Co. Ltd., Guangzhou, China; frequency range: 50–20000 Hz) and a glass window (20

× 40 cm). During the recording sessions, a male bird was placed in a cage near the

window within the recording studio and able to see a female bird positioned outside.

Each bird's vocalizations were recorded per day between 8:00 am and 11:00 am. Song

capturing was performed using Cool Edit Pro 2.0 (sampling rate: 44100 Hz; channels:

stereo; resolution: 16-bit).

1375

In vivo microinjection

Birds were anesthetized using isoflurane inhalation (1.5%-2%) and placed in a

stereotaxic apparatus. Cannulas were implanted into bilateral RA (M/L: ± 2.47; A/P:

0.7; D/V: left 2.8, right 3) (n = 16 birds). After birds recovered from surgery,

microjection injectors were inserted into cannulas and connected to a syringe pump

(R462, RWD, Guangdong, China) through flexible tubing. Outflow was continually

monitored throughout the duration of the experiment. Solutions were exchanged to

either phosphate buffered saline (PBS, for control experiments) or 1 mM CAR; 1 mM

OM; 1 mM DMPP, after a two to three-hour baseline period (flow rate held constant

at 1-1.5 µL/ min; solutions exchanged at the same time each day across experiments).

1386

Analysis of song features

We quantified song stereotypy by calculating the similarity to own songs and entropy

of the motifs using Sound Analysis Pro 2011. Thirty or Sixty motifs within the song

were used to assess the percentage similarity and entropy in Pre and during drug

administration.Higher similarity and lower entropy indicate greater stereotypy.[121, 122,

123]

1393

Quantification and statistical analysis

The data were acquired using Clampfit 10.7 through a Digidata 1550B (Molecular

Devices, CA, USA) with 10 kHz sampling frequency. Analysis of mEPSCs was

performed using Clampfit 10.7, MiniAnalysis 6.0 and Origin Pro 8.0 (OriginLab, MA,

USA). The average values of inter-event interval (IEI) and amplitude for events in the

control and drug-administered groups were compared. Paired t-tests were conducted

to statistically compare the mean averaged values for each group, unless otherwise

specified. The significance levels of IEI and amplitude shifts were determined by
employing the nonparametric Kolmogorov-Smirnoff (K-S) test to assess cumulative
probability distributions. The data are presented as mean ± SEM. The software tools
employed for the creation and analysis of experimental diagrams were Adobe
Illustrator 2021 (Adobe, CA, USA), Origin Pro 8.0, and Graphpad Prism 9.5
(GraphPad Software, CA, USA).

1436

1437

Figure 1 Cholinergic modulation of RAPNs' mEPSCs in adult male zebra finches

(A) Experiment schematic. In vitro whole-cell patch clamp recording of RAPNs' mEPSCs in adult male zebra finches.

(B) Top, example traces of adult male RAPNs' mEPSCs in Pre (gray) and CAR (blue); Bottom, quantification of mEPSCs' frequency (left) and amplitude (right) in Pre and CAR, n = 9.

(C-F) Similar to (B). OM (C), n = 8; DMPP (D), n = 9; CAR+ATR (E), n = 7; CAR+MEC (F), n = 8. ** $p \leq 0.01$, *** $p \leq 0.001$, paired t-test. Plots represent mean ± SEM.

1447

Figure 2 Cholinergic modulation of RAPNs' mEPSCs in adult female zebra finches

(A) Experiment schematic. In vitro whole-cell patch clamp recording of RAPNs' mEPSCs in adult female zebra finches.

(B) Top, example traces of adult female RAPNs' mEPSCs in Pre (black) and CAR (red); Bottom, quantification of mEPSCs' frequency (left) and amplitude (right) in Pre and CAR, n = 9.

(C-F) Similar to (B). OM (C), n = 8; DMPP (D), n = 9; CAR+ATR (E), n = 9; CAR+MEC (F), n = 8. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, paired t-test. Plots represent mean ± SEM.

1458

Figure 3 Cholinergic modulation of HVC-RAPN eEPSCs in adult male zebra finches

(A) Experiment schematic. In vitro whole-cell patch clamp recording of HVC-RAPN eEPSCs in adult male zebra finches.

(B) Top, example traces of adult male HVC-RAPN eEPSCs in Pre (black) and CAR (red). Quantification of eEPSCs' amplitude in Pre and CAR, n = 10; Bottom, example traces of eEPSCs in paired pulse stimulation. Quantification of paired-pulse ratio in Pre and CAR, n = 7.

(C) Similar to (B). OM: Top, n = 8; Bottom, n = 7.

(D-F) Example traces of adult male HVC-RAPN eEPSCs in Pre (black), DMPP (D, red), CAR+ATR (E, red) and CAR+MEC (F, red). Quantification of eEPSCs' amplitude in Pre, DMPP (D, n = 11), CAR+ATR (E, n = 8), and CAR+MEC (F, n = 11). * $p \leq 0.05$, *** $p \leq 0.001$, paired t-test. Plots represent mean ± SEM.

1472

Figure 4 Cholinergic modulation of LMAN-RAPN eEPSCs in adult male zebra finches

(A) Experiment schematic. In vitro whole-cell patch clamp recording of LMAN-RAPN eEPSCs in adult male zebra finches.

(B) Top, example traces of adult male LMAN-RAPN eEPSCs in Pre (gray) and CAR (blue). Quantification of eEPSCs' amplitude in Pre and CAR, n = 9; Bottom, example traces of eEPSCs in paired pulse stimulation. Quantification of paired-pulse ratio in Pre and CAR, n = 8.

(C) Example traces of adult male LMAN-RAPN eEPSCs in Pre (gray) and OM (blue). Quantification of eEPSCs' amplitude in Pre and OM, n = 8.

(D) Similar to (B). DMPP: Top, n = 8; Bottom, n = 8.

(E-F) Similar to (C). CAR+ATR (E), n = 10; CAR+MEC (F), n = 8. $* p \le 0.05$, $** p \le 0.01$, $*** p \le 0.001$, paired t-test. Plots represent mean $\pm$ SEM.

Figure 5 Cholinergic modulation of HVC-RAPN and LMAN-RAPN eEPSCs in adult female zebra finches

(A) Experiment schematic. In vitro whole-cell patch clamp recording of HVC-RAPN and LMAN-RAPN eEPSCs in adult female zebra finches.

(B) Example traces of adult female HVC-RAPN eEPSCs in Pre (black) and CAR (red). Quantification of eEPSCs' amplitude in Pre and CAR, n = 10.

(C-D) Similar to (B). DMPP (C), n = 8; OM (D), n = 9.

(E) Top, example traces of adult female LMAN-RAPN eEPSCs in Pre (gray) and CAR (blue). Quantification of eEPSCs' amplitude in Pre and CAR, n = 10; Bottom, example traces of eEPSCs in paired pulse stimulation. Quantification of paired-pulse ratio in Pre and CAR, n = 7.

(F) Similar to (E). OM: Top, n = 8; Bottom, n = 7.

(G) Example traces of adult female LMAN-RAPN eEPSCs in Pre (gray) and DMPP (blue). Quantification of eEPSCs' amplitude in Pre and DMPP, n = 8.

(H-I) Similar to (G). CAR+ATR (H), n = 10; CAR+MEC (I), n = 9. $* p \le 0.05$, $** p \le 0.01$, $*** p \le 0.001$, paired t-test. Plots represent mean $\pm$ SEM.

Figure 6 Cholinergic signaling within RA enhances song stability of adult male zebra finches via mAChRs rather than nAChRs

(A) Experiment schematic. In vivo pharmacological microinjection onto bilaterally RA and song recording.

(B) Left, a free-moving adult male zebra finch in a recording studio with cannulas bilaterally implanted over RA; Right, cannula tracks in a coronal section.

(C) Example spectrograms in Pre and CAR, OM, DMPP or PBS. 'iabcde' indicates the syllables constituting a song motif.

(D) Similarity to own songs in Pre and CAR (black, n = 4), Pre and OM (red, n = 4), Pre and DMPP (gray, n = 4), Pre and PBS (blue, n = 4).

(E) Song entropy in Pre and CAR (black, n = 4), Pre and OM (red, n = 4), Pre and DMPP (gray, n = 4), Pre and PBS (blue, n = 4). $** p \le 0.01$, $*** p \le 0.001$, paired t-test. Plots represent mean $\pm$ SEM.

Figure 7 Mechanism hypotheses on the cholinergic modulation of song premotor
circuits in male and female zebra finches
(A) A hypothesis regarding the mechanism through which cholinergic signaling
influences song production by regulating song premotor circuits in male zebra finches.
(B) A hypothesis regarding the mechanism through which cholinergic signaling
impacts the perception and/or selection of male songs by regulating song premotor
circuits in adult female zebra finches. Details in "DISCUSSION".