Olfactory Mucosa-Derived Mesenchymal Stem Cells Differentiate Towards a Schwann Cell-Like Phenotype Towards Sourcing for Peripheral Nerve Regeneration

Katelyn Neuman[1], Abigail N. Koppes[1,2,3], and Ryan A. Koppes[1,2]

[1]Department of Chemical Engineering, Northeastern University, Boston, Massachusetts, 02115, USA. [2]Department of Bioengineering, Northeastern University, Boston, Massachusetts, 02115, USA. [3]Department of Biology, Northeastern University, Boston, Massachusetts, 02115, USA.

Mesenchymal stem cells (MSCs) are a promising source of stem cells for treating peripheral nerve injuries. Here, we present the first investigation of differentiation of olfactory mucosa-derived MSC (OM-MSC) towards a Schwann Cell (SC)-like phenotype. OM-MSCs are an advantageous potential source of SCs for peripheral nerve repair, as isolation can be accomplished with a minimally invasive procedure compared to autologous nerve harvest and isolation. Here, Schwann Cell Conditioned Media (SCCM) or a defined growth factor supplemented media (GF) was applied to OM-MSC for twenty-one days. The differentiation process and resulting populations were characterized by immunocytochemistry and RT-qPCR. Functionality of differentiated populations was assessed in an *in vitro* co-culture model to evaluate interaction with sensory neurons (dorsal root ganglia) juxtaposed to native SCs. Compared to undifferentiated MSCs, differentiation protocols resulted in significant changes in morphology, gene expression, and functionality using SCCM and GF media, representing key characteristics of SCs. Specifically, differentiated populations exhibit elongated, spindle-like morphologies, a high degree of eccentricity, increased S-100, CD44, and NGF expression, and colocalization of myelin basic proteins with neurites in the co-culture model. In conclusion, this work highlights the potential of OM-MSCs to be expanded and differentiated to SCs to improve synthetic scaffolds or for use in decellularized allografts for nerve repair.

Injuries to the peripheral nervous system are often caused by trauma to the extremities. Leading causes include car accidents, gunshot injuries, and stretching/crush injuries, with a high (83%) prevalence in people under the age of 55[1]. Peripheral nerve injuries (PNI) often result in loss of sensation, function, pain, and paralysis[2,3]. Despite current treatment options, many patients suffer from chronic pain and a reduced functional capacity[3-6]. Surgical interventions are needed for large gap injuries (nerve gaps greater than 3 cm). The gold standard of treatment remains to be the implantation of a nerve autograft harvested from the sural nerve to bridge the gap between the distal and proximal nerve stumps. The fundamental justification for the lack of success of engineered grafts on the market is the lack of supportive Schwann Cells (SCs). Without SCs in injury gaps greater than 1 cm, axons do not regenerate sufficiently across the gap, and patients never see a functional recovery. In large gap injuries, the native SCs (from the distal/proximal nerve stumps) become stressed or exhausted, unable to keep up with the high proliferation demand[7]. These cells then negatively affect the microenvironment, impinging regeneration by releasing inflammatory cytokines[7]. Further, autografting is undesirable, as it requires secondary surgery, damaging a .

secondary nerve and increasing the patient's risk for infection. Therefore, an abundant source of a patient's SCs is paramount to the success of new repair technologies.

Decellularized cadaver tissue allografts provide topographical cues similar to that of the native peripheral nerve extracellular matrix (ECM) and are growing in popularity for small gap injuries[8]. While removing the need for secondary surgery, these allografts have no support cells, including SCs. SCs reside in the peripheral nerve and are fundamental in forming and repairing damaged tissue, producing neurotrophic factors, and remyelinating[9]. In the presence of injury, the phenotypes of SCs alter from their myelinated state to a progenitor-like cell[10-12]. This change promotes SCs to secrete neurotrophic factors, clear damaged myelin, express adhesive and axonal cues, initiate an inflammatory response, and proliferate[10]. These characteristics support nerve repair and regeneration after an injury has occurred. While beneficial, SCs are also invasive to retrieve - with similar drawbacks of harvesting an autograft, requiring a nerve biopsy, which often results in donor site morbidity[13]. Because SCs are limited, the cells are cultured and expanded in vitro for extended periods[14]. When sufficient cells are obtained, these cells can be delivered to the injury site in combination with a scaffold material to facilitate nerve regeneration.

Schwann-like cells derived from stem cells have been investigated as an alternative. Recent research has focused on mesenchymal stem cells (MSCs) collected from different tissue sources such as adipose tissue, human dental pulp, bone marrow, olfactory neuroepithelial tissue, and skeletal muscle tissue. MSCs are attractive as support cell options for peripheral nerve repair because of their anti-inflammatory properties and ability to differentiate into neural tissue[15,16]. Mesenchymal stem cells (MSCs), both undifferentiated and differentiated, have been demonstrated to modulate the immune response in ongoing and completed clinical trials and are thought to be immune-privileged, allowing for allogeneic use[17-20].

Fig. 1. Characteristic immunocytochemistry of OM-MSCs, isolated from rat tissue, display flat, fibroblast-like morphology typical of mesenchymal cells. (A) Green fluorescence denotes S100β, (B) Red fluorescence denotes Nestin, and (C) Displays both S100β and Nestin expression merged with cell nuclei (shown in blue, counterstained with DAPI). (D) Orange fluorescence denotes CD105, (E)Purple fluorescence denotes βIII tubulin, and (F) Displays both CD105 and βIII Tubulin expression merged with cell nuclei (shown in blue, counterstained with DAPI). Scale Bar = 200 μm.

Though these advantages of MSCs have been reported, the original tissue location, donor age, culture conditions, and isolation method must be investigated to predict cell behavior 15. The most widely used MSCs are derived from bone marrow (BM-MSCs) and adipose tissue (AD-MSCs). BM-MSCs have been shown to support nerve regeneration[21,22]. However, the procedure for obtaining bone marrow-derived stem cells (BMSCs) is invasive and painful, and it has a low yield of stem cells[21,23]. AD-MSCs are more abundant and less invasive to retrieve, although they have not been reported to express neural/glial genes[24,25]. Therefore, BM-MSCs and AD-MSCs may not be ideal for PNIs.

In this work, we utilized olfactory mucosa-derived mesenchymal stem cells (OM-MSCs). Due to their unique embryology (derived from neural crest vs. mesoderm) and niche (or original tissue location), we hypothesize that OM-MSCs have a higher potential of differentiation towards an SC-like phenotype and an increased tendency to promote peripheral nerve regeneration[26-28]. These cells have a high

.

proliferation rate and are easily accessible from human patients with a minimally invasive outpatient procedure[29-31]. This study aims to investigate, for the first time, the differentiation of OM-MSCs towards an SC-like cell using Schwann Cell Conditioned Media (SCCM) and a growth factor cocktail (GF). This study uses immunocytochemistry, flow cytometry, and RT-qPCR to characterize the differentiation process and resulting populations. The ability to support neurite extension and exhibit the first steps towards myelination in vitro highlight the utility of OM-MSCs in driving an SC phenotype for the treatment of PNI.

Results

Immunocytochemistry Characterization
ICC characterization of freshly isolated OM-MSC exhibits an expected flat, fibroblast-like morphology typical of mesenchymal stem cells. OM-MSCs have robust expressions of S100β, cluster of differentiation 105 (CD105), and actin filaments (Figure 1). Most OM-MSCs express Nestin, and

Fig. 2. Characteristic flow cytometry of OM-MSCs. (A-C) Display gating strategy employed prior to analysis. The selection of (A) cell population and removal of debris, (B) single cells, and then (C) live cell population is shown. (D-G) Display the distribution of positive and negative cells for the four markers of interest, (D) CD45, (E) CD44, (F) CD90, and (G) nestin. The gray histogram represents the negative control, and the blue histogram represents the cells of interest. (H-I) Display the co-localization of (H) CD44 and nestin expression, (I) CD90 and nestin expression, and (J) CD44 and CD90 expression.

all cells appear to have low-level expression of βIII tubulin and GFAP (Supplementary Fig. 1). SCs (Figure 3) and astrocyte (Supplementary Fig. 2) populations for the following experiments were determined to be homogenous.

OM-MSCs Express Mesenchymal Stem Cell Markers.

OM-MSCs isolated from female adult Sprague Dawley rats were assessed in a Beckman Coulter CytoFLEX for markers outlined in Table 2. Cell populations were selected by removing debris, doublets, and dead cells before analysis (Figure 2A-C). Less than 1% of OM-MSCs expressed cluster of differentiation 45 (CD45), a hemopoietic stem cell marker and negative control (Figure 2D). 99.8% and 94% of OM-MSCs expressed mesenchymal stem cell markers .

CD44 and CD90, respectively (Figure 2E-F). 60% of OM-MSCs expressed Nestin, a marker of stemness (Figure 2G). All cells that expressed Nestin were also CD44$^+$ (Figure 2H). <3% of cells that were Nestin$^+$, did not express CD90 (Figure 2J). Similar results were recorded for OM-MSCs isolated from male Sprague Dawley rats (Supplementary Fig. 3).

Differentiated OM-MSCs Resemble SCs.

Cultured for 21 days in both SCCM and GF media, dOM-MSCs exhibited cell morphology and Nestin expression changes. The dOM-MSCs retained S100β expression (Figure 2). Other media formulations were then evaluated to optimize the differentiation process. OM-MSCs did not

Fig. 3. Differentiated OM-MSC morphology resembles the small, spindle shape of SCs. (A) 21-day differentiation timeline of OM-MSCs. Created with Biorender.com (B-D) Representative images of SCs with (B) denoting S100β (green fluorescence), (C) denoting nestin (red fluorescence), and (D) displaying both S100β and nestin expression merged with cell nuclei (shown in blue, counterstained with DAPI). (E-G) Representative images of dOM-MSCs with SCCM with (E) denoting S100β (green fluorescence), (F) denoting nestin (red fluorescence), and (G) displaying both S100β and nestin expression merged with cell nuclei (shown in blue, counterstained with DAPI). (H-J) Representative images of dOM-MSCs with GF media with (H) denoting S100β (green fluorescence), (I) denoting nestin (red fluorescence), and (J) displaying both S100β and nestin expression merged with cell nuclei (shown in blue, counterstained with DAPI). Scale bar = 100 μm.

differentiate when subjected to a previously published chemical induction protocol applied to BM-MSCs 39. Also, less forskolin in the growth factor media resulted in a reduction in morphology changes and Nestin expression at day 14 (Supplementary Fig. 4). Cell Profiler™ was used to quantify these changes in cell morphology and compare them directly to undifferentiated OM-MSCs and SCs. The cell area of SCs, OM-MSCs, dOM-MSCs (SCCM), and dOM-MSCs (GF) were measured to be 490 ± 270 μm^2, 3900 ± 2500 μm^2, 400 ± 210 μm^2, and 1000 ± 410 μm^2

respectively (Figure 4A). All groups were significantly different from each other with all comparisons, having a p-value < 0.0001, except SC vs. dOM-MSCs (SCCM, p = 0.009). The eccentricity was also measured. SCs, OM-MSCs, dOM-MSCs (SCCM), and dOM-MSCs (GF) were estimated to be 0.90 ± 0.11, 0.76 ± 0.14, 0.81 ± 0.16, and 0.88 ± 0.12, respectively (Figure 4B). Again, all groups were significantly different, with a p-value < 0.0001, apart from SC vs dOM-MSCs (GF, p = 0.003). The median radius of each cell body of SCs, OM-MSCs, dOM-

.

Fig. 4. Changes in cell shape in differentiated OM-MSCs (dOM-MSCs) for both SCCM and GF media conditions were compared to SCs and OM-MSCs using Cell Profiler40. Results for (A) cell area, (B) eccentricity, (C) median cell radius, and (D) cell perimeter. Ten images were analyzed for each condition (n = 190, 210, 1434, 1621). A Krustal-Wallis test (non-parametric ANOVA) with Dunn's multiple comparisons test was used to assess significance. ** = p< 0.01 and **** = p<0.0001. Error bars display the standard deviation.

MSCs (SCCM), and dOM-MSCs (GF) measured to be 2.1 ± 0.6 μm, 5.4 ± 2.2 μm, 2.0 ± 0.5 μm, and 2.8 ± 0.6 μm, respectively (Figure 4C). SCs and dOM-MSCs (SCCM) were not significantly different from each other (p >0.9999). All others were significantly different from each other (p < 0.0001). Finally, the cell perimeter was also measured with SCs, OM-MSCs, dOM-MSCs (SCCM), and dOM-MSCs (GF) having perimeters of 160 ± 87 μm, 710 ± 410 μm, 140 ± 79 μm, and 240 ± 99 μm, respectively (Figure 4D). SCs and dOM-MSCs (SCCM) were not significantly different (p = 0.32). All others were significantly different from each other (p < 0.0001).

Gene expression of dOM-MSCs in SCCM resembles that of SCs.

.

Relative to undifferentiated OM-MSCs, the degree of expression of ten different genes across five experimental groups was evaluated with RT-qPCR (Figure 5). Relative expression of CD44 was upregulated in DRG (log2fold change of 4.5 ± 3.1) compared to OM-MSCs (log2fold change of 0.0 ± 0.5, p = 0.03) and dOM-MSCs (GF, log2fold change of -1.2 ± 2.5, 0.005, Figure 5A). No significant differences in relative gene expression of CD90 across all groups were measured (Figure 5B). The relative expression of GDNF was significantly downregulated in astrocytes (log2fold change of -13.4 ± 4.3) compared to all other groups (p < 0.0001, Figure 5C). GFAP was upregulated in astrocytes (log2fold change of 13.1 ± 0.8), which was significantly more than DRG (log2fold change of 4.2 ± 2.3, p = 0.0003), OM-MSCs (log2fold change of 0.0 ± 2.1, p < 0.0001), dOM-MSCs (GF, log2fold change of 6.0 ± 4.2, p = 0.004), and dOM-MSCs (SCCM, log2fold

Fig. 5. Relative gene expression of dOM-MSCs in SCCM closely resembles that of SCs. Relative log2fold changes in gene expression of (A) CD44, (B) CD90, (C) GDNF, (D) GFAP, (E) NGF, (F) P75NTR, (G) S100β, (H) SOX10, (I) Tubb3, and (J) VEGF. Gene expression relative to Olfactory Mucosa-derived Mesenchymal Stem Cells (OM-MSCs) was compared across five experimental groups: Dorsal Root Ganglia, Astrocytes, Schwann Cells (SCs), differentiated OM-MSCs (SCCM), and differentiated OM-MSCs (GF). Four independent samples were assessed per condition. Significance was assessed with a one-way ANOVA and Tukey's multiple comparison test. Significance is denoted with *, **, ***, and **** which represent p values of < 0.05, <0.01, <0.001, and <0.0001. Error bars indicate standard deviation.

change of -6.8 ± 1.5, $p < 0.0001$). No significant difference in relative gene expression in GFAP was observed between astrocytes and SCs (log2fold change of 9.0 ± 0.7, $p = 0.16$).

Compared to OM-MSCs, relative GFAP expression was upregulated in SCs ($p = 0.0003$) and dOM-MSCs (GF, $p = 0.02$). Relative GFAP expression was significantly downregulated in dOM-MSCs (SCCM) compared to DRG ($p < 0.0001$), SCs ($p < 0.0001$), OM-MSCs ($p = 0.006$), and dOM-MSCs (GF, $p < 0.0001$, Figure 5D). Relative expression of NGF was observed to be downregulated in DRG (log2fold change of -3.6 ± 2.3, $p = 0.03$), dOM-MSCs (SCCM, log2fold change of -6.8 ± 1.5, $p < 0.0001$), dOM-MSCs (GF, log2fold change of -4.2 ± 1.2, $p = 0.01$) compared to OM-MSCs (log2fold change of 0.0 ± 1.2). When compared to SCs, the relative expression of NGF was

significantly downregulated in DRG ($p = 0.003$), dOM-MSCs (SCCM, $p < 0.0001$), and dOM-MSCs (GF, $p = 0.001$). NGF was significantly upregulated in astrocytes (log2fold change of -0.9 ± 1.4) compared to dOM-MSCs (SCCM, $p = 0.0003$, Figure 5E).

P75NTR was upregulated in dOM-MSCs (log2fold change of 0.7 ± 1.6) compared to DRG (log2fold change of -3.6 ± 0.4, $p = 0.02$) and SCs (log2fold change of -3.1 ± 2.4, $p = 0.05$, Figure 5F). No significant differences in relative gene expression of S100β were observed, although SCs and dOM-MSCs (SCCM) had similar levels of relative expression, with log2fold changes of 4.5 ± 1.1 and 4.2 ± 2.8 respectively ($p > 0.9999$, Figure 5G). SOX10 was upregulated in SCs (log2fold change of 13.3 ± 1.7), dOM-MSCs (SCCM, log2fold change of 11.8 ± 2.9), and dOM-MSCs (GF, log2fold change of 9.4 ± 3.9) compared to OM-

Fig. 6. Differentiated OM-MSCs exhibit co-localized expression of MBP on DRG neurons comparable to SCs. (A) 21-day experimental timeline. Created with Biorender.com. (B-D) Representative images of SCs with dissociated DRG with (B) denoting neurofilament-heavy (NF-H; green fluorescence), (C) denoting myelin basic protein (MBP; red fluorescence), and (D) displaying both NF-H and MBP expression merged with cell nuclei (shown in blue, counterstained with DAPI). (E-G) Representative images of OM-MSCs with dissociated DRG with (E) denoting NF-H (green fluorescence), (F) denoting MBP (red fluorescence), and (G) displaying both NF-H and MBP expression merged with cell nuclei (shown in blue, counterstained with DAPI). (H-J) Representative images of dOM-MSCs (SCCM) with dissociated DRG with (H) denoting NF-H (green fluorescence), (I) denoting MBP (red fluorescence), and (J) displaying both NF-H and MBP expression merged with cell nuclei (shown in blue, counterstained with DAPI). (K-M) Representative images of dOM-MSCs (GF) with dissociated DRG with (K) denoting NF-H (green fluorescence), (L) denoting MBP (red fluorescence), and (M) displaying both NF-H and MBP expression merged with cell nuclei (shown in blue, counterstained with DAPI). Scale bar = 100 μm. (N) Displays the percent of neurites that had co-localized expression of MBP. (O) Displays the Mander's coefficient for each of the four experimental groups. Three independent trials were conducted with at least 36 images analyzed for each condition (n = 36, 49, 29, 49). A Krustal-Wallis test (non-parametric ANOVA) with Dunn's multiple comparisons test was used to assess significance. ** = p< 0.01 and *** = p<0.001. Error bars display the standard deviation.

MSCs (log2fold change of 0.0 ± 3.2, p = 0.0003, 0.001, 0.008). The upregulation of SOX10 expression observed in SCs and dOM-MSCs (SCCM) was also significantly different than the relative expression level in astrocytes (log2fold change of 2.0 ± 5.4, p = 0.002, 0.006, Figure 5H). DRG were observed to have a significant upregulation in the relative expression of Tubb3 (log2fold change of 4.5 ± 3.1) compared to astrocytes (log2fold change of -1.5 ±1.6, p = 0.002), SCs (log2fold change of 0.1 ± 0.8, p = 0.02), OM-MSCs (log2fold change of 0.0 ± 1.1, p = 0.02), and dOM-MSCs (GF, log2fold change of -1.0 ± 1.2, p = 0.004). A trending up regulation of Tubb3 expression was also observed in DRG compared to dOM-MSCs (SCCM, log2fold change of 0.6 ± 1.7, p = 0.05, Figure 5I). VEGF was significantly upregulated in SCs (log2fold change of 6.7 ± 2.3) when compared to the relative expression levels in astrocytes (log2fold change of -3.1 ± 1.9, p = 0.005) and DRG (log2fold change of -3.6 ± 0.4, p = 0.003). VEGF was also significantly upregulated in dOM-MSCs (SCCM, log2fold change of 7.6 ± 2.7) when compared to astrocytes (p = 0.002), DRG (p = 0.001), and OM-MSCs (log2fold change of 0 ± 4.7, p = 0.04, Figure 5J).

Differentiated OM-MSCs exhibit Co-Localized Expression of MBP with extending Neurites from dissociated sensory neurons.

The ability of the differentiated cells to perform a crucial function of SCs and myelinate DRG was assessed (Figure 6). Representative images of each of the four groups assessed, SCs, OM-MSCs, dOM-MSCs (SCCM), and dOM-MSCs (GF) are displayed in Figure 6B-M. The percent of MBP overlapping NF-H of each was measured to be 67 ± 17%, 47 ± 23%, 64 ± 19%, and 65 ± 20%, respectively. The groups with SCs, dOM-MSCs (SCCM), and dOM-MSCs (GF) as support cells were all significantly different from those with OM-MSCs (p = 0.0002, 0.003, 0.0006). Those three groups had no significant differences (p > 0.9999, Figure 6N). The Mander's coefficient of each group was also measured and found to be 0.68 ± 0.17, 0.48 ± 0.23, 0.64 ±.19, and 0.65 ± 0.20, respectively. Like the previous analysis, the groups with SCs, dOM-MSCs (SCCM), and dOM-MSCs (GF) as support cells were all significantly different from those with OM-MSCs (p = 0.0001, 0.004, 0.0008). There were no significant differences between SCs, dOM-MSCs (SCCM), and dOM-MSCs (GF, p >0.9999, Figure 6O).

Discussion

The olfactory system, which is responsible for our sense of smell, has uniquely evolved to continually support nerve regeneration due to its constant exposure to the outside .

environment[46-49]. This constant neurogenesis is fueled by stem cells in the surrounding tissue[50]. Multipotent stem cells that reside in the lamina propria of the olfactory mucosa have been identified previously as OM-MSCs due to their predicted neuroectoderm embryonic origin[31,50-53]. Isolating the lamina propria of the olfactory mucosa yielded plastic-adherent cells with fibroblast-like morphology (Figure 1, Supplementary Fig. 1), typical of MSCs from various sources and species[54-57]. OM-MSCs exhibited expression of three known MSC stem cell markers, CD44, CD90, and CD105, but not a hematopoietic stem cell marker, CD45 (Figure 1, Figure 2)[51,54,56,58-61]. CD44 is a cell surface receptor for glycosaminoglycan hyaluronan that mediates cell adhesion to the extracellular matrix[62]. CD90 (Thy-1) is a cell membrane glycoprotein that may play a role in MSCs self-renewal and differentiation[63,64]. CD105 (endoglin) is another cell membrane protein component of the receptor complex of transforming growth factor-beta. This protein is involved in cell proliferation, differentiation, and migration[65]. Lastly, CD45 is a cell surface marker of tyrosine phosphatase, an exclusive marker of hematopoietic lineage cells[66]. Here, our isolated OM-MSCs were confirmed to be characteristically MSCs.

In addition to the traditional MSCs, we identified the expression of several neural and glial cell-specific proteins expressed by the OM-MSCs before differentiation. There was a low-level expression of βIII tubulin (also known as Tuj-1), a standard neuronal marker, a microtubule found in the cytoskeletal of developing and regenerating axons in undifferentiated OM-MSCs (Figure 1)[67]. GFAP was also minimally expressed in OM-MSCs (Supplementary Fig. 1). GFAP is a known marker for mature astrocytes, one type of central nervous system glial cell[68]. Strong S100β and P75NTR expression was observed in the undifferentiated OM-MSCs, two common SC markers (Figure 1, Figure 5F)[69,70]. S100β and other S100 proteins involved in calcium binding regulate various cellular functions[71]. In the peripheral nervous system, P75NTR is an NGF receptor crucial for regulating axonal growth and myelination, among its other functions[72,73]. Using flow cytometry and immunocytochemistry, we also observed that most OM-MSCs are made of Nestin[+] (Figure 1, Figure 2). As a neuroepithelial stem cell protein, Nestin is a cytoskeletal intermediate filament thought to play a role in stemness and differentiation, particularly for neural crest-derived stem cells[74,75]. Previous work has also demonstrated that OM-MSCs express a wide variety of neuronal/glia markers such as S100A4, GFAP, P75NTR, MAP2, βIII tubulin, Nestin, NGF, and GDNF[51,55,58-60,76]. Notably, our population of Nestin[+] cells was substantially lower than that previously identified by Lindsay et al. (100%) and Delorme et al. (100%)[29,31]. The decrease in expression may be due to differences in cell culture conditions. Specifically, serum in

cell media inhibits Nestin expression or the use of earlier passage cells[77,78]. The expression of these neural-lineage-specific genes highlight the potential for OM-MSCs to be an ideal source of SC-like cells to improve peripheral nerve repair.

Undifferentiated OM-MSCs have been demonstrated to have neurodegenerative properties. Conditioned media and OM-MSCs enhanced the myelination of motor neurons in vitro; BM-MSCs, on the other hand, did not[79]. A clinical trial that transplanted whole olfactory mucosa biopsies (containing OM-MSCs and olfactory ensheathing cells) to lesion sites in the spinal cord observed recovering sensory and motor function[80]. We sought to differentiate OM-MSCs toward an SC-like cell to improve injury response. The multipotency of OM-MSCs has been demonstrated previously by inducing differentiation towards non-neuronal lineages such as liver, heart, skeletal muscle, bone, and adipose tissue.

OM-MSCs have also been demonstrated to differentiate towards neurons and central nervous system glia, astrocytes, and oligodendrocytes[50,81]. To our knowledge, only one other study investigated differentiating OM-MSCs towards an SCs-like cell using a fibrin matrix to differentiate them, which was defined by an increase in MBP expression in vitro[76]. We aimed to improve upon this by utilizing defined media components to differentiate these cells and assess their ability to function as SCs with a co-culture of sensory neurons.

Recent work has shown that conditioned media from SCs can, in varying conditions, modify angiogenesis, modify cell survival, increase neuritogenesis, and induce stem cell differentiation[82-85]. SCCM was able to alter the morphology, gene expression, and functionality of OM-MSCs to a population that can better support nerve regeneration. These results were promising but still require the harvest and expansion of SCs, which, as mentioned above, requires a nerve harvest to retrieve. So, we sought to develop a defined GF media cocktail that could mimic the resulting differentiation of the SCCM by supplementing SC media with three well-known growth factors, NGF, BDNF, and GDNF, that are widely known to be secreted by SCs[37]. Additionally, we supplemented the media with forskolin as it can increase cAMP levels[38]. This method also induced SC-like cell morphology, gene expression, and functionality compared to OM-MSCs.

Changes in the relative expression of various genes were used to characterize the dOM-MSCs further. In addition to SCs and OM-MSCs, DRG and astrocyte samples were utilized as controls. In whole DRG samples containing neuronal, glial, and additional stromal cells, βIII tubulin

was upregulated compared to the other experimental groups (Figure 5I). As expected, relative gene expression of GFAP in the astrocyte population, the glial/support cells of the central nervous system, was upregulated compared to all other experimental populations (Figure 5D). We also see the expected relative expression of CD90 and CD44 (Figure 5A-B).

The relative expression of two SC markers, S100β and SOX10, were evaluated with RT-qPCR. No significant changes in the relative expression of S100β, a standard SC marker preferentially expressed in myelinating SCs, were measured across the various experimental populations[69]. SCs and dOM-MSCs (SCCM) were observed to have similar log2fold changes in relative expression (Figure 5G)[69]. SOX10 is a transcription factor that is a reliable marker of SCs in vitro. It is the only known marker expressed throughout the SC development process, with immature SCs, precursor SCs, non-myelinating SCs, and myelinating SCs all expressing SOX10[86]. Compared to undifferentiated OM-MSCs, both differentiated cell populations significantly upregulated SOX10, similar to SCs (Figure 5H). The co-upregulation in the relative expression of both S100β and SOX10 suggests that dOM-MSCs (SCCM) are induced towards myelinating SC-like cells.

Interestingly, no changes in relative gene expression of P75NTR, which is generally expressed by non-myelinating SCs and immature SCs, was observed between the various stem cell populations (OM-MSCs, dOM-MSCs [SCCM], dOM-MSCs [GF], Figure 5F)[87]. P75NTR has recently been investigated as an MSC marker, with recent works demonstrating that the subset of MSCs that are P75NTR+ proliferate faster and have a more significant differentiation potential[88-90]. Between 2-40% of the MSC population was reported to be positive for P75NTR+ in BM-MSCs and AD-MSCs. Future work will evaluate the expression of this marker in OM-MSCs as it may play a role in the enhanced differentiation potential[91].

NGF and GDNF are important neurotrophic factors that play a crucial role in the survival, growth, and differentiation of neurons in the peripheral nervous system[92-95]. VEGF is an important growth factor that guides re-vascularization and blood vessel formation, which is crucial after injury[96]. The expression of important SC markers and growth factors in the dOM-MSCs, particularly (SCCM), makes these cells a promising candidate for supporting peripheral nerve regeneration. In an uninjured peripheral nerve, SCs are responsible for the myelin sheath, a fatty protein facilitating signal propagation along the nerve axon[97,98]. Thus, the ability to produce MBP is a crucial indicator of cells that can function similarly to SCs. Here, this functionality was quantified with two metrics,

.

percent co-localization, and the Mander's coefficient, with co-localization measuring the overlap of MBP only on the neurites and the Mander's coefficient measuring the overall overlap of both the neurites and MBP. Both dOM-MSCs (SCCM and GF) expressed MBP and co-localized with neurite extensions significantly more than undifferentiated OM-MSCs but not significantly different than SCs (Figure 6). Additionally, the expression of MBP further demonstrates the differentiation towards myelinating SC-like cells[86]. Our in vitro co-culture experiment illustrates the promise and potential for these cells to function as SCs themselves do.

While both dOM-MSCs (SCCM and GF) had comparable co-localization of myelin basic protein on neurites, cell morphology, and gene expression differences suggest potential for improving the GF cocktail. In contrast, the GF cocktail did not impact the relative expression of SC marker S100β compared to OM-MSCs. The GF cocktail significantly upregulates SOX10, a second SC marker, compared to undifferentiated OM-MSCs. The dOM-MSCs (GF) also expressed MBP after co-culture with DRG. The difference in cell morphology and gene expression is likely due to unidentified SCCM components contributing to more complete differentiation. For example, BM-MSCs that differentiated in neurons with SCCM were found to be induced by microRNA pathways[83].

Additionally, other molecules produced by SCs may be inducing differentiation. Future work will further optimize our growth factor media to include other growth factors produced by SCs, such as VEGF, which we see is upregulated in SCs via RT-qPCR analysis (Figure 5J). Further optimization is vital to eliminate the need for SCs to differentiate OM-MSCs and develop a protocol conducive to scalability. A defined media is paramount to translating these cells to the clinic, allowing a xeno-free and scalable process.

In conclusion, we differentiated OM-MSCs towards SC-like cells and characterized their function and phenotype. Our in vitro co-culture experiment illustrates the promise and potential for these cells to function as SCs themselves do. Future work will assess the functionality of these cells in vivo and work towards translating this protocol towards differentiating human-derived OM-MSCs for enhancing peripheral nerve regeneration in patients.

Materials and Methods

Animal Statement
All animal work was performed with the approval of Northeastern University's Institutional Animal Care and Use Committee (NU-IACUC; Protocol #20-0207R) and followed the NIH guide for animal use.

.

Primary Cell Isolations and Immunocytochemistry Characterization
Olfactory Mucosa Derived Mesenchymal Stem Cells (OM-MSCs) were isolated from explants removed from the nasal cavities of 6-week-old Sprague Dawley rats and used to collect multipotent OM-MSCs according to a previously established protocol 32. Briefly, after euthanasia, facial muscle, and excess tissue were removed, exposing the nasal cavity. Once exposed, the olfactory turbinates were gently lifted to uncover the olfactory mucosa between the ceiling of the nasal cavity, the arc of the perpendicular plate, and the cribriform plate. Using a 26-gauge needle and forceps, the mucosa was removed and transferred into a petri dish filled with Dulbecco's Modified Eagle Medium/Nutrient Mixture F-12 (DMEM/F-12; Invitrogen). A similar protocol has been established for humans in which a small biopsy can be taken from the patient's nose in a quick outpatient procedure[32].

The rat tissue was digested in dispase II solution (2.4 IU/mL; Sigma-Aldrich) for one hour at 37°C, then transferred to a collagenase I (4 mg/mL; Sigma-Aldrich) solution for ten minutes. The tissue was then dissociated with a pipette, and the reaction was terminated with phosphate-buffered saline (PBS; Sigma-Aldrich). The samples were then centrifuged at 300 g for five minutes. After removing the supernatant, the cell pellet was resuspended in DMEM/F-12 supplemented with 10% (v/v) fetal bovine serum (FBS; Corning) and 50 U/mL -penicillin/streptomycin (P/S; Sigma-Aldrich) and plated in a 12-well plate (Thermo Fisher). Cell medium (1 mL) was added to each well and exchanged every 2-3 days. The cells were incubated in standard cell culture conditions throughout all experiments (37°C, 5% CO_2). After cells reached 70-90% confluency (about 1-2 weeks), the OM-MSCs were passaged according to standard cell culture procedures with 0.25% Trypsin-EDTA (Gibco) and plated either in BioLite 25 cm² vented flasks (Thermo Scientific) to expand and use for further experimentation or on laminin-coated glass coverslips in a 12 well-plate for immunocytochemistry (ICC).

Glass coverslips (12 mm; Fisherbrand) were sterilized with ultraviolet light for six minutes per side and treated with plasma for two minutes using a Harrick Expanded Plasma Cleaner (PDC-002). Then, a 6.7 μg/mL mouse laminin (Corning) solution, diluted in Hank's Balanced Salt Solution (HBSS; Sigma Aldrich), was applied to the coverslips and incubated at 37°C for 30 minutes. After, the excess solution was removed, and the coverslips were rinsed with sterile water two times. The cells were then seeded on the coverslips, and 1 mL of media was added. The cells were incubated for three days before performing ICC.

SCs were isolated from postnatal day 2 (p2) Sprague Dawley rats based on previously established methods 14. Sciatic nerves were isolated and plated in basic media (Dulbecco's Modified Eagle Medium [DMEM; Hyclone], 10% (v/v) FBS, 50 U/mL P/S, and 2 mM of L-glutamine [Sigma Aldrich]). SCs were purified from contaminating fibroblasts with an anti-mitotic agent, $10-5$ M cytosine arabinoside (ara-CC; Sigma-Aldrich), and complement-mediated cell lysis (anti-CD90/Thy 1.1, Invitrogen). SCs were seeded on laminin-coated coverslips to assess purity, and ICC staining for S100β (Invitrogen: BSM-52506R, Invitrogen: A11035) was performed as detailed below. Spindle morphology and purity >98% was confirmed. Cells were expanded in BioLite 25 cm² vented flasks (Thermo Scientific) in growth media (DMEM, 10% (v/v) FBS, 50 U/mL P/S, 2 mM of L-glutamine, 10 μg/mL bovine pituitary extract [BPE; Gibco], 6.6 μM forskolin [Sigma-Aldrich]) in standard conditions (5% CO_2, 37°C). Media was exchanged every 2-3 days.

Central nervous system glia astrocytes were isolated from postnatal day 2 (p2) Sprague Dawley rats based on previously established methods 33,34. Neonatal brains were isolated, and the olfactory bulbs, cerebellum, and meninges were discarded. Brain tissue was suspended in a dissociation media containing DMEM, P/S, and papain (20 U/mL; MP Biomedicals) and manually triturated. After allowing the brain matter to settle for 15 minutes at standard cell culture conditions (5% CO_2, 37°C), the liquid supernatant was removed and diluted 1:1 with media (DMEM, 10% FBS,

and P/S) to deactivate the papain. The brain matter was resuspended in a second dissociation media containing DMEM, P/S, papain (20 U/mL), and DNase 1 (20 U/mL; Thermo Scientific) and again manually triturated. The liquid supernatant was removed after 15 minutes of incubation and diluted in the same manner as the previous extraction. Nine additional extractions were performed with the first dissociation media (no DNase 1). A total of ten extractions were performed, equal to the total number of brains harvested. The extracted cells were centrifuged at 400 g for five minutes, resuspended in media, and plated on a poly-l-lysine-coated T75 flask (5,000 cells/cm2). After two weeks of culture, the flask was shaken at 200 RPM for four hours to remove microglia on top of the astrocyte monolayer. After removal, the astrocytes were cultured and expanded. ICC staining for GFAP (Invitrogen: PA1-10004, Invitrogen: A11035) was performed to confirm a purity of >98%.

Neuronal explants, dorsal root ganglia (DRG), were isolated from postnatal (p2) Sprague Dawley Rats via previously reported methods 35,36. Whole spines were removed, and connective tissue was trimmed off. Spines were split medially with scissors, and the spinal cord was removed. DRG explants were lifted from the spinal column with forceps and trimmed excess connective tissue with a 15-blade scalpel. DRG were stored in 15 mL tubes filled with Hibernate-A Medium (Gibco) and wrapped with parafilm at 4°C for no more than one week before use.

Immunocytochemistry
For ICC, all samples were fixed with 4% paraformaldehyde (PFA; Sigma-Aldrich) for 20 minutes and permeabilized with 0.1% (v/v) Triton-X (Sigma-Aldrich) diluted in HBSS for ten minutes at room temperature. Non-specific binding was blocked with freshly prepared 2.5% (v/v) goat serum (Sigma-Aldrich) diluted in HBSS for two hours at room temperature. Primary antibodies and secondary antibody dilutions are listed in Table 1. Primary antibodies were diluted in the 2.5% goat serum solution and incubated at room temperature for one hour, followed by a triplicate rinse of HBSS. Following this step, secondary antibodies were diluted in the 2.5% goat serum solution and incubated at room temperature for one hour. Before imaging, the samples were washed three times with HBSS and mounted on a glass slide with ProLong™ Gold Antifade Mountant with DAPI (Invitrogen). Images were captured with an inverted light microscope (Zeiss Axio Observer).

Characterizing OM-MSCs with Flow Cytometry
Passage 5 OM-MSC cell populations were characterized using a Beckman Coulter CytoFLEX. A flow buffer solution was prepared in 50 mL of HBSS with 200 μL EDTA (Corning), 50 mg sodium azide (Sigma-Aldrich), and 50 mg of saponin (Sigma-Aldrich). This flow buffer was kept at 4°C for up to four weeks. Cells were passaged with 0.25% Trypsin-EDTA (Gibco) and pelleted. The cell pellet was resuspended in a solution containing a fixable viability stain (Invitrogen; 1 μL per mL of HBSS) and incubated at 4°C for 30 minutes in the dark. The cells were centrifuged at 350 g for five minutes, the supernatant was removed, and the cells were rinsed by resuspension in HBSS to remove excess stain. After centrifuging again to remove the HBSS, the flow buffer supplemented with 5% (v/v) mouse serum (Invitrogen) was added to the rinsed cell pellet and mixed well by gentle pipetting. The cells were incubated in this solution for 10 minutes at 4°C to limit non-specific binding and to permeabilize the cells. Conjugated antibodies (Table 2) were added to the cell solution and mixed well with the suspended cell solution with a pipette. The cells were incubated at 4°C for 30 minutes and then centrifuged at 350 g for five minutes. The supernatant was removed, and the cells were resuspended in a flow buffer. This centrifugation and resuspension were repeated. The cell solution was then run through the flow cytometer. Compensation was established with single fluorescence and unstained controls. Fluorescence minus one (FMO) controls were utilized to determine gating parameters. Analysis was performed with Flowjo™ Software.

Differentiation of OM-MSCs towards SC-like cells
OM-MSCs were differentiated with two separate media compositions: an SC-conditioned media (SCCM) and a growth factor media (GF). OM-MSCs .

were plated at 5,000 cells/cm² density for both differentiation protocols, kept at standard culture conditions (5% CO_2, 37°C) for 21 days, and had complete media exchanges every 2-3 days (Figure 3A). The first protocol utilizes SCCM or media taken directly from SC cultures after incubation for 2-3 days. The second protocol aims to mimic the SCCM by supplementing SC media (DMEM, 10% (v/v) FBS, 50 U/mL P/S, 2 mM of L-glutamine, 10 μg/mL BPE, and 6.6 μM forskolin) with additional forskolin (final concentration of 16.5 μM in media) to increase cyclic adenosine monophosphate (cAMP) levels and three well-known growth factors secreted natively by SCs: 2.5 ng/mL nerve growth factor (NGF; Gibco #13257-019), 2.5 ng/mL glial cell line-derived neurotrophic factor (GDNF; Gibco #PHC7045), 2.5 ng/mL brain-derived neurotrophic factor (BDNF; Gibco #450-02)37,38. Other media formulations were attempted to optimize the differentiation process, such as a previously established differentiation protocol for BM-MSCs to SCs and varying concentrations of forskolin[39] (Supplementary Fig. 4). ICC was performed with S100β and Nestin primary antibodies to evaluate changes in expression and cell morphology.

To quantify differences in cell shape, CellProfiler™ software was used 40. Ten immunofluorescent image sets (composed of individual channel images of the exact location) were imported in batches per condition to CellProfiler™ and processed through a custom-designed pipeline (Supplemental Figure 5). The Otsu thresholding method identified the cell nuclei from the DAPI channel. In combination with the identified nuclei and a secondary channel (S100β), the cell body was identified through a propagation method and Otsu thresholding. The "MeasureObjectSizeShape" function characterized each cell body, and results were outputted into an Excel file.

Gene Expression Evaluation with qPCR
Gene expression was evaluated using RT-qPCR, comparing RNA extracted from DRG, astrocytes, SCs, OM-MSCs, and dOM-MSCs (SCCM and GF media). RNA was collected and isolated from cell pellets with an RNeasy® Mini Kit (Qiagen: 74106) following the manufacturer's instructions. In place of the gDNA eliminator column, the RNase-Free DNase Set (Qiagen: 79254) was used to ensure sample purity. Ethanolytic precipitation was performed after isolating the RNA to remove excess salts and contaminants. Two volumes of 100% ethanol and 0.1 volumes of 3M sodium acetate (Invitrogen: AM9740) were added to each sample. After vortexing, the samples were kept at -80°C. Before translating the RNA to complementary DNA (cDNA), the RNA was pelleted by centrifugation for 20 minutes (>8000 x g, 4°C). Excess liquid was removed, and the pellet was washed with 100% ethanol and centrifuged again for 10 minutes (>8000 x g, 4°C). The pellet was washed and spun down one more time. After removing all liquid from the pellet, the RNA was allowed to air dry on ice for 10-15 minutes. RNASE-DNASE-free water (12 μL) was added to the pellet to resuspend the RNA. Yield and purity were quantified with a Nanodrop 1 (Thermo Scientific, Supplemental Table 1). Samples with an A260/A280 value between 1.95 and 2.1 were translated to cDNA. The Quantitect Reverse Transcription Kit (Qiagen: 205311) was used according to the manufacturer's instructions to translate RNA to cDNA, and the cDNA was stored at -20°C.

Primers for genes of interest were designed by inputting the National Center for Biotechnology Information (NCBI) reference sequence into Primer-BLAST41 (Table 3, Supplementary Table 2). Preference was given to primers with a PCR product of 70-200 base pairs (bp) and a melting temperature between 58-62°C. Lyophilized, salt-free primer pairs were then ordered through Eurofin. These primers were then resuspended in Tris-EDTA (TE; Fisher Scientific) to a stock concentration of 10 μM. A standard curve was performed to validate primers with a 10x serial dilution of cDNA starting at a concentration of 1 ng/μL. Optimal primer concentrations were determined based on standard curve efficiency (Supplementary Table 3).

The expected PCR product size was calculated using AmplifX 42, and the PCR product produced was verified with gel electrophoresis (Supplemental Table 3, Supplemental Figure 6A-B). Briefly, 2% (w/v) agarose gel was made by heating and dissolving agarose (Invitrogen) in 1x Tris Acetate EDTA (TAE; Thermo Scientific). SYBR® Safe DNA gel stain (10,000x: Invitrogen) was added to the solution to a final concentration of 1x, and the solution was then poured into a gel mold (Invitrogen), and a well comb was added. After solidification, the well comb was removed, and the gel was placed in the horizontal gel electrophoresis system. 1x TAE buffer with DNA gel stain was used to submerge the gel. Samples and a 50 bp DNA ladder were loaded into wells with Loading Dye (Invitrogen). The gel was run at 60 V at 0.03 A for two hours with a PowerEase® 300W (Life Technologies). Gels were imaged immediately after the run with a Gel Doc™ XR+ (Bio-Rad). Image Lab 6.1 (Bio-Rad) software was used to calculate the PCR product size.

Eight reference gene primers were designed and validated: beta-2-microglobin (B2M), glyceraldehyde-3-phosphate dehydrogenase (GAPDH), glucuronidase beta (GUSB), hypoxanthine-guanine phosphoribosyl transferase (HPRT1), ornithine decarboxylase antizyme 1 (OAZ1), peptidylprolyl isomerase A (PPIA), ribosomal protein lateral stalk subunit P0 (RPLP0), and ribosomal protein S13 (RSP13) (Supplementary Table 2). A gene study was performed across experimental groups using CFX Maestro software (Bio-Rad) with five validated reference genes (GAPDH, HPRT1, GUSB, RSP13, and B2M) to confirm stability as a reference gene. Samples from the six experimental groups were pooled together to assess the stability. GAPDH and HPRT1 had the highest stability and were chosen as reference genes to use in the experimental quantification plates (Supplementary Fig. 6C). Ten genes of interest were designed, validated, and evaluated: S100 calcium-binding protein B (S100β; glial cell marker), SRY-box transcription factor 10 (SOX10; glial cell marker), glial fibrillary acidic protein (GFAP; glial marker), nerve growth factor receptor (P75NTR; neurotrophin receptor), tubulin beta 3 class III (TUBB3; neural marker), nerve growth factor (NGF; neurotrophic factor), glial cell-derived neurotrophic factor (GDNF; neurotrophic factor), vascular endothelial growth factor (VEGF; growth factor), cluster of differentiation 90 (CD90/Thy-1 cell surface antigen; mesenchymal stem cell marker), and cluster of differentiation 44 (CD44; (mesenchymal stem cell marker) (Supplementary Table 3).

Three 384-well (Thermo Fisher) quantification plates were designed and executed, each containing the two reference genes and experimental genes to minimize intraplate variability. 4 ng of cDNA, 5 µL of SsoAdvanced™ Universal SYBR® Green Supermix, and one set of forward/reverse primer pairs (250-500 nM) were added to each well. All reactions were run in triplicate with four samples per experimental group. After the plate setup, the plate was sealed with Absolute qPCR Plate Seals (Thermo Scientific) and spun down for one minute. PCR was run with a CFX384 thermal cycler (Bio-Rad). 40 cycles of denaturation (98°C, 10 seconds), annealing (60°C, 20 seconds), and extension (72°C, 20 seconds) were performed. A melt curve was also generated with a temperature gradient of 65-95°C at 0.1°C per second. Gene expression was normalized to the two reference genes, GAPDH and HPRT1, and changes in relative expression (log2fold) were quantified using the ΔΔCq method43.

Co-culture with DRG to Assess the Functionality of Differentiated Cells

Dissociated DRG were cultured with SCs, OM-MSCs, dOM-MSCs (SCCM), and dOM-MSCs (GF) to assess the ability of our differentiated cells to myelinate neurons. The procedure is detailed in the timeline in Figure 6A and is based on previous protocols 44,45. DRG explants were dissociated in 1 mg/mL collagenase (Gibco), 1 mg/mL dispase (Gibco), 15 U/mL papain (Sigma-Alrich) in HBSS for 45 minutes and then triturated 20-40 times with a glass fire-polished pipette. The solution was then diluted at a 1:1 ratio with Neurobasal Media-A (Gibco) to stop the enzymatic activity and centrifugated at 400 g for five minutes. The supernatant was removed, and the dissociated cells were resuspended in Neurobasal-A media, supplemented with 2 mM L-glutamine, 50 U/mL P/S, 1x B-27
.

(Gibco), 25 ng/mL NGF, and 50 μg/mL of ascorbic acid (Sigma-Aldrich) and plated on laminin-coated coverslips in 24 well plates at a density of 40,000 cells per well. After 12 hr, dissociated DRG were treated with $10-5$ M ara-C to reduce the number of native support cells (SCs and fibroblasts). After 24 hr, the ara-C was removed with a full media exchange. SCs, OM-MSCs, and dOM-MSCs (SCCM), or dOM-MSCs (GF), were added to the DRG cultures at a density of 50,000 cells per well. The experiment was maintained in standard conditions (37°C, 5% CO_2), and a half media exchange was performed every 3-4 days. The DRG were cultured for 21 days and then fixed and stained as detailed above with mouse anti-neurofilament (NF-H; Invitrogen: MA1-2012) and rabbit anti-myelin basic protein (MBP; Invitrogen: MA5-42370) primary antibodies. Image analysis was performed using a custom-designed CellProfiler™ pipeline, detailed in Supplementary Fig. 7, and the percent pixel overlap of MBP on NF-H was quantified using the "MeasureImageOverlap" function[40]. The Manders coefficient was measured using the "MeasureColocalization" function. Three independent trials were conducted.

Statistical Analysis

Statistical analysis was performed with Prism GraphPad. The normality of each data set was assessed with the Shapiro-Wilk Test. For parametric (gene expression) data, significance was evaluated with one-way ANOVA and Tukey's multiple comparison test. For nonparametric data (CellProfiler™ data analysis), a Kruskal Wallis test with a post hoc Dunn's multiple comparison test was performed. Error bars on graphs represent the standard deviation (SD). *, **, ***, and **** represent a p-value of less than 0.05, 0.01, 0.001, and 0.0001, respectively.

Data availability statement

The data supporting this study's findings are available upon reasonable request from the authors.

Acknowledgments:

The authors would like to thank Dr. Rebecca Willits and Dr. McKay Cavanaugh for their assistance and use of their flow cytometer and Dr. David Diaz for his help isolating astrocytes and Schwann cells. Additionally, we would like to thank the Center for Research Innovation (CRI) and Spark Fund program at Northeastern University for all their support.

Funding

We thank the Northeastern University Spark Fund and the Department of Chemical Engineering.

Author Contributions

K.N. and R.K. conceived the project. K.N. performed all experimental work analysis and wrote the manuscript. R.K. and A.K. provided analysis support. A.K. and R.K. provided intellectual input and advice. All authors edited and provided feedback on the manuscript. R.K. supervised the work.

Conflict of Interest

The authors declare no competing interests.

ORCID IDs

Katelyn Neuman: https://orcid.org/0000-0002-0038-0432
Abigail Koppes: https://orcid.org/0000-0003-0433-9290
Ryan Koppes: https://orcid.org/0000-0002-3376-6358

References:

1. Taylor, C. A., Braza, D., Rice, J. B. & Dillingham, T. The incidence of peripheral nerve injury in extremity trauma. Am J Phys Med Rehabil 87, 381-385 (2008). https://doi.org:10.1097/PHM.0b013e31815e6370

2. Hussain, G. et al. Current Status of Therapeutic Approaches against Peripheral Nerve Injuries: A Detailed Story from Injury to Recovery. Int J Biol Sci 16, 116-134 (2020). https://doi.org:10.7150/ijbs.35653

3. Bergmeister, K. D. et al. Acute and long-term costs of 268 peripheral nerve injuries in the upper extremity. PLOS ONE 15, e0229530 (2020). https://doi.org:10.1371/journal.pone.0229530

4. Huckhagel, T., Nüchtern, J., Regelsberger, J., Lefering, R. & TraumaRegister, D. G. U. Nerve injury in severe trauma with upper extremity involvement: evaluation of 49,382 patients from the TraumaRegister DGU® between 2002 and 2015. Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine 26, 76 (2018). https://doi.org:10.1186/s13049-018-0546-6

5. Bruyns, C. N. et al. Predictors for return to work in patients with median and ulnar nerve injuries. J Hand Surg Am 28, 28-34 (2003). https://doi.org:10.1053/jhsu.2003.50026

6. Torres, R. Y. & Miranda, G. E. Epidemiology of Traumatic Peripheral Nerve Injuries Evaluated by Electrodiagnostic Studies in a Tertiary Care Hospital Clinic. Bol Asoc Med P R 107, 79-84 (2015).

7. Saheb-Al-Zamani, M. et al. Limited regeneration in long acellular nerve allografts is associated with increased Schwann cell senescence. Exp Neurol 247, 165-177 (2013). https://doi.org:10.1016/j.expneurol.2013.04.011

8. Azouz, S. M., Lucas, H. D., Mahabir, R. C. & Noland, S. S. A Survey of the Prevalence and Practice Patterns of Human Acellular Nerve Allograft Use. Plast Reconstr Surg Glob Open 6, e1803-e1803 (2018). https://doi.org:10.1097/GOX.0000000000001803

9. Mathot, F. et al. Gene expression profiles of differentiated and undifferentiated adipose derived mesenchymal stem cells dynamically seeded onto a processed nerve allograft. Gene 724, 144151 (2020). https://doi.org:https://doi.org/10.1016/j.gene.2019.144151

10. Clements, M. P. et al. The Wound Microenvironment Reprograms Schwann Cells to Invasive Mesenchymal-like Cells to Drive Peripheral Nerve Regeneration. Neuron 96, 98-114.e117 (2017). https://doi.org:https://doi.org/10.1016/j.neuron.2017.09.008

11. Gomez-Sanchez, J. A. et al. After Nerve Injury, Lineage Tracing Shows That Myelin and Remak Schwann Cells Elongate Extensively and Branch to Form Repair Schwann Cells, Which Shorten Radically on Remyelination. The Journal of Neuroscience 37, 9086-9099 (2017). https://doi.org:10.1523/jneurosci.1453-17.2017

12. Syu, W.-Z. et al. Adipose-Derived Neural Stem Cells Combined with Acellular Dermal Matrix as a Neural Conduit Enhances Peripheral Nerve Repair. Cell Transplant 28, 1220-1230 (2019). https://doi.org:10.1177/0963689719853512

13. Hood, B., Levene, H. B. & Levi, A. D. Transplantation of autologous Schwann cells for the repair of segmental peripheral nerve defects. Neurosurgery Focus 26 (2009). https://doi.org:10.3171/FOC.2009.26.2.E4

14. Morrissey, T. K., Kleitman, N. & Bunge, R. P. Isolation and functional characterization of Schwann cells derived from adult peripheral nerve. Journal of Neuroscience 11, 2433-2442 (1991).

15. Kozlowska, U. et al. Similarities and differences between mesenchymal stem/progenitor cells derived from various human tissues. World J Stem Cells 11, 347-374 (2019). https://doi.org:10.4252/wjsc.v11.i6.347

16. da Silva Meirelles, L., Fontes, A. M., Covas, D. T. & Caplan, A. I. Mechanisms involved in the therapeutic properties of mesenchymal stem cells. Cytokine & Growth Factor Reviews 20, 419-427 (2009). https://doi.org:https://doi.org/10.1016/j.cytogfr.2009.10.002

17. Kot, M. et al. The Importance of HLA Assessment in "Off-the-Shelf" Allogeneic Mesenchymal Stem Cells Based-Therapies. Int J Mol Sci 20 (2019). https://doi.org:10.3390/ijms20225680

18. Le Blanc, K., Tammik, C., Rosendahl, K., Zetterberg, E. & Ringdén, O. HLA expression and immunologic properties of differentiated and undifferentiated mesenchymal stem cells. Exp Hematol 31, 890-896 (2003). https://doi.org:10.1016/s0301-472x(03)00110-3

19. Wang, Y., Yi, H. & Song, Y. The safety of MSC therapy over the past 15 years: a meta-analysis. Stem Cell Research & Therapy 12, 545 (2021). https://doi.org:10.1186/s13287-021-02609-x

20. Lalu, M. M. et al. Safety of cell therapy with mesenchymal stromal cells (SafeCell): a systematic review and meta-analysis of clinical trials. PLoS One 7, e47559 (2012). https://doi.org:10.1371/journal.pone.0047559

21. McGrath, A. M. et al. Long-Term Effects of Fibrin Conduit with Human Mesenchymal Stem Cells and Immunosuppression after Peripheral Nerve Repair in a Xenogenic Model. Cell Med 10, 2155179018760327 (2018). https://doi.org:10.1177/2155179018760327

22. Kaka, G., Arum, J., Sadraie, S. H., Emamgholi, A. & Mohammadi, A. Bone Marrow Stromal Cells Associated with Poly L-Lactic-Co-Glycolic Acid (PLGA) Nanofiber Scaffold Improve Transected Sciatic Nerve Regeneration. Iran J Biotechnol 15, 149-156 (2017). https://doi.org:10.15171/ijb.1576

23. Fernandes, M. et al. Bone marrow-derived mesenchymal stem cells versus adipose-derived mesenchymal stem cells for peripheral nerve regeneration. Neural regeneration research 13, 100-104 (2018). https://doi.org:10.4103/1673-5374.224378

24. Fernandes, M. et al. Bone marrow-derived mesenchymal stem cells versus adipose-derived mesenchymal stem cells for peripheral nerve regeneration. Neural Regen Res 13, 100-104 (2018). https://doi.org:10.4103/1673-5374.224378

25. Georgiou, M., Golding, J. P., Loughlin, A. J., Kingham, P. J. & Phillips, J. B. Engineered neural tissue with aligned, differentiated adipose-derived stem cells promotes peripheral nerve regeneration across a critical sized defect in rat sciatic nerve. Biomaterials 37, 242-251 (2015). https://doi.org:10.1016/j.biomaterials.2014.10.009

26. Salehi, M. et al. Alginate/chitosan hydrogel containing olfactory ectomesenchymal stem cells for sciatic nerve tissue engineering. Journal of Cellular Physiology 234, 15357-15368 (2019). https://doi.org:https://doi.org/10.1002/jcp.28183

27. Roche, P. et al. Olfactory Derived Stem Cells Delivered in a Biphasic Conduit Promote Peripheral Nerve Repair In Vivo. STEM CELLS Translational Medicine 6, 1894-1904 (2017). https://doi.org:https://doi.org/10.1002/sctm.16-0420

28. Lindsay, L., Johnstone, A., McGrath, A., Mallinson, D. & Barnett, C. Comparative miRNA-Based Fingerprinting Reveals Biological Differences in Human Olfactory Mucosa- and Bone-Marrow-Derived Mesenchymal Stromal Cells. Stem Cell Reports 6, 729-742 (2016).

29. Lindsay, S. L. & Barnett, S. C. Are nestin-positive mesenchymal stromal cells a better source of cells for CNS repair? Neurochem Int 106, 101-107 (2017). https://doi.org:10.1016/j.neuint.2016.08.001

30. Féron, F., Perry, C., Girard, S. D. & Mackay-Sim, A. in Neural Progenitor Cells: Methods and Protocols (eds Brent A. Reynolds & Loic P. Deleyrolle) 107-114 (Humana Press, 2013).

31. Delorme, B. et al. The human nose harbors a niche of olfactory ectomesenchymal stem cells displaying neurogenic and osteogenic properties. Stem Cells Dev 19, 853-866 (2010). https://doi.org:10.1089/scd.2009.0267

32. Girard, S. D. et al. Isolating Nasal Olfactory Stem Cells from Rodents or Humans. JoVE, e2762 (2011). https://doi.org:10.3791/2762

33. Zarei-Kheirabadi, M., Mirsadeghi, S., Vaccaro, A. R., Rahimi-Movaghar, V. & Kiani, S. Protocol for purification and culture of astrocytes: useful not only in 2 days postnatal but also in adult rat brain. Molecular Biology Reports 47, 1783-1794 (2020). https://doi.org:10.1007/s11033-020-05272-2

34. Calvo, B., Rubio, F., Fernández, M. & Tranque, P. Dissociation of neonatal and adult mice brain for simultaneous analysis of microglia, astrocytes and infiltrating lymphocytes by flow cytometry. IBRO Reports 8, 36-47 (2020). https://doi.org:https://doi.org/10.1016/j.ibror.2019.12.004

35. Seggio, A. M., Ellison, K. S., Hynd, M. R., Shain, W. & Thompson, D. M. Cryopreservation of transfected primary dorsal root ganglia neurons. Journal of Neuroscience Methods 173, 67-73 (2008). https://doi.org:https://doi.org/10.1016/j.jneumeth.2008.05.017

36. Puzan, M. L., Legesse, B., Koppes, R. A., Fenniri, H. & Koppes, A. N. Bioactive Organic Rosette Nanotubes Support Sensory Neurite Outgrowth. ACS Biomaterials Science & Engineering 4, 1630-1640 (2018). https://doi.org:10.1021/acsbiomaterials.8b00326

37. Jessen, K. R., Mirsky, R. & Lloyd, A. C. Schwann Cells: Development and Role in Nerve Repair. Cold Spring Harb Perspect Biol 7, a020487 (2015). https://doi.org:10.1101/cshperspect.a020487

38. Seamon, K. B., Padgett, W. & Daly, J. W. Forskolin: unique diterpene activator of adenylate cyclase in membranes and in intact cells. Proc Natl Acad Sci U S A 78, 3363-3367 (1981). https://doi.org:10.1073/pnas.78.6.3363

39. Xue, J. et al. Differentiation of Bone Marrow Stem Cells into Schwann Cells for the Promotion of Neurite Outgrowth on Electrospun Fibers. ACS Applied Materials & Interfaces 9, 12299-12310 (2017). https://doi.org:10.1021/acsami.7b00882

40. Stirling, D. R. et al. CellProfiler 4: improvements in speed, utility and usability. BMC Bioinformatics 22, 433 (2021). https://doi.org:10.1186/s12859-021-04344-9

41. Ye, J. et al. Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. BMC Bioinformatics 13, 134 (2012). https://doi.org:10.1186/1471-2105-13-134

42. AmplifX v. 2.1.1 (Aix-Marseille University, 2020).

43. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods 25, 402-408 (2001). https://doi.org:10.1006/meth.2001.1262

44. Liu, Q. et al. Human neural crest stem cells derived from human ESCs and induced pluripotent stem cells: induction, maintenance, and differentiation into functional schwann cells. Stem Cells Transl Med 1, 266-278 (2012). https://doi.org:10.5966/sctm.2011-0042

45. Kim, H.-S., Kim, J. Y., Song, C. L., Jeong, J. E. & Cho, Y. S. Directly induced human Schwann cell precursors as a valuable source of Schwann cells. Stem Cell Research & Therapy 11, 257 (2020). https://doi.org:10.1186/s13287-020-01772-x

46. Escada, P. A., Lima, C. & Madeira da Silva, J. The human olfactory mucosa. European Archives of Oto-Rhino-Laryngology 266, 1675-1680 (2009). https://doi.org:10.1007/s00405-009-1073-x

47. Graziadei, P. P. & Graziadei, G. A. Neurogenesis and neuron regeneration in the olfactory system of mammals. I. Morphological aspects of differentiation

and structural organization of the olfactory sensory neurons. J Neurocytol 8, 1-18 (1979). https://doi.org:10.1007/bf01206454

Calof, A. L. & Chikaraishi, D. M. Analysis of neurogenesis in a mammalian neuroepithelium: Proliferation and differentiation of an olfactory neuron precursor in vitro. Neuron 3, 115-127 (1989). https://doi.org:https://doi.org/10.1016/0896-6273(89)90120-7

Mackay-Sim, A. & Kittel, P. Cell dynamics in the adult mouse olfactory epithelium: a quantitative autoradiographic study. J Neurosci 11, 979-984 (1991). https://doi.org:10.1523/jneurosci.11-04-00979.1991

Murrell, W. et al. Multipotent stem cells from adult olfactory mucosa. Developmental Dynamics 233, 496-515 (2005). https://doi.org:https://doi.org/10.1002/dvdy.20360

Tomé, M., Lindsay, S. L., Riddell, J. S. & Barnett, S. C. Identification of Nonepithelial Multipotent Cells in the Embryonic Olfactory Mucosa. STEM CELLS 27, 2196-2208 (2009). https://doi.org:https://doi.org/10.1002/stem.130

Lindsay, S. L., Riddell, J. S. & Barnett, S. C. Olfactory mucosa for transplant-mediated repair: A complex tissue for a complex injury? Glia 58, 125-134 (2010). https://doi.org:https://doi.org/10.1002/glia.20917

Lin, Y. et al. Proliferation and pluripotency potential of ectomesenchymal cells derived from first branchial arch. Cell Prolif 39, 79-92 (2006). https://doi.org:10.1111/j.1365-2184.2006.00374.x

Sung, J. H. et al. Isolation and Characterization of Mouse Mesenchymal Stem Cells. Transplantation Proceedings 40, 2649-2654 (2008). https://doi.org:https://doi.org/10.1016/j.transproceed.2008.08.009

Veron, A. D. et al. Isolation and characterization of olfactory ecto-mesenchymal stem cells from eight mammalian genera. BMC Veterinary Research 14, 17 (2018). https://doi.org:10.1186/s12917-018-1342-2

Pittenger, M. F. et al. Multilineage potential of adult human mesenchymal stem cells. Science 284, 143-147 (1999). https://doi.org:10.1126/science.284.5411.143

Tropel, P. et al. Isolation and characterisation of mesenchymal stem cells from adult mouse bone marrow. Exp Cell Res 295, 395-406 (2004). https://doi.org:10.1016/j.yexcr.2003.12.030

Entezari, M. et al. Human Olfactory Ecto-mesenchymal Stem Cells Displaying Schwann-cell-like Phenotypes and Promoting Neurite Outgrowth in Vitro. Basic Clin Neurosci 14, 31-42 (2023). https://doi.org:10.32598/bcn.2021.3542.1

Jaloux, C. et al. Human nasal olfactory stem cells, purified as advanced therapy medicinal products, improve neuronal differentiation. Frontiers in Neuroscience 16 (2022). https://doi.org:10.3389/fnins.2022.1042276

Alvites, R. D. et al. Rat Olfactory Mucosa Mesenchymal Stem/Stromal Cells (OM-MSCs): A Characterization Study. International Journal of Cell Biology 2020, 2938258 (2020). https://doi.org:10.1155/2020/2938258

Xun, C. et al. Insight into the proteomic profiling of exosomes secreted by human OM-MSCs reveals a new potential therapy. Biomedicine & Pharmacotherapy 131, 110584 (2020). https://doi.org:https://doi.org/10.1016/j.biopha.2020.110584

Lesley, J., Hascall, V. C., Tammi, M. & Hyman, R. Hyaluronan binding by cell surface CD44. J Biol Chem 275, 26967-26975 (2000). https://doi.org:10.1074/jbc.M002527200

Williams, A. F. & Gagnon, J. Neuronal Cell Thy-1 Glycoprotein: Homology with Immunoglobulin. Science 216, 696-703 (1982). https://doi.org:doi:10.1126/science.6177036

Maleki, M., Ghanbarvand, F., Reza Behvarz, M., Ejtemaei, M. & Ghadirkhomi, E. Comparison of Mesenchymal Stem Cell Markers in Multiple Human Adult Stem Cells. International Journal of Stem Cells 7, 118 - 126 (2014).

Aslan, H. et al. Osteogenic Differentiation of Noncultured Immunoisolated Bone Marrow-Derived CD105+ Cells. Stem Cells 24, 1728-1737 (2006). https://doi.org:10.1634/stemcells.2005-0546

Trowbridge, I. S. & Thomas, M. L. CD45: an emerging role as a protein tyrosine phosphatase required for lymphocyte activation and development. Annu Rev Immunol 12, 85-116 (1994). https://doi.org:10.1146/annurev.iy.12.040194.000505

Lee, M. K., Tuttle, J. B., Rebhun, L. I., Cleveland, D. W. & Frankfurter, A. The expression and posttranslational modification of a neuron-specific β-tubulin isotype during chick embryogenesis. Cell Motility 17, 118-132 (1990). https://doi.org:https://doi.org/10.1002/cm.970170207

Eng, L. F., Ghirnikar, R. S. & Lee, Y. L. Glial Fibrillary Acidic Protein: GFAP-Thirty-One Years (1969–2000). Neurochemical Research 25, 1439-1451 (2000). https://doi.org:10.1023/A:1007677003387

Mata, M., Alessi, D. & Fink, D. J. S100 is preferentially distributed in myelin-forming Schwann cells. Journal of Neurocytology 19, 432-442 (1990). https://doi.org:10.1007/BF01188409

Liu, Z. et al. Specific Marker Expression and Cell State of Schwann Cells during Culture In Vitro. PLOS ONE 10, e0123278 (2015). https://doi.org:10.1371/journal.pone.0123278

Zimmer, D. B., Cornwall, E. H., Landar, A. & Song, W. The S100 protein family: History, function, and expression. Brain Research Bulletin 37, 417-429 (1995). https://doi.org:https://doi.org/10.1016/0361-9230(95)00040-2

Tomita, K. et al. The neurotrophin receptor p75NTR in Schwann cells is implicated in remyelination and motor recovery after peripheral nerve injury. Glia 55, 1199-1208 (2007). https://doi.org:https://doi.org/10.1002/glia.20533

Roux, P. P. & Barker, P. A. Neurotrophin signaling through the p75 neurotrophin receptor. Prog Neurobiol 67, 203-233 (2002). https://doi.org:10.1016/s0301-0082(02)00016-3

Mignone, J. L., Kukekov, V., Chiang, A. S., Steindler, D. & Enikolopov, G. Neural stem and progenitor cells in nestin-GFP transgenic mice. J Comp Neurol 469, 311-324 (2004). https://doi.org:10.1002/cne.10964

Isern, J. et al. The neural crest is a source of mesenchymal stem cells with specialized hematopoietic stem cell niche function. eLife 3, e03696 (2014). https://doi.org:10.7554/eLife.03696

Chen, Q. et al. A fibrin matrix promotes the differentiation of EMSCs isolated from nasal respiratory mucosa to myelinating phenotypical Schwann-like cells. Mol Cells 38, 221-228 (2015). https://doi.org:10.14348/molcells.2015.2170

Morscheck, C. et al. Gene expression of Nestin, collagen type I and type III in human dental follicle cells after cultivation in serum-free medium. Oral and Maxillofacial Surgery 12, 89-92 (2008). https://doi.org:10.1007/s10006-008-0111-y

Liu, Y. et al. Recovery and maintenance of NESTIN expression in umbilical cord-MSC using a novel culture medium. AMB Express 10, 132 (2020). https://doi.org:10.1186/s13568-020-01067-7

Lindsay, S. L. et al. Human mesenchymal stem cells isolated from olfactory biopsies but not bone enhance CNS myelination in vitro. Glia 61, 368-382 (2013). https://doi.org:https://doi.org/10.1002/glia.22440

Lima, C. et al. Olfactory mucosa autografts in human spinal cord injury: a pilot clinical study. J Spinal Cord Med 29, 191-203; discussion 204-196 (2006). https://doi.org:10.1080/10790268.2006.11753874

Xiang, W. et al. Therapeutic Efficiency of Nasal Mucosa-Derived Ectodermal Mesenchymal Stem Cells in Rats with Acute Hepatic Failure. Stem Cells International 2023, 6890299 (2023). https://doi.org:10.1155/2023/6890299

Bampton, E. T. W. & Taylor, J. S. H. Effects of Schwann cell secreted factors on PC12 cell neuritogenesis and survival. Journal of Neurobiology 63, 29-48 (2005). https://doi.org:https://doi.org/10.1002/neu.20119

Wei, Z. J. et al. MicroRNA changes of bone marrow-derived mesenchymal stem cells differentiated into neuronal-like cells by Schwann cell-conditioned medium. Neural Regen Res 14, 1462-1469 (2019). https://doi.org:10.4103/1673-5374.253532

Huang, D. et al. Schwann cell-conditioned medium inhibits angiogenesis. Cancer research 60, 5966-5971 (2000).

Kwiatkowski, J. L., Rutkowski, J. L., Yamashiro, D. J., Tennekoon, G. I. & Brodeur, G. M. Schwann cell-conditioned medium promotes neuroblastoma survival and differentiation. Cancer research 58, 4602-4606 (1998).

Jessen, K. R. & Mirsky, R. The origin and development of glial cells in peripheral nerves. Nat Rev Neurosci 6, 671-682 (2005). https://doi.org:10.1038/nrn1746

Arroyo, E. J., Bermingham, J. R., Jr., Rosenfeld, M. G. & Scherer, S. S. Promyelinating Schwann cells express Tst-1/SCIP/Oct-6. J Neurosci 18, 7891-7902 (1998). https://doi.org:10.1523/jneurosci.18-19-07891.1998

Quirici, N. et al. Isolation of bone marrow mesenchymal stem cells by anti-nerve growth factor receptor antibodies. Experimental Hematology 30, 783-791 (2002). https://doi.org:10.1016/S0301-472X(02)00812-3

Latifi-Pupovci, H. et al. In vitro migration and proliferation ("wound healing") potential of mesenchymal stromal cells generated from human CD271+ bone marrow mononuclear cells. Journal of Translational Medicine 13, 1-9 (2015).

Yamamoto, N. et al. Isolation of multipotent stem cells from mouse adipose tissue. Journal of dermatological science 48, 43-52 (2007).

Zha, K. et al. Nerve growth factor (NGF) and NGF receptors in mesenchymal stem/stromal cells: Impact on potential therapies. Stem Cells Translational Medicine 10, 1008-1020 (2021). https://doi.org:10.1002/sctm.20-0290

Piirsoo, M., Kaljas, A., Tamm, K. & Timmusk, T. Expression of NGF and GDNF family members and their receptors during peripheral nerve development and differentiation of Schwann cells in vitro. Neurosci Lett 469, 135-140 (2010). https://doi.org:10.1016/j.neulet.2009.11.060

Mearow, K. M. The Effects of NGF and Sensory Nerve Stimulation on Collateral Sprouting and Gene Expression in Adult Sensory Neurons. Experimental Neurology 151, 14-25 (1998). https://doi.org:https://doi.org/10.1006/exnr.1998.6791

Cintron-Colon, A. F., Almeida-Alves, G., VanGyseghem, J. M. & Spitsbergen, J. M. GDNF to the rescue: GDNF delivery effects on motor neurons and nerves, and muscle re-innervation after peripheral nerve injuries. Neural Regen Res 17, 748-753 (2022). https://doi.org:10.4103/1673-5374.322446

Boyd, J. G. & Gordon, T. Neurotrophic factors and their receptors in axonal regeneration and functional recovery after peripheral nerve injury. Mol Neurobiol 27, 277-324 (2003). https://doi.org:10.1385/mn:27:3:277

Cattin, A.-L. et al. Macrophage-Induced Blood Vessels Guide Schwann Cell-Mediated Regeneration of Peripheral Nerves. Cell 162, 1127-1139 (2015). https://doi.org:10.1016/j.cell.2015.07.021

Boggs, J. M. Myelin basic protein: a multifunctional protein. Cell Mol Life Sci 63, 1945-1961 (2006). https://doi.org:10.1007/s00018-006-6094-7

Poitelon, Y., Kopec, A. M. & Belin, S. Myelin Fat Facts: An Overview of Lipids and Fatty Acid Metabolism. Cells 9 (2020). https://doi.org:10.3390/cells9040812

.

.

Additional Information

Supplementary Information accompanies this paper attached below.

.

Supplementary Information

Table 1. Displays Antibodies used for Immunocytochemistry.

Name	Classification	Dilution	Host Species	Brand	Product #
CD105	Mesenchymal Marker	1:250	Rabbit	BIOSS	BS-0579R
S100β	Glia Marker	1:300	Rabbit	BIOSS	BSM-52506R
Nestin	Stemness Marker	1:500	Mouse	Abcam	Ab254048
Beta III Tubulin	Neuronal Marker	1:300	Mouse	Sigma	T8660
GFAP	Neuronal Marker	1:250	Rabbit	Invitrogen	PA1-10004
Neurofilament-Heavy	Neurite Outgrowth	1:300	Mouse	Invitrogen	MA1-2012
Myelin Basic Protein	Glia Functionality	1:300	Rabbit	Invitrogen	MA5-42370
Alexa Fluor™ 647	Secondary	1:1000	Goat Anti-Mouse	Invitrogen	A32728
Alexa Fluor™ 546	Secondary	1:1000	Goat Anti-Rabbit	Invitrogen	A11035
Alexa Fluor™ 488	Secondary	1:1000	Goat Anti-Rabbit	Invitrogen	A27034
Phalloidin 546	Cytoskeleton	1:1000	-	Invitrogen	A22283
DAPI	Nuclei	1:500	-	Invitrogen	D1306

.

Table 2. Displays Antibodies/Markers for Flow Cytometry**.**

Excitation	E^m Filter	Marker	Color	Host/Target	Isotype	Clone	Brand	Catalog #	Dilution Factor
488	585/42	CD44	PE	Mouse α-Rat	IgG1	OX-50	Bio-Rad	MCA643PE	1:10
488	525/40	Nestin	Alexa 488	Mouse α-Rat	IgG1	Rat-401	Abcam	ab197495	1:100
488	695/40	CD90	PerCP	Mouse α-rat	IgG1	MRC OX-7	Abcam	ab216354	1:100
638	660/10	CD45	Alexa Fluor 647	Mouse α-Rat	IgG1	OX-1	Bio-Rad	MCA43A647	1:10
638	780/60	eFluor 780 Viability	eFluor 780	All Species	-	-	Invitrogen	65-0865-14	1:1000

Table 3: Displays Gene Targets and Primer Sequences

Gene Target	Classification	Reference Sequence	Forward Primer	Reverse Primer
GAPDH	Reference Gene	NM_017008.4	AACGACCCCTTCATTGACC	TCCACGACATACTCAGCAC
HPRT1	Reference Gene	NM_012583.2	CCCAGCGTCGTGATTAGTGA	GGCCTCCCATCTCCTTCATG
S100β	Glial Marker	NM_013191.2	AGCATCCTAACTCACGACC	AAAGACCTGACACTCTCTCC
SOX10	Glial Marker	NM_019193.2	AGCAGGCTGGACACTAAACC	AAAGGGGCAGCGATGTGTTA
GFAP	Glial Marker	NM_017009.2	AAATTGCTGGAGGGCGAAGA	GGGCACACCTCACATCACAT
P75NTR	Neuronal/Glial Marker	NM_012610.2	CACACTTCTCTCTCACACAC	GACCAGCCCTTGAACAAAC
Tubb3	Neuronal Marker	NM_139254.2	CATGAACGACCTGGTGTCTG	GTGCCCTGAAGAGCTGGTAG
NGF	Neurotrophic Factor	NM_001277055.1	ATAAGACCACAGCCACGGAC	ACGCCTTGACAAAGGTGTGA
GDNF	Neurotrophic Factor	NM_001401780.1	TCCTTCACGCCCTCAGAAAC	GTCACTGGTGACCACGCATA
VEGF	Growth Factor	NM_031836.3	GGAGTCTGTGCTCTGGGATTT	GAGTGAAGGAGCAACCTCTCC
CD90	Mesenchymal Marker	NM_012673.2	TGCCTTCTAAGCCAGATGCC	ACCCCTTTCCCCAACATTCC
CD44	Mesenchymal Marker	NM_012924.3	TGGGAATCAAGACAGTGGAG	CAGACGGCAAGAATCAGAG

.

Supplemental Figure 1. Characteristic immunocytochemistry of OM-MSCs. (A) Purple fluorescence denotes GFAP, (B) Green fluorescence denotes actin filaments, and (C) Displays both GFAP and actin filament expression merged with cell nuclei (shown in blue, counterstained with DAPI). Scale Bar = 200 µm.

Supplemental Figure 2. Characteristic immunocytochemistry of isolated astrocytes. Purple fluorescence denotes GFAP expression, and blue fluorescence denotes cell nuclei (counterstained with DAPI). Scale Bar = 200 µm.

Supplemental Figure 3. Characteristic flow cytometry of OM-MSCs isolated from adult male Sprague-Dawley rats. (A-C) Display gating strategy employed before analysis. The selection of (A) cell population and removal of debris, (B) single cells, and then (C) live cell population is shown. (D-G) Display the distribution of positive and negative cells for the four markers of interest: (D) CD45, (E) CD44, (F) CD90, and (G) nestin. The gray histogram represents the negative control, and the blue histogram represents the cells of interest. (H-I) Display the co-localization of (H) CD44 and nestin expression, (I) CD90 and nestin expression, and (J) CD44 and CD90 expression.

Supplemental Figure 4. Morphology and expression of S100β and Nestin for (A) Schwann cells, (B) OM-MSCs after culture in SCCM for 14 days, (C) OM-MSCs after a chemical induction protocol, (E) OM-MSCs after culture in GF media with reduced forskolin for 14 days, and (F) OM-MSCs after culture in GF media for 14 days. Green fluorescence indicates S100β, red fluorescence indicates Nestin, and blue fluorescence indicates the cell nuclei after counterstaining with DAPI. Scale bar = 200 μm.

.

Supplemental Figure 5. Cell Profiler™ Pipeline examples for (A-D) SCs, (E-F) OM-MSCs, and (G-H) dOM-MSCs. The cell nuclei are identified from the DAPI channel and used to identify and analyze the cell body.

Supplemental Table 1. mRNA Yield and Purity for RT-qPCR

Sample Name	RNA Yield (ng)	A260/A280
SC 1	11,126	2.1
SC 2	15,230	2.1
SC 3	10,199	2.06
SC 4	15,127	2.07
DRG 1	5,915	2.05
DRG 2	5,300	2.03
DRG 3	9,400	2.08
DRG 4	2,652	1.98
Astrocyte 1	2,929	1.95
Astrocyte 2	10,600	2.04
Astrocyte 3	9,900	2.02
Astrocyte 4	19,153	2.09
OM-MSC 1	13,500	2.12
OM-MSC 2	6,100	2.04
OM-MSC 3	5,400	1.98
OM-MSC 4	7,908	1.97
dOM-MSC 1 (SCCM)	1,844	1.9
dOM-MSC 2 (SCCM)	1,266	2.0
dOM-MSC 3 (SCCM)	2,974	1.98
dOM-MSC 4 (SCCM)	1,670	2.08
dOM-MSC 1 (GF)	3,380	2.01
dOM-MSC 2 (GF)	3,380	1.95
dOM-MSC 3 (GF)	4,350	2.05
dOM-MSC 4 (GF)	12,000	1.98

.

Supplemental Table 2. Reference Genes and Primer Sequences

Gene Target	Reference Sequence	Forward Primer	Reverse Primer
B2M	NM_012512	CACACGCAGTCTGAAAACCC	AAAGGAGCTTTGGGGACACA
GAPDH	NM_017008.4	AACGACCCCTTCATTGACC	TCCACGACATACTCAGCAC
GUSB	NM_017015.2	CAGGACACCGACAGGAGAGT	GGTCAGTGTGTTGTTGATGGC
HPRT1	NM_012583.2	CCCAGCGTCGTGATTAGTGA	GGCCTCCCATCTCCTTCATG
OAZ1	NM_001297557.1	CGAGAGTTCTAGGGTTGCCC	TTGTCGTTAGACGTCGGCTC
PPIA	NM_017101.1	CAGACGCCGCTGTCTCTTTTC	CGTGATGTCGAAGAACACGGT
RPLP0	NM_022402.2	GTCGGAGGAATCCGATGAGG	TTAAGCAGGCTGACTTGGTGT
RSP13	NM_130432.2	TGTCAGAAAGCACCTTGAGAGG	ATTTCCAGTTGGGTGGGAGC

.

Supplemental Table 3. Validation of Primer Pairs

Gene Target	Primer Concentration (nM)	STD Curve Efficiency (%)	R²	Expected Band Size (bp)	Estimated Band Size (bp)
B2M	500	92.5	0.995	184	167
GAPDH	500	96.5	0.996	191	190
GUSB	250	108.7	1.000	192	194
HPRT1	500	95.8	0.999	163	158
OAZ1	500	105.6	0.997	196	196
PPIA	500	91.3	0.991	70	67
RPLP0	500	105.8	0.997	74	73
RSP13	500	91.3	0.998	131	132
S100β	500	97.1	0.990	121	121
SOX10	500	92.9	0.998	116	129
GFAP	500	90.3	0.984	207	199
P75NTR	250	104.8	0.998	251	249
Tubb3	375	96.7	1.00	239	228
NGF	750	90.2	0.995	198	190
GDNF	500	95.5	0.991	171	168
VEGF	500	102.5	0.978	161	132
CD90	500	100.7	0.996	279	284
CD44	500	98.1	0.991	117	119

**STD Curve = Standard Curve

.

Supplemental Figure 6. **(A)** Band size verification of PCR products for reference genes with gel electrophoresis. **(B)** Band size verification of PCR products for experimental genes with gel electrophoresis. **(C)** The stability of reference genes were assessed with CFX Maestro software (Bio-Rad).

.