Beyond urbanization metrics: Using graphical causal models to investigate mechanisms in urban ecology and evolution

Jesse B. Borden[1], James P. Gibbs[2], John P. Vanek[1,3], Bradley J. Cosentino[1*]

[1]Department of Biology, Hobart and William Smith Colleges, 300 Pulteney Street, Geneva, NY, 14456, USA

[2]Department of Environmental Biology, The State University of New York College of Environmental Science and Forestry, 1 Forestry Drive, Syracuse, NY 13210, USA

[3]New York Natural Heritage Program, The State University of New York College of Environmental Science and Forestry, 625 Broadway, Albany, NY 12207, USA

[*]Corresponding author: Bradley J. Cosentino (cosentino@hws.edu)

Abstract

As the fastest-growing form of land use, urbanization drives profound environmental change that

reshapes biodiversity. Studies often characterize biodiversity patterns along urban-rural gradients

with statistical models that include one or more generic indices of urbanization. Such models

may be useful for prediction, but they do not permit explicit tests of causal hypotheses by which

urbanization mediates ecological and evolutionary processes. Here, we show how a graphical

causal modeling framework with directed acyclic graphs (DAGs) can be used to design clear

conceptual models and inform appropriate statistical analysis to better evaluate mechanistic

hypotheses about the effects of urbanization on biodiversity. We first introduce the basic

structure of DAGs and illustrate their value with simulated datasets. We then apply the

framework to a case study on coat color variation in eastern gray squirrels (*Sciurus carolinensis*)

along an urbanization gradient in Syracuse, New York, USA. We show how statistical models

ungrounded in causal assumptions are difficult to interpret and can lead to misleading

conclusions about mechanisms in urban ecology and evolution. In contrast, DAGs – by making

causal assumptions transparent – help researchers identify appropriate control variables for

statistical models to estimate the effects of interest. When applied to our case study, statistical

models informed by a DAG revealed a surprising finding: although squirrel melanism was more

prevalent in urban than rural populations, the prevalence of melanism was constrained by

components of environmental change common to cities, namely roads, forest loss, and predator

activity, contrary to expectations. Managing biodiversity in an increasingly urbanized world will

require a mechanistic understanding of how urbanization impacts biodiversity patterns; graphical

causal models such as DAGs can provide a powerful approach to do so.

Introduction

Urban areas represent the fastest growing land use type and have profound effects on biodiversity (Aronson et al. 2014; Ritchie and Roser 2018; Szabó et al. 2023). Urbanization consists of a suite of abiotic and biotic changes driven by high concentrations of humans and the physical environment they build (Ritchie and Roser 2018). The ecological and evolutionary consequences of urbanization are often studied using the urban-rural gradient approach, in which components of biodiversity are compared along a landscape-scale sampling area from the urban core to rural habitats (McDonnell and Pickett 1990; McDonnell and Hahs 2008). The urban-rural gradient approach has been used to document associations of population size, species distributions, community composition, and genetic and phenotypic trait variation with urbanization (Winchell et al. 2016; Moll et al. 2019; Cosentino and Gibbs 2022; Santangelo et al. 2022).

Although the urban-rural gradient method has been helpful to describe spatial patterns of biodiversity, reliance on univariate metrics of urbanization (e.g., impervious cover) can be problematic when the research goal is to identify processes underlying those patterns (Moll et al. 2019, Poisson et al. 2024). Moreover, the use of statistical models with multiple covariates (e.g., multiple linear regression) does not guarantee valid causal inference. Interpreting parameter estimates from statistical models as estimated causal effects requires explicit assumptions about the causal relationships among covariates in the model. In many studies those relationships are unstated, and covariates are included in models without regard to the causal connections between them (McElreath 2020).

Understanding mechanisms by which urbanization generates patterns in ecology and

evolution is complicated by at least two factors. First, environmental change associated with

urbanization can affect biodiversity through multiple causal pathways (Alberti et al. 2020). For

example, road networks can *directly* affect biodiversity through vehicular collisions and noise

pollution (Francis et al. 2012; Bennett 2017; Kok et al. 2023), but roads also have *indirect* effects

by contributing to habitat fragmentation and pollution runoff, altering predator-prey dynamics, or

facilitating the spread of invasive species (Bowman et al. 2010; Bennett 2017; Mumma et al.

2019; Cerqueira et al. 2021). When a univariate model is used to test how a biodiversity

component is related to road cover, the estimated effect combines the direct and indirect effects.

In some cases, these effects can offset one another, obscuring different causal mechanisms and

leading to the conclusion that road cover has no impact.

Second, many environmental conditions covary along urbanization gradients – for

example, road extent with housing density or light pollution – confounding the interpretation of

the effect of any one axis of environmental variation (Moll et al. 2019). Including multiple

covariates in a regression model without a causal rationale can introduce bias. For example, if

light level mediates the effect of road cover on biodiversity, including light as a covariate blocks

part of the causal pathway and underestimates the total effect of road cover. Conversely, failing

to include confounding variables that affect both road cover and biodiversity can lead to spurious

associations. Statistical models designed without regard to causal assumptions often yield

parameter estimates that conflate true causal effects with noncausal confounding, rendering a

mechanistic interpretation nearly impossible.

One way to improve causal inference is to use *graphical causal models*, which have been

shown effective across the sciences (e.g., Rohrer 2018; McElreath 2020; Arif and MacNeil 2023;

Arif and Massey 2023). Our goal is to show how these models can be leveraged to strengthen

inferences about urbanization effects on biodiversity. Specifically, we highlight one type of

graphical causal model, the *directed acyclic graph* (DAG), as an approach for specifying causal

assumptions about urban systems, including direct effects, indirect effects via mediators, and

confounding paths. We first provide a basic introduction to DAGs in the context of urban

ecology and evolution, using simulated datasets to show how the causal meaning of statistical

estimates depends on assumptions specified in the DAG. We then apply the method to a case

study in urban evolution, building a DAG to represent our causal assumptions and guide

statistical models to test hypotheses about urbanization effects on coat color variation in eastern

gray squirrels (*Sciurus carolinensis*).

Directed acyclic graphs for urbanization effects

Directed acyclic graphs are visual diagrams representing hypothesized causal relationships

among variables. DAGs help clarify causal assumptions and inform decisions about study design

and statistical analysis when the research goal is to understand mechanisms driving observed

patterns (Arif and Massey 2023). DAGs consist of variables connected by arrows, with each

arrow representing a direct causal effect. Importantly, causal pathways in DAGs are

probabilistic: the path $X \rightarrow Y$ means X influences the probability distribution of Y, not that Y is a

deterministic function of X.

Consider a hypothetical example, in which an urban ecologist is interested in the effects

of housing density on bird species diversity. We created an example DAG that might represent a

researcher's causal assumptions related to the research question (Fig. 1). The DAG shows

housing density is expected to affect bird diversity via two pathways: 1) the **direct effect**

Housing density → Species diversity, perhaps representing the expectation that housing

structures can provide nesting habitat for some bird species, and 2) the **indirect effect** *Housing*

density → Green space → Species diversity, which might reflect an ecologist's expectation that

housing structures reduce green space, while more green space increases species diversity. Here,

green space functions as a **mediator**, transmitting an effect of housing density on species

diversity. The **total effect** of a variable on an outcome is simply the sum of its direct and indirect

effects. Thus, based on our example DAG, the total effect of housing density includes its direct

effect plus the effect mediated by green space.

Fig. 1. Hypothetical directed acyclic graph (DAG) for a study of the causal effects of housing density on
bird species diversity. Boxes are measured variables, and arrows represent assumed direct causal effects
between variable pairs. Panels include an unlabeled version (A) and a labeled version (B) identifying the
explanatory (green box) and response (blue box) variables of interest, the causal pathways of interest
(green arrows), and important causal structures to identify when estimating the causal pathways of
interest. Estimating effects of housing density on species diversity requires adjusting for the confounding
effect of human population density (orange box) that opens a backdoor path (orange arrows). In contrast,
protected land is a collider and does not cause bias unless conditioned on. Estimation of the total effect of
housing density requires adjusting for the confounder only, whereas estimation of the direct effect
requires adjusting for the confounder and the mediator (green space).

Two additional causal structures are essential to identify in DAGs: confounders and

colliders. In our example DAG, the variables human population density and novel food are on a

backdoor path connecting housing density and species diversity: *Housing density ← Human*

population density → Novel food → Species diversity (Fig. 1). This path might reflect an

ecologist's expectation that human population density increases both housing density and novel

food (e.g., bird feeders), and that novel food increases species diversity. A backdoor path is any

non-causal pathway involving a variable Z that causally affects an explanatory variable X and the

response variable Y (directly or indirectly), having the general structure $X ← Z → Y$. In this case,

Z is a **confounder** because it affects both X and Y, which can create a spurious association or

mask a true causal relationship between them. When considering the effect of housing density on

species diversity, human population density is a confounder because it affects housing density

directly and species diversity indirectly via novel food.

Another important causal structure in the example DAG involves the variable protected

land: *Green space → Protected land ← Species diversity* (Fig. 1). Here, the amount of protected

land is influenced by green space and species diversity, perhaps reflecting an ecologist's

expectation that protected status is most likely to be prioritized for locations with high species

diversity or extensive green space. In this example, protected land is a **collider**. A collider has the

form $X → Z ← Y$, where the collider Z is affected by X and Y.

The causal structures highlighted here are essential to identify when using statistical

models to estimate causal effects. To correctly estimate the effects of interest, researchers need to

parameterize statistical models in a specific way based on causal assumptions expressed in the

DAG. For example, to correctly estimate the direct and total effects of housing density on species

diversity, it is necessary to adjust for the confounding pathway, typically by including human

population density or novel food (or both) as covariates in a regression model. But care must be

taken to decide which covariates to include in a model. If the direct effect of housing density was

of interest, one needs to adjust not only for the confounding effect of human population density,

but also the mediating effect of green space. In contrast, green space should *not* be included as a

covariate when estimating the total effect of housing density, as it functions as a mediator of

housing density and therefore contributes to the total effect. Similarly, colliders do not cause bias

by default and should not be included as a covariate when they are on a path from the

explanatory variable to the response variable. Including protected land in a regression model

with housing density would bias estimates of the housing density effects.

We used a data simulation exercise to explicitly demonstrate how valid estimation of

causal effects with statistical models depends on causal assumptions in a DAG. We first used our

example DAG (Fig. 1) to generate synthetic datasets in R (R Core Team 2024). On their own,

DAGs do not imply a functional form of causal relationships between variables (e.g., positive-

negative, linear-nonlinear, additive-interactive). For the simulation, we assumed all relationships

were linear and additive, and we defined the direction and strength of each direct effect in the

DAG as standardized regression coefficients (Table 1). We then used the assumed effects to

simulate data for 100 sampling locations, with each variable standardized to mean = 0 and SD =

1. All effects were probabilistic, such that the observed value of each variable at each sampling

location was a function of its causal ancestors plus a random error term drawn from a normal

distribution with mean = 0 and SD = 0.5. We generated a total of 1000 simulated datasets.

For each simulated dataset, we fit statistical models to estimate the effects of housing

density on species diversity. We used the *dagitty* package (Textor et al. 2016) to identify

appropriate **adjustment sets** to estimate the direct and total effects of housing density on species

diversity. An adjustment set consists of the covariates from a DAG that should be included in a

statistical model to produce unbiased estimates of a causal effect of interest (Textor and

Liśkiewicz 2011). There are often multiple possible adjustment sets for a given causal effect, and

in our example, the algorithm used by *dagitty* identified two valid adjustment sets for the direct

effect and two valid adjustment sets for the total effect. For each simulated dataset, we fit a

regression model corresponding to each adjustment set. For comparison, we fit two additional

models commonly used in the literature: 1) a univariate model with housing density as the only

predictor of species diversity, and 2) a full model including all covariates.

The simulation results showed the total effect of housing density on species diversity was

accurately estimated when adjusting for the confounding effect of human population density,

which required including either human population density or novel food in the regression model

(Fig. 2A, B). The direct effect of housing density was accurately estimated when adjusting for

the same confounding pathway, as well as the mediating effect of green space (Fig. 2C, D). In

contrast, the univariate model and full model both produced biased estimates of the housing

density effects (Fig. 2E, F). The univariate model conflates the direct and indirect effects of

housing density with the noncausal associations with human population density and novel food,

leading to a biased estimate that recovers neither of the true effects. The full model approximated

the direct effect of housing density because it adjusts for the confounder (human population

density) and mediator (green space), but it underestimated the direct effect because it also

included the collider, protected land. Overall, the data simulation illustrates how the meaning of

an estimated regression coefficient depends on both the assumed causal relationships among

variables specified in a DAG *and* the specific parameterization of a statistical model.

Fig 2. Histograms of the estimated housing density coefficient from regression models fit to 1000 simulated datasets generated from our example DAG (Fig. 1, Table 1). Each panel (A-F) shows the distribution of the estimated housing density coefficient from a single regression model. Model parameterization is shown as a formula with species diversity (Diversity) as a function of (~) housing density (Housing) and other covariates: human population density (Human), novel food (Food), green space (Green), and protected land (Protected). Vertical dashed lines show the true total and direct effects of housing density on species diversity. Histogram colors distinguish models that produce accurate estimates of the causal effect (green) from models that produce biased effects (orange).

DAGs are useful to make causal assumptions clear and to inform statistical analysis, but their use does not guarantee valid causal inference. A critical assumption when using a DAG is that it specifies the correct causal structure of a system. In practice, researchers should apply

prior knowledge of their study system and consider whether non-trivial causes are missing from

a DAG, especially variables that lie on confounding pathways between the explanatory and

response variables of interest. Importantly, confounding effects should be included in DAGs

even when they involve unmeasurable variables, as there are approaches to address unobserved

confounders (Byrnes and Dee 2025).

At the same time, not all variables need to be included in a DAG. Nearly all causal

pathways can be decomposed into more detailed steps (Shipley 2016), and, like any model,

DAGs are simplifications of reality. If a variable functions solely as a mediator between two

variables, it can often be removed from the DAG without changing the implied causal structure.

Objections may be raised about omitted variables, but we argue that debate about the structure of

DAGs is exactly what makes them so valuable. By forcing researchers to state their causal

assumptions explicitly, DAGs promote transparency, constructive criticism, and ideally stronger

inferences.

Case study: applying DAGs to test causal hypotheses about coat color variation in eastern

gray squirrels

Study species and hypothesized urbanization effects

Eastern gray squirrels (*Sciurus carolinensis*) provide an excellent study species to show how

DAGs can be used to help disentangle mechanisms by which urbanization affects ecological and

evolutionary outcomes. Eastern gray squirrels are common across urban and rural landscapes in

their native range of eastern North America. Individuals tend to have one of two coat color

morphs: gray or black (melanic) inherited in a simple Mendelian fashion (McRobie et al. 2009).

The prevalence of melanism tends to be greater in cities than adjacent rural forests (Gibbs et al.

2019; Cosentino and Gibbs 2022; Cosentino et al. 2023). Multiple hypotheses have been

proposed about the ecological and evolutionary processes maintaining these urban-rural clines

(Cosentino and Gibbs 2022; Cosentino et al. 2023). For example, the melanic morph is visually

more conspicuous than the gray morph, such that predation may select for the more cryptic gray

morph (Proctor et al. 2025). Thus, both predator activity and habitat context (e.g., sources of

refuge from predators, including trees and buildings) could alter the prevalence of melanism.

Additionally, road mortality has been hypothesized to maintain clines in melanism, as the

melanic morph is more visible to humans and underrepresented among roadkill compared to the

gray morph (Gibbs et al. 2019; Proctor et al. 2025; Parlin et al. 2025).

Generating an initial DAG

We created a DAG to represent our causal assumptions for how predator activity, components of

habitat structure that affect refuge availability (building cover, forest cover, and forest

fragmentation), and road cover affect spatial variation in the prevalence of melanism along an

urbanization gradient (Fig. 3A). We expected melanism would be greatest in areas of refugia

from predators (high building cover, high forest cover, low forest fragmentation) and high road

cover, and lowest in areas of high predator activity. Critically, while these variables are all

hypothesized to affect the prevalence of melanism directly (Table 2), they can covary along

urban-rural gradients and confound one another, highlighting the necessity of using a DAG to

identify potential confounds and appropriate adjustments to estimate the effects of interest. For

example, forest cover and forest fragmentation may directly affect the prevalence of melanism

by altering crypsis or availability of refuge, but such habitat alteration could also indirectly affect

the prevalence of melanism by changing predator activity (Bateman et al. 2012).

Fig. 3. Directed acyclic graphs (DAG) representing our scientific model, showing hypothesized causal

relationships between variables that could affect variation in melanism in eastern gray squirrels (*Sciurus*

carolinensis) along urbanization gradients. An initial DAG was generated representing our original

hypotheses (A), and the DAG was revised by dropping direct effects among explanatory variables that

were not empirically supported (B) (see *Statistical analysis* below). Arrows indicate hypothesized direct

effects. Boxes represent measured variables, and circles represent unobserved variables. "U" represents

unobserved effects of people on the prevalence of squirrel melanism, including translocations and hunting

pressure, which are known to occur but remain unmeasured. Arrow color distinguishes pathways that

were estimable with measured variables (black) from pathways that were not estimable due to unobserved

processes (purple).

The foundation of our DAG was human population density, operating on the assumption

that environmental change along urbanization gradients is ultimately driven by people. We

included a direct effect of human population density on building cover (Glover and Simon 1975).

Because roads are the main way to access buildings, we included a direct effect of building cover

on road cover. Buildings and roads were assumed to be direct causes of impervious cover, and

we included a direct effect of human population density on impervious cover to represent other

types of impervious surface that people build (e.g., sidewalks, parking lots). We included a direct

effect of impervious cover on forest cover, as the creation of infrastructure requires tree removal

(Liu et al. 2014). Forest cover extent is known to decrease forest fragmentation (Fahrig 2003,

2017), so we included a direct effect of forest cover on fragmentation. We also included a direct

effect of impervious cover on forest fragmentation, as landscape elements like buildings and

roads cause discontinuities in forest cover, regardless of their effects on the absolute amount of

forest (Liu et al. 2014; Liu et al. 2019). Human population density was included as a direct effect

of forest cover and fragmentation to account for other ways that people remove and fragment

forested areas (e.g., gardening and open parks). We included direct effects of road cover, forest

cover, and fragmentation on mammalian predator activity, as all three have been linked to urban

carnivore distribution and behavior (Ordeñana et al. 2010; Bateman et al. 2012).

291   Although DAGs are useful for identifying and adjusting for confounders, unobserved or

unmeasured confounders still create the potential for bias (Byrnes and Dee 2025). For example,

the high prevalence of melanism among squirrels in some cities may be due in part to historical

introductions by people (i.e., propagule pressure) or reduced hunting pressure in cities

(Cosentino et al. 2023), variables that were not possible to measure. Including unobserved

variables like these in a DAG can be important to make gaps in knowledge explicit and identify

confounding with other measured variables. As such, we added an unobserved variable to our

DAG to represent unmeasured mechanisms by which people could mediate coat color change

("U"). We included a direct effect of human population density on U and a direct effect of U on

the prevalence of melanism (Fig. 3A). By including this unobserved variable in the DAG (Fig.

3A), we were able to identify how backdoor paths which include U as a confounder could be

closed by adjusting for human population density (Table 3).

Study area and sampling

To parameterize the DAG, we used data on coat color variation in eastern gray squirrels across

an urban-rural gradient in Syracuse, New York, USA (43.0469° N, -76.1444° W). Syracuse had

an estimated human population of 146,000 (in 2022) and occurs in a humid continental climate

(Köppen 1936) with four distinct seasons, including hot summers and cold winters. Tree cover

ranges from 4.5% to 47% across the city, averaging 27% (Nowak et al. 2016). The surrounding

rural landscape is a mosaic of suburban areas, small villages, agricultural lands, wetlands, and

woodlands (Supplemental Fig. S1).

We used camera traps and point count surveys to estimate prevalence of color morphs at

49 sites across the urban-rural gradient between September 2021 and October 2022

(Supplemental Fig. S1; Cosentino et al. 2023). Sites were selected in urban greenspaces and rural

woodlots with mature deciduous trees, spanning the urban-rural gradient (range = 1.3–23.7 km

from the city center) and separated by at least 1 km. A single camera trap (Browning Strikeforce

Pro XD) was fastened to a tree 30-50 cm off the ground at each site and activated for 46-379

days (median = 263 days), and daily detection histories were generated for each color morph. We

also conducted 2-7 (median = 5) visual point count surveys at each site, recording the number of

squirrels seen of each color morph during a 3-min period. See Cosentino et al. (2023) for detailed

field methods.

Quantifying measured variables

All measured variables in the DAG other than predator activity were measured via remotely

sensed datasets with the *raster* and *exactextractr* packages in R (Table 2; Baston et al. 2022;

Hijmans et al. 2023). We used a 300-m buffer to summarize variables around each site to

encompass a typical home range size of gray squirrels (<5 ha) (Koprowski et al. 2016).

Mammalian predator activity was indexed using our camera trap imagery and calculated as the

percentage of camera trap days mammalian predators were detected at each site (Bu et al. 2016).

Predators detected by camera traps included coyote (*Canis latrans*), gray fox (*Urocyon*

cinereoargenteus), red fox (*Vulpes vulpes*), fisher (*Pekania pennanti*), domestic dog (*Canis*

familiaris) and domestic cat (*Felis catus*). Given our primary goal was to illustrate the broader

use of DAGs for causal inference, we limited our analysis of predator effects to a general index

of activity across mammalian predator species, acknowledging that the effects of mammalian

predators on the prevalence of squirrel melanism may be species-specific, and that we are not

accounting for avian predation.

Statistical analyses

We tested hypotheses from our DAG (Fig. 3A) in two stages. First, we used Bayesian

generalized linear models to estimate direct effects among measured variables other than squirrel

melanism, selecting appropriate distributions for response variables and adjustment covariates

based on the DAG using the *dagitty* R package. For example, to estimate the direct effect of

building cover on impervious cover, we assumed impervious cover follows a beta distribution

and modeled it as a function of building cover, along with road cover as an adjustment. Models

were fit with STAN via the R package *brms* (Bürkner 2017) with noninformative priors and

standardized predictors. We generated a revised DAG (Fig. 3B) that dropped direct effects when

their posterior distributions broadly overlapped 0 (Supplemental Table S1). Next, we estimated

the direct and total effects of each variable on squirrel melanism using an integrated hierarchical

model combining point count and camera trap data (Kéry and Royle 2020; Cosentino et al.

2023). This model included a process submodel to estimate morph-specific abundances and

proportion melanic at each site, and a detection submodel accounting for temperature effects on

activity. We estimated direct and total effects of measured variables in our DAG on proportion

melanic with linear models, adjusting for DAG-informed covariates as needed. We also fit

univariate models with each measured variable as a predictor of melanism to compare inferences

with direct and total effects. Hierarchical models were fit in JAGS via the *jagsUI* R package

(Plummer 2017, Kellner and Meredith 2024) using mostly noninformative priors, but also with

informative priors for detection and abundance intercepts based on prior knowledge. All models

were run with four chains and 4000 retained iterations after enough warm-up or burn-in to reach

convergence, which was confirmed visually with traceplots and R-hat < 1.01 (Gelman and Hill

2007). Posterior distributions were summarized with means and 95% credible intervals. Details

on statistical models are provided in Supplemental Text S1.

Results and discussion

Human population density had a positive total effect on the prevalence of squirrel melanism,

with prevalence of melanism greatest in more urbanized areas (Fig. 4, Table 3). While this aligns

with previous studies showing a strong cline along the urbanization gradient (Cosentino and

Gibbs 2022; Cosentino et al. 2023), our DAG-informed analyses provided multiple new insights

into the potential mechanisms maintaining the urban-rural cline. Remarkably, the total effects of

building cover, road cover, and impervious cover (all metrics of human impact commonly used

to represent "urbanization effects") on the prevalence of melanism were each *negative* despite

squirrel melanism being most prevalent in the city (Fig. 4, Table 3). These negative total effects

suggest the distribution of the melanic morph is limited by various aspects of physical

infrastructure, even in more urban areas where melanic individuals are generally most prevalent.

Considering the positive total effect of human population density combined with negative total

effects of physical infrastructure on squirrel melanism, our results suggest there remain

unmeasured cause(s) represented by U in our DAG (Fig. 5) that play an important role in

maintaining the urban-rural cline by favoring melanics in the city.

Fig. 4 Parameter estimates for direct effects (green), total effects (blue), and univariate models (orange)

for predictors of coat color (proportion melanic) among eastern gray squirrels (*Sciurus carolinensis*). The

complete posterior distribution is shown for each parameter, along with the mean (black circles) and 95%

credible intervals. All parameter estimates were made with standardized predictors. Not all predictors

have both direct and total effects based on our directed acyclic graph (Fig. 3B).

The statistical models informed by our DAG provided novel insights into specific

mechanisms maintaining the urban-rural cline in melanism. For example, prior research showed

melanic individuals are underrepresented among roadkill, relative to the gray morph—a finding

attributed to humans more easily detecting and avoiding melanics against gray asphalt when

driving (Gibbs et al. 2019, Parlin et al. 2025, Proctor et al. 2025). However, our findings show

road cover had strong negative direct and total effects on the prevalence of melanism across the

landscape (Figs. 4, 5; Table 3). This suggests melanic individuals may be avoiding areas with

greater road cover, and thus previously observed underrepresentation of the melanic morph

among roadkill (Gibbs et al. 2019, Parlin et al. 2025) may be due in part to habitat selection or

behavioral avoidance of roads by melanics, rather than being a sole function of differences in

visibility to people while driving. Isolating the direct and total effects of road cover while

considering confounds (e.g., building cover) provided this new insight into why melanics are

underrepresented among roadkill and contributes more evidence that within cities the prevalence

of squirrel melanism is constrained by high disturbance.

Fig. 5 Direction of estimated direct effects overlaid on our directed acyclic graph. Blue solid lines

indicate positive direct effects, red solid lines indicate negative direct effects, and black dotted lines

indicate effects for which the posterior distribution of the parameter estimate broadly overlapped zero.

Purple arrows show causal pathways which could not be estimated due to unobserved processes.

Parameter estimates for effects on squirrel melanism can be found in Table 3, and parameter estimates

among measured environmental variables can be found in Supplemental Table S1.

Another key finding from our models was the strong negative direct effect of predator

activity on the prevalence of melanism (Figs. 4, 5; Table 3). A previous translocation experiment

showed survival is lower for melanic morph than the gray morph in rural woodlands (Cosentino

et al. 2023), and we know the melanic morph is generally more conspicuous than the gray morph

(Proctor et al. 2025). As such, crypsis and predation pressure could contribute to patterns in

melanism across the urban-rural gradient. This combination of forces might help explain the

direct negative effect of predator activity *and* positive total effect of forest fragmentation on

melanism. Although we did not find strong evidence of a direct effect of fragmentation on

melanism, our analyses support the idea that fragmentation mediates the prevalence of melanism

through its negative effect on predator activity (Fig. 5, Supplemental Table S1), leading to a

positive total effect of fragmentation on melanism (Fig. 4, Table 3). The melanic morph may

benefit from a suppressive effect of fragmentation on predator activity locally, but it is important

to note there was a strong positive direct effect of human population density on predator activity

(Fig. 5, Supplemental Table S1), indicating predator activity was greatest in the city where

melanics are most common. It is possible the melanic morph uses behavioral strategies to

mitigate their predation risk in densely populated areas of the city where predator activity is

greatest (Sarno et al. 2015; Gaynor et al. 2019; Engel et al. 2020). Such strategies could include

different activity patterns in space (including vertically from ground to canopy) and time (diel

patterns), resulting in differing susceptibility to predation. Because vertical niche shifts have

been linked to competition and predation, climate change, and urbanization (Rankin et al. 2018;

Basham and Scheffers 2020; Borden et al. 2021; Gámez and Harris 2022), future research should

investigate differences in spatial-temporal behavioral patterns between color morphs and how it

may impact the landscape of fear. Additional analyses are also needed to examine the possibility

that predator impacts on squirrel melanism vary among predator species.

Despite these novel insights regarding mechanisms contributing to the urban-rural cline

in melanism, it is intriguing that melanism is most prevalent in Syracuse yet constrained by

components of physical infrastructure that are widespread in cities. This seemingly contradictory

pattern may be due in part to the interaction of historic and contemporary forces, including

ecological, evolutionary, and social (Des Roches et al. 2021). For example, the melanic morph,

once common across the forested landscape, was extirpated in many areas during the period of

European colonialism when extensive hunting accompanied deforestation and logging in old

growth forests near settlements (Robertson 1973; Cronon 1991; Benson 2013; Thompson et al.

2013). Evidence suggests urban areas may have functioned as refugia for eastern gray squirrels

in general, including the melanic morph (Benson 2013; Gibbs et al. 2019). Indeed, changing

attitudes about nature led to efforts to deliberately introduce squirrels to cities starting in the late

1800s, celebrating them as "our most loved" species in cities, and establishing a social attitude of

tolerance and even admiration (Benson 2013). Despite well-known cases of introductions of the

melanic morphs to cities (e.g., Washington DC, Fischman et al. 2021), many squirrel

translocations are undocumented. We strongly suspect human-mediated transport and bans on

hunting in densely populated cities, including Syracuse, have played important roles in

contributing to the maintenance of urban-rural clines in squirrel melanism.

Although such social forces can be difficult to measure, our DAG illustrates it can be

important to include unmeasured processes in a causal modeling framework. As just one

example, consider the direct negative effect of predator activity on melanism inferred from our

analyses. Based on the DAG in Fig. 3B, the adjustment set we used to estimate the direct effect

of predator activity on melanism included human population density and fragmentation.

Adjusting for human population density was needed in part to close the backdoor path *Predator*

activity ← Human population density → U → Squirrel melanism. If the *Human population*

density → U → Squirrel melanism pathway was *not* included in the DAG, the direct effect of

predator activity on squirrel melanism could be estimated without adjusting for human

population density. Doing so would lead to a biased estimate of the direct effect of predator

activity due to confounding with the unmeasured variable, U. Indeed, when we estimate the

direct effect of predator activity on melanism without adjusting for human population density,

the estimated effect is no longer negative (posterior mean = 0.04, 95% CI = -0.22, 0.35). This

positive bias is due to a positive effect of people on predator activity and the residual positive

association between human population density and melanism operating through the unobserved

processes. Graphical causal models, such as DAGs, highlight one way in which unobserved

processes in urban systems – known or unknown – can be explicitly considered when estimating

effects of interest.

A striking demonstration of the value of graphical causal models is seen when comparing

estimated effects when appropriately adjusted based on a DAG versus univariate models

commonly seen in urban ecology and evolution. For example, had we relied solely on univariate

models to estimate the association between squirrel melanism and common urbanization metrics,

we might have erroneously concluded that melanism is unrelated to *any* of the measured

variables in our DAG other than human population density, as credible intervals for all univariate

predictors other than human density overlapped zero (Fig. 4; Table 3). This finding raises

significant concerns about using single-variable indices of urbanization. For example,

impervious cover is commonly used to represent urbanization (Szulkin et al. 2020), yet our

univariate model with impervious cover resulted in a posterior distribution that was largely

negative, implying melanism is lowest in areas of high impervious cover (Fig. 4; Table 3). Had

we used impervious cover alone as an index of urbanization, we might have concluded there was

no evidence for an urban-rural cline, or even weak evidence for a reversed cline where melanism

is greater in rural areas. Indeed, our own previous studies using impervious surface as a surrogate

for urbanization likely underestimated true clines in melanism (Gibbs et al. 2019; Cosentino and

Gibbs 2022). If using single variables as proxies of urbanization to describe biodiversity patterns

along urbanization gradients cannot be avoided, our results suggest it is best to use variables that

are causal ancestors of environmental change in cities, namely the density of people (Fig. 5).

Distance to city center or composite indices can also be used to describe spatial patterns of

biodiversity along urbanization gradients (Moll et al. 2019, Alberti and Wang 2022), but we urge

caution about using these variables for causal inference as they often aggregate multiple

underlying mechanisms.

Conclusion

Overall, our research demonstrates that urban impacts on biodiversity are complex and involve a

network of causal pathways. Environmental variables commonly used to represent urbanization

may at best describe coarse patterns with respect to urbanization effects in ecology and

evolution, but they are nearly impossible to interpret mechanistically (Moll et al. 2019). We show

that graphical causal models, such as DAGs, can be an effective tool to clarify causal

assumptions and inform the design of statistical models to make inferences about a variable's

mechanistic effects. Causal modeling frameworks present great opportunity for disentangling the

mechanisms at play in urban systems and understanding the causal drivers of biodiversity within

and among cities.

Acknowledgments

505   **Table 1.** Standardized regression coefficients representing the direction and magnitude of direct effects in

506   our hypothetical DAG (Fig. 1) used to simulate observed datasets.

Explanatory variable	Response variable	Standardized coefficient
Human population density	Novel food	0.9
Human population density	Housing density	0.8
Housing density	Green space	-1.0
Housing density	Species diversity	0.3
Novel food	Species diversity	0.4
Green space	Species diversity	0.8
Green space	Protected land	1.0
Species diversity	Protected land	0.8

507   **Table 2.** Environmental variables hypothesized to affect the prevalence of melanism in eastern gray

508   squirrels (*Sciurus carolinensis*) and data used to measure those variables.

Environmental variable	Rationale	Hypothesized Direction	References	Variable data source
Human population density	The foundational driver of urbanization, increasing building and road cover, driving deforestation and habitat fragmentation, and causing many unmeasured changes.	(+)	(Glover and Simon 1975)	NASA's CIESIN gridded human population dataset (30 arcsec; Center For International Earth Science Information Network-CIESIN-Columbia University 2017)
Building cover	Alters habitat structure and availability by reducing tree cover, but provides nesting sites, novel microclimates, and shading that may be important for crypsis.	(+)	(Benson 2013; Parker and Nilon 2008)	Cornell University Geospatial Information Repository
Road cover	Roadkill is a major driver of squirrel mortality in urban contexts particularly roads with $\geq$ 30 mph speed limit. Melanic individuals are more visible to drivers on roads, and they have lower mortality rates from vehicular collisions than the gray morph.	(+)	(McCleery et al. 2008; Gibbs et al. 2019; Parlin et al. 2025; Proctor et al. 2025)	Cover of roads with at least 30 mph speed limit from NYS GIS Resources repository for public data (https://data.gis.ny.gov)
Forest cover	Increases resource availability, including food, nests, and refuge from predators.	(+)	(Parker and Nilon 2008; Koprowski et al. 2016; Merrick et al. 2016)	ESA WorldCover dataset derived from Sentinel-1 and Sentinel-2 satellite imagery (10 m resolution; Zanaga et al. 2022)
Forest fragmentation	Restricts movements and may affect fitness of color morphs differently by altering predator activity and crypsis of each morph assuming fragmented areas are more visually open.	(-)	(Parker and Nilon 2008; Koprowski et al. 2016)	Measured as disjunct core area density based on forest cover from ESA WorldCover dataset derived from Sentinel-1 and Sentinel-2 satellite imagery (Zanaga et al. 2022)

| Mammalian predator activity | Mammalian predator activity alters predation risk as well as the landscape of fear. Differences in crypsis or behavior may cause morphs to be affected differently by predator activity. | (-) | (Gibbs et al. 2019; Cosentino and Gibbs 2022; Proctor et al. 2025) | Quantified from camera trap detections |

Table 3. Posterior means, 95% credible intervals, and adjustment sets for parameter estimates of direct effects, total effects, and univariate models for predictors of coat color (proportion melanic) among eastern gray squirrels (*Sciurus carolinensis*). Separate models were fit to estimate direct and total effects of predictor variables on proportion melanic with appropriate adjustment covariates based on our directed acyclic graph (Fig. 3B). Univariate estimates were based on models with only the predictor variable and no adjustment covariates. Predictor variables included human population density (Human), building cover (Building), road cover (Road), impervious cover (Impervious), forest cover (Forest), forest fragmentation (Frag), and mammalian predator activity (Predator).

Direct Effects				
Predictor	*Mean*	*LCL*	*UCL*	*Adjustments*
Human	-	-	-	-
Building	0.02	-0.29	0.32	Human, Road, Forest, Frag
Road	**-0.69**	**-1.01**	**-0.39**	Human, Building, Forest, Frag
Impervious	-	-	-	-
Forest	**0.47**	**0.17**	**0.78**	Human, Building, Road, Frag
Frag	0.28	-0.07	0.64	Human, Impervious, Forest, Predator
Predator	**-0.54**	**-0.83**	**-0.26**	Human, Frag

Total Effects				
Predictor	*Mean*	*LCL*	*UCL*	*Adjustments*
Human	**0.42**	**0.18**	**0.67**	-
Building	**-0.33**	**-0.54**	**-0.10**	Human
Road	**-0.28**	**-0.54**	**-0.01**	Building
Impervious	**-0.80**	**-1.42**	**-0.19**	Human, Building, Road
Forest	0.16	-0.08	0.42	Impervious
Frag	**0.41**	**0.07**	**0.74**	Human, Impervious, Forest
Predator	**-0.54**	**-0.83**	**-0.26**	Human, Frag

Univariate Estimates				
Predictor	*Mean*	*LCL*	*UCL*	*Adjustments*
Human	**0.42**	**0.18**	**0.67**	**-**
Building	-0.04	-0.22	0.15	**-**
Road	-0.15	-0.34	0.02	**-**
Impervious	-0.15	-0.35	0.04	**-**
Forest	0.20	-0.01	0.42	**-**
Frag	0.15	-0.05	0.35	**-**
Predator	-0.14	-0.39	0.12	**-**

Literature cited

Alberti M, Palkovacs EP, Roches SD, De Meester L, Brans KI, et al (2020) The complexity of

urban eco-evolutionary dynamics. BioScience 70(9):772–793

Alberti M, Wang T (2022) Detecting patterns of vertebrate biodiversity across the

multidimensional urban landscape. Ecology Letters 25:1027–1045

Arif S, MacNeil MA (2023) Applying the structural causal model framework for observational

causal inference in ecology. Ecological Monographs 93(1):e1554

Arif S, Massey MDB (2023) Reducing bias in experimental ecology through directed acyclic

graphs. Ecology and Evolution 13(3):e9947

Aronson MFJ, La Sorte FA, Nilon CH, Katti M, Goddard MA, et al (2014) A global analysis of

the impacts of urbanization on bird and plant diversity reveals key anthropogenic drivers.

Proc R Soc B 281(1780):20133330–20133330

Basham EW, Scheffers BR (2020) Vertical stratification collapses under seasonal shifts in

climate. J Biogeogr 47:1–11

Baston D, ISciences, LLC (2022) exactextractr: Fast extraction from raster datasets using

polygons (version 0.9.1) [computer software]. https://cran.r-

project.org/web/packages/exactextractr/index.html

Bateman PW, Fleming PA, Fleming T, Comber SL (2012) Big city life: Carnivores in urban

environments. J Zool 287(1):1–23

Bennett VJ (2017) Effects of road density and pattern on the conservation of species and

biodiversity. Curr Landsc Ecol Rep 2(1):1–11

Benson E (2013) The urbanization of the eastern gray squirrel in the United States. J Am Hist

100(3):691–710

Borden JB, Bohlman S, Scheffers BR (2021) Niche lability mitigates the impact of invasion but

not urbanization. Oecologia 198:1–10

Bowman J, Ray JC, Magoun AJ, Johnson DS, Dawson FN (2010) Roads, logging, and the large-

mammal community of an eastern Canadian boreal forest. Can J Zool 88(5):454–467

Bu H, Wang F, McShea WJ, Lu Z, Wang D, et al (2016) Spatial co-occurrence and activity

patterns of mesocarnivores in the temperate forests of Southwest China. PLoS ONE

11(10):e016427

Bürkner P-C (2017) brms: An R package for Bayesian multilevel models using Stan. J Stat Softw

80:1–28

Byrnes JEK, Dee LE (2025) Causal inference with observational data and unobserved

confounding variables Ecol Lett 28:e70023

Center for International Earth Science Information Network-CIESIN-Columbia University

(2017) Gridded population of the world, version 4 (GPWv4): Population density, revision 11

[dataset]. Palisades, NY: Socioeconomic Data and Applications Center (SEDAC)

Cerqueira RC, de Rivera OR, Jaeger JAG, Grilo C (2021) Direct and indirect effects of roads on

space use by jaguars in Brazil. Sci Rep 11:22617

Cosentino BJ, Gibbs JP (2022) Parallel evolution of urban–rural clines in melanism in a

widespread mammal. Sci Rep 12:1752

Cosentino BJ, Vanek JP, Gibbs JP (2023) Rural selection drives the evolution of an urban–rural

cline in coat color in gray squirrels. Ecol Evol 13(10):e10544

Cronon W (1991) Nature's metropolis: Chicago and the Great West. WW Norton & Company,

New York

Des Roches S, Brans KI, Lambert MR, Rivkin LR, Savage AM, et al (2021) Socio-eco-

evolutionary dynamics in cities. Evol Appl 14(1):248–267

Engel LD, Carlen EJ, Losos JB, Winchell KM (2020) Eastern gray squirrels (*Sciurus*

carolinensis) differ in abundance and response to humans across urban habitats of St. Louis.

Urb Nat 33: 1–16

Fahrig L (2003) Effects of habitat fragmentation on biodiversity. Annu Rev Ecol Evol Syst

34(1):487–515

Fahrig L (2017) Ecological responses to habitat fragmentation per se. Annu Rev Ecol Evol Syst

48:1–23

Fischman B, Cosentino B, Gibbs J (2021) City squirrels look different. Is evolution driving a

color change? Discover Magazine, City Science Salon.

https://www.discovermagazine.com/planet-earth/city-squirrels-look-different-is-evolution-

driving-a-color-change

Francis CD, Kleist NJ, Ortega CP, Cruz A (2012) Noise pollution alters ecological services:

Enhanced pollination and disrupted seed dispersal. Proc R Soc B 279(1739):2727–2735

Gámez S, Harris NC (2022) Conceptualizing the 3D niche and vertical space use. Trends Ecol

Evol 37:953-962

Gaynor KM, Brown JS, Middleton AD, Power ME, Brashares JS (2019) Landscapes of fear:

Spatial patterns of risk perception and response. Trends Ecol Evol 34(4):355–368

Gelman A (2006) Prior distributions for variance parameters in hierarchical models (comment on

article by Browne and Draper). Bayesian Anal 1(3):515–534

Gelman A, Hill J (2007) Data analysis using regression and multilevel/hierarchical models.

Cambridge University Press

Gibbs JP, Buff MF, Cosentino BJ (2019) The biological system—urban wildlife, adaptation, and

evolution: Urbanization as a driver of contemporary evolution in gray squirrels (*Sciurus*

carolinensis). In: Hall MP, Balogh S (eds) Understanding Urban Ecology. Springer, pp 269–

286

Glover DR, Simon JL (1975) The effect of population density on infrastructure: The case of road

building. Econ Dev Cult Change 23(3):453–468

Hijmans RJ, van Etten J, Sumner M, Cheng J, Baston D, et al (2023) raster: Geographic data

analysis and modeling (version 3.6-20) [computer software].

Kellner K, Meredith M (2024) jagsUI: A wrapper around "rjags" to streamline "JAGS" analyses

(version 1.6.2) [computer software]. https://cran.r-

project.org/web/packages/jagsUI/index.html

Kéry M, Royle JA (2020) Applied hierarchical modeling in ecology: Analysis of distribution,

abundance and species richness in R and BUGS: Volume 2: Dynamic and advanced models.

Academic Press

Kok ACM, Berkhout BW, Carlson NV, Evans NP, Khan N, et al (2023) How chronic

anthropogenic noise can affect wildlife communities. Front Ecol Evol 11

Köppen W (1936) The geographical system of climate. In: Koppen W, Geiger R (eds) Handbook

of climatology. Gebruder Borntraeger, Berlin

Koprowski J, Munroe K, Edelman A (2016) Gray not grey: The ecology of *Sciurus carolinensis*

in their native range in North America. In: Shuttleworth M, Lurz P, Gurnell J (eds) Grey

Squirrels: Ecology and Management of an Invasive Species in Europe. European Squirrel

Initiative

Lemoine NP (2019) Moving beyond noninformative priors: Why and how to choose weakly

informative priors in Bayesian analyses. Oikos 128(7):912–928

Liu J, Coomes DA, Gibson L, Hu G, Liu J, et al (2019) Forest fragmentation in China and its

effect on biodiversity. Biol Rev 94(5):1636–1657

Liu S, Dong Y, Deng L, Liu Q, Zhao H, et al (2014) Forest fragmentation and landscape

connectivity change associated with road network extension and city expansion: A case study

in the Lancang River Valley. Ecol Indic 36:160–168

McCleery RA, Lopez RR, Silvy NJ, Gallant DL (2008) Fox squirrel survival in urban and rural

environments. J Wild Mgmt 72:133–137

McDonnell MJ, Hahs AK (2008) The use of gradient analysis studies in advancing our

understanding of the ecology of urbanizing landscapes: Current status and future directions.

Landsc Ecol 23:1143–1155

McDonnell MJ, Pickett STA (1990) Ecosystem structure and function along urban-rural

gradients: An unexploited opportunity for ecology. Ecology 71(4):1232–1237

McElreath R (2020) Statistical rethinking: A Bayesian course with examples in R and STAN.

CRC Press

McRobie H, Thomas A, Kelly J (2009) The genetic basis of melanism in the gray squirrel

(*Sciurus carolinensis*). J Hered 100(6):709–714

Merrick M, Evans K, Bertolino S (2016) Urban grey squirrel ecology, associated impacts, and

management challenges. In: Shuttleworth M, Lurz P, Gurnell J (eds) Grey Squirrels: Ecology

and Management of an Invasive Species in Europe. European Squirrel Initiative,

Warwickshire, UK

Moll RJ, Cepek JD, Lorch PD, Dennis PM, Tans E, et al (2019) What does urbanization actually

mean? A framework for urban metrics in wildlife research. J Appl Ecol 56(5):1289–1300

Mumma MA, Gillingham MP, Johnson CJ, Parker KL (2019) Functional responses to

anthropogenic linear features in a complex predator-multi-prey system. Landsc Ecol

34(11):2575–2597

Nowak DJ, Hoehn RE, Bodine AR, Greenfield EJ, O'Neil-Dunne J (2016) Urban forest

structure, ecosystem services and change in Syracuse, NY. Urban Ecosyst 19(4):1455–1477

640 Ordeñana MA, Crooks KR, Boydston EE, Fisher RN, Lyren LM, et al (2010) Effects of

641  urbanization on carnivore species distribution and richness. J Mammal 91(6):1322–1331

642 Parker TS, Nilon CH (2008) Gray squirrel density, habitat suitability, and behavior in urban

643  parks. Urban Ecosyst 11(3):243–255

644 Parlin AF, Cosentino BJ, Lehtinen RM, McDonald JE, Sinclair ECC, Gibbs JP (2025) Road

645  mortality contributes to the evolution of an urban-rural cline in squirrel coat color.

646  Evolutionary Applications 18(5):e70109.

647 Plummer M (2017) JAGS version 4.3.0 user manual. https://sourceforge.net/projects/mcmc-

648  jags/files/Manuals/4.x/

649 Poisson MK, Gebresenbet F, Butler AR, Tate P, Bergeron DH, Moll RJ (2024) The way

650  "urbanization" is defined has strong implications for its effects on mammal abundance.

651  Urban Ecosyst 27:2367-2380.

652 Proctor J, Bryan A, Cosentino BJ, Gibbs JP (2025) Crypsis in a polymorphic mammal along an

653  urbanization gradient. Urban Ecosyst 28:94

654 R Core Team (2024) R: A language and environment for statistical computing. R Foundation for

655  Statistical Computing. Vienna, Austria. https://www.R-project.org/

656 Rankin EEW, Knowltonid JL, Gruner DS, Flaspohler DJ, Giardina CP, et al (2018) Vertical

657  foraging shifts in Hawaiian forest birds in response to invasive rat removal. PLoS ONE 10

658 Ritchie H, Roser M (2018) Urbanization. Our World in Data.

659  https://ourworldindata.org/urbanization

Robertson GI (1973) Distribution of color morphs of *Sciurus carolinensis* in eastern North

America. MS Thesis, University of Western Ontario, London

Rohrer JM (2018) Thinking clearly about correlations and causation: Graphical causal models

for observational data. Adv Methods Pract Psychol Sci 1(1):27–42

Santangelo JS, Ness RW, Cohan B, Fitzpatrick CR, Innes SG, et al (2022) Global urban

environmental change drives adaptation in white clover. Science 375:1275–1281

Sarno RJ, Parsons M, Ferris A (2015) Differing vigilance among gray squirrels (*Sciurus*

carolinensis) along an urban–rural gradient on Long Island. Urban Ecosyst 18(2):517–523

Shipley B (2016) Cause and Correlation in Biology. 2nd ed. Cambridge University Press

Szabó B, Korányi D, Gallé R, Lövei GL, Bakonyi G, et al (2023) Urbanization decreases species

richness, and increases abundance in dry climates whereas decreases in wet climates: A

global meta-analysis. Sci Total Environ 859:160145

Szulkin M, Garroway CJ, Corsini M, Kotarba A, Dominoni D (2020) How to quantify

urbanization when testing for urban evolution? In: Szulkin M, Munshi-South J, Charmantier

A (eds) Urban Evolutionary Biology. Oxford University Press

Textor J, Liśkiewicz M (2011) Adjustment criteria in causal diagrams: An algorithmic

perspective. Proceedings of the 27[th] Conference on Uncertainty in Artificial Intelligence

(UAI 2011). AUAI Press, pp. 681–688

Textor J, van der Zander B, Gilthorpe, MS, Liskiewicz M, Ellison GT (2016) Robust causal

inference using directed acyclic graphs: the R package '*DAGitty*'. Int J Epidemiol

45(6):1887–1894

Thompson JR, Carpenter DN, Cogbill CV, Foster DR (2013) Four centuries of change in

northeastern United States forests. PLoS ONE 8(9):e72540

Winchell KM, Reynolds RG, Prado-Irwin SR, Puente-Rolón AR, Revell LJ (2016) Phenotypic

shifts in urban areas in the tropical lizard *Anolis cristatellus*. Evolution 70(5):1009–1022

Zanaga D, Van De Kerchove R, Daems D, De Keersmaecker W, Brockmann C, et al (2022) ESA

WorldCover 10 m 2021 v200 (version v200) [dataset]. Zenodo

https://doi.org/10.5281/zenodo.7254221

Statements and Declarations

Funding

This research was funded by the U.S. National Science Foundation (DEB-2018140, DEB-

2018249).

Data Availability

Data and code are available in the following repository:

https://github.com/bcosentino/urbanCausalInference