Ecological and evolutionary impacts of urbanization: applying a causal modeling

framework

Jesse B. Borden[1*], James P. Gibbs[2], John P. Vanek[3], Bradley J. Cosentino[1]

[1]Biology Department, Hobart and William Smith Colleges, 300 Pulteney Street, Geneva, NY,

14456, USA

[2]Department of Environmental Biology, The State University of New York College of

Environmental Science and Forestry, 1 Forestry Drive, Syracuse, NY 13210, USA

[3]New York Natural Heritage Program, The State University of New York College of

Environmental Science and Forestry, 625 Broadway, Albany, NY 12207, USA

[*]**Corresponding author email:** jessebborden@gmail.com

Abstract

Urbanization, as the fastest growing land use type, shapes biodiversity through a suite of abiotic

and biotic changes driven by the profound alteration of environmental conditions in built

environments. Understanding the mechanisms behind biodiversity patterns in urban areas

requires disentangling the direct and indirect pathways through which urbanization impacts

ecological and evolutionary dynamics. We show how a structural causal modeling framework

can be used to provide insight into processes generating biodiversity patterns in urban contexts

by building structural equation models to disentangle direct and indirect pathways by which

urbanization affects abundance and coat color variation in eastern gray squirrels (*Sciurus*

carolinensis). Integrating camera traps and point count surveys across 49 sites, we observed

pronounced urban-rural gradients in squirrel abundance and color variation. Human population

density affected abundance of color morphs by mediating resource availability, predator activity,

habitat amount, and fragmentation. But markedly different mechanisms drove abundance of

color morphs across the landscape. Our findings reveal how explicit modeling of multiple

pathways of environmental change can deepen our mechanistic understanding of how

urbanization alters biodiversity patterns.

Key words

Urban evolution, urbanization, causal modeling, path analyses, melanism, trait variation, urban

mechanisms, gray squirrel, *Sciurus carolinensis*

Authors' contributions

Jesse B. Borden: writing - original draft (lead); conceptualization, investigation, methodology, data curation, visualization, writing - reviewing and editing (equal). **James P. Gibbs:** conceptualization, funding acquisition, investigation, methodology, writing- reviewing and editing (equal). **John P. Vanek:** conceptualization, investigation, methodology, writing – review and editing. **Bradley J. Cosentino:** conceptualization, funding acquisition, investigation, methodology, formal analysis, project administrator, supervision, writing – review and revision (lead)

1. Introduction

Urbanization represents the fastest growing land use type and poses formidable challenges to biodiversity (Aronson et al., 2014; Ritchie & Roser, 2018; Szabó et al., 2023). At a fundamental level, urbanization consists of a suite of abiotic and biotic changes driven by high concentrations of people and the physical environment they build (Ritchie & Roser, 2018) (Fig. 1a). The ecological and evolutionary consequences of urbanization are often studied using the urban-rural gradient approach in which components of biodiversity are compared along a landscape-scale sampling area from the urban center to rural environments (Mcdonnell & Hahs, 2008; McDonnell & Pickett, 1990). This approach has been used to document associations of population size, species distributions, community composition, and genetic and phenotypic trait variation with urbanization. Although the urban-rural gradient approach has been helpful to clarify spatial patterns of biodiversity associated with urbanization, reliance on coarse metrics of urbanization (e.g., distance to city center, human population density, impervious cover; Moll et al. 2019) provides little insight into the underlying mechanisms driving urbanization effects on biodiversity.

To understand the mechanisms underlying biodiversity patterns in urban areas, it is helpful to consider that changes associated with urbanization cascade through multiple pathways, impacting biodiversity in complex ways (Alberti et al., 2020). For example, road networks, which are most dense in urban centers with high human settlement densities, can directly affect biodiversity through vehicular collisions and noise pollution (Bennett, 2017; Francis et al., 2012; Kok et al., 2023). Road networks can also *indirectly* affect biodiversity in multiple ways— through increasing habitat fragmentation and pollution runoff, altering predator-prey dynamics, constraining space use through habitat loss, or facilitating the spread of invasive species

(Bennett, 2017; Bowman et al., 2010; Cerqueira et al., 2021; Mumma et al., 2019). As such, axes

of environmental change along urbanization gradients are composed of a system of interacting

and cascading effects which directly and indirectly drive ecological and evolutionary patterns

(Fig. 1). Understanding the mechanisms generating spatial patterns of biodiversity along urban-

rural gradients requires approaches that disentangle the direct and indirect causal pathways

linking biodiversity to people.

Our study had two goals: first, to test hypotheses about how multiple environmental

changes interact to generate spatial patterns in abundance and trait variation along urbanization

gradients; and second, to illustrate how a structural causal model framework can effectively be

used to do so. We use eastern gray squirrels (*Sciurus carolinensis*) as our study organism.

Eastern gray squirrels are ubiquitous in urban and rural habitats across their native range of

eastern North America. Previous studies have shown strong urban-rural gradients in abundance

and coat color in eastern gray squirrels, with both abundance and the prevalence of melanism

greater in more urban environments than adjacent rural areas (Cosentino et al., 2023; Cosentino

& Gibbs, 2022; Gibbs et al., 2019). Yet, the processes generating these spatial patterns are

unclear, with multiple hypotheses having been proposed. For example, a translocation

experiment revealed mortality in rural areas was greater for the melanic than the gray morph,

leading to the hypothesis that selection pressures on coat color differ between morphs in rural

areas (Cosentino et al., 2023). Differential selection between color morphs may be due to the

melanic morph being generally more visible than the gray morph, which could amplify its

predation risk, but also decrease the risk of vehicular collisions due to increased visibility to

vehicle drivers (Gibbs et al., 2019). Thus, evidence suggests changes in crypsis may alter

predation pressure and road mortality in ways that contribute to urban-rural clines in melanism

(Cosentino & Gibbs, 2022; Gibbs et al., 2019). A mechanistic understanding of how these factors

operate in concert requires a method to disentangle direct and indirect effects among predation

pressure, road mortality risk, and related components of environmental change along the

urbanization gradient.

Causal inference in observational designs typical of urban ecology and evolution first

requires a scientific model that clearly defines hypothesized causal relationships among variables

in the system based on prior knowledge (Arif & MacNeil, 2023; Arif & Massey, 2023;

McElreath, 2020). Our research illustrates how directed acyclic graphs (DAG) can be used to

define a scientific model based on prior knowledge of the urban system. Cities are complex

systems that have a variety of interacting and covarying environmental gradients, so DAGs are

essential for communicating the causal relationships assumed about the system. We apply

structural equation models (Grace, 2006; Grace et al., 2012), as one possible statistical approach,

to estimate the relationships (or lack thereof) among variables implied by the DAG. We think

this approach can be harnessed and elaborated to better understand the mechanisms generating

spatial patterns of biodiversity along urbanization gradients. Because abiotic and biotic factors

are spatially heterogeneous across the urban-rural gradient, disentangling their relationships can

also help elucidate the processes generating fine-scale patterns of biodiversity variation within

cities.

2. Methods

2.1 Study site

This research was conducted across an urban-rural gradient surrounding Syracuse, New York,

USA (43.0469° N, -76.1444° W). Syracuse has ~150,000 people and is characterized by a humid

continental climate (Koppen, 1936) with four distinct seasons, including hot summers and cold

winters. Tree cover ranges from 4.5% to 47% across the city, averaging 27% (Nowak et al.,

2016), and the surrounding rural landscape is a mosaic agriculture and forest (Fig. S1). .

2,2 Study species and hypothesized drivers

Eastern gray squirrels are an excellent study organism to disentangle direct and indirect paths by

which urbanization impacts ecological (e.g., abundance) and evolutionary (e.g., trait variation)

outcomes. Eastern gray squirrels are common across urban and rural landscapes in eastern north

America, and individuals tend to have one of two common coat colors – gray or black (melanic)

— inherited in a simple Mendelian fashion (H. McRobie et al., 2009; H. R. McRobie et al.,

2019). Eastern gray squirrels repeatedly show clines in both abundance and prevalence of

melanism across urban-rural gradients (Cosentino & Gibbs, 2022; Merrick et al., 2016),

providing an excellent opportunity to disentangle mechanisms underlying such patterns. We

hypothesize these patterns would be driven by multiple, covarying axes of biotic and abiotic

variation. Specifically, we hypothesized human population density, building cover, forest cover,

forest fragmentation, the distance to roads (road distance), and mammalian predator activity

would affect squirrel abundance and patterns in trait variation across the landscape. We selected

these drivers based on previous research, and hypothesized mechanisms by which they affect

each other and have downstream impacts on squirrel abundance and coat color prevalence (Tab.

1). For example, human population density influences resources available to squirrels (through

food provided through bird feeders and nesting sites in buildings) (Benson, 2013) and building

cover can also indirectly affect squirrel abundance by reducing forest cover. Forest cover and

fragmentation are linked to nesting availability, food resources and habitat connectivity (Merrick

et al., 2016; Parker & Nilon, 2008). Proximity to roads increases road mortality risk, which plays

an important role in squirrel mortality in urban environments (Gibbs et al., 2019). Lastly,

mammalian predator activity drives predation risk and influences the landscape of fear,

potentially altering squirrel behavior and distribution (Gaynor et al., 2019).

2.3 Camera trapping and point count surveys

We used camera traps and point count surveys to estimate the abundance of gray squirrel color

morphs at 49 sites between 16 September 2021 and 13 October 2022 (Cosentino et al., 2023).

Because squirrel populations are limited by tree cover, sites were selected in urban greenspaces

and rural woodlots with mature deciduous trees (Merrick et al., 2016; Parker & Nilon, 2008).

Site locations were chosen to span the urban-rural gradient ranging 1.3–23.7 km from the city

center (median = 9.3 km), and separated by at least 1 km. At each site we deployed a single

camera trap (Browning Strikeforce Pro XD) between September 2021 and March 2022, fastening

the camera to a tree 30-50 cm off the ground. Cameras were active for 46-379 days (median =

263 days), with daily detection histories generated for each color morph by a single observer. We

also conducted visual point count surveys at each site between October 2021 and October 2022.

During each survey, a single observer recorded the number of squirrels seen of each color morph

during a 3-min period. Sites were surveyed 2-7 times each (median = 5).

2.4 Environmental covariates

The environmental covariates selected for our analyses (Tab.1) were measured using the

raster and *exactextractr* packages in R (Baston et al., 2022; Hijmans et al., 2023). We used a

300-m buffer to summarize environmental covariates around each site to encompass a typical

home range size of gray squirrels (<5 ha) (Koprowski et al., 2016). Human population density

was calculated using NASA's CIESIN gridded human population dataset (Center For

International Earth Science Information Network-CIESIN-Columbia University, 2017). Building

cover was calculated using the building footprint dataset from Cornell University Geospatial

Information Repository. Road distance was taken by measuring the distance to the nearest road

with a speed limit greater than 30 mph and measured using the NYS GIS Resources repository

for public data (https://data.gis.ny.gov). We isolated roads with high-speed limits because

vehicular collisions are rare on roads with speed limits <30 mph (unpublished data). Forest cover

was measured using the tree cover band from ESA WoldCover dataset derived from Sentinel-1

and Sentinel-2 satellite imagery (Zanaga et al., 2022). We quantified forest fragmentation using

the WoldCover dataset by calculating the number of disjunct cores within a 1-km radius of each

site using the *landscapemetrics* package in R (Hesselbarth et al., 2019). We used a 1-km radius

to represent potential fragmentation effects as this scale is sufficient to capture fragmentation (a

landscape-scale process) and it encompasses possible long distance dispersal events typical of

males during mating seasons (Cosentino & Gibbs, 2022). Mammalian predator activity was

indexed using our camera trap imagery and calculated as the percentage of camera trap days

mammalian predators were detected at each site (Bu et al., 2016). Predators detected by camera

trap included coyote (*Canis latrans*), gray fox (*Urocyon cinereoargenteus)*, red fox (*Vulpes

vulpes*), fisher (*Pekania pennanti*), domestic dog (*Canis familiaris*) and domestic cat (*Felis

catus*).

2.5 Modeling framework

We built a structural equation model (SEM; Grace, 2006) to test how urbanization affects color

morph prevalence and squirrel abundance via multiple direct and indirect pathways. We

generally followed the guidelines for graph-theoretic implementation of SEM with local

estimation (Grace et al., 2012), including development of a metamodel (Fig. 1), a causal graph

represented as a DAG (Fig. 2), and assessment of causal assumptions by testing the model for

missing links (Fig. 2), followed by re-specification and final model estimation (Fig. 3 & 4). Our

SEM consisted of two submodels. The first submodel (our "urbanization" submodel) included

direct and indirect pathways linking biotic and abiotic covariates that could affect squirrel

abundance (Fig. 2). The second submodel consisted of an integrated hierarchical model

combining camera detection data and visual point counts to estimate squirrel abundances by

color morph at each site (Cosentino et al., 2023; Kéry & Royle, 2020). This model included an

observation model to estimate detection probability of squirrels during surveys, and an

ecological process model to test how abundance of each color morph was related to each

environmental covariate. Integrating the urbanization and the hierarchical model allowed us to

quantify direct and indirect pathways by which urbanization drives color morph frequencies

while accounting for imperfect detection of individual squirrels at sampled sites.

2.5.1 Urbanization submodel

The urbanization submodel consisted of linear regressions with identity-link functions to

estimate the hypothesized causal relationships among human population density, building cover,

road distance, percent forest cover, forest fragmentation, and predator activity (Fig. 2). Although

many of these variables have been used as indices of urbanization either alone or as part of a

composite index, we developed a model to represent hypothesized relationships linking these

variables. The foundation of our urbanization submodel was human population density,

operating on the assumption that environmental change along urbanization gradients ultimately

is driven by people. We included direct effects of human population density on building cover

and road distance, assuming that buildings and roads are primarily infrastructure used by people.

Because roads are the main way to access buildings, we included a direct effect of building cover

on road distance. Because buildings and roads require tree removal, we included direct effects of

building cover and road distance on forest cover. Forest fragmentation has a known negative

relationship with forest cover (Fahrig, 2003, 2017), so we included a direct effect of forest cover

on fragmentation. We also included direct effects of building cover and road distance on forest

fragmentation, under the assumption that these landscape elements should cause discontinuities

in forest cover, irrespective of their effects on absolute forest cover. Each environmental variable

was included as a direct effect on predator activity as urbanization can have profound effects on

mammalian predator activity (Bateman et al., 2012; Ordeñana et al., 2010).

Urban systems are characterized by many covarying environmental gradients, which

creates high potential for confounding variables. Unobserved confounders can cause spurious

relationships between the environmental variables in our model and squirrel abundances. To

mitigate this concern, we included distance to city center as an additional covariate in the model,

with a direct effect on human population density. Including distance to city center allowed us to

estimate residual spatial variation in squirrel abundance and morph frequencies along the

urbanization gradient. This variable was intended to index axes of environmental variation along

the urbanization gradient that were not included in our model, controlling for possible

unobserved confounders (Cosentino et al., 2023; Dee et al., 2023).

2.5.2 Squirrel abundance submodel

Our integrated hierarchical model of squirrel abundance consisted of an ecological process

submodel to estimate the abundance of each color morph (N_{ik}, where $k = 1$ for the melanic morph

and $k = 2$ for the gray morph), and an observation process submodel to estimate individual

detection probabilities (p_{ijk}) based on repeated surveys j of color morph k at each location i. For

the observation process, we modeled the counts y_{ijk} of each morph from point count surveys as

$y_{ijk}|N_{ik} \sim Binomial(N_{ik}, p_{ijk})$ (Royle, 2004), and we modeled the binary detections y_{ijk} of each

morph from trail cameras as $y_{ijk}|N_{ik} \sim Bernoulli(P^*_{ijk})$, where $P^*_{ijk} = 1 - (1 - p_{ijk})^{N_{ik}}$ (Royle and

Nichols 2003). Because squirrel activity is influenced by ambient temperatures, we modeled

individual detection probability as a function of daily ambient temperature (Cosentino et al.,

2023; Wassmer & Refinetti, 2019):

$$\text{logit}(p_{ijk}) = a0_k + a1_k * \text{temperature}_{ij} + a2_k * \text{temperature}^2_{ij},$$

where for each color morph k, $a0_k$ represents the intercept, $a1_k$ represents a linear effect of daily

temperature, and $a2_k$ represents a quadratic effect of temperature. A quadratic effect was included

as squirrel activity peaks with the intermediate temperatures observed during fall and spring

(Cosentino et al., 2023).

For the ecological process submodel, we described the abundance of each color morph as

$N_{ik} \sim Poisson(\lambda_{ik})$, where λ_{ik} is the expected total abundance of squirrels of each color morph k at

location i. We modeled the expected abundance of each color morph as:

$$\log(\lambda_{ik}) = \beta 0_k + \beta 1_k * \text{distance}_i + \beta 2_k * \text{population}_i + \beta 3_k * \text{building}_i + \beta 4_k * \text{road}_i + \beta 5_k * \text{forest}_i$$

$$+ \beta 6_k * \text{frag}_i + \beta 7_k * \text{predator}_i,$$

where for each color morph k, $\beta 0$ represents the intercept and $\beta 1$- $\beta 7$ represent slopes for effects

of distance to city center, human population density, building cover, road distance, forest cover,

forest fragmentation, and predator activity.

2.5.3 Model estimation process

We used Bayesian estimation to fit the SEM with JAGS 4.3.0 (Plummer, 2017) and the package

jagsUI (Kellner & Meredith, 2024) in R (Version 4.2.3, R Core Team 2023). We generally used

noninformative prior distributions for intercept and slope parameters (Gelman & Hill, 2007). An

informative prior was specified for intercepts for individual detection probability and abundance

of each color morph because we had prior knowledge of mean detection probability and squirrel

abundance (Cosentino et al., 2023; Lemoine, 2019). Half-Cauchy priors were used for variance

parameters, as they are appropriate when setting parameters for variance in hierarchical models

(Gelman, 2006). We applied log transformations to population density, road distance, and

fragmentation, and we applied a logit transformation to building cover. All environmental

covariates were standardized to zero mean and unit standard deviation. For each step of the

estimation process (see below), we ran three chains with 20,000 adaptation iterations followed

by 4000 iterations, discarding the first 1000 iterations as burn-in and thinning the remaining

iterations by three. This resulted in 3000 iterations to describe posterior distributions.

Convergence of parameter estimates was confirmed with the Gelman-Rubin statistic (R-hat <

1.1; (Gelman & Hill, 2007).

Model fitting occurred in multiple steps, starting with fitting the initial urbanization

submodel. We applied d-separation tests to examine the urbanization submodel for missing links

(Shipley, 2009). These tests consisted of a linear regression between each pair of variables X_i and

X_j that were not directly connected in the initial model. Other variables $\{Z\}$ that were direct

effects on X_i or X_j were included as covariates in each regression. We tested each independence

claim $(X_i$ and $X_j)|\{Z\}$ by examining whether a 90% credible interval for the slope relating X_i and

X_j included 0. If the 90% CI excluded 0, we added a direct path between X_i and X_j to the

urbanization submodel. After all missing links were added based on the d-separation tests, we

refit the urbanization submodel (Fig. 3). We examined 90% CIs for each slope term in the refit

model, and we dropped terms when the 90% CI included 0. We used a 90% threshold to err on

the side of caution and including possibly important links in the final model. The final SEM

consisted of fitting the final urbanization submodel (Fig. 3) together with the squirrel abundance

submodel (Fig. 4, Fig. S2). We used 95% CIs to guide interpretation of the final SEM. Model

residuals were visually inspected at each step of the estimation process to confirm regression

assumptions were met.

For each response variable in the final SEM, we quantified explanatory power (R^2) for

Bayesian regression models as the variance of the predicted values divided by the sum of the

variance of the predicted values and variance of the residuals (Fig. S2; Gelman et al., 2019).

Residuals for each linear regression in the urbanization submodel were quantified as $y_i - \hat{y}_i$,

where y_i is the observed value and $\hat{y}_i$ is the predicted value at each site i. For squirrel

abundances, we defined residuals as $N_{ik} - \lambda_{ik}$ for each color morph k (Wright et al., 2019).

Because all urbanization covariates were standardized for analysis, regression

coefficients in the urbanization submodel represent standardized coefficients (Tab. S1), whereas

the raw regression coefficients in the squirrel abundance submodel were unstandardized (Tab.

S2). We used the observation-empirical approach to quantify standardized regression coefficients

in the squirrel abundance submodel (Lefcheck et al., 2018). We used the standardized

coefficients to quantify direct, indirect, and total effects of predictors on each response variable.

2.5.4 Residual spatial variation in proportion melanic and total abundance

Using the model estimates for each morph abundance, we derived total squirrel

abundance ($N\_total_i = N_{i1} + N_{i2}$) and the proportion of melanic squirrels (N_{i1}/N_total_i). We then

compared the direct and total effects of distance to city center on morph abundance, total

abundance, and proportion melanic. The direct effect of distance to city center represents

residual spatial variation along the urbanization gradient unexplained by explanatory variables in

our urbanization submodel, whereas the total effect represents residual spatial variance plus

variation explained by the urbanization submodel (i.e., indirect effects of distance to city center

mediated by covariates in the urbanization submodule). The difference between the direct and

total effect isolates variation explained by the urbanization submodel (Fig. 5).

2.5.5 Comparison of effects between morphs

To compare the direct effects of each explanatory variable on abundance between color

morphs, we derived the posterior distribution of the difference in unstandardized regression

coefficients between morphs for each predictor. We used unstandardized coefficients to compare

direct effects between morphs because standardized coefficients were conflated by differences in

variance in abundance for each morph. Thus, we used standardized coefficients to compare the

relative strength of pathways *within* each morph abundance model (Fig. 4a & 4b), whereas

unstandardized coefficients to compare effects *between* morphs (Fig. 4c).

3. Results

3.1 Urbanization submodel

Our urbanization submodel demonstrated how human population density affects abiotic and

biotic factors through multiple causal pathways (Fig. 3). Human population density directly

increased building cover, which subsequently decreased forest cover. When forest cover

decreased, forest fragmentation increased. Thus, population density indirectly decreased forest

cover and increased forest fragmentation through its effect on building cover. We also found

evidence for a direct positive effect of population density on forest fragmentation that was not

included in our initial model (Fig. 2; Tab. S1). Mammalian predator activity was positively

related to human population density, independent of effects of population density on building

cover, forest cover, and fragmentation. Population density was negatively related to distance to

city center as expected, and d-separation tests identified a residual negative relationship between

forest fragmentation and distance to city center. None of the variables in our model predicted

distance to high-speed roads.

3.2 Morph abundances

Our model revealed markedly different pathways by which urbanization affects color

morph abundances of gray squirrels (Fig. 4, Tab. S2). Human population density had a strong

direct effect on abundance of the gray morph, with abundance higher in more densely populated

areas. In contrast, there was no direct effect of human population density on melanic abundance.

Both gray and melanic abundances were indirectly related to human population density via its

cascading effects on building cover, forest cover, forest fragmentation, and predator activity.

Forest cover, fragmentation, and predator activity all had direct effects on each morph, but the

nature and magnitude of these effects varied between morphs. For example, forest cover had a

negative direct effect on gray abundance but a positive direct effect on melanic abundance.

Fragmentation had a negative direct effect on both morph abundances, but the effect was

stronger for the melanic than gray morph (Fig. 4c). Melanic abundance was negatively related to

predator activity, whereas gray abundance was *positively* related to predator activity. Road

distance had a positive direct effect on melanic abundance but no effect on gray abundance.

Building cover had no direct effect on the abundance of either morph.

In addition to direct effects, our model revealed many indirect and cascading paths by

which urbanization impacted each morph abundance (Tab. S2). The direction of most indirect

effects of urban covariates on abundance was consistent with direct effects, but in some cases the

direct and indirect effects were opposite. For example, forest cover and building cover had

different direct and total effects. Forest cover directly drove lower gray morph abundance but

had a weak indirect positive effect through its reducing fragmentation, resulting in no total effect

of forest cover on gray morph abundance (Tab. S2). Building cover had no direct effect on either

morph, but indirectly drove lower melanic abundance, resulting in a negative total effect of

building cover on melanic abundance (Tab. S2).

Importantly, there was no direct effect of distance to city center on gray morph

abundance, but there was a negative total effect (Fig. 5, Tab. S2). This indicates that the variation

in gray abundance along the urbanization gradient was adequately explained by variables in our

urbanization submodel. In contrast, we found both direct and total negative effects of distance to

city center on melanic abundance (Fig. 4 & 5, Tab. S2). The strong direct effect of distance to

city center on melanic abundance indicates there was substantial residual variation in melanic

abundance along the urbanization gradient that remained unexplained by the covariates in the

urbanization submodel. Derived quantities (proportion melanic and total squirrel abundance)

showed strong negative clines with distance to city center (Fig. 5b & 5c), outcomes that align

with previous research showing this pattern across numerous cities (Cosentino et al., 2023;

Cosentino & Gibbs, 2022). However, our models decompose this finding between morphs and

reveal the cline in total abundance was driven largely by the decline of melanic abundance as

distance to city center increased, a finding in line with previous work showing melanic survival

declines further from the city center (Cosentino et al., 2023, Fig. 5).

Discussion

377   Our findings reveal a complex network of effects by which the urban environment alters

abundance and trait variation in gray squirrels. We show markedly different mechanistic

pathways driving abundance of two distinct phenotypes, and these distinct processes together

determine urban-rural clines in species abundance and trait variation. Human population density

impacted the two morphs in different ways, having a direct positive effect on gray morph

abundance, a pattern likely facilitated by novel resources associated with people (i.e., bird

feeders), but only indirectly impacted melanic morph abundance through its effect on landscape

structure (i.e., through forest cover and fragmentation). For some covariates (e.g., forest

fragmentation) the direction of the effect on color morph abundances was the same, but the

magnitude of effect differed. But, for other covariates (e.g., predator activity and forest cover)

the effects were entirely opposite (Fig. 3). Thus, our study reveals phenotypic specific

mechanisms and complex processes underlying known urban-rural clines in biodiversity patterns

(Fig. 2 & 3). Our findings also demonstrate urbanization is comprised of a network of processes

that should be considered in concert to understand and mitigate their impacts.

Developing a scientific model with hypothesized causal relationships between drivers—

represented in our DAG—is critical for informing appropriate statistical modeling in ecological

systems (Arif & MacNeil, 2023). Statistical models built from DAGs enable researchers to

differentiate between direct and indirect effects, control for confounders, and identify spurious or

masked effects to better understand the multiple causal processes at play (Grace et al., 2012;

McElreath, 2020). For example, our models revealed forest cover had a direct negative effect and

an indirect positive effect through fragmentation on gray morph abundance, leading to no total

effect. Without modeling the relationships between forest cover and fragmentation in our system,

we would have erroneously concluded that forest cover has no effect on gray morph abundance,

when, in fact greater forest cover on its own likely caused a decline in gray morph abundance,

but this effect was countered by the positive indirect effect via reduced fragmentation. Thus, our

work highlights how DAGs can be used in studies in urban ecology and evolution to disentangle

interacting mechanisms at play in each system and break down their effects into their various

components (Arif & MacNeil, 2023; McElreath, 2020).

Our results also provide important insights into how urban effects can be mediated by

phenotypic trait variation, with some trait variants being more sensitive to anthropogenic impacts

than others. In our system, melanic abundance was more sensitive to landscape change via

deforestation, fragmentation, and distance to high speed roads than gray abundance (Fig. 4).

While many studies have demonstrated phenotypic changes across the urban-rural gradient

(Campbell-Staton et al., 2020; Santangelo et al., 2018, 2020; Winchell et al., 2016), our study

provides a clear example of anthropogenic impacts operating differently on two phenotypes.

These differences reveal a suite of mechanisms and processes underlying observed trait variation

across the landscape. For example, our results show divergent sensitivity to predation between

phenotypes. Indeed, greater predator activity drove lower melanic abundance, but not gray

abundance. Thus, modeling the abundances of each phenotype separately revealed important

differences in mechanisms at play affecting both phenotypes and shows both habitat change and

predator activity play key roles in determining the observed urban-rural clines, albeit by

operating differently on each phenotype.

419   The difference in effects of predator activity on each morph's abundance—with predator

activity negatively affecting melanic morphs, but positively affecting gray morphs—is consistent

with the hypothesis that melanic squirrels are more visible to predators and thus exposed to

greater predation risk (Bryan, 2020; Gibbs et al., 2019). The melanic morph is generally less

cryptic than the gray morph in urban and rural habitats (Bryan, 2020), so crypsis is likely a key

factor driving patterns in melanism across urban-rural gradients. However, in addition to

differences in crypsis, it is possible the two morphs differ in behavior or activity patterns in ways

that alter their predation risk (Engel et al., 2020; Sarno et al., 2015). For example, the melanic

and gray morph may exhibit different diel activity patterns, or activity patterns across vertical

space (from ground and canopy) resulting in differing susceptibility to predation or thermal

conditions (Bryan, 2020; Ciurej et al., 2020; Ducharme et al., 1989). One hypothesis for such

differences in spatial or temporal activity patterns is that each morph's thermal constraints differ

(Ciurej et al., 2020). Indeed, evidence suggests melanism can provide a thermal benefit in

squirrels, particularly in colder winter months, allowing them to be more active during cold times

(Ciurej et al., 2020; Ducharme et al., 1989). As thermal microclimates vary both throughout the

day and across vertical strata of arboreal habitats (Gámez & Harris, 2022; Nakamura et al.,

2017), differences in thermal constraints between morphs could result in differential activity

patterns. Thus, future research should investigate how thermal conditions and squirrel activity

patterns may contribute to patterns in melanism across the landscape.

While our models showed forest loss and fragmentation did not directly alter predator

activity patterns (Fig. 3), it is likely the effects of such habitat alteration and predator activity are

still connected. Many studies have shown fragmentation can increase exposure to predation risk

(Fischer & Lindenmayer, 2007; Robinson et al., 1995; Schneider et al., 2012). Thus, given

melanic squirrels were found to be more susceptible to predation pressure (Bryan, 2020), it may

explain one reason they are also more susceptible to landscape change, as it could expose them

to more predation and reduce their ability to camouflage. Indeed melanic squirrels are more

easily detected visually in urban landscapes (Gibbs et al., 2019) and thus, landscape change and

predator activity could synergistically drive lower melanic abundance. Our models do not

explicitly test for synergistic effects and future studies should explicitly test for synergies arising

in such systems.

The sensitivity of melanic abundance to multiple common axes of environmental change

in cities, such as habitat loss and fragmentation, raises an important question about what

maintains urban-rural clines in melanism. Indeed, the environmental variables in our

urbanization submodel showed that melanic abundance had an indirect positive relationship with

distance to city center, which should result in greater melanic abundance further from the city

center (Tab. S2). However, these indirect effects were counteracted by a strong, negative direct

effect of distance to city center on melanic abundance, ultimately resulting in a total negative

effect, with melanic abundance declining further from the city center (Tab. S2). The direct

negative effect of distance to city center on melanic abundance indicates our urbanization

submodel is missing some variables or processes that could cause melanic abundance to be high

in cities (Dee et al., 2023).

The intriguing and seemingly contradictory pattern, in which melanic abundance is more

sensitive to urban environmental changes but found in greater abundance closer to the city

center, may be due to the interaction of historic and contemporary forces, including ecological,

evolutionary, and social forces (Des Roches et al., 2021). For example, melanic squirrels, once

common across the landscape, were largely extirpated with the rise of European colonial

settlement, the accompanied deforestation and logging in surrounding old growth forests, and

extensive hunting (Benson, 2013; Cronon, 1991; Robertson, 1973). Evidence suggests urban

areas functioned as refugia for gray squirrels in general, including the melanic morph (Benson,

2013; Gibbs et al., 2019). Additionally in the 1930s some naturalists promoted feeding gray

squirrels, introducing them to urban areas, and celebrated them as "our most loved" species in

cities, establishing a social attitude of tolerance and even celebration (Benson, 2013). Thus,

urban-rural clines in abundance in gray squirrels likely arose from both ecological and social

processes, which together can drive patterns in present day urban biodiversity (Cosentino et al.,

2023; Cosentino & Gibbs, 2022; Des Roches et al., 2021). But, with nearly two centuries of

forest recovery (Thompson et al., 2013) and a decline in hunting across the United States (Price

Tack et al., 2018), it is striking that a cline with distance to city center persists for melanic

abundance, but not the gray morph (Fig. 4a & 4b). Thus, urban-rural clines in melanism are

likely maintained by novel contemporary forces, perhaps by a combination of predation pressure

(including human hunting), social forces (such as humans providing food via bird feeders), and

dispersal constraints, more strongly impacting melanics than gray morphs, despite much

surrounding forest recovery (Cosentino et al., 2023). These findings point to the importance of

considering historical social and ecological context when seeking to understand present day

patterns associated with urbanization and highlights historic and contemporary forces often

operate together determine current biodiversity patterns (Des Roches et al., 2021).

One surprising finding from our work is that predator activity had a positive effect on

gray morph abundance but a negative effect on melanic abundance. There are at least two

possible explanations for the positive effect of predators in gray morphs. First, there may be

unobserved confounders driving this association, for example a particular unmeasured habitat

characteristic or resource that benefits both the gray morph and predators. Alternatively, it is

possible this positive association is evidence of unmodeled indirect paths by which predator

activity drives greater gray morph abundance (Abrams, 2009; Holt & Bonsall, 2017). For

example, predator activity could reduce competition pressure if it disproportionately negatively

impacted competing species, thus indirectly allowing greater gray morph abundance (Abrams &

Matsuda, 1996). However, it should be noted that our methods only allowed us to account for

mammalian predators and future research should incorporate raptor activity, as they are also

important squirrel predators. Incorporating raptor predation risk may further elucidate complex

predator-prey dynamics in urban systems.

Another mechanism hypothesized to contribute to urban-rural clines in melanism of

eastern gray squirrels is road mortality (Gibbs et al., 2019). Prior research shows melanic

squirrels are underrepresented among road kill, relative to the gray morph—a finding attributed

to humans more easily detecting and avoiding melanic squirrels against gray asphalt when

driving (Gibbs et al., 2019). Our findings provide important insights into this hypothesis showing

melanic abundance is lower closer to roads (Fig. 4b). This finding suggests underrepresentation

of melanic squirrels among roadkill may be due in part to behavioral avoidance of roads by

melanic squirrels, rather than simply a reduced risk of mortality through human detections while

driving. However, these mechanisms are not mutually exclusive, as reduced road mortality risk

for melanic morphs may be driven by both behavioral avoidance of roads and vehicle drivers

being more likely to see and avoid hitting melanic individuals.

An important consideration for future research is camera trap placement, as our camera

traps were placed near ground level, limiting our ability to account for arboreal activity (Moore

et al., 2020). However, considering a species' vertical niche is critical when studying arboreal

species, as vertical niche lability has been linked to mitigating changes in climate, avoiding

predation, reducing competition between species, and can even change with urbanization

(Basham & Scheffers, 2020; Borden et al., 2021; Gámez & Harris, 2022; Rankin et al., 2018).

How urbanization affects the vertical stratification and movement of species through arboreal

space is poorly understood (Borden et al., 2021; Lakatos et al., 2022). Incorporating point count

surveys in which an observer visually counted squirrels across the available vertical habitat

likely mitigated the ground-level-bias in our study by accounting for squirrels in trees.

Nevertheless, some of the observed patterns in morph abundances could be influenced by shifts

in vertical niche, particularly if vertical niche use differs between morphs. For example, if one

morph shifts to a more arboreal niche in the presence of certain predators or under certain habitat

conditions (Borden et al., 2021; Neilly et al., 2017), it would alter the detection probability

between morphs in ways we cannot presently account for. Thus, future camera trap studies in

cities targeting arboreal species should consider including canopy camera traps to distinguish

between changes in abundance or activity, and the possibility that species are instead shifting

vertically across their vertical niche along the urban-rural gradient (Borden et al., 2021;

Culbertson & Herrmann, 2019).

Overall, our research demonstrates how urban effects on abundance and phenotypic trait

variation are complex, involving a system of direct and indirect mechanisms. We illustrate how

scientific models represented with DAGs and statistical analysis with structural equation models

can be used to disentangle the effects of environmental variables commonly used to broadly

represent urbanization. By applying a structural causal model framework, this study allowed us

to characterize the structure of relationships between drivers in our system and mechanistically

understand how this system drives ecological and evolutionary patterns in a ubiquitous mammal

(Arif & MacNeil, 2023). There is great opportunity for research in urban ecology and evolution

to incorporate structural causal modeling frameworks to disentangle the mechanisms at play in

urban systems and understand the causal drivers of intra and inter-city variation. By elucidating

these complex systems, we will gain valuable insight into the role distinct urban drivers play in

determining ecological and evolutionary patterns both within and between cities.

Acknowledgements

Field assistance was provided by Joelee Tooley and Richard Rich. Access to field sites was

provided by the New York Department of Environmental Conservation, City of Syracuse, and

select private and municipal landowners.

Funding: This research was funded by the U.S. National Science Foundation (DEB-2018140,

DEB-2018249).

Data Availability: Data and code are available in the following repository:

https://github.com/bcosentino/urbanCausalInference

Conflict of Interest Statement: We, the authors, declare no conflict of interest in conducting

and writing this work.

Figures
555

556
Figure 1. The conceptual metamodel of the complex network of effects from urbanization and
the many potential direct and indirect paths of impact to ecological and evolutionary processes.
a) Urbanization consists of a suite of interacting anthropogenic impacts, which can represent
distinct mechanisms of impact to biotic systems. This suite of changes directly alters b) human
wildlife interactions, c) habitat quality and configuration, d) ecological processes, and e)
evolutionary processes. However, there are also many indirect pathways by which urbanization
can impact ecological and evolutionary processes.
564

565
566
Figure 2. Diagram of our full scientific model, showing the hypothesized causal relationships
between drivers and morph abundances. Gray arrows show the initial paths we hypothesized, and
orange arrows indicate paths added after D-separation test revealed missing important causal
paths.

Table 1. List of initial hypothesized causal paths by which drivers interact and impact squirrel
ecology in our system. Arrows in the first column indicate the hypothesized direction of effect
and bolded rows show hypothesized direct causal effects on abundance and morph prevalence in
eastern gray squirrels.
575

Effect	Expected direction	Rationale	References
Human population density → building density	(+)	Greater human population drives land conversion, including expansion of urban areas and greater building cover	(Ehrlich et al., 2018; Güneralp et al., 2020; Xu et al., 2018)
Human population density → road distance	(-)	Greater human population is linked to greater infrastructure including higher road density, reducing the distance to roads	(Xu et al., 2018)
Human population density → mammalian predator activity	(-)	Greater human activity alters mammalian predator activity through altered human-wildlife interactions, niche shifts, habitat and resource alteration	(Cervantes et al., 2023; Gilbert et al., 2022; Tucker et al., 2021)
Building cover → forest cover	(-)	Greater building density, by definition, reduces forest cover, and is a major driver of deforestation in urban settings.	(Güneralp et al., 2020)
Building cover → road distance	(-)	Greater building cover requires greater infrastructure resulting in smaller distance to roads as building cover increases	(Bi et al., 2010)
Building cover → mammalian predator activity	(+/-)	Urbanization and building cover can alter landscape connectivity for mammalian predators, reduce habitat availability, and alter prey availability	(Bateman et al., 2012; Cervantes et al., 2023; Li et al., 2022; Tucker et al., 2021)
Building cover → forest fragmentation	(+)	Building cover replaces natural habitats and therefore drives forest loss and fragmentation	(Güneralp et al., 2020)
Forest cover → fragmentation	(-)	Forest loss and fragmentation, while different, are closely correlated and greater forest cover reduces fragmentation	(Fahrig, 2017; Fletcher et al., 2018)
Forest cover → mammalian predator activity	(+/-)	Forest cover alters mammalian predator activity and can be either negative or positive depending on the species and context.	(Dijak & Thompson, 2000; Regolin et al., 2017)

Human population density → squirrel/morph abundances	(+/-)	Human population density can drive greater squirrel abundance through provisions of novel resources (bird feeders, buildings for nesting), and through indirect paths altering predator activity and habitat availability. Effects may differ by morph.	(Benson, 2013; Mccleery et al., 2007; Reed & Bonter, 2018)
Building cover → squirrel/morph abundances	(+/-)	Alters habitat availability by reducing tree cover but also providing novel nesting sites in buildings	(Benson, 2013; Parker & Nilon, 2008)
Road distance → squirrel/morph abundances	(+/-)	Roadkill is a major driver of squirrel mortality in urban contexts and road mortality rates differ between morphs	(Gibbs et al., 2019)
Forest cover → squirrel/morph abundances	(+/-)	Reduces fragmentation and increases connectivity. Increases available natural resources (including food and nesting resources), increases predator activity	(Koprowski et al., 2016; Merrick et al., 2016; Parker & Nilon, 2008)
Forest fragmentation → squirrel/morph abundances	(+/-)	Reduces habitat connectivity and available natural resources, and constrains movement. May affect the morphs differently.	(Koprowski et al., 2016; Parker & Nilon, 2008)
Mammalian predator activity → squirrel/morph abundances	(+/-)	Mammalian predator activity alters predation risk as well as the landscape of fear, likely altering squirrel abundance patterns. Differences in crypsis or behavior may cause morphs to be affected differently by predator activity.	(Cosentino & Gibbs, 2022; Gibbs et al., 2019)

576
577

Figure 3. Results from the urbanization covariate sub model. This model reveals the complex and interactive structure and causal relationships between the urbanization variables in our system. Arrows represent effects for which the confidence intervals did not overlap zero and no arrows are included for effects for which the confidence interval did overlap zero. Arrow thickness represents the strength of the effect, and colors and symbols shows the direction of the effect (blue = positive, red = negative). Distance to city center represents a non-mechanistic spatial variable which accounts for unmeasured variance in our system, likely due to unmeasured covariates.

591
Figure 4. Models revealed markedly different direct and indirect pathways driving abundance
patterns in (a) gray morph, and (b) melanic morph, eastern gray squirrels. Specifically,
abundance of (a) the gray morph was positively affected by human population density and
predator activity, and negatively affected by forest cover and forest fragmentation. Abundance of
(b) the melanic morph was positively affected by road distance and greater forest cover, but
negatively affected by distance to city center, predator activity, and forest fragmentation.
Differences in model coefficients between morphs (c) revealed effects of distance to city center,
tree cover, fragmentation and predator activity all varied significantly between morphs. For all
panels, red and blue represent negative and positive values, respectively. Gray points (c)
represent values for which the credible interval overlaps zero.

Figure 5. Predicted (a) melanic and (b) gray morph abundances, (c) proportion melanic, and (d)
total squirrel abundance in relationship to distance to city center for eastern gray squirrels
(*Sciurus carolinensis*) in Syracuse, NY, USA. Differences between direct (orange shading) and
total (blue shading) effects in each panel provides a visual representation of the degree to which
the environmental covariates in our study accounted for observed variation in abundance and
color morph proportions.

References

615   **References**
616
Abrams, P. A. (2009). When does greater mortality increase population size? The long

history and diverse mechanisms underlying the hydra effect. *Ecology Letters*,

12(5), 462–474. https://doi.org/10.1111/j.1461-0248.2009.01282.x

Abrams, P. A., & Matsuda, H. (1996). Positive Indirect Effects Between Prey Species

that Share Predators. *Ecology*, *77*(2), 610–616. https://doi.org/10.2307/2265634

Alberti, M., Palkovacs, E. P., Roches, S. D., De Meester, L., Brans, K. I., Govaert, L.,

Grimm, N. B., Harris, N. C., Hendry, A. P., Schell, C. J., Szulkin, M., Munshi-

South, J., Urban, M. C., & Verrelli, B. C. (2020). The Complexity of Urban Eco-

evolutionary Dynamics. *BioScience*, *70*(9), 772–793.

https://doi.org/10.1093/BIOSCI/BIAA079

Arif, S., & MacNeil, M. A. (2023). Applying the structural causal model framework for

observational causal inference in ecology. *Ecological Monographs*, *93*(1), e1554.

https://doi.org/10.1002/ecm.1554

Arif, S., & Massey, M. D. B. (2023). Reducing bias in experimental ecology through

directed acyclic graphs. *Ecology and Evolution*, *13*(3), e9947.

https://doi.org/10.1002/ece3.9947

Aronson, M. F. J., La Sorte, F. A., Nilon, C. H., Katti, M., Goddard, M. A., Lepczyk, C.

A., Warren, P. S., Williams, N. S. G., Cilliers, S., Clarkson, B., Dobbs, C., Dolan,

R., Hedblom, M., Klotz, S., Kooijmans, J. L., Kuhn, I., MacGregor-Fors, I.,

McDonnell, M., Mortberg, U., … Winter, M. (2014). A global analysis of the

impacts of urbanization on bird and plant diversity reveals key anthropogenic

638  drivers. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1780),

639  20133330–20133330. https://doi.org/10.1098/rspb.2013.3330

640 Basham, E. W., & Scheffers, B. R. (2020). Vertical stratification collapses under

641  seasonal shifts in climate. *Journal of Biogeography*, *September 2019*, 1–11.

642  https://doi.org/10.1111/jbi.13857

643 Baston, D., ISciences, & LLC. (2022). *exactextractr: Fast Extraction from Raster*

644  *Datasets using Polygons* (0.9.1) [Computer software]. https://cran.r-

645  project.org/web/packages/exactextractr/index.html

646 Bateman, P. W., Fleming, P. A., Fleming, T., & Comber, S. L. (2012). Big city life:

647  Carnivores in urban environments. *Journal of Zoology*, *287*(1), 1–23.

648  https://doi.org/10.1111/J.1469-7998.2011.00887.X

649 Bennett, V. J. (2017). Effects of Road Density and Pattern on the Conservation of

650  Species and Biodiversity. *Current Landscape Ecology Reports*, *2*(1), 1–11.

651  https://doi.org/10.1007/s40823-017-0020-6

652 Benson, E. (2013). The Urbanization of the Eastern Gray Squirrel in the United States.

653  *Journal of American History*, *100*(3), 691–710.

654  https://doi.org/10.1093/jahist/jat353

655 Bi, X., Wang, H., & Ge, J. (2010). The changes of densities and patterns of roads and

656  rural buildings: A case study on Dongzhi Yuan of the Loess Plateau, China.

657  *Environmental Monitoring and Assessment*, *164*(1), 549–560.

658  https://doi.org/10.1007/s10661-009-0912-y

Borden, J. B., Bohlman, S., & Scheffers, B. R. (2021). Niche lability mitigates the impact

of invasion but not urbanization. *Oecologia*, *0123456789*.

https://doi.org/10.1007/s00442-021-05039-x

Bowman, J., Ray, J. C., Magoun, A. J., Johnson, D. S., & Dawson, F. N. (2010). Roads,

logging, and the large-mammal community of an eastern Canadian boreal forest.

Canadian Journal of Zoology, *88*(5), 454–467. https://doi.org/10.1139/z10-019

Bryan, A. (2020). *Effects of urbanization on the evolution of coat color in eastern gray*

squirrels (Sciurus carolinensis). Hobart and William Smith Colleges.

Bu, H., Wang, F., McShea, W. J., Lu, Z., Wang, D., & Li, S. (2016). Spatial Co-

Occurrence and Activity Patterns of Mesocarnivores in the Temperate Forests of

Southwest China. *PLOS ONE*, *11*(10), e0164271.

https://doi.org/10.1371/journal.pone.0164271

Campbell-Staton, S. C., Winchell, K. M., Rochette, N. C., Fredette, J., Maayan, I.,

Schweizer, R. M., & Catchen, J. (2020). Parallel selection on thermal physiology

facilitates repeated adaptation of city lizards to urban heat islands. *Nature*

Ecology & Evolution, 1–7. https://doi.org/10.1038/s41559-020-1131-8

Center For International Earth Science Information Network-CIESIN-Columbia

University. (2017). *Gridded Population of the World, Version 4 (GPWv4):*

Population Density, Revision 11 [Dataset]. Palisades, NY: Socioeconomic Data

and Applications Center (SEDAC). https://doi.org/10.7927/H49C6VHW

Cerqueira, R. C., de Rivera, O. R., Jaeger, J. A. G., & Grilo, C. (2021). Direct and

indirect effects of roads on space use by jaguars in Brazil. *Scientific Reports*,

11(1), Article 1. https://doi.org/10.1038/s41598-021-01936-6

Cervantes, A. M., Schooley, R. L., Lehrer, E. W., Gallo, T., Allen, M. L., Fidino, M., &

Magle, S. B. (2023). Carnivore coexistence in Chicago: Niche partitioning of

coyotes and red foxes. *Urban Ecosystems*. https://doi.org/10.1007/s11252-023-

01368-2

Ciurej, A. K., Oblander, A., Swift, A. W., & Wilson, J. A. (2020). Melanism as a potential

thermal benefit in eastern fox squirrels (Sciurus niger). *European Journal of*

Ecology, *5*(2), 79–87. https://doi.org/10.2478/eje-2019-0013

Cosentino, B. J., & Gibbs, J. P. (2022). Parallel evolution of urban–rural clines in

melanism in a widespread mammal. *Scientific Reports*, *12*(1), 1–7.

https://doi.org/10.1038/s41598-022-05746-2

Cosentino, B. J., Vanek, J. P., & Gibbs, J. P. (2023). Rural selection drives the evolution

of an urban–rural cline in coat color in gray squirrels. *Ecology and Evolution*,

13(10), e10544. https://doi.org/10.1002/ece3.10544

Cronon, W. (1991). Nature's metropolis: Chicago and the Great West WW Norton &

Company. *New York*.

Culbertson, K. A., & Herrmann, N. C. (2019). Asymmetric interference competition and

niche partitioning between native and invasive Anolis lizards. *Oecologia*, *190*(4),

811–820. https://doi.org/10.1007/s00442-019-04466-1

Dee, L. E., Ferraro, P. J., Severen, C. N., Kimmel, K. A., Borer, E. T., Byrnes, J. E. K.,

Clark, A. T., Hautier, Y., Hector, A., Raynaud, X., Reich, P. B., Wright, A. J.,

Arnillas, C. A., Davies, K. F., MacDougall, A., Mori, A. S., Smith, M. D., Adler, P.

B., Bakker, J. D., … Loreau, M. (2023). Clarifying the effect of biodiversity on

productivity in natural ecosystems with longitudinal data and methods for causal

inference. *Nature Communications*, *14*(1), Article 1.

https://doi.org/10.1038/s41467-023-37194-5

Des Roches, S., Brans, K. I., Lambert, M. R., Rivkin, L. R., Savage, A. M., Schell, C. J.,

Correa, C., De Meester, L., Diamond, S. E., Grimm, N. B., Harris, N. C., Govaert,

L., Hendry, A. P., Johnson, M. T. J., Munshi-South, J., Palkovacs, E. P., Szulkin,

M., Urban, M. C., Verrelli, B. C., & Alberti, M. (2021). Socio-eco-evolutionary

dynamics in cities. *Evolutionary Applications*, *14*(1), 248–267.

https://doi.org/10.1111/eva.13065

Dijak, W. D., & Thompson, F. R. (2000). Landscape and Edge Effects on the

Distribution of Mammalian Predators in Missouri. *The Journal of Wildlife*

Management, *64*(1), 209–216. https://doi.org/10.2307/3802992

Ducharme, M. B., Larochelle, J., & Richard, D. (1989). Thermogenic Capacity in Gray

and Black Morphs of the Gray Squirrel, Sciurus carolinensis. *Physiological*

Zoology, *62*(6), 1273–1292. https://doi.org/10.1086/physzool.62.6.30156213

Ehrlich, D., Kemper, T., Pesaresi, M., & Corbane, C. (2018). Built-up area and

population density: Two Essential Societal Variables to address climate hazard

impact. *Environmental Science & Policy*, *90*, 73–82.

https://doi.org/10.1016/j.envsci.2018.10.001

Engel, L. D., Carlen, E. J., Losos, J. B., & Winchell, K. M. (2020). *Eastern Gray*

Squirrels (Sciurus carolinensis) Differ in Abundance and Response to Humans

Across Urban Habitats of St. Louis. *33*.

Fahrig, L. (2003). Effects of Habitat Fragmentation on Biodiversity. *Annual Review of*

Ecology, Evolution, and Systematics, *34*(1), 487–515.

https://doi.org/10.1146/annurev.ecolsys.34.011802.132419

Fahrig, L. (2017). Ecological Responses to Habitat Fragmentation Per Se. *Annu. Rev.*

Ecol. Evol. Syst. 2017, *48*, 1–23. https://doi.org/10.1146/annurev-ecolsys-110316

Fischer, J., & Lindenmayer, D. B. (2007). Landscape modification and habitat

fragmentation: A synthesis. *Global Ecology and Biogeography*, *16*(3), 265–280.

https://doi.org/10.1111/j.1466-8238.2007.00287.x

Fletcher, R. J., Didham, R. K., Banks-Leite, C., Barlow, J., Ewers, R. M., Rosindell, J.,

Holt, R. D., Gonzalez, A., Pardini, R., Damschen, E. I., Melo, F. P. L., Ries, L.,

Prevedello, J. A., Tscharntke, T., Laurance, W. F., Lovejoy, T., & Haddad, N. M.

(2018). Is habitat fragmentation good for biodiversity? *Biological Conservation*,

226(April), 9–15. https://doi.org/10.1016/j.biocon.2018.07.022

Francis, C. D., Kleist, N. J., Ortega, C. P., & Cruz, A. (2012). Noise pollution alters

ecological services: Enhanced pollination and disrupted seed dispersal.

Proceedings of the Royal Society B: Biological Sciences, *279*(1739), 2727–2735.

https://doi.org/10.1098/rspb.2012.0230

Gámez, S., & Harris, N. C. (2022). Conceptualizing the 3D niche and vertical space use.

Trends in Ecology & Evolution, 1–10. https://doi.org/10.1016/j.tree.2022.06.012

Gaynor, K. M., Brown, J. S., Middleton, A. D., Power, M. E., & Brashares, J. S. (2019).

Landscapes of Fear: Spatial Patterns of Risk Perception and Response. *Trends*

in Ecology & Evolution, *34*(4), 355–368.

https://doi.org/10.1016/J.TREE.2019.01.004

Gelman, A. (2006). Prior distributions for variance parameters in hierarchical models

(comment on article by Browne and Draper). *Bayesian Analysis*, *1*(3), 515–534.

https://doi.org/10.1214/06-BA117A

Gelman, A., Goodrich, B., Gabry, J., & Vehtari, A. (2019). R-squared for Bayesian

Regression Models. *The American Statistician*, *73*(3), 307–309.

https://doi.org/10.1080/00031305.2018.1549100

Gelman, A., & Hill, J. (2007). *Data Analysis Using Regression and*

Multilevel/Hierarchical Models. Cambridge University Press.

Gibbs, J. P., Buff, M. F., & Cosentino, B. J. (2019). The Biological System—Urban

Wildlife, Adaptation, and Evolution: Urbanization as a Driver of Contemporary

Evolution in Gray Squirrels (Sciurus carolinensis). In M. H. P. Hall & S. B. Balogh

(Eds.), *Understanding Urban Ecology: An Interdisciplinary Systems Approach*

(pp. 269–286). Springer International Publishing. https://doi.org/10.1007/978-3-

030-11259-2_12

Gilbert, N. A., Stenglein, J. L., Pauli, J. N., & Zuckerberg, B. (2022). Human disturbance

compresses the spatiotemporal niche. *Proceedings of the National Academy of*

Sciences of the United States of America, *119*(52), e2206339119.

https://doi.org/10.1073/PNAS.2206339119/SUPPL_FILE/PNAS.2206339119.SA

PP.PDF

Grace, J. B. (2006). *Structural Equation Modeling and Natural Systems*. Cambridge

University Press.

Grace, J. B., Schoolmaster Jr., D. R., Guntenspergen, G. R., Little, A. M., Mitchell, B.

R., Miller, K. M., & Schweiger, E. W. (2012). Guidelines for a graph-theoretic

772   implementation of structural equation modeling. *Ecosphere*, *3*(8), art73.

773   https://doi.org/10.1890/ES12-00048.1

774 Güneralp, B., Reba, M., Hales, B. U., Wentz, E. A., & Seto, K. C. (2020). Trends in

775   urban land expansion, density, and land transitions from 1970 to 2010: A global

776   synthesis. *Environmental Research Letters*, *15*(4), 044015.

777   https://doi.org/10.1088/1748-9326/ab6669

778 Hesselbarth, M. H. K., Sciaini, M., With, K. A., Wiegand, K., & Nowosad, J. (2019).

779   landscapemetrics: An open-source R tool to calculate landscape metrics.

780   *Ecography*, *42*(10), 1648–1657. https://doi.org/10.1111/ecog.04617

781 Hijmans, R. J., Etten, J. van, Sumner, M., Cheng, J., Baston, D., Bevan, A., Bivand, R.,

782   Busetto, L., Canty, M., Fasoli, B., Forrest, D., Ghosh, A., Golicher, D., Gray, J.,

783   Greenberg, J. A., Hiemstra, P., Hingee, K., Ilich, A., Geosciences, I. for M. A., …

784   Wueest, R. (2023). *raster: Geographic Data Analysis and Modeling* (3.6-20)

785   [Computer software]. https://cran.r-project.org/web/packages/raster/index.html

786 Holt, R. D., & Bonsall, M. B. (2017). *Annual Review of Ecology, Evolution, and*

787   *Systematics Apparent Competition*. https://doi.org/10.1146/annurev-ecolsys-

788   110316

789 Kellner, K., & Meredith, M. (2024). *jagsUI: A Wrapper Around "rjags" to Streamline*

790   *"JAGS" Analyses* (1.6.2) [Computer software]. https://cran.r-

791   project.org/web/packages/jagsUI/index.html

792 Kéry, M., & Royle, J. A. (2020). *Applied hierarchical modeling in ecology: Analysis of*

793   *distribution, abundance and species richness in R and BUGS: Volume 2:*

794   *Dynamic and advanced models*. Academic Press.

https://books.google.com/books?hl=en&lr=&id=fyk0DwAAQBAJ&oi=fnd&pg=PP1

&dq=kery+and+royle&ots=p3B1YI2aSu&sig=tS4IT7w1GbrmKDt_V9wUMsXVB_

A

Kok, A. C. M., Berkhout, B. W., Carlson, N. V., Evans, N. P., Khan, N., Potvin, D. A.,

Radford, A. N., Sebire, M., Shafiei Sabet, S., Shannon, G., & Wascher, C. A. F.

(2023). How chronic anthropogenic noise can affect wildlife communities.

Frontiers in Ecology and Evolution, *11*.

https://doi.org/10.3389/fevo.2023.1130075

Koppen, W. (1936). The Geographical System of Climate. *Koppen W, Geiger Reds.*

Handbook of Climatology. Gebruder Borntraeger, Berlin, 1–50.

Koprowski, J., Munroe, K., & Edelman, A. (2016). *Gray not grey: The ecology of Sciurus*

carolinensis in their native range in North America (pp. 1–18).

Lakatos, T., Chamberlain, D. E., Garamszegi, L. Z., & Batáry, P. (2022). No place for

ground-dwellers in cities: A meta-analysis on bird functional traits. *Global*

Ecology and Conservation, *38*, e02217.

https://doi.org/10.1016/j.gecco.2022.e02217

Lefcheck, J., Byrnes, J., & Grace, J. (2018). *piecewiseSEM: Piecewise structural*

equation modeling. An r package version 2.1. 0.

Lemoine, N. P. (2019). Moving beyond noninformative priors: Why and how to choose

weakly informative priors in Bayesian analyses. *Oikos*, *128*(7), 912–928.

https://doi.org/10.1111/oik.05985

Li, G., Fang, C., Li, Y., Wang, Z., Sun, S., He, S., Qi, W., Bao, C., Ma, H., Fan, Y.,

Feng, Y., & Liu, X. (2022). Global impacts of future urban expansion on terrestrial

818   vertebrate diversity. *Nature Communications*, *13*(1), 1628.

819   https://doi.org/10.1038/s41467-022-29324-2

820   Mccleery, R. A., Lopez, R. R., Silvy, N. J., & Kahlick, S. N. (2007). Habitat Use of Fox

821   Squirrels in an Urban Environment. *The Journal of Wildlife Management*, *71*(4),

822   1149–1157. https://doi.org/10.2193/2006-282

823   Mcdonnell, M. J., & Hahs, A. K. (2008). *The use of gradient analysis studies in*

824   *advancing our understanding of the ecology of urbanizing landscapes: Current*

825   *status and future directions*. https://doi.org/10.1007/s10980-008-9253-4

826   McDonnell, M. J., & Pickett, S. T. A. (1990). Ecosystem Structure and Function along

827   Urban-Rural Gradients: An Unexploited Opportunity for Ecology. *Ecology*, *71*(4),

828   1232–1237. https://doi.org/10.2307/1938259

829   McElreath, R. (2020). *Statistical Rethinking: A Bayesian Course with Examples in R and*

830   *STAN*. CRC Press.

831   McRobie, H. R., Moncrief, N. D., & Mundy, N. I. (2019). Multiple origins of melanism in

832   two species of North American tree squirrel (Sciurus). *BMC Evolutionary Biology*,

833   *19*(1), 1–14. https://doi.org/10.1186/s12862-019-1471-7

834   McRobie, H., Thomas, A., & Kelly, J. (2009). The Genetic Basis of Melanism in the Gray

835   Squirrel (Sciurus carolinensis). *Journal of Heredity*, *100*(6), 709–714.

836   https://doi.org/10.1093/jhered/esp059

837   Merrick, M., Evans, K., & Bertolino, S. (2016). *Urban grey squirrel ecology, associated*

838   *impacts, and management challenges* (pp. 57–77).

839   Moll, R. J., Cepek, J. D., Lorch, P. D., Dennis, P. M., Tans, E., Robison, T., Millspaugh,

840   J. J., & Montgomery, R. A. (2019). What does urbanization actually mean? A

framework for urban metrics in wildlife research. *Journal of Applied Ecology*,

56(5), 1289–1300. https://doi.org/10.1111/1365-2664.13358

Moore, J. F., Pine, W. E., Mulindahabi, F., Niyigaba, P., Gatorano, G., Masozera, M. K.,

& Beaudrot, L. (2020). Comparison of species richness and detection between

line transects, ground camera traps, and arboreal camera traps. *Animal*

Conservation, *23*(5), 561–572. https://doi.org/10.1111/ACV.12569

Mumma, M. A., Gillingham, M. P., Johnson, C. J., & Parker, K. L. (2019). Functional

responses to anthropogenic linear features in a complex predator-multi-prey

system. *Landscape Ecology*, *34*(11), 2575–2597. https://doi.org/10.1007/s10980-

019-00905-9

Nakamura, A., Kitching, R. L., Cao, M., Creedy, T. J., Fayle, T. M., Freiberg, M., Hewitt,

C. N., Malhi, Y., Mitchell, A., Novotny, V., Ozanne, C. M. P., Song, L., Wang, H.,

& Ashton, L. A. (2017). Forests and Their Canopies: Achievements and Horizons

in Canopy Science. *Trends in Ecology & Evolution*, *32*, 438–451.

https://doi.org/10.1016/j.tree.2017.02.020

Neilly, H., Nordberg, E. J., Vanderwal, J., & Schwarzkopf, L. (2017). Arboreality

increases reptile community resistance to disturbance from livestock grazing.

Journal of Applied Ecology, *July 2017*, 786–799. https://doi.org/10.1111/1365-

2664.12982

Nowak, D. J., Hoehn, R. E., Bodine, A. R., Greenfield, E. J., & O'Neil-Dunne, J. (2016).

Urban forest structure, ecosystem services and change in Syracuse, NY. *Urban*

Ecosystems, *19*(4), 1455–1477. https://doi.org/10.1007/s11252-013-0326-z

Ordeñana, M. A., Crooks, K. R., Boydston, E. E., Fisher, R. N., Lyren, L. M., Siudyla, S.,

864   Haas, C. D., Harris, S., Hathaway, S. A., Turschak, G. M., Miles, A. K., & Vuren,

865   D. H. V. (2010). Effects of urbanization on carnivore species distribution and

866   richness. *Journal of Mammalogy*, *91*(6), 1322–1331. https://doi.org/10.1644/09-

867   MAMM-A-312.1

868   Parker, T. S., & Nilon, C. H. (2008). Gray squirrel density, habitat suitability, and

869   behavior in urban parks. *Urban Ecosystems*, *11*(3), 243–255.

870   https://doi.org/10.1007/s11252-008-0060-0

871   Plummer, M. (2017). *JAGS Version 4.3.0 user manual.*

872   https://sourceforge.net/projects/mcmc-jags/files/Manuals/4.x/

873   Price Tack, J. L., McGowan, C. P., Ditchkoff, S. S., Morse, W. C., & Robinson, O. J.

874   (2018). Managing the vanishing North American hunter: A novel framework to

875   address declines in hunters and hunter-generated conservation funds. *Human*

876   *Dimensions of Wildlife*, *23*(6), 515–532.

877   https://doi.org/10.1080/10871209.2018.1499155

878   Rankin, E. E. W., Knowltonid, J. L., Gruner, D. S., Flaspohler, D. J., Giardina, C. P.,

879   Leopold, D. R., Buckardt, A., Pitt, W. C., & Fukami, T. (2018). Vertical foraging

880   shifts in Hawaiian forest birds in response to invasive rat removal. *PLoS ONE*,

881   *10*. https://doi.org/10.1371/journal.pone.0202869

882   Reed, J. H., & Bonter, D. N. (2018). Supplementing non-target taxa: Bird feeding alters

883   the local distribution of mammals. *Ecological Applications*, *28*(3), 761–770.

884   https://doi.org/10.1002/eap.1683

Regolin, A. L., Cherem, J. J., Graipel, M. E., Bogoni, J. A., Ribeiro, J. W., Vancine, M.

H., Tortato, M. A., Oliveira-Santos, L. G., Fantacini, F. M., Luiz, M. R., Castilho,

P. V. de, Ribeiro, M. C., & Cáceres, N. C. (2017). Forest cover influences

occurrence of mammalian carnivores within Brazilian Atlantic Forest. *Journal of*

Mammalogy, *98*(6), 1721–1731. https://doi.org/10.1093/jmammal/gyx103

Ritchie, H., & Roser, M. (2018). Urbanization. *Our World in Data*.

https://ourworldindata.org/urbanization

Robertson, G. I. (1973). *Distribution of color morphs of Sciurus carolinensis in eastern*

North America. Unpubl [PhD Thesis]. MS Thesis, Univ. Western Ontario, London.

Robinson, S. K., Thompson, F. R., Donovan, T. M., Whitehead, D. R., & Faaborg, J.

(1995). Regional Forest Fragmentation and the Nesting Success of Migratory

Birds. *Science*, *267*(5206), 1987–1990.

https://doi.org/10.1126/science.267.5206.1987

Royle, J. A. (2004). N-Mixture Models for Estimating Population Size from Spatially

Replicated Counts. *Biometrics*, *60*(1), 108–115. https://doi.org/10.1111/j.0006-

341X.2004.00142.x

Santangelo, J. S., Miles, L. S., Breitbart, Sophie T., Murray-Stoker, D., Rivkin, L. R.,

Johnson, Marc T., Ness, R. W., Szulkin, M., Munshi-South, J., & Charmantier, A.

(2020). Urban environments as a framework to study parallel evolution. In *Urban*

Evolutionary Biology (p. 36).

Santangelo, J. S., Rivkin, L. R., & Johnson, M. T. (2018). The evolution of city life. *Proc.*

R. Soc. B, *285*, 20181529–20181529. https://doi.org/10.1098/rspb.2018.1529

Sarno, R. J., Parsons, M., & Ferris, A. (2015). Differing vigilance among gray squirrels

(Sciuridae carolinensis) along an urban–rural gradient on Long Island. *Urban*

Ecosystems, *18*(2), 517–523. https://doi.org/10.1007/s11252-014-0414-8

Schneider, N. A., Low, M., Arlt, D., & Pärt, T. (2012). Contrast in Edge Vegetation

Structure Modifies the Predation Risk of Natural Ground Nests in an Agricultural

Landscape. *PLOS ONE*, *7*(2), e31517.

https://doi.org/10.1371/journal.pone.0031517

Shipley, B. (2009). Confirmatory path analysis in a generalized multilevel context.

Ecology, *90*(2), 363–368. https://doi.org/10.1890/08-1034.1

Szabó, B., Korányi, D., Gallé, R., Lövei, G. L., Bakonyi, G., & Batáry, P. (2023).

Urbanization decreases species richness, and increases abundance in dry

climates whereas decreases in wet climates: A global meta-analysis. *Science of*

The Total Environment, *859*, 160145.

https://doi.org/10.1016/j.scitotenv.2022.160145

Thompson, J. R., Carpenter, D. N., Cogbill, C. V., & Foster, D. R. (2013). Four

Centuries of Change in Northeastern United States Forests. *PLOS ONE*, *8*(9),

e72540. https://doi.org/10.1371/journal.pone.0072540

Tucker, M. A., Santini, L., Carbone, C., & Mueller, T. (2021). Mammal population

densities at a global scale are higher in human-modified areas. *Ecography*,

44(1), 1–13. https://doi.org/10.1111/ecog.05126

Wassmer, T., & Refinetti, R. (2019). Individual Daily and Seasonal Activity Patterns in

Fox Squirrels (Sciurus niger) Quantified by Temperature-Sensitive Data Loggers.

Frontiers in Ecology and Evolution, *7*. https://doi.org/10.3389/fevo.2019.00179

Winchell, K. M., Reynolds, R. G., Prado-Irwin, S. R., Puente-Rolón, A. R., & Revell, L. J.

(2016). Phenotypic shifts in urban areas in the tropical lizard Anolis cristatellus.

Evolution, *70*(5), 1009–1022. https://doi.org/10.1111/evo.12925

Wright, W. J., Irvine, K. M., & Higgs, M. D. (2019). Identifying occupancy model

inadequacies: Can residuals separately assess detection and presence?

Ecology, *100*(6), e02703. https://doi.org/10.1002/ecy.2703

Xu, H., Shi, T., Wang, M., Fang, C., & Lin, Z. (2018). Predicting effect of forthcoming

population growth–induced impervious surface increase on regional thermal

environment: Xiong'an New Area, North China. *Building and Environment*, *136*,

98–106. https://doi.org/10.1016/j.buildenv.2018.03.035

Zanaga, D., Van De Kerchove, R., Daems, D., De Keersmaecker, W., Brockmann, C.,

Kirches, G., Wevers, J., Cartus, O., Santoro, M., Fritz, S., Lesiv, M., Herold, M.,

Tsendbazar, N.-E., Xu, P., Ramoino, F., & Arino, O. (2022). *ESA WorldCover 10*

m 2021 v200 (Version v200) [Dataset]. Zenodo.

https://doi.org/10.5281/zenodo.7254221

945
946
947