Multiple LPA3 receptor agonist binding sites evidenced under docking and functional studies

K. Helivier Solís[1], M. Teresa Romero-Ávila[1], Ruth Rincón-Heredia[1], José Correa-Basurto[2], and J. Adolfo García-Sáinz[1]*

[1] Departamento de Biología Celular y Desarrollo and Unidad de Imagenología, Instituto de Fisiología Celular, Universidad Nacional Autónoma de México. Ciudad Universitaria. Ap. Postal 70-600, Ciudad de México 04510. México

[2] Laboratorio de Diseño y Desarrollo de Nuevos Fármacos e Innovación Biotecnológica, Escuela Superior de Medicina, Instituto Politécnico Nacional. Plan de San Luis y Díaz Mirón S/N, Casco de Santo Tomas, Miguel Hidalgo, Ciudad de México11340, México.

* Correspondence: J. A. García-Sáinz. Inst. Fisiología Celular, UNAM Ciudad Universitaria, Ap. Postal 70-600
Ciudad de México 04510 México Email: agarcia@ifc.unam.mx

Abstract: Comparative studies using (lysophosphatidic acid) LPA and the synthetic agonist, oleoyl-methoxy glycerophosphothionate, OMPT, in cells expressing the LPA3 receptor revealed differences in the action of these agents. The possibility that more than one recognition cavity might exist for these ligands in the LPA3 receptor was considered. We performed agonist docking studies exploring the whole protein to obtain tridimensional details of the ligand-receptor interaction. Functional in-cellulo experiments using mutants were also executed. Our work includes blind docking using the unrefined and refined proteins subjected to hot spot predictions. Distinct ligand protonation (charge -1 and -2) states were evaluated. One LPA recognition cavity is located near the lower surface of the receptor close to the cytoplasm (Lower Cavity). OMPT displayed an affinity for an additional identification cavity detected in the transmembrane and extracellular regions (Upper Cavity). Docking targeted to Trp102 favored binding of both ligands in the transmembrane domain near the extracellular areas (Upper Cavity), but the associating amino acids were not identical due to close sub-cavities. In the in-cellulo studies, LPA action was much less affected by the distinct mutations than that of OMPT (which was almost abolished). Docking and functional data suggest distinct agonist binding cavities in the LPA3 receptor

Keywords: Lysophosphatidic acid; LPA; LPA3 receptor; oleoyl-methoxy glycerophosphothionate; OMPT; biased agonism; ligand docking

1. Introduction

Lysophosphatidic acid (LPA) is a bioactive lipid that exerts many of its actions through six G protein-coupled receptors (GPCRs) that constitute the LPA subfamily [1-3]. Among them is the LPA3 receptor, which mainly couples to Gαi/o and Gαq/11, participating in diverse signaling events that lead to various physiological and pathophysiological effects [4, 5]. Structurally, the LPA3 receptor is a membrane protein of 353 amino acids containing 7 transmembrane domains (TM), which are connected by three intracellular (IL) and three extracellular (EL) loops; it comprises an extracellular amino terminus and an intracellular carboxyl end [2, 4, 5]. Though it has not been crystallized yet, it has a similarity of approximately 53% with the LPA1 receptor, the best-studied member of this group [3, 6-8]. It is worth mentioning that the domains that regulate LPA1 receptor function are not conserved in the LPA3 receptor; these include a dileucine motif, a mono ubiquitination residue, and a PDZ pattern, among others [5, 9]. Structural analysis of the LPA3 receptor showed that it conserves motifs/domains typical of the A family of GPCRs, which comprise the CWxP, EHR, PIF, and NPxxY areas, each having a specific function during and after agonist attachment [2, 5, 10].

It is generally accepted that the receptor's recognition cavities are flexible, and that ligand binding promotes conformational changes that impact their spatial configuration (https://pubmed.ncbi.nlm.nih.gov/33076289/). These dynamic events allow displacements of the intracellular loops and the TM regions, which promote G-protein recruitment and exposure of sites that can be phosphorylated by distinct protein kinases, including G protein-coupled receptor kinases (GRKs) and second messenger-activated protein kinases [10-13]. Receptor phosphorylation favors β-arrestin-receptor interaction, triggering its internalization and subsequent signaling events [14-16]. Determining how a ligand interrelates with a given receptor provides fundamental insights into the search for drugs that could effectively regulate its activity and the participating mechanisms. In this sense, molecular docking simulations have become a method that allows a better knowledge of the receptor regions in which the ligand could bind, suggesting the possible residues involved [17].

Pioneer structural work with the LPA3 receptor identified amino acids that interact with the polar head of LPA [7, 18-21]. When these residues were mutated, its function was altered, particularly those in the TM domains 3, 4, 5, and 7, mainly involving the following sites: Arg105 (3.28), Gln106 (3.29), Trp153 (4.64), Arg209 (5.38), and Lys296 (7.35) (in parentheses are the Ballesteros and Weinstein [22] numbering scheme; these numbers were obtained from the server: https://gpcrdb.org/protein/lpar3_human/ [23]. It is relevant that mutation of residues Trp153 (4.64) and Lys296 (7.35) directly impacts ligand binding and activation of the LPA3 receptor, i.e., LPA potency and efficacy were decreased in RH7777 cells transfected with mutant receptors when compared to those expressing the wild type [20].

No specific agonist for the LPA3 receptor subtype is available since most also activate the LPA1 and LPA2 receptors. A ligand that shows high potency for LPA3 is 1-oleoyl-2-methyl-sn-glycero-3-phosphothionate (OMPT) [24]. OMPT and related analogs exhibit affinity for different LPA receptors, but they have a weak selectivity for the LPA receptor subtypes [25-27]; despite these selectivity problems, OMPT remains a reference for studies on LPA3 receptors. To our knowledge, the binding site for OMPT in LPA receptors has not been explored.

It was recently reported that LPA and OMPT exhibit marked pharmacodynamic differences in LPA3-expressing cells [28]. In particular, two findings were astonishing: firstly, OMPT exerts its

effects within two very distinct concentration ranges (1-10 nm and > 100 nM), and secondly, LPA induced a "receptor refractory state" that blocks calcium response to re-challenging with this natural agonist; however, under the same conditions, OMPT was able to produce a conspicuous response [28]. Both groups of data suggest the possibility that distinct binding cavities might exist for these agonists in the LPA3 receptor: one showing a very high affinity for OMPT and another one, likely shared by LPA and OMPT (which are structural analogs) with a lower affinity for both agonists in the range of 100-300 nM [28]. Intrigued by such findings and lacking structural information on the OMPT-LPA3 receptor interaction, we performed ligand-receptor docking studies to get insights into their protein binding pocket(s). The docking examination and the hot spot predictions indicate the presence of at least two distinct ligand recognition cavities, which might explain the observed functional responses to OMPT. In addition, receptor mutants were constructed and expressed, and their response to these agonists was studied. One of the mutants showed activity in reaction to LPA and essentially no effect on OMPT, which is consistent with the proposal of multiple binding pockets for these ligands [28]. This report documents distinct agonist identification cavities via docking simulations and in-cellulo experiments with receptor mutants. The findings open new experimental possibilities and might help better understand LPA3 receptor-ligand interaction and aid in developing new agonists and antagonists of potential therapeutic use.

2. Results

2.1 LPA3 receptor model without refinement.

The LPA3 receptor has not been crystallized yet; therefore, its most likely 3D structure was obtained using different servers for configuration prediction (additional information is provided in section 4.1). Supplementary Table S1 enlists the values obtained using these servers and programs. Protein arrays from the distinct patterns were subjected to sequence alignments (Supplementary Fig. S1), and the Swiss Model was preferred because the best structural score values were obtained, and the resulting sequence was identical to the UniProt Q9UBY5 entry.

The unrefined 3D pattern from the Swiss Model program was employed to determine the binding parameters (free energy values (ΔG), Kd, and residue interaction) for the LPA3 receptor agonists, LPA and OMPT (a 126 Å3 grid box centered on the protein with (x, y, and z) coordinates in grid center for both agonists: 6.014, -0.124, and -2.385). Molecular docking simulation showed that the LPA3 receptor recognition cavities for LPA and OMPT were different (Supplementary Table S2), and the binding energy value was lower for OMPT than for LPA. The residues participating in ligand-receptor interaction are also indicated in Supplementary Table S2; please notice that the amino acids involved are listed in the order indicated by the program, considering their importance in the ligand-receptor interaction. Supplementary Fig. S2 depicts such interrelations using the unrefined model. OMPT (red) associates with amino acids located in TM regions 2 and 7 and the EL 1, 2 (we name this broad region Upper Cavity; see Supplementary Fig S2), whereas LPA (blue) reached a distinct site (broad region named Lower Cavity) of the LPA3 receptor structure with molecules preferentially located in IL 2 and IL 3, very close to the cytoplasm (Supplementary Fig. S2). The cartoon shown in Supplementary Fig. S3 depicts the LPA-interacting residues (blue) and OMPT-associating amino acids (red) (notice the remarkably different distribution of such binding cavities).

The 2D maps of the ligand-receptor interrelations obtained from the Discovery Studio program are also shown (Supplementary Figs. S4 (LPA) and S5 (OMPT)). This methodology allows adaptations of the ligand to obtain the best possible interaction with the receptor and performs the necessary adjustments to achieve the most favorable association; it also permits determining the agonist orientation, identifying the receptor's amino acids involved, and defining the type of interaction (Supplementary Table S2). Notice that the phosphate group of LPA is fully protonated (i.e., net charge 0; Supplementary Figure S4), but in contrast, the phosphorothioate group of OMPT is partially deprotonated (net charge -1; Supplementary Figure S5) when interacting with the distinct LPA3 receptor-ligand recognition cavities.

The polar head groups of agonists such as LPA and OMPT seem important when these ligands bind to LPA3 receptors and for their activation [7, 18-20, 25]. LPA's phosphate and OMPT's phosphorothionate can be protonated (net charge 0), partially deprotonated (net charge -1), or fully deprotonated (net charge -2) at distinct pH values. Experimentally defined pKa values for these compounds have not been reported, but those of the analog, phosphatidic acid, are available (pKa1=3.0 and pKa2=8.0; https://avantiresearch.com/tech-support/physical-properties/ionization-constants). Assuming that the pKa values of LPA and OMPT could be similar to those of phosphatidic acid, at a pH of 7.4, the protonated species (charge 0) would represent < 0.1% of the total amount of the compound, whereas the partially deprotonated (charge -1) and fully deprotonated (charge -2) species would represent ≈ 79% and ≈ 20% of the total, respectively. The docking simulations using the unrefined receptor structure employed the protonated agonist species (charge 0). For all the following simulations, the three molecular species were employed (i.e., species with net charges 0, -1, and -2).

2.2 LPA3 refined Model.

The molecular docking studies of protonated (charge 0) LPA (Fig. 1; blue) show that it occupied the Lower Site of the receptor, interacting with the IL 2 and 3, which, like in the LPA3 unrefined model, was very close to the cytoplasm. On the other hand, OMPT (charge 0) (Fig. 1; red) associates with the amino acids located in the Upper cavity, including TM 5, 6, 7, and 8, as well as the extracellular loop 3 (Fig. 1). Interestingly, simulation with both charge -1 ligands resulted in inversion of the agonist position in these cavities (Fig. 1). Simulation with charge -2 derived in interaction patterns similar to those observed with the uncharged ligands (Fig. 1). Tables 1 (for LPA) and 2 (for OMPT) provide the ΔG and Kd values and the list of interacting amino acids defined for the distinct conditions. Simulation with LPA (charge -2) resulted in a more negative ΔG and lower Kd values compared to the other conditions (Table 1); in contrast, modification of OMPT charge had only a minor impact on these parameters (Table 2). Cartoons showing the interrelating amino acids under these conditions are presented in Supplementary Fig. S6, and the 2D schemes showing the interacting residues and the type of association are presented in Supplementary Figs. S7.

LPA₃
LPA (Blue) , OMPT (Red)

Charge 0   Charge -1   Charge -2

Figure 1. 3D images resulting from the LPA3-LPA (blue) and LPA3-OMPT (red) docking analysis using PyMol. The refined structure was employed. Ligand net charges are indicated.

Table 1. Thermodynamic parameters and interacting amino acids were determined in the distinct docking simulations using LPA as the ligand.

Focus (Charge)	ΔG (Kcal/mol)	Kd (μM)	Interacting Amino acids
NONE (0)	-5.01	≈ 210	Val219, Leu220, Ser221, Lys216, Arg230, Pro234, Val136
NONE (-1)	-4.10	≈ 995	Arg276, Trp277, Val22, His273, Val274, Lys 29, Trp25, Ile 32, Leu 280, Val33, Val 36, Tyr83, Gly 37, Leu283, Met 87, Leu86, Cys41
NONE (-2)	-7.18	≈ 5	Lys216, Thr217, Ile131, Met134, Arg230, Pro234, Leu237, Arg127, Val136
Trp102 (0)	-5.99	≈ 40	Leu283, Leu280, Leu279, Met87, Val33, Thr90, Trp25, Arg276, Lys29
Trp102 (-1)	-4.10	≈ 990	Gln106, Arg105, Lys275, Leu86, Tyr14, Leu109, Trp102, Ala180, Leu279, Thr90, Asp23, Gly91,

Focus (Charge)	ΔG (Kcal/mol)	Kd (μM)	Interacting Amino acids
			Arg276, Val36, Val33, Leu280, Trp277, Trp225, Lys29, Ile32
Trp102 **(-2)**	-8.93	≈ 0.30	Trp277, Leu279, Arg276, Gly37, Val33, Trp25, Lys29, Thr90, Leu86, Met87
Tyr293 **(0)**	-5.79	≈ 60	Leu220, Lys216, Thr217, Ile131, Val213, Pro234, Arg135, Met134, Val136, Asn139, His137, Phe63
Tyr293 **(-1)**	-2.88	≈ 8000	Thr146, Lys142, Val145, Trp153, Ile149, Leu150, Tyr67, Val55, Ile56, His62, Ala70, Asn71
Tyr293 **(-2)**	-7.62	≈ 3	Trp277, Leu280, Leu279, Leu283, Tyr83, Arg276, Gly37, Val33, Trp25, Leu86, Met87

Table 2. Thermodynamic parameters and interacting amino acids were determined in the distinct docking simulations using OMPT as the ligand.

Focus (Charge)	ΔG (Kcal/mol)	Kd (μM)	Interacting Amino acids
NONE **(0)**	-5.67	≈ 70	Trp277, Leu279, Leu280, Leu283, Arg276, Tyr14, Asp23, Trp25, Ser94, Thr90, Arg105, Leu86, Val33
NONE **(-1)**	-5.36	≈ 120	Lys216, Val213, Ile131, Met134, Arg230, Val136, Arg135
NONE **(-2)**	-5.85	≈ 50	Ser94, Arg105, Leu86, Asp23, Lys275, Arg276, Leu279, Val36, Trp277, Val33, Ile32
Trp102 **(0)**	-5.15	≈ 170	Leu283, Leu280, Leu279, Tyr83. Val36, Met87, Val33, Thr90, Trp25, Arg276, Trp277, Leu86
Trp102 **(-1)**	-7.75	≈ 6	Leu256, Leu159, Leu188, Phe278, Lys275, Trp191, Leu179, Ala180, Tyr183, Tyr187, Gln106, Arg105, asp110, Ala282, Leu279, Trp252

Trp 102 (-2)	-7.11	≈ 6	Leu280, Leu279, Leu283, Ile32, Gly37, Val36, Val33, Arg276, Trp25, Met87
Tyr293 (0)	-4.64	≈ 395	Arg230, Leu220, Lys216, Thr217, Arg231, Ile131, His137, Val136, Thr233, Pro234
Tyr293 (-1)	-5.76	≈ 60	Arg232, Met235, Lys239, Lys236, Thr240, Tyr295, Glu298, Ser294, Tyr301, Ile291
Tyr293 (-2)	-5.64	≈ 75	Lys296, Lys236, Thr240, Lys305, Ile308, Tyr295, Glu298, Tyr301, Ser294, Met304,Ile291

2.3 Trp102-focused model.

Trp102 is located near the Upper Cavity or the LPA3 receptor configuration (Supplementary Fig. 8). As previously mentioned, according to 3D structural analysis with PyMol., this residue was consistently found in the hot spot predictions and visualized as exposed and centered in a cavity. Trp102-focused docking was performed for LPA and OMPT with distinct ligand protonations as indicated. Independently of the molecule charge, both agonists interacted with the receptor's Upper cavity (Fig. 2). The uncharged ligands interrelated with the TM regions 1, 2, and 7. LPA (charge -1) interconnects with TM domains 1, 2, 3, and 7 and EL2, whereas OMPT (charge -1) binds to amino acids in TM domains 3, 5, 6, and 7 and EL 3. Simulations with both charge -2 ligands lead to their interaction with residues in TM 1 and 7 and EL 2. Cartoons showing the participating amino acids under these conditions are presented in Supplementary Fig. S9, and the 2D schemes showing the interacting residues and their type of association are illustrated in Supplementary Fig. S10. For LPA (Charge -2), Trp102-focused docking resulted in a more negative ΔG and lower Kd values (Table 1); similarly, OMPT (charges -1 and -2) also derived more favorable

thermodynamic values (Table 2). The cartoons and the 2D images (Figs. S9 and S10) evidenced that both ligands shared some amino acids under these conditions.

Figure 2. Images resulting from the docking analysis of the refined 3D structure of LPA3 receptor (Trp102 focused) interacting with LPA (blue) and OMPT (red). Ligand net charges are indicated.

2.4 Tyr293-focused model.

The localization of Tyr293 in the LPA3 structure is shown in Supplementary Fig. S8. This residue participates in the attachment of the LPA antagonist, Kil6425, and an extensive series of pharmacological agents used in therapeutics [29]. Figure 3 presents the non-bond interaction of LPA (blue) and OMPT (red) with the LPA3 receptor when the docking analysis is focused on Tyr293. Using ligands with charge 0 for the docking simulation showed that the LPA binding cavities were located near the cytoplasm in the intracellular loops 1, 2, and 3. OMPT-LPA3 receptor interaction occurred in the Upper Cavity involving TM domains 1, 7, and EL 2 and 3. When simulations were performed using both ligands with charge -1, the agonists interrelated with the lower part of the receptor (LPA) IL1, IL2, and TM 4; for OMPT binding residues at IL3, Helix 8 and TM 7 played a role. With the receptor focused on Tyr 293, both molecules with charge -2 interacted with a pattern opposite to that observed at charge 0. Cartoons illustrating the partaking amino acids under these conditions are presented in Supplementary Fig S11, and the 2D schemes showing the participating residues and their type of association are depicted in Supplementary Fig12. Table 1 shows that LPA (charge -2) exhibited a relevant negative ΔG and a small Kd value. OMPT charge causes minor modifications of these thermodynamic parameters.

LPA₃ Tyr293
LPA (Blue) , OMPT (Red)

Charge 0   Charge -1   Charge -2

Figure 3. Images resulting from the docking analysis of the refined 3D structure of the LPA₃ receptor (Tyr293 focused) interacting with LPA (blue) and OMPT (red). Ligand net charges are indicated.

2.5 In cellulo signaling experiments

Mutant receptors in which the ligand-interacting residues (unrefined model) in the cavities were substituted for alanine were expressed, and their calcium response to 1 µM LPA and 1µM OMPT was determined. The reaction of cells manifesting the wild-type receptor was considered the reference and was compared to the mutants in the upper cavity, lower cavity, and both. Representative calcium tracings were presented in Fig. 4, and the data analysis is shown in Fig. 5. In the control cells, as anticipated, 1µM LPA or 1µM OMPT triggered an almost immediate increase in intracellular calcium. In contrast, a very weak response to OMPT was observed with the distinct mutants. The reaction to LPA was diminished but not blocked, and the Upper Cavity mutant showed a robust effect. These data indicate that some residues are much more relevant for the action of OMPT than for that of LPA. Similar findings were obtained with these mutants when ERK 1/2 phosphorylation was studied (i.e., the response to LPA was diminished in cells expressing the mutant receptors, while that to OMPT was almost abolished) (unpublished data).

Figure 4. Representative calcium tracings were obtained in cells expressing

the wild-type LPA$_3$ receptors or mutants with substitutions in Cavity 1 (C1 mutant), Cavity 2 (C2 mutant), or both Cavities (C1+2 mutant). Cells were challenged with 1 μM LPA (right) or 1 μM OMPT (left).

Figure 5. Calcium tracings were obtained in cells expressing the wild-type LPA$_3$ receptors or mutants with substitutions in Cavity 1 (C1 mutant), Cavity 2 (C2 mutant), or both Cavities (C1+2 mutant). Cells were challenged with 1 µM LPA or 1 µM OMPT. The means are plotted, and vertical lines indicate the SEM of 4-5 experiments performed on different days using distinct cell cultures. **p < 0.01, and ****p < 0.0001 vs. cell expressing the wild-type receptor

3. Discussion

The present work was designed to define if LPA and OMPT bind to a single cavity or different agonist recognition pockets in the LPA3 receptor. The possible existence of more than one ligand-identification region was suggested in experiments using cultured cells [28]. Blind docking studies were performed using the predicted configuration of LPA3 (UniProt, Q9UBY5 entry). 3D structures of refined and focused (to Tyr293 and Trp102) docking analyses were carried out employing the distinct protonation states of the ligands. As indicated, Tyr293 was identified as a critical amino acid for several agents interacting with the LPA3 receptor [29], and Trp102 was consistently detected in the hot spot prediction.

In our work, we observed the docking of LPA and OMPT to two principal LPA3 receptor regions named, for this work, Upper and Lower Cavities. This pattern was noticed in both the unrefined and refined docking analysis and with changes in the net charge of the ligands. In fact, under most of these conditions LPA was localized in the Lower Cavity. This location was reversed when a charge of -1 was assigned to both ligands. Previous work [7, 18-21] has detected a central binding cavity with emplacement similar to the Upper Cavity reported here. In these studies, a box of 60 Å3 was employed, whereas in our work, we used a 126 Å3 box and a grid space of 0.375 Å3. Therefore, it seems likely that space limitations (i.e., using a box of 60 Å3) might have impeded the Lower Cavity detection. Avoiding expanse restrictions has positive and negative aspects. When the ligand recognition cavity(s) is not known, it seems convenient to explore the entire protein surface by docking, a procedure named "blind docking", which implies using a large box [30]; however, focusing on predicted binding regions is known to improve docking accuracy and efficiency [30].

It should be mentioned that the Upper Cavity is in the TM core of the receptor, where most agonists associate with class A GPCRs. The Lower Cavity observed in this work included ligand interaction with ICLs and hardly any contact with the TM domains, which is an uncommon feature. This Lower Cavity might represent a gating point for the molecules rather than an agonist-activating region. However, with the current information, determining the functional relevance of these Cavities is impossible. The operational data obtained with the mutants of these pockets indicate that amino acid substitution in any of them markedly altered the action of OMPT but, to a much lesser extent, that of LPA. These findings support the idea that the effect of these two agonists does not involve the same receptor associations but rather that distinct residues participate in receptor activation.

Previous work, also using docking/computational modeling and mutants, identified that the LPA3 receptor interacted mainly with the polar head of LPA at Arg105 (3.28), Gln106 (3.29), Trp153 (4.50), Arg209 (5.60), Arg276 (7.36) and Lys275 (7.35) [7, 18-21]. These amino acids were considered critical for LPA binding to the receptor and their mutation decreased agonist efficacy and potency and as expected, such modifications eliminated ionic interactions in the modeled LPA3-ligand prototype [7, 18-21]. It is important to mention that the same amino acids were identified in our docking studies. The residues detected were Arg276 (9 times), Arg105 (4 times), Lys275 (3 times), Gln106 (2 times), and Trp153 (1 time); this information includes data for LPA and OMPT. Other interacting amino acids frequently found in the docking simulations for both ligands include: Val33

(9 times), Trp25 (7 times), Trp277 (7 times), Leu280 (7 times), Leu86 (7 times), and Leu283 (6 times) among others.

The similarities of our present findings with those published earlier [7, 18-21] suggest the possibility that the LPA binding site observed in the Upper Cavity might correspond to that previously determined [7, 18-21]. No prior docking data for LPA3-OMPT interaction has been published. Our findings indicate that OMPT occupies the same Upper Cavity as LPA. Interestingly, the functional information formerly published [28] and that using mutants presented here strongly suggest that differences among these agonist-recognizing regions exist; therefore, they should be considered distinct, although some interacting residues could be shared. Exploration of the LPA3 receptor cavities, using PyMol, showed many capable of ligand binding (Supplementary Fig. S13). The localization of the agonists was included in the analysis, employing the Trp102 docking simulations (charges -1 and -2). It can be observed that LPA and OMPT occupy close but not identical positions; in addition, it seems likely that molecules with distinct charges (i.e., mixtures of agonists with charges -1 and -2) might interact with LP3. Very recent data using single-particle cryo-EM structures of human S1P1 receptor bound to Siponimod and heterotrimeric Gi complexes, as well as human LPA1 receptor and Gi complexes in the presence of LPA indicated that the ligands adopt different conformations to interact with their cognate GPCRs [31]. Such distinct molecule configurations might participate in the agonist-induced LPA3 activation. Similar structural approaches likely would be required to define the precise residues that interact with the different ligands and their diverse conformations. The present findings open new research possibilities to explore this crucial step in LPA3 receptor signaling, considering the physiological and physiopathological events in which it is involved [4, 5].

4. Materials and Methods

4.1 Modeling of LPA3 receptor structure, Gene ID:23566.

The LPA3 receptor has not yet been crystallized. Therefore, we were forced to predict its most likely 3D structure using the amino acid sequence obtained from the Uniprot database (Q9UBY5 entry) (https://www.uniprot.org/uniprotkb/Q9UBY5/entry). Three of the most frequently utilized servers for structure prediction were employed: SWISS-MODEL (https://swissmodel.expasy.org/), I-TASSER (https://zhanggroup.org/I-TASSER/), and AlphaFold database (https://alphafold.ebi.ac.uk/). Supplementary Table S1 enlists the values obtained using these servers and programs. Subsequently, protein arrays from the three models were subjected to sequence alignments (Supplementary Fig. S1) employing the PROMALS3D server (http://prodata.swmed.edu/promals3d/promals3d.php) to determine the gaps or residue insertions that improve the arrangement identity with the 3D protein templates during the folding of the molecule. Although the data obtained showed that the 3D construction built with AlphaFold was the best evaluated structurally, the alignment revealed that several mutations or deletions improved the 3D configuration quality, possibly eliminating residues that can interact with the ligands. Therefore, the Swiss Model was preferred because the best values were estimated by ERRAT and a second evaluation of Ramachandran maps; additionally, the sequence obtained was identical to the UniProt Q9UBY5 entry.

4.2 Refined and validation of the LPA3 receptor 3D structure.

Protein 3D structure refinement was carried out to perform accurate docking simulations. The arrangement obtained from the Swiss Model server was subjected to molecular refinement using ReFOLD (https://www.reading.ac.uk/bioinf/ReFOLD/ReFOLD3_form.html). The refined protein with the best score was used for the docking study to explore receptor cavities capable of ligand recognition.

4.3 Search for ligand-binding cavities on the LPA3 receptor.

The hot spot prediction was performed to identify possible ligand-recognition cavities using the BetaCavityWeb (http://voronoi.hanyang.ac.kr/betacavityweb/) and the Fpocketweb (http://durrantlab.com/fpocketweb) servers. The protein from the Swiss Model server, as well as that from ReFOLD, were used. Both servers calculate these cavities for a given molecular structure and a defined spherical probe and report their geometrical properties, such as volume, boundary area, buried area, and other parameters described on their websites. In proteins, cavities can be predicted mainly at the barrel end built by an alpha secondary structure, a pattern observed in the refined and unrefined molecule.

4.4 LPA and OMPT Docking at the LPA3 receptor binding sites.

The 3D configurations of the ligands studied were obtained from PubChem (ID 5497152 for LPA and ID 16078994 for OMPT), and their molecular structure minimization was carried out using the Avogadro program (https://avogadro.cc/docs/tools/auto-optimize-tool/); which allows the net inter-atomic force on each atom to be acceptably close to zero and the position on the potential energy surface is such that the improved structures correspond to the configurations found in nature. The lengths and angles of the links and junctions and their geometry were appropriated and transmitted with optimized energy.

After ligand structure minimization, a blind docking study of the native protein (unrefined) was performed, allowing us to explore the whole molecule. It is worth noticing that it has been reported that Tyr293 participates in the binding of the LPA antagonist, Kil6425, as well as of an extensive series of pharmacological agents used in therapeutics [22]. Residue Trp102 was also particularly interesting because it was consistently found in the hot spot predictions and visualized as exposed and centered in a cavity according to 3D structural analysis with PyMol (https://www.pymol.org/). Therefore, focused docking simulations on the alpha carbon of Tyr293 and Trp102 were also performed. Subsequently, the files of ligands and protein employed to perform the docking analysis were obtained using the program Autodock tools 1.5.6 (ADT) (https://autodocksuite.scripps.edu/adt/). Autogrid4 and Autodock4 programs procured the grid maps and docking results. The PyMol program was also employed to explore the possible ligand-binding cavities in the LPA3 receptor.

Blind docking simulations consist of adjusting the grid box (search area) to the maximum expanse the program allows, i.e., 126 Å3 and a grid space of 0.375 Å3, centered by default on the proteins. All simulations were performed using the LINUX operating system (Centus) and the Lamarckian genetic algorithm, which was chosen to search for the best ligand configurations coupled to the LPA3 receptor. This process generates the most probable results of the ligand

conformations within the receptor search area, with a maximum number of 1×10^7 energy evaluations and a population of 100 randomized individuals (represented by the 100 best evaluations). For the focused docking, the grid box was 60 Å^3 with a grid space of 0.375 Å^3 centered on the alpha carbons of Tyr293 or Trp102, as mentioned previously. The 2D maps showing the receptor's residues interacting with the ligands were obtained using the Discovery Studio Visualizer (https://discover.3ds.com/discovery-studio-visualizer-download).

4.5 Reagents, plasmids, and cells.

1-Oleyl lysophosphatidic acid (LPA) and 2S-(1-oleoyl-2-O-methyl-glycerophosphothionate) (OMPT) were from Cayman Chemical Co. (Ann Arbor, MI, USA). Dulbecco's modified Eagle's medium, trypsin, Lipofectamine 2000, streptomycin, penicillin, amphotericin B, blasticidin, hygromycin B, doxycycline hyclate, and Fura-2 AM were purchased from Invitrogen-Life Technologies (Carlsbad, CA, USA). Our previous publications indicate the source of other materials [28, 32, 33]. Parental Flp-In T-Rex HEK293 cells were obtained from Invitrogen (Carlsbad, CA, USA). The LPA3 receptor sequence was fused at the carboxyl terminus (Ctail) with a green fluorescent protein and cloned into the pCDNA5/FRT/TO plasmid to employ the inducible Flp-In TREx expression system [28, 32, 33]. Mutants of this plasmid were obtained commercially (Bioinnovatise, Inc., Rockville, MD, USA), and the changes were confirmed by sequencing. Those amino acids identified to interact with the ligands in the docking simulations were substituted by alanine. Two main agonist-binding cavities were detected (Cavity 1 and 2, see Results). The residues replaced by alanine for the Upper Cavity mutant were (Thr25, Leu86, Thr90, Arg105, Gln106, Asp110, Leu113, Leu179, Trp191, Trp252, Gly255, Leu259, Lys275, Arg276, Phe278, and Leu279). For the Lower Cavity mutant, the amino acids substituted were (Ile131, Met134, Arg135, Val136, Val213, Lys216, Thr217, Val219, Leu220, Arg230, Pro234, Lys275, Arg276, and Phe278). Only the Lower Cavity's last three amino acids (i.e., Lys275, Arg276, and Phe278) were present in the Upper Cavity. A third mutant was also obtained in which all the residues detected in these binding pockets were substituted by alanine (Upper+Lower Cavity mutant). T-Rex HEK293 cells were transfected with plasmids for the expression of the wild-type and mutant receptors and were subjected to selection as described before [28, 32, 33].

4.6 Intracellular Calcium Concentration.

Determinations were performed as previously described [28, 32, 33]. In brief, the cells were serum-starved and treated for 12 h with 100 ng/mL doxycycline hyclate to induce LPA3 expression; all receptors (wild-type and mutants were tagged with the green fluorescent protein [28, 32, 33]; the density was similar as reflected by fluorescence microscopy). The cells were loaded with 2.5 μM Fura-2 AM for 1 h. Cells were carefully detached from the Petri dishes, washed to eliminate unincorporated dye, and maintained in suspension. Two excitation wavelengths (340 and 380 nm) and an emission wavelength of 510 nm were employed. Intracellular calcium levels were calculated as described by Grynkiewicz et al.[34].

4.7 Statistical Analyses.

The data are presented as the means ± standard error of the means. Statistical analyses were performed using ordinary one-way ANOVA with the Bonferroni post-test using the software

included in the GraphPad Prism program (version 10.2.3). A p-value < 0.05 was considered statistically significant.

Supplementary Materials: The following supporting information can be downloaded at: www.mdpi.com/xxx/s1, Figure S1: title; Table S1: title; Video S1: title.

Author Contributions: Conceptualization: K.H.S., J.C.-B., J.A.G.-S.; Methodology: K.H.S., J.C.-B.; Investigation: K.H.S., M.T.R.-A., R.R.-H., J.C.-B.; Visualization: K.H.S., M.T.R.-A., R.R.-H., J.C.-B.; Writing – original draft: K.H.S., J.C.-B., J.A.G.-S.; Writing – review & editing: K.H.S., M.T.R.-A., R.R.-H., J.C.-B., J.A.G.-S.

Funding: This research was partially supported by Grants from CONAHCYT (Fronteras 6676) and DGAPA (IN201221 and IN201924).

Data Availability Statement: The data are available from the corresponding author upon reasonable request.

Acknowledgments: We thank Dr. A. César Poot-Hernández for his suggestions to improve the manuscript. The advice and technical support of the following members of the indicated Service Units of our Institute is gratefully acknowledged: Dr. Héctor Malagón and Dr. Claudia Rivera (Bioterio); Juan Barbosa and Gerardo Coello (Cómputo); and Aurey Galván and Manuel Ortínez (Taller). K. Helivier Solís is a student of the Programa de Doctorado en Ciencias Biomédicas (UNAM) of Universidad Nacional Autónoma de México (account 520014983).

Conflicts of Interest: The authors declare that there is no conflict of interest regarding the publication of this article.

References

1. Choi, J. W.; Herr, D. R.; Noguchi, K.; Yung, Y. C.; Lee, C. W.; Mutoh, T.; Lin, M. E.; Teo, S. T.; Park, K. E.; Mosley, A. N.; Chun, J., LPA receptors: subtypes and biological actions. *Annu. Rev. Pharmacol. Toxicol.* **2010,** 50, 157-86. https://doi.org/10.1146/annurev.pharmtox.010909.105753

2. Yung, Y. C.; Stoddard, N. C.; Chun, J., LPA receptor signaling: pharmacology, physiology, and pathophysiology. *J. Lipid Res.* **2014,** 55, (7), 1192-1214. https://doi.org/10.1194/jlr.R046458

3. Fukushima, N.; Chun, J., The LPA receptors. *Prostaglandins Other Lipid Mediat.* **2001,** 64, (1-4), 21-32. https://doi.org/10.1016/s0090-6980(01)00105-8

4. Hama, K.; Aoki, J., LPA(3), a unique G protein-coupled receptor for lysophosphatidic acid. *Prog. Lipid Res.* **2010,** 49, (4), 335-42. https://doi.org/10.1016/j.plipres.2010.03.001

5. Solís, K. H.; Romero-Ávila, M. T.; Guzmá-Silva, A.; García-Sáinz, J. A., The LPA(3) Receptor: Regulation and Activation of Signaling Pathways. *Int J Mol Sci* **2021,** 22, (13), 6704. https://doi.org/10.3390/ijms22136704

6. Contos, J. J.; Chun, J., The mouse lp(A3)/Edg7 lysophosphatidic acid receptor gene: genomic structure, chromosomal localization, and expression pattern. *Gene* **2001,** 267, (2), 243-53. https://doi.org/10.1016/s0378-1119(01)00410-3

7. Fells, J. I.; Tsukahara, R.; Liu, J.; Tigyi, G.; Parrill, A. L., Structure-based drug design identifies novel LPA3 antagonists. *Bioorg. Med. Chem.* **2009,** 17, (21), 7457-64. https://doi.org/10.1016/j.bmc.2009.09.022

8. Bandoh, K.; Aoki, J.; Taira, A.; Tsujimoto, M.; Arai, H.; Inoue, K., Lysophosphatidic acid (LPA) receptors of the EDG family are differentially activated by LPA species. Structure-activity relationship of cloned LPA receptors. *FEBS Lett.* **2000,** 478, (1-2), 159-65. https://doi.org/10.1016/s0014-5793(00)01827-5

9. Urs, N. M.; Kowalczyk, A. P.; Radhakrishna, H., Different mechanisms regulate lysophosphatidic acid (LPA)-dependent versus phorbol ester-dependent internalization of the LPA1 receptor. *J. Biol. Chem.* **2008,** 283, (9), 5249-57. https://doi.org/10.1074/jbc.M710003200

10. Hauser, A. S.; Kooistra, A. J.; Munk, C.; Heydenreich, F. M.; Veprintsev, D. B.; Bouvier, M.; Babu, M. M.; Gloriam, D. E., GPCR activation mechanisms across classes and macro/microscales. *Nat. Struct. Mol. Biol.* **2021,** 28, (11), 879-888. https://doi.org/10.1038/s41594-021-00674-7

11. Flock, T.; Hauser, A. S.; Lund, N.; Gloriam, D. E.; Balaji, S.; Babu, M. M., Selectivity determinants of GPCR-G-protein binding. *Nature* **2017,** 545, (7654), 317-322. https://doi.org/10.1038/nature22070

12. Hilger, D.; Masureel, M.; Kobilka, B. K., Structure and dynamics of GPCR signaling complexes. *Nat. Struct. Mol. Biol.* **2018,** 25, (1), 4-12. https://doi.org/10.1038/s41594-017-0011-7

13. Zhou, Q.; Yang, D.; Wu, M.; Guo, Y.; Guo, W.; Zhong, L.; Cai, X.; Dai, A.; Jang, W.; Shakhnovich, E. I.; Liu, Z. J.; Stevens, R. C.; Lambert, N. A.; Babu, M. M.; Wang, M. W.; Zhao, S., Common activation mechanism of class A GPCRs. *Elife* **2019,** 8. https://doi.org/10.7554/eLife.50279

14. Arkin, M. R.; Randal, M.; DeLano, W. L.; Hyde, J.; Luong, T. N.; Oslob, J. D.; Raphael, D. R.; Taylor, L.; Wang, J.; McDowell, R. S.; Wells, J. A.; Braisted, A. C., Binding of small molecules to an adaptive protein-protein interface. *Proc. Natl. Acad. Sci. U. S. A.* **2003,** 100, (4), 1603-8. https://doi.org/10.1073/pnas.252756299

15. Stank, A.; Kokh, D. B.; Fuller, J. C.; Wade, R. C., Protein Binding Pocket Dynamics. *Acc Chem Res* **2016,** 49, (5), 809-15. https://doi.org/10.1021/acs.accounts.5b00516

16. Martínez-Morales, J. C.; Romero-Ávila, M. T.; Reyes-Cruz, G.; García-Sáinz, J. A., Roles of Receptor Phosphorylation and Rab Proteins in G Protein-Coupled Receptor Function and Trafficking. *Mol. Pharmacol.* **2022,** 101, (3), 144-153. https://doi.org/10.1124/molpharm.121.000429

17. Rosell, M.; Fernandez-Recio, J., Docking-based identification of small-molecule binding sites at protein-protein interfaces. *Comput Struct Biotechnol J* **2020,** 18, 3750-3761. https://doi.org/10.1016/j.csbj.2020.11.029

18. Fells, J. I.; Tsukahara, R.; Fujiwara, Y.; Liu, J.; Perygin, D. H.; Osborne, D. A.; Tigyi, G.; Parrill, A. L., Identification of non-lipid LPA3 antagonists by virtual screening. *Bioorg. Med. Chem.* **2008,** 16, (11), 6207-17. https://doi.org/10.1016/j.bmc.2008.04.035

19.   Fujiwara, Y.; Sardar, V.; Tokumura, A.; Baker, D.; Murakami-Murofushi, K.; Parrill, A.; Tigyi, G., Identification of residues responsible for ligand recognition and regioisomeric selectivity of lysophosphatidic acid receptors expressed in mammalian cells. *J. Biol. Chem.* **2005,** 280, (41), 35038-50. https://doi.org/10.1074/jbc.M504351200

20.   Valentine, W. J.; Fells, J. I.; Perygin, D. H.; Mujahid, S.; Yokoyama, K.; Fujiwara, Y.; Tsukahara, R.; Van Brocklyn, J. R.; Parrill, A. L.; Tigyi, G., Subtype-specific residues involved in ligand activation of the endothelial differentiation gene family lysophosphatidic acid receptors. *J. Biol. Chem.* **2008,** 283, (18), 12175-87. https://doi.org/10.1074/jbc.M708847200

21.   Balogh, B.; Pazmany, T.; Matyus, P., Analysis of Edg-Like LPA Receptor-Ligand Interactions. *Curr. Pharm. Des.* **2015,** 21, (24), 3533-47. https://doi.org/10.2174/1381612821666150216120500

22.   Ballesteros, J. A.; Weinstein, H., Integrated methods for the construction of three-dimensional models and computational probing of structure-function relations in G protein coupled receptors. *Methods Neurosci.* **1995,** 25, 366-428. https://doi.org/https://doi.org/10.1016/S1043-9471(05)80049-7

23.   Pandy-Szekeres, G.; Munk, C.; Tsonkov, T. M.; Mordalski, S.; Harpsoe, K.; Hauser, A. S.; Bojarski, A. J.; Gloriam, D. E., GPCRdb in 2018: adding GPCR structure models and ligands. *Nucleic Acids Res.* **2018,** 46, (D1), D440-D446. https://doi.org/10.1093/nar/gkx1109

24.   Hasegawa, Y.; Erickson, J. R.; Goddard, G. J.; Yu, S.; Liu, S.; Cheng, K. W.; Eder, A.; Bandoh, K.; Aoki, J.; Jarosz, R.; Schrier, A. D.; Lynch, K. R.; Mills, G. B.; Fang, X., Identification of a phosphothionate analogue of lysophosphatidic acid (LPA) as a selective agonist of the LPA3 receptor. *J. Biol. Chem.* **2003,** 278, (14), 11962-9. https://doi.org/10.1074/jbc.M209168200

25.   Qian, L.; Xu, Y.; Simper, T.; Jiang, G.; Aoki, J.; Umezu-Goto, M.; Arai, H.; Yu, S.; Mills, G. B.; Tsukahara, R.; Makarova, N.; Fujiwara, Y.; Tigyi, G.; Prestwich, G. D., Phosphorothioate analogues of alkyl lysophosphatidic acid as LPA3 receptor-selective agonists. *ChemMedChem* **2006,** 1, (3), 376-83. https://doi.org/10.1002/cmdc.200500042

26.   Kajitani, N.; Okada-Tsuchioka, M.; Inoue, A.; Miyano, K.; Masuda, T.; Boku, S.; Iwamoto, K.; Ohtsuki, S.; Uezono, Y.; Aoki, J.; Takebayashi, M., G protein-biased LPAR1 agonism of prototypic antidepressants: Implication in the identification of novel therapeutic target for depression. *Neuropsychopharmacology* **2024,** 49, (3), 561-572. https://doi.org/10.1038/s41386-023-01727-9

27.   Jiang, G.; Inoue, A.; Aoki, J.; Prestwich, G. D., Phosphorothioate analogs of sn-2 radyl lysophosphatidic acid (LPA): metabolically stabilized LPA receptor agonists. *Bioorg. Med. Chem. Lett.* **2013,** 23, (6), 1865-9. https://doi.org/10.1016/j.bmcl.2013.01.002

28.   Solís, K. H.; Romero-Ávila, M. T.; Rincón-Heredia, R.; Martínez-Morales, J. C.; García-Sáinz, J. A., LPA3: Pharmacodynamic Differences Between Lysophosphatidic Acid and Oleoyl-Methoxy Glycerophosphothionate: Biased

Agonism, Two Sites. *Receptors* **2024,** 3, (4), 555-573. https://doi.org/10.3390/receptors3040029

29. Malar, D. S.; Verma, K.; Prasanth, M. I.; Tencomnao, T.; Brimson, J. M., Network analysis-guided drug repurposing strategies targeting LPAR receptor in the interplay of COVID, Alzheimer's, and diabetes. *Sci Rep* **2024,** 14, (1), 4328. https://doi.org/10.1038/s41598-024-55013-9

30. Ghersi, D.; Sanchez, R., Improving accuracy and efficiency of blind protein-ligand docking by focusing on predicted binding sites. *Proteins* **2009,** 74, (2), 417-24. https://doi.org/10.1002/prot.22154

31. Liu, S.; Paknejad, N.; Zhu, L.; Kihara, Y.; Ray, M.; Chun, J.; Liu, W.; Hite, R. K.; Huang, X.-Y., Differential activation mechanisms of lipid GPCRs by lysophosphatidic acid and sphingosine 1-phosphate. *Nature Communications* **2022,** 13, (1). https://doi.org/10.1038/s41467-022-28417-2

32. Solís, K. H.; Romero-Ávila, M. T.; Rincón-Heredia, R.; García-Sáinz, J. A., LPA(3) Receptor Phosphorylation Sites: Roles in Signaling and Internalization. *Int J Mol Sci* **2024,** 25, (10), 5508. https://doi.org/10.3390/ijms25105508

33. Solís, K. H.; Romero-Ávila, M. T.; Rincón-Heredia, R.; García-Sáinz, J. A., Lysophosphatidic Acid Receptor 3 (LPA3): Signaling and Phosphorylation Sites. *Int J Mol Sci* **2024,** 25, (12). https://doi.org/10.3390/ijms25126491

34. Grynkiewicz, G.; Poenie, M.; Tsien, R. Y., A new generation of Ca2+ indicators with greatly improved fluorescence properties. *J. Biol. Chem.* **1985,** 260, (6), 3440-50. https://doi.org/10.1016/S0021-9258(19)83641-4

SUPPLEMENTARY MATERIAL

Supplementary Figure S1. Alignment of the FASTA sequence of the LPA₃ receptor (Q9UBY5) with the 3D model programs I-Tasser (model 01.pdb chainA_s001), Swiss Model (model1) and AlphaFold (AF-A v4).

Colored PROMALS3D alignment (sequences in aligned order)

```
Conservation:              99999999 999999999 99 9 9 99 999999 999999999 999999999 99 999 9999999
AF-A0A1W7RGA1-F1-model_v4  1 MNECHYDKRMDFFYNRSNADTADEWSGTTLVIVLCFGTFFCLFIFLSNSLVIAAVVKNKKFHYPFYYLLA  70
sp_Q9UBY5_LPAR3_HUMAN_Lys  1 MNECHYDKHMDFFYNRSNTDTVDDWTGTKLVIVLCVGTFFCLFIFFSNSLVIAAVIKNRKFHFPFYYLLA  70
model_01.pdb_chainA_s001   1 MNECHYDKHMDFFYNRSNTDTVDDWTGTKLVIVLCVGTFFCLFIFFSNSLVIAAVIKNRKFHFPFYYLLA  70
model1.pdb_chainA_s002     1 MNECHYDKHMDFFYNRSNTDTVDDWTGTKLVIVLCVGTFFCLFIFFSNSLVIAAVIKNRKFHFPFYYLLA  70
Consensus_aa:                MNECHYDK+MDFFYNRSNhDThD-WoGTpLVIVLChGTFFCLFIFhSNSLVIAAVlKN+KFH@PFYYLLA
Consensus_ss:                         eeee              hhhhhhhhhhhhhhhhhhhhhhhhhhh      hhhhhhh

Conservation:              9999999999999999999999999 99 999999999999 999999 9999999999 99   9999999
AF-A0A1W7RGA1-F1-model_v4 71 NLAAADFFAGIAYVFLMFNTGPVSRTLSVNRWFLRQGLLDTSLTASLANLLVIAVERHISIVKMRVHSNL 140
sp_Q9UBY5_LPAR3_HUMAN_Lys 71 NLAAADFFAGIAYVFLMFNTGPVSKTLTVNRWFLRQGLLDSSLTASLTNLLVIAVERHMSIMRMRVHSNL 140
model_01.pdb_chainA_s001  71 NLAAADFFAGIAYVFLMFNTGPVSKTLTVNRWFLRQGLLDSSLTASLTNLLVIAVERHMSIMRMRVHSNL 140
model1.pdb_chainA_s002    71 NLAAADFFAGIAYVFLMFNTGPVSKTLTVNRWFLRQGLLDSSLTASLTNLLVIAVERHMSIMRMRVHSNL 140
Consensus_aa:                NLAAADFFAGIAYVFLMFNTGPVS+TLoVNRWFLRQGLLDoSLTASLhNLLVIAVERHhSIh+MRVHSNL
Consensus_ss:                hhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhh

Conservation:              999999 9999 999999999999999999 99 9999999999999 9999999  9 999999 999
AF-A0A1W7RGA1-F1-model_v4 141 TKKRVTFLILLIWAIAIFMGAVPTLGWNCLCDISTCSSLAPIYSRSYLIFWTVSNLGVFFIMVVVYIRIY 210
sp_Q9UBY5_LPAR3_HUMAN_Lys 141 TKKRVTLLILLVWAIAIFMGAVPTLGWNCLCNISACSSLAPIYSRSYLVFWTVSNLMAFLIMVVVYLRIY 210
model_01.pdb_chainA_s001  141 TKKRVTLLILLVWAIAIFMGAVPTLGWNCLCNISACSSLAPIYSRSYLVFWTVSNLMAFLIMVVVYLRIY 210
model1.pdb_chainA_s002    141 TKKRVTLLILLVWAIAIFMGAVPTLGWNCLCNISACSSLAPIYSRSYLVFWTVSNLMAFLIMVVVYLRIY 210
Consensus_aa:                TKKRVThLILLlWAIAIFMGAVPTLGWNCLCsIShCSSLAPIYSRSYLlFWTVSNL.hFhIMVVVYlRIY
Consensus_ss:                    hhhhhhhhhhhhhhhhhhhh          eeeeee    hhheeehhhhhhhhhhhhhhhhhh

Conservation:               99 99999999999999999 9999999999 99999 99999999999999999  9 99 9999999
AF-A0A1W7RGA1-F1-model_v4 211 MYVQRKTNVLSPHTSGSISRRKTPMKLMKTVMAVLGAFIVCWTPGLVVLLLDGLNCTRCEVQNVKRWFLL 280
sp_Q9UBY5_LPAR3_HUMAN_Lys 211 VYVKRKTNVLSPHTSGSISRRRTPMKLMKTVMTVLGAFVVCWTPGLVVLLLDGLNCRQCGVQHVKRWFLL 280
model_01.pdb_chainA_s001  211 VYVKRKTNVLSPHTSGSISRRRTPMKLMKTVMTVLGAFVVCWTPGLVVLLLDGLNCRQCGVQHVKRWFLL 280
model1.pdb_chainA_s002    211 VYVKRKTNVLSPHTSGSISRRRTPMKLMKTVMTVLGAFVVCWTPGLVVLLLDGLNCRQCGVQHVKRWFLL 280
Consensus_aa:                hYVpRKTNVLSPHTSGSISRR+TPMKLMKTVMhVLGAFlVCWTPGLVVLLLDGLNCppC.VQpVKRWFLL
Consensus_ss:                hhhhhhhhhhhh      hhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhhh        hhhhhhhhh

Conservation:              999999  999999 99  9  99  9999 9      99   9999 99 99 9  999 9  999
AF-A0A1W7RGA1-F1-model_v4 281 LALLNSLMNPIIYSCKDDEMSNTMRRMICCSFEERSQERWSLRIPSAVLGRSIDSSSQYLENGISQVTIS 350
sp_Q9UBY5_LPAR3_HUMAN_Lys 281 LALLNSVVNPIIYSYKDEDMYGTMKKMICC-FSQENPERRPSRIPSTVLSRS-DTGSQYIEDSISQGAVC 348
model_01.pdb_chainA_s001  281 LALLNSVVNPIIYSYKDEDMYGTMKKMICC-FSQENPERRPSRIPSTVLSRS-DTGSQYIEDSISQGAVC 348
model1.pdb_chainA_s002    281 LALLNSVVNPIIYSYKDEDMYGTMKKMICC-FSQENPERRPSRIPSTVLSRS-DTGSQYIEDSISQGAVC 348
Consensus_aa:                LALLNSlhNPIIYShKD--M.sTM++MICC.Fppcs.ER.s.RIPShVLtRS.DotSQYlEstISQshlt
Consensus_ss:                hhhhhhhhhhhhhhh  hhhhhhhhhhhh                              eeeeee

Conservation:               9
AF-A0A1W7RGA1-F1-model_v4 351 GKENT  355
sp_Q9UBY5_LPAR3_HUMAN_Lys 349 NKSTS  353
model_01.pdb_chainA_s001  349 NKSTS  353
model1.pdb_chainA_s002    349 NKSTS  353
Consensus_aa:                sKpso
Consensus_ss:
```

Supplementary Figure S2. Images resulting from the LPA$_3$-LPA (blue) and LPA$_3$-OMPT (red) docking analysis using PyMol.

Supplementary Figure S3. Cartoon showing the LPA₃ receptor sites interacting with LPA (blue) and OMPT (red).

LPA₃ UNREFINED

Supplementary Figure S4. The 2D diagram shows interactions observed at the LPA₃ binding site (Lower Cavity) with LPA (Discovery Studio Visualizer); critical amino acids contributing to interactions are shown in circles.

Supplementary Figure S5. The 2D diagram shows interactions at the LPA₃ binding site (Upper Cavity) with OMPT (Discovery Studio Visualizer). Critical amino acids contributing to interactions are shown in circles.

Supplementary Figure S6. Cartoons showing the refined LPA₃ receptor sites interacting with LPA (blue) and OMPT (red). Charges are indicated in the Figures

CHARGE -2

LPA ▇

OMPT ▇

Supplementary Figure S7. The 2D diagram shows interactions at the LPA$_3$ binding site with LPA or OMPT. In the individual images Ligand and Charge are indicated (Discovery Studio Visualizer). Critical amino acids contributing to interactions are shown in circles.

LPA(-1)

![red]	Unfavorable Positive-Positive
![lightgreen]	Van Der Waals
![red]	Unfavorable Donor-Donor
![green]	Conventional Hydrogen Bond

![lightgreen]	Carbon Hydrogen Bond
![purple]	Alkyl
![purple]	Pi- Alkyl

LPA(-2)

Van Der Waals

Attractive Charge

Conventional Hydrogen Bond

Carbon Hydrogen Bond

Alkyl

OMPT(0)

▇ (light green)	Van Der Waals	
▇ (green)	Conventional Hydrogen Bond	
▇ (green)	Carbon Hydrogen Bond	
▇ (pink)	Alkyl	
▇ (pink)	Pi-Alkyl	

OMPT(-1)

Van Der Waals

Carbon Hydrogen Bond

Salt Bridge

Unfavorable Positive-Positive

Conventional Hydrogen Bond

Alkyl

OMPT(-2)

■	Van Der Waals	■	Carbon Hydrogen Bond
■	Salt Bridge	■	Unfavorable Positive-Positive
■	Attractive Charge	■	Alkyl
■	Conventional Hydrogen Bond	■	Pi-Alkyl

Supplementary Figure S8. Localization of the amino acids at the LPA$_3$ receptor to be focused. Trp102, orange; Tyr293, green.

Supplementary Figure S9. Cartoons showing the LPA₃ receptor (Trp102-focused) sites interacting with LPA (blue) and OMPT (red).Charges are indicated in the Figures

Trp102 CHARGE 0

Trp102 CHARGE -1

Trp102 CHARGE -2

Supplementary Figure S10. The 2D diagram shows interactions of the LPA$_3$ (Trp102) structure with LPA or OMPT. In the individual images Ligand and Charge are indicated (Discovery Studio Visualizer). Critical amino acids contributing to interactions are shown in circles.

Trp102
LPA(-1)

Van Der Waals
Conventional Hydrogen Bond
Unfavorable Positive-Positive

Alkyl
Pi-Alkyl

Trp102
LPA(-2)

🟩	Van Der Waals		🟩	Carbon Hydrogen Bond
🟨	Salt Bridge		🟪	Alkyl
🟧	Attractive Charge		🟪	Pi-Alkyl

Trp102

OMPT(0)

🟩 Van Der Waals	🟪 Alkyl
🟩 Carbon Hydrogen Bond	🟪 Pi-Alkyl
🟥 Unfavorable Positive-Positive	

Trp102
OMPT(-1)

🟩 Van Der Waals	🟥 Unfavorable Positive-Positive
🟧 Attractive Charge	🟪 Pi-Sigma
🟢 Conventional Hydrogen Bond	🟪 Alkyl
🟩 Carbon Hydrogen Bond	🟪 Pi-Alkyl

Trp102
OMPT(-2)

Van Der Waals
Salt Bridge
Attractive Charge

Carbon Hydrogen Bond
Alkyl
Pi-Alkyl

Supplementary Figure S11. Cartoons showing the LPA₃ receptor (Tyr293-focused) sites interacting with LPA (blue) and OMPT (red).Charges are indicated in the Figures

Tyr293 CHARGE 0

Tyr293 CHARGE -1

Tyr293 CHARGE -2

Supplementary Figure S12. The 2D diagram shows interactions of the LPA₃ (Tyr293) structure with LPA or OMPT. In the individual images Ligand and Charge are indicated (Discovery Studio Visualizer). Critical amino acids contributing to interactions are shown in circles.

Tyr293
LPA(-1)

Van Der Waals

Conventional Hydrogen Bond

Pi-Cation

Alkyl

Pi-Alkyl

Tyr293
LPA(-2)

Van Der Waals
Salt Bridge
Attractive charge
Carbon Hydrogen Bond
Pi-Alkyl
Alkyl

Tyr 293
OMPT(0)

Van Der Waals

Conventional Hydrogen Bond

Carbon Hydrogen Bond

Alkyl

Tyr293
OMPT(-1)

Van Der Waals

Attractive Charge

Conventional Hydrogen Bond

Unfavorable Positive-Positive
Alkyl
Pi-Alkyl

Tyr293
OMPT(-2)

	Van Der Waals			Pi-Sigma
	Salt Bridge			Alkyl
	Carbon Hydrogen Bond			Pi-Alkyl
	Unfavorable Positive-Positive			

Supplementary Figure S13. LPA$_3$ receptor cavities were evidenced using PyMol. In the left panel the cavities found are indicated in yellow. In the middle (Trp102-focused, Charge -1) and right (Trp102-focused, Charge -1) panel the position of the ligands (LPA, blue, and OMPT, red) are indicated. Change in ligand color is due to merge with the yellow color of the cavity.

Supplementary Table S1. Structural evaluation of the 3D models generated using the three most frequently used homology modeling servers (in green, the best values).

Server	MolProbity: Protein Geometry/ Ramachandran favored	SAVESv6.0: ERRAT	SAVESv6.0: VERIFY
SWISS-MODEL	91.45%	94.8052	33.14%
I-TASSER	78.92%	83.4302	38.81%
AlphaFold	95.18%	94.3038	29.01%

Supplementary Table S2. Blind docking for LPA and OMPT at the unrefined LPA3 receptor structure with a 126 Å3 grid box centered on the protein.

Ligand	ΔG (kcal/mol)	Kd (μM)	Sites of ligand-receptor interaction
LPA	-3.98	1200	Leu220, Thr217, Arg230, Lys216, Val213, Met134, Ile131, Arg135, Pro234, Asn139, Val136, His137
OMPT	-5.31	128	Leu86, Leu283, Leu279, Leu280, Lys275, Val33, Val36, Thr90, Trp25, Arg276, Trp277