Circuit function is more robust to changes in synaptic than intrinsic conductances.

Zachary Fournier [a)], Leandro M. Alonso [b)] and Eve Marder [c)]

Volen Center and Biology Department, Brandeis University, Waltham, MA, 02454,

USA.

(Dated: 30 August 2024)

Circuit function results from both intrinsic conductances of network neurons and the synaptic conductances that connect them. In models of neural circuits, different combinations of maximal conductances can give rise to similar activity. We compared the robustness of a neural circuit to changes in their intrinsic versus synaptic conductances. To address this, we performed a sensitivity analysis on a population of conductance-based models of the pyloric network from the crustacean stomatogastric ganglion (STG). The model network consists of three neurons with nine currents: a sodium current (Na), three potassium currents (Kd, KCa, A-type), two calcium currents (CaS and CaT), a hyperpolarization-activated current (H), a non-voltage-gated leak current (leak), and a neuromodulatory current (MI). The model cells are connected by seven synapses of two types, glutamatergic and cholinergic. We produced one hundred models of the pyloric network that displayed similar activities with values of maximal conductances distributed over wide ranges. We evaluated the robustness of each model to changes in their maximal conductances. We found that individual models have different sensitivities to changes in their maximal conductances, both in their intrinsic and synaptic conductances. As expected the models become less robust as the extent of the changes increase. Despite quantitative differences in their robustness, we found that in all cases, the model networks are more sensitive to the perturbation of their intrinsic conductances than their synaptic conductances.

[a)] Current Affiliation: Department of Biochemistry and Molecular Biology, The University of Chicago, Chicago, IL, 60637

[b)] Electronic mail: lalonso@brandeis.edu

7 Manuscript information

8   *Keywords:* neuronal oscillators, ion channels, inhibitory synapses, pyloric rhythm, central pattern

9   generators.

10   *Running title: network sensitivity*

11   *Funding acknowledgments: NIH R35 NS097343, R01 MH046742 (E.M.)*

c)Electronic mail: marder@brandeis.edu

I. Introduction

Since the early studies of Hopfield and Hebb (Hebb, 1949; Hopfield, 1982; Sejnowski and Tesauro, 1989) it has been often assumed that changes in synaptic strength provide the salient mechanism by which circuit dynamics are altered in learning (Siegelbaum and Kandel, 1991; Dayan and Abbott, 2001; Tsodyks et al., 1998; Abbott and Nelson, 2000; Sjöström et al., 2001). This tradition was reinforced by years of extracellular recordings of spikes in brain circuits, and by a myriad of computational studies using simplified model neurons such as rate models or integrate and fire neurons (Kempter et al., 1999; Fusi, 2002; Caporale and Dan, 2008; Fiebig and Lansner, 2017). While all of these studies have been extremely informative, they have led to an underestimate of the role of intrinsic membrane currents in network dynamics in larger networks.

At the same time, it is now patently clear that most, if not all, brain circuits display many voltage and time-dependent currents that give their neurons interesting and complex cellular dynamics (McCormick and Huguenard, 1992; Markram, 1997; Ranjan et al., 2020; Grienberger and Magee, 2022). Despite this, there are still prevailing beliefs that changes in synaptic strength that occur as a consequence of activity, represent the dominant mechanism for altering circuit dynamics. In contrast, those who have been studying the role of synaptic and intrinsic currents in circuit dynamics in small central pattern generating circuits have been acutely aware of the importance of modulating intrinsic currents (Calabrese and De Schutter, 1992; Swensen and Marder, 2000; Marder and Bucher, 2001; Harris-Warrick, 2011; Yang et al., 2022) and in some cases the relative insignificance of modifications of synaptic strength (Prinz et al., 2003b). Nonetheless, to the best of our knowledge, there have not been systematic sensitivity analyses in circuits with easily measurable outputs to compare the relative importance of modification of synaptic and intrinsic conductances for network performance.

In this study, we take advantage of the stereotyped motor patterns of the crustacean pyloric rhythm (Maynard, 1972; Selverston, 1976) to assess its stability to changes in its conductances. Because we did not want these results to be an idiosyncratic consequence

40 of the set of starting parameters, we generated 100 degenerate models (different sets of
41 parameters with similar outputs) to explore the generality of the results in this basic circuit,
42 and then evaluated the robustness of each of these models to changes in circuit parameters.

II. Results

A. Feature extraction to classify model network activity

The pyloric rhythm in crustaceans is produced by the sequential bursting of the PD
(pyloric dilator), LP (lateral pyloric), and PY (pyloric) neurons. The AB (anterior burster)
neuron is electrically coupled to the PD neurons and together they form a pacemaker kernel.
In these simulations the AB and PD neurons are represented by a single compartment cell,
which we call PD. We modified the model of the pyloric network in Prinz et al. (2004)
to include a neuromodulatory current, I_{MI}, in each cell (Golowasch and Marder, 1992b;
Swensen and Marder, 2000). Figure **1A** shows a schematic of the network studied in this
work. Each cell is modeled as a single compartment with multiple currents (Golowasch
and Marder, 1992a; Buchholtz et al., 1992; Goldman et al., 2001). The interactions
in the network are mediated by seven chemical synapses of two types: glutamatergic and
cholinergic (Prinz et al., 2004). Each neuron has a sodium current, I_{Na}; transient and slow
calcium currents, I_{CaT} and I_{CaS}; a transient potassium current, I_A; a calcium-dependent
potassium current, I_{KCa}; a delayed rectifier potassium current, I_{Kd}; a hyperpolarization-
activated inward current, I_H; a leak current, I_{leak}; and a neuromodulatory current I_{MI}.

The voltage traces in Figure **1B** are an example of normal triphasic pyloric activity
produced by one of the models in this study. To evaluate the activity of the network, we
performed a number of measurements on the voltage traces of each cell. The colored lines
correspond to the quantities used to calculate features of the network's activity. Duty cycles
are calculated as the burst duration divided by the bursting period, τ_b, of the cell. The phases
of the rhythm at which the cells begin and end their bursts, are measured using the start
of the PD burst as a reference (Figure **1B**). Because the network activity is approximately
periodic, it is possible to normalize the times of these events by the network period τ_n. In

this way, we computed a number between 0 and 1 that corresponds to the ON and OFF

phases for each cell. By computing these quantities for several periods of the activity, we

obtained a distribution of values of duty cycles, burst periods, and ON/OFF phases, for each

cell. The mean values and standard deviations of these distributions were used as *features*

for classification purposes. For example, a cell that produces irregular bursts will yield a

distribution of duty cycles with a standard deviation that would be larger than in the case

of a cell that bursts regularly in a normal triphasic rhythm.

The activity of the network depends on the values of the maximal conductances $\bar{g}_i$. In

this study we compare the sensitivity of the network to changes in intrinsic versus synaptic

conductances. Figure **1C** shows a normal (control) triphasic rhythm in black. The blue

trace in Figure **1C** corresponds to the activity of the *same* model when each of the intrinsic

conductances are changed by a random amount. In this case, altering the intrinsic con-

ductances results in a pattern of activity that is different from the control activity and is

not "pyloric" because the cells fail to produce bursts of action potentials. Figure **1D** shows

control activity in black, and activity when the synaptic conductances are changed in red.

In this case the pattern is similar to control activity, but there are some differences. Despite

only changing the synaptic conductances, the network's activity becomes slower and the

phases of the cycle at which cell fires are slightly different. Figure **1E** shows the control

values of the *intrinsic* conductances in black, and the changed values of the conductances

in blue used in the simulations in Figure **1C**. For simplicity, we only show how the $\bar{g}_{Na}$

and $\bar{g}_{Kd}$ were changed, but all intrinsic conductances were changed by a similar *percent*. In

the LP cell, the sodium $\bar{g}_{Na}$ conductance decreased while the delayed-rectifier potassium

$\bar{g}_{Kd}$ conductance increased, leading to a loss of spiking activity. The sodium $\bar{g}_{Na}$ maximal

conductance in the PY cell also decreased but $\bar{g}_{Kd}$ decreased even further, thus maintaining

spiking activity. These effects are due to the relative magnitude of $\bar{g}_{Na}$ and $\bar{g}_{Kd}$ in the cell,

as these are the two largest maximal conductances. Finally Figure **1F** shows the control

values of the synaptic maximal conductances in black and the perturbed values in red.

B. A model database of degenerate solutions

Conductance based models of neuronal activity are often employed to test and generate hypotheses about the functioning of individual neurons and/or circuits. One difficulty in working with these models is that tuning them to perform specific tasks can be challenging. Finding values of the parameters, such as the maximal conductances $\bar{g}_i$, so that a given circuit will display a specific set of behaviors has been done in the past by painstakingly hand-tuning the models' parameters (Traub et al., 1991; Nadim et al., 1995; Buchholtz et al., 1992). One alternative to hand-tuning consists of evaluating all possible combinations of parameter values within a specified domain, building a model database, and filtering the solutions of the models that satisfy a set of criteria (Prinz et al., 2003a, 2004; Crasto et al., 2007). Another alternative was introduced recently in Alonso and Marder (2019). For any given set of parameters, it is possible to simulate the model, and score the solutions depending on their features. The approach consists of defining a *cost function* that takes values of conductances as inputs, and returns a score. The goal then is to define a cost function such that low scores correspond to solutions of the model that display the target or desired behavior. Thus, *training* the model corresponds to optimizing —finding local minima —of the cost function. This approach was employed successfully to produce models of single neurons and pyloric networks that can operate robustly over a range of temperatures (Alonso and Marder, 2020). In this study we followed the approach in Alonso and Marder (2020). We modified the model in Prinz et al. (2004) to include a modulatory current I_{MI} and we also modified the *cost function* in Alonso and Marder (2020) to enforce a greater voltage separation between slow oscillations and spikes (see methods). In this work, optimization of the *cost function* yields values of conductances such that the circuit displays a pyloric rhythm as in Figure **1B**. We used a genetic algorithm to generate a database of $N = 100$ models with different values of maximal conductances (Goldberg and Holland, 1988). All models in the database display pyloric rhythms whose features —such as frequency and burst duration —are consistent with experimental measures performed in the pyloric circuit in the crustacean STG (Hamood et al., 2015).

The values of the conductances in our model database are widely distributed over many-
fold ranges. For example, Figure **2A** and **2B** shows the distributions of the PD A-current
conductances and the PD to PY glutamatergic synapse conductances. These distributions
span a broad range and are not normally distributed ($p < 0.05$ Shapiro-Wilk test (Shapiro
and Wilk, 1965)). In this study, each model is fully specified by its set of maximal con-
ductances $\bar{g}_i$, while all other parameters such as reversal potentials, activation functions,
membrane capacitance, were kept fixed and are the same across models. The total number
of conductances in a model is $N_g = 34$. To try to visualize the conductances in the database
we employed a popular technique (t-SNE) to visualize high-dimensional vectors. Figure **2C**
shows a t-SNE plot of the model conductances, where we reduced each models' conductance
set for visualization in a two-dimensional space (Hinton and Roweis, 2002). The clusters
in this plot indicate that there are groups of models with similar underlying conductance
relationships in the model database.

The models in the database were constrained to produce solutions whose features were
similar to the experimental measurements in the STG, but their activities are not identical.
For example, Figures **2D** and **2E** show the distributions of two features: the duty cycle of
the LP cell, and the phase of the cycle at which the LP cell begins to fire. Interestingly these
features are normally distributed ($p > 0.05$). Similarly to the values of the conductances,
the *features* we extracted from the activity also can be matched to a high dimensional
vector with a total number of features of $N_F = 11$. Figure **2F** shows a t-SNE plot of the
models' features. In this case it is not possible to identify clusters, suggesting that there
are no underlying relationships between the features in the model database. Even though
the conductances are broadly distributed over many-fold ranges, the output of the circuits
result in tight yet uncorrelated distributions.

C.   Model networks with similar behavior and different underlying currents.

Computational and experimental studies suggest that neurons can produce similar ac-
tivities employing different contributions of each ionic current. This observation is known
as *degeneracy* and one way to illustrate this was introduced recently in Alonso and Marder
(2019). A currentscape is a representation of neural dynamics that displays the percent con-

tribution of each ionic current to the total inward and outward currents over time. Figure **3**
shows the currentscapes for the PD, LP, and PY neurons in two different models. The voltage
traces of each network are plotted above the currentscapes. The total outward and inward
current of each cell are shown as black-filled curves above and below each currentscape plot
on a logarithmic scale. The dashed lines correspond to reference current values. The percent
contributions of each ionic current are represented in colors. The outward currents are on
top and inward currents on the bottom.

Figure **3** shows the currentscapes of each cell in two model networks. The currentscapes
show visible differences in each cell within a model as well as across the two different models,
yet the activity of the networks is similar. For example, there is a visibly larger contribution
of the I_{MI} current to the PD cell in Model B as compared to Model A. Likewise, the
contribution of the I_{Kd} current is more prominent in Model B's LP cell than Model A's
LP cell. In addition, there is little to no contribution of the KCa current in the Model
A PY cell, while contributions of this current are more visible in the Model B PY cell.
Furthermore, there is a small contribution of the I_{CaS} current in the Model A PD cell, but
a larger contribution in the Model B PD cell. Because the activity is driven by different
combinations of currents, the excitability properties of the cells are different, and this in
turn results in differential sensitivities to perturbations.

D.   Model networks are more sensitive to perturbation of intrinsic than synaptic
conductances.

We compared the robustness of intrinsic and synaptic conductances by modifying these
two sets separately. For this, we independently varied the values of each maximal conduc-
tance $\bar{g}_i$ by some random amount within a range (Giovannini et al., 2013; Zang and Marder,
2023). For each conductance, this variation range is defined as a *percent* deviation from its
starting or control value. The larger this percentage, the larger the change in maximal
conductances. The values of the modified conductances are obtained by sampling from a
uniform distribution over each range. When the range is small and there is a small percent
deviation from the control values, the expectation is that the activity will change very little
and stay pyloric in most cases. As the variation range is increased, corresponding to larger

percent deviations from control values, it is reasonable to expect that the activity will be
disrupted in a greater proportion of cases.

We explored changes to maximal conductances between 0 to 100 percent. The change
size or variation range was defined as $\delta \in [0, 1]$. For each conductance we perform the change
$\bar{g}_i \rightarrow \bar{g}_i * \gamma$ with $\gamma = 1 + \delta U[-1, 1]$, a random number drawn from a uniform distribution
and scaled by δ. In this way values of $\delta \approx 0$ correspond to small percent changes and the
limit case $\delta = 1$ allows for changes of 100% of the value of a given conductance. For each
value of δ we sampled $N = 1000$ sets of conductances. For each set of conductances we
simulated the model, extracted features, and classified the activity as *pyloric* or *not pyloric*
using a random forest classifier (Ho, 1995). We then computed the percentage of traces that
displayed pyloric activity, and evaluated how this percentage changes as the variation range
δ increases. This procedure was performed independently for both intrinsic and synaptic
conductances, allowing us to compare the sensitivity to changes of each set. We repeated
the procedure for all one hundred models in our database.

Figure **4** shows the result of this analysis on ten models. In all panels, the x-axis cor-
responds to the variation range δ. The y-axis indicates the fraction of traces that were
classified as pyloric activity. By definition, when the variation range is zero ($\delta = 0$), all
"perturbations" result in the same model, and therefore, the activity is the same as in
control. As expected, the fraction of traces that display pyloric activity decreases as the
variation range (δ) increases. This is true for both intrinsic (blue) and synaptic conductances
(red). The synaptic sensitivity curve is above the intrinsic sensitivity curve for all values
of the variation range δ. This means that the models are more resilient to changes in their
synaptic conductances.

To quantify the observation that the intrinsic conductances are more sensitive to per-
turbations than the synaptic conductances, we calculated the average sensitivity curves in
Figure **5A**. The plot shows the average fraction of pyloric traces across all models for all
values of the variation range δ. To further quantify these observations we fitted each sen-
sitivity curve with a sigmoid function (Figure **5B**), $S(\delta) = c \cdot \left(1 + \frac{1}{1+e^{-(\delta-b)/a}}\right)$, that best
approximates it. The parameters are a, which controls the width, and b, which determines

the sharpness. Parameter c controls the overall amplitude of the sigmoid and was kept fixed
at $c = 100$, since by definition it must be $S(0) = 100$.

The sigmoid functions approximate the sensitivity curves well, with an average correlation
coefficient of $r^2 = 0.98$, and an average error of approximately 1% for each value of δ. The
fits to the synaptic sensitivity curves were marginally better than in the intrinsic case as the
mean error for the synaptic was 0.55% ($r^2 = 0.98$), whereas it was 1.54% ($r^2 = 0.98$) for the
intrinsic.

To compare the intrinsic and synaptic sensitivity curves, we inspected the distributions of
the sigmoidal fit parameters (Figure **5C, 5D**). These sets of parameters were not normally
distributed ($p < 0.05$) except for the midpoint parameter in the intrinsic curves ($p = 0.13$).
To evaluate if two distributions were statistically different, we used Student's t-Test (Student,
1908). The midpoint parameter distributions are different, and there is little to no overlap
between them ($p < 0.001$). In addition, we found that the intrinsic conductances have much
smaller midpoint parameters, consistent with the observation that their sensitivity curves
decrease faster as the size of the conductance changes δ increases.

Although the distributions of midpoint parameters are visibly different, the width pa-
rameter distributions largely overlap. However, these distributions are significantly different
($p < 0.001$) because the red distribution is long-tailed. A smaller midpoint parameter and a
smaller width parameter both indicate that a curve decreases more quickly. This difference
in the midpoint parameters indicates that the synaptic conductances must be varied further
from their starting values to disrupt pyloric activity.

To further compare the intrinsic and synaptic sensitivity curves, we calculated the areas
under the curves (Figure **5E**). The areas under the synaptic curves are significantly larger
($p < 0.001$), consistent with the previous observation that the network is more robust to
perturbations in their synaptic conductances.

To characterize the variability in the sensitivity curves, we performed principal component
analysis (PCA). The first component captures 97.6% of the variance, which indicates that
the curves are similar and predominantly vary along a single dimension. The distribution of
the eigenvalues required to reconstruct the curves from the principal component are shown

in Figure **5F**. The eigenvalues for the intrinsic and synaptic curves result in visibly different 239 distributions ($p < 0.05$) with means of opposite signs. These eigenvalues are normally 240 distributed in the case of the intrinsic curves ($p = 0.82$), and not normally distributed in the 241 synaptic case ($p < 0.05$). The intrinsic and synaptic curves can be reasonably approximated 242 by the same sigmoid-looking curve, but flipping the sign in each case.

243 **E. Relative robustness of models visualized in conductance space**

244   These 3 neuron models require the specification of $N_i = 27$ intrinsic conductances and 245 $N_s = 7$ synaptic conductances. There are multiple combinations of $\bar{g}_i$ that result in a proper 246 pyloric rhythm. This means that there are regions in conductance space G for which the 247 activity is pyloric, G_P. The shape and properties of G_P have been explored and studied in 248 the past (Taylor et al., 2006). To gain some intuition about the properties of these regions 249 we exploited the trained classifier, and evaluated its output over 2D ranges of conductance 250 space.

251   Figure **6** shows 2D projections of the regions of conductance space for which the activity 252 is pyloric G_P, for two different models. The top rows show the projection of G_P over 253 pairs of intrinsic conductances, $\bar{g}_{Na}$ and $\bar{g}_{Kd}$. The blue regions correspond to values of 254 the conductances for which the activity is pyloric (G_P), and the white cross indicates the 255 "control" or initial values of the conductances for that model. The white box indicates the 256 maximum range of the perturbations used in the sensitivity analyses ($\delta = 1$).

257   The shape of G_P is intricate and complex. Despite this, some basic properties of G_P are 258 revealed. The top panels of Figure **6** show the projection of G_P over the same pair of intrinsic 259 conductances, across cell types. The blue regions across panels are different, indicating that 260 the same perturbation in different cell types may result in drastically different outputs: cell 261 identity matters. The white cross is closer to the edge in the case of the LP cell, meaning that 262 this particular model is more sensitive to perturbations in the LP cell than to perturbations 263 in the other cells.

264   The second row in Figure **6**, model A, shows the projection of G_P over pairs of *synaptic* 265 conductances. The red areas correspond to values of the conductances for which the activity 266 is pyloric (G_P), and the white cross indicates the starting or "control" values of this particular

model. As before the shape of these projections are intricate and complex. They also are different across pairs of conductances. Notice that with the exception of the cholinergic synapses (first panel), the initial value of the conductances (white cross) is farther away from the edge than in the intrinsic cases, consistent with the finding that the model is more robust to perturbations in its synaptic conductances. In the same vein, note that the proportion of area of the white box ($\delta = 1$) occupied by the red (synaptic) region is visibly larger than the proportion of area occupied by the blue (intrinsic) region.

Figure **6** (bottom) shows the same calculation in a different model (model B). As before, the blue regions and the red regions are different across cells and pairs of synapses. The shape of the projections of G_P are different across models. In the case of the intrinsic conductances the shapes are similar overall, but there are visible differences, such as the white cross in the LP cell being farther away from the edge as in model A. The projection of G_P on pairs of synaptic conductances is very different in this model. In this case, almost the totality of the area enclosed by the white boxes ($\delta = 1$) corresponds to G_P, suggesting that model B is more robust than model A to changes in synaptic conductances. This is consistent with the finding that the synaptic sensitivity curve is steeper in model A (Model #1 in Figure **4**) than in model B (Model #2 in Figure **4**). These results provide an intuition of the complexities of G_P. Not only are these regions hard-to-impossible to characterize in one circuit, but they are also different across circuits.

III.   Discussion

A major goal of much of neuroscience is to understand how experience alters the function of brain circuits, both in the context of development and learning. At the same time, we are aware that damage to the brain can produce altered brain structure. It is therefore important to understand the resilience of brain circuits to damage, and to understand the plasticity mechanisms that foster learning and recovery from damage. Thus, it is critical to understand which changes in circuit parameters are most influential in altering circuit function.

Despite the enormous body of work on the importance of alterations of synaptic strength in numerous preparations, there are also many well-documented cases of neurons that change their intrinsic excitability during development or as a function of neuromodulation. For example the Rohan-Beard cells first show a slow Ca^{2+}- dependent spike, before eventually acquiring a rapid Na^+-dependent action potential (Baccaglini and Spitzer, 1977) and ACh and biogenic amine neuromodulators directly alter the intrinsic properties of thalamic and other neurons (McCormick and Pape, 1990; McCormick et al., 1991; McCormick and Prince, 1986, 1987, 1988; McCormick and Wang, 1991). There are myriad other examples in which neuromodulators completely transform the dynamics of neuronal activity (Bargmann, 2012; Marder, 2012; Nusbaum and Blitz, 2012). A neuromodulator that changes a neuron's activity profile from silent, to tonic firing, to producing plateau potentials or oscillatory bursts of activity is likely to appreciably alter the activity of the circuits in which these neurons are found. Because altering an intrinsic current is often accompanied by changes in the time scales of neuronal activity, even modest changes in an intrinsic conductance can result in major alterations in the dynamics of a target neuron and the circuits in which it functions. In a sense, we biased the results of this study in manner that underestimates the influence of intrinsic activity by selecting for neurons that were within a restricted region of the degenerate range of solutions.

We expect that results similar to these will be found in many brain regions that show strong propensities for plateau and oscillatory properties, especially in response to neuro-modulation. It remains to be seen whether cortical or hippocampal regions, typically thought to be modified by changes in synaptic strength (Hasselmo et al., 1995), are less sensitive to modulation of intrinsic properties.

In this study we evaluated the role of changes in inhibitory synaptic strengths, as in the STG synaptic connections are inhibitory. Because increases in inhibitory synaptic con-ductances push the membrane potential closer to reversal potential, there can be a dimin-ishing effect of increasing the inhibitory synaptic strength, although the increase in the post-synaptic conductance still decreases the efficacy of all other currents, and changes in inhibitory synaptic currents can saturate (Prinz et al., 2003b).

₃₂₃ Although increasing the strength of excitatory synaptic currents also is somewhat di-
₃₂₄ minished as the membrane potential of neurons moves closer to the reversal potential of an
₃₂₅ EPSP, the effect of increasing the EPSP may be amplified as voltage dependent currents
₃₂₆ are activated. Therefore, it is possible that a circuit dominated by excitation may be more
₃₂₇ dominated by changes in synaptic than intrinsic currents. Nonetheless, a large body of work
₃₂₈ now argues that much cortical function depends on a balance of excitatory/inhibitory func-
₃₂₉ tion (Atallah and Scanziani, 2009; Isaacson and Scanziani, 2011; Lim and Goldman, 2013;
₃₃₀ Sadeh and Clopath, 2021), so a similar study such as this needs to be done in a modeled
₃₃₁ cortical or hippocampal circuit. That said, this study benefits greatly from a simple mea-
₃₃₂ sure of circuit performance that can be used to determine circuit robustness, and this is less
₃₃₃ obvious in many other circuits.

IV.   Methods

A.   The Model

⁣ₓ₃₃₆   We employed a model circuit inspired by the STG of crustaceans that was developed and studied previously by multiple authors (Golowasch et al., 1999; Prinz et al., 2004; Alonso and Marder, 2020). The model network is composed of three cells named PD, LP and PY, each modeled by a single compartment with nine currents: all cells have a sodium current, I_{Na}; a delayed rectifier potassium current, I_{Kd}; transient and slower calcium currents, I_{CaT} and I_{CaS}; a transient potassium current, I_A; a calcium-dependent potassium current, I_{KCa}; a hyperpolarization-activated inward current, I_H; a leak current, I_{leak}; and a neuromodulatory current I_{MI} (Golowasch and Marder, 1992b; Swensen and Marder, 2000). This model of the pyloric network is identical to previous ones, except for the inclusion of the neuromodulatory current I_{MI}.

⁣   The membrane potential of the cells evolves according to a Hodgkin-Huxley equation for a single compartment,

$$C\frac{dV}{dt} = I_e - \sum_{i=1}^{9} I_i. \tag{1}$$

The terms in the sum are the ionic currents $I_i = \bar{g}_i m_i^{p_i} h_i^{q_i}(V - E_i)$ and I_e is injected current. The maximal conductance of a channel is given by $\bar{g}_i$. The variables m_i and h_i correspond to the activation and inactivation processes. The integers p_i and q_i are the number of gates in each channel. The reversal potential of the ion associated with the $i-th$ current is given by E_i.

⁣   The reversal potential of the Na, K, H, leak, and I_{MI} currents were kept fixed at $E_{Na} = 30mV$, $E_K = -80mV$, $E_H = -20mV$, $E_{leak} = -50mV$, and $E_{IMI} = -10mV$. The reversal potential of the calcium currents E_{Ca} was updated using the Nernst equation assuming an extracellular calcium concentration of $3 \times 10^3 \mu M$. The kinetic equations describing the seven voltage-gated conductances were modeled as in Liu et al. (1998),

$$\tau_{m_i}(V)\frac{dm_i}{dt} = m_{\infty_i}(V) - m_i \tag{2}$$
$$\tau_{h_i}(V)\frac{dh_i}{dt} = h_{\infty_i}(V) - h_i.$$

358 The functions $\tau_{m_i}(V)$, $m_{\infty_i}(V)$, $\tau_{h_i}(V)$ and $h_{\infty_i}(V)$ are based on the experimental work of
359 Turrigiano et al. (1995). The activation functions of the K_{Ca} current require a measure of the
360 internal calcium concentration $[Ca^{+2}]$ (Liu et al., 1998). The dynamics of the intracellular
361 calcium concentration are given by,

$$\tau_{Ca}\frac{d[Ca^{+2}]}{dt} = -Ca_F(I_{CaT} + I_{CaS}) - [Ca^{+2}] + Ca_0. \tag{3}$$

Here $Ca_F = 0.94 \ \frac{\mu M}{nA}$ is a current-to-concentration factor and $Ca_0 = 0.05 \ \mu M$. The total
capacitance of the cell is $C = 1nF$.

The interactions in the network consist of 7 chemical synapses and are similar to Prinz
et al. (2004). The synaptic current is given by $I_s = \bar{g}_s s(V_{post} - E_s)$, where $\bar{g}_s$ is the synaptic
strength, V_{post} is the membrane potential of the postsynaptic neuron and E_s is the reversal
potential of the synapse. The activation of a synapse $s(t)$ is given by

$$\frac{ds}{dt} = \frac{s_\infty(V_{pre}) - s}{\tau_r + \tau_s} \tag{4}$$

368 with,

$$s_\infty(V_{pre}) = \frac{1}{1 + exp((V_{th} - V_{pre})/\Delta)}, \tag{5}$$

369 and

$$\tau_s = \frac{1 - s_\infty(V_{pre})}{k_-}. \tag{6}$$

370   These equations are identical to Prinz et al. (2004) except for the inclusion of a bound
371 for the timescale of activation $\tau_r = 20msec$ (Alonso and Marder, 2020).

B. Generating the model database.

We generated 100 different models of the pyloric network following the approach outlined in Alonso and Marder (2019, 2020), which we briefly review here. Each model is specified by a set of maximal conductances, while all other parameters (such as reversal potentials) are the same across models. Our approach consists of defining a *target* or *objective* function such that upon minimization, it will yield sets of conductances that result in normal or control pyloric activity.

The *target function* takes values of conductances as its input. The model is integrated for 20 seconds and the first 10 seconds are dropped to minimize transient activity. We then compute several quantities: the average ($<>$) burst frequency $< f_b >$, the average duty cycle $< dc >$, the number of crossings with a slow wave threshold $\#_{sw} = -50mV$, the number of bursts $\#_b$, and the average lags between bursts $< \Delta_{PD-LP} >$ and $< \Delta_{PD-PY} >$. In this study we also sought solutions that would display a large voltage separation between the slow wave and the spike troughs. This was done by defining a second *slow wave threshold* at $\#_{sw} = -45mV$ and checking that the number of crossings is the same as the number of bursts. Finally, to penalize unstable solutions we compute the standard deviation of the burst frequency and duty cycle distributions; a solution was penalized if $std(\{f_b\}) \geq < f_b > \times 0.1$ or $std(\{dc\}) \geq < dc > \times 0.2$. If a solution is not discarded we can use these quantities to score its similarity to the target, pyloric behavior. We define the *errors* as follows,

$$
\begin{aligned}
E_f &= \sum_{i=\text{cell}} (f_{tg} - < f_b >_i)^2 \\
E_{dc} &= \sum_{i=\text{cell}} (dc_{tg} - < dc >_i)^2 \\
E_{sw1} &= \sum_{i=\text{cell}} (\#_{sw} - \#_b)^2 \\
E_{sw2} &= \sum_{i=\text{cell}} (\#_{sw} - \#_b)^2 \\
E_{ph} &= (\Delta_{PD-LPtg} - \frac{< \Delta_{PD-LP} >}{\tau_b})^2 + (\Delta_{PD-PYtg} - \frac{< \Delta_{PD-PY} >}{\tau_b})^2 .
\end{aligned}
$$

(7)

(8)

E_f measures how far the bursting frequency of each cell is to a target frequency $f_{tg} = 3$,

E_{dc} accounts for the duty cycle, E_{sw} accounts for the difference between the number of bursts

and the number of crossings with the slow wave threshold at $t_{sw} = -50mV$ (if $\#_{sw} \neq \#_b$ then

E_{sw} is large), E_{sw2} is the same as E_{sw} but for a different threshold value ($t_{sw2} = -45mV$),

E_{ph} compares the lags between bursts (in units of the bursting period $\tau_b = \frac{1}{<f_b>}$) to a target

lag Δ_{tg}. These measures are discussed in more detail in Alonso and Marder (2019, 2020).

For each set of maximal conductances $\bar{\boldsymbol{g_i}}$ we can then define an objective function,

$$E(\boldsymbol{G}) = \alpha E_f + \beta E_{dc} + \gamma E_{sw} + + \gamma_2 E_{sw2} + \eta E_{ph}. \tag{9}$$

The coefficients $(\alpha, \beta, \gamma, \gamma_2, \eta)$ determine which features are more important. In this work

we used $\alpha = 10$, $\beta = 1000$, $\gamma = 1$, $\gamma_2 = 1$, $\eta = 10$. We found that using these weights results

in a *target function* that produces useful sets of conductances (that produce normal pyloric

triphasic activity). All penalties E_i were calculated using $T = 10$ secs. The simulations were

done using an *RK4* integration routine with $dt = 0.05$ msecs.

The *target control behavior* is defined as all cells bursting with 20% duty cycle for PD

($dc_{tg_{PD}} = 0.2$) and 25% for the LP and PY cells ($dc_{tg_{LP,PY}} = 0.25$). The lag between bursts

was targeted to be $\Delta_{PD-LPtg} = 0.5$ and $\Delta_{PD-PYtg} = 0.75$. The target burst frequency of

all cells was set to $f_{tg} = 1Hz$ (Bucher et al., 2006; Tang et al., 2010; Hamood et al., 2015).

Minimization of the *target function* successfully produces sets of maximal conductances $\boldsymbol{g_i}$

that produce normal or control triphasic rhythms.

We defined a search space of allowed values listed here: for each cell we searched for $\bar{g}_{Na} \in$

$[0, 10^2]$ ($[\mu S]$), $\bar{g}_{CaT} \in [0, 10^1]$, $\bar{g}_{CaS} \in [0, 10^1]$, $\bar{g}_A \in [0, 10^1]$, $\bar{g}_{KCa} \in [0, 10^1]$, $\bar{g}_{Kd} \in [0, 10^2]$,

$\bar{g}_H \in [0, 10^1]$, $\bar{g}_L \in [0, 1]$, $\bar{g}_{MI} \in [0, 10]$. All synaptic conductances were searched in the range

$\bar{g}_{syn} \in [0, 1]$ ($[\mu S]$). We discretized the search space by taking 10^3 equally spaced values for

each parameter. The objective function (9) was minimized using a custom genetic algorithm

(Holland, 1992) on a desktop computer with 32 cores. We ran the genetic algorithm using

1000 initial individuals taken at random in the search space for 300 generations. Finding a

successful population of models takes several hours. We repeated this process to produce

₄₁₇ 100 different models with values of conductances distributed over large ranges.

₄₁₈ **C.   Extracting features from model traces.**

₄₁₉   The models were simulated for 10 seconds, and the final 5 seconds of the simulation ₄₂₀ were utilized. We calculated spike times by finding the local maxima of the time series of ₄₂₁ the membrane potential of each cell. Some subthreshold depolarizations resulted in local ₄₂₂ maxima that were not spikes. To isolate the spikes from these subthreshold depolarizations, ₄₂₃ local maxima were filtered to only include spike times within $5ms$ of the local maxima of ₄₂₄ the derivative of the voltage $\frac{dV}{dt}$. This is because local maxima of the voltage derivative will ₄₂₅ typically be within $5ms$ of the local maxima of the voltage if the rise in voltage is very rapid, ₄₂₆ which is the case for spikes produced by rapid sodium activation. In addition, we computed ₄₂₇ the maximum value of voltage in the 5 seconds of simulation that we analyzed (V_{max}), and ₄₂₈ we only kept spikes that were less than $25mV$ below this value. To prevent spike times ₄₂₉ from being detected in non-firing cells with fast but small oscillations, we imposed that if ₄₃₀ $V_{max} < -25mV$, then the spike threshold was set to $-25mV$. Models with cells that fire ₄₃₁ spikes beneath $-25mV$ do exist but were not included in this work.

₄₃₂   To calculate the bursts' start and end times, we set a threshold to distinguish inter-burst ₄₃₃ intervals (IBIs) from inter-spike intervals (ISIs) ISI_{th}. In the case of a regularly bursting ₄₃₄ cell, the ISI distribution is bimodal, with one peak corresponding to the intraburst ISI, and ₄₃₅ the other peak corresponding to the IBIs. We calculated the ISI threshold by first detecting ₄₃₆ the peaks in the histograms of ISIs and then chose the value halfway between these peaks.

₄₃₇   In some cases the distribution of ISIs was not bimodal due to irregular bursting or tonic ₄₃₈ spiking. This made it difficult to determine an ISI threshold to correctly identify bursts. ₄₃₉ In cases where the threshold could not be estimated, or it was estimated to be larger than ₄₄₀ $300mS$ or smaller than $100mS$, we set $ISI_{th} = 100mS$. This allowed us to reliably calculate ₄₄₁ phases by counting tonic spikes in the triphasic rhythm as burst starts and burst ends while ₄₄₂ also detecting spikes with large ISIs at the end of a burst. Using this method led to errors in ₄₄₃ burst identification in some cases, and this is a source of error and a limitation of the study.

₄₄₄   The features extracted from each model include the mean and standard deviations of the ₄₄₅ distributions of duty cycles that each cell produces over the period of 5 seconds. We also

446 calculated the means of the distributions of the phases of the rhythm at which each cell
447 begins and ends a burst. We calculated the duty cycles for each cell as the duration of the
448 burst B_d divided by the bursting period (τ_b),

$$\mathrm{DC} = \frac{B_d}{\tau_b} \tag{10}$$

449 The phases were calculated as the difference between the cell's on/off times and the start of
450 burst in the PD cell PD_{on} (taken as a reference) divided by the period of the rhythm (τ_N),

$$\theta = \frac{cell_{on/off} - PD_{on}}{\tau_n} \tag{11}$$

451 We defined τ_n as the bursting period τ_b of the PD cell, thus when PD either did not fire or
452 fired irregularly phases were not calculated and these quantities were set to 0.

D. Classification of model traces.

The activity of each trace was evaluated by first building a vector containing features of
its activity. The features are: the means of the distributions of duty cycles for each cell,
the standard deviations of the distributions of duty cycles for each cell, and the means of
the distributions of phases for each cell. We attempted classification of the vectors using a
number of algorithms including random forest classifiers and gradient boosting machines (Ho,
1995; Friedman, 2001), and we chose to use random forest classifiers to avoid the possibility
that gradient boosting machines might over fit to the training data for our purposes (Natekin
and Knoll, 2013). A random forest classifier is a machine learning method used to classify
items based on a list of the item's features (Ho, 1995). The classifier we created consisted of
500 decision trees (max depth = 4), where each tree was fed a different subset of the features
for classification. A winner-takes-all vote between the trees chooses the output label of the
classifier. The classifier must first be trained with a set of data that was manually labeled,
with the help of experimentalists that study the pyloric network in the crustacean STG.
When the cells were bursting in a triphasic pattern in the order *PD-LP-PY*, with at least
two spikes per burst, we considered the activity to be pyloric (see Figure **1**). We trained the

classifier on a database of 1200 model solutions or traces featuring a wide variety of wave-
forms. Any other type of activity was considered 'not pyloric'. In some cases we excluded
rhythms that displayed pyloric characteristics but also exhibited irregular features, such as
abnormally large duty cycles, cells with low amplitude spikes, and inconsistent firing and/or
bursting patterns.

We found that the trained classifier generalized with $\geq 95\%$ accuracy for many, but not
all models. To verify that the classifier properly classified the models we chose, we manually
checked thousands of traces. Only models that were classified with this level of accuracy
after perturbation were used in this study. On rare occasions, the classifier misclassified
models that are clearly pyloric as not pyloric even though the list of features fed to the
classifier were calculated properly. The percentage of misclassifications was approximately
$2 - 3\%$, which is consistent with the accuracy of the classifier.

E.   Sigmoidal fits

To analyze the sensitivity curves quantitatively, we fit them with sigmoid functions. We
optimized for the sigmoidal midpoint and width parameters by minimizing an error function
using a brute force approach. The sigmoid function is given by,

$$S(\delta) = c \cdot \left(1 + \frac{1}{1 + e^{-(\delta - b)/a}} \right) \tag{12}$$

where the a parameter controls the width of the sigmoid, the b parameter determines the
midpoint of the sigmoid, and the c parameter corresponds to the maximum asymptote of
the sigmoid function. Variable δ represents the variation range parameter. Parameter c was
fixed at 100% because this is the maximum percentage of models that can be pyloric, and
by definition all curves start at 100% when the size of the perturbation is $\delta = 0$ (i.e. we
impose $S(\delta = 0) = 100$). To find the values of a and b that best approximate the sensitivity
curves we minimize an error function defined as follows,

$$E(a, b) = \sum_{i=0}^{N_\delta} \left((S(\delta_i, a, b) - y(\delta_i))^2 \right) \tag{13}$$

492   where $delta_i = [0, 0.1, ...1]$ are the values of δ for which we computed the sensitivity

493 curves ($N_\delta = 11$), and $y(\delta_i)$ are the values of a sensitivity curve for each value of δ_i. We

494 evaluated all possible combinations of the a and b parameters over a 100×100 2D grid of

495 possible pairs with (-0.5 to -0.01 for a, 0 to 2 for b), and kept the values of a and b that

496 minimized the error function E.

497 F. Conductance Ranges

There are regions in conductance space where the activity is a normal pyloric rhythm.

Visualizing and characterizing these regions is challenging because they are high dimensional

objects, and we are limited to visualizing their two-dimensional projections. We evaluated

the output of the classifier over 2-dimensional ranges in conductance space. For this we

chose a pair of conductances $(\bar{g}_x, \bar{g}_y)$ and picked 100 values between 0 and $\bar{g}_{max}$ for each

pair. We then evaluated network activity using the classifier on the 10^4 possible pairs of

$(\bar{g}_x, \bar{g}_y)$ over this range. The values of $(\bar{g}_x, \bar{g}_y)$ that produce pyloric activity were plotted in

color (blue, red) and the values of $(\bar{g}_x, \bar{g}_y)$ that result in non-pyloric activity were plotted

in grey. For the intrinsic conductances, we chose to visualize the $\bar{g}_{Na}$ and $\bar{g}_{Kd}$ maximal

conductances projection for each cell. For the synaptic conductances, we visualized three

pairs: the cholinergic inputs to LP and PY from PD, the glutamatergic inputs to LP from

PD and PY, and the glutamatergic inputs from PD to LP and LP to PD. We performed

this procedure on two different models. For *Model A*, $\bar{g}_{max}$ for $\bar{g}_{Na}$ and $\bar{g}_{Kd}$ was $250nS$. In

the case of the synaptic conductances $\bar{g}_{max}$ was set to $0.45nS$ on the x-axis and $1.75nS$ on

the y-axis. For *Model B*, $\bar{g}_{max}$ was $200nS$ for $\bar{g}_{Na}$ and $175nS$ for $\bar{g}_{Kd}$. $\bar{g}_{max}$ for the synaptic

conductances was $0.15nS$ on the x-axis and $1.75nS$ on the y-axis.

References

Abbott, L. F. and Nelson, S. B. (2000). Synaptic plasticity: taming the beast. *Nature neuroscience*, 3(11):1178–1183.

Alonso, L. M. and Marder, E. (2019). Visualization of currents in neural models with similar behavior and different conductance densities. *Elife*, 8:e42722.

Alonso, L. M. and Marder, E. (2020). Temperature compensation in a small rhythmic circuit. *Elife*, 9:e55470.

Atallah, B. V. and Scanziani, M. (2009). Instantaneous modulation of gamma oscillation frequency by balancing excitation with inhibition. *Neuron*, 62(4):566–577.

Baccaglini, P. I. and Spitzer, N. C. (1977). Developmental changes in the inward current of the action potential of rohon-beard neurones. *The Journal of physiology*, 271(1):93–117.

Bargmann, C. I. (2012). Beyond the connectome: how neuromodulators shape neural circuits. *Bioessays*, 34(6):458–465.

Bucher, D., Taylor, A. L., and Marder, E. (2006). Central pattern generating neurons simultaneously express fast and slow rhythmic activities in the stomatogastric ganglion. *Journal of neurophysiology*, 95(6):3617–3632.

Buchholtz, F., Golowasch, J., Epstein, I. R., and Marder, E. (1992). Mathematical model of an identified stomatogastric ganglion neuron. *Journal of Neurophysiology*, 67(2):332–340.

Calabrese, R. L. and De Schutter, E. (1992). Motor-pattern-generating networks in invertebrates: modeling our way toward understanding. *Trends in neurosciences*, 15(11):439–445.

Caporale, N. and Dan, Y. (2008). Spike timing–dependent plasticity: a hebbian learning rule. *Annu. Rev. Neurosci.*, 31(1):25–46.

Crasto, C. J., Koslow, S. H., and Prinz, A. A. (2007). Computational exploration of neuron and neural network models in neurobiology. *Neuroinformatics*, pages 167–179.

Dayan, P. and Abbott, L. F. (2001). *Theoretical Neuroscience*. Cambridge, MA: MIT Press.

Fiebig, F. and Lansner, A. (2017). A spiking working memory model based on hebbian short-term potentiation. *Journal of Neuroscience*, 37(1):83–96.

Friedman, J. H. (2001). Greedy function approximation: a gradient boosting machine. *Annals of statistics*, pages 1189–1232.

Fusi, S. (2002). Hebbian spike-driven synaptic plasticity for learning patterns of mean firing rates. *Biological cybernetics*, 87(5):459–470.

Giovannini, A., Severi, S., and Sobie, E. (2013). Parametric sensitivity analysis of the most recent computational models of rabbit cardiac pacemaking. *Biophysical Journal*, 104(2):296a.

Goldberg, D. E. and Holland, J. H. (1988). Genetic algorithms and machine learning. *Machine Learning*, 3(2):95–99.

Goldman, M. S., Golowasch, J., Marder, E., and Abbott, L. (2001). Global structure, robustness, and modulation of neuronal models. *Journal of Neuroscience*, 21(14):5229–5238.

Golowasch, J., Casey, M., Abbott, L., and Marder, E. (1999). Network stability from activity-dependent regulation of neuronal conductances. *Neural computation*, 11(5):1079–1096.

Golowasch, J. and Marder, E. (1992a). Ionic currents of the lateral pyloric neuron of the stomatogastric ganglion of the crab. *Journal of Neurophysiology*, 67(2):318–331.

Golowasch, J. and Marder, E. (1992b). Proctolin activates an inward current whose voltage dependence is modified by extracellular ca^{2+}. *Journal of Neuroscience*, 12(3):810–817.

Grienberger, C. and Magee, J. C. (2022). Entorhinal cortex directs learning-related changes in CA1 representations. *Nature*, 611(7936):554–562.

Hamood, A. W., Haddad, S. A., Otopalik, A. G., Rosenbaum, P., and Marder, E. (2015). Quantitative reevaluation of the effects of short-and long-term removal of descending modulatory inputs on the pyloric rhythm of the crab, cancer borealis. *Eneuro*, 2(1).

Harris-Warrick, R. M. (2011). Neuromodulation and flexibility in central pattern generator networks. *Current opinion in neurobiology*, 21(5):685–692.

Hasselmo, M. E., Schnell, E., and Barkai, E. (1995). Dynamics of learning and recall at excitatory recurrent synapses and cholinergic modulation in rat hippocampal region CA3. *Journal of Neuroscience*, 15(7):5249–5262.

Hebb, D. (1949). *The Organization of Behavior.* Wiley. New York.

Hinton, G. E. and Roweis, S. (2002). Stochastic neighbor embedding. *Advances in Neural Information*
Processing Systems, 15.

Ho, T. K. (1995). Random decision forests. *Proceedings of 3rd International Conference on Document*
Analysis and Recognition, 1:278–282 vol.1.

Holland, J. H. (1992). Genetic algorithms. *Scientific American*, 267(1):66–73.

Hopfield, J. J. (1982). Neural networks and physical systems with emergent collective computational abilities.
Proceedings of the national academy of sciences, 79(8):2554–2558.

Isaacson, J. S. and Scanziani, M. (2011). How inhibition shapes cortical activity. *Neuron*, 72(2):231–243.

Kempter, R., Gerstner, W., and Van Hemmen, J. L. (1999). Hebbian learning and spiking neurons. *Physical*
Review E, 59(4):4498.

Lim, S. and Goldman, M. S. (2013). Balanced cortical microcircuitry for maintaining information in working
memory. *Nature neuroscience*, 16(9):1306–1314.

Liu, Z., Golowasch, J., Marder, E., and Abbott, L. (1998). A model neuron with activity-dependent con-
ductances regulated by multiple calcium sensors. *Journal of Neuroscience*, 18(7):2309–2320.

Marder, E. (2012). Neuromodulation of neuronal circuits: back to the future. *Neuron*, 76(1):1–11.

Marder, E. and Bucher, D. (2001). Central pattern generators and the control of rhythmic movements.
Current biology, 11(23):R986–R996.

Markram, H. (1997). A network of tufted layer 5 pyramidal neurons. *Cerebral cortex (New York, NY: 1991)*,
7(6):523–533.

Maynard, D. M. (1972). Simpler networks. *Annals of the New York Academy of Sciences*, 193(1):59–72.

McCormick, D., Pape, H.-C., and Williamson, A. (1991). Actions of norepinephrine in the cerebral cortex
and thalamus: implications for function of the central noradrenergic system. *Progress in brain research*,
88:293–305.

McCormick, D. and Wang, Z. (1991). Serotonin and noradrenaline excite gabaergic neurones of the guinea-
pig and cat nucleus reticularis thalami. *The Journal of physiology*, 442(1):235–255.

McCormick, D. A. and Huguenard, J. R. (1992). A model of the electrophysiological properties of thalam-
ocortical relay neurons. *Journal of neurophysiology*, 68(4):1384–1400.

McCormick, D. A. and Pape, H.-C. (1990). Noradrenergic and serotonergic modulation of a
hyperpolarization-activated cation current in thalamic relay neurones. *The Journal of physiology*,
431(1):319–342.

McCormick, D. A. and Prince, D. A. (1986). Acetylcholine induces burst firing in thalamic reticular neurones
by activating a potassium conductance. *Nature*, 319(6052):402–405.

McCormick, D. A. and Prince, D. A. (1987). Actions of acetylcholine in the guinea-pig and cat medial and
lateral geniculate nuclei, in vitro. *The Journal of physiology*, 392(1):147–165.

McCormick, D. A. and Prince, D. A. (1988). Noradrenergic modulation of firing pattern in guinea pig and
cat thalamic neurons, in vitro. *Journal of neurophysiology*, 59(3):978–996.

Nadim, F., Olsen, Ø. H., De Schutter, E., and Calabrese, R. L. (1995). Modeling the leech heartbeat ele-
mental oscillator i. interactions of intrinsic and synaptic currents. *Journal of computational neuroscience*,
2:215–235.

Natekin, A. and Knoll, A. (2013). Gradient boosting machines, a tutorial. *Frontiers in neurorobotics*, 7:21.

Nusbaum, M. P. and Blitz, D. M. (2012). Neuropeptide modulation of microcircuits. *Current opinion in*
neurobiology, 22(4):592–601.

Prinz, A. A., Billimoria, C. P., and Marder, E. (2003a). Alternative to hand-tuning conductance-based
models: construction and analysis of databases of model neurons. *Journal of Neurophysiology*, 90(6):3998–
4015.

Prinz, A. A., Bucher, D., and Marder, E. (2004). Similar network activity from disparate circuit parameters.
Nature Neuroscience, 7(12):1345.

Prinz, A. A., Thirumalai, V., and Marder, E. (2003b). The functional consequences of changes in the
strength and duration of synaptic inputs to oscillatory neurons. *Journal of Neuroscience*, 23(3):943–954.

Ranjan, R., Logette, E., Marani, M., Herzog, M., Tache, V., Scantamburlo, E., Buchillier, V., and Markram,
H. (2020). A kinetic map of the homomeric voltage-gated potassium channel (Kv) family. *Biophysical*
Journal, 118(3):108a.

Sadeh, S. and Clopath, C. (2021). Inhibitory stabilization and cortical computation. *Nature Reviews*
Neuroscience, 22(1):21–37.

Sejnowski, T. J. and Tesauro, G. (1989). The hebb rule for synaptic plasticity: algorithms and implementations. In *Neural models of plasticity*, pages 94–103. Elsevier.

Selverston, A. I. (1976). Neuronal mechanisms for rhythmic motor pattern generation in a simple system. *Neural control of locomotion*, pages 377–399.

Shapiro, S. S. and Wilk, M. B. (1965). An analysis of variance test for normality (complete samples). *Biometrika*, 52(3-4):591–611.

Siegelbaum, S. A. and Kandel, E. R. (1991). Learning-related synaptic plasticity: LTP and LTD. *Current opinion in neurobiology*, 1(1):113–120.

Sjöström, P. J., Turrigiano, G. G., and Nelson, S. B. (2001). Rate, timing, and cooperativity jointly determine cortical synaptic plasticity. *Neuron*, 32(6):1149–1164.

Student (1908). The probable error of a mean. *Biometrika*, pages 1–25.

Swensen, A. M. and Marder, E. (2000). Multiple peptides converge to activate the same voltage-dependent current in a central pattern-generating circuit. *Journal of Neuroscience*, 20(18):6752–6759.

Tang, L. S., Goeritz, M. L., Caplan, J. S., Taylor, A. L., Fisek, M., and Marder, E. (2010). Precise temperature compensation of phase in a rhythmic motor pattern. *PLoS biology*, 8(8):e1000469.

Taylor, A. L., Hickey, T. J., Prinz, A. A., and Marder, E. (2006). Structure and visualization of high-dimensional conductance spaces. *Journal of neurophysiology*, 96(2):891–905.

Traub, R. D., Wong, R. K., Miles, R., and Michelson, H. (1991). A model of a ca3 hippocampal pyramidal neuron incorporating voltage-clamp data on intrinsic conductances. *Journal of neurophysiology*, 66(2):635–650.

Tsodyks, M., Pawelzik, K., and Markram, H. (1998). Neural networks with dynamic synapses. *Neural computation*, 10(4):821–835.

Turrigiano, G., LeMasson, G., and Marder, E. (1995). Selective regulation of current densities underlies spontaneous changes in the activity of cultured neurons. *Journal of Neuroscience*, 15(5):3640–3652.

Yang, J., Shakil, H., Ratté, S., and Prescott, S. A. (2022). Minimal requirements for a neuron to coregulate many properties and the implications for ion channel correlations and robustness. *Elife*, 11:e72875.

Zang, Y. and Marder, E. (2023). Neuronal morphology enhances robustness to perturbations of channel densities. *Proceedings of the National Academy of Sciences*, 120(8):e2219049120.

FIG. 1. **Feature extraction to classify model network activity (A)** Schematic of the model used in Prinz et al. (2004). The three cells are connected by seven inhibitory chemical synapses of two types: the red curves are cholinergic synapses and the grey curves are glutamatergic synapses. **(B)** Pyloric network activity. Each cell displays periodic bursting activity and the cells fire in a sequence *PD-LP-PY*. The color lines indicate the measures used for calculating duty cycle and firing phases. **(C)** Membrane voltages of control network activity (in black), overlaid with the network activity that results from changing the intrinsic conductances (blue). **(D)** Membrane voltages of control network activity (in black) overlaid with network activity that results from changing the synaptic conductances (in red). **(E)** Changes in intrinsic maximal conductance values. The black dots show the unperturbed or control conductances and the blue dots correspond to the changed conductances that produced the blue voltage traces in Figure **1C**. Only 2 maximal conductances per cell are shown for simplicity, but all $9 \times 3 = 27$ intrinsic conductances were changed. **(F)** Synaptic maximal conductance values for the control network (in black) and the perturbed network (in red) in Figure **1D**.

FIG. 2. **A model database of degenerate solutions. (A, B)** Distributions of gA maximal conductance in PD cells and $PD-PY_{glut}$ glutamatergic synapse maximal conductance ($N=100$). **(C)** Dimensionality reduction (t-SNE) and visualization of models' maximal conductances. **(D, E)** Distributions of features: LP duty cycle and LP-ON phase. **(F)** Dimensionality reduction (t-SNE) and visualization of models' conductances models' features

FIG. 3. **Model networks produce similar behavior with different underlying currents.** Currentscapes for two different models. The voltage traces for both networks are plotted at the top. The filled curves on top and bottom of the currentscapes for each cell type show the total inward (outward) current over time in logarithmic scale. The colored filled curves indicate the percent contribution of each current over time.

FIG. 4. **Model networks are differentially sensitive to modification of intrinsic and synaptic conductances.** Curves representing the decrease in the percentage of pyloric models as a function of the variation range. Curves from intrinsic conductance perturbations are shown in blue, and synaptic in red. Ten sets of curves are shown for ten models with different combinations of maximal conductances.

FIG. 5. **Model networks are more sensitive to changes in intrinsic conductances (A)** Average sensitivity curves across models. The plots show the average percentage of pyloric models for each value of δ (intrinsic curves show in blue, synaptic in red). **(B)** Example sigmoidal fits for one set of intrinsic and synaptic sensitivity curves. **(C)** Distributions of midpoint parameters for all fits of the sensitivity curves (intrinsic in blue, synaptic in red, 20 bins). **(D)** Distributions of width parameters for all fits of the sensitivity curves (intrinsic in blue, synaptic in red, 20 bins). **(E)** Distributions of the areas under the curves (intrinsic in blue, synaptic in red, 20 bins). **(F)** Distribution of eigenvalues for the first PCA component (97.6%) (intrinsic in blue, synaptic in red, 20 bins).

FIG. 6. **Relative robustness of models visualized in conductance space.** Ranges of conductance space where the activity of the network is pyloric, for two different models (model #1 and model #2). All panels show the output of the classifier over 2D ranges of conductance space. The top panels show pairs of intrinsic maximal conductances ($\bar{g}_{Na}$ and $\bar{g}_{Kd}$ of each cell). The blue dots correspond to values of the conductances that produce pyloric activity, while the gray dots indicate values were the network activity is not pyloric. The bottom panels show pairs of synaptic maximal conductances for which the activity is pyloric in red. The white cross indicates the values of the conductances for the models. The white boxes indicate the extent of the perturbations when they are allowed to be as large as a 100 deviation from their control values ($\delta = 1$).