Gradients of Recognition Molecules Shape Synaptic Specificity of a Visuomotor Transformation

Mark Dombrovski[1], Yixin Zang[2#], Giovanni Frighetto[3#], Andrea Vaccari[4#], Hyojong Jang[5#], Parmis S. Mirshahidi[1#], Fangming Xie[1], Piero Sanfilippo[1], Bryce W. Hina[5], Aadil Rehan[1], Roni H. Hussein[1], Pegah S. Mirshahidi[1], Catherine Lee[1], Mark A. Frye[3], Catherine R. von Reyn[5], Yerbol Z. Kurmangaliyev[6], Gwyneth M. Card[2*] & S. Lawrence Zipursky[1*]

[1]Department of Biological Chemistry, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, USA
[2]Department of Neuroscience, Howard Hughes Medical Institute, The Mortimer B. Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY, USA
[3]Department of Integrative Biology and Physiology, University of California, Los Angeles, Los Angeles, CA, USA
[4]Department of Computer Science, Middlebury College, Middlebury, VT, USA
[5]School of Biomedical Engineering, Science and Health Systems, Drexel University, Philadelphia, PA, USA
[6]Department of Biology, Brandeis University, Waltham, MA, USA

[#]Equal contributions

*Correspondence: S. Lawrence Zipursky (lzipursky@mednet.ucla.edu), Gwyneth M. Card (gc3017@columbia.edu),

Abstract

47
48
Converting sensory information into motor commands is fundamental to
most of our actions[1,2]. In *Drosophila*, visuomotor transformations are mediated by
Visual Projection Neurons (VPNs)[3,4]. These neurons convert object location and
motion into directional behaviors downstream through a synaptic gradient
mechanism[5]. However, the molecular origins of such graded connectivity remain
unknown. We addressed this question in a VPN cell type called LPLC2[6], which
integrates looming motion and transforms it into an escape response through two
parallel dorsoventral synaptic gradients at its inputs and outputs. We identified two
corresponding dorsoventral expression gradients of cell recognition molecules
within the LPLC2 population that regulate this synaptic connectivity. Dpr13
determines synaptic outputs of LPLC2 axons by interacting with its binding partner
DIP-ε expressed in the Giant Fiber, a neuron that mediates escape[7]. Similarly,
beat-VI regulates synaptic inputs onto LPLC2 dendrites by interacting with Side-II
expressed in upstream motion-detecting neurons. Behavioral, physiological, and
molecular experiments demonstrate that these coordinated molecular gradients
control differential synaptic connectivity, enabling the accurate transformation of
visual features into motor commands. As within-neuronal-type continuous variation
in gene expression is also observed in the mammalian brain[8], graded expression
of cell recognition molecules may represent a common mechanism underlying
synaptic specificity.

Main

Every day, we engage in a multitude of visually guided behaviors that rely on the conversion of an object's location in eye coordinates into directional body movements[9,10]. The brain regions and neural circuits responsible for these visuomotor transformations have been characterized in both vertebrates[11–13] and invertebrates[14,15]. These studies have revealed that learned visuomotor tasks[16–18] and innate visuomotor responses[19,20] rely on precise neuronal connectivity, which is established during development, or through experience-dependent circuit plasticity.

In the fly brain, visuomotor transformation occurs between Visual Projection Neurons (VPNs) and Descending Neurons[21] (DNs, Fig. 1a). Although VPNs encompass a broad range of neurons that connect the optic lobe to the central brain[22], here we focus on well-characterized VPN sub-classes of columnar neurons[4], including LC (lobula columnar) and LPLC (lobula plate/lobula columnar). There are about 30 VPNs of these types, each comprising 20 to 200 neurons per brain hemisphere[3,4]. For simplicity, we will refer to these collectively as VPNs. The dendrites of individual neurons within a VPN type each span 20 to 40 degrees of visual space, and as a population, they uniformly cover the optic lobe (Fig. 1a), creating a feature-detecting retinotopic map[23–25]. The axon terminals of each VPN type bundle together in the central brain, forming optic glomeruli, where some elaborate synapses onto the dendrites of DNs. For most of the VPN types, axonal retinotopy is lost[4,5]. We recently demonstrated that the transformation from visual input to motor output in VPN-DN circuits relies on a synaptic gradient mechanism, which functions independently of axonal retinotopy[5].

We define a synaptic gradient as a connectivity pattern in which the number of synapses connecting individual neurons within a presynaptic population (e.g., a VPN type) to postsynaptic targets (e.g., DNs), exhibits a retinotopically biased systematic variation. For example, we found that within a population of looming-responsive VPNs, those with anterior receptive fields make more synapses onto DNs that drive backward movements, while neurons with posterior receptive fields make more synapses onto DNs that drive forward movements and takeoff[5]. This anterior-posterior gradient of synapses enables the fly to detect looming threats and execute a rapid escape response[5]. Thus, the synaptic gradient mechanism provides a non-spatial, abstract representation of visual space, serving as an example of retinotopically guided, yet spatially independent, within-cell-type synaptic specificity. While cell recognition molecules have been described that regulate synaptic specificity between different neuronal types[26–28], how neurons of the same type form varying numbers of synapses with their targets remains unclear.

To uncover the molecular basis of synaptic gradients, we focused on a specific type of VPN called lobula plate/lobula columnar 2 (LPLC2), a population of about 100 neurons per hemibrain, that each respond selectively to dark, radially-expanding motion stimuli centered on their receptive field, and, therefore act as local looming detectors[6,23]. The axons of LPLC2 neurons convey visual information to the Giant Fiber (GF), a DN that elicits a rapid escape takeoff in response to looming threats[7,29,30] (Fig. 1b). In our previous work, we analyzed electron

microscopy (EM)-based connectomic reconstructions from two brain volumes
(Janelia "hemibrain[31]" and "FAFB-Flywire[32,33]") and showed that LPLC2 neurons
elaborate a dorsoventral (DV) gradient of synapses onto the GF. Specifically,
LPLC2 neurons with dendrites sampling the dorsal part of the visual field form
more synapses with the GF than their ventral counterparts (Fig. 1c-d).

Here, we investigated the molecular basis of the LPLC2-GF synaptic
gradient by combining single cell RNA sequencing (scRNA-seq), spatial
transcriptomics and genetics with morphological, behavioral, and physiological
studies. We identified two cell recognition molecules expressed in a gradient
across the LPLC2 population. One of them, dpr13, regulates connections of LPLC2
axons and the GF dendrites. We also discovered a second recognition molecule,
beat-VI, which controls the formation of a synaptic gradient onto the dendrites of
LPLC2 neurons. This shows that varying levels of cell recognition molecules within
a defined neuronal type can specify different numbers of synapses both pre- and
post-synaptically. Our findings represent a new mechanism of regulating synaptic
specificity.
155

156 LPLC2-Giant Fiber Synaptic Gradient Shapes Escape Behavior

To determine whether the DV synaptic gradient between LPLC2 and the GF
is functional, we quantified the escape response to looming stimuli from different
parts of the visual field. Previous work showed that flies exhibit two modes of
takeoff – a short mode using only leg extension, and a long mode using
coordinated wing depression and leg extension[34] – and that short-mode takeoffs
result solely from activation of the GF[7]. We therefore predicted that if LPLC2
neurons with more dorsal receptive fields made more synapses with the GF than
ventral LPLC2 neurons, then we should observe more short-mode takeoffs in
response to higher, compared to lower, elevations of a looming stimulus. To test
this hypothesis, we analyzed thousands of previously collected[35] high-speed
videos of takeoffs elicited by looming stimuli at different elevations to determine
whether takeoff responses were short- or long-mode. As predicted, we found that
the percentage of short-mode takeoffs steadily increased as the elevation of the
looming stimulus increased from -30° to 77° (Fig. 1e, Extended Data Fig. 1a), as
would be expected from higher engagement of the GF in response to more dorsal
looming. Because the GF receives direct visual input from only one other VPN type
(LC4[36]) in addition to LPLC2, and LC4 neurons do not form a DV synaptic gradient
with the GF[5], the most parsimonious interpretation of our analysis is that this bias
toward short-mode takeoffs when looming occurs in the dorsal visual field results
directly from the synaptic gradient between LPLC2 and the GF. This bias
potentially enables the fly to escape more quickly from overhead predator attacks.

LPLC2 forms synaptic gradients without axonal retinotopy[4,5,37]. This means
dendrites of the GF establish differential connectivity with axons of different
individual LPLC2 neurons that are intermingled (Fig. 1f). We hypothesized that this
pattern could be achieved through differential molecular recognition: individual
LPLC2 neurons sampling different regions of visual space could express different
levels of the same cell recognition molecules. This mechanism would specify the

number of LPLC2-GF synapses independent of the spatial arrangement of LPLC2
axons.
186

scRNA-seq Uncovers Molecular Heterogeneity in VPNs

To investigate whether LPLC2 neurons exhibit molecular variation that
correlates with the synaptic gradient, we examined the LPLC2 transcriptome at
three developmental stages during and after synaptogenesis (48, 72, and 96 hours
after puparium formation (hAPF); Fig. 1g). In addition to LPLC2, we profiled its
sister cell type, LPLC1, which also forms synaptic gradients without axonal
retinotopy[5], and LC4, one of the few VPN cell types with axons arranged in a
retinotopic fashion[5]. Given that neuronal transcriptomes are very dynamic during
development and can be affected by genetic background[38], we introduced internal
controls at each time point to account for transcriptional heterogeneity driven by
these factors (Fig. 1g).
Briefly, we employed a genetic multiplexing strategy to perform pooled
single-cell profiling across multiple biological replicates[38–40] (including different
genetic backgrounds and developmental stages, see Methods). Our final dataset
included approximately 600 high-quality single-cell transcriptomes per cell type
and time point (Fig. 1h, Extended Data Fig. 1b). We validated the identity of each
VPN cell type (LPLC2, LPLC1, or LC4) based on the expression of known marker
genes[38], such as transcription factors (Fig. 1i, Extended Data Fig. 1b). This
resulted in per-cell-type coverage roughly 30 times higher than in the existing
single-cell atlases of the *Drosophila* optic lobes[38,41].
To explore heterogeneity in gene expression across each VPN cell type, we
performed Principal Component Analysis (PCA) separately for each cell type and
time point (Fig. 1 j-l, Extended Data Fig. 1c-e, 2a-b). In LPLC2 neurons at 48h APF,
the first principal component (PC1) captured a significant group of genes that were
expressed in a graded manner across individual neurons (Fig. 1j). For example,
the most differentially expressed gene across LPLC2 was dpr13. It displayed the
highest magnitude of within-cell-type variation of expression, but there was no
clear boundary separating neurons with high versus low dpr13 expression levels.
Many of the most differentially expressed genes were cell recognition molecules:
members of the Immunoglobulin Superfamily (IgSF) (e.g., dpr13, tei, beat-VI),
Leucine-rich-repeat (e.g., trn) and Cadherin (e.g., Cad87A) families[28,42](Fig, 1j).
We investigated whether this transcriptional heterogeneity reflected distinct
neuronal subtypes or a continuous variation, by forcing LPLC2 neurons to form
discrete clusters, and shuffling gene expression levels within and across clusters
(Extended Data Fig. 2c-g). Our analysis revealed that the heterogeneity was not
discrete, indicating that LPLC2 neurons form a continuum within a transcriptionally
similar neuronal type, rather than comprising several distinct subtypes. The
distribution of neurons along PC1 did not correlate with developmental age, sex,
genetic background, or mRNA coverage (Fig. 1k), indicating that the observed
within-cell-type continuous molecular heterogeneity had a different origin. The
graded expression of top differentially expressed genes associated with PC1 at
48h APF (including IgSF molecules dpr13, beat-VI, and tei, as well as SiaT

(Sialyltransferase)) was persistent throughout development and was also captured
by PC1 at 72h and 96h APF (Fig. 1l).
231   In LPLC1 neurons, PC1 also captured gradients of IgSF recognition
molecules (e.g., DIP-kappa, CG33543, dpr3, sdk) that were persistent throughout
development (Extended Data Fig. 1c-d). Similarly to LPLC2, molecular gradients
in LPLC1 could not be explained by differences in age, sex, genetic background,
or mRNA coverage across individual neurons (Extended Data Fig. 1e, 2a-b). In
contrast, molecular heterogeneity in LC4 neurons did not appear biologically
relevant. PC1 in LC4 correlated with the technical quality of single-cell
transcriptomes (number of transcripts captured per each neuron, Extended Data
Fig. 1e, 2a-b).
240   In summary, our analysis uncovered persistent expression gradients of
241   recognition molecules within LPLC1 and LPLC2 neurons. The absence of such
242   gradients in LC4 neurons suggests that molecular heterogeneity within cell types
243   may be a feature exclusive to VPNs lacking axonal retinotopy.
244
Spatial Transcriptomics Validates Molecular Gradients

246   We sought to independently verify within-cell-type gene expression
gradients in LPLC2. To do this, we used Single-Molecule Hairpin Chain Reaction
Fluorescent in Situ Hybridization[43] (HCR-FISH), a spatial transcriptomics method
that allows for the quantification of gene expression in individual cells. Combined
with Expansion-Assisted Light Sheet Microscopy[44], this approach enabled us to
visualize and count individual transcripts within neuronal somas (Fig. 2a). We
observed that individual LPLC2 somas, often positioned adjacent to one another,
exhibited striking differences in the numbers of mRNA transcripts for dpr13 and
SiaT, two of the most differentially expressed genes in the LPLC2 dataset (Fig. 2b-
c). Similar patterns were observed for beat-VI and dpr17, two differentially
expressed IgSF recognition molecules (Extended Data Fig. 3a-b).
257   To broaden our analysis to other genes, we tested pairs of genes that
exhibited correlated or anticorrelated expression patterns based on scRNA-seq
data (Fig. 1j) to determine if these relationships were replicated using HCR-FISH.
We quantified differential gene expression using Flyseg[45], an automated
volumetric instance segmentation algorithm we previously developed (see
Methods). The results of scRNA-seq and HCR-FISH were similar for genes
exhibiting antiparallel (e.g., dpr13 and SiaT, Fig. 2d, top) and parallel expression
patterns (e.g., dpr13 and beat-VI, Fig. 2d, bottom). Most of these relationships
remained consistent throughout development (Fig. 2e, Supplemental Table 2),
supporting our findings from scRNA-seq data. One exception was mbl
(*muscleblind*): although this gene exhibited graded expression in scRNA-seq (Fig.
1j), it showed low, uniform expression in HCR-FISH. The reason for this
discrepancy is unclear. Together, these data confirmed the presence of gene
expression gradients within the LPLC2 population.
271
Molecular Gradients Align with Synaptic Gradients

273   The cell bodies of the LPLC2 neurons where we measured different levels
of transcripts were arranged in salt-and-pepper fashion (Fig. 2c). LPLC2 dendrites

are, however, retinotopically arranged. We next investigated whether there was a
correlation between the retinotopic position of LPLC2 dendrites and the expression
level of genes. To do this, we used a genetic intersection method to selectively
visualize neurons within a specific cell type that express a particular gene at a
given time point (Fig. 2f-g). Recognition molecules dpr13, beat-VI, and Cad87A,
which displayed correlated expression profiles in the scRNA-seq data (Fig. 1j),
were predominantly expressed by LPLC2 neurons with dendrites in the dorsal
region of the lobula at 48h APF (Fig. 2h-i, Extended Data Fig. 4a-b). Conversely,
the expression of SiaT, dpr17, CG03419, Tsp42Ef, and stacl was limited to the
ventral region at the same developmental stage (Fig. 2h-i, Extended Data Fig. 4c-
d). We validated this DV heterogeneity using HCR-FISH, confirming that sparsely
labeled dorsal LPLC2 contained significantly more dpr13 and beat-VI transcripts
than their ventral counterparts (Extended Data Fig. 3c-d). Heterogeneous
expression of recognition molecules DIP-kappa, dpr17 and sdk in LPLC1 neurons
had similar retinotopic correlates (Extended Data Fig. 4e-f), suggesting that such
retinotopically biased expression gradients are a general feature of many VPN cell
types.
In summary, we found that the dendrites of individual neurons within a
single VPN cell type, sampling different regions of the visual space, are molecularly
heterogeneous. Across a VPN population, the neurons express gradients of
recognition molecules that match the orientation of synaptic gradients in these
VPNs. This observation suggests that even though the axons of these neurons are
spatially intermingled, they maintain their molecular differences. Our next objective
was to determine if these molecular and synaptic gradients have functional
correlates.
300

The Dpr13::DIP-ε gradient forms a looming escape synaptic gradient

Above we showed that LPLC2 neurons express IgSF recognition molecules
in a DV gradient, with dpr13 and beat-VI having higher expression levels in dorsal
LPLC2 neurons, and dpr17 higher expression in ventral ones. We hypothesized
that this DV gradient could provide the developmental origins for the gradient in
synapse number between LPLC2 neurons and the GF observed in adult flies
(Extended Data Fig. 5a). For this to be true, the GF would need to express cell
surface proteins that bind one or more of the three recognition molecules we
identified. This would allow the differential molecular expression in LPLC2 neurons
to be converted into differential cell adhesion between individual LPLC2 neurons
and the GF. Interaction maps for the IgSF molecules have previously been
established and include interactomes of Dpr and DIP proteins (DIP/Dpr[42,46]), as
well as interactomes of Beat and Side proteins (Beat/Side[47]; Fig. 3a). To test our
hypothesis, we assessed expression levels of IgSF binding partners in the GF
using HCR-FISH. DIP-ε, a binding partner of Dpr13, was the only gene showing
strong expression in the GF during development (Fig. 3b-c).
To determine if DIP-ε played a role in forming connections between LPLC2
axons and GF dendrites, we next looked at the interaction between LPLC2 axons
and GF dendrites in DIP-ε-deficient transgenic flies during development. In animals

with no DIP-ε (*DIP-ε^null* allele), we observed a ~10-fold decrease in axo-dendritic
overlap volume between LPLC2 axons and GF dendrites (Fig. 3d-e). Wild type
levels of axo-dendritic overlap could be restored in these *DIP-ε^null* flies by
overexpression of DIP-ε in just the GF in *DIP-ε^null* flies (Fig. 3f, Extended Data Fig.
5d). Knockdown of DIP-ε specifically in the GF using two different RNAi lines also
significantly reduced LPLC2-GF axo-dendritic overlap (Fig. 3f, Extended Data Fig.
5c). Thus, DIP-ε is required in the GF to establish connectivity with LPLC2 axons.
If DIP-ε is required to form functional LPLC2-GF synapses, then we would
expect that, compared to controls, DIP-ε-deficient flies shown looming stimuli
would have decreased GF responses and exhibit fewer GF-driven "short-mode"
takeoffs. Furthermore, we anticipate that, if too few LPLC2-GF synapses form to
make a DV gradient, then the number of short-mode takeoffs would no longer
increase with increasing elevation of the looming stimulus. To check these
hypotheses, we first performed whole-cell patch-clamp recordings in the GF while
presenting looming stimuli. As predicted, we observed a reduction in the peak
magnitude of the GF response in *DIP-ε^null* animals and in those expressing DIP-ε
RNAi in the GF compared to controls (Fig. 3h-i; Extended Data Fig. 6). We next
tested DIP-ε-deficient flies in our FlyPEZ behavior apparatus and found that both
DIP-ε^null animals and animals expressing DIP-ε RNAi in the GF displayed a
dramatic reduction in short-mode takeoff percentage at all stimulus elevations, and
a reduced difference between 0° and 77° elevations compared to the wild-type
control, indicative of reducing the functional synaptic gradient (Fig. 3j-k, Extended
Data Fig. 7a-b). Together with our anatomical findings, these functional data
further support a role for GF-expressed DIP-ε in establishing graded synaptic
connectivity with LPLC2.
We next sought to test the causal role of the dpr13::DIP-ε-defined gradient
on takeoff behavior by retaining strong LPLC2-GF connectivity but eliminating the
connectivity gradient. To achieve this, we increased the level of dpr13 uniformly
across the LPLC2 population (heretofore referred to as "overexpression").
Compared to controls, animals overexpressing dpr13 in LPLC2 exhibited a higher
percentage of short-mode takeoffs mediated by the GF (Fig. 3l, Extended Data
Fig. 7c). This gain-of-function was most pronounced at lower (0°) elevations,
decreased at 45°, and was absent at higher (77°) stimulus elevations, resulting in
no significant change in short-mode escape frequency with elevation. Thus, when
dpr13 is uniformly expressed across LPLC2 neurons, we observe an elimination
of the DV gradient of short-mode takeoffs that is seen in control animals (Fig. 3l).
Overexpression of dpr13 also led to an increase in LPLC2-GF axo-dendritic
overlap (Fig. 3m, Extended Data Fig. 5e-f). These data support a causal
relationship between the level of dpr13 and the number and graded distribution of
LPLC2 synapses onto the GF.
We next sought to examine the consequences of removing dpr13 on the
LPLC2-GF interaction. We examined LPLC2-GF axo-dendritic overlap in dpr13
null mutant (*dpr13^null* allele) animals (Extended Data Fig. 5g-h) and found no
difference from control animals. Our scRNA-seq data had revealed that LPLC2
neurons also express four other DIP-ε-interacting dprs (dpr14, dpr18, dpr19, and
dpr20; Extended Data Fig. 5i-j), suggesting there is indeed some functional

redundancy among the dprs expressed in LPLC2. However, unlike dpr13, the
expression of these other dprs is not graded (Extended Data Fig. 5k). To address
this, we ectopically expressed DIP-ε in LPLC2 with the intent of promoting *cis*-
interactions between DIP-ε and all five Dpr paralogs, thereby inhibiting their *trans*-
interactions with DIP-ε in the GF (an effect observed for other DIP/Dpr pairs[47,48]).
This genetic manipulation reduced LPLC2-GF axo-dendritic overlap (Fig. 3n,
Extended Data Fig. 5l-m), mimicking the phenotype of DIP-ε removal from the GF.
Together, these findings indicate that multiple Dprs binding to DIP-ε collectively
promote synaptic connectivity between LPLC2 axon terminals and GF dendrites in
a redundant manner, while dpr13 specifically contributes to establishing the
connectivity gradient.
In summary, DIP-ε and Dpr13 act as a ligand-receptor pair to control
synaptic connectivity between LPLC2 and the GF. Our data supports a model in
which a DV gradient of Dpr13::DIP-ε interactions, shaped by graded expression of
dpr13 across LPLC2 neurons, specifies the formation of the DV gradient of
synaptic weights (Extended Data Fig. 5l).
383

Beat-VI::Side-II gradient forms a motion detection synaptic gradient
We next investigated the functional role of other genes that displayed
gradients of expression across LPLC2 neurons (Fig. 1h). Like dpr13, beat-VI,
another IgSF recognition molecule[47], was also expressed in a DV gradient in
LPLC2 (higher expression in dorsal, and lower in ventral LPLC2 – Fig. 2i, Extended
Data Fig. 8a). However, neither the loss of beat-VI nor its binding partner side-II[46]
affected the overlap between LPLC2 axon terminals and the GF dendrites
(Extended Data Fig. 8b-c). This prompted us to investigate whether the beat-VI
gradient might function in a different neuronal compartment by regulating graded
inputs to LPLC2 dendrites instead.
We previously showed that beat-VI regulates synaptic specificity in the
Drosophila motion detection circuit[50]; beat-VI and its interacting partner side-II[47]
are required for matching motion-detecting T4d/T5d neurons with their
postsynaptic partners in layer 4 of the lobula plate (LoP4), which encodes
downward motion. Because LPLC2 is a part of this circuit, receiving excitatory
inputs from T4/T5 neurons in all four LoP layers[6] (Fig. 4a, top), we hypothesized
that the beat-VI gradient might similarly influence LPLC2 dendritic connectivity.
Each LPLC2 neuron has four dendritic branches in the lobula plate, one in
each layer, extending in four cardinal directions, corresponding to motion
sensitivity in each layer[6] (Fig. 4a, bottom). The response at the LPLC2 axon is a
non-linear sum of the activity in each of these four branches. Individual LPLC2
neurons thus serve as "looming detectors". They respond most strongly to looming
stimuli that originate at their receptive field center extending radially outward
simultaneously generating motion in all four cardinal directions, thereby maximally
exciting each of the four dendritic branches[6]. Using "FAFB-Flywire"[32,33]
connectome reconstruction, we found that two of the four classes of T4/T5
neurons, T4c/T5c and T4d/T5d, form antiparallel DV synaptic gradients onto
LPLC2 dendrites in Lop3 and Lop4, respectively (Fig. 4b).

These findings suggest that dorsal and ventral LPLC2 neurons within one
animal may receive differential amounts of upward and downward motion
information, potentially linked to the molecular gradient of beat-VI. Supporting this
idea, RNAi knockdown of beat-VI differentially affected LPLC2 dendrites
depending on their retinotopic location. LoP4 branches were missing in dorsal
LPLC2, while central LPLC2 exhibited a milder phenotype, and ventral LPLC2
were unaffected (Fig. 4c-d – single cell sparse visualization, Extended Data Fig.
8d-e – whole population labeling). Similarly, removing side-II from T4/T5 neurons
produced a graded phenotype, while overexpression of beat-VI in LPLC2 caused
abnormal LoP4 dendritic branching in dorsal, but not ventral, LPLC2 neurons (Fig.
4e, Extended Data Fig. 8f-g). These results support the idea that DV-graded
variations in Beat-VI and Side-II interactions differentially regulate the connectivity
between T4d/T5d axons and LPLC2 dendrites.

To determine whether beat-VI deficiency in LPLC2 had functional effects,
we performed a directional tuning experiment combined with calcium imaging in
individual LPLC2 neurons. We imaged from the unbranched neurite that connects
an LPLC2 neuron's dendrites in the LoP and Lo (see Methods, Fig. 4f-h, Extended
Data Fig 9), which has previously been shown to have weak responses to a small
bar moving in each of four cardinal directions (i.e., stimuli exciting LPLC2 dendrite
branches in a single LoP layer) and a much larger response to looming (i.e., stimuli
exciting LPLC2 dendritic branches in all four LoP layers simultaneously)[6].

Our results showed that wild type dorsal and ventral LPLC2 neurons
exhibited different patterns of responses to directional stimuli. Ventral LPLC2
showed little to no response to downward motion, as would be expected if they
received little input from downward sensing T4d/T5d neurons in LoP layer 4 (Fig.
4i, bottom, Extended Data Fig. 9b-e). As a result, responses to stimuli presented
in all directions showed a bias towards upward motion (Fig. 4j). In contrast, dorsal
LPLC2 neurons were more omnidirectional, as expected if synaptic inputs from all
four LoP layers was more evenly balanced (Fig 4i-j, Extended Data Fig. 9b-e).
Notably, after beat-VI knockdown, dorsal LPLC2 neuron responses became
biased for upward motion, similar to their wild-type ventral counterparts, consistent
with down selective T4d/T5d input being reduced. Beat-VI knockdown had little
effect on ventral LPLC2 neurons, which remained biased for upward motion (Fig.
4j, Extended Data Fig. 9b-c, e). A heatmap summarizing the experimental effects
showed that beat-VI knockdown disrupted the normal directional tuning biases of
LPLC2 neurons (Fig. 4k) without significantly affecting their responses to looming
stimuli (Fig. 4l, Extended Data Fig. 9f).

Together, these findings suggest that Beat-VI and Side-II act as a ligand-
receptor pair to control synaptic connectivity between T4d/T5d and LPLC2
neurons. Our data support a model in which the DV gradient of Beat-VI::Side-II
interactions contributes to the formation of the DV gradient of synaptic weights
between T4d/T5d and LPLC2 neurons, decreasing the response to downward
motion by LPLC2 neurons with ventral receptive fields.
455
456
457

Discussion

458
In this study, we identified the molecular origins of two parallel dorsoventral
synaptic gradients in the axons and dendrites of LPLC2, the looming-detecting
VPN cell type. These synaptic gradients translate retinotopic information from the
fly's visual field into specific motor outputs. We demonstrate that a gradient of
Beat-VI::Side-II IgSF recognition proteins regulates the graded synaptic inputs
from T4d/T5d neurons onto LPLC2, sculpting differential integration of the
downward motion component in the looming stimulus across different regions of
the eye. Similarly, a gradient of DIP-ε::Dpr13, another pair of IgSF recognition
molecules, shapes the graded synaptic outputs of LPLC2 onto the Giant Fiber,
biasing escape towards faster takeoffs in response to looming stimuli from the
dorsal visual field. These synaptic gradients, however, are distinct from each other.
For the dendritic synaptic gradient, there is a retinotopic match between
populations of input neurons (T4d/T5d) and the postsynaptic LPLC2 neurons (i.e.,
many-to-many with retinotopy). By contrast, multiple LPLC2 axons form different
numbers of synapses onto the dendrites of a single cell (GF) (i.e., *many-to-one
without retinotopy*). Irrespective of these differences, our findings suggest a
molecular model of synaptic specificity, in which the expression levels of
recognition molecules establish synapse numbers.
What is the relationship between the parallel gradients of beat-VI and dpr13
expression and the number of synaptic inputs to and outputs from LPLC2? In a
beat-VI loss-of-function experiment, the severity of functional consequences was
directly proportional to beat-VI expression levels in individual LPLC2 neurons. The
morphological phenotypes were similar whether beat-VI was removed from LPLC2
or side-II from T4/T5 neurons, with the reduction in LoP4 dendritic branching also
corresponding to beat-VI expression levels. However, it remains unclear whether
this phenotype arises from a failure to extend dendrites, or to stabilize them. We
propose that beat-VI and side-II promote adhesion between T4d/T5d axons and
LPLC2 dendrites, with the strength of this adhesion correlating with beat-VI
expression levels. This protein interaction could influence synapse number either
indirectly, by increasing the area of contact between T4d/T5d axons and LPLC2
dendrites, or directly, by controlling synapse formation, with decreased synapse
numbers leading to reduced axo-dendritic contact.
Dpr13 and DIP-ε are essential for establishing synaptic connections
between LPLC2 neurons and the GF. Loss of DIP-ε function reduced the overlap
between LPLC2 axons and the GF dendrites, leading to significant defects in GF
responses to looming stimuli, as demonstrated by both behavioral and
electrophysiological assays. Analyzing the loss of dpr13 function is complicated by
the presence of four additional Dpr paralogs in LPLC2, interacting with DIP-ε with
similar binding affinities. By ectopically expressing DIP-ε in LPLC2, we inhibited all
DIP-ε-interacting Dprs in LPLC2. This phenocopied the DIP-ε loss-of-function
phenotype in the GF, providing additional support for the interactions between DIP-
ε in the GF dendrites with its cognate binding Dprs in LPLC2 axons. The role of
multiple Dpr paralogs interacting with DIP-ε may be to ensure baseline synaptic
connectivity in the crucial escape circuit, while the formation of the synaptic
gradient depends specifically on dpr13. In support of this, increasing dpr13

expression uniformly across the LPLC2 population eliminated the graded
behavioral response to dorsal versus ventral looming stimuli, along with increasing
the LPLC2-GF axo-dendritic overlap. These observations support the idea that
Dpr13::DIP-ε interactions control the number of synapses formed between two
neurons, resulting in the formation of a synaptic gradient between LPLC2 and the
GF. However, the precise mechanism by which this occurs remains unclear.
Future studies should investigate the development of these synapses in detail, to
assess the consequences of changing dpr13 levels in individual neurons using
mosaic animals, and employing EM reconstruction or expansion microscopy to
elucidate how dpr13 and DIP-ε contribute to synaptic gradient formation at a
mechanistic level.
The dpr13::DIP-ε gradient functions differently from Eph and Ephrin
gradients in vertebrate retinotectal maps[51,52]. Both gradients relay visual-spatial
information from the eye to the brain but use distinct strategies. Ephrins and Ephs
form topographic maps by maintaining the positional information of neurons in the
retina to arrangement of their axons in the midbrain[53] (i.e., retinotopy). In contrast,
the dpr13::DIP-ε gradient translates positional information from VPN dendrites into
synapse numbers in the central brain, independent of axon retinotopy. This
represents a new molecular mechanism of visuomotor transformation.
As most VPNs form synaptic gradients[5], varying levels of adhesion
molecules may represent a general strategy for regulating synapse numbers.
Notably, altering levels of SynCAM1, another IgSF adhesion molecule, also affects
synapse numbers in the mammalian hippocampus[54]. Together, these data suggest
that not only do the cell recognition molecules determine synaptic specificity by
qualitatively matching synaptic partners[26–28] (a classic notion established through
decades of genetic studies), but that their expression levels may also influence the
number of synapses formed between matched neurons with functional and
behavioral consequences. This variation holds an enormous potential for fine-
tuning ethologically relevant behavioral responses driven by complex neural
circuits.
Our work has uncovered graded molecular heterogeneity within single VPN
cell types. This feature is distinct from discrete[38,41] or stochastic[55,56] variations in
gene expression within neuronal types previously described in *Drosophila*. We
identified many graded genes in VPNs beyond cell recognition molecules,
suggesting that molecular gradients may regulate other cellular and circuit
functions. These findings are particularly interesting as cell types in the mammalian
cortex also exhibit continuous transcriptomic variation[8,57–59]. These data raise the
possibility that within-cell-type molecular heterogeneity could be a common
mechanism for generating neuronal diversity which, in turn, contribute to synaptic
specificity.
In conclusion, we describe a new molecular strategy regulating neural
circuit assembly. This was enabled through detailed EM-connectomics of the fly
brain[22,31,33] and rapid advances in single-cell sequencing technologies. We
anticipate that merging these approaches to study the mammalian brain[60,61] will
uncover new molecular mechanisms underlying wiring specificity.
549

References

550
1. Shin, S., Crapse, T. B., Angeles, L., Mayo, J. P. & Sommer, M. A.
Visuomotor Integration. *Encycl. Neurosci.* (2009).
2. Andersen, R. A., Snyder, L. H., Li, C. S. & Stricanne, B. Coordinate
transformations in the representation of spatial information. *Curr. Opin.*
Neurobiol. **3**, 171–176 (1993).
3. Otsuna, H. & Ito, K. Systematic analysis of the visual projection neurons of
Drosophila melanogaster. I. Lobula-specific pathways. *J. Comp. Neurol.*
497, 928–958 (2006).
4. Wu, M. *et al.* Visual projection neurons in the Drosophila lobula link feature
detection to distinct behavioral programs. *Elife* **5**, 1–43 (2016).
5. Dombrovski, M. *et al.* Synaptic gradients transform object location to
action. *Nature* **613**, 534–542 (2023).
6. Klapoetke, N. C. *et al.* Ultra-selective looming detection from radial motion
opponency. *Nature* **551**, 237–241 (2017).
7. Von Reyn, C. R. *et al.* A spike-timing mechanism for action selection. *Nat.*
Neurosci. **17**, 962–970 (2014).
8. Cembrowski, M. S. & Menon, V. Continuous Variation within Cell Types of
the Nervous System. *Trends Neurosci.* **41**, 337–348 (2018).
9. Ghahramani, Z., Wolpert, D. M. & Jordan, M. I. Generalization to local
remappings of the visuomotor coordinate transformation. *J. Neurosci.* **16**,
7085–7096 (1996).
10. Fujiwara, Y., Lee, J., Ishikawa, T., Kakei, S. & Izawa, J. Diverse coordinate
frames on sensorimotor areas in visuomotor transformation. *Sci. Reports*
2017 71 **7**, 1–9 (2017).
11. Buneo, C. A., Jarvis, M. R., Batista, A. P. & Andersen, R. A. Direct
visuomotor transformations for reaching. *Nature* **416**, 632–636 (2002).
12. Helmbrecht, T. O., dal Maschio, M., Donovan, J. C., Koutsouli, S. & Baier,
H. Topography of a Visuomotor Transformation. *Neuron* **100**, 1429-
1445.e4 (2018).
13. Kim, J. H., Ma, D. H., Jung, E., Choi, I. & Lee, S. H. Gated feedforward
inhibition in the frontal cortex releases goal-directed action. *Nat. Neurosci.*
2021 2410 **24**, 1452–1464 (2021).
14. Huston, S. J. & Jayaraman, V. Studying sensorimotor integration in insects.
Curr. Opin. Neurobiol. **21**, 527–534 (2011).
15. Cruz, T. L. & Chiappe, M. E. Multilevel visuomotor control of locomotion in
Drosophila. *Curr. Opin. Neurobiol.* **82**, 102774 (2023).
16. Wang, T. S. L., Martinez, M., Festa, E. K., Heindel, W. C. & Song, J. H.
Age-related enhancement in visuomotor learning by a dual-task. *Sci.*
Reports 2022 121 **12**, 1–12 (2022).
17. Dombrovski, M. *et al.* A Plastic Visual Pathway Regulates Cooperative
Behavior in Drosophila Larvae. *Curr. Biol.* (2019)
doi:10.1016/j.cub.2019.04.060.
18. Zada, D. *et al.* Development of neural circuits for social motion perception
in schooling fish. *Curr. Biol.* **0**, (2024).
19. Wei, P. *et al.* Processing of visually evoked innate fear by a non-canonical

596   thalamic pathway. *Nat. Commun. 2015 61* **6**, 1–13 (2015).

20. Zhao, P. *et al.* The visuomotor transformations underlying defensive
behaviors and hunting. *bioRxiv* 2024.05.07.592863 (2024)
doi:10.1101/2024.05.07.592863.

21. Cande, J. *et al.* Optogenetic dissection of descending behavioral control in
Drosophila. *Elife* **7**, (2018).

22. Nern, A. *et al.* Connectome-driven neural inventory of a complete visual
system. *bioRxiv* 2024.04.16.589741 (2024)
doi:10.1101/2024.04.16.589741.

23. Klapoetke, N. C. *et al.* A functionally ordered visual feature map in the
Drosophila brain. *Neuron* **110**, 1700-1711.e6 (2022).

24. Aptekar, J. W., Keleş, M. F., Lu, P. M., Zolotova, N. M. & Frye, M. A.
Neurons forming optic glomeruli compute figure–ground discriminations in
Drosophila. *J. Neurosci.* **35**, 7587–7599 (2015).

25. Keleş, M. F. & Frye, M. A. Object-Detecting Neurons in Drosophila. *Curr.*
Biol. **27**, 680–687 (2017).

26. Margeta, M. A. & Shen, K. Molecular mechanisms of synaptic specificity.
Mol. Cell. Neurosci. **43**, 261–267 (2010).

27. Südhof, T. C. Towards an Understanding of Synapse Formation. *Neuron*
100, 276 (2018).

28. Sanes, J. R. & Zipursky, S. L. Synaptic Specificity, Recognition Molecules,
and Assembly of Neural Circuits. *Cell* **181**, 536–556 (2020).

29. Bacon, J. P. & Strausfeld, N. J. The dipteran 'Giant fibre' pathway: neurons
and signals. *J. Comp. Physiol. A* **158**, 529–548 (1986).

30. Jang, H., Goodman, D. P., Ausborn, J. & von Reyn, C. R. Azimuthal
invariance to looming stimuli in the Drosophila giant fiber escape circuit. *J.*
Exp. Biol. **226**, (2023).

31. Scheffer, L. K. *et al.* A connectome and analysis of the adult drosophila
central brain. *Elife* **9**, 1–74 (2020).

32. Zheng, Z. *et al.* A Complete Electron Microscopy Volume of the Brain of
Adult Drosophila melanogaster. *Cell* **174**, 730-743.e22 (2018).

33. Schlegel, P. *et al.* Whole-brain annotation and multi-connectome cell typing
quantifies circuit stereotypy in Drosophila. *bioRxiv* 2023.06.27.546055
(2023) doi:10.1101/2023.06.27.546055.

34. Card, G. & Dickinson, M. H. Visually Mediated Motor Planning in the
Escape Response of Drosophila. *Curr. Biol.* **18**, 1300–1307 (2008).

35. Williamson, R., Peek, M. Y., Breads, P., Coop, B. & Card, G. M. Tools for
Rapid High-Resolution Behavioral Phenotyping of Automatically Isolated
Drosophila. *Cell Rep.* **25**, (2018).

36. Ache, J. M. *et al.* Neural Basis for Looming Size and Velocity Encoding in
the Drosophila Giant Fiber Escape Pathway. *Curr. Biol.* **29**, 1073-1081.e4
(2019).

37. Moreno-Sanchez, A. *et al.* Morphology and synapse topography optimize
linear encoding of synapse numbers in Drosophila looming responsive
descending neurons. *bioRxiv Prepr. Serv. Biol.* (2024)
doi:10.1101/2024.04.24.591016.

38.   Kurmangaliyev, Y. Z., Yoo, J., Valdes-Aleman, J., Sanfilippo, P. &
Zipursky, S. L. Transcriptional Programs of Circuit Assembly in the
Drosophila Visual System. *Neuron* **108**, 1045-1057.e6 (2020).
39.   Kang, H. M. *et al.* Multiplexed droplet single-cell RNA-sequencing using
natural genetic variation. *Nat. Biotechnol. 2017 361* **36**, 89–94 (2017).
40.   MacKay, T. F. C. *et al.* The Drosophila melanogaster Genetic Reference
Panel. *Nature* **482**, 173–178 (2012).
41.   Özel, M. N. *et al.* Neuronal diversity and convergence in a visual system
developmental atlas. *Nature* **589**, 88–95 (2021).
42.   Özkan, E. *et al.* An extracellular interactome of immunoglobulin and LRR
proteins reveals receptor-ligand networks. *Cell* **154**, 228 (2013).
43.   Choi, H. M. T. *et al.* Third-generation in situ hybridization chain reaction:
multiplexed, quantitative, sensitive, versatile, robust. *Development* **145**,
(2018).
44.   Lillvis, J. L. *et al.* Rapid reconstruction of neural circuits using tissue
expansion and light sheet microscopy. *Elife* **11**, 1–36 (2022).
45.   Vaccari, A. & Dombrovski, M. FlySeg: An Automated Volumetric Instance
Segmentation Algorithm for Dense Cell Populations in Drosophila
Melanogaster Nervous System. *Conf. Rec. - Asilomar Conf. Signals, Syst.*
Comput. 1474–1478 (2023)
doi:10.1109/IEEECONF59524.2023.10476819.
46.   Cosmanescu, F. *et al.* Neuron-Subtype-Specific Expression, Interaction
Affinities, and Specificity Determinants of DIP/Dpr Cell Recognition
Proteins. *Neuron* **100**, 1385-1400.e6 (2018).
47.   Li, H. *et al.* Deconstruction of the beaten path-sidestep interaction network
provides insights into neuromuscular system development. *Elife* **6**, (2017).
48.   Menon, K. P., Kulkarni, V., Shin-Ya, T., Anaya, M. & Zinn, K. Interactions
between dpr11 and dip-y control election of amacrine neurons in
drosophila color ision circuits. *Elife* **8**, 1–32 (2019).
49.   Morano, N. C. *et al.* Cis inhibition of co-expressed DIPs and Dprs shapes
neural development. *bioRxiv* 2024.03.04.583391 (2024)
doi:10.1101/2024.03.04.583391.
50.   Yoo, J. *et al.* Brain wiring determinants uncovered by integrating
connectomes and transcriptomes. *Curr. Biol.* **33**, 3998-4005.e6 (2023).
51.   Brown, A. *et al.* Topographic mapping from the retina to the midbrain is
controlled by relative but not absolute levels of EphA receptor signaling.
Cell **102**, 77–88 (2000).
52.   Triplett, J. W. & Feldheim, D. A. Eph and ephrin signaling in the formation
of topographic maps. *Semin. Cell Dev. Biol.* **23**, 7 (2012).
53.   Kita, E. M., Scott, E. K. & Goodhill, G. J. Topographic wiring of the
retinotectal connection in zebrafish. *Dev. Neurobiol.* **75**, 542–556 (2015).
54.   Robbins, E. M. *et al.* SynCAM 1 Adhesion Dynamically Regulates Synapse
Number and Impacts Plasticity and Learning. *Neuron* **68**, 894–906 (2010).
55.   Wernet, M. F. *et al.* Stochastic spineless expression creates the retinal
mosaic for colour vision. *Nature* **440**, 174–180 (2006).
56.   Miura, S. K., Martins, A., Zhang, K. X., Graveley, B. R. & Zipursky, S. L.

Probabilistic splicing of Dscam1 establishes identity at the level of single
neurons. *Cell* **155**, 1166 (2013).
57. Cembrowski, M. S. & Spruston, N. Heterogeneity within classical cell types
is the rule: lessons from hippocampal pyramidal neurons. *Nat. Rev.*
Neurosci. 2019 204 **20**, 193–204 (2019).
58. Cheng, S. *et al.* Vision-dependent specification of cell types and function in
the developing cortex. *Cell* **185**, 311-327.e24 (2022).
59. Yao, Z. *et al.* A taxonomy of transcriptomic cell types across the isocortex
and hippocampal formation. *Cell* **184**, 3222-3241.e26 (2021).
60. Sievers, M. *et al.* Connectomic reconstruction of a cortical column. *bioRxiv*
2024.03.22.586254 (2024) doi:10.1101/2024.03.22.586254.
61. Yao, Z. *et al.* A high-resolution transcriptomic and spatial atlas of cell types
in the whole mouse brain. *Nature* **624**, 317–332 (2023).
62. Gramates, L. S. *et al.* FlyBase: a guided tour of highlighted features.
Genetics **220**, (2022).
63. Kang, H. M. *et al.* Multiplexed droplet single-cell RNA-sequencing using
natural genetic variation. *Nat. Biotechnol. 2017 361* **36**, 89–94 (2017).
64. Li, H. A statistical framework for SNP calling, mutation discovery,
association mapping and population genetical parameter estimation from
sequencing data. *Bioinformatics* **27**, 2987–2993 (2011).
65. Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable
single-cell analysis. *Nat. Biotechnol.* **42**, 293–304 (2024).
66. Gabbiani, F., Krapp, H. G. & Laurent, G. Computation of Object Approach
by a Wide-Field, Motion-Sensitive Neuron. *J. Neurosci.* **19**, 1122–1141
(1999).
67. Frighetto, G. & Frye, M. A. Columnar neurons support saccadic bar
tracking in Drosophila. *Elife* **12**, (2023).
68. Reiser, M. B. & Dickinson, M. H. A modular display system for insect
behavioral neuroscience. *J. Neurosci. Methods* **167**, 127–139 (2008).
69. Zhao, A. *et al.* Eye structure shapes neuron function in Drosophila motion
vision. *bioRxiv* 2022.12.14.520178 (2022) doi:10.1101/2022.12.14.520178.
70. DeBruine, L. M. & Barr, D. J. Understanding Mixed-Effects Models Through
Data Simulation. *Adv. Methods Pract. Psychol. Sci.* **4**, (2021).
71. Saravanan, V., Berman, G. J. & Sober, S. J. Application of the hierarchical
bootstrap to multi-level data in neuroscience. *Neurons, Behav. data Anal.*
theory **3**, (2020).
72. Xu, S. *et al.* Interactions between the Ig-Superfamily Proteins DIP-α and
Dpr6/10 Regulate Assembly of Neural Circuits. *Neuron* **100**, 1369-1384.e6
(2018).
73. Xu, H. *et al.* Sequence determinants of improved CRISPR sgRNA design.
Genome Res. **25**, 1147–1157 (2015).
74. Ren, X. *et al.* Optimized gene editing technology for Drosophila
melanogaster using germ line-specific Cas9. *Proc. Natl. Acad. Sci. U. S. A.*
110, 19012–19017 (2013).
75. Wassie, A. T., Zhao, Y. & Boyden, E. S. Expansion microscopy: principles
and uses in biological research. *Nat. Methods* **16**, 33–41 (2019).

76. Wang, Y. *et al.* EASI-FISH for thick tissue defines lateral hypothalamus
spatio-molecular organization. *Cell* **184**, 6361-6377.e24 (2021).
77. Lee, P. T. *et al.* A gene-specific T2A-GAL4 library for Drosophila. *Elife* **7**,
(2018).
78. Nern, A., Pfeiffer, B. D. & Rubin, G. M. Optimized tools for multicolor
stochastic labeling reveal diverse stereotyped cell arrangements in the fly
visual system. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E2967–E2976 (2015).
79. Matsliah, A. *et al.* Neuronal "parts list" and wiring diagram for a visual
system. *bioRxiv* (2024) doi:10.1101/2023.10.12.562119.

743
744
745

Materials and Methods

Experimental model details

Flies were reared under standard conditions at 25 °C and 50% humidity with
a 12-h light/12-h dark cycle on a standard cornmeal fly food. Males and females 1-
3 days after eclosion were used for all experiments except where specified.
Supplementary Table 1 provides detailed descriptions of fly genotypes used in
each experiment and origins of transgenic stocks. For developmental experiments,
white pre-pupae (0h APF) were collected and incubated at 25 °C for the indicated
number of hours.

scRNA-seq experiment

Virgin females carrying LPLC2, LPLC1 or LC4 split-GAL4 driver lines were
crossed to males expressing a nuclear GFP reporter (UAS-His2A-GFP) and
carrying unique 3rd chromosomes from the isogenic wild-type strains with known
genotypes (DGRP, *Drosophila* Reference Genetic Panel[40]). Each experimental
condition (cell type and time point) was crossed to 6 unique DGRP strains (see
Supplemental Table 1 and Source Data for details; LC4 driver was crossed into 4
DGRP strains for 72 and 96h APF time points). Each of these crosses was
considered an independent biological replicate. F1 generation animals (males and
females) were collected at 0h APF and incubated for either 48h, 72h or 96 hours
at 25°C. To introduce a control for developmental age, we split 6 DGRPs (for each
cell type and time point) into equal "early" and "late" groups (Fig. 1g, see Source
data for a detailed breakdown). Animals from "early" DGRPs were continuously
staged and collected within the 2h time window, after which we continuously
staged and collected animals from "late" DGRPs within the next 2h time window.
This ensured that by the time brains were dissected (after 48, 72 or 96 hours),
animals from the "early" group were on average 2h older than their "late"
counterparts. Brains were dissected in ice-cold Schneider's *Drosophila* Medium
(GIBCO #21720-024), collected in three Eppendorf tubes (each time point was
processed as a separate sample) and kept on ice. Brains were incubated in mix of
papain (Worthington #LK003178, 1000 units/ml) and Liberase™ (Sigma-Aldrich
#5401119001, 0.18 Wu/ml) at 25°C for 30 min shaking at 1000rpm, washed twice

with 2% Bovine Serum Albumin (BSA, Sigma #B6917) in Schneider's Medium,
followed by two washes in 1% BSA in PBS. Finally, brains were transferred to
0.04% BSA in PBS and dissociated mechanically by pipetting. The cell suspension
was filtered through a 20 µm cell-strainer (Corning #352235) and sorted by FACS
(BD FACS Aria II) to purify GFP+ cells. Single-cell suspensions were used to
generate scRNA-seq libraries using the 10X Genomics Chromium Next GEM
Single Cell 3'-kit (v3.1) following the manufacturer's protocol. Each sample
(corresponding to a single time point) was loaded to a single lane of 10X
Chromium. Three scRNA-seq libraries were sequenced using one lane of
NovaSeq 6000 SP platform (28bp + 91 bp). The library preparation and
sequencing were performed by the Technology Center for Genomics and
Bioinformatics at UCLA.
791

scRNA-seq data processing and analysis

Raw scRNA-Seq reads were processed using Cell Ranger (10X Genomics,
version: 7.1.0). The reference genome and gene annotations were downloaded
from FlyBase[62] (release 6.29). Each time point was processed separately. Six
biological replicates were tagged with a unique wild-type chromosome, and
demultiplexed based on a unique wild-type chromosome using demuxlet[63] (version
2, https://github.com/statgen/popscle). Demultiplexing was performed using
single-nucleotide polymorphisms (SNPs) from 7 DGRP strains used in
experiments (see supplemental table 1 for the full list of genotypes and Source
Data for the list of DGRP strains used for each genotype and time point) and 3
additional DGRP strains as negative controls (line_129, line_427, line 712). SNPs
were filtered using the following criteria: (1) only bi-allelic SNPs on the 3rd
chromosome without missing data (called in all 10 strains); (2) non-reference allele
only in 1 of 10 strains. We quantified allelic counts for filtered SNPs using samtools
mpileup[64] (version 1.10). SNPs with a minimum total coverage of 10 in all three
samples (time points) and a maximum non-reference allele frequency of 0.25 were
kept for downstream analysis (34,655 SNPs). Only a few cells (0.1-0.2%) were
erroneously assigned to negative controls; 5–15% of cells were classified as
"doublets" and "ambiguous" (mostly cells with low transcript coverage). We
removed cells with less than 10,000 or more than 50,000 transcripts per cell (and
more than 10% of mitochondrial transcripts). The final dataset included 2,595 cells
for 48h APF, 2,369 cells for 72h APF, and 1,039 cells for 96h APF.

The scRNA-seq analysis was performed using Seurat[65] package (5.0.1). The
analysis was performed separately for each time point using the standard Seurat
workflow: raw transcript counts were normalized, 1000 highly variable genes were
scaled and used for PCA, first 5 PCs were used for clustering (resolution: 0.05)
and to calculate t-SNE projections. We use t-SNE projection only for the summary
visualization of the dataset. Clusters were annotated based on known marker
genes[38] (Fig. 1h-i and Extended Data Fig. 1d). Most of the cells corresponded to
LPLC2, LPLC1, and LC4 neurons. To explore further heterogeneity within VPN
type, we subsetted each cell type at each time point, identified 500 highly variable
genes, and repeated PCA. We plotted known biological and technical covariates

along each analyzed PC (Fig. 1k and Extended Data Fig. 1-2), including
developmental age, DGRP genotype, sex, and coverage (i.e., transcripts per cell).
PC1s in LPLC2 and LPLC1 did not correlate with any of these covariates and were
driven by similar sets of genes (Fig. 1j, l and Extended Data Fig. 1e-f). Further *in*
vivo validations using orthogonal approaches confirmed that these PCs captured
true molecular heterogeneity within each of these cell types (Fig. 2).

Behavioral experiments
High-throughput takeoff assay
High-throughput takeoff assay was performed with the FlyPEZ system, which
allows for the near-automated collection of fly behaviors in response to visual
stimulation in large sample sizes. FlyPEZ experiments were performed as
previously described[35]. Briefly, untrained flies (3-5 days old, both males and
females) were individually dispensed onto a 5 mm by 5 mm glass recording
platform via an automated "FlyGate" module[35], which controls the opening and
closing of the gate using a laser light sheet and linear photodetector. Once the fly
reaches the platform, four mirrors allow simultaneous recording of two views of the
fly (a bottom view and a side view) onto a single camera lens, which enables real-
time detection of the center of mass and heading of the fly via the "FlyDetect"
algorithm. The heading direction is then used to generate looming stimulation in
fly coordinates, and the behavioral response of the fly is recorded by a high-speed
camera at 6,000 frames per second under infrared illumination. A single stimulus
was presented per fly. All behavioral experiments were performed 4 hours before
incubator lights were switched off, which coincides with the flies' activity peak in
the afternoon light cycle.

Visual stimulation for behavioral assay
A 7-inch-diameter back-projection coated dome is placed centered over the glass
platform to present visual stimulation. Specifically, dark looming disks that
approach the fly from azimuth of 0° (front looms) or 90° (side looms), at elevations
of -30°, 0°, 23°, 45°, or 77° in fly coordinates were used. Looming stimuli were
generated using the same equation as described for calcium imaging experiments
(see below). All looming stimuli have l/v=40ms. Experiments in Figure 1e and Ext.
Fig 1a show trials that were performed in the past (from 2014 to 2024) using control
flies. Both looming disks expanding from 10° to 180° and 5° to 90° were combined
to get enough trials. Experiments in Figure 3 include trials with looming disks
expanding from 10° to 180° only.

Behavioral data analysis
To quantify the duration of the takeoff sequence, videos were manually annotated
to identify the start of the sequence (the first frame of wing-rising) and the end of
the sequence (the last frame that shows T2 legs in contact with the platform).
Takeoff sequence durations between 0ms to 7ms were considered "short-mode
takeoffs," and takeoff sequence durations longer than 7ms were considered "long-
mode takeoffs," as described previously[7]. The total takeoff percentage is

calculated by the number of takeoffs divided by the total number of trials. Short-
mode takeoff percentage is calculated by the number of short-mode takeoffs
divided by the total number of takeoffs. For experiments in Figure 1d, 1e, and Ext.
Fig 1c, takeoff sequence duration longer than 50ms was eliminated as outliers. All
takeoff sequence durations were less than 50ms for experiments in Figure 3.
874

Statistical analysis

Statistical comparison of the percentages of short-mode takeoffs was performed
with the Chi-squared test, with post-hoc Bonferroni correction for multiple
comparisons. Statistical comparison of takeoff sequence distributions between two
samples was performed with the Mann-Whitney U test. Statistical comparison of
takeoff sequence distributions between more than two samples was performed
with the Kruskal-Wallis test, with post-hoc Dunn correction for multiple
comparisons. Analysis and plotting were conducted with custom scripts in
MATLAB 2022b, and Scipy and Seaborn in Python 3.
884

Electrophysiological experiments

886
Electrophysiological recordings

In vivo whole-cell, current-clamp electrophysiology was performed on behaving,
tethered flies as previously described[30]. Flies were anesthetized at 4°C and their
head and thorax were attached to polyether ether ketone plates with UV glue
(Loctite 3972). The T1 legs were severed at the femur to prevent cleaning of the
head and obstruction of the eyes. The proboscis was secured in its retracted
position to minimize brain movement during the recording. The antennae were also
fixed with UV glue to restrict stimulation of antennal afferents to the GF. To access
the GF for recordings, the cuticle and trachea on the posterior side of the head
above the GF soma were removed and the brain was perfused with standard
extracellular saline (NaCl 103 mmol, KCl 3 mmol, TES 5 mmol, trehalose-$2H_2O$ 8
mmol, glucose 10 mmol, $NaHCO_3$ 26 mmol, NaH_2PO_4 1 mmol, $CaCl_2 \cdot 2H_2O$ 1.5
mmol, and $MgCl_2 \cdot 6H_2O$ 4 mmol; Gouwens and Wilson 2009). Osmolarity was
adjusted to 270–275 mOsm and bubbled with 95% O2/5% CO2 to maintain a pH
of 7.3. All experiments were conducted at room temperature (20–22°C). A brief,
localized application of collagenase (0.5% in extracellular saline) with a glass
electrode was used to disrupt the brain sheath and access the soma for recording.
GFP-labeled soma was then targeted by patch-clamp electrodes (4–6 MΩ)
containing intracellular saline (potassium aspartate 140 mmol, KCl 1 mmol, Hepes
10 mmol, EGTA 1 mmol, Na_3GTP 0.5 mmol, MgATP 4 mmol, Alexafluor-568 5
μmol, 265 mOsm, pH 7.3). In vivo whole-cell recordings were performed in current-
clamp mode using a MultiClamp 700B amplifier, and digitized (NI-DAQ, National
Instruments) at 20 kHz. All data were acquired using the open-source software
Wavesurfer (https://wavesurfer.janelia.org/) running in MATLAB (MathWorks).
Traces were not corrected for a 13-mV liquid junction potential (Gouwens and
Wilson, 2009). Recordings were considered acceptable when the initial seal
resistance was >2 GΩ before rupture, the resting membrane potential was less

than −50 mV, and the input resistance was >50 MΩ, as the typical input resistance
of the GF has been reported to be in the range of 50 to 100 MΩ.
916

<u>Visual stimulation for electrophysiology and data analysis</u>
Visual stimuli were back-projected onto a 4.5-inch diameter mylar cylindrical
screen covering 180 degrees in azimuth via two DLP projectors (Texas
Instruments Lightcrafter 4500) as previously described[33]. The projected images
were displayed in 6-bit grayscale at 240Hz, above the flicker fusion frequency of
Drosophila (100Hz; Niven et al., 2003). All stimulus frames were generated in
MATLAB and presented using Psychtoolbox (http://psychtoolbox.org/). Looming
stimuli were generated with the background set to white and the expanding disk
(looming stimulus) set to black on the projector (5500 lx-white and 15 lx-black,
measured at the position of the fly). The angular size (θ) of the stimulus subtended
by the approaching object was calculated in time (t) by the following equation[66]:

$$\theta(t) = 2tan^{-1}\left(\frac{r}{vt}\right),$$

where t<0 before collision and t=0 at collision for an approaching object with a
radius (r) and constant velocity (v). As previously described[66], r/v (ms) is the radius
to approach speed ratio. The smaller the r/v, the more abrupt the stimulus
expansion. The visual stimulus set consisted of four looming stimuli (r/v=10, 20,
40, 80 ms) starting at 10 degrees and expanding to 63 degrees in angular size and
held for 1 second, displayed at −45 degrees in azimuth, 0 degrees in elevation.
Each visual stimulus was presented once per trial, in a randomized order.
Interstimulus intervals were set to 30 seconds to allow enough time for the GF
membrane potential to return to baseline and avoid habituation. The responses to
the same stimulus from two successive trials were then averaged for each fly.
Electrophysiology data distributions were evaluated using a Kolmogorov-Smirnov
test. If the data were found not to follow a normal distribution, the appropriate non-
parametric test was selected. Statistical comparison of GF's looming responses
between two groups was performed with the Mann-Whitney U test.
944

Two-photon calcium imaging experiments
<u>Imaging setup</u>
Calcium imaging was performed with a VIVO Multiphoton Open (Intelligent
Imaging Innovation, Inc.) system based on a Movable Objective Microscope
(MOM, Sutter Instruments). The excitation of the sample was delivered by a
Ti:Sapphire laser (Chameleon Vision I, Coherent) tuned to 920 nm with power
ranging from 15 to 30 mW (depending on imaging depth). A dual axis mirror
galvanometer was used for x-y laser scanning (RGG scanbox, Sutter Instrument).
We imaged with a 20x water-immersion objective (W Plan-Apochromat 20x/1.0
DIC, Zeiss) and a band-pass filter (Semrock 525/40 nm) was placed in front of the
photomultiplier tube (H11706P-40, Hamamatsu) to reduce the blue light from the
visual display. Microscope and data acquisition were controlled by Slidebook 2024
(Intelligent Imaging Innovation, Inc.). Single plane images were sampled at 9 Hz
with a spatial resolution of approximately 180 x 180 pixels (95.7 x 95.7 µm, pixel

size $\cong$ 0.53 µm, dwell time $\cong$ 2 µs). Images and external visual stimuli were
synchronized *a posteriori* using frame capture markers (TTL pulses from Slidebook
2024) and stimulus events (analog outputs from the visual display) sampled with
a data acquisition device (USB-6229, National Instruments) at 10 kHz.
964

Fly tethering and preparation for imaging
Flies were prepared and head-fixed to a custom objective stage fly holder as
previously described[67]. The cuticle above the right optic lobe was removed and the
brain bathed in isotonic saline. The holder with the tethered fly was placed under
the objective at the center of the visual display in the horizontal plane. GCaMP7f
responses of dendritic branches from individual LPLC2 neurons were recorded
from a posterior view. The fly head was pitched forward pointing down at the visual
display so that the fly eye's equator held a pitch angle of approximately 60° relative
to the imaging plane. For each fly, we identified the most dorsocaudal dendritic
arbors in the lobula plate and then moved the focal plane ~10 µm below them to
start mapping the receptive field (RF) centers of dorsal LPLC2 neurons, or moved
~50 µm to probe ventral LPLC2 RFs, similarly to previous calcium imaging
experiments in LPLC2[6]. Random steps ($\pm$ 5 µm) between these two bracketed Z-
planes were used to probe the RF centers of dorsoventrally intermediate LPLC2
neurons. Unstable recordings or recordings from preparations that did not respond
during the RF scanning trials were not included in the data set.
981

Visual stimuli for imaging
A visual display composed of 48 8x8 dot matrix LED panels arranged in a semi-
cylinder[68] was used for visual stimulation as previously described[67]. Four layers of
filter (071, LEE Filters) were placed over the display to reduce its light intensity. To
compensate for the angle of the eye's equator and optimize the extension of the
surrounding visual context, the display was tilted forward at an angle of 30° from
the horizontal plane. Visual presentation was confined to the right half of the visual
field, ipsilateral to the recording site. Visual stimuli were generated and controlled
using custom-written MATLAB (MathWorks) scripts that communicated to the
display through the microcontroller serial port. Looming stimuli simulated an object
approaching the fly at a constant velocity, equivalent to twice the inverse tangent
of the ratio between object's half-size and object's approach speed (see
description of electrophysiological experiments). The display background was set
to 70% maximum intensity while foreground objects (looming or moving bars) were
set to 0%. The set of visual stimuli was presented in random block design and
repeated 2 times. Each visual stimulation lasted 4 s and was composed by 0.5 s
of uniform background, and 0.5 s of visual motion followed by 3 s of static pattern.
Each trial was followed by 3 s of rest in which flies faced the visual background.
1000

Receptive field center and directional tuning
We identified an active neurite from a single neuron in the multiphoton field of view
(Extended Data Fig. 9a, bottom). The RF center of that neurite was identified in
real-time and subsequently scanned for directional sensitivity. We developed a
custom GUI in MATLAB, which allowed for real-time modifications to stimulus

positions on the visual display. This interface enabled hand-triggered looming
stimuli and the visual inspection of GCaMP responses. To identify the RF center
of an LPLC2 dendritic branch, we created a rectangular grid of 48 positions across
the right half of the visual display.
The positions were spaced every 5 LEDs in both horizontal and vertical directions,
with each LED covering approximately 2.2° on the retina at the eye's equator.
Using the GUI, the experimenter presented a looming stimulus centered at each
grid position. The looming stimulus simulated a circular object with a 0.5 cm radius,
starting from a distance of 50 cm and traveling at 62.5 cm/s. This caused the object
to expand from 0.6° to 14° with a loom velocity (l/v) of 8 ms. If a response was
visually detected, the surrounding grid positions were probed next. The position
with the highest peak response was taken as the RF center for the subsequent
directional tuning experiment. We tested directional selectivity by moving a dark
edge outward from the center of the RF in 24 different directions (Fig. 4g, top;
Extended Data Fig. 10). The edges moved at 20°/s with orientations ranging from
0° to 345° in 15° increments. Each edge subtended 15° at the eye's equator and
swept 15° orthogonal to its orientation, filling a 15° black square upon completion.
Additionally, a looming stimulus centered within the RF, with the same dynamics
as those used for the RF scans, was included in this battery of dark moving edges.
1025
LPLC2 position's mapping and directional sensitivity index
We sampled neurons along the DV and AP axes of the lobula and confirmed their
anatomical locations by mapping the RF centers onto the fly eye (Fig. 4g, bottom).
To identify the putative individual LPLC2 neurons stimulated by the RF center
scans, we mapped the horizontal coordinates of their retinal positions onto the 2D
retinal ommatidia lattice. We identified specific dorsal and ventral retinal ommatidia
and their corresponding columnar LPLC2 in the fly connectome, verifying the
recorded LPLC2 neurons' locations (Fig. 4h, Extended Data Fig. 9a).
Coordinates were calculated using a 3D reconstruction of the fly head, holder, and
visual display in AutoCAD (Autodesk). We estimated the fly ommatidia with
overlapping horizontal coordinates through the following steps:
(1) Identified the locations of the ommatidia pointing to positions 16 and 40 based
on a Mollweide projection of 3D ommatidia directions from a microCT scan[69];
(2) Mapped these ommatidia locations onto identified visual columns of the male
optic lobeconnectome[22];
(3) Used T4 neurons included in these visual columns to identify downstream
LPLC2 neurons[22].
For each recording, the direction sensitivity index (DSI) was computed as follows:
$$DSI = (R_{up} - R_{down})/(R_{up} + R_{down})$$
where R_{up} is the peak response to an upward moving edge (0° direction) and
R_{down} is the peak response to a downward moving edge (180° direction). The
index ranges from -1 to 1, with negative values indicating downward sensitivity and
positive values indicating upward sensitivity. The heatmap of the DSI for the tested
positions was smoothed with a Gaussian filter (σ = 1).
1050
Imaging data analysis

Images exported from Slidebook 2024 were processed following established
protocols[67]. We used a custom MATLAB toolbox developed by Ben J. Hardcastle
(available at https://github.com/bjhardcastle/SlidebookObj) to correct for motion
artifacts in the x-y plane and to delineate regions of interest (ROI) around individual
LPLC2 neurites within the dendritic tree. For each recording, a time-series was
generated by calculating the mean fluorescence intensity of pixels within the ROI
(Ft) in each frame. These mean values were then normalized to a baseline value
using the formula:

$$\Delta F/F = (Ft - F_0)/F_0$$

1061 where F_0 is the mean of Ft during the 0.5 seconds preceding stimulus onset. This
1062 approach ensures accurate correction for motion artifacts and reliable
1063 quantification of fluorescence intensity changes in LPLC2 neurites.
1064
<u>Statistical analysis of calcium imaging data</u>
Given the repeated sampling and unbalanced sample sizes between groups and
conditions, we employed Linear Mixed Effects (LME) models to fit the DSI values.
This method maintains statistical power by avoiding averaging procedures and
provides more accurate estimates of model parameters, including both fixed and
random effects. The fixed effects were defined by the interaction between
genotype (control/RNAi) and condition (dorsal/ventral), while the random effects
were attributed to individual flies[70,71]. We modeled the data using the R package
`**lme4**` assuming residuals followed a Gaussian distribution. Pairwise post-hoc
comparisons of the fixed effects were conducted using t-tests with Bonferroni
adjustments, implemented through the R package `**emmeans**`. Additionally, to
estimate the mean DSI differences across groups and conditions without assuming
a specific distribution, we performed standard bootstrap simulations with 10,000
replicates using the R package `**boot**`. Smoothed heatmaps were generated with
the R package '**spatstat**'.
1080

Generation of 5xUAS-DIP-ε, -dpr13 and -beat-VI transgenic flies
The coding sequences of DIP-ε, dpr13 (isoform RB), and beat-VI were cloned into
a modified pJFRC5 vector (Addgene: 5XUAS-IVS-mCD8::GFP, plasmid #26218)
by replacing the mCD8::GFP coding sequence. Synthesis and cloning were carried
out by Genewiz, Inc. Plasmids and sequences are available upon request. Flies
were generated by injecting the plasmid into embryos for recombination into attP1
sites (BDSC stock #8621) by BestGene, Inc.
1088

Generation of DIP-ε and dpr13 null alleles
DIP-ε^null allele was generated as previously described[72]. Briefly, two sgRNAs were
used to generate a frame-shift deletion in the DIP-ε coding sequence. High-score
spacer sequences were chosen using the SSC tool[73]. Each sgRNA was cloned
into pU6-2-sgRNA-short (Addgene 41700) plasmid and two plasmids were co-
injected into vas-Cas9 line (BDSC # 51324) by Bestgene, Inc. Injected larvae were
crossed with balancer lines, and PCR-screened in F1 for single flies carrying the
deletion. A mutant stock was established from this single F1.
sgRNA target sequence used for *DIP-ε^null* allele generation:

DIP-ε^null gRNA1: GCTGTTCTGTGGTCATACGATAGC,
DIP-ε^null gRNA2: CTTCAATCGATTGACGGTGGAGC
1100
dpr13^null allele was similarly generated. SgRNA sequences were identified with an
efficiency score above 5, as defined by the CRISPR Efficiency Predictor
(https://www.flyrnai.org/evaluateCrispr/). The sgRNA sequence oligos were
synthesized (Integrated DNA Technologies) and cloned into the pU6b-sgRNA-
short vector[74] to generate a large ~30kb deletion spanning most of the dpr13
genomic region. All pU6 vectors generated were verified by Sanger sequencing.
Two plasmids were co-injected into vas-Cas9 line (BDSC# 51323) in Bestgene,
Inc. Injected larvae were crossed with balancer lines, and PCR-screened in F1 for
single flies carrying the deletion. A mutant stock lacking the entire coding sequence
of dpr13 was established from this single F1.
sgRNA target sequence used for *dpr13^null* allele generation:
dpr13^null gRNA1: CGATATAATCCACTTGATGC,
dpr13^null gRNA2: ACGTAGCAGCTCCAGGATGT.
Detailed protocols are available upon request.
1115

Immunohistochemistry, tissue clearing and DPX mounting

Brains were dissected in ice-cold Schneider's Drosophila Medium (GIBCO
#21720-024) and fixed in PBS containing 4% paraformaldehyde (PFA, Electron
Microscopy Sciences 15710) for 30 minutes at room temperature (RT) or in acid-
free glyoxal (Addax Biosciences) containing 5% sucrose (Sigma S9378) overnight
at 4°C. Brain tissues were washed three times with PBST (PBS containing 0.5%
Triton X-100, Sigma T9284) and incubated in blocking solution (PBST containing
10% Normal Goat Serum, Sigma G6767) for at least 1 hour at RT before incubation
with the primary antibody. Brains were incubated in primary antibody diluted in
blocking solution for two nights at 4°C, washed three times in PBST for 2 hours at
RT, then incubated in secondary antibodies diluted in blocking solution for two
nights at 4°C. Brains were washed three times in PBST for 2 hours at RT.
Subsequent mounting in DPX (EMS 13510) followed the Janelia Adult Drosophila
CNS   DPX   mounting   protocol   (https://www.janelia.org/project-
team/flylight/protocols). After washing from the secondary antibodies, brains were
fixed with PBS containing 4% PFA for 30 minutes at RT, rinsed with PBS, and
mounted on 22x22mm square No.1.5H cover glass (Thorlabs CG15CH2)
previously coated with poly L-Lysine (0.078% solution in deionized water, Sigma
P1524) with added 0.2% Kodak Photo-Flo 200 Solution (Electron Microscopy
Sciences 74257), followed by a quick 1-2 second rinse with MilliQ water. Brains
were dehydrated by placing the cover glass into baths with successively increasing
ethanol (Sigma 459844) concentrations (30-50-75-95-100-100-100%, 10 minutes
each), followed by three successive baths of xylene (Thermo Fisher Scientific X5-
500), 5 minutes each. The glass was uniformly covered with 5-6 drops of DPX
(Electron Microscopy Sciences 13510) and placed on a prepared slide between
spacers made of two 22x22mm square No.2 cover glasses (Fisher Scientific 12-
540B). The slide was left to dry for 24 hours in the hood, then transferred to room
temperature and imaged at least 24 hours later.

Antibody information

Primary antibodies and dilutions used in this study: chicken anti-GFP (1:1000, Abcam #13970), rabbit anti-dsRed (1:200, Clontech #632496), mouse anti-Nc82 (1:20, Developmental Studies Hybridoma Bank (DSHB) Nc82), chicken anti-V5 (1:200, Fortis Life Sciences #A190-118A), mouse anti-V5 (1:500, Abcam #27671), rabbit anti-HA (1:200, Cell Signaling Technology #3724), rabbit anti-FLAG (1:200, Abcam #205606), rat anti-N-Cadherin (1:40, DSHB MNCD2), and anti-GFP nanobody (1:200 for expansion microscopy, 1:500 for confocal microscopy, NanoTag Biotechnologies #N0304-At488-L). Secondary antibodies and dilutions used in this study: goat anti-chicken AF488 (1:500, Invitrogen #A11039), goat anti-mouse AF488 (1:500, Invitrogen #A21131), goat anti-rabbit AF568 (1:500, Invitrogen #A11011), goat anti-mouse AF568 (1:500, Thermo Fisher #A11031), goat anti-mouse AF647 (1:500, Jackson ImmunoResearch #115-607-003), goat anti-mouse CF633 (1:500, Biotium #20250-1), and goat anti-rat AF647 (1:500, Jackson ImmunoResearch #112-605-167).

Confocal image acquisition and processing

Immunofluorescence images were acquired using a Zeiss LSM 880 confocal microscope with 488 nm, 561 nm, and 633 nm lasers using Zen digital imaging software with a Plan-Apochromat 63x/1.4 Oil DIC M27 objective. Serial optical sections were obtained from whole-mount brains with a typical resolution of 1024µm × 1024µm × 0.5µm. Image stacks were exported to either Fiji software or Imaris 10.1 (Oxford Instruments) for level adjustment, cropping, removal of off-target brain regions and background noise, and 3D volume reconstructions.

Analysis of neuroanatomical data from confocal image stacks

To measure the axo-dendritic overlap volume between LPLC2 axons and GF dendrites, confocal image stacks of colocalized LPLC2 glomeruli and GF dendrites were imported into Imaris 10.1 for 3D reconstruction using the Surfaces tool to create masks for membranes of pre- and postsynaptic neurons from the corresponding channels. A Surfaces Detail value of 1 micron was used for both LPLC2 and GF surfaces to ensure accurate reconstruction. Background subtraction was applied with a diameter of the largest sphere that fits into the object set to 1 micron to minimize noise and non-specific signals. For experiments using GF-GAL4, the background subtraction threshold was set to 1 for the GF and 2 for LPLC2, while for experiments using LPLC2-GAL4, the threshold was set to 0.7 for the GF and 2 for LPLC2. The overlap volume between the two reconstructed surfaces was then assessed to quantify the spatial relationship between the LPLC2 axons and GF dendrites. To measure LoP4 dendritic branch length in sparsely labeled LPLC2 neurons, corresponding image stacks were imported into Fiji, rotated so that LoP3 and LoP4 branches were seen as antiparallel, and the distance from the point of bifurcation to the most distal tip of the LoP4 branch was measured along Lop4.

ExLSM and HCR-FISH

Tissue staining, gelation and expansion for ExLSM

Brains were dissected in ice-cold Schneider's Drosophila Medium (GIBCO #21720-024) and fixed in PBS containing 4% paraformaldehyde (PFA, Electron Microscopy Sciences 15710) for 30 minutes on ice. Brains were then washed three times with PBST (PBS containing 0.5% Triton X-100, Sigma T9284) on ice and left in PBST containing anti-GFP nanobody (NanoTag Biotechnologies #N0304-At488-L) overnight at 4°C. All samples were processed using a protein and RNA retention ExM protocol with minor modifications[75] and adjustments for the fly brain[44]. All solutions used for ExLSM and HCR-FISH were prepared with DNase/RNase free distilled water (Thermo Fisher #10-977-023). After staining, brains were washed three times with PBST on ice and left in MOPS buffer3 (20 mM γ-(N-Morpholino) propanesulfonic acid, Fisher Scientific cat# AC327661000, pH 7.7) containing 1:100 AcX stock solution (acryloyl-X, SE Invitrogen, Cat#A20770 at 10 mg/mL in anhydrous DMSO (Sigma-Aldrich, Cat# 276855)) and 1:2 Melphalan X[76] solution (made by combining 2.5 mg/mL Melphalan, Cayman Chemical Company #16665 with 10 mg/mL AcX overnight shaking at room temperature) overnight at 37°C. Brains were then washed three times with 1xPBS. A gelation chamber was created by applying a Press-to-Seal silicone gasket (Invitrogen, Cat# P24740) to a glass slide, which was then coated with the PLL solution. AcX-treated brains were immobilized on the PLL surface, up to nine per gasket. Gelation solution was prepared just before gelation to prevent premature gel polymerization on ice by adding 10 µL each of 4-hydroxy-TEMPO at 0.5% (w/v) (Sigma-Aldrich, Cat# 176141), TEMED (N,N,N′,N′-Tetramethylethylenediamine at 10% (v/v) (Sigma-Aldrich, Cat# T7024), and ammonium persulfate at 10% (w/v) (Sigma-Aldrich, Cat# A3678) to 470 µL of monomer stock solution (11.5 mL sodium acrylate stock at 4M, prepared by neutralizing 5.5 mL acrylic acid (99% purity; Sigma-Aldrich, Cat# 147230) with 10N NaOH using a water bath and fume hood, in a total volume of 20 mL; 2.5 mL acrylamide stock at 50% (w/v), Sigma-Aldrich, Cat# A9099; 7.5 mL bis-acrylamide stock at 1% (w/v), Sigma-Aldrich, Cat# M7279; 18 mL 5 M NaCl, Sigma-Aldrich, Cat# S5150). Brains were washed with gelation solution, and then the gelation chamber was filled with ~200 µL of gelation solution and incubated on ice for 25 minutes. The gelation chamber was then sealed with a cover slip and placed in a 37°C incubator to gel and cure for 2 hours. Gelation chambers were disassembled, and individual gels were trimmed close to each brain. Gels were trimmed to a right trapezoid shape to ease specimen orientation. Gels were transferred into 12-well plates (Laguna Scientific #4624-12) and incubated with proteinase K (NEB, Cat# P8107S) diluted 1:100 (v/v) in proteinase K digestion buffer (0.5% Triton X-100, 50 mM NaCl, 1 mM EDTA, and 50 mM Tris pH 8) with shaking overnight at room temperature. This resulted in ~2x gel expansion.

HCR-FISH

The HCR-FISH protocol was adapted from Wang et al[76] with minor optimizations for the fly brain[44]. Following digestion with proteinase K, gels with embedded brains were washed 3 times with PBS, transferred into 24-well plates (Laguna Scientific #4624-24), and digested with DNase diluted

in RDD buffer (RNase-Free DNase Set, Qiagen #79254) to limit DAPI signal
to RNA only and facilitate subsequent analysis for 2 hours at 37°C. After
three washes in PBS, gels were equilibrated in Probe Hybridization Buffer
(Molecular Instruments, Inc.) for 30 minutes at 37°C, and then transferred to
new 24-well plates containing custom-designed probes (Molecular
Instruments, Inc.) diluted in pre-warmed Probe Hybridization Buffer (1 µL of
1 µM stock probe solution per 200 µL of buffer) and left shaking overnight at
37°C. The following day, gels were washed 4 times with pre-warmed Probe
Wash Buffer (Molecular Instruments, Inc.) for 20 minutes at 37°C, then
washed twice for 5 minutes with SSCT buffer (SSC, Thermo Fisher
#AM9763 with 0.05% Triton X-100) at room temperature and transferred to
new 24-well plates with HCR Amplification buffer (Molecular Instruments,
Inc.) for equilibration. Hairpins (HCR Amplifiers, Molecular Instruments, Inc.)
conjugated with AF546 or SeTau647 dyes were diluted in Amplification
Buffer (2 µL of each hairpin per 100 µL of buffer), heat-activated in a thermal
cycler (90 seconds at 95°C), removed, and kept for 30 minutes at room
temperature in the dark. After 30 minutes, the hairpins were added to the
24-well plates with gels (300 µL per well) and incubated with shaking at room
temperature in the dark for 3 hours. The hairpin solution was then removed,
and the gels were washed 4 times with SSCT and 2 times with SSC for 10
minutes at RT in the dark. Gels were subsequently stored at 4°C in SSC
until final expansion.
1258
<u>Sample mounting</u>
Samples were expanded to ~3X in 0.5x PBS containing 1:1000 SYTO-DAPI
(Thermo Fisher S11352) at room temperature for 2 hours before mounting onto
PLL-coated coverslips (see description for DPX mounting above). The coverslips
were then bonded with Bondic UV-curing adhesive (Bondic starter kit, Bondic) onto
a custom fabricated sample holder (Janelia Tech ID 2021-021) to be suspended
vertically in the imaging chamber. Mounted samples were imaged in 0.5x PBS with
1:10,000 SYTO-DAPI after a minimum of 1 hour of equilibration in the imaging
chamber. Unexpanded gels were stored at 4°C in 1X PBS + 0.02% sodium azide
(Sigma-Aldrich, Cat# S8032) for up to 14 days before final expansion and imaging.
1269
<u>Light-sheet microscopy</u>
Images were acquired on a Zeiss LS7 microscope equipped with 405 nm, 488 nm,
561 nm, and 638 nm lasers. Illumination optics with a 10x/0.2 NA were used for
excitation (Zeiss, Cat# 400900-9020-000). Detection was performed using a W
Plan-Apochromat 20x/1.0 DIC M27 water immersion objective (Zeiss, Cat#
421452-9700-000). The LS7 optical zoom was set to 2.5x, resulting in a total
magnification of 50x. DAPI and AF546 dyes were simultaneously excited by the
405 nm and 561 nm laser lines, and emission light was separated by a dichroic
mirror SBS LP 510 with emission filters BP 420-470 (Zeiss, Cat# 404900-9312-
000) and a modified BP 527/23 (Chroma, Cat# ET672/23m). Similarly, AF488 and

SeTau647 dyes were simultaneously excited via 488 nm and 638 nm, and the emission was split through a dichroic SBS LP 560 with emission filters BP 505-545 and LP 660 (Zeiss, Cat# 404900-9318-000). To eliminate laser transmission, a 405/488/561/640 laser blocking filter (Zeiss, Cat# 404900-9101-000) was added to the emission path. Images were captured using dual PCO.edge 4.2 detection modules (Zeiss, Cat# 400100-9060-000) with a 50 msec exposure time. Filter and camera alignment were manually calibrated prior to each imaging session. Image volumes were acquired at optimal Z-step and light-sheet thickness, and the Pivot Scan feature was used to reduce illumination artifacts by sweeping the light-sheet in the xy-plane. The LS7 microscope was operated using ZEN Black 3.1 (v9.3.6.393).

Analysis of HCR-FISH data from ExLSM Image Stacks

The full details of our analysis are available in our previous publication[45]. The acquired z-stacks, stored in CZI format, were imported and pre-processed to remove noise and artifacts generated by the imaging modality. These artifacts included limited channel contrast, variations in contrast from image to image within a dataset, background noise fluctuations due to both intra-channel variations and inter-channel crosstalk, and localized brightness changes from varying fluorophore concentrations within and among stained nuclei. Pre-processing involved the following steps: full-scale contrast stretching (FSCS) to normalize luminosity across different channels, local background removal using a 3D Gaussian filter, a second FSCS to adjust for any contrast loss due to background removal, and a final median filter to remove any remaining localized noise. These pre-processed stacks served as the starting point for instance segmentation of the nuclei. First, the nuclei centers were identified using a Laplacian-of-Gaussian (LoG) filter. Then, the imaged volume was subdivided into 3D Voronoi cells, using the detected centers as seeds and Euclidean distance. Each cell contained one nucleus, which was segmented using a threshold obtained by minimizing an energy functional designed to find the optimal surface separating the nucleus from the surrounding cytoplasm. Once each nucleus was segmented, the FISH puncta were identified within the 3D volume associated with it using a LoG filter, and only the puncta within the nucleus region and its immediate surrounding volume were counted. Pre-processed products, segmented nuclei features, associated FISH puncta, and their features were stored for further analysis. Puncta counts were normalized by the maximum count for each brain. The significance of gene expression relationships inferred from HCR-FISH (Fig. 2) was assessed using both a linear regression model and a multi-level negative binomial generalized linear mixed model, accounting for inter-animal heterogeneity: random effects were attributed to individual animals (Supplemental Table 2). The package was written in Python, is open-source, and available for download at https://github.com/avaccari/DrosophilaFISH.

Genetic Intersection of cell types and genes

To assess gene expression levels within LPLC2 and LPLC1 cell types, we used a combination of GAL4 and LexA binary expression systems. Expression of a

1326 LexAop-FRT-stop-FRT-membrane marker controlled by a cell-type-specific LexA
1327 driver was blocked until the FRT-stop-FRT cassette was excised by a Flippase,
1328 driven by a gene-specific T2A-GAL4 driver line[77]. Additionally, an unconditional
1329 membrane marker controlled by the same LexA driver visualized the entire cell
1330 population. The list of fly stocks used is available in Supplemental Table 1.
1331

Sparse labeling of neuronal populations

To visualize single cell morphology of LPLC2 dendrites in the lobula plate
(Fig 4) and for dorsoventral HCR-FISH analysis (Ext. Data Fig. 3), we used
MultiColor FlpOut (MCFO)[78], a genetic tool for sparse labeling individual cells
within a neuronal population acting downstream of a given Gal4 driver. 24h pupae
were heat shocked in a 37°C water bath for ten minutes to achieve the labeling of
one LPLC2 neuron per hemibrain om average.
1339

Analysis of the Flywire Connectome reconstruction

To analyze the *Flywire* connectome, we developed an open-source Python
package, available at https://github.com/avaccari/DrosophilaCon. The primary
library used by *DrosophilaCon* to interface with the *FlyWire* connectome is
fafbseg[32,33] (version 3.0.0). This package enables users to specify the labels of
"source" and "target" neurons and generates a connectivity diagram where target
neurons are color-coded based on the total count of synapses with the source
neurons. Once the labels are specified, the package queries the latest available
annotations to identify all neurons matching (or containing) the labels. The primary
sources of annotation used to identify the neurons are:
(1) Free-form community annotations provided through the *neuroglancer* user
interface (https://github.com/google/neuroglancer).
(2) Systematic annotations for the entire brain[33].
(3) Systematic cell types for the right optic lobe[79].
Next, the latest version of the *FlyWire* production dataset is queried for the
adjacency matrix representing the connectivity between each neuron in the source
and each neuron in the target for the selected side of the brain. This information is
returned as an adjacency table, providing the counts of synapses between each
source-target pair. There are two versions of the synapse datasets: one filtered by
synaptic cleft and one unfiltered. We used the unfiltered dataset because the
filtered version applied a fixed threshold for distance, resulting in reduced synapse
counts. The adjacency table was used to evaluate the total synapse counts for
each target neuron. These counts were normalized by the maximum count
observed across all target neurons. The mesh representation for each identified
target neuron was downloaded from the *FlyWire* dataset, skeletonized to optimize
processing, and displayed with color-coding corresponding to the normalized
synapse count, allowing for comparison across different source-target pairs.
1367

Statistical analysis

1369 All statistical analyses were performed using RStudio 1.4.1103, MATLAB,
1370 or Prism 9.2.0 (GraphPad). Significance levels were defined as follows: $p < 0.05$,
1371 *$p < 0.01$, **$p < 0.001$, ***$p < 0.0001$ for all figures. Statistical tests were chosen

based on data distribution, which was assessed using the Kolmogorov-Smirnov test in R with a p-value threshold of <0.05 for normality. Two groups of normally distributed data sets were tested for statistically significant differences using unpaired t-tests with Welch's correction for non-identical variance. For comparisons involving more than two groups, we employed either one-way ANOVA followed by Tukey's HSD test for post-hoc pairwise comparisons or the Kruskal-Wallis test followed by Dunn's multiple comparisons post-hoc test with Bonferroni correction. Binary data were compared using chi-squared tests. Detailed statistical analyses for behavioral data, HCR-FISH data, and neuroanatomical data are described in Supplementary Table 2. All other statistical tests, number of replicates, significance levels, and other elements of the statistical analysis are reported within the corresponding Methods and Results sections and/or Figure Legends. No data were excluded from the analysis unless specified in the corresponding Methods section. All measurements were taken from distinct samples.

Data availability

The datasets generated and/or analyzed during the current study are available from the corresponding authors upon reasonable request. For further information regarding any resources and reagents, please contact S. Lawrence Zipursky (lzipursky@mednet.ucla.edu) and Gwyneth M. Card (gc3017@columbia.edu). Source Data are provided with this paper.

Code availability

The code for single-cell RNA-seq analysis is available at https://github.com/kurmangaliyev-lab. Our code for the analysis of Flywire and "hemibrain" connectome reconstructions is available at https://github.com/avaccari/DrosophilaCon and https://github.com/avaccari/DrosophilaVPNWiring, respectively. Code used to analyze HCR-FISH data is available at https://github.com/avaccari/DrosophilaFISH. Code related to calcium imaging experiments is available at https://osf.io/z7xfk/. All other code can be made available from the corresponding authors upon request.

Acknowledgements

We thank the Bloomington Stock Center (NIH P40OD018537), Vienna *Drosophila* Research Center and Janelia Fly Facility for providing fly stocks. We also thank the UCLA BSCRC Sequencing Core and the UCLA TCGB core for help with library preparation and scRNA-seq; the UCLA BSCRC FACS core for assistance with FACS purification. We thank members of the Flywire Consortium for connectome annotation in the FAFB dataset. We thank all members of the Zipursky lab for advice and comments on the manuscript. This work was supported by the HHMI-Helen Hay Whitney Foundation (M.D.), NEI K99EY036123 (M.D.), NEI R01EY026031 (M.F.), and NINDS R01NS118562 (C.R.v.R) S. L. Z. and G.M.C. are investigators of the Howard Hughes Medical Institute.

Author Contributions

Conceptualization and Methodology: M.D., Y.Z.K, C.R.v.R., M.A.F., G.M.C., S.L.Z.; Investigation & Data collection: M.D., Y.Z., G.F., H.J., Parmis S. Mirshahidi, A.R., R.H.H., C.L. and Pegah S. Mirshahidi; Data analysis: M.D., Y.Z.K., A.V., Y.Z., G.F., H.J., F.X., B.W.H.; Writing, reviewing and editing.; M.D., Y.Z.K., Y.Z., G.F., H.J., C.R.v.R., G.M.C. and S.L.Z.; Supervision and funding acquisition: S.L.Z, G.M.C., C.R.v.R., M.A.F., M.D.

Ethics Declarations
Competing interests
The authors declare no competing interests

Figure 1

Fig. 1: Gradients of recognition molecules correlate with synaptic gradients and behavior in Visual Projection Neurons

a-d, Anatomy

a, Visual Projection Neurons (VPN) have dendrites that cover the lobula (Lo), lobula plate (LoP) or both neuropils of the fly optic lobe. Axons converge on optic glomeruli in the central brain innervated by dendrites of target neurons, e.g., Descending Neurons (DN); D-dorsal, V-ventral, L-lateral.

b, Confocal projection (posterior view) of LPLC2 VPNs and the Giant Fiber DN.

c, Top: EM-based reconstructions of LPLC2 neurons (posterior view) from the "hemibrain" volume, color-coded by the number of synapses their axons make onto the GF. Bottom: linear regression of LPLC2-GF synapse numbers as a function of LPLC2 dendrite position along the DV axis of the visual space. Dots, individual LPLC2 neurons (colored as above). Shaded region, 95% confidence interval).

d, same as (**e**, top), inferred from the Flywire volume (left and right sides)

e, **Behavior**: GF-mediated short-mode (<7 ms sequence duration) takeoffs in response to frontal (0°) and lateral (90°) looms at various elevations. Error bars, 95% confidence intervals.

f, **Summary**: Transformation of retinotopic dendritic maps into axonal synaptic gradients between LPLC2 and the GF without axonal retinotopy.

g-l, **Identification of gene expression gradients in VPNs**.

g, Single-cell RNA-sequencing experimental design (See Methods).

h-i, t-SNE plots of 48h APF dataset (**h**) LPLC2, LPLC1, and LC4 neurons were annotated based on the expression of known marker genes (**i**). X, unknown ectopic cell type.

j, Heatmaps of the expression patterns of the top 30 genes with the highest contribution (loading) to Principal Component 1 (PC1) across LPLC2 neurons at 48h APF (PCA was performed separately on LPLC2 neurons); genes (rows) ordered by their loadings to PC1; cells (columns) ordered by their PC1 scores; expression patterns colored by scaled expression levels (left) and original log-normalized expression values (right).

k, Distributions of LPLC2 neurons at 48h APF along PC1 and PC2 (PCA plots) colored by developmental age, genotype, sex (male-specific transcript), and coverage.

l, PCA plots of LPLC2 neurons at 48, 72, and 96h APF, colored by the expression levels of the select genes from (**j**). Differential expression of these genes is consistent over time.

Figure 2

a LPLC2>GFP.NLS (green), dpr13 mRNA (blue), SiaT mRNA (red), DAPI (all RNA) (gray). D, L. 3x tissue expansion. Scale bar 100μm.

b dpr13 (blue), SiaT (red). r=-0.42. Axes: log(TP10k+1) vs PC_1.

c high dpr13 low SiaT (blue arrowheads), high SiaT low dpr13 (red arrowheads).

d scRNA-seq and FISH panels. SiaT vs dpr13: r = -0.42 (scRNA-seq), r=-0.41 (FISH). beat-VI vs dpr13: r = 0.59 (scRNA-seq), r=-0.48 (FISH). Genotype: 399, 535, 555, 730, 799, 805. Brain: 1-9.

e Spearman's r at 48h APF, 72h APF, 96h APF. scRNA-seq (blue), FISH (red).

48h APF: n=865, 534, 395, 360, 364, 449, 448, 526, 532, 540, 306, 548.
72h APF: n=589, 331, 279, 371, 324, 407, 356, 383, 421, 347, 439.
96h APF: n=436, 358, 532, 354, 415, 363, 430, 341, 539, 460, 465.

Categories: dpr13/beat-VI, fz2/beat-VI, tail/dpr13, nAChRd5/beat-VI, dpr17/dpr13, beat-VI/dpr17, SiaT/dpr13, beat-VI/SiaT, SiaT/Cad87A, CG30419/beat-VI, CG30419/CG30419, dpr13/CG30419, mbl/dpr13.

f Genetic intersection (□) of cell types and genes. LPLC2-LexA, LexAop, FRT, STOP, FRT, myr::sm-HA/V5. Gene-of-interest GAL4, UAS FLP, FLP. Gene of interest ∩ Cell Type.

g positive control. All LPLC2 (LexA), □ LPLC2-GAL4, merge. D, M. Scale bar 20μm. N=5 brains.

h RNA-seq. dpr13 (blue), beat-VI (red): r=0.59. dpr17 (blue), SiaT (red): r=-0.3. Axes: log(TP10k+1) vs PC 1.

i "dorsal" genes and "ventral" genes.
All LPLC2 (green), LPLC2 □ dpr13 (red), side view. NCad. 48h APF. D, M. D, A. N=11 brains. Scale bar 10μm.
All LPLC2, LPLC2 □ SiaT, side view. N=10.
All LPLC2, LPLC2 □ beat-VI, side view. N=8.
All LPLC2, LPLC2 □ dpr17, side view. N=7.

Fig. 2: Gradients of recognition molecules align with synaptic gradients

a-e, FISH analysis.

a, Light sheet projection of the expanded *Drosophila* optic lobe (48h APF) with labeled LPLC2 nuclei and transcripts of dpr13 and SiaT.

b, Anti-parallel expression of dpr13 and SiaT across LPLC2 neurons at 48h APF (inferred from scRNAseq, along PC1). Fitted lines, trends of gene expression changes along PC1; shaded areas, 95% confidence intervals. r, Spearman's rank correlation coefficient.

c, Single slice (0.5μm thick) from the projection in (**a**) (zoomed into the dashed rectangular region); arrows indicate individual LPLC2 neurons expressing markedly different levels of dpr13 and SiaT.

d, Comparison of scRNA-seq (left) and FISH (right) measuring correlation in expression for two pairs of graded genes: dpr13/SiaT (top) and dpr13/beat-VI (bottom) across LPLC2 neurons at 48h APF. Fitted lines, linear relationship between the expression levels of two genes for each brain (FISH) or each wild-type genotype (scRNA-seq). Shaded regions, 95% confidence intervals.

e, Comparison of RNA-seq and HCR-FISH measuring correlation in expression for twelve pairs of genes at three time points across LPLC2 neurons. Individual dots represent Spearman's rank correlation coefficients for each brain (FISH) and each genotype (scRNA-seq). Error bars, means ±95% confidence intervals; n, total number of neurons tested using FISH.

f-i, GAL4-reporter analysis.

f, Genetic approach to visualize a subset of neurons within a VPN cell type population expressing a specific gene at a particular time point.

g, Positive control for (**f**). LPLC2-LexA genetically intersected with LPLC2-GAL4, results in labeling of all LPLC2 neurons.

h, Correlation between expression levels of dpr13/beat-VI and dpr17/SiaT (inferred from scRNA-seq), along PC1 across the LPLC2 population at 48h APF.

i, Subsets of LPLC2 neurons expressing dpr13, dpr17, beat-VI and SiaT at 48h APF. D, dorsal; M, medial; A, anterior.

Figure 3

Fig. 3: A gradient of DIP-ε::Dpr13 recognition molecules controls a looming escape synaptic gradient in LPLC2 neurons

a, Protein interaction map showing molecular binding partners of most differentially expressed recognition molecules in LPLC2.

b, Light sheet projections of the GF (nuclear marker). Red puncta, mRNA of designated genes assessed through HCR-FISH.

c, Candidate gene expression levels (FISH puncta counts) in the GF throughout development. Error bars, means with ±95% confidence intervals.

d, Confocal projections of LPLC2 axon terminals and the GF dendrites in wild type and *DIP-ε^null^* animals throughout development. M, medial; D, dorsal.

e-g, Comparison of LPLC2-GF axo-dendritic overlap volume (see Methods) between controls and in *DIP-ε^null^* animals (**e**), DIP-ε rescue in the GF (**f**), and GF expressing two different DIP-ε RNAi (**g**).

h, Whole-cell patch-clamp recordings in the GF during presentations of looming stimuli. Left: GF responses to looming stimuli, r/v = 40ms. Control (average: black, individual fly: grey) and *DIP-ε^null^* (average: orange, individual fly: light orange) traces are overlayed. Looming stimulus profile over time is displayed below the GF responses. Right: Quantification of expansion peak amplitudes from individual flies; each data point represents the mean value of two recordings per animal.

i, Same as (**h**) for controls (grey) and animals overexpressing DIP-ε RNAi in the GF (light blue).

j, Top: Violin plots of takeoff sequence durations for lateral stimuli (90°) at different elevations (0°, 45°, 77°) in wild-type and *DIP-ε^null^* animals. Lines, single takeoffs. Short-mode (<7 ms, grey for wild-type, orange for *DIP-ε^null^*) and long-mode (≥7 ms, grey) durations separated by a red dashed line. N = number of flies. Bottom: short-mode takeoff percentages at different elevations. Error bars, 95% confidence intervals. Dashed lines, wild-type DV gradient (grey) and eliminated DV gradient in *DIP-ε^null^* (orange).

k, Same as (**j**) for controls (grey) and animals overexpressing DIP-ε RNAi in the GF (light blue). .

l, Same as (**j**) for controls (grey) and animals overexpressing dpr13 in LPLC2 neurons (green).

m, Comparison of LPLC2-GF axo-dendritic overlap volume in control animals, and animals expressing overexpressing dpr13 in LPLC2

n, Same as (**m**), for control animals, and animals overexpressing DIP-ε in LPLC2.

panels **e**, **f**, **g**, **m**, **n**: dots, individual brains. Error bars, means ±95% confidence intervals.

Figure 4

a LPLC2 dendrites omnidirectionally detect motion

nc82   LPLC2 population (~100 cells)

nc82   Single LPLC2 neuron

Lo dendrites   LoP dendrites

b Dendritic Synaptic Gradients

T5c>LPLC2   T5d>LPLC2

synapses

T4c>LPLC2   T4d>LPLC2

c LPLC2>control   LPLC2>beat-VI RNAi

dorsal   N=12 neurons   10μm

central   N=11   N=13

ventral   N=9   N=9

d
control
beat-VI RNAi

LoP 4 branch length, μm

dorsal ****   central ****   ventral n.s.

e T4/T5>side-II RNAi

N=6   dorsal

N=6 neurons   ventral

f
920 nm

g
45°
165°
90°
20 °s⁻¹

positions mapped onto fly eye

elevation   azimuth

h putative neurons corresponding to two grid positions

LPLC2 (bodyID: 28871)   25 μm

LPLC2 (bodyID: 30207)

i
16   dorsal LPLC2
5% ΔF/F
40   ventral LPLC2

j
LPLC2 > control   LPLC2 > beat-VI RNAi

Directional Selectivity Index

***   n.s.

N=7 flies   N=4   N=9   N=4
dorsal   ventral   dorsal   ventral

k
LPLC2 > control

LPLC2 > beat-VI RNAi

directional sensitivity
downward   upward

l
LPLC2 > control
n.s.
Peak ΔF/F (%)

LPLC2 > beat-VI RNAi
n.s.
dorsal   ventral

Fig. 4: A gradient of Beat-VI::Side-II recognition molecules controls a motion detection synaptic gradient in LPLC2 neurons

a-b, LPLC2 dendritic synaptic gradients

a, Confocal projections of the LPLC2 population (top) and an individual LPLC2 neuron (bottom) in dorsal view, highlighting the dendritic branches in the lobula plate (LoP). Numbers, LoP layers. Inset: posterior view of a single LPLC2 LoP dendrite with branches extending into only one of the four cardinal directions in each layer. D, dorsal; V, ventral; M, medial; P, posterior.

b, EM-based reconstructions (Flywire connectome) of LPLC2 neurons (posterior view), color-coded by the number of inputs each LPLC2 neuron receives from T4c, T4d, T5c and T5d neurons. Insets, schematics of single LPLC2 (red) and T4/T5c-d neurons, overlaid on lobula/lobula plate outlines.

c-e, beat-VI differentially regulates LPLC2 dendrites along the LoP4 DV axis.

c, Confocal projections of individual LoP dendrites (posterior view) in dorsal, central and ventral LPLC2 in control and LPLC2>beat-VI RNAi flies. Numbers, LoP layers. Dashed ovals, LoP4 dendritic branches.

d, Length of LoP4 dendritic branches for dorsal, central and ventral LPLC2 neurons in control and LPLC2>beat-VI RNAi flies. Dots, individual neurons. Error bars, means ±95% confidence intervals.

e, Confocal projections of individual LoP dendrites (posterior view) of dorsal and ventral LPLC2 neurons in control and T4/T5>side-II RNAi flies. Dashed ovals, LoP4 dendritic branches.

f-l, beat-VI differentially regulates integration of motion by LPLC2 neurons

f, Schematic of the fly eye relative to the surrounding LED display for visual stimulation during two-photon calcium imaging.

g, Positions on the LED display probing LPLC2 receptive field by presentation of dark edges moving in 24 (0° to 345°) different orientations (top). Yellow line, fly eye's equator (eq) projected onto the LED display. Mollweide projection of the outermost positions (black boxes) relative to the fly eye during Ca2+ imaging experiments (bottom).

h, EM-based reconstructions of LPLC2 neurons stimulated by grid positions 16 and 40 (see Methods). LPLC2 neurites viewed during Ca2+ imaging with single-cell regions of interest (color-coded by positions) overlaid on them (insets, scale bar: 10 μm).

i, Polar plots of the peak responses to moving dark edges in dorsal and ventral LPLC2 neurons in a representative fly.

j, Directional sensitivity index (DSI) for dorsal and ventral LPLC2 neurons in control and beat-VI RNAi flies. Dots, individual recordings. Error bars, means ±95% confidence intervals.

k, Filtered heatmap of the DSI for all positions tested in control and beat-VI RNAi flies.

l, Average peak responses to a looming stimulus above (dorsal) or below (ventral) the eye's equator in control and beat-VI RNAi flies. Error bars, means ±95% confidence intervals.

Extended Data Figure 1 (related to Fig. 1)

Extended Data Figure 1 (related to Fig. 1). Synaptic gradients are associated with molecular heterogeneity across VPN cell types

a, Histograms showing the distribution of takeoff sequence durations in response to frontal (left) and lateral (right) looming stimuli at different elevations (detailed breakdown of data summarized in Fig. 1e). Short-mode and long-mode takeoffs are distinguished by the red dashed lines.

b, t-SNE plots of 72h and 96h APF datasets (left). LPLC2, LPLC1, and LC4 neurons were annotated based on the expression of known transcription factors (right). X indicates unknown ectopic cell types.

c, Heatmaps of the expression patterns of the top 30 genes with the highest contribution (loading) to differentially expressed genes along Principal Component 1 (PC1) across LPLC1 neurons at 48h APF (see Fig. 1 legend for details). Genes encoding cell recognition molecules (IgSF superfamily) are highlighted in bold.

d, PCA plots of LPLC1 neurons at 48, 72, and 96h APF, colored by the expression levels of two cell recognition molecules from (**c**).

e, PCA plots of LPLC2, LPLC1, and LC4 neurons at 48h APF colored by genotype, age (early vs late collection), genotype (DGRP line), sex (male-specific transcript), and coverage. PC1-6 for LPLC1 and LC4, and PC3-6 for LPLC2 are shown.

Extended Data Figure 2 (related to Fig. 1)

Extended Data Figure 2 (related to Fig. 1). Further analysis of within-cell-type heterogeneity in gene expression across LPLC2, LPLC1 and LC4

a-b, PCA plots of LPLC2, LPLC1 and LC4 neurons at 72h APF (**a**) and 96h APF (**b**) colored by genotype, age (early vs late collection), genotype (DGRP line), sex (male-specific transcript), and coverage. Shown are PC1-6.

c-e, Scatter plots of LPLC2 cells (48h APF) embedded in the first two principal components (PC1 and 2) colored by their cluster labels (n=2, inferred from K-means clustering). PCs are calculated using the top 1000 highly variable genes based on the actual data (**c**), after shuffling gene expression levels for each gene independently across all cells (**d**), and after shuffling each gene only within each cluster (**e**), respectively. Shuffling gene expression across all cells (**d**) disrupts this gradient, indicating that the observed differences reflect coordinated gene expression rather than uncorrelated variation. Shuffling within each cluster (**e**) disrupts the internal gradient of each cluster, creating artificial gaps not present in the original data (**c**), which suggests that the continuous gradient observed in the original data cannot be represented as a set of discrete clusters.

f, Scatter plots of cells in PCA embeddings after shuffling genes within each cluster for an increasing number of fine-grained clusters (n=2,5,30).

g, Gap size as a function of the number of clusters. Gap size is defined as the minimum distance in PC1 and PC2 space needed to connect 90% of cells into a single graph component. The gap size decreases and plateaus as the number of clusters increases. This means that a continuum can be approximated by a large number of discrete clusters. However, a small number of clusters (like 2) is insufficient to capture the continuous nature of the data.

Extended Data Figure 3 (related to Fig. 2)

a LPLC2>GFP.NLS (green), dpr17 mRNA (cyan), Beat-VI mRNA (red), DAPI (all RNA) (grey), 3x tissue expansion, 48h APF, 50µm, D/L

high dpr17 low beat-VI, high beat-VI low dpr17, 10 µm

b FISH
n=429 cells
N=7 brains
r=-0.25
Dpr17, norm puncta count vs Beat−VI, norm puncta count
Brain: 1, 2, 3, 4, 5, 6, 7

scRNA-seq
n=1445 cells
N=90 brains
r=-0.36
dpr17, log2(TP10K+1) vs beat-VI, log2(TP10K+1)
Genotype: line_399, line_535, line_555, line_730, line_799, line_805

c dpr13 mRNA (cyan), LPLC2 (green), 20µm, D/M
Dpr13, puncta #: Dorsal LPLC2, Ventral LPLC2
Dorsal 1: 43 puncta, 5µm
Central 2: 37 puncta
Ventral 3: 14 puncta

d beat-VI mRNA (cyan), LPLC2 (green), 20µm
Beat-VI, puncta #: Dorsal LPLC2, Ventral LPLC2
Dorsal 1: 52 puncta, 5µm
Central 2: 37 puncta
Ventral 3: 19 puncta

Extended Data Figure 3 (related to Fig. 2). HCR-FISH validates gradients of cell recognition molecules in LPLC2.

a, Light sheet projection of the expanded *Drosophila* optic lobe (48h APF) with labeled LPLC2 nuclei and transcripts of dpr17 and beat-VI (top) and a single slice (0.5μm) from the light sheet projection (bottom, zoomed into the yellow dashed rectangular region); arrows indicate individual LPLC2 somas expressing markedly different levels of dpr17 and beat-VI. D, dorsal; L, lateral.

b, Comparison of scRNA-seq (top) and HCR-FISH (bottom) measuring the correlation in expression for dpr17 and beat-VI across LPLC2 neurons at 48h APF. Fitted lines represent the linear relationship between the two genes for each brain (FISH) or each wild type genotype (scRNA-seq); shaded regions represent 95% confidence intervals. r, Spearman's rank correlation coefficient.

c, dpr13 expression levels using HCR-FISH in individual sparsely labeled LPLC2 neurons. Right, zoomed-in cell bodies of dorsal, ventral, and central LPLC2 neurons, indicating the number of dpr13 transcripts per cell. Inset, comparison of dpr13 puncta counts in dorsal and ventral LPLC2 neurons. Dots represent individual neurons (1 neuron per brain). Error bars represent means ±95% confidence intervals.

d, Same as in (**c**), for beat-VI

Extended Data Figure 4 (related to Fig. 2)

a More examples of "dorsal" genes

All LPLC2 · LPLC2 · Cad87A-GAL4
NCad · 48h APF · 20 µm · D L · N=4

All LPLC2 · LPLC2 · CG44422-GAL4
48h APF · N=4

b
Cad87A · dpr13 · r=0.49
log(TP10K+1) · PC_1

CG44422 · dpr13 · r=0.5
log(TP10K+1) · PC_1

c More examples of "ventral" genes

LPLC2 · StacI-GAL4 · LPLC2 · Tsp42Ef-GAL4 · LPLC2 · CG30419-GAL4
NCad · 48h APF · D A
nc82 · 96h APF

d
SiaT · StacI · r=0.31
SiaT · Tsp42Ef · r=0.35
CG30419 · SiaT · r=0.36
log(TP10K+1) · PC_1

e
48h APF (side view) · 48h APF · 72h APF · 96h APF
All LPLC1 · NCad · nc82

LPLC1 · DIP-kappa · D L · N=9 · 20 µm · N=7 · N=6

LPLC1 · sdk · N=8 · N=9 · N=5

LPLC1 · dpr17 · N=4 · N=3 · N=3

D A

f
DIP-kappa · sdk · r=0.55
DIP-kappa · dpr17 · r=-0.33
log(TP10K+1) · PC_1

Extended Data Figure 4 (related to Fig. 2). Retinotopic correlates of molecular gradients in LPLC2 and LPLC1

a, Subsets of LPLC2 neurons expressing CG44422 and Cad87A at 48h APF. White arrows indicate ventral regions of the lobula lacking expression of both CG44422 and Cad87A. Dashed ovals, partial overlap of expression in somas, i.e., different LPLC2 neurons express different levels of CG44422 and Cad87A. N, number of brains tested. D, dorsal; L, lateral.

b, Positive correlation between expression levels of Cad87A, CG44422 and dpr13 (inferred from scRNA-seq), along PC1 across the LPLC2 population at 48h APF, indicating that both CG44422 and Cad87A can be considered "dorsal" genes.

c, Additional examples of genes expressed by ventral subpopulation of LPLC2 neurons. Red, dendrites of LPLC2 (lateral view of the lobula) that express stacl, Tsp42Ef and CG30419 at 48h (top) and 96h (bottom) APF. D, dorsal; P, posterior.

d, Positive correlation between expression levels of stacl, Tsp42Ef and CG30419 (inferred from scRNA-seq), along PC1 across the LPLC2 population at 48h APF.

e, Retinotopically biased gene expression across LPLC1 neurons. Red, subsets of LPLC1 neurons expressing DIP-kappa, sdk and dpr17 throughout development (posterior and lateral views of the lobula). N, number of brains tested. D, dorsal; M, medial; P, posterior.

f, Positive correlation between expression levels of DIP-kappa and sdk (top), and negative correlation between expression levels of DIP-kappa and dpr17 (bottom), (inferred from scRNA-seq), along PC1 across the LPLC1 population at 48h APF, reflecting the retinotopically biased expression of these genes in (**e**).

Extended Data Figure 5 (related to Fig. 3). Synaptic gradient between LPLC2 and the GF is established through a gradient of DIP-ε::Dpr13 molecular interactions.

a, To establish a synaptic gradient with LPLC2 based on dorsoventral expression gradient of any of the candidate molecules (Dpr13, Beat-VI, Dpr17), the GF needs to express a molecular binding partner to recognize one or more of these molecules.

b, Validation of VT049479-GAL4 expression in LPLC2 using GMR75G12-LexA as a reference. A complete overlap confirms that VT049479-GAL4 targets the entire LPLC2 population. D, dorsal; M, medial.

c, Confocal projections of LPLC2 axon terminals and Giant Fiber dendrites in wild-type animals, as well as animals expressing control RNAi and two different DIP-ε RNAi in the Giant Fiber.

d, Same as (**c**). Rescue experiment: overexpression of DIP-ε cDNA in the GF in *DIP-ε^{null}* background (right) compared to two negative controls (left and middle).

e, Confocal projections of LPLC2 axon terminals and the Giant Fiber dendrites in control animals and animals overexpressing dpr13 in LPLC2

f, Comparison of dpr13 mRNA levels in LPLC2 (assessed using HCR-FISH) in control animals and animals overexpressing dpr13. Dots, individual brains. Error bars, means ±95% confidence intervals.

g-h, Confocal projections of LPLC2 axon terminals and the Giant Fiber dendrites in wild type and *dpr13^{null}* animals **(g)** and comparison of axo-dendritic overlap volume between LPLC2 and the GF for these conditions **(h)**; Dots, individual brains. Error bars, means ±95% confidence intervals.

i, Protein interaction map showing binding strength (affinity values are inversely proportional to edge thickness) between DIP-ε and multiple Dprs expressed in LPLC2.

j, expression levels of DIP-ε binding Dprs in LPLC2 at 48h and 96h APF (inferred from scRNA-seq data generated in this study).

k, Correlation between expression levels of dpr13 and other DIP-ε binding dprs in LPLC2 at 48h and 96h APF (inferred from scRNA-seq), along PC1 across the LPLC2 population at 48h and 96h APF.

l-m, Confocal projections of LPLC2 axon terminals and the Giant Fiber dendrites in control animals and animals overexpressing DIP-ε in LPLC2 (**l**) and comparison of axo-dendritic overlap volume between LPLC2 and the GF for these conditions (**m**); dots, individual brains. Error bars, means ±95% confidence intervals.

n, Suggested model: a dorsoventral gradient of dpr13 expression across LPLC2 population determines how many synapses each LPLC2 axon will make with the GF via binding with DIP-ε on the GF dendrites.

Extended Data Figure 6 (related to Fig. 3)

Extended Data Figure 6 (related to Fig. 3). Electrophysiology of the GF
a, GF responses to looming stimuli, in r/v = 10ms (top), r/v = 20ms (middle), r/v = 80ms (bottom). Control (average: black, individual fly: grey) and *DIP-ε^{null}* (average: orange, individual fly: light orange) traces are overlayed. Looming stimulus profile over time is displayed below the GF responses.
b, Quantification of expansion peak amplitudes in (**a**) from individual flies.
c, Same as (**a**) for controls (grey) and animals overexpressing DIP-ε RNAi in the GF (light blue).
d, Quantification of expansion peak amplitudes in (**c**) from individual flies.

Extended Data Figure 7 (related to Fig. 3). Functional relevance of DIP-ε::Dpr13 molecular gradient regulating the LPLC2-GF synaptic gradient.

a, Histograms showing the distribution of takeoff sequence durations at different stimulus elevations (0°, 45°, and 77°) in wild-type (left, grey) and *DIP-ε^null^* flies (right, orange). Short-mode and long-mode takeoffs are distinguished by red dashed lines. N, number of flies that performed a takeoff.

b, Same as (**a**) for controls (empty-Gal4 driving DIP-ε RNAi expression, left, grey) and flies expressing DIP-ε RNAi in the GF (right, blue).

c, Same as (**a**) for controls (empty-Gal4 driving expression of dpr13, left, grey) and flies overexpressing dpr13 in LPLC2 neurons (right, green).

a DV gradient of beat-VI expression in LPLC2

All LPLC2   LPLC2 ∩ beat-VI

48h APF

D ↕ V

NCad   50 µm

96h APF

nc82

b Side-II not required for LPLC2-GF connectivity

c Beat-VI is not required for LPLC2-GF connectivity

axo-dendritic overlap volume, µm³

ns

wild type   side-II^null

n.s.   n.s.

LPLC2>Control (KK)   >beat-VI RNAi (KK)   LPLC2>Control (GD)   >beat-VI RNAi (GD)

d

LPLC2>Control   LPLC2>Beat-VI RNAi

dorsal population

LPLC2   D   L↔   1 2 3 4   P   M   10µm

n=16   n=18

central population

ventral population

e   VT049479-GAL4   OL0048b split-GAL4

>control   >beat-VI RNAi #2   >control RNAi   >beat-VI RNAi (KK)   >Beat-VI RNAi (GD)

Dorsal   1 2 3 4   A ↕ P   10 µm

Central

Ventral   N=9   N=7   N=8   N=6   N=4

f

FISH puncta counts

**   UAS-Control   5xUAS-beat-VI

g   LPLC2>Control   >UAS-beat-VI   >UAS-beat-VI, >UAS-beat-VI RNAi

dorsal   ventral   dorsal   ventral   dorsal

Increased Lop4 density   rescue

Extended Data Figure 8 (related to Fig. 4). Graded expression of Beat-VI across LPLC2 neurons differentially affects the LoP4 dendritic wiring.

a, Dorsally biased expression of beat-VI across LPLC2 population. Red, subsets of LPLC2 neurons expressing beat-VI at 48h APF (top) and 96h APF (bottom). Green, all LPCL2 neurons. D, dorsal; V, ventral.

b-c, Comparison of axo-dendritic overlap volume between LPLC2 and the GF in *side-II^null* animals (**b**), and in animals with LPLC2 expressing beat-VI RNAi (**c**). Dots, individual brains. Error bars, means ±95% confidence intervals.

d, Confocal projections of LPLC2 dendrites (entire LPLC2 population labeled) in the lobula plate in control animals (left) and in animals with beat-VI RNAi expressed in LPLC2 (riht). Leftmost panels, posterior views of the LPLC2 population with dashed rectangles indicating the location of cross-sections for dorsal, central, and ventral subsets of LPLC2. Numbers indicate LoP layers. Yellow arrows indicate LoP4 layer. D, dorsal; M, medial; L, lateral; P, posterior.

e, Same as (**d**) for two different LPLC2 GAL4 driver lines and two different beat-VI RNAi lines.

f, Comparison of beat-VI mRNA levels in LPLC2 (assessed using HCR-FISH) in control animals and animals overexpressing beat-VI in LPLC2. Dots, individual brains. Error bars, means ±95% confidence intervals.

g, Confocal projections of LPLC2 dendrites (entire LPLC2 population labeled, dorsal and ventral cross-sections compared) in the lobula plate in control animals (left), animals overexpressing beat-VI cDNA (middle), and animals overexpressing beat-VI cDNA while also expressing beat-VI RNAi (right). Yellow arrowheads indicate a change in LoP4 dendritic density.

Extended Data Figure 9 (related to Fig. 4)

directional
sensitivity
=
$$\frac{D0° - D180°}{D0° + D180°}$$

Extended Data Figure 9 (related to Fig. 4). Beat-VI::Side-II molecular gradient regulates LPLC2-T4d/T5d synaptic gradient.

a, 3D reconstruction of LPLC2 neurons from a Z-stack taken using the two-photon microscope in a head-fixed fly (lateral view, scale bar: 10 μm). Dashed lines highlight the two Z-planes where LPLC2 dendrites responded to dark looming at positions 16 (red) and 40 (blue) on the LED display (top). White solid lines approximately define the reference neuropils. Schematic of the procedure used to identify the putative neurons in the fly connectome stimulated by two representative grid positions (bottom, see Methods).

b, Polar plots of the average peak responses to moving dark edges in dorsal (16) and ventral (40) regions (color-coded by position) for control and beat-VI RNAi flies. Error band, s.e.m.

c, Bootstrap of the mean of the directional sensitivity ratio in dorsal (16) and ventral (40) regions (color-coded by position) for control and beat-VI RNAi flies (10,000 repetitions).

d, Polar plots of the peak responses to moving dark edges in dorsal (14) and ventral (30) regions in two representative flies (color-coded by position).

e, Polar plots of the average peak responses to moving dark edges in aggregated dorsal (above the equator) and ventral (below the equator) regions for control and beat-VI RNAi flies. Error band, s.e.m.

f, Average calcium transient in response to a dark looming in dorsal (above the equator) and ventral (below the equator) regions for control and beat-VI RNAi flies. Error band, s.e.m.

g, Average calcium transients in response to dark edges moving in 24 (from 0° to 345°) different orientations for control and LPLC2>beat-VI RNAi flies. Dorsal (left) and ventral (right) regions. Error band, s.e.m.