Crystal structure and catalytic mechanism of PL35 family glycosaminoglycan lyases with an ultrabroad substrate spectrum

Lin Wei[1,6], Hai-Yan Cao[2,6], Ruyi Zou[1], Min Du[1], Qingdong Zhang[3], Danrong Lu[3], Xiangyu Xu[1], Yingying Xu[1], Wenshuang Wang[1], Xiu-Lan Chen[4,5], Yu-Zhong Zhang[2,4,5,]*, Fuchuan Li[1,5]*

[1] National Glycoengineering Research Center and Shandong Key Laboratory of Carbohydrate Chemistry and Glycobiology, Shandong University, Qingdao, China.

[2] MOE Key Laboratory of Evolution and Marine Biodiversity, Frontiers Science Center for Deep Ocean Multispheres and Earth System & College of Marine Life Sciences, Ocean University of China, Qingdao, China.

[3] School of Life Science and Technology, Weifang Medical University, Weifang, China.

[4] Marine Biotechnology Research Center, State Key Laboratory of Microbial Technology, Shandong University, Qingdao, China.

[5] Joint Research Center for Marine Microbial Science and Technology, Shandong University and Ocean University of China, Qingdao, China.

[6] These authors contributed equally to this work.

* Corresponding authors. E-mail addresses: zhangyz@sdu.edu.cn (Y. Z.); fuchuanli@sdu.edu.cn (F. L.)

Abstract

Recently, a new class of glycosaminoglycan (GAG) lyases (GAGases) belonging to PL35 family has been discovered with an ultrabroad substrate spectrum that can degrade three types of uronic acid-containing GAGs (hyaluronic acid, chondroitin sulfate and heparan sulfate) or even alginate. In this study, the structures of GAGase II from *Spirosoma fluviale* and GAGase VII from *Bacteroides intestinalis* DSM 17393 were determined at 1.9 and 2.4 Å resolution, respectively, and their catalytic mechanism was investigated by the site-directed mutant of their crucial residues and molecular docking assay. Structural analysis showed that GAGase II and GAGase VII consist of an N-terminal $(\alpha/\alpha)_7$ toroid multidomain and a C-terminal two-layered β-sheet domain with Mn^{2+}. Notably, although GAGases share similar folds and catalytic

mechanisms with some GAG lyases and alginate lyases, they exhibit higher structural homology

with alginate lyases than GAG lyases, which may present a crucial structural evidence for the

speculation that GAG lyases with $(\alpha/\alpha)_n$ toroid and antiparallel β-sheet structures arrived by a

divergent evolution from alginate lyases with the same folds. Overall, this study not only solved

the structure of PL35 GAG lyases for the first time and investigated their catalytic mechanism,

especially the reason why GAGase III can additionally degrade alginate, but also provided a key

clue in the divergent evolution of GAG lyases that originated from alginate lyases.

Introduction

Glycosaminoglycans (GAGs) are a class of linear polyanionic polysaccharides ubiquitously

distributed on the cell surface and in the extracellular matrix (ECM) of animal tissues (***Cohen***

and Merzendorfer,***2019***), and they participate in various physiological and pathological

processes through interacting with a series of chemokines (***Dyer, et al.***,***2015***; ***Irie, et al.***,***2008***),

growth factors (***Sirko, et al.***,***2010***; ***Zhang, et al.***,***2019***) or other ECM components. Except for

keratan sulfate (KS), which does not contain hexuronic acid (HexUA) residues, other GAGs are

composed of repeating disaccharides consisted of HexUA (D-glucuronic acid (GlcUA) or L-

iduronic acid (IdoUA)) and hexosamine (HexNAc) (D-*N*-acetylgalactosamine (GalNAc) or D-*N*-

acetylglucosamine (GlcNAc)). Based on the disaccharide composition and the type of glycosidic

bonds between disaccharides units, HexUA-containing GAGs are classified into three classes:

hyaluronan (HA), chondroitin sulfate (CS)/dermatan sulfate (DS) and heparan sulfate

(HS)/heparin (Hep) (***Cohen and Merzendorfer***,***2019***). HA is the only non-sulfated GAG

composed of repeating -GlcUAβ1-3GlcNAc- disaccharides linked by β1-4 glycosidic bonds. In

contrast, the structures of CS/DS and Hep/HS are highly complex due to sulfation and

epimerization modifications. The backbone of CS is composed of repeating -GlcUAβ1-

3GalNAc- disaccharides linked by β1-4 glycosidic bonds, and is further sulfated at C-2 of

GlcUA and C-4/C-6 of GalNAc by sulfotransferases; meanwhile, the GlcUA residues in CS can

be epimerized into IdoUA residues by glucuronyl C-5 epimerase to form DS (***Kusche-Gullberg***

and Kjellen,***2003***). Similarly, HS/Hep composed of repeating -4GlcUAβ1-4GlcNAcα1- units

can be sulfated at C-2 of GlcUA and N/C-2/C-3/C-6 of GlcNAc/deacetylated GlcN, and GlcUA

residues are often epimerized to IdoUA residues, especially in Hep (***Cohen and***

Merzendorfer,*2019*). Such complex modification through sulfation and epimerization endows
sulfated GAGs with diverse biological functions.

GAG lyases, belonging to the superfamily of polysaccharide lyases (PLs), specifically
catalyze the degradation of HexUA-containing GAGs (*Charnock, et al.*,*2002*; *Garron and*
Cygler,*2010*). These reactions generate a plethora of oligosaccharides that contain an unsaturated
double bond between C-4 and C-5 on uronic acids at their nonreducing ends, which is generated
via a β-elimination mechanism (*Garron and Cygler*,*2010*). As a complimentary mechanistic
strategy to glycoside hydrolases (GHs), GAG lyases degrade HexUA-containing GAGs without
the participation of a water molecule and involved in the polysaccharide metabolism of
microorganisms (*Garron and Cygler*,*2010*; *Ndeh, et al.*,*2020*). For the past many years, an
increasing number of gene sequences containing PL catalytic modules and some ancillary
modules have been annotated as PLs (*Lombard, et al.*,*2010*). Based on their sequence
homologies, 43 families (and an unclassified PL0 family) have been hierarchically classified in
the CAZy database (http://www.cazy.org/), and at least one member of each family has been
subjected to activity identification and biochemical property analysis (*Drula, et al.*,*2022*). The
identified GAG lyases are widely distributed in the PL6, PL8, PL12, PL13, PL15, PL16, PL21,
PL23, PL29, PL30, PL33, PL35 and PL37 families (*Drula, et al.*,*2022*). Based on substrate
specificity, GAG lyases are classified as HA-specific lyases, which can degrade HA only, CS/DS
lyases (chondroitinase, CSases), which degrade CS/DS as well as HA in most cases, and Hep/HS
lyases (heparinases, Hepases), which specifically cleave Hep/HS. All identified GAG lyases
have substrate specificity to strictly distinguish one or two kinds of GAGs based on saccharide
composition and glycosidic bonds between each HexUA-HexNAc disaccharide units, while
other uronic acid-containing polysaccharides are usually unable to serve as substrates for these
enzymes.

With the evolution of host polysaccharides (*Csoka and Stern*,*2013*; *Popper, et al.*,*2011*),
the bacterial enzymes that degrade these polysaccharides have also evolved to varying degrees in
response to complex environments. As a kind of unbranched anionic extracellular polysaccharide
produced by lower organisms algae and bacteria (such as brown alga (*Popper, et al.*,*2011*) and
bacteria belonging to *Pseudomonas* (*Evans and Linker*,*1973*) and *Azotobacter* (*Clementi*,*1997*),
alginate containing no hexosamine and sulfation possesses a simpler structure than GAGs in
animals and should emerge earlier than GAGs during the course of evolution. Like GAGs,

alginate is a linear polyanionic polysaccharide that contains HexUA residues; however, alginate
possesses a completely different chemical structure, which is composed of D-mannuronate (M)
and its C5 epimer L-guluronate (G), and M and G residues often alternate randomly to form
heteropolyuronic blocks, including polyM composed of M residues linked by β1-4 bonds, polyG
composed of G residues linked by α1-4 bonds, and polyMG/GM composed of alternating M and
G linked by β/α1-4 bonds (*Pawar and Edgar,2012*). Likewise, a series of alginate lyases
essential for the metabolism of alginate were identified from microorganisms and algae as well
as lower marine animals, which belong to the PL5, PL6, PL7, PL8, PL14, PL15, PL17, PL18,
PL31, PL32, PL34, PL36, PL38, PL39 and PL41 families (*Drula, et al.,2022*). Based on their
specificity, these lyases are also divided into the following categories: M block-specific lyases
(*Itoh, et al.,2019*; *Zhu, et al.,2015*), G block-specific lyases (*Matsubara, et al.,1998*; *Yamasaki,*
et al.,2005) and MG-specific alginate lyases (*Jagtap, et al.,2014*; *Yamasaki, et al.,2004*). From
an evolutionary perspective in previous reports, GAG lyases are thought to have originated from
alginate lyases by a divergent evolution through adaptation to the evolution of substrate
polysaccharides, which is mainly attributed to some homology in the sequences and folds of
these two types of enzymes, such as the $(\alpha/\alpha)_n$ toroid and antiparallel β-sheet folds of GAG
lyases in PL8 (hyaluronate lyase and chondroitin lyase), PL12 (heparinase III), PL21 (heparinase
II) and PL23 (chondroitin lyase) families and alginate lyases in PL15, PL17 and PL39 families,
the β-jelly roll fold of heparinase I in PL13 family and alginate lyases in PL7 and PL18 families,
and the β-helix fold of chondroitinase B and alginate lyase in PL6 family (*Garron and*
Cygler,2014). Considering that the alginate with simpler structure from brown alga or bacteria
might appear earlier, it is possible that the ancestral enzymes originally used alginate as a
substrate. However, enzymes with transitional characteristics in activity and structure have not
been discovered to support this speculation.
Recently, a new class of GAG lyases belonging to PL35 family was discovered with an
ultrabroad substrate spectrum (*Wei, et al.,2024*). Unlike the identified GAG lyases (HA-specific
lyases, CSases and Hepases) with quite strict substrate specificities, these novel GAG lyases can
degrade three types of uronic acid-containing GAGs (HA, CS and HS), and thus were named
GAGases. More interestingly, one of the eight identified GAGases (GAGase III) can even
degrade alginate. Analysis of substrate structure preference represented by GAGase I showed
that GAGases selectively act on GAG domains composed of non/6-*O*-/*N*-sulfated hexosamines

and D-glucuronic acid, among which GAGase III can also selectively degrade polyM blocks composed of D-mannuronic acids but not polyG blocks composed of L-guluronic acids in alginate. Furthermore, the primary structure alignment and phylogenetic analysis showed that GAGases have considerable degree of homology with not only GAG lyases from PL15, PL21 and PL33 families but also alginate lyases from PL15, PL17 and PL34 families (***Wei, et al.,2024***), indicating that GAGases may exhibit some transitional features from alginate lyases to GAG lyases in structure.

In this study, crystal structures of two GAGases (GAGase II and GAGase VII) were determined for the first time, and their catalytic mechanism was further elucidated through the site-directed mutagenesis of their crucial site residues and molecular docking. Structural alignment showed that GAGases are more structural similarity to some alginate lyases rather than GAG lyases, suggesting that PL35 family GAGases might originate from alginate lyases possessing an N-terminal $(\alpha/\alpha)_n$ toroid domain and a C-terminal antiparallel β-sheet domain. Moreover, one of the reasons why GAGase III can additionally degrade M-block alginate are also resolved by structural modeling, structural alignment and site-directed mutagenesis. This study not only helps to resolve the catalytic mechanism of the PL35 family lyases in particular GAGases, but also provides a potential structural evidence for the divergent evolution of GAG lyases.

Results

Overall crystal structures of GAGase II and GAGase VII

To explore the substrate recognition and catalytic mechanism of the GAGases, the eight enzymes with different sequence identity (*Supplemental Table S1*) were individually used to prepare their crystals for structural analysis. Finally, the structures of GAGase II from *Spirosoma fluviale* (GenBank accession number: SOD82962.1, 81.2% sequence identity with GAGase I), GAGase VII from *Bacteroides intestinalis* (GenBank accession number: EDV05210.1, 44.8% sequence identity with GAGase I) and selenomethionine (SeMet)-labelled GAGase II (Se-GAGase II) in their ligand-free states were solved at 1.9 Å, 2.4 Å and 2.0 Å resolution, respectively (*Table 1*). Both GAGase II and GAGase VII are monomers consisting of two domains: a N-terminal $(\alpha/\alpha)_7$ toroid domain and a C-terminal two-layered β-sheet domain (*Figure 1A*).

The N-terminal domain (Met^{34}-Ala^{365}) of GAGase II is composed of 16 large or small α-
helices (α1-α16) to form an $(α/α)_7$ toroid domain. Specifically, the N-terminal 16 α-helices
constitute seven complete α-helix pairs (α1-2, α3-4, α5-7, α8-9, α10-12, α13-14 and α15-16),
forming seven hairpin structures, which eventually embrace into the $(α/α)_7$ toroid structure. The
transition between α6 to α8 and between α10 to 12 is discontinuous, and both transitions possess
a small α-helix α7 and α11, respectively. In each hairpin structure, two antiparallel α-helices are
connected by loops composed of 9-12 residues, and each hairpin contains a turn structure
composed of 1-4 residues. The N-terminal of this toroid domain is connected to a signal peptide
that has been removed during cloning expression, and its C-terminal is connected to the two-
layered β-sheet domain through a 17-residue linkage (Trp^{350}-Met^{365}). Its inner layer is formed by
helices α1, α3, α5, α8, α10, α13 and α16, while its outer layer is formed by α2, α4, α6+α7, α9,
α11+α12, α14 and α15. Moreover, the α-helix pairs α13-14 and α15-16 are significantly shorter
than the first five sets, and α15 and α16 are connected by a short β-turn consisting of only 3
residues, which makes the toroid structure closed (*Figure 1A*).

The C-terminal domain (Met^{366}-Ser^{612}) of GAGase II is composed of 17 β-strands (β1-β17)
to form a two-layered β-sheet domain with 4 small α-helices located between β5-β6 and β8-β9.
Both layers of this C-terminal β-sheet domain are composed of eight groups of β-strands, namely
β1-β5, β11 and β15-β16 and β7-β10, β12-β14 and β17. The two layers are roughly parallel to
each other and twisted to form an angle of approximately 60°. The first five β-strands (β1-β5) of
the C-terminal domain are arranged in anti-parallel configuration. Subsequently, there is an Ω-
loop structure wrapping around a metal ions (Mn^{2+}), which consists of three small α-helices and
a small β-strands, with many positively charged amino acid residues (such as His^{401}, His^{403},
Tyr^{427}, Glu^{431}, Trp^{438}, Arg^{446} and so on) on top, which might serve as potential substrate-binding
sites. Finally, the C-terminal domain terminates at β17 in the second layer of the bilayer structure.
Taken together, these structural elements constitute the C-terminal anti-parallel bilayer β-sheet
domain of GAGase II (*Figure. 1A*).

Similar to GAGase II, GAGase VII also has an N-terminal $(α/α)_7$ toroid domain (Leu^{35}-
Arg^{360}) composed of 16 α-helices, but its α1 helix consists of only three residues (Glu^{42}-Val^{44})
and a long loop chain (*Figure. 1B*). Moreover, the C-terminal two-layered β-sheet domain
(Met^{361}-Met^{616}) of GAGase VII is also composed of 17 β-strands (β1-β15, β16a, β16b and β17).
However, unlike GAGase II, its β16 strands is interrupted into two small β-strands (β16a, β16b)

by a randomly coiled loop chain (*Figure. 1B*). Compared with GAGase II, these additional loop

regions may make GAGase VII less structurally stable than GAGase II, and may be an important

factor in why the activity of GAGase VII is far lower than GAGase II. Nevertheless, structural

alignment further shows that GAGase II has good superposition with GAGase VII and other

GAGases, indicating that GAGases are highly conserved in structures (*Supplemental Figure S1*).

Mn binding site of GAGase II and GAGase VII

Unlike the GAG lyases identified from PL8, PL15 and PL21 families, which contain Ca^{2+}

or Zn^{2+} in their overall crystal structures (***Shaya, et al.,2008***; ***Shaya, et al.,2006***; ***Zhang, et***

al.,2021), a Mn^{2+} was detected in C-terminal domains of both GAGase II and VII by inductively

coupled plasma-mass spectrometry (ICP-MS) analysis (*Figure 1B*). Structurally, the Mn^{2+} is

located inside the long Ω loop after the β1-β5 antiparallel β-sheet, and coordinates with the

nearby nitrogen and oxygen atoms of several residues such as Asp and His in both GAGase VII

and GAGase II (GAGase II also has coordination with three surrounding water molecules), with

distances of 2.0-2.5 Å. Such interaction should stabilize the nearby loop structures and may form

a substrate-binding site relevant for substrate selectivity. To investigate the roles of these

adjacent residues interacting with the Mn^{2+}, His^{403}, Asp^{405}, Asp^{421} and His^{457} in GAGase II were

individually mutated to alanine (Ala), and the results showed that these mutations resulted in

complete or substantial inactivation of GAGase II (*Figure 2A*). In addition, the inhibitory effects

of the chelator (EDTA) and the stimulation of Mn^{2+} on the enzymatic activities also support the

critical role of Mn^{2+} on GAGases (*Figure 2B*).

Figure 1. Structural description of GAGase II and GAGase VII. (A), Overall structures of GAGase II
(*left*) and GAGase VII (*right*) are shown in carton. The α-helix, β-strand and random coil are colored with
yellow, green and gray, respectively. The N-terminal $(\alpha/\alpha)_7$ toroid domain and C-terminal two-layered β-
sheet domain were circled and the secondary structure elements are marked nearby. **(B)**, Mn binding site
of GAGase II and GAGase VII. The *purple* sphere presents Mn^{2+}. Mn^{2+} binding site of GAGase II (*left*)
and GAGase VII (*right*) is shown in stick. A cut off distance of 3.0 Å was carried out to choose
neighboring residues.

Table 1. Data collection and refinement statistics.

	GAGase II	GAGase VII	SeMet-GAGase II
PDB entry code	8KHV	8KHW	-
Data collection			
Wavelength (Å)	0.9792	0.9792	0.9792
Resolution range	44.01-1.90 (1.94-1.90)	19.87-2.40 (2.49-2.40)	40.91-2.00 (2.071-2.00)
Space group	$P2_1$	$P4_32_12$	$P2_1$
Unit cell	a=45.917 Å	a=98.4075 Å	a=46.39 Å

	b=82.096 Å	b=98.4075 Å	b=81.819 Å
	c=79.8854 Å	c=134.75 Å	c=79.6576 Å
	$\alpha=90°$	$\alpha=90°$	$\alpha=90°$
	$\beta=106.59°$	$\beta=90°$	$\beta=106.879°$
	$\gamma=90°$	$\gamma=90°$	$\gamma=90°$
Unique reflections	86421 (4092)	49305 (2737)	38212 (3738)
Completeness (%)	98.55 (92.04)	99.68 (99.85)	99.15 (98.47)
R_{meas}	0.100 (0.230)	0.100 (0.737)	0.084 (0.238)
$R_{p.i.m}$	0.038 (0.100)	0.020 (0.158)	0.032 (0.093)
Mean $I/\sigma(I)$	12.9 (6.1)	20.6 (4.3)	17.9 (8.2)
Refinement statistics			
R_{work}	0.1584 (0.1712)	0.2079 (0.2639)	-
R_{free}	0.1919 (0.2267)	0.2722 (0.3622)	-
RMSD bond length (Å)	0.006	0.008	-
RMSD bond angle (°)	0.79	0.99	-
Ramachandran plot (%)			
Favored	97.75	94.31	-
Allowed	2.25	5.00	-
Outliers	0.00	0.69	-
Rotamer outliers (%)	0.00	0.40	-
Clashscore	2.78	6.89	-
Average B-factor ($Å^2$)	13.51	54.14	-
Macromolecules	11.66	54.20	-
Ligands	21.71	45.54	-
Solvent	25.75	47.24	-

Statistics for the highest-resolution shell are shown in parentheses.

Figure 2. Activity of GAGase II and its mutants. (A), Activity of site-directed mutants of His and Asp

nearby the Mn^{2+}. His and Asp residues nearby the Mn^{2+} were individually mutated to Ala and the relative

activity of each mutant was measured as described in "*Materials and methods*". All of the residual

activities were evaluated and shown as the relative intensity compared with that of wild type GAGase II.

***: $p<0.001$. **(B)**, The effects of Mn^{2+} and chelating reagent of GAGase II were determined using HA (1

mg/ml) as substrate. Error bars represent averages of triplicates ± S.D.

Multiple structural alignments of GAGase II and its structurally similar GAG lyases and alginate lyases

The ultrabroad substrate degradation spectra of GAGases indicate that they should be

structurally homologous with some other PLs with different substrate specificities. By using

GAGase II as a representative, a structural alignment assay showed that the PL35 GAGases are

structurally similar to various N-terminal $(\alpha/\alpha)_n$ toroid domain and C-terminal antiparallel β-

sheet domain-containing GAG/alginate lyases from PL8 (***Fethiere, et al.,1999***; ***Huang, et***

al.,2003; ***Li and Jedrzejas,2001***; ***Lunin, et al.,2004***; ***Shaya, et al.,2008***), PL12 (***Hashimoto, et***

al.,2014; ***Ulaganathan, et al.,2017***), PL15 (***Ochiai, et al.,2010***; ***Zhang, et al.,2021***), PL17 (***Park,***

et al.,2014), PL21 (***Shaya, et al.,2006***), PL23 (***Sugiura, et al.,2011***) and PL39 (***Ji, et al.,2019***).

Notably, GAGase II shares very low sequence identity with PL21 heparinase II from *Pedobacter*

heparinus (***Shaya, et al.,2006***) (23.1% identity with 29% query coverage), PL12 heparinase III

from *Bacteroides thetaiotaomicronin* (***Ulaganathan, et al.,2017***) (23.6% identity with 32%

query cover) and PL39 alginate lyase from *Defluviitalea phaphyphila* (***Ji, et al.,2019***) (26.5%

identity with 31% coverage), and no significant sequence identity with PL17 alginate lyases

(***Park, et al.,2014***), PL15 exoHep (***Zhang, et al.,2021***), PL8 chondroitin ABC lyase (***Shaya, et***

al.,2008) and chondroitin AC lyase (***Lunin, et al.,2004***); however, all are relatively well

superimposable in their overall structures and highly conserved in their triplet active site residues

(*Figure 3*). Among them, GAGases show more structural similarity to alginate lyases, such as

those from the PL15 (PDB code: 3a0o) (***Ochiai, et al.,2010***), PL17 (PDB code: 4NEI) (***Park, et***

al.,2014) and PL39 (PDB code: 6JP4) (***Ji, et al.,2019***) families, than various GAG lyases (*Table*

2), indicating that GAGases might originate from alginate lyase rather than GAG lyase.

Reasonable support for this speculation could be the significant alginate-degrading activity of

GAGase III. Therefore, the broad substrate spectrum of GAGases may be because they are

evolutionarily transitional types with the enzymatic features of alginate lyases and GAG lyases.

Figure 3. Multiple structural alignments of GAGase II and its structurally similar proteins. GAGase II (8KHV, *blue*) was aligned with structurally identified GAGs and alginate lyases, including PL17 family alginate lyase (4NEI, *purple*), PL39 family alginate lyase (6JP4, *olive*), PL15 family alginate lyase (3A0O, *red*), PL12 family heparinase III (4MMH, *salmon*), PL15 family exoHep (6LJA, *gray*), PL21 family heparinase II (2FUQ, *magenta*), PL8 family chondroitin sulfate AC lyase II (1RWA, *green*), PL23 family chondroitinase (3VSM, *yellow*) and PL8 family chondroitin sulfate ABC lyase II (2Q1F, *pink*). The detailed views of the crucial active site residues are shown in stick mode. The root-mean-square deviation (RMSD) between these structures and GAGase II were calculated as 1.342, 1.344, 1.299, 1.242, 1.145, 1.352, 1.235, 1.138 and 1.405 Å based on 171, 148, 141, 111, 99, 87, 60, 22 and 21 pruned atoms, respectively.

Table 2. Structural homology of GAGase II/GAGase VII with GAG/alginate lyases analysed using DALI.

	Chain	Z-score	RMSD (Å)	lali	nres	Description	PL family
GAGase	3a0o-A	29.4	3.2	539	764	Alginate lyase	15
II	6jp4-A	28.9	3.0	523	770	Alginate lyase	39

	4nei-A	27.3	3.1	523	705	Alginate lyase	17
	4mmh-A	28.6	4.4	481	637	Heparinase III	12
	6lja-A	25.7	3.9	540	841	Exo-type heparinase	15
	2fuq-A	25.1	4.9	528	743	Heparinase II	21
	1rwa-A	15.2	5.4	470	754	Chondroitin AC lyase	8
	3vsm-a	14.3	5.3	369	633	Chondroitinase	23
	2q1f-a	11.1	5.3	434	991	Chondroitin ABC lyase	8
GAGase VII	3a0o-A	29.2	3.6	545	764	Alginate lyase	15
	6jp4-A	28.0	3.3	530	770	Alginate lyase	39
	4nei-A	27.2	3.2	529	705	Alginate lyase	17
	4mmh-A	26.4	4.4	492	637	Heparinase III	12
	2fuq-A	23.1	5.4	531	747	Heparinase II	21
	1rwa-A	14.3	4.5	473	754	Chondroitin AC lyase	8
	3vsm	12.9	4.3	365	633	Chondroitinase	23
	2q1f-A	10.5	4.6	407	991	Chondroitin AC lyase	8

Catalytic center and substrate binding sites of GAGases

Based on the sequence and structural alignment with homogenous alginate/GAG lyases, the conserved residues Tyr^{246}, His^{401}, Asn^{192} of GAGase II, Tyr^{243}, His^{398}, Asn^{189} of GAGase III, and Tyr^{241}, His^{397}, Asn^{187} of GAGase VII may be the key triplet residues involved in their β-elimination catalysis active sites (*Figure 3*). To confirm this hypothesis, a site-directed mutagenesis assay was performed in which these residues were individually replaced by Ala; as a result, the activity of all of mutants toward CS-C was completely lost (*Figure 4A*). These results indicate that GAGases act through a general acid–base catalytic mechanism by using His/Tyr as a Brønsted acid/base, similar to that observed for various other identified GAG/alginate lyases (*Table* 3). Compared with their homologous GAG/alginate lyases, GAGases contain a shorter catalytic cavity, which may explain why they can accommodate and degrade a variety of substrates with very different structures. However, we were unsuccessful in preparing the cocrystals of GAGases/their inactive mutants with substrates to reveal their substrate binding and degradation mechanisms. Alternatively, molecular docking experiments of GAGase II with several structurally defined hexasaccharide (Hexa) substrates were performed as shown in *figure 4B*. Based on the docking results obtained for GAGase II with an HA Hexa (GlcUA1–3GlcNAc)$_3$ (PDB code: 1HYA) (**Winter, et al.,1975**) (*Figure 4B*), the nonreducing end of the substrate at the "-1" and "-2" subsites is near some positively charged amino acids, including Arg^{87}, Arg^{88}, Arg^{94}, His^{136}, Asn^{191} and Asn^{192}, in the vicinity of the catalytic cavity of

GAGase II, which could facilitate the binding of negatively charged substrates. Besides, several acidic amino acids (Glu^{242}, Asp^{299} and Glu^{431}) are very close to the "+1" and "+2" subsites, which could promote the release of the reducing end product via charge repulsion. There are some other residues surrounding around the catalytic cavity, such as Tyr^{240}, Gln^{307}, Arg^{342}, Tyr^{427}, Glu^{428}, Glu^{431}, Trp^{438}, Met^{440} and Arg^{446}, which may play a key role in stabilization of the catalytic cavity and overall structure. Site-directed mutagenesis of these residues individually replaced with Ala showed that all resulting mutants were partially or completely inactivated (*Figure 4C*), confirming that these residues play important roles in the binding and release of substrate and structural stabilization.

Figure 4. Catalytic center and substrate binding sites of GAGases. (A), The crucial catalytic site-directed mutagenesis of GAGase II, GAGase III and GAGase VII. The conserved crucial residues of GAGase II, GAGase III and GAGase VII were individually mutated to Ala. CS-C and alginate were used as substrates for the activity evaluation of GAGase II, GAGase III, GAGase VII and its mutants. The activity of each mutant was detected using gel filtration HPLC on a Superdex Peptide column as described in "*Materials and methods*"; **(B)**, Molecular docking of GAGase II with a HA hexasaccharide. The molecule docking was carried out with GAGase II and a HA hexasaccharide (PDB code: 1HYA) to predict the substrate binding sites. The binding site residues (*green*) and hexasaccharide ligand (*gray*) are showed as sticks. **(C)**, The putative substrate-binding site residues surrounding the docking substrate were individually mutated to Ala and the relative activity of each mutant was shown as the relative intensity compared with that of wild type GAGase II. *: $p<0.01$, **: $p<0.001$, ***: $p<0.0001$, #: the activities of the mutants were too low to be accurately detected and almost completely inactivated.

Table 3. Catalytic residues comparison of identified GAGs and alginate lyases shared similar structure with GAGase II and GAGase VII

	PDB code	PL family	Substrates	Sequence identity (Query cover/Per. Ident)*	Catalytic residues	
					General base/acid	Neutralizer
GAGase II	8KHV	PL35	HA, CS and HS	100%/100%	Tyr246 His401	Asn192
GAGase VII	8KHW	PL35	HA, CS and HS	93%/44.67%	Tyr241, His397	Asn187
GAGase III	-	PL35	HA, CS, HS and alginate (M-specific)	98%/64.08%	Tyr243,His398,	Asn189 His188
Chondroitin sulfate AC lyase II	1RWA	PL8	HA and CS	N.D.**	His233 Tyr242,	Asn183
Chondroitin sulfate ABC lyase II	2Q1F	PL8	HA, CS and DS	N.D.	His345,Tyr461,His454	Asp398
Heparinase III	4MMH	PL12	HS	8%/30.91%	Tyr294, His424	Asn240
Alginate lyase	3A0O	PL15	Alginate (M/G specific)	43%/23.93%	His311,Tyr365,His531	Arg314
ExoHepase	6LJA	PL15	Hep and HS	N.D.	His337,Tyr390,His555	Asn336
Alginate lyase	4NEI	PL17	Alginate (M-specific)	15%/25.51%	Tyr258, His413	Asn201,His200
Heparinase II	2FUQ	PL21	Hep and HS	29%/23.13%	His202, Tyr257, His406	Glu205
Chondroitin lyase	3VSM	PL23	CS	N.D.	Tyr299,His291	Asn236
Alginate lyase	6JP4	PL39	Alginate (M/G specific)	31%/26.53%	Tyr239, His405	Asn186,His187

* Sequence identity means the sequence similarity compared with GAGase II.

**N.D. means no significant identity is detected.

Substrate selectivity of GAGases

As mentioned above, GAGase III can degrade not only HA, CS and HS but also alginate, even though the activity on alginate is relatively low (about 1/50 of the activity on HA). To investigate the reason why this enzyme is capable to act on alginate, its structure was predicted using Robetta, and aligned with GAGase II and GAGase VII. Results showed that GAGase III has a unique His[188] residue in the catalytic cavity of GAGase III, which is conserved in alginate lyases from PL17 (*Park, et al.,2014*), PL39 (*Ji, et al.,2019*), PL38 (*Ronne, et al.,2023*), PL5 (*Pandey, et al.,2021*) and PL15 (*Ochiai, et al.,2010*) families, while other GAGases have a conserved asparagine residue such as the Asn[191] of GAGase II and the Asn[186] of GAGase VII in

the corresponding position (*Figure 5A-B*). To verify the function of this histidine, His[188] was

replaced with alanine and asparagine, respectively, and the results of substrate-specificity

analysis showed that the alginate-degrading activity of GAGase III-H188A and H188N was

completely abolished but only partially affect its GAG-degrading activity, indicating that the

His[188] residue is essential for the alginate-degrading activity of GAGase III (*Figure 5C-D*).

However, when the asparagine residue at the corresponding position of other GAGases was

muted to histidine residue the obtained mutants did not exhibit any alginate-degrading activity

(*Supplemental Figure S2*), suggesting that other structural factors that are vital to the alginate-

degrading activity of GAGase II remains to determine excepting for the His[188] residue.

Figure 5. Analysis of a key residue for the alginate-degrading activity of GAGase III

(A), Multiple structural alignment of GAGase II, III and VII. GAGase II (8KHV, *green*) and GAGase VII

(8KHW, *skyblue*) were aligned with GAGase III (model, *gray*). The detailed views of crucial catalytic

residues are shown in stick mode; (B), Conversed catalytic residues of alginate lyases from PL17 (4NEI,

purple), PL39 (6JP4, *yellow*), PL38 (8BDQ, *slate*), PL5 (7FHU, *cyan*) and PL15 (3A0O, *pink*) family.

Residues are shown in stick; (C), Activity assay of GAGase III-H188N and GAGase III-H188A toward

CS-C and alginate. The crucial site His[188] was mutated to alanine and asparagine, respectively. The

activity of each mutant against CS-C and alginate was assessed using gel filtration HPLC on a Superdex

Peptide column, as described under "*Materials and methods*"; (D), Relative activity of GAGase III and its

mutants . Three types of GAGs (HA, CS-C and HS) were individually treated with GAGase III and its mutants (GAGase III-H188N and GAGase III-H188A) at 40 °C for 1 h; relative activities of enzymes were determined by detecting the absorbance at 232 nm. Data are shown as the percentage of the activity relative to the wild-type GAGase III. *, p<0.5; **, p<0.01; ***, p<0.001.

GAGases show preference for some specific sulfation patterns (*Wei, et al.,2024*). For example, GAGases prefer to degrade CS domains composed of non-/6-*O*-sulfated but not 4-*O*-sulfated GalNAc residues and D-GlcA residues. To further investigate the reason why these enzymes have such structural preference for substrates, we tried to prepare the co-crystal of GAGase II or VII with various structure-defined oligosaccharide substrates, but ultimately failed. Therefore, we used molecular docking analysis to preliminarily explore the possible reasons why GAGases have substrate structural preference. To this end, docking assays of GAGase II with a nonsulfated CS Hexa $(GlcUA\beta1-3GalNAc)_3$ (PDB code: 2KQO) and a tri-4-*O*-sulfated CS Hexa $(GlcUA\beta1-3GalNAc(4S))_3$ (PDB code: 1C4S) (*Sattelle, et al.,2010*) were carried out, respectively. In the docked model, the monosaccharide residues of the substrate were named "+1" and "+2" toward reducing end, and "-1" and "-2" toward non-reducing end from the β1-4 cleavage site following the nomenclature (*Davies, et al.,1997*). The results show that for the nonsulfated CS Hexa, the GlcUA residue at the "+1" subsite is closely approached by the Tyr[246] residue as a Brønsted base (*Figure 6 left*), which facilitates the uptake of the C5 proton of GlcUA by Tyr[246], and in the case of tri-4-*O*-sulfated CS Hexa, the 4-*O*-sulfate group in GalNAc residue at the "+2" subsite interacts with the neutralizer Asn[192] and the catalytic residue Tyr[246] via hydrogen bonds etc., the C5 proton of GlcUA at the "+1" subsite is away from the key catalytic residue Tyr[246] (*Figure 6 right*, *Supplemental Figure S3*), which hinders the degradation of CS with such sulfation pattern, as we detected in the biochemical analysis above. Of course, the results from docking assay need to be further confirmed by more structural and biochemical evidences.

Figure 6. Molecular docking of GAGase II with hexasaccharide ligands. Molecular docking of GAGase II with CS ligands. The molecule docking was carried out with GAGase II and CS ligands, including nonsulfated (PDB code: 2KQO) (*left*) and 4-*O*-trisulfated (PDB code: 1C4S) (*right*) CS hexasaccharide, to investigate the substrate selectivity. The catalytic triplet residues (*green*) and CS ligand (*yellow*) are showed as sticks.

Catalytic mechanism of GAGases

In summary, taking GAGase II as representative, the catalytic process of GAGases is proposed as follows: the negatively charged GAG substrate binds to several positively charged residues, such as Arg[87], Arg[88], Arg[94], His[136], Asn[191] and Asn[192], near the catalytic triad residues in the catalytic cavity; the Asn[191] and Asn[192] residues interact with the carboxyl group of GlcUA at the +1 subsite to reduce the pKa and promote the protonation of the C5 proton; His[401] works as a proton receptor (Brønsted base) to abstract protons from the C5 position of GlcUA and create an enolate tautomeric; the protonated Tyr[246] (Brønsted acid) provides a proton to the C-4–O-1 glycosidic bond between the "-1" and "+1" subsite; the glycosidic bond is cleaved as an electron transfer occurs from the carboxyl group to form a double bond between C4 and C5 of GlcUA; and a new reducing end and a new unsaturated nonreducing end are recreated; moreover, the degradation of polyM by GAGase III requires the involvement of His[188] residue, which may involve in the neutralization of carboxyl groups in M-blocks in alginate (*Figure 7*).

Figure 7. Proposed catalytic mechanism of GAGase II and GAGase III. Take the HA degradation by GAGase II and the polyM degradation by GAGase III for example. Briefly, the substrate is firstly binding to the negatively charged residues near the catalytic sites, the carboxylate group is neutralized by Asn[192], Asn[191] in GAGase II and His[188], Asn[189] in GAGase III; His[398] in GAGase II and His[401] in GAGase III are proposed to work as general base to abstract a proton from the C5 position of GlcUA at +1 position, and Tyr[246] in GAGase II and Tyr[243] in GAGase III work as general acid to donate the leaving group at -1 position a proton. The arrows indicate the direction of electron transfer, and the C5-C6 double bond on the middle panel indicate the enolate anion intermediate created by proton abstraction at C-5 position.

Discussion

GAG lyases derived from microorganisms, as GAG-specific degrading enzymes, are classified into 13 PL families in the CAZy database (***Drula, et al.,2022***). They are not only critical for microorganisms to degrade and utilize GAGs as carbon sources, but also become useful tools for structural and functional studies of GAGs. In contrast, alginate lyases, as alginate-specific degrading enzymes, are classified into 15 PL families in the CAZy database (***Drula, et al.,2022***). They are essential for the metabolism of alginate in microorganisms and algae as well as lower marine animals, and have completely different substrate specificities than GAG lyases. Due to the similarity between GAG lyases and alginate lyases in folds, GAG lyases are thought to have originated from the divergent evolution of alginate lyases (***Garron and***

Cygler,*2014*), but enzymes that exhibit transitional features in function and structure, which
would provide direct evidence to support this view, are lacking. On the basis of previous
functional studies (***Wei, et al.***,***2024***), structural studies of GAGases with HA, CS, HS and even
alginate-degrading activities should further provide substantial evidence for the evolution of
GAG lyases.

Prior to this study, no member of the PL35 family had been structurally characterized. Here,
the structures of GAGase II and VII were determined at resolutions of 1.9 Å and 2.4 Å,
respectively, and both exhibit a highly homologous structure of "N-terminal $(\alpha/\alpha)_7$ toroid and C-
terminal two-layer antiparallel β-sheet", which is also the fold adopted by most structurally
known GAG lyases except CSase B in PL6 (***Huang, et al.***,***1999***), Hepase I in PL13 (***Han, et***
al.,***2009***) and hyaluronan lyase in PL16 (***Martinez-Fleites, et al.***,***2009***). Structural alignment has
shown that GAGases share considerable degrees of homology with all GAG/alginate lyases with
the $(\alpha/\alpha)_n$ toroid and antiparallel β-sheet fold in PL8 (***Fethiere, et al.***,***1999***), PL12 (***Hashimoto,***
et al.,***2014***), PL15 (***Ochiai, et al.***,***2010***; ***Zhang, et al.***,***2021***), PL17 (***Park, et al.***,***2014***), PL21
(***Shaya, et al.***,***2006***), PL23 (***Sugiura, et al.***,***2011***) and PL39 (***Ji, et al.***,***2019***) families, especially
the key catalytic triplet residues (Tyr, His and Asn) in their active centers, which are highly
conserved; thus, they may share a similar Brønsted base/acid catalytic mechanism (***Garron and***
Cygler,***2010***; ***Garron and Cygler***,***2014***). Notably, GAGases are more structurally homologous to
alginate lyases from PL15 (***Ochiai, et al.***,***2010***), PL17 (***Park, et al.***,***2014***) and PL39 (***Ji, et***
al.,***2019***) family than to various GAG lyases, which further structurally supports the speculation
that GAGases originated from alginate lyases.

As mentioned previously, many different GAG/alginate lyases share similar catalytic center
sites and the same catalytic machinery (***Garron and Cygler***,***2010***; ***Garron and Cygler***,***2014***), but
their differences in substrate selectivity and endo-/exolytic manner are mainly attributed to subtle
differences in the substrate-binding sites. The recognition of appropriate substrates by PLs
mainly depends on the interaction between the positively charged side chains of basic amino acid
residues in their substrate-binding sites and the negatively charged carboxyl groups of HexUA
residues in substrates. Moreover, compared to the structurally and biochemically characterized
members of the PL8, PL12, PL15, PL21 and PL39 families, PL35 family proteins possess a more
open and shorter catalytic cavity; thus, PL35 family proteins can more easily accommodate
various substrates with different structures but may exhibit a weaker ability to bind and degrade

specific substrates. The activities of GAGases towards various GAGs and alginate are much
lower than those of GAG or alginate-specific lyases.

In addition, GAGases tend to act on GlcUA-containing HA, CS and HS but not IdoUA-
containing DS and Hep, which should result from the absence of a His residue functioning as an
anti-base in the catalytic cavity (***Garron and Cygler*,*2010***; ***Shaya, et al.*,*2008***; ***Shaya, et***
al.*,*2006). The His residue as an *anti*-base can substitute Tyr to absorb the C5 proton in the
IdoUA and ensure that β-elimination is carried out successfully, which is detected in chondroitin
sulfate ABC lyase (2Q1F) from PL8 family (His[454]) (***Shaya, et al.*,*2008***) and heparinase II
(2FUQ) from PL21 family (His[202]) (***Shaya, et al.*,*2006***), and endows these enzymes with a
bifunctional activity against GlcUA- and IdoUA-containing GAGs. Notably, GAGase III, the
only identified GAGase with alginate-degrading activity, has a unique His[188] residue also
conversed in various alginate lyases from PL5, PL15, PL17, PL38 and PL39 families. Mutation
assay showed that the His[188] is a key residue for the alginate-degrading activity of GAGase III.
However, when a conversed asparagine residue at the corresponding position of other GAGases
was replaced by histidine, the muted GAGases still could not act on alginate, indicating that
some other unidentified structural factors are also involved in the alginate degradation of
GAGase III. Overall, while the substrate-degrading features of GAGases can be preliminarily
elucidated structurally, more direct evidence remains to be obtained by detailed structural and
biochemical approaches.

In addition, the identification of GAGases may help clarify an inappropriate naming issue
regarding the functional module "Hepar_II_III superfamily". This which functional module is
commonly found in not only GAG lyases but also alginate lyases with an "N-terminal (α/α)$_n$
toroid + C-terminal β-sandwich" structure. Interestingly, however, a large number of identified
alginate lyases with the functional module "Hepar_II_III superfamily" do not have the ability to
degrade Hep/HS. Historically, due to the importance of Hep as an anticoagulant in clinical
application, Hep/HS-related studies including heparinases have received much higher attention
than alginate-related studies including alginate lyases, which may be the reason why some
sequences in alginate lyases that are homologous to heparinase are annotated named the
"Hepar_II_III superfamily" module. From an evolutionary perspective, this "Hepar_II_III
superfamily" module might be a conserved functional domain in the divergent evolution process
from alginate lyase to GAG lyase. Therefore, it might be more reasonable and accurate for this

functional domain to be named as the "alginate lyase superfamily" module, and the presence of
this functional module in GAGases with evolutionary transitional characteristics may well
support this view.

In conclusion, the first determination of the structure of GAGases not only facilitates the
elucidation of the catalytic mechanism of PL35 family enzymes, especially the substrate
selection mechanism of GAGases, but also provides potential evidence for the hypothesis that
GAG lyase evolved from alginate lyase.

Materials and methods
Materials
HA from *Streptococcus equi*, CS-C from shark cartilage, alginate sodium from brown algae,
polyethylene glycol (PEG) 6000, PEG 400, calcium chloride ($CaCl_2$), tris-(hydroxymethyl)
aminomethane (Tris) and hydroxyethyl piperazine ethanesulfonic acid (HEPES) were purchased
from Sigma–Aldrich. Lysine, phenylalanine, threonine, leucine, isoleucine, valine and L-
selenomethionine were purchased from Solarbio Co.,Ltd. (Beijing, China).

Heterologous expression and purification of GAGases and GAGase II selenomethionine
derivant
The cloning and induced expression of PL35 family GAGases were described in previous
study (***Wei, et al.*,*2024***). Briefly, the synthesized expression plasmid pET-30a-GAGases were
transformed into *E. coli* BL21 (DE3) cells and culture in LB broth at 37 °C. Recombinant
enzymes with a 6His-tag were expressed at 16 °C for 16 h by inducing with isopropyl 1-thio-β-
D-galactopyranosid (IPTG) (final concentration of 0.05 mM) at the cell density of OD_{600}=0.6–0.8.
Furthermore, to obtain the Se-GAGase II, cultured cells harboring the pET-30a-GAGase II were
centrifuged at 3,700 g at 4 °C for 5 min, washed with M9 culture medium (17.2 g/L Na_2HPO_4, 3
g/L KH_2PO_4, 2.5 g/L NaCl and 5 g/L NH_4Cl), transferred into M9 culture medium supplemented
with 1 mM $MgSO_4$, 0.1 mM $CaCl_2$, 0.4% (w/v) glucose and cultured at 37 °C until the OD_{600}
reached 0.6. Then, the culture medium was cooled to 22 °C. Additional essential amino acids,
including lysine (100 mg/L), phenylalanine (100 mg/L), threonine (100 mg/L), leucine (50
mg/L), isoleucine (50 mg/L), valine (50 mg/L), and L-selenomethionine (L-SeMet) (50 mg/L),
were added and incubated at 22 °C for 15 min. The proteins contained L-SeMet were also
induced and expressed at 16 °C for 16 h by supplementing with 5 mM IPTG.

After further induced cultivation, cells were harvested, resuspended and disrupted by

sonication, the recombinant proteins in the supernatant of cell lysates are primarily purified by

loading on a nickel affinity column, washing with buffer A containing 10 mM imidazole to

remove impurities, and finally eluting using buffer A containing 250 mM imidazole. After nickel

affinity chromatography, the elutes of GAGase II and GAGase VII were diluted and loaded on a

Q-Sepharose FF column (GE Healthcare) eluted with a gradient concentration of NaCl from 0 to

1 M, and the target proteins were desalted and concentrated through ultrafiltration and further

sub-fractionated by gel filtration on a Superdex G-200 column (GE Healthcare) for crystal

culture. The concentration of each protein was determined using BCA Protein Assay Kit with

bovine serum albumin (BSA) as reference protein (Cwbio, Shanghai).

Activity assay of GAGase II, III, VII and their mutants toward various GAGs and alginate

To evaluate the activity of GAGase II, III, VII and their mutants, CS-C or alginate (30 μg)

was treated with each wild-type enzyme or its mutant (6 μg) in the optimal buffer (50 mM Tris-

HCl, pH 7.0) for 12 h at 40 °C. After inactivated by boiling for 10 min, the products were

analyzed by gel filtration HPLC using a Superdex^{TM} Peptide 10/300 GL column eluted with 0.20

M NH$_4$HCO$_3$ at a flow rate of 0.4 ml/min and monitored at 232 nm using a UV detector, and

online analysis was conducted using LCsolution version 1.25.

To evaluate the activity of GAGase II and GAGase III mutants from the site-directed

mutagenesis of some potential substrate-binding residues, HA, CS-C, or HS (150 μg) was treated

with each wild-type enzyme or its mutant (6 μg) in the optimal buffer (50 mM Tris-HCl, pH 7.0)

for 1 h at 40 °C, and the absorbance of each resultant were determined at 232 nm using a UV

spectrophotometer.

Crystallization, X-ray diffraction and data collection of GAGase II, GAGase VII and Se-
Met-GAGase II

The purified GAGase II (20 mg/ml) was crystallized in 0.1 M HEPES (pH 7.0), 12% (w/v)

PEG 6000 and 0.2 M CaCl$_2$. The purified Se-Met-GAGase II (5 mg/ml) was crystallized in 0.1

M Tris (pH 7.5), 20% (w/v) PEG 6000 and 0.2 M CaCl$_2$. The purified GAGase VII (10 mg/ml)

was crystallized in 0.1 M HEPES (pH 7.0) and 36% (v/v) PEG 400. The crystallization of all the

above proteins was carried out at 18 °C using hanging drop vapor diffusion method in 24-well

plates. Crystals were harvested using nylon loops and soaked in cryoprotectant consisting of 20%

glycerol and 80% mother liquor. X-ray diffraction data were collected at Shanghai Synchrotron

Radiation Facility (SSRF) BL18U1 beamline. Diffraction datasets were processed using XDS

(**Kabsch**,**2010**). Data collection statistics are summarized in *Table 1*.

Structure determination and refinement

The structure of GAGase II was solved by selenium single-wavelength anomalous

dispersion (Se-SAD) using AutoSol from Phenix suite (**Adams, et al.**,**2011**). The structure of

GAGase VII was determined by molecular replacement using Phaser (**McCoy, et al.**,**2007**) with

the structure of GAGase II as the search model. Initial model building was carried out using

AutoBuild from Phenix suite. Then several rounds of refinement and manual building were

performed alternately using Phenix.Refine and Coot (**Emsley and Cowtan**,**2004**), respectively.

The ion type was confirmed by inductively coupled plasma-mass spectrometry (ICP-MS)

analysis. For each sample, 10 ml of GAGase II or GAGase VII (10 mg) was mixed with 10 ml of

nitric acid in equal proportions. The mixture was then placed in a metal bath at 120 °C and

heated until clarified. Subsequently, the clarified mixture was filtered through a 0.22 µm

membrane and analyzed using ICP-MS (NexlON, PerkinElmer). The structures of other

GAGases were predicted using Robetta (**Baek, et al.**,**2021**). Structure alignments were performed

using ChimeraX with point accepted mutation (PAM)-120 mtrix (**Pettersen, et al.**,**2021**). All of

the structure illustrations were generated using ChimeraX (https://www.cgl.ucsf.edu/chimerax/)

and Pymol (https://www.pymol.org/2/). A structure-based homology search for GAGase II and

VII were performed using the DALI server (http://ekhidna2.biocenter.helsinki.fi/dali/) (**Holm**

and Rosenstrom,**2010**). The classification of identified enzyme were confirmed based on the

database of carbohydrate-active enzymes (CAZy) (http://www.cazy.org/) (**Drula, et al.**,**2022**)

and the structures of identified PLs were download from the RCSB Protein Data Bank (PDB)

(https://www.rcsb.org/) (**Berman, et al.**,**2000**).

Site-directed mutagenesis of GAGases

Some potentially important residues located in and around the active center were

individually replaced with alanine (Ala) by using the Fast Mutagenesis Kit V2 from Vazyme

Biotech Co., Ltd (Nanjing, China) and these mutants were expressed and examined for activity

against various substrates (HA, CS-C, HS, and alginate) to confirm their crucial role in enzyme

catalysis. The primer pairs of these mutants are shown in the *Table S2*. The activities of the

mutants were evaluated as described above.

Molecular docking

Tertiary structure-defined GAG or alginate hexasaccharide was used to dock into the active cavity of GAGase II by using the Autodock Vina program (***Eberhardt, et al.,2021***; ***Trott and Olson,2010***). Docking was performed on a search space of 15.00×18.19×19.53 Å covering the active centre (x, y, z = -2.60, -12.57, 34.13). The binding energies using non-sulfated and 4-O-trisulfated hexasaccharides are calculated as -5.874 and -5.163 kcal/mol, respectively.

Data availability

All data supporting the findings of this study are available within the paper (and supporting information files). All relevant data generated during this study or analysed in this published article are available from the corresponding author on reasonable request. The atomic coordinates and structure factors of the structures in this study have been deposited in the Protein Data Bank (PDB codes: 8KHV and 8KHW). The sequences of GAGase II (GenBank: SOD82962.1) and GAGase VII (GenBank: EDV05210.1) have already existed in NCBI database.

Acknowledgment

This work was supported by the National Key R&D Program of China (No. 2021YFC2103100), National Natural Science Foundation of China (Nos. 31971201, 32330001, 31570071, 31800665), Major Scientific and Technology Innovation Project (MSTIP) of Shandong Province (No. 2019JZZY010817), Natural Science Foundation of Shandong Province (No. ZR2023MC017) and the Program of Shandong for Taishan Scholars (tspd20181203).

Author contributions

Wei Lin, Investigation, Validation, Visualization, Writing-Original Draft; Cao Hai-Yan, Investigation, Validation, Visualization, Writing-Original Draft; Zou Ruyi, Investigation, Validation; Du Min, Investigation, Validation; Zhang Qingdong, Investigation; Lu Danrong, Resources; Xu Xiangyu, Investigation; Xu Yingying, Investigation; Wang Wenshuang, Investigation; Chen Xiu-Lan, Investigation; Zhang Yu-Zhong, Conceptualization, Methodology, Writing-Reviewing and Editing; Li Fuchuan, Conceptualization, Methodology, Writing-Reviewing and Editing.

References:
Adams PD, Afonine PV, Bunkóczi G, Chen VB, Echols N, Headd JJ, Hung L, Jain S, Kapral GJ, Grosse Kunstleve

RW, McCoy AJ, Moriarty NW, Oeffner RD, Read RJ, Richardson DC, Richardson JS, Terwilliger TC, Zwart PH.

2011. The Phenix software for automated determination of macromolecular structures. *Methods* **55**:94-106.

doi:10.1016/j.ymeth.2011.07.005

Baek M, DiMaio F, Anishchenko I, Dauparas J, Ovchinnikov S, Lee GR, Wang J, Cong Q, Kinch LN, Schaeffer

RD, Millan C, Park H, Adams C, Glassman CR, DeGiovanni A, Pereira JH, Rodrigues AV, van Dijk AA, Ebrecht

AC, Opperman DJ, Sagmeister T, Buhlheller C, Pavkov-Keller T, Rathinaswamy MK, Dalwadi U, Yip CK, Burke

JE, Garcia KC, Grishin NV, Adams PD, Read RJ, Baker D. 2021. Accurate prediction of protein structures and

interactions using a three-track neural network. *Science* **373**:871-876. doi:10.1126/science.abj8754

Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, Shindyalov IN, Bourne PE. 2000. The protein

data bank. *Nucleic Acids Res* **28**:235-242. doi:10.1093/nar/28.1.235

Charnock SJ, Brown IE, Turkenburg JP, Black GW, Davies GJ. 2002. Convergent evolution sheds light on the anti-

beta-elimination mechanism common to family 1 and 10 polysaccharide lyases. *P Natl Acad Sci Usa* **99**:12067-

12072. doi:10.1073/pnas.182431199

Clementi F. 1997. Alginate production by *Azotobacter vinelandii*. *Crit Rev Biotechnol* **17**:327-361.

doi:10.3109/07388559709146618

Cohen E, Merzendorfer H. 2019. Extracellular sugar-based biopolymers matrices. *Springer International Publishing*

AG, Cham doi:10.1007/978-3-030-12919-4

Csoka AB, Stern R. 2013. Hypotheses on the evolution of hyaluronan: A highly ironic acid. *Glycobiology* **23**:398-

411. doi:10.1093/glycob/cws218

Davies GJ, Wilson KS, Henrissat B. 1997. Nomenclature for sugar-binding subsites in glycosyl hydrolases. *Biochem*

J **321 (Pt 2)**:557-559. doi:10.1042/bj3210557

Drula E, Garron M, Dogan S, Lombard V, Henrissat B, Terrapon N. 2022. The carbohydrate-active enzyme

database: functions and literature. *Nucleic Acids Res* **50**:D571-D577. doi:10.1093/nar/gkab1045

Dyer DP, Salanga CL, Volkman BF, Kawamura T, Handel TM. 2015. The dependence of chemokine–

glycosaminoglycan interactions on chemokine oligomerization. *Glycobiology*:cwv100. doi:10.1093/glycob/cwv100

Eberhardt J, Santos-Martins D, Tillack AF, Forli S. 2021. AutoDock Vina 1.2.0: New docking methods, expanded

force field, and python bindings. *J Chem Inf Model* **61**:3891-3898. doi:10.1021/acs.jcim.1c00203

Emsley P, Cowtan K. 2004. Coot: model-building tools for molecular graphics. *Acta Crystallographica Section D*

Biological Crystallography **60**:2126-2132. doi:10.1107/S0907444904019158

Evans LR, Linker A. 1973. Production and characterization of the slime polysaccharide of *Pseudomonas*

aeruginosa. *J Bacteriol* **116**:915-924. doi:10.1128/JB.116.2.915-924.1973

Fethiere J, Eggimann B, Cygler M. 1999. Crystal structure of chondroitin AC lyase, a representative of a family of

glycosaminoglycan degrading enzymes. *J Mol Biol* **288**:635-647. doi:10.1006/jmbi.1999.2698

Garron ML, Cygler M. 2010. Structural and mechanistic classification of uronic acid-containing polysaccharide

lyases. *Glycobiology* **20**:1547-1573. doi:10.1093/glycob/cwq122

Garron ML, Cygler M. 2014. Uronic polysaccharide degrading enzymes. *Curr Opin Struc Biol* **28**:87-95.

doi:10.1016/j.sbi.2014.07.012

Han Y, Garron M, Kim H, Kim W, Zhang Z, Ryu K, Shaya D, Xiao Z, Cheong C, Kim YS, Linhardt RJ, Jeon YH,

Cygler M. 2009. Structural snapshots of heparin depolymerization by heparin lyase I. *J Biol Chem* **284**:34019-

34027. doi:10.1074/jbc.M109.025338

Hashimoto W, Maruyama Y, Nakamichi Y, Mikami B, Murata K. 2014. Crystal structure of *Pedobacter heparinus*

heparin lyase Hep III with the active site in a deep cleft. *Biochemistry-Us* **53**:777-786. doi:10.1021/bi4012463

Holm L, Rosenstrom P. 2010. Dali server: conservation mapping in 3D. *Nucleic Acids Res* **38**:W545-W549.

doi:10.1093/nar/gkq366

Huang W, Lunin V, Li Y, Suzuki S, Sugiura N, Miyazono H, Cygler M. 2003. Crystal structure of *Proteus vulgaris*

chondroitin sulfate ABC lyase I at 1.9 Å resolution. *J Mol Biol* **328**:623-634. doi:10.1016/S0022-2836(03)00345-0

Huang W, Matte A, Li Y, Kim YS, Linhardt RJ, Su H, Cygler M. 1999. Crystal structure of chondroitinase B from

Flavobacterium heparinum and its complex with a disaccharide product at 1.7 A resolution. *J Mol Biol* **294**:1257-

1269. doi:10.1006/jmbi.1999.3292

Irie F, Okuno M, Matsumoto K, Pasquale EB, Yamaguchi Y. 2008. Heparan sulfate regulates ephrin-A3/EphA

receptor signaling. *P Natl Acad Sci Usa* **105**:12307-12312. doi:10.1073/pnas.0801302105

Itoh T, Nakagawa E, Yoda M, Nakaichi A, Hibi T, Kimoto H. 2019. Structural and biochemical characterisation of a

novel alginate lyase from *Paenibacillus* sp. str. FPU-7. *Sci Rep-Uk* **9**. doi:10.1038/s41598-019-51006-1

Jagtap SS, Hehemann J, Polz MF, Lee J, Zhao H. 2014. Comparative biochemical characterization of three exolytic

oligoalginate lyases from *Vibrio splendidus* reveals complementary substrate scope, temperature, and pH

adaptations. *Appl Environ Microb* **80**:4207-4214. doi:10.1128/AEM.01285-14

Ji S, Dix SR, Aziz AA, Sedelnikova SE, Baker PJ, Rafferty JB, Bullough PA, Tzokov SB, Agirre J, Li F, Rice DW.

2019. The molecular basis of endolytic activity of a multidomain alginate lyase from *Defluviitalea phaphyphila*, a

representative of a new lyase family, PL39. *J Biol Chem* **294**:18077-18091. doi:10.1074/jbc.RA119.010716

Kabsch W. 2010. XDS. *Acta Crystallogr D Biol Crystallogr* **66**:125-132. doi:10.1107/S0907444909047337

Kusche-Gullberg M, Kjellen L. 2003. Sulfotransferases in glycosaminoglycan biosynthesis. *Curr Opin Struc Biol*

13:605-611. doi:10.1016/j.sbi.2003.08.002

Li S, Jedrzejas MJ. 2001. Hyaluronan binding and degradation by *Streptococcus agalactiae* hyaluronate lyase. *J Biol*

Chem **276**:41407-41416. doi:10.1074/jbc.M106634200

Lombard V, Bernard T, Rancurel C, Brumer H, Coutinho PM, Henrissat B. 2010. A hierarchical classification of

polysaccharide lyases for glycogenomics. *Biochem J* **432**:437-444. doi:10.1042/BJ20101185

Lunin VV, Li Y, Linhardt RJ, Miyazono H, Kyogashima M, Kaneko T, Bell AW, Cygler M. 2004. High-resolution

crystal structure of *Arthrobacter aurescens* chondroitin AC lyase: an enzyme–substrate complex defines the

catalytic mechanism. *J Mol Biol* **337**:367-386. doi:10.1016/j.jmb.2003.12.071

Martinez-Fleites C, Smith NL, Turkenburg JP, Black GW, Taylor EJ. 2009. Structures of two truncated phage-tail

hyaluronate lyases from *Streptococcus pyogenes serotype* M1. *Acta Crystallogr Sect F Struct Biol Cryst Commun*

65:963-966. doi:10.1107/S1744309109032813

Matsubara Y, Iwasaki K, Muramatsu T. 1998. Action of poly (alpha-L-guluronate) lyase from *Corynebacterium* sp.

ALY-1 strain on saturated oligoguluronates. *Biosci Biotech Bioch* **62**:1055-1060. doi:10.1271/bbb.62.1055

McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ. 2007. Phaser crystallographic

software. *J Appl Crystallogr* **40**:658-674. doi:10.1107/S0021889807021206

Ndeh D, Baslé A, Strahl H, Yates EA, McClurgg UL, Henrissat B, Terrapon N, Cartmell A. 2020. Metabolism of

multiple glycosaminoglycans by *Bacteroides thetaiotaomicron* is orchestrated by a versatile core genetic locus. *Nat*

Commun **11**doi:10.1038/s41467-020-14509-4

Ochiai A, Yamasaki M, Mikami B, Hashimoto W, Murata K. 2010. Crystal structure of exotype alginate lyase

Atu3025 from *Agrobacterium tumefaciens*. *J Biol Chem* **285**:24519-24528. doi:10.1074/jbc.M110.125450

Pandey S, Mahanta P, Berger BW, Acharya R. 2021. Structural insights into the mechanism of pH-selective

substrate specificity of the polysaccharide lyase Smlt1473. *J Biol Chem* **297**:101014. doi:10.1016/j.jbc.2021.101014

Park D, Jagtap S, Nair SK. 2014. Structure of a PL17 family alginate lyase demonstrates functional similarities

among exotype depolymerases. *J Biol Chem* **289**:8645-8655. doi:10.1074/jbc.M113.531111

Pawar SN, Edgar KJ. 2012. Alginate derivatization: A review of chemistry, properties and applications.

Biomaterials **33**:3279-3305. doi:10.1016/j.biomaterials.2012.01.007

Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, Morris JH, Ferrin TE. 2021. UCSF

ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci* **30**:70-82.

doi:10.1002/pro.3943

Popper ZA, Michel G, Hervé C, Domozych DS, Willats WGT, Tuohy MG, Kloareg B, Stengel DB. 2011. Evolution

and diversity of plant cell walls: from algae to flowering plants. *Annu Rev Plant Biol* **62**:567-590.

doi:10.1146/annurev-arplant-042110-103809

Ronne ME, Tandrup T, Madsen M, Hunt CJ, Myers PN, Moll JM, Holck J, Brix S, Strube ML, Aachmann FL,

Wilkens C, Svensson B. 2023. Three alginate lyases provide a new gut *Bacteroides ovatus* isolate with the ability to

grow on alginate. *Appl Environ Microb* **89**:e0118523. doi:10.1128/aem.01185-23

Sattelle BM, Shakeri J, Roberts IS, Almond A. 2010. A 3D-structural model of unsulfated chondroitin from high-

field NMR: 4-sulfation has little effect on backbone conformation. *Carbohyd Res* **345**:291-302.

doi:10.1016/j.carres.2009.11.013

Shaya D, Hahn B, Bjerkan TM, Kim WS, Park NY, Sim J, Kim Y, Cygler M. 2008. Composite active site of

chondroitin lyase ABC accepting both epimers of uronic acid. *Glycobiology* **18**:270-277.

doi:10.1093/glycob/cwn002

Shaya D, Hahn BS, Park NY, Sim JS, Kim YS, Cygler M. 2008. Characterization of chondroitin sulfate lyase ABC

from *Bacteroides thetaiotaomicron* WAL2926. *Biochemistry-Us* **47**:6650-6661. doi:10.1021/bi800353g

Shaya D, Tocilj A, Li Y, Myette J, Venkataraman G, Sasisekharan R, Cygler M. 2006. Crystal structure of

heparinase II from *Pedobacter heparinus* and its complex with a disaccharide product. *J Biol Chem* **281**:15525-

15535. doi:10.1074/jbc.M512055200

Sirko S, von Holst A, Weber A, Wizenmann A, Theocharidis U, Gotz M, Faissner A. 2010. Chondroitin sulfates are

required for fibroblast growth factor-2-dependent proliferation and maintenance in neural stem cells and for

epidermal growth factor-dependent migration of their progeny. *Stem Cells* **28**:775-787. doi:10.1002/stem.309

Sugiura N, Setoyama Y, Chiba M, Kimata K, Watanabe H. 2011. Baculovirus envelope protein ODV-E66 is a novel

chondroitinase with distinct substrate specificity. *J Biol Chem* **286**:29026-29034. doi:10.1074/jbc.M111.251157

Trott O, Olson AJ. 2010. AutoDock Vina: improving the speed and accuracy of docking with a new scoring

function, efficient optimization, and multithreading. *J Comput Chem* **31**:455-461. doi:10.1002/jcc.21334

Ulaganathan T, Shi R, Yao D, Gu R, Garron M, Cherney M, Tieleman DP, Sterner E, Li G, Li L, Linhardt RJ,

Cygler M. 2017. Conformational flexibility of PL12 family heparinases: structure and substrate specificity of

heparinase III from Bacteroides thetaiotaomicron (BT4657). *Glycobiology (Oxford)* **27**:176-187.

doi:10.1093/glycob/cww096

Wei L, Zou R, Du M, Zhang Q, Lu D, Xu Y, Xu X, Wang W, Zhang YZ, Li F. 2024. Discovery of a class of

glycosaminoglycan lyases with ultrabroad substrate spectrum and their substrate structure preferences. *J Biol Chem*

300:107466. doi:10.1016/j.jbc.2024.107466

Winter WT, Smith PJ, Arnott S. 1975. Hyaluronic acid: structure of a fully extended 3-fold helical sodium salt and

comparison with the less extended 4-fold helical forms. *J Mol Biol* **99**:219-235. doi:10.1016/s0022-2836(75)80142-

2

Yamasaki M, Moriwaki S, Miyake O, Hashimoto W, Murata K, Mikami B. 2004. Structure and function of a

hypothetical *Pseudomonas aeruginosa* protein PA1167 classified into family PL-7. *J Biol Chem* **279**:31863-31872.

doi:10.1074/jbc.M402466200

Yamasaki M, Ogura K, Moriwaki S, Hashimoto W, Murata K, Mikami B. 2005. Crystallization and preliminary X-

ray analysis of alginate lyases A1-II and A1-II′ from *Sphingomonas* sp. A1. *Acta Crystallogr F* **61**:288-290.

doi:10.1107/S174430910500299X

Zhang F, Zheng L, Cheng S, Peng Y, Fu L, Zhang X, Linhardt R. 2019. Comparison of the interactions of different

growth factors and glycosaminoglycans. *Molecules* **24**:3360. doi:10.3390/molecules24183360

Zhang Q, Cao H, Wei L, Lu D, Du M, Yuan M, Shi D, Chen X, Wang P, Chen X, Chi L, Zhang Y, Li F. 2021.

Discovery of exolytic heparinases and their catalytic mechanism and potential application. *Nat Commun*

12doi:10.1038/s41467-021-21441-8

Zhu B, Huang L, Tan H, Qin Y, Du Y, Yin H. 2015. Characterization of a new endo-type polyM-specific alginate

lyase from Pseudomonas sp. *Biotechnol Lett* **37**:409-415. doi:10.1007/s10529-014-1685-0

A

GAGase II

N-terminal (α/α)₇ toroid

C-terminal two-layered β-sheet

GAGase VII

N-terminal (α/α)₇ toroid

C-terminal two-layered β-sheet

B

GAGase II

GAGase VII

(PL17, 4NEI)

(PL39, 6JP4)

(PL15, 3A0O)

(PL12, 4MMH)

(PL15, 6LJA)

(PL21, 2FUQ)

(PL8, 1RWA)

(PL23, 3VSM)

(PL8, 2Q1F)

A

Asn192
Asn187
Asn189

Tyr246
Tyr241
Tyr243

His398
His397
His401

Asn191
His188

GAGase II
GAGase III
GAGase VII

Tyr424
Tyr423
Tyr427

B

Asn201
Tyr258   His200
        His413
Tyr450

PL17 (4NEI)

His187
    Asn186
Tyr239
    His405
Tyr428

PL39 (6JP4)

His243
    Asn242
Tyr298
Tyr91   Arg292

PL38 (8BDQ)

His168
    Asn167
Tyr39   Tyr222
    Arg215

PL5 (7FHU)

Arg314   His311
        Tyr365
        His531
Tyr555

PL15 (3A0O)

C

CS-C

GAGase III

GAGase III-H188A

GAGase III-H188N

Ultraviolet intensity (10^4)

Time (min)

Alginate

GAGase III

GAGase III-H188A

GAGase III-H188N

Ultraviolet intensity (10^4)

Time (min)

D

HA

Relative activity (%)

GAGaseIII   H188N   H188A
***   ***

CS-C

Relative activity (%)

GAGaseIII   H188N   H188A
                *

HS

Relative activity (%)

GAGaseIII   H188N   H188A
            *   **

Non-sulfated CS hexasaccharide   4-O-trisulfated CS hexasaccharide

GAGase II

GAGase III