DNMT1-Mediated Regulation of Inhibitory Interneuron Migration Impacts Cortical Architecture and Function

Julia Reichard[1, 2], Philip Wolff[1], Jian Du[1], Camila L. Fullio[3,4], Lilli Dorp[1], Can Bora Yildiz[1,2], Jenice Linde[1,2], Johanna Vollmer[1], Shirley Kempf[1], Severin Graff[2,5], Gerion Nabbefeld[2,6], Minali Singh[7], Chao-Chung Kuo[8], K. Naga Mohan[7], Tanja Vogel[3], Simon Musall[2,5] and Geraldine Zimmer-Bensch[1, 2]

Affiliations
[1] RWTH Aachen University, Division of Neuroepigenetics, Institute of Zoology (Biology 2), Worringerweg 3, 52074 Aachen, Germany.
[2] Research Training Group 2416 Multi Senses – Multi Scales, RWTH Aachen University, 52074 Aachen, Germany
[3] Institute for Anatomy and Cell Biology, Department of Molecular Embryology, Faculty of Medicine, Albert-Ludwigs-University Freiburg, 79104 Freiburg, Germany
[4] Faculty of Biology, Albert-Ludwigs-University Freiburg, 79104 Freiburg, Germany
[5] Institute of Biological Information Processing (IBI-3) Bioelectronics, Forschungszentrum Jülich, Germany
[6] RWTH Aachen University, Division of Neurophysiology, Institute of Zoology (Biology 2), Worringerweg 3, 52074 Aachen, Germany.
[7] Molecular Biology and Genetics Laboratory, Department of Biological Sciences, BITS Pilani, Hyderabad Campus, Hyderabad, India
[8] Genomics Facility, Interdisciplinary Center for Clinical Research (IZKF), RWTH Aachen University, 52074 Aachen, Germany

Abstract
The fine-tuned establishment of neuronal circuits during the formation of the cerebral cortex is pivotal for its functionality. Developmental abnormalities affecting the composition of cortical circuits, which consist of excitatory neurons and inhibitory interneurons, are linked to a spectrum of neuropsychiatric disorders. Excitatory neurons originate in cortical proliferative zones, while inhibitory interneurons migrate from discrete domains of the basal telencephalon into the cortex. This migration is intricately governed by intrinsic genetic programs and extrinsic cues. Our current study reveals the role of the DNA methyltransferase 1 (DNMT1) in controlling expression of key genes implicated in mouse cortical interneuron development and in guiding the migration of somatostatin-expressing interneurons within the developing cortex. *Dnmt1* deletion causes interneurons to exit prematurely from the superficial migratory stream. In addition to the perturbed migration pattern and altered gene expression signatures, *Dnmt1*-deficient interneurons had a discernible non-cell autonomous effect on cortical progenitors, which culminated in nuanced alterations of layer thicknesses in the adult cortex. Our study reveals that an epigenetic mechanism governs the migration of cortical interneurons and through this, their instructive role in sculpting the intricate cortical layer architecture by signaling to cortical progenitors, with pronounced effects on cortical network function.

Introduction

The intricate neuronal circuitry of the mammalian cerebral cortex, composed of excitatory and inhibitory neurons, empowers its enormous cognitive capabilities. The circuit formation involves coordinated developmental steps: proliferation, differentiation, migration, morphological maturation, and synapse formation[1].

Excitatory cortical projection neurons originate from radial glial cells (RGCs) and intermediate progenitors (IPCs) located in the cortical proliferative zones, through direct and indirect neurogenesis. Symmetric divisions of RGCs initially expand the progenitor pool, later shifting to asymmetric divisions to produce neurons and RGCs or IPCs [2;3]. EOMES-expressing IPCs localize to the subventricular zone (SVZ) and divide into neurons for all cortical layers after few self-renewal divisions[3]. Cortical excitatory neurons are produced in a temporally regulated manner, with early-born neurons forming deep layers and later-born neurons forming superficial layers, coming with unique functional features and connectivity[4]. The timing of cortical neuron generation seems to be regulated by both intrinsic and extrinsic factors[2;3]. Intrinsic factors include (epi)genetic programs within progenitor cells that dictate their developmental potential. Extrinsic factors encompass signaling molecules from neighboring cells, postmitotic cells in the cortical plate, embryonic cerebrospinal fluid secreted by the choroid plexus, and incoming thalamic afferents[5–9]. These signals influence the self-renewal and neurogenic potential of cortical progenitors, ensuring the proper temporal patterning and layering of the cortex.

Despite composing only 10-20% of the neuronal population, cortical inhibitory neurons (cINs), including parvalbumin (PV) and somatostatin (SST) interneuron subtypes, crucially modulate cortical activity by inhibiting both excitatory and other inhibitory neurons, shaping cortical circuits for precise information processing[10]. Disruptions in their development unbalance excitation and inhibition, causing abnormal cortical activity and cognitive impairments in various neurological and neuropsychiatric disorders, such as schizophrenia or epilepsy[11–14]. The different types of cINs originate in three subpallial structures: the medial ganglionic eminence (MGE), the caudal ganglionic eminence (CGE), and the preoptic area (POA)[10]. The MGE gives rise to PV+ and SST+ interneurons, whereas the CGE generates Serotonin 3A receptor (5HTR3a) subtypes[15,16]. The POA produces mostly neurogliaform cells and Neuropeptide Y (NPY) expressing multipolar interneurons[17,18]. Postmitotic cINs migrate tangentially through the basal telencephalon into the cortex, guided by various extrinsic cues[19–22]. They invade the cortex along the marginal zone (MZ) and subventricular zone (SVZ)/intermediate zone (IZ), where they spread tangentially before switching to radial migration to enter the cortical plate. This migration pattern is under the influence of a diverse set of intrinsic and extrinsic cues[12]. For example, the transcription factors ARX and LHX6, expressed in MGE-derived interneurons, directly and indirectly regulate the expression of cytokine receptors that mediate the attraction to CXCL12, which is expressed in the MZ migratory route of the developing cortex[23,24]. After tangentially spreading over the cortical areas, cINs invade the cortical plate. The cortical plate expresses NRG2, which exerts attractive effects on ERBB4-expressing interneurons[25].

Epigenetic mechanisms, including histone modifications and DNA methylation, modulate intrinsic transcriptional programs[14,26]. It was previously reported that the DNA methyltransferase 1 (DNMT1) regulates gene expression programs in POA-derived cINs, promoting their proper migration and survival[27] through non-canonical interactions with histone

modifications[27]. MGE-derived cINs, which give rise to SST[+] and PV[+] interneuron subtypes also
express DNMT1[27]. However, DNMT1 function in these cIN subsets is unknown so far. SST-
expressing interneurons constitute about 30% of all cINs and play vital roles in inhibitory
control, network synchronization, circuit plasticity, and cognitive functions[28]. SST[+] cINs
encompass two main subtypes: Martinotti cells, which migrate along the marginal zone (MZ)
and form axon collaterals in layer I, and non-Martinotti cells, which disperse via the
subventricular/intermediate zones (SVZ/IZ) and have different layer distributions as well as
targeting properties[12,28]. Expression of transcription factors, including ARX, LHX6, and MAFB,
correlates with the two different migratory paths[23,29,30]. MAFB marks Martinotti cells, mediating
together with ARX[23] and LHX6[29] the migration of cINs along the MZ[30]. The migration of SST[+]
cINs along the MZ is linked to the axonal arbor formation in layer I[30]. Despite the critical roles
of SST[+] interneurons in cortical circuitry and their likely reliance on epigenetic regulation for
proper development, the specific mechanisms by which DNMT1 influences SST[+] cIN migration
and its subsequent impact on cortical structure and function remain poorly understood.
Additionally, given the temporally and spatially orchestrated migration patterns of SST[+] cINs
within the cortex, it remains unclear how the migration of this subpopulation might influence
the proliferation and differentiation of cortical progenitors. Addressing this gap will help to
understand how cell type-specific epigenetic regulation orchestrates the precise formation of
cortical networks.
Recognizing the functional relevance of SST[+] cINs, we here investigated the role of DNMT1 in
their development. We found that reduced *Dnmt1* expression in SST[+] interneurons altered the
transcriptional and DNA methylation profiles of key genes for cIN development such as *Arx*.
Thereby, *Dnmt1* deletion altered cIN migration along the MZ in the developing cortex. This
impacted on the excitatory cortical progenitor pool and neurogenesis at specific developmental
stages, and led to changes in cortical layering, functionality, and behavioral abnormalities in
adulthood.
117
118 **Results**
119
DNMT1 regulates the expression of key genes during SST[+]-cIN development
To examine the role of DNMT1 specifically in the SST-expressing subtypes of cINs, we created
a conditional knockout mouse model (*Sst-Cre/tdTomato/Dnmt1 loxP[2]*) with *Dnmt1* deletion
being induced by *Cre*-expression in post-mitotic SST[+] interneurons primarily originating in the
MGE[31]. *Sst-Cre/tdTomato* mice served as controls (**Extended Data Fig. E1a**). The validation
of the mouse models is shown in **Extended Data Fig. E1b-h**. We first focused on embryonic
day (E) 14.5, a critical time when extensive migration of SST[+] cINs occurs through the basal
telencephalon, marking the beginning of their invasion into the cerebral cortex[31]. To reveal
putative target genes of DNMT1, we conducted RNA sequencing and methyl-sequencing of
FACS-enriched E14.5 *Sst-Cre/tdTomato/Dnmt1 loxP[2]* and *Sst-Cre/tdTomato* cells, to
determine changes in transcriptional and DNA methylation profiles, respectively. We found
1080 differentially expressed genes (DEGs) and 9919 differentially methylated sites (DMSs)
in E14.5 *Dnmt1* deficient *Sst-Cre/tdTomato* cells (**Fig. 1a-c, Supplementary Table 1**). As
differentially methylated regions (DMRs) encompass broader genomic regions, such as
promoters, enhancers, and other regulatory elements[32,33], they often hold greater regulatory
significance than single DMSs. To account for this, we also calculated DMRs, identifying a

total of 23622 regions. Correlating these DMRs to the DEGs, we identified 600 genes that exhibit both alterations in gene expression (up- and downregulation) and DNA methylation in *Sst-Cre/tdTomato/Dnmt1 loxP2* cells (**Fig. 1d, Supplementary Table 2 and 3**). 349 of these 600 genes increased in expression upon *Dnmt1* deletion (**Fig. 1d**), pointing to putative targets of DNMT1-dependent repressive DNA methylation in SST+ interneurons of control mice. In support of this, 66 of these 349 genes exhibited DNMT1 binding motifs that we have identified through additional ChIP-sequencing experiments (**Fig. 1d, g, Supplementary Table 4**).

Gene Ontology (GO) analysis on genes, which increased significantly in *Sst-Cre/Dnmt1* KO cells and concomitantly exhibited changes in DNA methylation, revealed a significant enrichment of genes that are functionally associated with nervous system development, neuron differentiation, cell-cell signaling, locomotion, neurogenesis, and cell differentiation (**Fig. 1f, Supplementary Table 5).** Similar results were observed for the entire set of significantly upregulated genes in *Dnmt1* KO samples, regardless of concurrent changes in DNA methylation (**Supplementary Table 6**). Notably, among the genes with altered expression and DNA methylation, we identified key regulators of cortical interneuron differentiation and maturation, such as *Arx, Zeb, Mef2c, Dlx1, Sox6, Mafb,* and *Lhx6*. In addition to these well-known transcription factors that govern cIN fate and development[12,31,34], we found locomotion and cell-cell signaling-related genes to be regulated by DNMT1 (**Fig. 1b, f, h**). These included *Erbb4, Sema3A,* and *Robo2,* as well as members of the Eph/ephrin family such as *Efnb2*, which encode proteins that are crucial for directional guidance and neuronal positioning[25,35,36,37]. Of note, *Erbb4, Mafb, Arx, Lhx6,* and *Sox6,* were amongst the most highly differentially expressed genes compared to controls (**Fig. 1a, b**). This positions DNMT1 as a crucial epigenetic regulator of these key genes in cIN development. Moreover, these genes along with numerous others that upregulated upon *Dnmt1* deletion in E14.5 SST+ cINs, were downregulated in embryonic stem cell (ESC)-derived *Dnmt1tet/tet* neurons that overexpressed *Dnmt1* (**Supplementary Table 6**). In general, a significant proportion of all differentially expressed genes upon *Dnmt1* deletion in E14.5 SST+ interneurons (up- and downregulated), were also dysregulated in an opposite direction in these neurons overexpressing DNMT1 (**Fig. 1e, Supplementary Table 6, 7**).

Remarkably, the DNMT1-regulated gene sets displayed significantly longer coding sequences, transcript lengths, genome spans as well as lengths of the 3´UTRs and 5`UTRs compared to the whole set of genes that were detected in SST+ cINs. Moreover, compared to the reference gene list of all detected transcripts, DNMT1-dependent DEGs contained significantly more protein-coding genes, as well as exons and isoforms (*Chi-squared test*; **Extended Data Fig. E2, E3, E4, E6**). Many genes involved in neuronal function and development are known to be particularly long[38], further supporting our finding that DNMT1-dependent DNA methylation regulates key neurodevelopmental genes (**Fig. 1a, f, Supplementary Table 1-7**). In line with this, the deletion of MeCP2, a reader of DNA methylation[39], also results in the upregulation of such long genes[38].

Dnmt1 deletion in SST+ interneurons resulted in differentially methylated sites near transcriptional start sites (TSS) as well as within intragenic regions and intergenic loci (**Figure 1i, Supplementary Table 2, Extended Data Fig. E4**). Intragenic DNA methylation associates with alternative promoter choice and alternative splicing, while DNA methylation at promoter regions represses transcription[40]. We therefore plotted changes in DNA methylation for the aforementioned key genes of interneuron development at the TSS due to its well-documented

impact on gene transcription (**Fig. 1i**). Consistent with previous studies[27,41], we observed sites
with decreased and increased CpG methylation in *Dnmt1* deficient interneurons, also in
specific regions around the TSS (**Fig. 1i, Supplementary Table 2**, **Extended Data Fig. 4**).
These changes in DNA methylation aligned with genes that increased or decreased
significantly in expression in both *Dnmt1* knockout and in *Dnmt1* overexpressing neurons (**Fig.**
1a-c, d, e; **Supplementary Table 7**)[27,41].
For *Arx,* we detected strongly reduced DNA methylation at the TSS, as well as in up- and
downstream regions in *Sst-Cre*/*Dnmt1* KO samples (**Fig. 1i, j**), which correlated well with
increased *Arx* expression detected in *Sst-Cre*/*Dnmt1* KO cells. Moreover, the regions around
the TSS of *Zeb2* and *Sox6* displayed reduced methylation (**Fig. 1i**), two genes that also
increased significantly in expression (**Fig. 1a, b, h**). These genes encode proteins with key
roles in (SST+) cIN development. ARX is crucial for cIN migration within the marginal zone
(MZ)[23], and SOX6 is required for the proper positioning of basket and Martinotti neurons[42].
ZEB2 promotes the differentiation and migration of cortical GABAergic interneurons[43,44].
Apart from these key transcription factors that orchestrate cIN differentiation and migration, we
observed elevated expression levels for *Erbb4* upon *Dnmt1* deletion **(Fig. 1a, b, i)**. Elevated
ERBB4 expression at protein level was confirmed by immunostaining in dissociated MGE cells
(E14.5) following 24-hour treatment with *Dnmt1* siRNA oligos (**Extended Data Fig. E6a-c**).
ERBB4 mediates the invasion of cortical interneurons into the cortical plate upon activation
by its ligand Neuregulin (NRG3)[25], which is expressed within the cortical plate. Hence, we
hypothesized that DNMT1 contributed to the proper migration of SST+ interneurons along the
cortical migratory streams and timing of the tangential to radial switch in migrating cIN by
limiting *Erbb4* expression levels.
In sum, these results strongly suggest that DNMT1 contributes to the establishment of proper
gene regulatory networks that direct the migration and differentiation of SST+ interneurons.
206
***Dnmt1* deletion in postmitotic SST+ interneurons causes a premature exit from the**
superficial migratory stream in the embryonic cortex
To investigate the function of DNMT1 in the development of SST+ cINs in more detail, we next
analyzed the migration of *Sst-Cre/tdTomato/Dnmt1 loxP²* and *Sst-Cre/tdTomato* cells at
different embryonic stages. We initially focused on embryonic stage (E)14.5, when first SST+
interneurons have reached the cerebral cortex (**Fig. 2a-e**, **Extended Data Fig. E6d, e**). In situ
examination of *tdTomato*-expressing cells in coronal brain sections from E14.5 revealed
overall comparable numbers of SST+ cINs across the cortex and within the basal
telencephalon in both genotypes (**Fig. 2b-e**). Yet, we observed a marked reduction of SST
interneurons within the marginal zone (MZ) in conditional *Dnmt1* knockout embryos, while the
proportion of *Sst-Cre/tdTomato/Dnmt1 loxP²* neurons within the cortical plate was significantly
increased (**Fig. 2f-h**). To ensure the comparability of the two genotypes and to exclude slight
age differences, we also analyzed the dimensions of all brain sections, which revealed no
variations (**see Extended Data Fig. E7**). Live-cell imaging of organotypic brain slices at E14.5
for 20 hours revealed an increased proportion of *Dnmt1*-deficient SST+ interneurons that
deviated from the superficial migratory stream and entered the cortical plate (CP; **Fig. 2i-k, o**
and p, Supplementary Movies 1 and 2), which aligned with the in situ observations (**Fig. 2f-**
h). Notably, *Dnmt1*-deficient SST+ interneurons displayed a significantly increased frequency
of directional changes during their cortical migration (**Fig. 2l**), suggesting a disruption in their

ability to remain confined within the MZ stream. Moreover, their migratory pace was enhanced, which was particularly evident for radially migrating cells, and cells migrating through the cortical plate (**Fig. 2m-p**). This aligns with an increased path length seen for the migrating KO cells in the living brain slices (**Extended Data Fig. E6d, e**). Of note, in contrast to *Dnmt1*-deficient POA-derived *Hmx3*-expressing interneurons[27], we did not observe morphological abnormalities for migrating *Sst-Cre/tdTomato/Dnmt1 loxP²* cells (see **Extended Data Fig. E6f**). This aligns with in vitro experiments aiming to assess the morphology of *Dnmt1* siRNA-treated MGE cells (E14.5 +1div), in which we did not detect morphological differences compared to control transfections (**Extended Data Fig. E6g-i**). In combination with the sequencing data, our results point to a DNMT1-dependent regulation of SST⁺ interneuron migration by ensuring their proper movement along the superficial stream along the MZ of the cortex. MZ migrating Martinotti cINs express distinct sets of genes compared to the SVZ migrating non-Martinotti cells[30]. SVZ cells show strong expression of e.g. *Sox1*, *Npy*, *Sp9, Tox,* and *Gabra2*, which in turn are repressed in MZ cells[30]. Moreover, *Erbb4* expression was recently described to be expressed in non-Martinotti cells[45]. Interestingly, we found these genes to be repressed by DNMT1 in SST⁺ cINs (**Supplementary** Table 1). This proposes that DNMT1 mainly influences the migration of Martinotti cells.

Although *Dnmt1* KO SST⁺ interneurons prematurely exited the superficial migratory stream and entered the cortical plate by E14.5 within 20 hours (**Fig. 2i-k**, **o and p**), we did not observe significant differences at later stages (E16.5 and E18.5). This was neither the case for the proportion of cells that reached the cortex nor for the absolute number, density, or distribution of *Sst-Cre/Dnmt1* KO cells within the cortical regions quantified (**Fig. 3**). The cortical dimensions were likewise similar between knockout and wildtype E16.5 and E18.5 embryos (**Extended Data Fig. E8**), confirming that the embryos analyzed were at comparable developmental stages. This suggests that the impact of *Dnmt1* deletion on the directional migration of SST⁺ interneurons within the MZ may be transient.

DNMT1-dependent regulation of SST⁺ interneuron migration affects cortical progenitors non-cell autonomously

The development of cIN and cortical excitatory neurons is intricately linked[46,47], and supernumerary cINs in the IZ were already described to elicit altered IPC numbers impacting the generation of upper layer excitatory neurons[47]. Observing an altered migration pattern of SST⁺ interneurons invading the cortical plate and proliferative zones prompted us to investigate its influence on cortical progenitors and neurogenesis. Moreover, we found altered expression of genes involved in cell-cell signaling, known to modulate cortical neurogenesis, such as *efnB2*[48]. EphrinB2 can bind to EPHA4[49], being expressed by RGCs[5]. Activation of EPHA4 receptors by ephrinA5, expressed in invading thalamic afferents, has been shown to regulate RGC division, impacting cortical layer formation[5]. The premature exit from the superficial marginal stream in combination with altered expression of signaling molecules in *Sst-Cre/tdTomato/Dnmt1 loxP²* could therefore alter cell-cell signaling between interneurons and cortical progenitors, potentially affecting neurogenesis.

To investigate this question, we employed immunostaining for TBR1, labeling early-born neurons of the preplate and layer VI[50,51], to assess whether the generation of these neurons is

altered in *Sst-Cre/tdTomato/Dnmt1 loxP²* embryos. Indeed, we found a significant increase in
TBR1⁺ cells within the cortical plate (CP) in E14.5 *Sst-Cre/Dnmt1* KO embryos (**Fig. 4 b, c**),
which became even more evident at E16.5 (**Fig. 4f, g**). At E18.5 the radial extension of both
layer VI (TBR1+ cells) and the remaining CP were significantly expanded in the conditional
Dnmt1 KO embryos compared to age-matched controls (**Fig. 4j-m**). EOMES⁺ IPCs represent
an intermediate progenitor stage from RGCs to neurons, proposed to give rise to most cortical
projection neurons of all cortical layers[52;53,54]. Altered cIN numbers in the embryonic cortex
have already been reported to induce altered numbers of IPCs[47]. In line with that, we detected
increased numbers of EOMES⁺ IPCs at E14.5 in *Dnmt1* deficient animals, aligning temporally
with the premature exit of *Dnmt1*-deficient SST interneurons from the MZ and the increased
numbers of TBR1⁺ neurons (**Fig. 4d, e**).
As no differences in the pool of IPCs were detected at E18.5 (**Fig. 4h, i**), the deletion of *Dnmt1*
therefore seems to elicit a temporally restricted non-cell autonomous effect on cortical IPCs
and the production of deep-layer neurons.
285
***Dnmt1*-deficient SST⁺ interneurons display changes in transcription of genes known to**
modulate cortical progenitors
The non-cell autonomous effect of *Dnmt1* deletion in SST⁺ cINs on cortical progenitors and the
generation of deep-layer neurons suggested, that cINs provide signaling molecules that act on
cortical progenitors, similar to what was reported for thalamic afferents that import ephrinA5,
which influences IPC generation from RGCs through the EPHA4 receptor[5]. *Dnmt1* deficiency
in SST⁺ cINs altered the expression of diverse signaling molecules that are known to modulate
cortical neurogenesis and progenitor division, including *Wnt7a*, *Notch1* and *Notch3*, *Robo2*,
efnB2 and *Sema3A* (**Supplementary Table 1**) [6,55,56;57].
To analyze in detail which signaling molecules play a role in potential interactions between
embryonic cortical interneurons and cortical progenitors, we took advantage of the *CellChat*
bioinformatic tool, which enables quantitative extraction of cell-cell communication networks
from single cell (sc)RNA-sequencing datasets. We exploited scRNA sequencing data from the
E14.5 dorsal telencephalon from control mice to predict key outgoing and incoming signals for
SST⁺ cINs and cortical progenitor communication. This analysis allowed to assess
quantitatively the probability of ligand-receptor interactions between specific cell types,
because each cell type exhibits distinct sets of ligands and receptors with varying strengths,
indicating their role as signaling senders and receivers. The probabilities of ligand-receptor
pairs predicted to mediate the communication between SST⁺ cINs with apical progenitors
(APs) as well as with IPCs at E14.5 are shown in **Fig. 4 n, o**. The highest probability for
interactions between SST⁺ cINs with both APs and IPCs was calculated for NRXN3 – NLGN1
signaling. *Nrxn3* was upregulated in E14.5 *Dnmt1*-deficient SST⁺ interneurons (and displayed
decreased transcript levels in *Dnmt1* overexpressing neurons; **Supplementary Table 1, 7**). In
addition, EFNB2/EPHA4 signaling was predicted to occur between *efnB2* expressing SST⁺
MGE-interneurons, and APs as well as IPCs at E14.5 (**Fig. 4 n-o**). *Efnb2* was also upregulated
in E14.5 *Dnmt1*-deficient SST⁺ interneurons and dysregulated in *Dnmt1* overexpressing
neurons (Fig. 1h, **Supplementary Table 1, 7**). EFNB2 acts as a pro-neurogenic cue in the
embryonic neocortex[48], and can interact with EPHA4 receptors[58]. EPHA4 is expressed by
cortical RGCs and known to be activated by EFNA5, which is imported by thalamic afferents.
This interaction regulates the AP division mode, and through this IPC generation and the

neuronal output for deep versus upper cortical neurons[5]. Interestingly, EFNB2-mediated increase in neuronal production is temporally restricted to the mid-corticogenesis stage (E15.5)[48]. As we observed a transient defect of MZ migrating *Dnmt1*-deficient SST[+] interneurons in the E14.5 cortex, and increased *efnB2* expression level in these cINs, we argue that EFNB2 imported by SST[+] interneurons could contribute to the regulation of the timed generation of cortical neurons. Thus, we hypothesized that the increased expression of *efnB2* in *Dnmt1*-deficient SST[+] interneurons could lead to increased differentiation of APs towards IPCs and TBR1[+] neurons at E14.5 and E16.5. To address this question, we performed a pair-cell assay according to a previous study[5], stimulating E14.5 dissociated cortical neurons plated at clonal density with recombinant ephrinB2-Fc or control-Fc protein for 24 h. Immunostaining against nestin and β-III-tubulin was used to identify APs and postmitotic neurons, respectively[5] (**Fig. 4p**). This assay confirmed an increased proportion of neurogenic RGC divisions upon ephrinB2-Fc stimulation (**Fig. 4q**). Moreover, we analyzed whether ephrinB2-Fc stimulation likewise impacts the generation of EOMES[+] IPCs from RGCs, by immunostaining against nestin and EOMES. Similar to the increase in asymmetric neurogenic divisions of RGCs, we observed an augmented proportion of IPC/RGC pairs (**Fig. 4r, s**). In sum, these results indicate that ephrinB2-Fc treatment stimulates RGCs to asymmetrically divide into neurons and IPCs. In summary, our results indicate an essential role for DNMT1 in the migration of SST[+] interneurons, regulating the expression of key genes orchestrating cIN development but also genes related to cortical neurogenesis. *Dnmt1* deletion in SST[+] interneurons alters their migration trajectory along the MZ causing a premature invasion into the cortical plate and of the cortical IZ-VZ. Likely in conjunction with increased expression of pro-neurogenic factors, the altered cIN migration seems to influence non-cell autonomously neural differentiation of RGCs towards IPCs and deep-layer neurons.

Dnmt1 deletion-induced changes at embryonic stages manifest in an altered cortical architecture at adult stages

Finally, we investigated whether the DNMT1-related changes in SST[+] interneuron migration, cortical progenitors and in deep layer neuron generation results in long-lasting structural changes in the adult cortex. We therefore conducted a comparative analysis of the distribution, density, and fate of *Sst-Cre/tdTomato* and *Sst-Cre/tdTomato/Dnmt1 loxP[2]* interneurons in adult mice. Immunohistochemistry in adult brain sections confirmed SST expression, and in addition the co-expression of NPY and calretinin in *Sst-Cre/tdTomato* cells of both genotypes (**Extended Data Fig. E9**), in line with other studies[59]. However, a difference between the genotypes was observed for the proportion of SST[+] cINs co-expressing PV, which was significantly increased in *Sst-Cre/tdTomato/Dnmt1 loxP[2]* mice compared to controls (**Fig. 5a-c**). This is in line with the elevated transcript levels of *Mef2c* found in E14.5 *Sst-Cre/tdTomato/Dnmt1 loxP[2]* cells (**Fig 1h**). MEF2C is essential for the development of PV[+] interneurons, as it initiates distinct transcriptional programs in immature PV[+] interneurons upon settling within the cortex by opening the chromatin landscapes at PV-specific loci[60,61]. Of note, *Mef2c* expression is driven by MAF and MAFB, which also increased in expression upon *Dnmt1* deletion[62] (**Fig 1h**). Hence, by repressing *Maf* and *Mefc2* expression DNMT1 appears to promote the fate of SST[+] cINs.

While the comparative analysis of *Sst-Cre/tdTomato* cIN numbers across all cortical layers
revealed no major differences between the genotypes (**Fig. 5d-f**), cell density in the superficial
cortical layers was significantly increased in *Sst-Cre/tdTomato/Dnmt1 loxP²* mice (**Fig. 5f**).
CUX1 immunostaining, labeling layer II-IV neurons, was utilized to discriminate between the
superficial layers II-IV from deep layers V and VI (**Fig. 5d, e**). This staining visualized a reduced
radial extension of the CUX1-positive superficial layers II-IV in *Sst-Cre/tdTomato/Dnmt1 loxP²*
mice (**Fig. 5g**), explaining the increased SST⁺ cIN densities (**Fig. 5f**). In contrast, the deep
layers V and VI were increased in their radial extension, while the overall cortical thickness did
not differ between the knockout and control mice (**Fig. 5g**). This shift in the radial extension of
the deep versus upper cortical layers is consistent with the increased numbers of TBR1 positive
neurons observed at E14.5 and E16.5. These findings suggest that the non-cell autonomous
effect of *Dnmt1* deletion in SST⁺ interneurons on cortical progenitors and the timed neuronal
production manifests in changes of the adult cortical architecture, which in turn affects SST⁺-
interneuron densities in the upper cortical layers.
374
***Sst-Cre/tdTomato/Dnmt1* KO mice display functional and behavioral abnormalities**
To analyze whether the observed structural changes in the adult cortex also translate into
functional changes, we conducted high-density Neuropixels recordings in the primary
somatosensory barrel cortex (S1) in awake head-fixed *Sst-Cre/tdTomato/Dnmt1 loxP²* and
Sst-Cre/tdTomato control mice. Mice were placed on a running wheel during passive tactile
stimulation, consisting of short air puffs to the whisker pad[63] (**Fig. 5h**). Neuropixels can
simultaneously capture the single-cell activity of many neurons across the entire cortical depth,
representing a powerful tool to investigate neural activity with high resolution and scale. Tactile
stimulation induced a clear transient response in average local field potentials (LFPs) across
all cortical layers of SST-Cre control mice. In contrast, LFP responses in conditional *Dnmt1*
KO mice were much weaker and temporally delayed (**Fig. 5i, k**), suggesting a disruption in
sensory processing. Moreover, LFP responses in *Sst-Cre/tdTomato/Dnmt1 loxP²* mice did not
exhibit a positive rebound after the initial negative deflection, potentially due to a lack of feed-
forward inhibition (**Fig. 5k**).
To better assess the spatiotemporal structure of synaptic inputs to the cortical column, we also
employed a current source density (CSD) analysis[64] (**Fig. 5j**). The earliest responses for *Sst-*
Cre/tdTomato mice (left) occurred in the granular layer at ~400 μm cortical depth and then
spread to layer 2/3 (100-300 μm) and layer 5 (500-900 *μ*m). These responses were in good
agreement with the literature[65], suggesting intact spatiotemporal processing of sensory
signals. In contrast, the CSD profile in *Sst-Cre/tdTomato/Dnmt1 loxP²* mice was strongly
disrupted and showed no clear spatial pattern across cortical layers in response to tactile
stimulation. These results strongly suggest that the DNMT1-related structural changes in the
adult cortex also cause a marked disruption of cortical network function.
We next used spike-sorting to extract the spiking activity of cortical S1 neurons to quantify their
temporal precision in response to tactile stimulation. Consistent with the LFP results, the
peristimulus time histogram (PSTH) showed weaker spiking responses for all neurons that
responded to the tactile stimulus in *Dnmt1* KO mice compared to control mice. This was
particularly evident in the first neural response within 30 ms after sensory stimulation (**Fig. 5l**).
Interestingly, sensory responses in *Dnmt1* KO mice were much longer lasting compared to
control mice, with neural responses extending up to 400 ms after sensory stimulation (**Fig. 5l,**

m, o). As indicated by our LFP results, this could be due to a lack of SST-related feedback inhibition which prevented the overextended activation of excitatory neurons in control mice. Moreover, tactile spiking responses were also significantly broader in *Dnmt1* KO mice, shown by the full width at half maximum (FWHM) of the initial responses to sensory stimulation (**Fig. 5m**), again pointing to a reduction in the temporal precision of sensory processing in *Dnmt1* KO versus control mice.

To further test for functional changes in the adult cortex of *Dnmt1*-KO mice, we also measured neural responses to 5s-long visual grating stimuli in the primary visual cortex (V1, **Fig. 5p, q**). Visual stimulation increased neural activity and resulted in strong cortical network oscillations in the gamma range, especially around 30 Hz and 70 Hz (**Fig. 5q**). Such gamma (γ) oscillations from 30-120 Hz are essential for integrating neural networks within and across brain structures during cognitive processes, and strongly rely on accurate function of SST⁺ cINs[66,67]. Abnormalities in γ oscillations are therefore a signature of dysfunction in neuronal networks and are often disturbed in cognitive diseases such as schizophrenia, Alzheimer´s disease and Fragile X syndrome[68].

Thus, changes in γ-oscillations represent a useful marker of function and dysfunction in cortical circuit operations[69]. In line with the impaired neural responses to tactile stimulation in S1, we found that gamma oscillations in *Dnmt1*-KO mice were strongly reduced, while responses in lower frequency bands were increased compared to control mice (**Fig. 5q**). These clear changes in cortical oscillations further demonstrate that DNMT1 deletion results in dysfunctional neural networks in the adult cortex, most likely due to an impairment of SST⁺ cIN function and structural changes in cortical network architecture.

Based on these observations, we next asked whether behavioral outputs were altered correspondingly. Moreover, we noted abnormal eye movements during the tests on the running wheel (**Supplementary Video 3**), and frequent jumping of *Sst-Cre/tdTomato/Dnmt1 loxP²* mice in their home cages (**Supplementary Video 4**). These behavioral abnormalities suggest defects related to neurological or neuropsychiatric diseases[70]. We additionally exploited the curated MGI function in ShinyGO, which enables insights into phenotypes, or disease associations linked to given gene sets. Analysis of the upregulated genes in *Dnmt1*-deficient SST⁺ interneurons using the curated MGI function resulted in numerous terms related to behavioral abnormalities and seizure phenotypes (**Extended Data Fig. E9e).** Given the reported implication of cortical SST⁺-interneuron dysfunction in various neuropsychiatric and neurological disorders, such as schizophrenia and autism spectrum disorders[71], we tested for disease-relevant behavior. We first scored nest-building performance, as defects in nest-building behavior can indicate general distress resulting from disease-related impairments[72]. In line with our functional results, we found that *Sst-Cre/tdTomato/Dnmt1 loxP²* mice showed a reduced performance and diminished interest in nest-building material (**Fig.6a-e**). Furthermore, changes in SST⁺ cIN activity and function have been reported to cause epileptic events[71], and γ oscillations are proposed to play a role in seizure generation[73]. Hence, we next conducted pharmacological interventions by administering PTZ (pentylenetetrazol), which is widely utilized to examine epilepsy in animal models and to assess changes in seizure responses[74]. Our findings revealed prolonged seizure durations for *Sst-Cre/tdTomato/Dnmt1 loxP²* mice, which point to alterations in cortical inhibition (**Figure 6f-h**, Supplementary Videos 5-11).

449 In summary, our cellular, functional, and behavioral results demonstrate that *Dnmt1* deletion
450 in postmitotic SST⁺ interneurons leads to altered cortical architecture and network function in
451 adult mice. This strongly suggests that DNMT1 plays a crucial role in the development of SST⁺
452 interneurons and has non-cell autonomous functions essential for the proper formation of the
453 cerebral cortex.
454
Discussion
Two novel and main features of cortical development arise from our experiments: (i) dispersion
of SST⁺ interneurons along the superficial migratory stream within the cortical MZ is regulated
by DNMT1, and (ii) invading cortical interneurons influence apical cortical progenitors and
through this proper cortical layering.
Dnmt1 deletion in SST-expressing interneurons caused a premature exit from the MZ at E14.5,
accompanied by increased numbers of EOMES⁺ IPCs at E14.5 and more TBR1-positive
postmitotic neurons at E14.5 and E16.5, likely resulting from the altered division mode of
RGCs. This led to an extended cortical plate at E18.5 and enlarged deep layers at adult stages,
while upper layers were reduced in radial extension. These changes in cortical architecture
and resulting altered densities of SST⁺ interneurons in adult mice were associated with clear
functional and behavioral deficits, demonstrating the importance of DNMT1 for establishing
accurate network function in the adult cortex.
468
Martinotti cells constitute the majority of SST⁺ interneurons and localize frequently in superficial
layers II/III and deep layers V and VI[59]. They migrate primarily along the MZ during
corticogenesis[30]. In contrast, non-Martinotti SST⁺ cells disperse rather via the SVZ/IZ and
express a distinct set of genes[30]. From the reported genes being distinctively expressed
between SST⁺ cells of the SVZ versus MZ migratory stream[30], we identified several genes to
be regulated by DNMT1. For example, genes like *Sox1*, *Robo1* and *Npy*[30], which are
expressed higher in SVZ-migrating SST⁺ cells compared to ones migrating along the MZ, were
repressed by DNMT1 in E14.5 SST⁺ cINs (Supplementary Table S1). Moreover, *Erbb4* was
reported as a marker gene for Martinotti SST⁺ cells[45], being repressed by DNMT1 in SST⁺
cells. In combination with the selective effects of *Dnmt1* deletion on MZ migrating SST⁺
interneurons, these findings propose DNMT1 to be relevant for proper migration of Martinotti
cells.
Apart from genes being distinctively expressed between MZ versus SVZ migrating SST⁺
interneurons, our global transcriptome and methylome analyses propose that DNMT1 controls
other essential key transcription factors of MGE-cortical interneuron development such as
LHX6, ARX, ZEB2, SATB1, SOX1, SOX6, and DLX5[12,31,34]. ARX has been reported to directly
and indirectly modulate other genes controlling different stages of cortical interneuron
development, including *Maf*, *Mafb*, *Mef2c*, *Nkx2*.1, *Lmo1*, *Cxcr4*, *Nrg1*, *Erbb4*, *Sema3*, *Npr1*,
and *EphA4*[23], many of which were altered in expression upon *Dnmt1* deletion in SST⁺
interneurons. The elevated *Mef2c* levels in *Sst-Cre/tdTomato/Dnmt1 loxP²* cells are in line with
the increased proportion of PV-positive SST⁺-tdTomato cells in adult *Dnmt1* KO mice, as
Mef2c was reported to drive the PV fate[60,61]. In addition to ARX, MAF and MAFB promote
Mef2c expression[62], which also increased in expression upon *Dnmt1* deletion.
In line with the observed transcriptional changes, *Lhx6, Maf, Mafb,* and *Arx* mutant mouse
brains display a defect in cIN migration within the marginal zone (MZ)[23,29,30]. Moreover, ERBB4

regulates the switch from tangential to radial migration and the distribution of interneurons
within the CP. Tangential migration of cINs within the cortex is promoted by the CXCR4
receptor that binds to CXCL12, the latter being enriched in the MZ and SVZ migratory
streams[24]. The ERBB4 receptor recognizes NRG3, which is expressed in the CP, promoting
the radial navigation into the CP[25]. Both the loss of CXCR4 receptor in migrating interneurons,
and *Nrg3* overexpression in the CP result in premature invasion of migrating interneurons into
the CP[25]. As we did not detect an altered expression of *Cxcr4* in *Dnmt1*-deficient SST$^+$
interneurons, the increased *Erbb4* expression likely contributes to the premature invasion of
the CP in *Sst-Cre/tdTomato/Dnmt1-loxP²* brains, by shifting the balance from CXCL12/CXCR4
towards NRG3/ERBB4 signaling. Our data thus strongly suggests that DNMT1-mediated
repression of *Erbb4* promotes the migration of SST$^+$ cINs along the superficial migratory
stream. As Martinotti cells were described to migrate primarily along the MZ, while non-
Martinotti cells disperve via the SVZ[30], DNMT1 potentially contributes to the proper integration
of SST$^+$ cIN subtypes within the cortical migratory routes.

The migration of Martinotti cells along the MZ is functionally associated with the proper
formation of their axon collaterals in layer I[30]. These axon collaterals extend horizontally for up
to 2000 *µ*m, making up approximately three-quarters (76%) of the GABAergic axons in this
layer[59]. This enables Martinotti cells to influence the activity of numerous pyramidal neurons,
which are their primary postsynaptic targets[59]. Hence, DNMT1 might also indirectly influence
the proper formation of these powerful axon collaterals in layer I, by promoting the migration
of SST$^+$ interneurons along the MZ. Lack of axon collaterals could contribute to the strong
functional and behavioral impairments that we observed for the adult *Sst-Cre/Dnmt1* KO mice,
despite the overall number not having been affected. This is in line with data from *Mafb* KO
mice, which also display a premature exit from the MZ and no changes in interneuron numbers
and distribution in adults. Yet, functional impairments were likewise observed[30].
Martinotti cells have densely branched axons within their originating layer and inhibit not only
pyramidal neurons but also other IN subtypes such as PV basket cells and vasoactive intestinal
peptide (VIP) bipolar cells. Together with the non-Martinotti cells, that are present across all
cortical layers, SST$^+$ cINs play crucial roles in modulating cortical circuitry, excitability, and
network synchronization[59]. It is therefore most likely that the altered SST$^+$ interneuron density,
evident in our mouse model due to shifts in cortical layer thicknesses, contributes to the
observed functional and behavioral deficits. However, *Dnmt1*-deletion may have also triggered
functional impairments in adult SST$^+$ interneurons. In PV interneurons, DNMT1 regulates
GABAergic transmission via DNA methylation-dependent transcriptional control of
endocytosis-related genes, which influences transmitter recycling[41]. *Dnmt1*-deletion in SST$^+$
interneurons could thus similarly result in disrupted synaptic signaling in these cells.

The *Dnmt1* deletion-induced changes in SST$^+$ interneuron migration were accompanied by
increased production of deep-layer neurons and altered IPC numbers at embryonic stages,
manifesting in larger deep and reduced upper layers in the adult cortex. This strongly implies,
that invading cINs influence cortical progenitors and the timed generation of cortical neurons.
As evidenced by Sessa et al. (2010)[46], the development of cortical interneurons and excitatory
neurons must have been effectively integrated throughout evolution, since an increase in the
production of excitatory neurons also necessitates a rise in interneuron generation to maintain

the proper balance of excitation and inhibition. Likewise, invading cINs could influence the
cortical progenitor cells. Indeed, it was already shown that cINs modulate EOMES[+] IPCs´
proliferation, and through this, act on the timed generation of cortical excitatory neurons[47]. This
aligns in part with our findings presented here, whereas the pair cell assay rather points to an
influence of cortical interneurons on RGCs through ephrinB2 - EPHA4 signaling, influencing
the generation of IPCs and neurons from these cells. External regulation of EPHA4-expressing
RGCs was already described for invading thalamic afferents, that import ephrinA5, another
ephrin ligand[5]. This signaling regulates the temporal balance of proliferative and neurogenic
divisions of apical progenitors. Both *efnA5* and *EphA4* deficiency cause an increased
production of deep-layer neurons due to a developmental change in the RGCs' neurogenic
potential[5].
Similarly, invading cINs are likely to import "external information" in the form of membrane-
bound or secreted signaling molecules, as we propose here. *Dnmt1* deletion in SST[+]
interneurons caused increased *efnb2* expression, which is a ligand known to modulate cortical
neurogenesis. While progenitor-restricted *efnb2* deletion results in a delay in neurogenesis,
boosted ephrinB2 triggered EPHB signaling leads to a transient increase in neurons[48]. Similar
to our observations, the changes in neurogenesis measured by TBR1-positive cells were
restricted to a specific temporal window in the *efnB2* knockout mice, which aligns with our
results. This increase in neuronal production from RGCs was resembled in vitro with the pair
cell assay, stimulating RGC clones with ephrinB2-Fc. That changes in neuronal production at
specific embryonic stages can lead to altered layer thicknesses was already shown by other
studies[5,75].
In sum, the altered expression of secreted and membrane-bound signaling molecules with
known potential to regulate cortical progenitor division and fate, in concert with the altered
migration pattern of *Dnmt1* deficient SST[+] interneurons could lead to the altered cortical layer
formation. This suggests a crucial role for cortical interneurons in modulating the temporal
production of distinct subtypes of excitatory cortical neurons, and a high level of crosstalk
between immature cortical interneuron precursors and cortical progenitors to optimize the
formation of neural circuits. These findings also establish a key role for DNMT1 in orchestrating
cortical development through cell autonomous and non-cell autonomous actions.
569
570

Acknowledgements
This work was supported by the Flow Cytometry Facility and the Sequencing Facility, two core
facilities of the Interdisciplinary Center for Clinical Research (IZKF) Aachen within the Faculty
of Medicine at RWTH Aachen University. We thank Dr. Mira Jakovcevski and Ananya
Sangeetha for her experimental support, and Sandra Brill for her support with mouse care and
genotyping.
577
578

Author contributions
J.R. performed experiments and data analysis, figure illustration, manuscript editing; P.W.
performed experiments and data analysis, figure illustration; J.D. performed experiments and
data analysis; C.L.F. performed experiments and data analysis; L.D. performed experiments
and data analysis; C.B.Y. performed experiments and data analysis; J.L. performed

experiments and data analysis; J.V. performed experiments and data analysis; S.K. performed
experiments and data analysis; S.G. performed experiments and data analysis; G.N. methods
support and data analysis; M.S. performed experiments and data analysis, C.C.K. data
analysis; K.N.M. Data analysis, data discussion; T.V. data discussion and analysis; S.M. data
analysis and discussion; G.Z.B. conceptual design, wrote the manuscript, data discussion.
589

Funding
This research was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation)—368482240/GRK2416, ZI-1224/13-1.

Data Availability
The sequencing datasets generated and/or analyzed for this study are available in the Gene
Expression Omnibus repository with the following accession numbers: GSE276510,
GSE276512, and GSE276516. High-quality versions of all figures including those from the
extended data are available under https://doi.org/10.6084/m9.figshare.26963338.
599

Figure Legends

Figure 1 – DNMT1 regulates hub genes of cIN development in E14.5 SST$^+$ interneurons
Differential gene expression and methylation analysis of *Sst-Cre/tdTomato* control (ctrl) and
Sst-Cre/tdTomato/Dnmt1 loxP^2 (*knockout;* KO) mice based on total RNA and enzymatic
methyl-sequencing of FAC-sorted *tdTomato$^+$* cells isolated from the E14.5 basal
telencephalon. For RNA- and methyl-sequencing, we separately processed two samples per
genotype, each consisting of cells pooled from multiple embryos (RNA sequencing: Ctrl $n = 23$
embryos; KO $n = 14$ embryos; methyl-sequencing: Ctrl $n = 15$ embryos; KO n = 11 embryos).
(a) Heatmap depicting the 50 genes with the lowest adjusted *p*-values, color-coded based on
relative expression. **(b)** Volcano plot depicting the differentially expressed genes (DEGs)
between control and KO samples. Genes annotated as significantly changed (adjusted *p*-value
< 0.05) are colored based on the fold change of the expression in the KO group. (**c**) Volcano
plot of differentially methylated sites (DMS) identified in KO cells compared to control samples
($p < 0.001$ for color-labeled genes). **(d)** Venn diagram illustrating the overlap between up- or
downregulated genes and genes associated with or containing a differentially methylated
region (DMR). The intersection of upregulated and differentially methylated genes after
knockout was tested for enrichment of the DNMT1 binding motif identified by DNMT1-ChIP-
sequencing, shown in g. **(e)** DEGs identified in *Dnmt1* KO samples were compared to DEGs
determined in embryonic stem cell-derived murine neurons overexpressing DNMT1 **(f)**
Selection of brain development-related gene ontology (GO) terms enriched in genes that were
both upregulated and differentially methylated in E14.5 *Sst-Cre/tdTomato/Dnmt1* KO cells. **(g)**
DNA motif enriched in DNMT1-interacting chromatin, detected by ChIP-sequencing ($N = 2$
biological replicates). (**h**) Scatter plot combining DNA methylation changes and differential
gene expression highlighting *Arx* as a potential target of DNA methylation-dependent
transcriptional control by DNMT1. Bismark Methylation calls were filtered for a minimum of 3
reads per CpG and averaged for the promoter of each gene that was associated with a DMR
and was either up or downregulated. Promoter regions were defined as 1kb upstream of the

TSS plus 100 bp into the gene body. **(i)** Position of differentially methylated regions relative to
the transcription start sites of genes related to cIN development. Horizontal bars indicate the
region's size, and color-coded is the mean methylation change for the respective region. **(j)**
RNA sequencing tracks combined with the methylation profile of the *Arx* gene locus obtained
from ctrl and KO samples*.*

634
635

Figure 2: DNMT1 regulates the migration of SST+ interneurons in the developing cortex
at E14.5. (a) Schematic illustration of a coronally sectioned E14.5 brain hemisphere, depicting
the migration routes of cortical interneurons. **(b, c)** Exemplary microphotographs of tdTomato-
positive interneurons in coronally sectioned (50 μm) hemispheres of *Som-Cre/tdTomato*
(control) **(b)** and of *Som-Cre/tdTomato/Dnmt1 loxP²* (KO) embryos **(c)**. Scale bars: 500 μm.
(d) Quantitative analysis of the proportional distribution of tdTomato-positive cells in the basal
telencephalon and the cerebral cortex in E14.5 coronal sections (two-tailed nested *t*-test*, n =*
12 sections for control and *n* = 8 sections for KO, *N* = 3 embryos per genotype). **(e)**
Quantification of tdTomato-positive interneuron number within the cortex of control and KO
embryos normalized to the area, analyzed in 50 μm coronal cryosections (two-tailed nested *t*-
test *n* = 14 sections for control and *n* = 8 sections for KO, *N* = 3 embryos per genotype). **(f, g)**
Magnified microphotographs of the cortices of control and KO embryos (E14.5) to illustrate the
distribution of tdTomato+ cells within the cortical zones (MZ: marginal zone, CP: cortical plate,
IZ-VZ: intermediate zone to ventricular zone). White asterisks point to cells migrating along the
MZ (superficial migratory stream) or deep stream. Yellow arrows indicate tdTomato+
interneurons located in the CP. Scale bars: 100 μm. **(h)** Quantitative analysis of the
proportional distribution of tdTomato+ cells within the cortical zones normalized to the overall
tdTomato+-cell count within the cortex of control and KO embryos, analyzed in 50 μm coronal
cryosections (two-tailed nested *t*-test, *n* = 13 sections for control and *n* = 14 sections for KO,
N = 3 embryos per genotype). **(i, j)** Temporal z-projections of migration tracks of tdTomato-
positive cells in organotypic brain slice cultures (350 μm thickness, coronal) from E14.5 *Som-*
Cre/tdTomato (control) and *Som-Cre/tdTomato/Dnmt1 loxP²* (KO) embryos, captured over
20 h by live-cell imaging. Scale bars: 500 μm. **k-n)** Quantification of the live-cell recordings in
organotypic brain slices. Quantitative analysis of the increase in fluorescence intensity in the
cortical plate due to invasion of tdTomato cells (first 12 h, two-tailed nested *t*-test, *n* = 4 slices
for control and *n* = 6 slices for KO embryos, *N* = 3 embryos per genotype) is shown in **(k)**, the
average number of directional changes (*n* = 134 cells across 5 slices for control (*N* = 4
embryos), *n* = 74 cells across 3 slices for KO (*N* = 3 embryos), two-tailed nested *t*-test) is
depicted in **(l)**. The velocity of all cells migrating within the cortex (*n* = 377 cells across 6 slices
for control (*N* = 5 embryos), *n* = 323 cells across 5 slices for KO (*N* = 3 embryos)) is shown in
(m), while the speed of cells within the area between CP and VZ is illustrated in **(n)** for control
(*n* = 193 cells across 6 slices, *N* = 5 embryos) and KO (*n* = 114 cells across 3 slices, *N* = 3
embryos); two-tailed nested *t*-test. **(o, p)** Selected frames extracted from live-cell imaging
recordings of organotypic brain slices (350 μm) derived from E14.5 control and KO embryos.
Depicted are tdTomato-positive cells migrating through the cortex. For each genotype, three
selected cells (dotted circles) were tracked. Scale bars: 50 μm.

Error bars represent the standard error of the mean (SEM). $p < 0.05$*, $p < 0.01$**, $p < 0.001$***;
n.s.: not significant. CP: cortical plate, VZ: ventricular zone, SVZ: subventricular zone, MZ:
marginal zone, IZ: intermediate zone.
675
Figure 3: The *Dnmt1* deletion-induced effect on SST+ interneuron migration is
temporarily restricted. (a) Exemplary microphotographs of tdTomato+ interneurons in
coronally sectioned (50 µm) hemispheres of E16.5 *Som-Cre/tdTomato* (control) and of *Som-*
Cre/tdTomato/Dnmt1 loxP² (KO) embryos; scale bars: 500 µm. TdTomato-positive cells are
labeled in red and DAPI in blue. Quantitative analysis of the proportional distribution of
tdTomato cells in the basal telencephalon and the cerebral cortex in E16.5 coronal sections
(two-tailed nested *t*-test, $n = 11$ sections for control and $n = 15$ sections for KO, $N = 4$ embryos
per genotype; error bars represent the standard error of the mean (SEM). **(c, d)** Magnified
microphotographs of the cortices of control and KO embryos from E16.5 coronal brain sections
(50 µm) to illustrate the distribution of tdTomato cells within the cortical zones. Scale bars: 100
µm. **(e)** Quantitative analysis of the proportional distribution of tdTomato cells within the E16.5
cortical zones normalized to the overall tdTomato-cell count within the cortex of control and
KO embryos, analyzed in 50 µm coronal cryosections. Two-tailed nested *t*-test, $n = 13$ sections
for control and $n = 14$ sections for KO, $N = 4$ embryos per genotype. **(f)** Quantitative analysis
of tdTomato cell density normalized to the given area of the respective cortical zones (E16.5,
two-tailed nested *t*-test, $n = 13$ sections for control and $n = 14$ sections for KO, $N = 4$ embryos
per genotype). **(g, h)** Exemplary microphotographs of tdTomato-positive interneurons in the
cortex of coronally sectioned (50 µm) brains of E18.5 *Som-Cre/tdTomato* (control) and *Som-*
Cre/tdTomato/Dnmt1 loxP² (KO) embryos; scale bars: 100 µm. **(i)** Quantitative analysis of the
proportional distribution of tdTomato cells within the E18.5 cortical zones normalized to the
overall tdTomato-cell count within the cortex of control and KO embryos, analyzed in 50 µm
coronal cryosections. Two-tailed nested *t*-test, $n = 11$ sections for control and $n = 9$ sections
for KO, $N = 4$ embryos per genotype, error bars represent the SEM. **(j)** Quantitative analysis
of tdTomato cell density normalized to the given area of the respective cortical zones (E18.5,
two-tailed nested *t*-test, $n = 11$ sections for control and $n = 9$ sections for KO, $N = 4$ embryos
per genotype). MZ: marginal zone, CP: cortical plate, IZ-VZ: intermediate zone to ventricular
zone. $p < 0.05$ *, $p < 0.01$ **, $p < 0.001$ ***, n.s.: not significant.
703
704
Figure 4: Conditional deletion of *Dnmt1* in SST interneurons affects IPCs and the
generation of TBR1+ cells non-cell autonomously. (a) Schematic illustration of a coronally
sectioned embryonic brain hemisphere, depicting the lateral and dorsal localization of the two
bin selections, used to quantify the density and distribution of EOMES+- and TBR1+-cells in
E14.5 and E16.5 brain sections. **(b-e)** TBR1 and EOMES immunostaining in coronally
sectioned (50 µm) brains of E14.5 *Sst-Cre/tdTomato* (control) and *Sst-Cre/tdTomato/Dnmt1*
loxP² (KO) embryos. **(b** and **d)** Exemplary microphotographs of **(b)** TBR1 immunostaining
(green) and **(d)** EOMES immunostaining (green); scale bars: 50 µm (DAPI is shown in blue).
(c and **e)** Quantitative analysis of the TBR1+ postmitotic neuron numbers **(c)** and EOMES+
intermediate progenitor cell (IPC) numbers **(e)** per bin in the E14.5 cortex of control and KO
embryos, analyzed in 50 µm coronal cryosections (two-way ANOVA with subsequent

Bonferroni correction*, n* = 13 control and *n* = 9 KO sections from *N* = 3 embryos for both genotypes). **(f)** TBR1 immunostaining (green) in coronally sectioned (50 *μ*m) brains of E16.5 control and KO embryos, scale bars: 50 *μ*m. **(g)** Quantitative analysis of the TBR1-positive postmitotic neuron numbers per bin in the E16.5 cortex of control and KO embryos, analyzed in 50 *μ*m coronal cryosections (two-way ANOVA with subsequent Bonferroni correction*, n* = 8 for control and KO from *N* = 3 embryos for both genotypes). **(h** and **j)** Exemplary microphotographs of (**h**) EOMES immunostaining (green) and **(j)** TBR1 immunostaining (green) in coronal sections of E18.5 control and KO brains; scale bars: 50 *μ*m (DAPI is shown in blue). **(i** and **k)** Quantitative analysis of the density of EOMES-positive IPCs **(i)** and TBR1-positive postmitotic neurons **(k)** in the E18.5 cortex of control and KO embryos, analyzed in 50 *μ*m coronal cryosections (two-tailed nested *t*-test*, n* = 11 sections for both genotypes from *N* = 4 embryos per genotype). **(l)** Schematic illustration of the cortex of a coronally sectioned embryonic brain (E18.5), depicting the parameters analyzed for the quantification shown in (**m**). Three regions (lateral (1), dorsolateral (2), and dorsal (3)) were averaged to obtain the mean thickness respectively for each layer (two tailed nested *t*-test*; n* = 12 control and *n* = 9 KO sections*. N* = 4 brains for both genotypes). **(n** and **o)** Panels depict significant ligand-receptor pairs involved in communication from SST⁺ cINs to cortical progenitors, filtered by differentially expressed genes (DEG) from single-cell RNA-sequencing data. Communication probability is represented by dot color, and *p*-value by dot size*,* with *p*-values being computed using a one-sided permutation test. Panel (**n**) shows the communication with different apical progenitor (AP) populations at E14.5. Panel **(o)** illustrates communication with IPCs at E14.5. Panels **(p and r)** present microphotographs of cell pair combinations formed from cortical neurons isolated from E14.5 C57BL6/J embryos, plated at clonal density, after 24 hours. Cell clones were either treated with control (ctrl.)- or recombinant efnB2-FC for 24 h, before immunocytochemical staining for Nestin (magenta), β-III-tubulin (TUBB3, green), EOMES (TBR2, yellow) and DAPI (blue). Scale bars: 5 *μ*m. **(q and s)** Quantification of the proportion of Nestin/Nestin (2xN), β-III-tubulin/β-III-tubulin (2xT), Nestin/β-III-tubulin (N-T), EOMES-EOMES (2xE), and Nestin-EOMES (N-E) positive cell pairs normalized to the overall number of these pairs. Unpaired, two-tailed Student's *t*-test with subsequent Welsh's correction (panel q: control-Fc: 66 cell pairs, efnB2-Fc: 58 cell pairs, from *N* = 3 experiments; panel s: control-Fc: 71 cell pairs, efnB2-Fc: 73 cell pairs, from *N* = 4 experiments). $p < 0.05$ *, $p < 0.01$ **, $p < 0.001$ ***. MZ: marginal zone, CP: cortical plate, IZ-VZ: intermediate zone-ventricular zone, ctrl.: control, n.s.: not significant, ctrl: control.

Figure 5: Adult *Dnmt1*-deficient mice show aberrant cortical architecture and functionality. (a-d) Immunostaining using an antibody directed against parvalbumin (PV) in 50 *μ*m brain sections of six-month-old *Sst-Cre/tdTomato* (control) and *Sst-Cre/tdTomato/Dnmt1 loxP²* (KO) mice show increased proportions of PV⁺ *Sst-Cre/tdTomato* neurons upon *Dnmt1* deletion. Exemplary microphotographs of immunostainings (tdTomato in red, PV in green, DAPI in blue) taken from the primary somatosensory cortex (S1) of coronal slices (Bregma 0.14, -0.22, and -0.34) from (**a**) control and (**b**) KO mice. Scale bars: 100 *μ*m (left panels) and 20 *μ*m (magnifications of marked selections). **(c)** Quantification of the proportion of PV-positive *Sst-Cre/tdTomato* cells over the whole S1 as well as in supra- and infragranular layers normalized to the total amount of all detected *Sst-Cre/tdTomato* cells. Two-

tailed nested *t*-test with *n* = 11 slices for control and *n* = 10 for KO from *N* = 3 brains for both

genotypes. **(d-g)** *Dnmt1* KO mice display changes in cortical layer thickness alongside

alterations in *Sst-Cre/tdTomato* cell density. **(d, e)** Exemplary microphotographs of

immunostainings of CUX1 and tdTomato combined with DAPI taken from S1 (sagittal slices of

Bregma 1.32 and 1.44, six-month-old male mice). The CUX1-positive upper layers II-IV can

be distinguished from deep layers V and VI. Scale bars: 100 μm. **(f, g)** Quantification of the

tdTomato-positive cell density normalized to the given area of the respective layers (**f**), the

cortical thickness, and the proportional thickness of the deep and upper layers (**g**). Two-tailed

nested *t*-test with *n* = 24 slices for control and *n* = 23 for KO from *N* = 3 brains for both

genotypes. **(h-q)** Neuropixels recordings in S1 of adult *Sst-Cre/tdTomato* (control) and *Sst-*

Cre/tdTomato/Dnmt1 loxP^2 (KO) mice (n = 8 sessions from 2 mice per genotype), while animals

were receiving either tactile stimulation by air puffs against their whiskers or visual stimuli **(h)**.

(i) Average local field potential (LFP) in S1 across the whole cortical depth, following a 20 ms

air puff stimulus (onset at white line) to the distal whisker pad. Blue colors indicate negative,

and yellow colors positive LFP responses. Across all layers, LFP responses were stronger and

more temporally precise in control mice (left) compared to KO mice (right). **(j)** Average current

source density (CSD) in S1 calculated from the measured LFP. Blue colors indicate excitatory

and yellow colors inhibitory currents. The earliest responses in control mice (left) occur in the

granular layer at ~400 μm cortical depth and then spread to layer II/III (100-300 μm) and layer

V (500-900 μm). In contrast, no clear spatial response profile is visible in KO mice, suggesting

a disruption of cortical network function. **(k)** Traces of LFP responses in S1 depicting averaged

LFPs across all cortical layers (0 - 1 mm) for recordings in control (gray) and KO (orange) mice

(shading indicates the standard error of the mean (SEM)) illustrate significantly larger and

faster tactile responses in control mice compared to KO mice (peak amplitude$_{control}$ = -159.09

± 34.37 μV, peak amplitude$_{KO}$ = -57.17 ± 7.49 μV; Wilcoxon *rank-sum* test, *p* = 0.0059; peak

time$_{control}$ = 26.85 ± 26.85 ms, peak time$_{KO}$ = 30.80 ± 1.10 ms, *p* = 0.0108). **(l-o)** Quantification

of the temporal precision of the spiking response of S1 neurons to tactile stimulation in control

(gray) and KO (orange) mice. **(l)** Peristimulus time histogram (PSTH) for spiking response of

S1 neurons to tactile stimulation in KO (orange) and control mice (gray). **(m)** The width of the

initial response, measured as the full width at half maximum (FWHM), was significantly larger

in KO compared to control mice (n = 298 clusters, Wilcoxon *rank-sum* test, *p* = 7x10^{-31}). **(n)**

Quantification of early tactile response amplitude in control versus KO mice. In the first 30 ms

after tactile stimulation, responses in control mice were significantly larger compared to KO

mice (mean firing rate$_{control}$ = 9.09 ± 0.31 Hz, *n* = 436 clusters, mean firing rate$_{KO}$ = 7.22 ± 0.32

Hz, *n* = 298 clusters, Wilcoxon rank-sum test, p = 0.000045). **(o)** Late tactile responses in KO

mice were significantly larger compared to control mice (mean firing rate$_{control}$ = 0.4 ± 0.10 Hz,

n = 436 clusters, mean firing rate$_{KO}$ = 3.85 ± 0.43 Hz, n̲ = 298 clusters, Wilcoxon *ranksum*

test, *p* = 4x10^{-37}). **(p)** Absolute spectral power density (ASPD) of LFP recordings across all

layers in primary visual cortex (V1) during full-field visual stimulation in control (left) and KO

mice (right). Hot colors indicate higher spectral power, units are on a log$_{-10}$ scale. Spectral

power falls off with frequency but shows increased activity in the higher frequency range >

20Hz between 0.3- and 0.7-mm cortical depth for control mice. In contrast, no clear modulation

of higher frequencies was found in KO mice. **(q)** Relative change in spectral power during

visual stimulation, computed as the percent difference relative to baseline. In control mice (left),

visual stimulation induced a strong increase in the gamma range (> 30Hz), especially at around

70 Hz between 0.3- and 0.7-mm cortical depth. In contrast, no clear modulation was found in
KO mice (right). $p < 0.05$ *, $p < 0.01$ **, $p < 0.001$ ***, n.s.: not significant.
808
809

Figure 6: Adult *Dnmt1*-deficient mice show behavioral abnormalities. (a-e) Nest building
test reveals a significant deterioration in using given material and forming proper nests in adult
Sst-Cre/tdTomato/Dnmt1 loxP[2] (KO) individuals compared to *Sst-Cre/tdTomato* (control) mice.
With the help of the scoring index depicted in **(a)** and formerly described[73], all mice were ranked
based on nest quality and amount of material used **(d, e)**. Representative photographs are
shown in **(b)** for control and **(c)** for KO mice. Tests were performed overnight and twice within
one week. Respective values were averaged for both trials. Unpaired, two-tailed Student's *t*-
test with subsequent Welch's correction. $N = 16$ male adult mice for both genotypes. **(f-h)**
Pentylenetetrazol (PTZ)-injections point to defects in cortical inhibition in KO mice. **(f)**
Schematic illustration defining PTZ-induced seizures in three-month-old males, that received
intraperitoneal injections of PTZ every ten minutes until displaying a final tonic-clonic seizure
(event IV) within a time window of a maximum of 120 minutes. Footages captured from video
monitoring of PTZ-induced convulsions defining criteria for small seizures (events I and II) and
tonic-clonic seizures (III and IV). **(g, h)** Quantification of seizure profiles of control ($N = 9$) and
KO individuals ($N = 6$). Unpaired, two-tailed Student's *t*-test with subsequent Welch's
correction. $p < 0.05$ *, $p < 0.01$ **, $p < 0.001$ ***.
826
827

828
829 **Methods:**
830
Animals
In the recent study transgenic mouse strains with a genetic C57BL/6J background (initially
obtained from the University Hospital UKA Aachen, Germany) were used. *Sst^tm2.l(cre)Zjh^*/J x
*B6.CgGt(ROSA)26Sor^tml 4(CAG-tdTomato)Hze^* (*Sst^+/-^-Cre/tdTomato*) served as control animals
whereas *Sst^tm2.l(cre)Zjh^/J* x B6.CgGt(ROSA)26Sor^tm14(CAG-tdTomato)Hze^ x *B6; 129S-Dnmt1tm4Jae*/J
(*Sst-Cre^+/-^/tdTomato/Dnmt1 loxP^2^)* were used as *Dnmt1* knockout (KO) model.
837

Generation of control animals was conducted by crossing a *Sst^tm2.l(cre)Zjh^*/J strain
(RRID:IMSR_JAX:013044, Jackson Laboratory, Bar Harbour, U.S.A.) with a t*dTomato* reporter
mouse model (*B6.CgGt(ROSA)26Sor^tml 4(CAG-tdTomato)Hze^,* obtained from Christian Hübner,
University Hospital Jena, Germany). An *internal ribosomal entry site* (*IRES*), a *Cre-
recombinase* sequence, a *polyA* sequence, and a *frt*-flanked neo cassette were inserted into
the 3' untranslated region (UTR) of the *Somatostatin (Sst)* locus on chromosome 16, limiting
the respective *Cre*-expression to *Sst*-positive cells. The t*dTomato* reporter strain was
characterized by the corresponding *tdTomato* sequence combined with a *loxP*-flanked stop
cassette, both located in the *Rosa26* locus. Expression of *Cre-recombinase* dependent on *Sst*-
promoter activity resulted in the deletion of the *loxP*-flanked stop cassette during recombination
and a final expression of *tdTomato* in *Sst-Cre*-positive cells. *Jackson Laboratory* recommends
creating and breeding *Sst-Cre* heterozygous mice[77], which received the *Cre*-allele from the

maternal side. This strategy avoids potential instability of the Cre-recombinase activity during
paternal germline recombination[77].
To generate the triple transgenic conditional KO-model *Sst-Cre+/-/tdTomato* females were bred
with male individuals derived from crossing *tdTomato* mice with a *Dnmt1 loxP²* strain (*B6;*
129Sv-Dnmt1tm4Jae/J, obtained from Rudolf Jaenisch, Whitehead Institute for Biomedical
Research Boston, U.S.A.) in which exon 4 and 5 of the *Dnmt1* gene are loxP-flanked, resulting
in a null allele of these loci and a subsequent DNMT1 deficiency[78]. Corresponding programs
and primers used for genotyping can be found in the supplementary information. An illustration
of control- and KO mice is shown in Extended Data Figure E1a.
859
860

Isolation of Embryonic and Adult Brains
Individuals of embryonic stage 14.5, 16.5 or 18.5 were isolated as described in Symmank et
al. (2018)[79]. Briefly, pregnant females were anesthetized by intraperitoneal administration of
ketamine/xylazine (200/25 mg/kg living weight per injection up to a maximum application of
600/75 mg ketamine/xylazine per kg living weight in total). Upon reaching surgical tolerance,
the abdominal cavity was opened to expose the uterine horns that were removed. Embryos
were isolated from the uteri and decapitated. For histological analyses of E14.5 and E16.5
brains, the heads were directly transferred to 4% paraformaldehyde (PFA)/1x phosphate-
buffered saline (PBS). For histological analyses at E18.5, brains were removed from the skull
before transferring them to 4% PFA/1x PBS. E14.5 heads were fixed for 5 hours, while E16.5
heads and E18.5 brains were fixed overnight at 4 °C on an orbital shaker (50 rpm). Afterwards,
stepwise cryopreservation was performed by overnight incubation first in 10% Sucrose/1x
PBS, followed by 30% sucrose/1x PBS at 4 °C on an orbital shaker (50 rpm). Tissue was
frozen in liquid nitrogen and stored at -80 °C.
875
For the isolation of adult brains, male mice were sacrificed with an overdose of 5% (v/v)
isoflurane. For histological analyses and identification of Neuropixels recording sites,
transcardial perfusion was conducted with 1x PBS (pH 7.4) followed by 4% PFA/1x PBS
(pH 7.4) with a pump[80]. After brain preparation, post-fixation was conducted in 4% PFA/1x
PBS for 24 h at 4°C on a roller mixer with constant rotation (approx. 50 rpm). Then,
cryopreservation was performed in 10% sucrose/1x PBS and 30% sucrose/1x PBS for 24 h
each at 4°C on a roller mixer with constant rotation (approx. 50 rpm) before brains were frozen
in liquid nitrogen and stored at -80°C.
884
Organotypic brain slices and single-cell preparations
MGE- and cortical single cells were prepared as described previously[19,41]. Briefly, embryonic
heads were collected and dissected in Gey's Balanced Salt Solution (GBSS, pH 7.4)/0.65%
D-glucose on ice. MGEs and cortices were cut out and collected in Hanks' Balanced Salt
Solution (HBSS, w/ phenol red, w/o calcium, w/o magnesium)/0.65% D-glucose on ice. The
tissue was treated with 0.04% of trypsin/EDTA (Thermo Fisher Scientific, U.S.A.) for 17 min at
37°C, before MGE cells were dissociated in cold Dulbecco's Modified Eagle Medium (DMEM)
with additional L-glutamine and 4.5 g/L D-glucose (Thermo Fisher Scientific, U.S.A.), 10% fetal
bovine serum (FBS; Biowest, U.S.A.), and 1% penicillin/streptomycin (P/S) (Thermo Fisher

Scientific, U.S.A.) by trituration with glass Pasteur pipettes and by subsequent filtering through
a nylon gauze (pore size 140 μm; Merck, U.S.A.). Dissociated single cells were seeded on
laminin (19 μg/ml; Sigma-Aldrich, U.S.A.)/poly-L-lysine- (10 μg/ml; Sigma-Aldrich, U.S.A.)
coated glass coverslips. For MGE cells, we used a density of 455 cells/mm^2. Cortical single
cells were seeded in clonal densities of 150 cells/mm^2. Transfection of MGE cells with siRNA
oligos occurred after 5–6 h of incubation (37°C, 5% CO_2 and 95% relative humidity) in
Neurobasal medium with phenol red, 1x B27™, and 0.25x GlutaMAX (Gibco, U.S.A.) for 24 h.
For this, the medium was supplemented with Lipofectamine™ 3000 in accordance with the
manufacturer's instructions (Thermo Fisher, U.S.A.) together with *Dnmt1* siRNA (30 nM, #sc-
35203, Santa Cruz, U.S.A.) and a scrambled control siRNA construct (15 nM, BLOCK-iT™
Fluorescent Oligo, #2013, Thermo Fisher, U.S.A.). After 24 h MGE cells were fixated with 4%
PFA/1x PBS for 10 min.
Incubation of cortical cells plated at clonal density occurred at 37°C, 5% CO_2 and 95% relative
humidity in culture medium (Neurobasal medium with phenol red, 1x B27™, 0.25x GlutaMAX
(Gibco, U.S.A.), and 0.4% methylcellulose (Sigma, U.S.A.)) for 24 h. Treatment of cortical
single cells was conducted 5–6 h after plating. For this, the medium was supplemented with
10 μM 5-ethynyl-2′-deoxyuridine (5-EdU, baseclick™, baseclick GmbH, Germany) and with
either 5 μg/mL of a recombinant human Fc control protein (Rockland Immunochemicals,
U.S.A.) or 5 μg/mL of a recombinant human efnB2-Fc (R&D Systems, U.S.A.). Both
recombinant proteins were pre-clustered with 10 μg/mL of an anti-human IgG antibody
(Thermo Fisher, U.S.A.). Cortical single cells were fixated with 4% PFA/1x PBS for 10 min.
For organotypic brain slice preparations[27], brains of E14.5 embryos were embedded in 4% low
melt agarose (37°C) in Krebs buffer (126 mM NaCl, 2.5 mM KCl, 1.2 mM NaH_2PO_4, 1.2 mM
$MgCl_2$ * $6H_2O$, 2.1 mM $CaCl_2$; pH 7.4; supplemented with 10 mM D-glucose and 12.5 mM
$NaHCO_3$, sterile-filtered). For sectioning of 350 μM coronal slices, a 5100mz-Plus vibrating
microtome (speed: 0.7–0.8 mm/s and blade oscillation frequency: 5 Hz; Camden Instruments,
United Kingdom) was used. Slices were transferred to ice-cold post-holding buffer (1x Krebs
buffer, 10 mM HEPES, 1% P/S, 0.2% gentamycin, pH 7.4). Slices were imaged in μ-slide 4
Well imaging plates (ibidi GmbH, Germany) coated with 19 μg/mL laminin (Sigma-Aldrich,
U.S.A.) and 10 μg/mL poly-L-lysine (Sigma-Aldrich, U.S.A.) in Neurobasal without phenol red,
1x B27™, 1% P/S, 0.5% D-glucose, 10 mM HEPES, using a confocal-like Leica DMi8
fluorescent microscope in combination with a THUNDER® imager unit (Leica, Germany) and
the corresponding software *LASX* (Leica, Germany) equipped with an incubation chamber
(37°C, 5% CO_2 and 95% relative humidity). Imaging was conducted with a wavelength of
544 nm (TRITC-channel) in tile scans including z-stacks of 8–10 μm step size every 15 min.
Post-processing of the respective footages was executed in the *LASX* software using the
"*Mosaic Merge*"- and "*Thunder Lightning (Large Volume)*"-tools. Finally, a maximum intensity
projection of the merged tile scans was processed using the *Fiji* software[81]. Analysis was
performed blindly using the manual track plugin from ImageJ (NIH, USA) to track cells with a
migration time of at least 6 h.
934
Histology and immunocytochemistry
For histology on embryonic brains, we used 50-μm coronal brain slices, cryosectioned with a
CM3050 S Cryostate (Leica, Germany), and collected on Superfrost®Plus (Avantor, U.S.A.)
object slides. On-slide immunohistochemistry was performed as described in Pensold et al.

(2017)[27]. Briefly, all samples were washed for 5x 20 min in 1x PBS /0.5% Triton X-100/0.5%
Tween®20 with constant horizontal shaking (50 rpm). After that all samples were treated with
blocking solution (4% bovine serum albumin (BSA) and 10% normal goat serum in 1x
PBS/0.5% Triton X-100/0.5% Tween®20) was applied for 2 h at room temperature (RT).
Primary antibody incubation diluted in blocking solution was conducted overnight at RT in a
humid chamber. After washing for 5x 20 min in 1x PBS/0.5% Triton X-100/0.5% Tween®20,
secondary antibodies were applied in blocking solution for 2h at RT. After washing for 3 x 20
min, the samples were stained with 4',6-diamidino-2-phenylindol dihydrochloride (DAPI,
1:10000 in PBS, Carl Roth, Germany) for 15 min at RT, followed by washing twice for 5 min
with PBS, sections were embedded in Mowiol.
For adult brains, immunohistochemistry was performed in free-floating sagittal and coronal
sections (30 μm), prepared using the CM3050 S Cryostate (Leica, Germany). Slices were
transferred to 1x PBS and stored at 4°C. For sagittal sections, Bregma 1.32 and 1.44 were
used, while Bregma 1.18, 1.10, 0.14, -0.22, -3.08, and -3.28 were taken for coronal sections.
The same procedure as for embryonic sections was applied, except that an antigen retrieval
with heated HistoVT One® (1x HistoVT One®/H$_2$O bidest., Nacalai, Japan) at 70°C for 20 min
was performed for staining against DNMT1. Sections that were stained for somatostatin,
calretinin, and NPY underwent an antigen retrieval with citrate buffer (95°C; 10 mM, pH 6.0,
supplemented with 0.5% Tween®20) for 15 min and subsequent cool-down for 30 min before
washing. Finally, all slices were transferred onto glass slides in 0.5% (w/v) gelatine/0.05%
(w/v) chromium(III) potassium sulfate dodecahydrate (KCrS$_2$O$_8$)/H$_2$O solution prior to
embedding in Mowiol.
Immunostaining on dissociated MGE and cortical single cells was conducted by washing
coverslips with 1x PBS/0.1% Triton X-100 for 3x 5 min and blocking with 1x PBS/0.1%
Triton X-100/4% BSA for 30 min at room temperature. Primary antibodies were applied for 2 h
at room temperature. After washing for 3x 5 min with 1x PBS/0.1% Triton X-100, the secondary
antibody was incubated for 1 h at room temperature. Phalloidin-647 diluted 1:1000 in 1x PBS
(#ab176759, Abcam, U.S.A.) was applied for 20 min after washing. Then a final washing step
with 1x PBS for 10 min and a DAPI staining (1:10000/1x PBS; Carl Roth, Germany) for 5 min
were performed and coverslips were mounted in Mowiol.

The applied primary antibodies were used: mouse anti-Calretinin (1:500; Swant, Switzerland,
#6B3), rabbit anti-NPY (1:2500; Immunostar, U.S.A., #2940), rabbit anti-TBR1 (1:200; Abcam,
U.S.A., #ab31940); rabbit anti-EOMES (1:500; Abcam, U.S.A., #ab23345); rat anti-
Somatostatin (1:100; Millipore, U.S.A., #MAB354), mouse anti-Parvalbumin (1:2000; Swant,
Switzerland, #235); rabbit anti-DNMT1 (1:100; Santa Cruz, U.S.A., # sc20701); rabbit anti-
CUX1 (CDP; 1:100, Santa Cruz, U.S.A., #sc13024); mouse anti-Nestin (1:100; Merck U.S.A.,
#MAB353), rabbit anti-ß-Tubulin III (1:500; Sigma Aldrich, U.S.A., #T2200), and rabbit anti-
ErB4 (1:1000; Proteintech, U.S.A., #22387-1-AP).
The following secondary antibodies conjugated with respective fluorophores were used at
1:1000 dilutions: Alexa488-goat anti-Rat IgG (Invitrogen, U.S.A., #A11006) Cy5-Goat anti-
Rabbit IgG (Life Technologies, U.S.A., #A10523), A488-Donkey anti-Mouse IgG (Jackson,
U.S.A., #15454150), Cy5-Goat anti-Mouse IgG (Jackson, U.S.A., #115175146), A488-Goat
anti-Rabbit IgG (Life Technologies, U.S.A., #A11008).
982

Transcriptome data on neurons overexpressing DNMT1

Dnmt1tet/tet mouse embryonic stem cells have been previously described[82]. The levels of DNMT1 in these cells are four times higher than the normal wild-type *R1* ESCs. Upon differentiation, the induced *Dnmt1tet/tet* neurons produced contain twice the DNMT1 levels than in *R1* neurons[82]. Differential expression gene data on *Tet/Tet* neurons was taken from Singh *et al.* (2023)[83] and genes with significantly altered transcript levels were compared with the transcriptome data on the tissues lacking DNMT1.

Preparation of E14.5 tdTomato cells for RNA and methyl-sequencing

For FACS-mediated enrichment of E14.5 *Sst-Cre/tdTomato* cells, the whole basal telencephalon was prepared from the embryos and subjected to cell dissociation. Nuclease-free reaction tubes were used during isolation and long-term storage of the resulting material. The basal telencephalons were collected in cold HBSS (w/ phenol red, w/o calcium, w/o magnesium)/0.65% D-glucose; 4 μg/μL (600 U) of DNAse I (AppliChem GmbH, Germany). After treatment with 0.04% trypsin for 17 min at 37 °C, HBSS was replaced by DMEM with additional L-glutamine and 4.5 g/L D-glucose, 10% FBS and 1% P/S to stop the trypsinization. Subsequently, the cells were pelletized, resuspended and triturated in cold HBSS (w/o phenol red, w/o calcium, w/o magnesium)/0.65% D-glucose, before being filtered through a nylon gauze (pore size 140 μm, Merck, U.S.A.) for FACS.

FACS was performed by the Flow Cytometry Facility (FCF, University Hospital RWTH Aachen, Germany). Respective parameters for the procedure using a BD FACS Aria Fusion (BD Biosciences, U.S.A) were defined as follows: 5-laser (FCS, SSC, PE, BV421), 18-color (3-6-2-4-3). TdTomato-positive cells were either collected in 100 μL of cold TRIzol™ (Thermo Fisher Scientific, U.S.A.) for subsequent RNA sequencing or in 100 μL of cold HBSS (w/o phenol red, w/o calcium, w/o magnesium)/0.65% D-glucose for further processing for DNA methylation analysis. Finally, all samples were stored at -80°C.

Total RNA seq

TdTomato-positive E14.5 MGE cells from *Sst-Cre/tdTomato or Sst-Cre/tdTomato/ Dnmt1 loxP2* were FAC sorted into 100 μL TRIzol™ reagent (Thermo Fisher Scientific, U.S.A.). Samples were mixed by inversion and stored at -80°C until processing. RNA was isolated from 50,000 cells pooled per genotype. The samples were filled up to 1 ml with TRIzol™ and dounced 25 times in a glass homogenizer using the small-clearance pestle. Samples were incubated for 5 min at room temperature, supplemented with 200 μL chloroform, and mixed by shaking vigorously until a homogenous milky solution appeared. Following another incubation at room temperature (until a visible phase separation appeared) samples were centrifuged at 13,000x *g* and 4°C for 15 min in a tabletop centrifuge with cooling function. For precipitation, the clear upper phase was collected and mixed with 70% ethanol. Samples were bound onto RNeasy mini columns (Qiagen, Germany) and washed according to the instructions in the manual. RNA was eluted with 30 μL nuclease-free water.

RNA concentrations were determined using a Qubit™ 4 flourometer (Thermo Fisher Scientific, U.S.A.) with RNA HS reagents (Thermo Fisher Scientific, U.S.A.). RNA integrity was checked on a Tape Station 4200 using the HS RNA kit (Agilent Technologies, U.S.A.).

Prior to library preparation, samples were digested with HL dsDNAseI (ArcticZymes Technologies ASA, Norway) for 10 min at 37°C to remove traces of gDNA. To avoid digestion

of the newly generated cDNA during library preparation, the dsDNAseI was heat-inactivated
for 5 min at 58°C. Thereafter, samples were subjected to library preparation using the TAKARA
SMARTer Stranded Total RNA-Seq Kit v3 - Pico Input Mammalian for Illumina (Takara Bio,
Japan). Fragmentation was done at 94°C for 4 min. Illumina adaptors and indices were ligated
to cDNA in a PCR reaction with 5 cycles (98°C for 15 s, 55°C for 15 s and 68°C for 30 s).
RRNA depletion was performed on the amplified libraries. Final libraries were subjected to an
upscale PCR with 13 cycles (98°C for 15 s, 55°C for 15 s and 68°C for 30 s). Amplified libraries
were cleaned with magnetic AMPure XP beads (Beckman Coulter, U.S.A.). Library
concentration was determined with a Qubit™ 4 fluorometer using the 1x dsDNA HS reagent
(Thermo Fisher Scientific, U.S.A.). Library size was assessed on a Tape Station 4200 using
the DNA D1000 kit (Agilent Technologies, U.S.A.).
Libraries were diluted, equimolarly pooled, denatured, and loaded for clustering onto an
Illumina NovaSeq 6000 SP (200 cycles) flow cell (Illumina, U.S.A.) in conjunction with 1% PhiX
control library, and run on an Illumina NovaSeq machine (Illumina, U.S.A.) in 75 bp paired-end
mode. The 1% PhiX control library was spiked in to improve base calling accuracy.
1043
FASTQ files were generated using *bcl2fastq* (Illumina, U.S.A.). To facilitate reproducible
analysis, samples were processed using the publicly available *nf-core/rnaseq* pipeline version
3.12[84] implemented in Nextflow 23.10.0[85] with minimal command. In brief, lane-level reads
were trimmed using *Trim Galore* 0.6.7[86] and aligned to the mouse genome (GRCm39) using
STAR 2.7.9a[87]. Gene-level and transcript-level quantification was done by *Salmon v1.10.1*[88].
All analysis was performed using custom scripts in *R versi*on 4.3.2 using the *DESeq2 v.1.32.0*
framework[89].
1051
1052
MethylSeq
TdTomato-positive E14.5 MGE cells from *Sst-Cre/tdTomato* and *Sst-Cre/tdTomato/ Dnmt1*
loxP² were FAC-sorted into 100 µL HBSS. Samples were mixed by inversion and stored at -
80°C until processing.
DNA was isolated from at least 50.000 pooled cells per sample using the PureLink genomic
DNA Mini Kit (Thermo Fisher Scientic, U.S.A.). Per genotype, the samples were initially pooled
by volume (200 uL each), supplemented with lysis buffer provided in the kit and incubated with
2 µL proteinase K (Thermo Fisher Scientific, U.S.A.) at 55°C for 10 min. DNA was precipitated
with 100% ethanol. Samples were bound to one column per genotype each, washed as
instructed, briefly dried and eluted with 35 µL Tris-EDTA (TE) buffer. DNA concentrations were
determined using the Qubit™ 4 flourometer with the 1x dsDNA HS reagent (Thermo Fisher
Scientic, U.S.A.). Quality/integrity was checked on a Tape Station 4200 using the Genomic
DNA kit (Agilent Technologies, U.S.A.).
Libraries were generated using the Enzymatic Methyl-seq kit for Illumina (#E7120, New
England Biolabs, U.S.A.). After enzymatic shearing for 25 min at 37°C with the NEBNext
UltraShear enzyme and buffer, followed by an inactivation at 65°C for 15 min, 5-
methylcytosines were oxidized by TET2 after adaptor ligation. Samples were denatured with
formamide and deaminated with APOBEC. Libraries were subjected to 7 cycles of upscale
PCR (98°C for 10 s, 62°C for 30 s, and 65°C for 60 s). Library concentrations were determined
with a Qubit™ 4 fluorometer and the 1x dsDNA HS reagent (Thermo Fisher Scientific, U.S.A.).

All recommended clean-up steps were performed with magnetic AMPure XP beads (Beckman
Coulter, U.S.A.). Library size was assessed on a Tape Station 4200 using the DNA D5000 kit
(Agilent Technologies, U.S.A.). Concentrations were determined with a Qubit™ 4 fluorometer
and the 1x dsDNA HS reagent (Thermo Fisher Scientific, U.S.A.).

Libraries were diluted, equimolarly pooled, denatured, and clustered onto an Illumina NextSeq
500/550 Mid Output Kit v2.5 (300 cycles) flow cell (Illumina, U.S.A.). The 1% PhiX control
library was spiked in to improve base calling accuracy. Paired-end sequencing was performed
with 151 cycles.

FASTQ files were generated using bcl2fastq (Illumina, U.S.A.). To facilitate reproducible
analysis, samples were processed using the publicly available nf-core/methylseq pipeline
version 2.6.0[84] implemented in Nextflow 23.10.0[85] with the minimal command. Briefly, FastQC
(http://www.bioinformatics.babraham.ac.uk/projects/fastqc) and Trimmomatic[90] were utilized
to perform quality control, and clean reads were aligned to the reference genome using
Bismark[91] to account for bisulfite conversion and identify methylated cytosines. Differentially
methylated sites and regions were called by the R package DSS[92] and annotated using the
GENCODE VM23 dataset. Individual sites were counted as associated with a gene if located
within the gene body or within 500 bp of the transcription start site (TSS). Regions were
similarly annotated with a margin of 1 kb in front of the TSS. Methylation tracks were converted
from mm10 to mm39 using the USCS LiftOver tool[93] for integration with RNA seq data and
visualized in IGV[94]. To investigate promoter methylation in particular, the mean methylation
was calculated from all covered CpG sites within 1 kbp upstream and 100 bp downstream of
the TSS for genes of interest using custom Python scripts. The sequence of the promoter
regions (obtained from Ensembl release 112[95]) within these limits was further analyzed for the
enrichment of previously identified DNMT1-binding motifs using the AME algorithm provided
within the MEME suite[96]. Gene sets of interest were analyzed for the enrichment of gene
ontology terms using ShinyGO 0.80 with the biological processes' dataset[97].

1099

Chromatin immunoprecipitation (ChIP) and sequencing
Cerebellar granule cells[98] were cultured and treated as previously described[99]. DNMT1-
interacting chromatin was immunoprecipitated via native chromatin immunoprecipitation
(ChIP)[99]. The sequencing libraries were prepared using the NEBNext® Ultra II DNA Library
Prep kit (#E7410S, New England Biolabs, U.S.A.) as per manufacturer's instructions. The input
controls were pooled across samples in equimolar proportions. The libraries were amplified
via PCR on a T100 Thermal Cycler (Bio-Rad, U.S.A.) with the following conditions: initial
denaturation at 98°C for 30 s, 15 cycles of denaturation at 98°C for 10 s and
annealing/extension at 65°C for 75 s, followed by a final extension at 65°C for 5 min. To
account for the laddering caused by the enzymatic digestion, the libraries were size selected
on a 2% (v/v) agarose gel and purified using the QIAquick Gel Extraction Kit (#28704, Qiagen,
Germany). The libraries were sequenced on the NextSeq platform using 75 bp single-end
reads at the IZKF Genomics Facility (University Hospital Aachen).

FASTQ files generated from the ChIP-seq experiment were processed using the nf-
core/chipseq pipeline (version 2.0.0) with the mm10 reference genome[84]. Differential peak
calling between the samples was performed using THOR with default settings[100]. The identified
peaks with -log10p > 1.4 in the control samples are listed in Supplementary Table 4.

Sequences within a 500 bp window surrounding these peaks were analyzed for motif detection
using MEME-ChIP[101].
1119
1120
1121

Single-cell RNA-sequencing
The dorsal telencephalons from E14.5 C57BL/6J mice were dissected. Single-cell
suspensions were prepared using a papain dissociation system (#LK003150; Worthington
Biochemical Corporation, U.S.A) and all centrifugation steps were done at 4°C. Cells were
manually counted using Trypan Blue exclusion, which consistently showed a viability greater
than 95%.
ScRNA-seq libraries were prepared using the 10x Genomics 3' Gene Expression Kit v3.1 (10x
Genomics, U.S.A). Each sample had a target of 10'000 cells and was independently loaded in
the Chromium Controller (10x Genomics, U.S.A). Quality control of the cDNA and final libraries
was conducted using the High Sensitivity RNA D5000 and D1000 ScreenTape assays (Agilent
Technologies, U.S.A.), respectively. Libraries were pooled and sequenced on a NovaSeq 6000
using the S2 Reagent Kit v1.5 with 100 cycles (Illumina, U.S.A).
FASTQ files were processed using *CellRanger v7.0.0*. All bioinformatics analysis was
performed in *R 4.4.0* running under *Ubuntu 20.04.6 LTS*. The *Seurat v5* package[102] was used
for filtering, quality control, clustering, and annotation of the data. *CellChat v2.1.2*[49] was used
to compute the cell communication probabilities among different clusters.
1138
1139

Neuropixels recordings and behavioral experiments
Surgical Procedures
Mice received systemic analgesia prior to surgical procedures in form of subcutaneous
injections of carprofen (4 mg/kg, Rimadyl, Zoetis GmbH, Germany) and buprenorphine
(0.1 mg/kg, Buprenovet sine, Bayer Vital GMBH, Germany) and were then anesthetized using
(1%–2.5%) isoflurane in oxygen. The eyes were covered with eye ointment (Bepanthen, Bayer
Vital GmbH, Germany) to prevent them from drying out. Following a local injection of
bupivacaine (0.08 ml of 0.25% Bucain 7.5mg/ml, Puren Pharma GmbH, Germany), the scalp
was incised and pushed outward to fix it in place using tissue adhesive (Vetbond, 3M). A
circular headbar was attached on top using dental cement (C&B Metabond, Parkell U.S.A.;
Ortho-Jet, Lang Dental). For electrophysiological recordings, craniotomies were performed to
inject a viral vector into the cortex (AAV1.shortCAG.dlox.hChR2(H134R).WPRE.hGHp, Zurich
Vector Core, viral titer = ~8x10^{12} vg/ml). We targeted primary somatosensory (S1) and visual
(V1) cortex (stereotactic coordinates; S1: -1.5 mm anteroposterior and -3.5 mm mediolateral
from Bregma; V1: -4 mm anteroposterior and -2 mm mediolateral from Bregma) and injected
a volume of 200 nl in bouts of 10 nl with a 14-s delay between bouts (Nanoject III, Drummond
Scientific, U.S.A.). For each coordinate, injections were performed at two cortical depths (300
and 600 μm), adding up to a total volume of 400 nl per injection site. The glass pipette was
removed at least 5 min after the full volume was injected to reduce the risk of backflow. After
viral injections a 3-mm wide round coverslip was placed inside the craniotomy and fixed with
light-curable dental cement (DE Flowable composite, Nordenta, Germany). The coverslips
contained a small opening (~0.2 mm) that was covered with silicon to allow access of the

Neuropixels probes to the cortex. After the surgery, all animals received the same injections
of carpofen and buprenorphine as in the beginning. They also received buprenorphine (0.009
mg/ml, Buprenovet sine, Bayer Vital GmbH, Germany) and enrofloxacin (0.0227 mg/ml, Baytril
5%, Bayer Vital GMBH, Germany) in their drinking water for three days following the surgery.
Behavioral training started after at least two weeks of recovery
1167

Neuropixels recordings

1168
Electrophysiological recordings were done with Neuropixels 1.0 Probes in head-fixed mice,
freely running on a wheel. For each mouse, we performed four recordings on consecutive
days. To later recover the probe position in each recording, the probes were painted with DiD
cell labeling solution (Invitrogen V22887, U.S.A.) before each recording. To record neural
responses to tactile stimulation across all cortical layers, the probes were inserted at an
orthogonal angle to the brain surface (~10°). The stimulation protocol was started 5–10 min
after insertion, we recorded high pass-filtered data above 300 Hz at 30 kHz and low pass-
filtered signals between 0.5–100 Hz at 2.5 kHz from the bottom 384 channels of the
Neuropixels probe (~3.8 mm active recording area). Signals were acquired with an external
Neuropixels PXIe card (IMEC, Belgium) used in a PXIe-Chassis (PXIe-1071, National
Instruments, USA). Triggers and control signals for different stimuli and wheel movements
were separately recorded as analog and digital signals using the SpikeGLX software (Janelia
Farm Research Campus, U.S.A.; Bill Karsh).
Visual stimulation to induce gamma oscillations consisted of 5-s long full-field drifting square
wave gratings at horizontal orientation and a spatial frequency of 0.04 cycles per degree. All
visual stimuli were presented at a 17 cm distance from the right eye on a gamma-corrected
LED-backlit LCD monitor (Viewsonic VX3276-2K-MHD-2, 32", 60Hz refresh rate). Tactile
stimulation was controlled by a microcontroller-based finite state machine (Bpod r2, Sanworks,
U.S.A.) using custom Matlab code (2020b, MathWorks). Tactile stimuli consisted of 20-ms long
air puffs at 0.03 bar pressure directed at the distal parts of the whiskers. In each trial, two air
puffs were presented at an inter-stimulus interval of 500 ms. The overall stimulation protocol
consisted of 50 tactile trials and 50 visual trials in randomized order. Inter-trial intervals were
also randomized between 3–5 seconds.
To extract spiking activity, channels that were broken or outside of the brain were detected
and removed from the recording using the SpikeInterface analysis toolbox. The remaining
channels were time-shifted and median-subtracted across channels and time. Corrected
channels were then spike-sorted with the Kilosort 2.5 software and manually curated using
phy. All sorted spiking and local field potential data were analyzed using custom Matlab code
(2020b, MathWorks).
Peri-event time histograms (PETHs) for sensory responses in each cluster were computed
with a bin size of 10 ms and baseline-subtracted by the mean activity within 1 second before
the laser onset (between 2 to 1 seconds before the first tactile stimulus). Trial-averaged local
field potentials (LFPs) in response to sensory stimulation were similarly baseline-corrected. To
compute current source densities (CSDs) from LFP signals, we used the inverse CSD method
by Pettersen et al. (2006)[64]. We applied the spline iCSD method, assuming a smoothly varying
CSD between electrode contacts, based on polynomial interpolation. We assumed a
homogeneous, isotropic conductivity of $\sigma = 0.3$ S/m within and directly above the cortex[64]. To
reduce spatial noise, the estimated CSD was subsequently convolved with a Gaussian spatial

filter with a standard deviation of 0.1 mm. For LFP and CSD analyses, the signals from
neighboring contacts at the same depth were averaged together to improve signal quality.
1209

Nest-building procedure
Nest-shredding and -building tests were performed with male mice between 12 and 24 weeks
of age. Both *Sst-Cre* strains were tested by placing respective individuals in separate housing
cages with ad libitum access to food and water but without further enrichment except for fresh
bedding and standardized nest-building material (standardized autoclaved nestlets with a
weight of 2–3 g (5 cm x 5 cm, Lexx, Germany). For each nestlet the weight was determined
before the experiment which took place during the murine active phase (5 p.m. to 9 a.m.). After
testing, all mice were transferred to their home cages and the percentual shredded material
as well as the corresponding nest building score[76] were recorded.
1219

Pentylenetetrazole (PTZ)-induced epileptic seizure protocol
Male *Sst-Cre/tdTomato* and *Sst-Cre/tdTomato/Dnmt1 LoxP²* mice (13- to 17-week-old)
underwent an intraperitoneal administration of pentylenetetrazole (PTZ, Sigma-Aldrich,
U.S.A.) with 10 mg/kg living weight. PTZ was diluted in sterile Ringer's solution (pH 7.3–7.4)
to a concentration of 5 μg/μL beforehand. All mice received an intraperitoneal injection of PTZ
(40 μL PTZ-solution per 20 g living weight) every ten minutes. All tested animals were
monitored in transparent cages with no additional enrichment. Experiments ended as soon as
the animal had a tonic-clonic seizure with significantly overstretched limbs and no further
immediate recovery (event IV) or after the time limit of 150 minutes was reached. Afterwards,
all mice were sacrificed with an overdose of 5% (v/v) isoflurane followed by cervical dislocation
after reaching surgical tolerance, verified by loss of withdrawal response to toe pinch.
1231

Statistics and Figure illustration
Statistical tests were performed using Matlab, Phyton, and GraphPad Prism. Unless otherwise
stated, normally distributed data was tested applying unpaired and two-tailed Student's *t*-test
including a subsequent Welsh's correction, two-tailed nested *t*-test, and two-way ANOVA
followed by Bonferroni correction. For non-normally distributed data we used a Wilcoxon *rank-*
sum test. Respective statistical tests can be found in the main text and individual captions.
Unless otherwise stated, all depicted error bars represent the standard error of the mean
(SEM). Violin plots show the distribution of the numeric data using density curves, with the
respective frequency of data points represented by the width. Straight lines represent the
median and dotted lines the corresponding interquartile range.
1242

References
1.   Vanderhaeghen, P. & Polleux, F. Developmental mechanisms underlying the
evolution of human cortical circuits. *Nat Rev Neurosci* **24**, 213–232 (2023).
2.   Delaunay, D., Kawaguchi, A., Dehay, C. & Matsuzaki, F. Division modes and physical
asymmetry in cerebral cortex progenitors. *Current Opinion in Neurobiology* **42**, 75–83 (2017).
3.   Lin, Y. *et al.* Behavior and lineage progression of neural progenitors in the
mammalian cortex. *Current Opinion in Neurobiology* **66**, 144–157 (2021).
4.   Harris, K. D. & Shepherd, G. M. G. The neocortical circuit: themes and variations. *Nat*
Neurosci **18**, 170–181 (2015).
5.   Gerstmann, K. *et al.* Thalamic afferents influence cortical progenitors via ephrin A5-
EphA4 interactions. *Development* **142**, 140–150 (2015).

6.   Gerstmann, K. *et al.* A balance of noncanonical Semaphorin signaling from the
cerebrospinal fluid regulates apical cell dynamics during corticogenesis. *Sci. Adv.* **8**,
eabo4552 (2022).
7.   Gerstmann, K. & Zimmer, G. The role of the Eph/ephrin family during cortical
development and cerebral malformations. (2018).
8.   Reillo, I. *et al.* A Complex Code of Extrinsic Influences on Cortical Progenitor Cells of
Higher Mammals. *Cerebral Cortex* **27**, 4586–4606 (2017).
9.   Villalba, A., Götz, M. & Borrell, V. The regulation of cortical neurogenesis. in *Current
Topics in Developmental Biology* vol. 142 1–66 (Elsevier, 2021).
10.   Kepecs, A. & Fishell, G. Interneuron cell types are fit to function. *Nature* **505**, 318–
326 (2014).
11.   Kessaris, N. & Denaxa, M. Cortical interneuron specification and diversification in the
era of big data. *Current Opinion in Neurobiology* **80**, 102703 (2023).
12.   Lim, L., Mi, D., Llorca, A. & Marín, O. Development and Functional Diversification of
Cortical Interneurons. *Neuron* **100**, 294–313 (2018).
13.   Wonders, C. P. & Anderson, S. A. The origin and specification of cortical
interneurons. *Nat Rev Neurosci* **7**, 687–696 (2006).
14.   Reichard, J. & Zimmer-Bensch, G. The Epigenome in Neurodevelopmental Disorders.
Front. Neurosci. **15**, 776809 (2021).
15.   Fogarty, M. *et al.* Spatial Genetic Patterning of the Embryonic Neuroepithelium
Generates GABAergic Interneuron Diversity in the Adult Cortex. *J. Neurosci.* **27**, 10935–
10946 (2007).
16.   Butt, S. J. B. *et al.* The Temporal and Spatial Origins of Cortical Interneurons Predict
Their Physiological Subtype. *Neuron* **48**, 591–604 (2005).
17.   Niquille, M. *et al.* Neurogliaform cortical interneurons derive from cells in the preoptic
area. *eLife* **7**, e32017 (2018).
18.   Gelman, D. M. *et al.* The Embryonic Preoptic Area Is a Novel Source of Cortical
GABAergic Interneurons. *J. Neurosci.* **29**, 9380–9389 (2009).
19.   Zimmer, G. *et al.* Bidirectional ephrinB3/EphA4 signaling mediates the segregation of
MGE and POA derived interneurons in the deep and superficial migratory stream. *J
Neuroscience* **50**, (2011).
20.   Zimmer, G. *et al.* Chondroitin Sulfate Acts in Concert with Semaphorin 3A to Guide
Tangential Migration of Cortical Interneurons in the Ventral Telencephalon. *Cerebral Cortex*
20, 2411–2422 (2010).
21.   Zimmer, G. *et al.* Ephrin-A5 acts as a repulsive cue for migrating cortical
interneurons. *Eur J of Neuroscience* **28**, 62–73 (2008).
22.   Zimmer-Bensch, G. Diverse facets of cortical interneuron migration regulation -
Implications of neuronal activity and epigenetics. *Brain research* **1700**, 160–169 (2018).
23.   Lim, Y. *et al.* ARX regulates cortical interneuron differentiation and migration. Preprint
at https://doi.org/10.1101/2024.01.31.578282 (2024).
24.   Wang, Y. *et al.* CXCR4 and CXCR7 Have Distinct Functions in Regulating
Interneuron Migration. *Neuron* **69**, 61–76 (2011).
25.   Bartolini, G. *et al.* Neuregulin 3 Mediates Cortical Plate Invasion and Laminar
Allocation of GABAergic Interneurons. *Cell Reports* **18**, 1157–1170 (2017).
26.   Yildiz, C. B. & Zimmer-Bensch, G. Role of DNMTs in the Brain. *Adv Exp Med Biol*
1389, 363–394 (2022).
27.   Pensold, D. *et al.* The DNA Methyltransferase 1 (DNMT1) Controls the Shape and
Dynamics of Migrating POA-Derived Interneurons Fated for the Murine Cerebral Cortex.
Cereb Cortex **27**, 5696–5714 (2017).
28.   Urban-Ciecko, J. & Barth, A. L. Somatostatin-expressing neurons in cortical networks.
Nat Rev Neurosci **17**, 401–409 (2016).
29.   Liodis, P. *et al. Lhx6* Activity Is Required for the Normal Migration and Specification of
Cortical Interneuron Subtypes. *J. Neurosci.* **27**, 3078–3089 (2007).
30.   Lim, L. *et al.* Optimization of interneuron function by direct coupling of cell migration
and axonal targeting. *Nat Neurosci* **21**, 920–931 (2018).
31.   Hu, J. S., Vogt, D., Sandberg, M. & Rubenstein, J. L. Cortical interneuron

development: a tale of time and space. *Development* **144**, 3867–3878 (2017).
32.　Bock, C. Analysing and interpreting DNA methylation data. *Nat Rev Genet* **13**, 705–
719 (2012).
33.　Lister, R. *et al.* Human DNA methylomes at base resolution show widespread
epigenomic differences. *Nature* **462**, 315–322 (2009).
34.　Mayer, C. *et al.* Developmental diversification of cortical inhibitory interneurons.
Nature **555**, 457–462 (2018).
35.　Gruner, H. N., Kim, M. & Mastick, G. S. Robo1 and 2 Repellent Receptors Cooperate
to Guide Facial Neuron Cell Migration and Axon Projections in the Embryonic Mouse
Hindbrain. *Neuroscience* **402**, 116–129 (2019).
36.　Talebian, A. *et al.* Autonomous and non-autonomous roles for ephrin-B in interneuron
migration. *Developmental Biology* **431**, 179–193 (2017).
37.　Marín, O., Yaron, A., Bagri, A., Tessier-Lavigne, M. & Rubenstein, J. L. Sorting of
striatal and cortical interneurons regulated by semaphorin-neuropilin interactions. *Science*
293, 872–875 (2001).
38.　Gabel, H. W. *et al.* Disruption of DNA-methylation-dependent long gene repression in
Rett syndrome. *Nature* **522**, 89–93 (2015).
39.　Nan, X. *et al.* Transcriptional repression by the methyl-CpG-binding protein MeCP2
involves a histone deacetylase complex. *Nature* **393**, 386–389 (1998).
40.　Boyes, J. & Bird, A. DNA methylation inhibits transcription indirectly via a methyl-CpG
binding protein. *Cell* **64**, 1123–1134 (1991).
41.　Pensold, D. *et al.* DNA Methylation-Mediated Modulation of Endocytosis as Potential
Mechanism for Synaptic Function Regulation in Murine Inhibitory Cortical Interneurons.
Cereb Cortex **30**, 3921–3937 (2020).
42.　Batista-Brito, R. *et al.* The Cell-Intrinsic Requirement of Sox6 for Cortical Interneuron
Development. *Neuron* **63**, 466–481 (2009).
43.　van den Berghe, V. *et al.* Directed Migration of Cortical Interneurons Depends on the
Cell-Autonomous Action of Sip1. *Neuron* **77**, 70–82 (2013).
44.　McKinsey, G. L. *et al.* Dlx1&2-Dependent Expression of Zfhx1b (Sip1, Zeb2)
Regulates the Fate Switch between Cortical and Striatal Interneurons. *Neuron* **77**, 83–98
(2013).
45.　Fisher, J. *et al.* Cortical somatostatin long-range projection neurons and interneurons
exhibit divergent developmental trajectories. *Neuron* **112**, 558-573.e8 (2024).
46.　Sessa, A. *et al.* Tbr2-positive intermediate (basal) neuronal progenitors safeguard
cerebral cortex expansion by controlling amplification of pallial glutamatergic neurons and
attraction of subpallial GABAergic interneurons. *Genes Dev.* **24**, 1816–1826 (2010).
47.　Silva, C. G. *et al.* Cell-Intrinsic Control of Interneuron Migration Drives Cortical
Morphogenesis. *Cell* **172**, 1063-1078.e19 (2018).
48.　Kischel, A., Audouard, C., Fawal, M.-A. & Davy, A. Ephrin-B2 paces neuronal
production in the developing neocortex. *BMC Dev Biol* **20**, 12 (2020).
49.　Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat
Commun* **12**, 1088 (2021).
50.　Hevner, R. F. *et al.* Tbr1 Regulates Differentiation of the Preplate and Layer 6.
Neuron **29**, 353–366 (2001).
51.　Bulfone, A. *et al.* T-Brain-1: A homolog of Brachyury whose expression defines
molecularly distinct domains within the cerebral cortex. *Neuron* **15**, 63–78 (1995).
52.　Vasistha, N. A. *et al.* Cortical and Clonal Contribution of Tbr2 Expressing Progenitors
in the Developing Mouse Brain. *Cereb. Cortex* **25**, 3290–3302 (2015).
53.　Kowalczyk, T. *et al.* Intermediate neuronal progenitors (basal progenitors) produce
pyramidal-projection neurons for all layers of cerebral cortex. *Cereb Cortex* **19**, 2439–2450
(2009).
54.　Mihalas, A. B. *et al.* Intermediate Progenitor Cohorts Differentially Generate Cortical
Layers and Require Tbr2 for Timely Acquisition of Neuronal Subtype Identity. *Cell Reports*
16, 92–105 (2016).
55.　Harrison-Uy, S. J. & Pleasure, S. J. Wnt Signaling and Forebrain Development. *Cold
Spring Harbor Perspectives in Biology* **4**, a008094–a008094 (2012).

56. Nian, F.-S. & Hou, P.-S. Evolving Roles of Notch Signaling in Cortical Development.
Front. Neurosci. **16**, 844410 (2022).
57. Cárdenas, A. *et al.* Evolution of Cortical Neurogenesis in Amniotes Controlled by
Robo Signaling Levels. *Cell* **174**, 590-606.e21 (2018).
58. Smith, A., Robinson, V., Patel, K. & Wilkinson, D. G. The EphA4 and EphB1 receptor
tyrosine kinases and ephrin-B2 ligand regulate targeted migration of branchial neural crest
cells. *Current Biology* **7**, 561–570 (1997).
59. Riedemann, T. Diversity and Function of Somatostatin-Expressing Interneurons in the
Cerebral Cortex. *IJMS* **20**, 2952 (2019).
60. Ward, C. *et al.* Developmental Disruption of Mef2c in Medial Ganglionic Eminence–
Derived Cortical Inhibitory Interneurons Impairs Cellular and Circuit Function. *Biological*
Psychiatry S000632232401360X (2024) doi:10.1016/j.biopsych.2024.05.021.
61. Allaway, K. C. *et al.* Genetic and epigenetic coordination of cortical interneuron
development. *Nature* **597**, 693–697 (2021).
62. Pai, E. L.-L. *et al.* Maf and Mafb control mouse pallial interneuron fate and maturation
through neuropsychiatric disease gene regulation. *eLife* **9**, e54903 (2020).
63. Musall, S., Kaufman, M. T., Juavinett, A. L., Gluf, S. & Churchland, A. K. Single-trial
neural dynamics are dominated by richly varied movements. *Nat Neurosci* **22**, 1677–1686
(2019).
64. Pettersen, K. H., Devor, A., Ulbert, I., Dale, A. M. & Einevoll, G. T. Current-source
density estimation based on inversion of electrostatic forward solution: Effects of finite extent
of neuronal activity and conductivity discontinuities. *Journal of Neuroscience Methods* **154**,
116–133 (2006).
65. Petersen, C. C. H. The Functional Organization of the Barrel Cortex. *Neuron* **56**, 339–
355 (2007).
66. Fries, P., Reynolds, J. H., Rorie, A. E. & Desimone, R. Modulation of Oscillatory
Neuronal Synchronization by Selective Visual Attention. *Science* **291**, 1560–1563 (2001).
67. Veit, J., Hakim, R., Jadi, M. P., Sejnowski, T. J. & Adesnik, H. Cortical gamma band
synchronization through somatostatin interneurons. *Nat Neurosci* **20**, 951–959 (2017).
68. Ahnaou, A., Huysmans, H., Van De Casteele, T. & Drinkenburg, W. H. I. M. Cortical
high gamma network oscillations and connectivity: a translational index for antipsychotics to
normalize aberrant neurophysiological activity. *Transl Psychiatry* **7**, 1285 (2017).
69. Yamamoto, J., Suh, J., Takeuchi, D. & Tonegawa, S. Successful Execution of
Working Memory Linked to Synchronized High-Frequency Gamma Oscillations. *Cell* **157**,
845–857 (2014).
70. Kas, M. J. H., Fernandes, C., Schalkwyk, L. C. & Collier, D. A. Genetics of
behavioural domains across the neuropsychiatric spectrum; of mice and men. *Mol Psychiatry*
12, 324–330 (2007).
71. Liguz-Lecznar, M., Dobrzanski, G. & Kossut, M. Somatostatin and Somatostatin-
Containing Interneurons—From Plasticity to Pathology. *Biomolecules* **12**, 312 (2022).
72. Jirkof, P. Burrowing and nest building behavior as indicators of well-being in mice.
Journal of Neuroscience Methods **234**, 139–146 (2014).
73. Perry, G., Brindley, L. M., Muthukumaraswamy, S. D., Singh, K. D. & Hamandi, K.
Evidence for increased visual gamma responses in photosensitive epilepsy. *Epilepsy*
Research **108**, 1076–1086 (2014).
74. Mandhane, S. N., Aavula, K. & Rajamannar, T. Timed pentylenetetrazol infusion test:
A comparative analysis with s.c.PTZ and MES models of anticonvulsant screening in mice.
Seizure **16**, 636–644 (2007).
75. Chenn, A. & Walsh, C. A. Regulation of Cerebral Cortical Size by Control of Cell
Cycle Exit in Neural Precursors. *Science* **297**, 365–369 (2002).
76. Deacon, R. M. Assessing nest building in mice. *Nat Protoc* **1**, 1117–1119 (2006).
77. Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization
system for the whole mouse brain. *Nat Neurosci* **13**, 133–40 (2010).
78. Jackson-Grusby, L. *et al.* Loss of genomic methylation causes p53-dependent
apoptosis and epigenetic deregulation. *Nat Genet* **27**, 31–39 (2001).
79. Symmank, J., Gölling, V., Gerstmann, K. & Zimmer, G. The Transcription Factor

LHX1 Regulates the Survival and Directed Migration of POA-derived Cortical Interneurons. *Cereb Cortex* **29**, 1644–1658 (2019).

80.   Hahn, A. *et al.* DNA Methyltransferase 1 (DNMT1) Function Is Implicated in the Age-Related Loss of Cortical Interneurons. *Frontiers in cell and developmental biology* **8**, 639 (2020).

81.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676–682 (2012).

82.   Saxena, S., Maroju, P. A., Choudhury, S., Anne, A. & Mohan, K. N. Analysis of transcript levels of a few schizophrenia candidate genes in neurons from a transgenic mouse embryonic stem cell model overexpressing DNMT1. *Gene* **757**, 144934 (2020).

83.   Singh, M., Saxena, S. & Mohan, K. N. DNMT1 downregulation as well as its overexpression distinctly affect mostly overlapping genes implicated in schizophrenia, autism spectrum, epilepsy, and bipolar disorders. *Front. Mol. Neurosci.* **16**, 1275697 (2023).

84.   Ewels, P. A. *et al.* The nf-core framework for community-curated bioinformatics pipelines. *Nat Biotechnol* **38**, 276–278 (2020).

85.   Di Tommaso, P. *et al.* Nextflow enables reproducible computational workflows. *Nat Biotechnol* **35**, 316–319 (2017).

86.   Krueger, F., James, F., Ewels, P., Afyounian, E. & Schuster-Boeckler, B. FelixKrueger/TrimGalore: v0.6.7 - DOI via Zenodo. Zenodo https://doi.org/10.5281/ZENODO.5127899 (2021).

87.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

88.   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods* **14**, 417–419 (2017).

89.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

90.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

91.   Krueger, F. & Andrews, S. R. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. *Bioinformatics* **27**, 1571–1572 (2011).

92.   Feng, H., Conneely, K. N. & Wu, H. A Bayesian hierarchical model to detect differentially methylated loci from single nucleotide resolution sequencing data. *Nucleic Acids Research* **42**, e69–e69 (2014).

93.   Hinrichs, A. S. The UCSC Genome Browser Database: update 2006. *Nucleic Acids Research* **34**, D590–D598 (2006).

94.   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29**, 24–26 (2011).

95.   Harrison, P. W. *et al.* Ensembl 2024. *Nucleic Acids Res* **52**, D891–D899 (2024).

96.   Buske, F. A., Bodén, M., Bauer, D. C. & Bailey, T. L. Assigning roles to DNA regulatory motifs using comparative genomics. *Bioinformatics* **26**, 860–866 (2010).

97.   Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals and plants. *Bioinformatics* **36**, 2628–2629 (2020).

98.   Fossale, E. *et al.* Membrane trafficking and mitochondrial abnormalities precede subunit c deposition in a cerebellar cell model of juvenile neuronal ceroid lipofuscinosis. *BMC Neurosci* **5**, 57 (2004).

99.   Yildiz, C. B. *et al.* EphrinA5 regulates cell motility by modulating Snhg15/DNA triplex-dependent targeting of DNMT1 to the Ncam1 promoter. *Epigenetics & Chromatin* **16**, 42 (2023).

100.   Allhoff, M., Seré, K., F Pires, J., Zenke, M. & G Costa, I. Differential peak calling of ChIP-seq signals with replicates with THOR. *Nucleic Acids Res* **44**, e153 (2016).

101.   Machanick, P. & Bailey, T. L. MEME-ChIP: motif analysis of large DNA datasets. *Bioinformatics* **27**, 1696–1697 (2011).

102.   Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-cell analysis. *Nat Biotechnol* (2023) doi:10.1038/s41587-023-01767-y.

Main Figures

Figure 1:

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

a
Score 1 Score 2 Score 3 Score 4 Score 5

b control control

c KO KO

d used nest material [%]
**
control KO

e nestbuilding score
***
control KO

f 1st PTZ injection
event I event II event III event IV
small seizure types tonic-clonic seizures
small seizures only small + tonic-clonic seizures
seizure-free period convulsion period

g convulsion period [min]
*
control KO

h small seizures only [min]
**
control KO

Extended Data Figure Legends

Extended Data Figure E1:

Validation of SST expression and reduced DNMT1 levels in tdTomato-positive cells. (a) Mating strategy to obtain *Sst-Cre/tdTomato* (control) and *Sst-Cre/tdTomato/Dnmt1 loxP²* (KO) mice. *Sst^{tm2.l(cre)Zjh}*/J mice were mated with *tdTomato* reporter mice (*B6.CgGt(ROSA)26Sor^{tml 4(CAG-tdTomato)Hze}*) resulting in *Sst-Cre/tdTomato* mice with an internal ribosomal entry site (*IRES*), a *Cre-recombinase* sequence, a *polyA* sequence, and a *frt*-flanked neo cassette located in the 3' untranslated region (UTR) of the *Somatostatin* (*Sst*) locus on chromosome 16, limiting the respective *Cre* expression to *Sst*-positive neurons. The *tdTomato* sequence and a loxP-flanked stop cassette are located in the *Rosa26* locus. Triple transgenic *Sst-Cre/tdTomato/Dnmt1 loxP²* mice were obtained by crossing control mice with a *Dnmt1 loxP²* strain (*B6; 129Sv-Dnmt1tm4Jae/J*; exon 4 and 5 of the *Dnmt1* gene are *loxP*-flanked, resulting in a null allele of these loci and a subsequent DNMT1 deficiency. **(b)** Exemplary microphotographs depicting SST expression in tdTomato-cells in sagittalbrain sections of a four-month-old male *Sst-Cre/tdTomato* mouse (Bregma 2.76). **(c)** Magnified images of the areas outlined in b. Scale bars: 100 *μ*m in b and 20 *μ*m c. **(d-g)** Quantification of DNMT1 expression by immunohistochemistry in *Dnmt1*-control **(d, e)** and *Dnmt1*-KO **(f, g)** male mice (six-month-old individuals) in 30 *μ*m, fixated coronal cryosections. Scale bars **(d, f)** 100 *μ*m and 10 *μ*m **(e,g)**. **(h)** Quantification of the ratio of DNMT1-positive to DNMT1-negative *tdTomato*-expressing cells (Bregma 1.18, 1.10, 0.14, -0.22, -2.92. -3.08, -3.28 were used for quantification). Two-tailed nested *t*-test, *(p < 0.001 ***, N = 3 for Sst-Cre/tdTomato (n = 18* slices) and *Sst-Cre/tdTomato/Dnmt1 loxP²* (*n =* 17 slices), respectively.

Extended Data Figure E2:

Characteristics of genes that were upregulated in expression as well as differentially methylated in FAC-sorted *Sst-Cre/tdTomato/Dnmt1 loxP²* neurons compared to *Sst-Cre/tdTomato* cells at E14.5 compared with the rest in the genome; Chi-squared and Student's *t*-tests were run to analyze if our gene set has special characteristics when compared with all the other genes (ShinyGO 0.80; http://bioinformatics.sdstate.edu/go/). **(a)** Gene ontology (GO), **(b)** bar plots, and **(c)** density plots.

Extended Data Figure E3:

(a) Gene ontology (GO) analysis for the set of genes that were upregulated in expression in *Sst*-positive neurons of *Sst-Cre/tdTomato/Dnmt1 loxP²* mice compared to *Sst-Cre/tdTomato* at E14.5. **(b)** Further characteristics are shown in bar plots (tested by Chi-squared test) and density plots **(c),** depicting the characteristics of this gene set compared with the entire set of detected transcripts (Student's *t*-tests).

Extended Data Figure E4:

RNA sequencing tracks combined with methylation profile for select genes obtained from FAC-sorted cells of *Sst-Cre/tdTomato* and *Sst-Cre/tdTomato/Dnmt1 loxP²* mice (E14.5). Forward and reverse tracks from both RNA sequencing runs were combined into one track per genotype. Similarly, both methylation tracks per genotype were combined. Differentially methylated regions identified by DSS are indicated below. Tracks were illustrated with IGV viewer V2.17.2, refseq mm39.

Extended Data Figure E5:

The genes upregulated in *Sst*-expressing interneurons of *Sst-Cre/tdTomato/Dnmt1-loxP²* mice partially overlap with genes that are downregulated after *Dnmt1* overexpression in neurons obtained from murine embryonic stem cells (ESCs). **(a)** Gene ontology (GO) analysis for the overlap. **(b)** Further characteristics are shown in bar plots (tested by Chi-squared test) and density plots **(c)**, depicting the characteristics of this gene set compared with the entire set of detected transcripts (Student's *t*-tests).

Extended Data Figure E6:

(a-c) ErbB4 level is increased after *Dnmt1* knockdown (KD) in MGE cells (E14.5 + 1div). Exemplary microphotographs showing control siRNA-treated (a) and *Dnmt1*-KD cells (b) stained for Phalloidin-647 and ErbB4. On the very right of each example, thermal-color-coded panels represent the respective fluorescent intensity of ErbB4 (therm. (thermal) LUT). A dark blue color indicates 0 and a red color represents 5000 fluorescent units. Scale bars: 10 μm. **(c)** Quantification of (a) and (b) comparing the integrated density (IntDens) of ErbB4 fluorescence signals of control and *Dnmt1*-KD MGEs. Unpaired, two-tailed Student's *t*-test with subsequent Welch's correction, $p < 0.001$ *** ($n = 88$ cells for control siRNA and 78 cells for *Dnmt1* siRNA, $N = 4$ experiments). **(d-f)** Live-cell imaging analysis of *Sst-Cre/tdTomato* and *Sst-Cre/tdTomato/Dnmt1 loxP²* cells in E14.5 organotypic brain slices (350 μm coronal sections, time-period of imaging: 20 h). Analysis of the migrated path length and the radial migration velocity are shown in **(d)** and **(e)**, respectively. **(f)** DNMT1 does not impact the migratory morphology of tdTomato-positive cells investigated in organotypic brain slices (350 μm coronal sections). Two-tailed nested *t*-test ($p < 0.01$ **) with $N = 3$ embryos and $n = 4$ slices for control and KO animals respectively, $n = 185$ cells for control, and $n = 219$ KO cells. **(g-i)** siRNA-mediated depletion of *Dnmt1* in MGE-derived single cells *in vitro* (E14.5 + 1div) does not affect the cellular morphology compared to control siRNA-transfected cells. Scale bars: 10 μm. Quantification is shown in **(i)**. Unpaired, two-tailed Student's *t*-test with subsequent Welsh's correction, $n = 60$ cells each for control siRNA and *Dnmt1* siRNA. siRNA: small-interfering RNA; LP: leading process, n.s.: not significant.

Extended Data Figure E7:

The cortical size and dimensions of its different regions do not differ between E14.5 *Sst-Cre/tdTomato/Dnmt1-loxP²* and E14.5 *Sst-Cre/tdTomato* brains. **(a)** Schematic illustration of the regions and extensions measured to analyze cortical extensions and areas of coronal, 50-*µ*m-thick, fixated slices. Respective quantifications are depicted in **(b)** and **(c)**. Two-tailed nested *t*-test, $N = 3$ embryos for both genotypes, $n = 13$ slices for *Sst-Cre/tdTomato* and $n = 9$ slices for *Sst-Cre/tdTomato/Dnmt1 loxP²*. MZ: marginal zone, IZ: intermediate zone, SVZ+VZ: subventricular and ventricular zone, not significant.

Extended Data Figure E8:

The cortical size and dimensions of its different regions are not different in E16.5 and E18.5 *Sst-Cre/tdTomato/Dnmt1 loxP²* embryos compared to age-matched *Sst-Cre/tdTomato* individuals. **(a, d)** Schematic illustration of the regions and extensions measured to analyze cortical dimensions and areas of coronal, 50-*µ*m-thick, fixated slices at E16.5 **(a)** and E18.5 **(d)**. Respective quantifications are depicted in **(b)**, **(c),** and **(e)**. **(b and c)** Quantification of the cortical dimension of embryos at E16.5. Unpaired nested *t*-test, $N = 4$ embryos for both genotypes, $n = 12$ slices for *Sst-Cre/tdTomato,* and $n = 15$ slices for *Sst-Cre/tdTomato/Dnmt1 loxP²*. **(e)** Quantification of the cortical dimension of embryos at E18.5. Unpaired nested *t*-test, $N = 4$ embryos for both genotypes, $n = 12$ slices for *Sst-Cre/tdTomato,* and $n = 9$ slices for *Sst-Cre/tdTomato/Dnmt1 loxP²*. MZ: marginal zone, CP: cortical plate, IZ: intermediate zone, SVZ+VZ: subventricular zone, n.s.: not significant.

Extended Data Figure E9:

TdTomato-expressing cells show immunoreactivity for neuropeptide Y and calretinin. Exemplary microphotographs of sagittal sections from a four-month-old *Sst-Cre/tdTomato* animal (Bregma 2.76). **(a** and **b)** Examples of immunohistochemical stainings using an anti-somatostatin antibody (green) and an anti-neuropeptide Y antibody (NPY, magenta). TdTomato-positive cells are depicted in red. **(c** and **d)** Examples of immunohistochemical stainings using an anti-somatostatin antibody (green) and an anti-calretinin antibody (magenta). TdTomato-positive cells are depicted in red. Scale bars in **(a)** and **(c)**: 100 µm. Scale bars in **(b)** and **(d)**: 20 µm (magnified views of sections outlined in **(a)** and **(c)**, respectively). **(e)** Mammalian phenotype ontology analysis for the enrichment of the genes upregulated in *Sst-Cre/tdTomato/Dnmt1 loxP²* animals at E14.5. ShinyGO 0.80, Curated.MGI pathway database.

Extended Data Figures

Extended Data Figure E1

Extended Data Figure E2

Extended Data Figure E3

Extended Data Figure E4

Extended Data Figure E5

Extended Data Figure E6

Extended Data Figure E7

Extended Data Figure E8

Extended Data Figure E9