CMPortal: a community-oriented, data-driven resource to inform protocol design for cardiac modelling from
human pluripotent stem cells

Chris Siu Yeung Chow[1], Sophie Shen[1], Amy Hanna[1], Sumedha Negi[1], Clarissa L. L. Tan[1], Shaine Chenxin Bao[1], Chen
Fang[1,2], James E. Hudson[3], Woo Jun Shim[1], Yuanzhao Cao[1,4], Nathan J. Palpant[1,4,*]

[1]*Institute for Molecular Bioscience, The University of Queensland, Brisbane, QLD, Australia*

[2]*Ochsner Clinical School-The University of Queensland School of Medicine, New Orleans, Louisiana*

[3]*Queensland Institute for Medical Research, Brisbane, QLD, Australia*

[4]*Co-senior authors*

[*]Corresponding author

Professor Nathan J. Palpant
Institute for Molecular Bioscience
The University of Queensland
306 Carmody Road, St. Lucia, QLD 4072 Australia
Email: n.palpant@uq.edu.au
16

Abstract

Protocol design and benchmarking is central to optimising model development using human pluripotent stem cell
derived cardiomyocytes (hPSC-CMs). By applying data mining to decades of research and hundreds of peer reviewed
studies, we evaluate how protocol variables associate with common properties of cardiac functional and physiological
maturation. This resource is publicly accessible through CMPortal, a community-oriented website that provides data-
driven tools for researchers to navigate leverage decades of knowledge for benchmarking protocol designs and outcomes
for their dedicated applications in developmental biology, disease modelling, and drug screening.

Human pluripotent stem cell-derived cardiomyocytes (hPSC-CMs) are a widely used model for cardiac research, enabling studies of development, physiology, disease mechanisms, and drug responses.[1] However, maturing hPSC-CMs toward adult-like phenotypes remains an ongoing challenge for modeling adult cardiac biology.[2] Despite development of maturation strategies involving specialized media, co-culture systems, and various types of matrix and biomaterials, optimizing protocol conditions suitable for modelling adult heart phenotypes remain challenging due to high variability in experimental design and outcomes.

Ewoldt et al.[3,4] recently reviewed 300 study designs and demonstrate that while maturity outcomes have improved over time, no significant differences exist between maturation strategies. They highlight difficulties in parsing individual effects of protocol variables due to complex design variables and inconsistent reporting, drawing attention to the need for data-driven approaches and community-oriented benchmarking strategies to improve the accuracy and efficiency of hPSC-CM protocol design.

Here, we develop a data-driven platform that offers user-guided analyses of protocol features and their association with functional outcomes. Rather than assuming direct correlations exist between protocol factors and physiological maturation, we use an unsupervised approach that enables detection of bidirectionality and context-dependency allowing parameters to be independently assessed.

We first expanded the Ewoldt database[3] to 322 studies, including 14 metabolic maturation studies and 8 publications to capture recent developments in the field. We also added protocol design features including cell line sex, ancestry, and transcriptomic markers of myofilament isoform switching. The final database has standardized nomenclature across 400 protocol features spanning five categories: experimental design variables (e.g., media composition factors, plating densities, matrix), analysis methods (e.g., imaging techniques, electrophysiology approaches), cell profiles (details about cell line including sex, ancestry, and source), study characteristics (e.g., publication year, journal), and measured outcomes (e.g., measured sarcomere length, contractile force, calcium handling, myofilament isoforms) (**Figure 1A**).

We applied a hybrid strategy combining conventional statistics and data mining to assess protocol features against 117 target parameters, such as 18 maturity indicators, cell profiles, and applications in disease and pharmacological modeling (**Tables S1 and S2**). Numerical measurements were binned into quantiles (e.g. Q1, Q2, etc.) to enable analysis of protocols ranked by their maturity outcomes. One-hot encoding was used to derive a data format suitable for machine learning, which also accounts for the significant missing data and assumes protocol variables, when not used or reported, are still potentially causal.

For data mining, we used random forest classifiers with 10,000 permuted feature sets to associate protocol designs with target parameters through Shannon entropy and Gini impurity, allowing the detection of statistically robust associations even in small and sparsely reported datasets (**Figure 1A** and **Table S3**). Dimensionality reduction via UMAP demonstrated high protocol heterogeneity across studies, consistent with reported findings[3] (**Figure 1B**). Despite this heterogeneity, polynomial regression confirmed that the database sufficiently captures protocol features for target endpoints relating to best and worst performing quantiles in a data-limited setting (**Figure 1C**). Quality checks using Jaccard index analysis confirmed that data transformation preserved feature distinctiveness (**Figure S1**).

Data mining revealed alignment between protocol variables and expected biological impacts on cardiomyocyte maturation (**Figures 1D-E**). For example, Q1 contractility protocols were significantly associated with Wnt modulation, fibroblast inclusion, and matrix stiffness which are well established factors impacting sarcomere assembly and force generation.[5,6] Importantly, we identified that identical maturation strategies could be linked to both best and worst maturation outcomes, demonstrating that this approach identifies context-dependent effects of protocol variables and not simply assume their generalized effects (**Figure 1F**). Specifically, the findings suggest that maturation parameters can be independently modulated by distinct sets of protocol variables, challenging the assumption that improving one cardiac maturity metric necessarily enhances all others.

We next used a spearman correlation analysis of all continuous variables in the database (n=31) to assess whether maturity measurements recapitulate expected physiological relationships (**Table S4**). Indeed, the data demonstrates that across hundreds of peer-reviewed studies, there are significant positive correlations between expected parameters including resting membrane potential and conduction velocity ($\rho=1$, $p<0.01$), as well as contractile force and 3D tissue

size (ρ=0.33, p<0.05) (**Figure 1G**). Interestingly, the database also identifies protocol correlations that provide important
considerations in co-culture design. Different cell type compositions revealed positive, null, and negative effects on
maturity indicators including calcium kinetics and sarcomere length (**Figure 1H**).

We integrated these findings into a community-oriented web resource, *CMPortal* (https://palpantlab.com/cmportal). The
Database Viewer enables searching of the 322 curated protocols based on five feature categories using techniques from
simple keyword matching to multi-criteria searches, with options to toggle column visibility, export customized datasets,
and evaluate relevant protocols of interest (**Figures 2A-C**). The Variable Search enables researchers to identify studies
and protocols based on topic areas and/or sets of protocol variables which enables unsupervised discovery of relevant
protocols based on desired and/or associated protocol features, even when the published protocols do not report the
selected metrics (**Figure 2D**). The Protocol Benchmarking tool enables researchers to evaluate differentiation and
maturation protocols against database standards. By uploading or manually selecting protocol features and the user's
own experimental measurements, the tool compares protocols through a radar chart that highlights maturity indicators
across multiple categories and visually distinguishes between predicted, normal-range, and outlier experimental values
(**Figures 2E-F**). Lastly, the Enrichment Browser allows researchers to identify protocol features significantly associated
with specific outcomes which serves as the data foundation for the search and benchmarking features. (**Figure 2G**).

We aimed to validate an unexpected finding from the database showing that contractility and sarcomere length are not
correlated maturation endpoints (**Figure 2H**), each having distinct protocol variable enrichments governing their
maturation (**Table S2**). This finding suggests that the Frank-Starling effect, one of the most fundamental principles
governing the relationship between heart pump function and load, develops through independent mechanisms. We
investigated this by first identifying enriched protocol characteristics of Q1 and Q5 contractility, choosing backbone
media compositions involving DMEM+fatty acids (FA) (Q1 contractility) and RPMI+B27 (Q5 contractility) (**Figure**
2I). DMEM+FA confers enhanced contractile maturity due to the high calcium concentration[7] and fatty acids[5] that
mimic postnatal development. As shown in data mining, impedance-based measurements showed significantly higher
peak contraction amplitude in DMEM+FA versus RPMI+B27 (**Figure 2J**), which was also supported by RNA-seq data
showing higher metabolic activity from glycolysis to mature fatty acid oxidation in DMEM+FA cells (**Figure 2K**).
Consistent with the database correlation, contractility changes occurred without significant variation in sarcomere length
(**Figure 2L**). This supports our observation that fundamental features of cardiac maturation, including interdependent
mechanisms controlling heart function like the Frank-Starling relationship, are not necessarily mechanistically linked in
development. *CMPortal* allows for unsupervised discovery of these context-dependent protocol variables, enabling
users to independently optimize protocols for specific functions.

CMPortal represents the first quantitative approach to overcome significant analytical difficulties of unreported, sparse
protocol data, successfully associating context-dependent protocol variables with cell modelling outcomes. These
findings challenge a fundamental but rarely challenged assumption that hPSC-CM maturity advances uniformly across
all metrics. Instead, the results support a modularised paradigm to develop protocols through selected metrics based on
experimental goals, aligning with recent literature demonstrating that attributes of hPSC-CM maturation can be
compartmentalized.[8] We translate these findings into user-friendly tools provided on a public website for researchers to
evaluate, design, and benchmark protocols. *CMPortal* provides a streamlined approach to literature review, inform
experimental design, and facilitate protocol benchmarking for understanding cardiac biology, diseases, and drug
responses as a community-oriented resource for hPSC-CM research.

Acknowledgements

123
124 This work has been supported by grant funding from the NHMRC (MRFCDDM000033 and 2007625 to NP), the Ian
125 Potter Foundation (31111380 to NP), and the National Heart Foundation of Australia (106721 to NP). J.E.H.
126 acknowledges the support of a Snow Medical Research Foundation Fellowship, Grant No. SMRF2019–060. Microscopy
127 was performed at the Institute for Molecular Bioscience Microscopy Facility which was established with the support of
128 the Australian Cancer Research Foundation (ACRF). We thank Meredith Redd for her assistance with project design.

Figure 1. *CMPortal* database construction and hybrid data mining approach.

(A) Schematic overview of the workflow for integrating and analysing 322 hPSC-CM studies to evaluate protocol designs and target parameters.

(B) UMAP visualizations project the database by five feature categories, enabling assessment of variability in protocol components and identification of shared trends, with annotations shown in color.

(C) Relationship between the number of significant enrichments and the number of training protocols. A polynomial model fitted to the best and worst quantiles of maturity indicators (pink and purple) suggests a plateau at 20–30 out of 413 protocol features significantly associated with maturity metrics. Grey indicates target parameters for intermediate maturity quantiles, cell profile, and study characteristics.

(D–E) Bar plots showing protocol feature metadata for the highest (D) and lowest (E) contraction force quantiles, based on enrichments with $p<0.025$. Yellow bars indicate the number of studies reporting the relevant variables; purple bars represent the confidence in true feature enrichment. Log-odds ratio (logOR) and Fisher's exact test (FET) estimate overall directionality of enrichment. Co-reporting bias is calculated as the percentage of enriched features co-published more often than expected by chance, using the mean Jaccard Index (JI) distribution for a matched number of features.

(F) Stacked bar plot compares maturation strategies revealing significant context-dependent effects on maturation properties.

(G) Table summarizing representative Spearman correlations among numerical protocol features, showing expected associations (full data in Table S4).

(H) Spearman correlation plots of cardiac cell ratios in 3D cultures show opposing relationships with sarcomere length and calcium relaxation time as maturity indicators.

Figure 2. *CMPortal* website interface and experimental use cases.

(A) Database Viewer allows users to explore 322 hPSC-CM protocols across 483 protocol features, and to install the filtered database.

(B) Pie charts of the database show media and substrate usage of metabolic maturation protocols.

(C) Example UMAPs show segregation of studies by the protocol feature categories. Cell line ancestry shows distinct trends of protocol variable and analysis method usage. Common media, like most features, show no clear trends.

(D) Variable Search enables protocols retrieval by any desired and/or significantly enriched features by target parameters.

(E) Protocol Benchmarking module enables users to upload and compare protocols against the database using experimental data and reference ranges estimated with the feature enrichments.

(F) Radar plot generated by *CMPortal* for comparing maturation outcomes between Protocol 80 and 250 which are amongst the most cited studies in the database. The plot shows protocol 80 being associated to properties of higher transcriptional and contractile maturation, but lower electrophysiological and calcium handling maturation. Hallow datapoints are estimated with enrichments when no experimental data is available.

(G) Enrichment Browser retrieves enrichments across the 117 target parameters.

(H) Spearman correlation of pairwise, co-reported values of contractility and sarcomere length in 322 protocols. No significance was found.

(I) *CMPortal*-informed experiment designed to compare the impact of media composition on contractility.

(J) Representative traces and bar plot of impedance-based contractile measurements in hPSC-CM across time show higher peak amplitude for DMEM+FA and RPMI+B27 cells.

(K) Bulk RNA-seq analysis of pathway-specific gene panels indicates metabolic reprogramming towards higher maturity in hPSC-CMs cultured in DMEM+FA compared to RPMI+B27.

(L) Representative alpha-actinin staining (green), DAPI (blue) in fixed hPSC-CM confirms independence of contractility from sarcomere length. Bar charts values are mean ± standard error whiskers with student's *t*-test significance (G, I). *P<0.05, **P<0.01,***P<0.001, ****P<0.0001, ns: non-significant. Study Char. : Study Characteristic; Meas. Endp.: Measured Endpoint; Sarco. Length: Sarcomere Length.

References

1. Murry, C. E. & Keller, G. Differentiation of Embryonic Stem Cells to Clinically Relevant Populations: Lessons from Embryonic Development. *Cell* **132**, 661–680 (2008).

2. Yang, X., Pabon, L. & Murry, C. E. Engineering Adolescence: Maturation of Human Pluripotent Stem Cell–Derived Cardiomyocytes. *Circulation Research* **114**, 511–523 (2014).

3. Ewoldt, J. K. *et al.* Induced pluripotent stem cell-derived cardiomyocyte in vitro models: benchmarking progress and ongoing challenges. *Nat Methods* **22**, 24–40 (2025).

4. Ewoldt, J. *et al.* Induced pluripotent stem cell-derived cardiomyocyte in vitro models: tissue fabrication protocols, assessment methods, and quantitative maturation metrics for benchmarking progress. 192537 bytes Dryad https://doi.org/10.5061/DRYAD.KSN02V7BH (2024).

5. Lyra-Leite, D. M. *et al.* A review of protocols for human iPSC culture, cardiac differentiation, subtype-specification, maturation, and direct reprogramming. *STAR Protocols* **3**, 101560 (2022).

6. Lee, S. *et al.* Contractile force generation by 3D hiPSC-derived cardiac tissues is enhanced by rapid establishment of cellular interconnection in matrix with muscle-mimicking stiffness. *Biomaterials* **131**, 111–120 (2017).

7. Tiburcy, M. *et al.* Defined Engineered Human Myocardium With Advanced Maturation for Applications in Heart Failure Modeling and Repair. *Circulation* **135**, 1832–1847 (2017).

8. Fetterman, K. A. *et al.* Independent compartmentalization of functional, metabolic, and transcriptional maturation of hiPSC-derived cardiomyocytes. *Cell Reports* **43**, (2024).

A ① Database Viewer

Search protocols most relevant to a target parameter

Explore 483 protocol features across five categories

	Protocol Variable	Analysis Method	...	Cell Profile	...	Study Characteristic	...	Measured Endpoint
	Maturation Strategy	Gene Analysis	...	hiPSC Cell Line	...	Disease Modelling	...	Sarco. Length
DOI 1		PCR	...		...		...	Q1
DOI 2	Metabolic Maturation		...	BJ1	...	Ischaemia	...	N/A
DOI 3	Metabolic Maturation		...	iCell	...	Ischaemia	...	Q2
DOI 4		PCR	...		...		...	Q1
...			...		...		...	
DOI N		PCR	...	BJ1, iCell	...		...	Q5

Review all 322 studies

Sort or filter protocols by features of interest

Numerical measures are binned into quantiles

B Understand media composition of selected studies

Base media
- DMEM (no glucose)
- RPMI+B27+Insulin
- RPMI+B27
- DMEM (low glucose)
- StemPro-34
- RPMI
- DMEM (high glucose)
- BPEL medium
- RPMI without Glucose
- RPMI GlutaMAX pluse Gem21
- α-MEM GlutaMAX
- Others

- 322 hiPSC-CM Studies
- Metabolic Maturation

Metabolic substrates
- Fatty Acid and Lipids
- Sugar and Carbohydrates

C Assess trends by protocol components with UMAP visualisation

| Protocol Variable | Analysis Method | Cell Profile | Study Characteristic | Measured Endpoint |

- Asian Ancestry
- Caucasian Ancestry

- RPMI-1640 (30)
- DMEM (21)
- Rare (4)
- F12 (7)
- Commercial Kit (5)

D ② Variable Search

Toggle features of interest with ease on CMPortal

Search by Matching... ▼
- Target's Enrichments ▶
- Protocol Features

Search target: Ischaemia Modelling ▼

Filter results by: IWP2 ▼

Feature categories containing enrichments for the search target

	Score	Target	Filter(s)	Other Features	Enrichment found in protocol:				
		Ischaemia Modelling	IWP2		Protocol Variable	Analysis Method	Cell Profile	Study Char.	Meas. Endp.
DOI 1	95	True	✓	...			✓	✓	
...	92	True	✓		✓				
...	89	Not Reported	✓		✓		✓		✓
...	...	...	...	...	...	...	...	...	...
DOI n	0	False	✓		✓	✓	✓		✓

Search engine can be toggled, searching by enrichments to identify relevant protocols designs regardless of their intended application and reported values

E ③ Protocol Benchmarking

Compare custom protocol designs to existing protocols

• • • • Empty data row

PDF User input using PDF template

A • C D Filled protocol details comparable to database

Maturity indicator comparison:

Contractile Force	Cell Size	**Beat Rate**
... Q1	Q1	Q1
... Q2	Q2	**Q2**
... Q3	Q3	Q3
... Q4	Q4	Q4
... Q5	Q5	Q5

Suggested indicator for optimal maturity

Reference value range for user's protocol design

F

Radar chart axes:
- Sarcomere Length
- Cell Area
- T-tubule Structure
- Contractile Force
- Contractile Stress
- Contraction Upstroke Velocity
- Calcium Flux Amplitude
- Time to Calcium Flux Peak
- Time from Calcium Peak to Relaxation
- Conduction Velocity from Calcium Imaging
- Action Potential Conduction Velocity
- Action Potential Amplitude
- Resting Membrane Potential
- Beat Rate
- Max Capture Rate of Paced CMs
- MYH7% (MYH6)
- MYL2% (MYL7)
- TNNI3% (TNNI1)

Legend
- Protocol 250
- Protocol 80
- Predicted Data
- Experimental Data (Within Range)
- Experimental Data (Below Worst Bound)

Maturity Indicator Categories
- Morphology
- Contractile Function
- Calcium Handling
- Electrophysiology
- Rhythmicity
- Transcriptomic Markers

G ④ Enrichment Browser

Query CMPortal to inform testable hypotheses for experimental considerations and designs

Select target... ▼
- Maturation Indicators ▶
  - Contractile Force
  - Ca Flux Amplitud...
  - MYH7 Percentage
  - Action Potential ...
- Physiological Descri...
- Cell Sex and Ancestry
- Disease Modelling
- Pharmacological Test

Query Outcomes:

Associated with BEST (Q1) maturation:
- hiPSC Backbone Media - DMEM
- Cell Line - BJ1
- ...

Associated with WORST (Q5) maturation:
- Maturation Media - RPMI-1640
- Maturation Strat. - Cell Alignment
- ...

H

ρ = -0.23
p = 0.41
n=15

Contractile Force (mN) vs Sarcomere Length (μm)

I

Day	-1	0	1	2	3	4	5	6	7	8
		Contractile Impedance (Ω)							Gene Expression	
Media	Replating Media	**RPMI+B27** and **DMEM+FA** Maturation Media Comparison								

Base Media	RPMI-1640	DMEM no glucose/10 mM HEPES
Supplements	2% B27 with insulin	2mM L-carnitine / 1mM nonessential amino acids
		5mM creatine / 1× insulin-transferrinselenium
		5mM taurine / 1× linoleic-oleic acid

J

RPMI+B27
DMEM+FA
Contractility (Ω) vs Time (s)

% Against RPMI+B27 — RPMI+B27 vs DMEM+FA **

K z-score 2 to -2

FAO: CPT1B, MLYCD, ACACB, ATP5F1A, PRKAA1, PPARGC1A, ACAA1, ACADSB, ACAT1, HADHB, LPIN1, PPARA, DGAT2

Glycolysis: HK2, GPI, PFKL, ALDOA, TPI1, GAPDH, PGK1, PGM2, ENO1, PKM, SLC2A1, LDHA

RPMI+B27 / DMEM+FA

L

DAPI α-actinin

RPMI+B27 DMEM+FA

50μm

Sarcomere Length (μm) — RPMI+B27 vs DMEM+FA ns