Planar cell polarity coordination in a cnidarian embryo provides clues to animal body axis evolution

Julie Uveira[1], Antoine Donati[2,3], Marvin Léria[1,4], Marion Lechable[1,5], François Lahaye[1], Christine Vesque[2], Evelyn Houliston[1] and Tsuyoshi Momose[1,6*]

Sorbonne Université, CNRS, Laboratoire de Biologie du Développement de Villefranche-sur-mer (LBDV- UMR7009), 181 Chemin du Lazaret, 06230, Villefranche-sur-Mer France
Sorbonne Université, CNRS UMR7622, INSERM U1156, Institut de Biologie Paris Seine (IBPS)-Developmental Biology Unit, 7-9 quai Saint Bernard, 75005, Paris, France
Present Address: Department of Cell and Developmental Biology, University of California San Diego, 92093 La Jolla, California, USA
Present Address: Aix Marseille Université, CNRS, IBDM-UMR7288, Turing Centre for Living Systems, Marseille, France
Present Address: Institute of Zoology, University of Innsbruck, 6020 Innsbruck, Austria
Lead Contact
* Correpondance: tsuyoshi.momose@imev-mer.fr

Summary

Body axis specification is a crucial event in animal embryogenesis and was an essential evolutionary innovation for founding the animal kingdom. It involves two distinct components that coordinate to establish the spatial organisation of the embryo: initiation of cascades of regionalised gene expression and orientation of morphogenetic processes such as body elongation. Intense interest in the first component has revealed Wnt/β-catenin signalling as ancestrally responsible for initiating regional gene expression, but the evolutionary origin of oriented morphogenesis has received little attention. Here, by addressing the cell and morphological basis of body axis development in embryos of the cnidarian *Clytia hemisphaerica,* we have uncovered a simple and likely ancestral coordination

mechanism between Wnt/β-catenin signalling and directed morphogenesis. We show that the ligand Wnt3, known to initiate oral gene expression via localised Wnt/β-catenin pathway activation, also has a key β-catenin-independent role in globally orienting planar cell polarity (PCP) to direct morphogenesis along the oral-aboral axis. This PCP orientation occurs in two distinct steps: local orientation by Wnt3 and global propagation by conserved core PCP protein interactions along the body axis. From these findings we propose novel scenarios for PCP-driven symmetry-breaking underlying the emergence of the animal body plan.

Abbreviations

PCP: planar cell polarity. EG: early gastrula, LG: late gastrula, MO: Morpholino, OA: Oral-Aboral

Keywords

Planar cell polarity, PCP, Wnt, Body Axis, Morphogenesis, Evolution, Cnidaria, Cilia

Introduction

The earliest animals are thought to have been multicellular assemblages such as the "choanoblastaea", which evolved from a unicellular ancestor (Brunet and King, 2017; Maldonado, 2004; Nielsen, 2008; Sebé-Pedrós et al., 2017). Subsequent evolutionary events that introduced a morphologically distinct primary body axis, i.e. breaking of tissue polarity symmetry, remain largely unaddressed (Fig.1A). In the animal lineage, Wnt/β-catenin-mediated specification of the primary body axis was likely an early innovation since it is found today across many species of both bilaterians (all of the animals' groups showing bilateral symmetry) and cnidarians (their sister group, comprising corals, sea anemones, jellyfish etc) (Holstein et al., 2011; Loh et al., 2016; Petersen and Reddien, 2009). Current evidence is insufficient to assess whether this innovation predated the branching of the more distantly

related ctenophore and sponge clades (Jager et al., 2013; Reid et al., 2018). Importantly, however, transcriptional activation by β-catenin and its partner TCF at one pole cannot alone explain the morphological component of axial tissue patterning. Rather, polarity information capable of directing axis development can be distributed globally across the whole embryo, as clearly illustrated by classic experimental manipulations in the hydrozoan *Clytia gregarium*, which led to the concept of global polarity (Freeman, 1981).

Freeman's global polarity concept retrospectively has much in common with planar cell polarity (PCP), the mechanism coordinating the cell polarity in the plane of epithelial tissues, later identified and unravelled genetically in *Drosophila* and vertebrates. PCP involves interactions between neighbouring cells, with each cell in a sheet, thus carrying polarity information. It is mediated by interactions of core PCP proteins: the Wnt receptor Frizzled (Fz), intracellular signal transducer Dishevelled (Dsh), transmembrane protein Strabismus/Van Gogh (Stbm), LIM-domain protein Prickle (Pk) and G-protein coupled receptor cadherin Flamingo (Fmi), to ensure tissue polarity and enable morphogenesis (Butler and Wallingford, 2017; Gray et al., 2011; May-Simera and Kelley, 2012; Seifert and Mlodzik, 2007; Vladar et al., 2009). A conserved feature of PCP is the polarised localisation of the Stbm-Pk-Fmi complex on one side of the apical cortex of each cell and Fz-Dsh-Fmi complex on the other side. These two complexes interact between adjacent cells to coordinate PCP orientation of neighbouring cells. The PCP proteins are conserved across bilaterians and cnidarians, as well as in some sponge groups (Hale and Strutt, 2015; Lapébie et al., 2011; Schenkelaars et al., 2016). Fz and Dsh are both a core PCP component and the receptor for Wnt ligands in initiating Wnt/β-catenin signalling, so Wnt, Fz and Dsh are potentially key elements in coupling the pathways.

Two crucial issues both for the developing embryo, and for the earliest animal ancestors, are i) to orient PCP and thus morphogenesis consistently and globally across the whole embryo, and ii) to coordinate PCP orientation with regionalised cell fate specification initiated by Wnt/β-catenin signalling. These issues have not been addressed in the context of the primary body axis specification, while current examples of PCP orienting mechanisms from other tissues and animals show considerable variability (Butler and Wallingford, 2017).

We address here the coupling of Wnt signalling and PCP during body axis specification in a cnidarian in order to shed light on the roles of PCP in the evolutionary emergence of the metazoan body axis, exploiting the jellyfish model species *Clytia hemisphaerica* (Houliston et al., 2022). Previously, we showed that maternal mRNAs for the ligand CheWnt3 (orthologue of bilaterian Wnt3, hereafter called Wnt3) and its receptors are localised with respect to the animal-vegetal polarity of the unfertilised *Clytia* egg (Fig. 1B), and activate Wnt/β-catenin signalling in a future oral (posterior) domain of the developing embryo to control gene expression along the developing primary body axis (oral-aboral: OA) (Momose et al., 2008; Momose and Houliston, 2007). In parallel, conserved core PCP proteins, including Stbm orthologue CheStbm (Stbm), are responsible for establishing tissue polarity (PCP) in the epidermis to support directed larval swimming via cilia beating along the oral-aboral axis, and are also required for elongation along the primary body axis during gastrulation (Momose et al., 2012). Stbm protein is localised on the aboral side of each cell (Momose et al., 2012) as expected from *Drosophila* and vertebrate PCP deployment mechanisms.

In the current work, we explore the hypothesis that Wnt3, as well as activating Wnt/β-catenin signalling (Momose et al., 2008), provides the initial cue to direct

global orientation of PCP and thereby define the *Clytia* embryonic body axis both molecularly and morphologically. A first hint of this from our previous work was that the characteristic and complete absence of embryonic axis elongation observed following Wnt3 knockdown by Morpholino (Wnt3-MO) injection could be restored by injection of synthetic Wnt3 mRNA into single blastomeres during early development (Momose et al., 2008). Importantly, Wnt3 is the sole Wnt mRNA detected before the early gastrula (EG) stage, the period when axial morphology and epidermal PCP are established, with other Wnt genes being expressed downstream of Wnt3 or at subsequent stages of larval development (Lapébie et al., 2014; Momose et al., 2008) (Fig.1B Fig. S1). This greatly facilitates the assessment of Wnt ligand function in directing PCP, in contrast to traditional model species where multiple Wnt ligands have parallel roles during embryogenesis. The studies reported here reveal how local Wnt3 cues orient the global axial PCP and PCP-driven oriented morphogenesis in the *Clytia* embryo. Two discrete steps define the PCP orientation: a local cueing step with orally localised Wnt3 signal and a global propagation and coordination step mediated by core PCP proteins. Interestingly, while the role of Wnt3 is the instructive cue for PCP orientation and thus morphological axis development during *Clytia* normal development, other cues, including mechanical strains, can orient PCP. A similar variety of PCP orientation cues has been reported in *Drosophila* and vertebrates. Our findings allow us to propose a 'PCP-first' model for body axis symmetry-breaking in the metazoan ancestors, in which the self-organising activity of the PCP pathway allowed body axis innovation in early animal ancestors.

Results

A local Wnt3 cue orients global PCP

We first analysed in detail PCP in Wnt3-depleted embryos at the EG stage, the earliest stage that shows morphological OA polarity (Fig.1B, S1). PCP manifests as structural and positional asymmetries of the ciliary basal bodies, which are consistently positioned towards the oral side of the apical cortex. As previously (Momose et al., 2012), we quantified epidermal PCP orientation by assessing the basal body positioning (Fig.1D). In unmanipulated or control-MO-injected embryos, cells showed coordinated PCP, with 85% of cells orienting within ±45° (s.d. 33 and 34° respectively) of the average orientation. In embryos injected with Morpholino antisense oligonucleotides (Wnt3-MO) to block Wnt3 translation, PCP was oriented nearly completely randomly (Fig.1E, F) (s.d. 84°, comparable to 95° for Stbm-MO embryos (Momose et al., 2012)), and individual cells were less polarised (Fig.1G), reflecting the complete lack of axial morphology. The apical cell boundary was wrinkled in Wnt3-MO-injected cells (Fig. 1E), as observed following Fz1-MO, Dsh-MO or Stbm-MO-injections (Momose et al., 2012), indicative of markedly reduced cortical tension. These analyses demonstrate that Wnt3 depletion causes PCP defects equivalent to the knockdown of the core PCP proteins (Momose et al., 2012). This supports the hypothesis that Wnt3 provides a local cue to orient global PCP, which is responsible for the development of cilia polarity and axially elongated morphogenesis in *Clytia*.

To test this hypothesis, we injected Wnt3 mRNA and fluorescently labelled Dextran into one blastomere at the four-cell stage in Wnt3-MO embryos, which we call the "Wnt3-rescue" experiment (Fig.1H, J). At the EG stage, we quantified PCP in a broad area (246 x 246 µm) and represented it by bars with a circular colour code

(Fig.1I). Control mock injections did not affect the PCP defects caused by Wnt3-MO. In contrast, coordinated PCP was restored by Wnt3 rescue in a non-autonomous manner (Fig. 1J). Restored ciliary alignment was observed at least 200 µm away from the Wnt3 mRNA-positive area. The Wnt3 source likely influences PCP at even further distances as a virtually complete morphological axis was restored by the late gastrula stage, with the Wnt3-mRNA lineage forming a new oral pole (Fig.1K) as shown previously (Momose et al., 2008). Locally expressed Wnt3 signal is thus necessary and sufficient to orient the PCP along the whole body axis.

Rapid global PCP development precedes basal body alignment

To understand the initial process by which PCP becomes coordinated by Wnt3, we examined stages preceding EG, again using basal body localisation as the read-out. We reasoned that if PCP propagation occurs through successive cell interactions, cell alignment might spread as a wave across the embryo. PCP was detectable from the mid-blastula stage (8-9 hpf Fig. 2A-D), slightly after the onset of the zygotic Wnt3 expression (Momose et al., 2008) and epithelialisation of the blastula cells (Kraus et al., 2020). We did not, however, detect any wavefront of local cell alignment; rather, the PCP gradually increased in coherence across the embryo over time (Fig.2A, B). This was confirmed by live imaging, injecting synthetic mRNAs in the egg to visualise cell contours with PH-Venus and basal bodies with Poc1-mCherry (Fig. 2E, Movie S1). The movies further revealed active mitosis in the blastodermal cells. A weak bias in cleavage orientation was observed along the OA axis (Fig. 2F). During mitosis, duplicated centrioles from the ciliary basal bodies are deployed in the mitotic spindles (Fig. 2E, Movie S1) such that basal body positioning becomes scrambled immediately after mitosis. Subsequently, the displaced basal bodies migrate

unidirectionally toward the oral side of each cell, where the cilium is re-established, a clear indication that PCP was already in place across the tissue (Fig 2E, G, H). Together these observations indicate that PCP is established at the mid-blastula stage, presumably via canonical localisation of core PCP proteins to opposite cortices of each cell through continuous Fz-Stbm interaction. The protein partitioning then manifests as asymmetric basal body positioning. Localised PCP proteins may attract basal bodies to the oral cortex of the cell as described in ascidian embryos (Negishi et al., 2016). During subsequent stages, PCP is either maintained (Bellaiche et al., 2004) in each cell during mitosis or rapidly restored through interactions with neighbouring cells (Devenport et al., 2011).

Wnt3 orients PCP in a β-catenin-independent manner.

We previously showed that local Wnt3 expression is necessary and sufficient to activate the Wnt/β-catenin pathway in the oral pole of the *Clytia* embryo (Momose et al., 2008). We next examined whether the action of Wnt3 for PCP orientation and morphogenesis is mediated by the Wnt/β-catenin pathway. First, we showed that PCP orientation by Wnt3 is independent of β-catenin: ectopic Wnt3 could orient PCP in the Wnt3-rescue experiment even in the presence of a dominant-negative form of CheTCF (dnTCF) (Fig.3A), which blocks downstream transcription of the β-catenin pathway. Gastrulation was completely prevented, confirming that the Wnt/β-catenin pathway was inactive (Fig. 3C, D, Fig. S2). Furthermore, body axis elongation was partially restored at the early gastrula stage in the absence of an active β-catenin pathway (Fig. 3C), although they did not develop pointed morphology of the oral pole nor elongate further (Fig. 3D, Fig.S2). These results support a model in which axial elongation requires both PCP coordination and gastrulation (migration and

intercalation of ingressed endodermal cells), as suggested previously (Kraus et al., 2020). Conversely, we found that constitutively active β-catenin (CA-β-cat) mRNA was not able to restore PCP coordination nor elongation of Wnt3-depleted embryos (Fig.3B) but did restore β-catenin-dependent gastrulation (Fig. S2). Taken together, these experiments demonstrate that global PCP orientation along the *Clytia* OA axis by Wnt3 is not mediated by the Wnt/β-catenin pathway. Wnt3 thus has dual independent roles: PCP orientation and Wnt/β-catenin activation.

Global PCP is established in two distinct steps.

We next tested the contribution of the core PCP proteins CheStbm and CheFmi (hereby called Stbm and Fmi) to linking PCP orientation to the local Wnt3 source. As expected, locally confined expression of Wnt3 mRNA did not restore global PCP coordination in Stbm-MO or Fmi-MO embryos (Fig.3E, F). Importantly, however, polarity was restored in the area immediately adjacent to the Wnt3 mRNA-positive cell patch. The effect covered a distance of a few cells, potentially derived from cells directly in touch with the Wnt source at earlier stages. This observation raised the possibility that global PCP orientation in *Clytia* embryos is established in two steps: Stbm/Fmi-independent PCP orientation by local Wnt3 at the oral end, followed by global PCP propagation through the intercellular interaction of core PCP proteins (two-step model), as predicted by the domineering non-autonomy property of mosaic PCP mutations (Amonlirdviman et al., 2005). Alternatively, Wnt3 may diffuse from the oral Wnt3-expressing cells (Fig.1A) to form a long-range gradient along the OA-axis and orient the PCP (Wnt-gradient model). Wnt activity gradients have been shown to control or orient PCP in mice (Gao et al., 2011; Minegishi et al., 2017).

To distinguish these two models, we created mosaic embryos with a small patch of core-PCP negative cells within Wnt3-rescued embryos.

To achieve this, we transplanted single 32- or 64-cell stage blastomeres from Stbm-MO/Wnt3-MO or Fmi-MO/Wnt3-MO injected embryos to Wnt3-rescued embryos with Dextran lineage markers (Fig. 4A-E). The two-step model predicts that the PCP orientation would make a detour around the Stbm-MO/Fmi-MO mosaic area; in contrast, under the Wnt-gradient model, PCP would be affected only in the mosaic area. We found that in the Stbm-MO or Fmi-MO mosaic Wnt3-rescued embryos, PCP was correctly oriented or slightly disturbed within the Stbm-MO patch while often de-polarised in the Fmi-MO patch. Outside the Sbtm-MO or Fmi-MO patches, PCP orientation was inverted on the aboral side of the patches unless these were in close proximity to the Wnt3-mRNA patches. Stbm- or Fmi-defected cells thus exhibited domineering non-autonomy effects. This is consistent with the intercellular Fz-Fmi and Sbm-Fmi interaction model shown in *Drosophila* and vertebrates and the localisation of Stbm protein on the aboral side of each epidermal cell in *Clytia* (Momose et al., 2012). PCP was oriented normally in the cells on the oral or lateral sides of the patches (Fig. 4B, D). PCP orientation from oral to aboral detouring around the Stbm-negative patches was similarly observed when mosaics were generated by injection of Stbm-MO into a single blastomere in a Wnt3-rescue embryo at the 16-cell stage (Fig.4F, G). These observations strongly support the two-step model in which Wnt3 acts locally to orient PCP, and PCP coordination is then transmitted from oral (i.e. Wnt3-positive lineage) to aboral through core PCP interactions. They are not compatible with the alternative model that a long-range Wnt3-gradient orients PCP.

Mechanical cues can orient PCP.

Transplantation of Wnt3-MO blastomeres into Wnt3-MO host embryos at the 32- or 64-cell stage did not cause any PCP defects in most cases. However, in a few cases (N=3 out of 8), reorientation of PCP toward the transplanted donor lineage was observed (Fig.5A-E). In these cases, the donor cells were incompletely incorporated into the host ectodermal epithelium (Fig. 5D), such that the donor cells remained as a protrusion with a "rosette" (radial) arrangement. In these embryos, host and donor-derived epithelial cells acquired an elongated morphology around the circumference of the smooth host-donor boundary (Fig.5B). Although the grafted tissue influenced the orientation of PCP in the surrounding host cells, the direction was opposite to that driven by Wnt3-expressing sources. The rosette-like transplants thus operate as artificial "aboral" cues, in contrast with Wnt3 sources, which provide an oral cue (Fig. 5C). Unlike Stbm-MO or Fmi-MO mosaic embryos, cell polarity was continuous across the host-transplant border in these Wnt3-MO to Wnt3-MO transplantations (Fig.5A), suggesting the core PCP protein interaction was not disturbed at the boundary and thus not likely due to the nonautonomous effect of PCP defects. Epithelial 'scarring' can thus act to orient PCP, perhaps via discontinuities in cell adhesion or mechanical strain. PCP re-orientation by mechanical strain has been reported in wing PCP repatterning in *Drosophila* and for the orientation of *Xenopus* epidermal motile cilia (Aigouy et al., 2010; Chien et al., 2022; Guirao et al., 2010; Mitchell et al., 2007; Olguin et al., 2011). We also performed transplantations into Wnt3-MO-only hosts but did not obtain any such rosettes, so we could not determine whether Wnt has a permissive role for PCP in this context. Nevertheless, we can conclude that any local PCP asymmetry can potentially act as a PCP orientation cue

via core-PCP proteins, and thus again is consistent with the two-step orientation model.

To explore further whether mechanical strain can orient PCP, we disrupted the morphology of the embryo by expressing dominant negative and constitutively active forms of *Clytia* RhoABC, the single cnidarian orthologue of the vertebrate Rho GTPases including RhoA. Late gastrula embryos expressing dominant negative RhoABC showed defects in axial elongation, consistent with a role in morphogenesis. Conversely, embryos injected with constitutively active RhoABC showed a scrambled axial morphology with ectopic protrusions induced along the major axis (Fig.5F). The induced protrusions appeared prior to gastrulation. PCP orientation was locally coordinated along the micro-axes of each protrusion so that they became extra aboral poles with respect to PCP (Fig.5G, H), while PCP orientation along the major body axis was globally maintained. Enhanced cortical actin fibres were arranged around the circumference of the induced protrusion (Fig. 5I). This method to provoke mechanical deformation of the embryo thus generated protrusions that mimicked the morphological extension and circumferential cell arrangement seen in the transplantation-induced rosettes, with in both cases PCP orientation aligning along the experimentally formed ectopic mini-axes.

Discussion

In this study, we have shown that during embryogenesis in the cnidarian *Clytia,* the ligand Wnt3, known to initiate oral gene expression via Wnt/β-catenin pathway activation, also provides β-catenin-independent cues to orient PCP globally in the embryo and thereby direct morphogenesis along the oral-aboral axis. PCP

orientation occurs in two distinct steps: local orientation by Wnt3 and global

propagation via conserved core PCP machinery. These findings demonstrate deep

conservation of PCP orientation and tissue morphogenesis mechanisms in

eumetazoans, and allow us to novel scenarios for PCP-driven axis symmetry-

breaking underlying the emergence of the animal body plan.

A single Wnt ligand coordinates molecular and morphological axis

determination

Wnt/β-catenin signalling (Holstein et al., 2011; Loh et al., 2016; Petersen and

Reddien, 2009) and PCP pathway components (Gray et al., 2011; Kumburegama et

al., 2011; Momose et al., 2012; Wallingford, 2012; Zallen, 2007) are involved

respectively in embryonic body axis specification and morphogenesis across many

metazoan species, but how these aspects are linked has been complicated to

address. For instance, in vertebrates, Wnt5a and Wnt11 play crucial roles during

body axis morphogenesis, (Heisenberg et al., 2000; Kilian et al., 2003; Tada and

Smith, 2000; Wallingford and Harland, 2001) but it is not straightforward to

demonstrate whether they provide positional cues to orient PCP along the body axis,

while they do in other contexts (eg. orientation of ciliated node cells and muscle

fibre) (Gros et al., 2009; Minegishi et al., 2017). During *Clytia* embryogenesis, a

single Wnt ligand directs both molecular and morphological aspects of axis formation.

Wnt3 translated from maternally localised mRNA controls axial gene expression

along the body axis via β-catenin regulation (Momose et al., 2008). This dual role of

the Wnt3 elegantly ensures the coherence of gene expression along the developing

OA axis with morphogenetic aspects such as embryo elongation during gastrulation

and beating of ectodermal cilia for directed locomotion. Maternal XWnt11 in *Xenopus*

activates the Wnt/β-catenin pathway (Tao et al., 2005), and it is also implicated in the β-catenin-independent role to control convergent extension (Tada and Smith, 2000). The dual roles of a Wnt ligand for PCP and Wnt/tβ-catenin pathway may be a common strategy in vertebrates and cnidarians.

Two-step PCP orientation, initiated by CheWnt3 and propagated by core PCP interactions.

The experiments presented here reveal two distinct steps of Wnt3-driven PCP and axial morphology in *Clytia* embryos (Fig.6A): A Wnt3-dependent local PCP orientation (cueing) step is followed by a propagation step mediated by core PCP proteins. This resembles the "domineering non-autonomy" phenomenon described in *Drosophila* mosaic mutants for core PCP genes (Amonlirdviman et al., 2005). How Wnt3 locally orients PCP in the cueing step in *Clytia* embryos remains to be understood. A likely candidate is CheFz1 (a single orthologue to vertebrate Fzd1/2/7/3/6), whose predicted localisation is on the oral side of each cell, opposite to the aboral accumulation of Stbm. An attractive hypothesis is that Wnt3 from the oral pole anchors CheFz1 at the oral side of the cells adjacent to the Wnt3 source in a mechanism similar to zebrafish Wnt11 that locally accumulates Fz7 in the plasma membrane (Witzel et al., 2006). A further implication of the two-step model and the domineering non-autonomy character of PCP proteins is that PCP orientation may act as a morphogen-independent body axis maintenance mechanism in certain regeneration contexts in cnidarians (Freeman, 1981; Livshits et al., 2017; Sinigaglia et al., 2020). Established local PCP coordination may be propagated to restore the body axis in regeneration. In this context, it would be interesting to investigate the mechanism by which core PCP modulates axial gene expression. In *Clytia*, PCP

defects resulting from Stbm knockdown led to changes in gene expression at the early gastrula stage, notably affecting cells in the oral pole that ingress during gastrulation (Lapébie et al., 2014).

Conservation of the PCP orientation and morphogenesis mechanisms in metazoans.

In the *Drosophila* wing disc, local membrane-tethered Wg/Wnt4 in the wing margin can globally orient the PCP (Wu et al., 2013; Yu et al., 2020). In vertebrate tissues, Wnt ligands either act as global cues for PCP orientation (Chu and Sokol, 2016; Gros et al., 2009; Minegishi et al., 2017), or triggers involved in the body elongation (Heisenberg et al., 2000; Kilian et al., 2003; Tada and Smith, 2000; Wallingford and Harland, 2001). The role of Wnt for PCP orientation is thus common across cnidarians and bilaterians. The Wnt subtypes employed in PCP orientation are, however, not conserved. The Wnt gene family gained its complexity prior to the bilaterian-cnidarian bifurcation (Kusserow et al., 2005). Then, different Wnt subtypes were likely co-opted in different taxa and developmental contexts. In zebrafish, PCP orientation by wnt11 and core PCP regulation are distinctive in mechanosensory hair cell orientation (Acedo et al., 2019). The modulable two-step mechanism of Wnt-driven PCP orientation is likely common in metazoans.

Other cues than Wnt also function as PCP orientation cues in the animal kingdom. In the developing *Drosophila* eye and abdomen, another signalling module comprising the atypical cadherins Fat(Ft) and Dachsous(Ds) and kinase Four-jointed(Fj) regulate the localisation of core PCP complexes via gradients of Ds and Fj (Strutt and Strutt, 2021; Yang et al., 2002). Cnidarians have conserved Ft and Ds ortholog genes. In hydra Fat-like protein (HyFat1) is implicated in epithelial cell alignment and adhesion (Brooun et al., 2020). In *Clytia,* inhibition of Fat-like (CheFat1) and Ds

(CheDs) disrupted global PCP coordination, and the Ds disruption was restored by Wnt3 expression (Fig.S3), suggesting that these molecules participate in global PCP orientation in a manner yet to be studied. Besides molecular signalling cues, mechanical constraints or fluid flow are also known to direct PCP (Aigouy et al., 2010; Chien et al., 2022; Guirao et al., 2010; Mitchell et al., 2007; Olguin et al., 2011) and can potentially act as the PCP symmetry-breaking factor. The availability of alternative PCP orientation cues might have allowed evolutionary flexibility to link PCP to available structures, such as the cell polarity scaffold of eggs. While tissue polarity orientation by PCP is strongly conserved, the cellular mechanisms underlying the morphogenetic movement are highly variable. Oriented cell division or convergent extension are likely major driving forces of blastula elongation prior to gastrulation in *Podocoryne carnea* (Momose and Schmid, 2006), another hydrozoan jellyfish. In contrast, in *Clytia*, elongation likely involves PCP-dependent intercalation of both ectoderm (Byrum, 2001) and ingressing endoderm (Sande et al., 2020) cells during gastrulation. In *Nematostella*, hydraulic pressure created by body wall muscle, organised along the body axis, drives the elongation (Stokkermans et al., 2022) (Fig. S4)

Body axis symmetry-breaking in the evolutionary history of metazoa

The dual role of Wnt3 and the two-step PCP orientation mechanism revealed here are particularly intriguing in the context of understanding the emergence of animal body axes during evolution. The metazoan ancestor is considered to have taken a multicellular state such as a "choanoblastaea" from a unicellular ancestor (Arendt et al., 2015; Nielsen, 2008) (Fig. 6B). Then metazoans acquired a body axis and germ layers and formed a hypothetical ancestral state such as Haeckel's "gastraera"

(Haeckel, 1873) or Hyman's "planuloid-acoeloid" (Hyman, 1951). Early multicellular ancestors thus acquired morphological axis patterning and coupled it with regional gene expression. It is tempting to speculate that the β-catenin/TCF-dependent local gene expression and morphogenetic patterning coordinated by core PCP proteins were independently acquired and coupled by co-opting Wnt ligands and signal transducers from one pathway to the other. For example, Wnt, Fz and Dsh might have first been employed in the Wnt/β-catenin pathway, then recruited to a primordial PCP regulatory system (Fig. 6B). Genome analyses indicate that PCP effectors and a part of core PCP components such as Cdc42, RhoA, ROCK, Prickle or Inversin are evolutionarily older than metazoans (Lapébie et al., 2011) (Fig. S5, Table S1). Wnt-Fz-Dsh would then have been recruited to create morphogenesis machinery directed by PCP based on the existing local gene expression by Wnt/β-catenin signalling. Alternatively, core PCP interactions and Wnt-driven PCP orientation may be more ancestral with Wnt-Fz-Dsh used in PCP orientation prior to its involvement in "canonical" Wnt signalling, acquired later though recruitment of the cell adhesion molecule β-catenin and its interaction with the DNA-binding TCF. Comparisons of the genomic repertoires for these components between species from across the metazoan phylogeny do not allow us to conclude which modules were acquired first (Fig.S5, S6, Table S1). In the outgroups of the cnidaria-bilateria clade, a full set of core PCP proteins are present in a sponge *Oscarella carmera* (Lapébie et al., 2011; Nichols et al., 2006; Schenkelaars et al., 2016), and full Wnt/β-catenin pathways were identified in both Porifera and Ctenophora, while multiple cases of loss of Stbm/Fmi occurred in Porifera and potentially in Ctenophora (Fig.S5, Table S1).

PCP is a fundamental character in multicellular tissue organisation seen wide across metazoans (Fig. S5, S6) and may have been particularly important in the metazoan ancestor with lower anatomical complexity. It is known that a rotational polarity of the flagellar basal body is coordinated across the spheroid of green alga *Volvox carteri*, which thus independently acquired PCP-like character (Hoops, 1993) (Fig.S6), suggesting that a PCP equivalent evolved independently in *Volvox*. PCP-driven axis evolution is an attractive scenario and can explain how an initial "choanoblastaea" multicellular ancestor acquired coordinated cilia/flagella flow. For example, PCP regulation might initially have improved filter-feeding efficiency in a choanoblastaea ancestor (Cavalier-Smith, 2017; Nielsen, 2008). A comparative cell biology, or "Cellular Evo-Devo" approach, of the PCP/tissue polarities in diverse metazoans, in particular "basal metazoans" including Cnidaria, Ctenophora and Porifera, will be the key to addressing how symmetry-breaking to create a body axis arose during evolution.

Acknowledgements

This work was supported by ANR DiploDevo (ANR-09-BLAN-0236-01), iMMEJ (ANR-17-CE13-0016-01) and the CNRS INSB Diversity of Biological Mechanisms program. We thank the "Service Aquariologie" of the CRB platform (IMEV, FR3761) supported by EMBRC-France and ANR funding (ANR-10-INBS-02) for access to facilities and resources and the PIM platform for imaging. We thank Brandon Weissbourd for the critical reading of the manuscript.

Author Contributions

TM designed the project. TM and EH wrote the manuscript. TM, AD, JU and MLé performed the experiments. MLe, FL prepared *Clytia* animal setup. FL prepared

plasmid vectors. TM, JU and MLé performed image data analysis. TM, EH and CV

acquired funding.

Declaration of Interests

The authors declare no competing interests.

Materials and Methods
Animal culture and embryo preparation.
Animal culture methods are described in (Lechable et al., 2020). *Clytia*

hemisphaerica Laboratory strains were provided by the Service Aquariologie of CRB

(IMEV-FR3761, Sorbonne Université/CNRS contact: Axel Duchene

axel.duchene@imev-mer.fr) as an EMBRC-Fr service (https://www.embrc-france.fr/).

Daily spawning was induced by light with a 24-hour day-night cycle. Eggs and sperm

were separately collected from jellyfish and transferred into 10 cm glass dishes two

hours after the light stimulation. Eggs were fertilised less than one hour after the

spawning. To acquire EG stages, embryos were incubated at 16°C for 18 hours,

which corresponds to the 12~13 hpf stage at the standard culture temperature

(18°C).

Microinjection

Morpholinos and mRNA were microinjected into unfertilised eggs or blastomeres at

4-cell ~ 16-cell stages using an Eppendorf Femtojet microinjector with

microcapillaries pulled with Narishige PN-30 needle puller. The tip of the needle was

broken by touching it to a glass surface scratched by a diamond pen to open the

needle tip to give an outer diameter of 2 to 4 μm. Eggs were aligned in a

microinjection chamber made by casting and polymerising PDMS in a negative

mould printed with a Form3 3D printer. 2~3% of the egg/cell volume of solution was

injected by inserting a microcapillary needle with constant back pressure (Femtojet

compensation pressure). It typically takes 0.5~1 seconds for each egg at 5~10 kPa

when the needle has an appropriate opening size. The volume of injection for each

injection was visually controlled under an Olympus SZX6 stereomicroscope with SZ-

ILLT oblique illumination. 3~5 mg/ml lysine fixable 10 kDa Dextran labelled with

Alexa647 or Cascade Blue were co-injected with Wnt3 mRNA and Stbm-MO/Fmi-

MO for the lineage tracking in the Wnt3-Rescue experiment and PCP mosaic

experiments.

Plasmid construction
Mutant and fluorescent protein templates were constructed in pCX3 vector, which

contains short 5' and 3'UTR sequences from CheStbm mRNA. Wildtype CheWnt3

with 3 mispaired nucleotide residues at the Morpholino target and CheTCF lacking

amino acid residues 2-27 (dnTCF) as a dominant negative form (lacking β-catenin

binding site) of human TCF-4 (Korinek et al., 1997) were amplified by PCR and

cloned into the vector. A constitutively active form of Che-β-catenin (CA-β-cat) was

created by introducing S97A T101A S105A mutations corresponding to the

constitutively active mice β-catenin mutant lacking a GSK3-β phosphorylation target

(Baba et al., 2006). Constitutively active forms (CA1: G14V, CA2: Q63L) that gave

identical phenotypes and a dominant form (DN: T19N) of CheRhoABC were

designed based on the identical vertebrate RhoA mutants with the known

phenotypes. These point mutations were introduced into wildtype cDNA clones in

pCX3 vector using the QuickChange site-directed mutagenesis kit (Stratagene).

Poc1-mCherry and PH-domain-Venus fusions were constructed in a transgenic

pMiniTol2-ACT2 vector (Weissbourd et al., 2021) using NEBuilder Assembly kit from

individually amplified PCR products or synthetic genes. mRNA was synthesised in

vitro using mMessage mMachine T3 kit and poly(A)-tailing kit (Thermo Fisher) from

PCR products amplified from the plasmids.

Immunostaining

Immunofluorescent visualisation of PCP by gamma-tubulin antibody and phalloidin

staining was described previously (Momose et al., 2012). List of the antibodies are in

the Supplementary method. In brief, the embryos were fixed with 4%

Paraformaldehyde in 0.1 M Hepes, 50 mM EGTA, 80 mM Maltose, 10 mM MgSO$_4$,

0.1% Triton X-100 and washed in PBST (1x PBS with 0.1%Triton X-100) then

incubated in antibody solution in 1% BSA fraction V in PBST, followed by three times

of washes for at least 5 minutes in PBST. After removing the detergent by washing in

PBS, actin was stained with fluorescently labelled phalloidin in PBS (0.13 µM). After

staining, embryos were cleared and mounted in Citiflour AF-1 diluted with an equal

volume of PBS and scanned using Leica SP5 or Stellaris confocal microscopes with

63x objectives. The surface of the ciliated epidermis was Z-scanned (up to 10 µm)

in the entire 246 x 246 µm field of view and z-stacked to visualise cell contour with

apical cortical actin and basal body, typically at 0.081 µm resolution in XY-axis and

0.5~1 µm resolution in Z-axis.

Image analysis

The x-y coordinates of the apical surface centroid (x_c, y_c) of each cell and basal body

of its cilium (x_b, y_b) are identified from the z-stacked confocal images. Detailed

procedures and ImageJ scripts are described in the supplementary method. The

polarity angle θ for each cell was calculated as

$$\theta = atan2(y_c - y_b, x_c - x_b)$$

The mean orientation (arrow in the radar plot) and circular standard deviation (circ.

s.d.) are calculated (Berens, 2009). A Matlab script was used to summarise the PCP

orientation and statistics (Supplementary Method). The morphology of embryos in

three dimensions (Figure. 5D, H) is reconstructed from a series of cortical actin

confocal images by building a surface of the volume containing the cortical actin

signals using Imaris software (Oxford Instruments).

Figure legends

Figure 1 Local Wnt3 globally orients the body axis tissue polarity and

morphogenesis

(A) A key unanswered question of animal evolution is the origin of body axis

patterning mechanisms, allowing initial multicellular assemblages to position

specialised cell types (red) and coordinate cell polarity (represented in this diagram

by oriented cilla and elongated shape) (B) Key stages of the early morphogenesis of

Clytia embryos in relation to Wnt3 mRNA localisation (red). At the early gastrula

(EG) stage, ectodermal cilia have formed, PCP has developed along the future oral-

aboral axis, and the pointed morphology of the animal pole morphology can first be

distinguished. Most axial elongation takes place during gastrulation. See also

Supplementary Figure 1. (C) Schematic representation of planar cell polarity in a

single epidermal cell at the EG stage. Basal bodies are positioned on the oral side at

the apical cortex (translational polarity). (D) Confocal image of the apical surface of a

single epidermal cell stained for actin (cyan, phalloidin labelling) and γ-tubulin

immunostaining (magenta), and a representation of the PCP vector, defined by the

basal body position and its mirror image across the apical surface centroid as the

initial and terminal points, respectively. (E) Local PCP coordination in Control-MO

and Wnt3-MO-injected embryos at the EG stage. bar= 10 μm. Top) Confocal images

as in D. Bottom) PCP vector representation of the cells in the images. Dots represent

cells with no clear polarisation or mitotic cells. Apical cell contours are wavy in Wnt3-

MO embryos. (F) Distribution of cell orientation in individual scans (lines) and their

average (histogram) for non-injected (373 cells, 7 images), control-MO injected (304 cells from 3 images) and Wnt3-MO-injected (225 cells from 6 images). The average orientation is defined as the origin (0°) for each scan. Horizontal bar: standard deviation. G) The PCP polarisation index (degree of the polarisation of cells) by the length of the PCP vector, normalised by the cell size. See supplementary method for the definition (**** p< 0.0001, Mann–Whitney U test). (H) Experimental procedures of the Wnt3-rescue experiment. Wnt3MO is injected into unfertilised eggs, then rescue mRNA solution with fluorescently labelled 10 kDa dextran is injected into a 4-cell stage blastomere. (I) Circular colour-code representation of the PCP. (J) Wide range PCP coordination observed in Wnt3-rescue experiments. Left: Mock rescue control by water injection (N=9). Right: Rescue by Wnt3 mRNA injection (N=27). Each set consists of three illustrations. top: colour-wheel representation of the PCP. Cells derived from the Wnt3 or water-injected blastomere are at the top-right of each image, indicated with grey shading. bar= 50 μm. bottom-left: confocal thumbnail image. cyan: actin; yellow: Dextran-Alexa647(injected blastomere derived cells). bottom-right: radar plot of the cell orientation. The arrows in the radar show the circular mean of the PCP orientation within the single scan. circ s.d.: circular standard deviation. (K) Axial morphology of normal, Wnt3-MO injected and Wnt3 rescued embryos at late gastrula (LG) stage. bar=50 μm. Elongated body axis morphology was restored so that cells with injected Wnt3 mRNA were located at the induced oral (posterior). See also Fig. 3C.

Figure 2 Wnt3-driven PCP orientation precedes visible cell alignment.

(A, B) PCP orientation in Wnt3 rescue embryo at the mid-blastula stage (A: 8 hpf, N=10) and late-blastula stage (B: 9 hpf, B, N=10). Graphical PCP representations and radar histogram plots are as in Fig. 1. (C, D) Radar plot of PCP in distal (C) and

proximal (D) areas of the region shown in (B). (E) Basal body displacements after cell divisions. Bar=10 µm. Cell membranes labelled by PH-Venus (green) and basal bodies with Poc1-mCherry (magenta). Arrows indicate basal bodies positioned on opposite sides of cleavage planes located after being engaged to the spindles. Aborally displaced basal bodies migrate toward the oral side. The average PCP defined the oral-aboral axis. (F) Distribution of the angles between the cleavage orientation axis to the OA axis defined by the PCP orientation (0-90°, n=131). Cleavages along the OA axis (0-45°) are favoured (* p<0.05 chi-square test). (G) Directional migration of the basal body toward the oral cell boundary after the cell cleavage. Each line indicates the distance of a single basal body to the oral edge of the cell after the last cleavage. Basal bodies migrate toward the oral cell boundary, suggesting PCP has been established by the time PCP becomes structurally manifest. (H) Subset of (G) where the initial distance to the oral edge is less than four micrometres after the cell division.

Figure 3 PCP orientation by Wnt3 is Wnt/β-catenin-independent, and its propagation across the body axis requires Stbm and Fmi.

(A) Wnt3-rescue with an additional injection of a dominant-negative form of TCF (dnTCF) into the egg, N=10. (B) Wnt3-rescue by the constitutively active form of β-catenin (CA-β-cat) instead of Wnt3 mRNA, N=7. (C) Onset of axial morphology in EG embryos. Localised Wnt3 is necessary and sufficient for the onset of the elongation toward its source. Green: actin (phalloidin), magenta: nuclei (To-Pro3) and blue: Wnt3-mRNA/Dextran injected cell lineage. (D) Roles of Wnt3 and β-catenin in the axial elongation in LG embryos. The elongation was measured by the elongation index (L/W): the primary axis length (L) divided by its perpendicular length (W), **** p< 0.0001, Mann–Whitney U test). Both local Wnt3 expression and β-

catenin/TCF-dependent gastrulation were required for axial elongation. See supplementary Figure 3. (E, F) Wnt3-rescue experiments with additional injections of (E) Stbm-MO (N=23) and (F) Fmi-MO (N=6). Radar plots of the PCP orientation show the manually defined proximal area (Proximal: between the dotted line and the Wnt3-positive region) and distal area (Distal: opposite area to the Wnt3-positive region across the dotted line). PCP is correctly polarised without Stbm or Fmi close to the Wnt3-injected cell clone but not in the distant area.

Figure 4. PCP behaviour in Stbm/Fmi-mosaic/ Wnt3-rescue experiments supports the two-step model for global PCP orientation

(A-D) Wnt3-rescued early gastrula embryos, including Stbm(−) or Fmi(−) mosaic patches made by blastomere transplantation. (A, B) blastomeres from Wnt3-MO/Stbm-MO double-injected embryos were transplanted as the doner into Wnt3-rescued host embryos at the 64-cell stage. They happened to be incorporated in close/proximal (A: 658 cells, N=4) or far/distal (B: 757 cells, N=4) positions with respect to the Wnt3 source, respectively. (C, D) The same experiment was conducted with Wnt3-MO/Fmi-MO-injected blastomeres as the donor, incorporated in proximal (C: 984 cells, N=3) and distal (D: 649 cells, N=3) positions, respectively. (E) Schematic drawing of the experimental procedure of the transplantation. (F) Wnt3-rescued early gastrula embryos with Stbm(−) mosaic patch generated by additional injection of Stbm-MO at the 16-cell stage (N=1). (G) Schematic drawing of the experimental procedures for (F). The flow of the PCP orientation is indicated by grey arrows in the bottom right drawings in (A-D) and right in (G). (H) Graphical summary of Stbm-MO and Fmi-MO mosaic experiments. The graphical representation and thumbnail confocal images indicate Wnt3-positive lineages and

Stbm-MO or Fmi-MO-injected lineages in yellow and blue. All vector representations are as in Figure 1.

Figure 5. Mechanical strains can reorient PCP.

(A-E) PCP orientation induced by transplantation from a Wnt3-MO donor into the Wnt3-MO background of a Wnt3-rescue host. A) Incomplete incorporation of host and donor cells into a smooth epidermis (N=3/8) causing long-range PCP orientation, as if the donor acts as an aboral cue, in addition to the Wnt3 oral cue. (A) PCP representation by colour-coded bars. See also the legends for Figure.1 (B) Confocal image of the apical cortex of the epidermis showing the rosette structure caused by incomplete incorporation of the donor cells. Both host and donor cells are elongated around the graft boundary. (C) Graphical summary of the PCP orientation: Wnt3-mRNA lineage in Wnt3-rescue host in yellow, Wnt3-MO graft in blue. (D) 3D reconstruction of the rosette structure from the confocal images by contouring cortical actin signals. (E) Schematic representation of the transplant experiment. (F) 3D representation of the late gastrula morphologies induced by mRNA injection of dominant negative form (RhoABC-DN) and constitutively active forms (RhoABC-CA2) of the PCP effector RhoABC, which caused reduced elongation and extra protrusions, respectively. The overall oral-aboral axis remained distinguishable based on morphology and gastrulation. (G-I) PCP is coordinated consistently with respect to the induced protrusions caused by RhoABC-CA2. (G) Colour-wheel representation of PCP in the RhoABC-CA2 injected early gastrula embryos; the presumed oral pole is to the top-right. (H) 3D reconstruction of the tissue morphology for the same specimen, with PCP flow represented by arrows. (I) Confocal image of the apical cortex of the epidermis at the induced protrusion, corresponding

rectangles in G and H. Cortical actin fibres are circumferentially organised around the protrusion. Bars are 50 μm except for 10 μm in B and I. See Figure 1 for the circular colour-code representation of PCP orientation. cyan: actin; magenta: γ-tubulin; blue: donor lineage marker Dextran for B and I

Figure 6. Models for body axis symmetry-breaking during *Clytia* embryogenesis and PCP-driven axis innovation scenarios in metazoan evolution

(A) Two-step PCP axis orientation by local Wnt3. Local gradients of Wnt3 at the oral end of the *Clytia* embryo orient epidermal cell polarity near the Wnt3-positive area. This process does not require Wnt/β-catenin signalling. Local PCP orientation propagates to the aboral side through the action of core PCP proteins between neighbouring cells. Wnt3 expression is likely maintained by Wnt3/β-catenin feedback regulation. (B) Two possible metazoan body axis evolution scenarios from a choanoblastaea, a multicellular and non-polarised metazoan common ancestor. In the "Wnt/β-catenin-first" scenario (bottom), the choanoblastaea first acquires a mechanism to activate the Wnt/β-catenin pathway locally. Stably locally maintained Wnt ligand expression is then recruited by preexisting but non-directed core Fz/PCP interactions to orient PCP and morphogenesis. In the "PCP-first scenario" (top), the self-organising capacity of core PCP protein interactions creates tissue polarity without any specific cue to specify the direction. The advantage of coordinated polarity of cilia movement for filter-feeding, swimming or morphology could have favoured directed symmetry breaking using reliable cues. These could have initially been provided by environmental or mechanical constraints before the recruitment of

localised Wnt binding to Fz to assure PCP orientation, and then the emergence of downstream beta-catenin signalling.

References

Acedo JN, Voas MG, Alexander R, Woolley T, Unruh JR, Li H, Moens C, Piotrowski T. 2019. PCP and Wnt pathway components act in parallel during zebrafish mechanosensory hair cell orientation. *Nature communications* **10**:3993–17. doi:10.1038/s41467-019-12005-y

Aigouy B, Farhadifar R, Staple DB, Sagner A, Röper J-C, Jülicher F, Eaton S. 2010. Cell flow reorients the axis of planar polarity in the wing epithelium of Drosophila. *Cell* **142**:773–786. doi:10.1016/j.cell.2010.07.042

Amonlirdviman K, Khare NA, Tree DRP, Chen W-S, Axelrod JD, Tomlin CJ. 2005. Mathematical modeling of planar cell polarity to understand domineering nonautonomy. *Science* **307**:423–426. doi:10.1126/science.1105471

Arendt D, Benito-Gutierrez E, Brunet T, Marlow H. 2015. Gastric pouches and the mucociliary sole: setting the stage for nervous system evolution. *Philosophical Transactions Royal Soc B Biological Sci* **370**:20150286. doi:10.1098/rstb.2015.0286

Baba Y, Yokota T, Spits H, Garrett KP, Hayashi S-I, Kincade PW. 2006. Constitutively active beta-catenin promotes expansion of multipotent hematopoietic progenitors in culture. *Journal of immunology (Baltimore, Md : 1950)* **177**:2294–2303. doi:10.4049/jimmunol.177.4.2294

Bellaiche Y, Beaudoin-Massiani O, Stuttem I, Schweisguth F. 2004. The planar cell polarity protein Strabismus promotes Pins anterior localization during asymmetric

division of sensory organ precursor cells in Drosophila. *Development (Cambridge, England)* **131**:469–478.

Berens P. 2009. CircStat: A MATLAB Toolbox for Circular Statistics. *Journal of Statistical Software* **31**.

Brooun M, Klimovich A, Bashkurov M, Pearson BJ, Steele RE, McNeill H. 2020. Ancestral roles of atypical cadherins in planar cell polarity. *Proc National Acad Sci* **117**:19310–19320. doi:10.1073/pnas.1917570117

Brunet T, King N. 2017. The Origin of Animal Multicellularity and Cell Differentiation. *Developmental cell* **43**:124–140. doi:10.1016/j.devcel.2017.09.016

Butler MT, Wallingford JB. 2017. Planar cell polarity in development and disease. *Nature Reviews Molecular Cell Biology* **18**:375–388. doi:10.1038/nrm.2017.11

Byrum CA. 2001. An analysis of hydrozoan gastrulation by unipolar ingression. *Developmental biology* **240**:627–640. doi:10.1006/dbio.2001.0484

Cavalier-Smith T. 2017. Origin of animal multicellularity: precursors, causes, consequences-the choanoflagellate/sponge transition, neurogenesis and the Cambrian explosion. *Philosophical transactions of the Royal Society of London Series B, Biological sciences* **372**:20150476. doi:10.1098/rstb.2015.0476

Chien Y-H, Kim S, Kintner C. 2022. Mechanical strain breaks planar symmetry in embryonic epithelia via polarized microtubules. *Cells Dev* **170**:203791. doi:10.1016/j.cdev.2022.203791

Chu C-W, Sokol SY. 2016. Wnt proteins can direct planar cell polarity in vertebrate ectoderm. *eLife* **5**:940. doi:10.7554/elife.16463

Devenport D, Oristian D, Heller E, Fuchs E. 2011. Mitotic internalization of planar cell polarity proteins preserves tissue polarity. *Nature cell biology* **13**:893–902. doi:10.1038/ncb2284

Freeman G. 1981. The role of Polarity in the Development of the Hydrozoan Planula Larva. *Wilhelm Roux's Archives* **190**:168–184.

Gao B, Song H, Bishop K, Elliot G, Garrett L, English MA, Andre P, Robinson J, Sood R, Minami Y, Economides AN, Yang Y. 2011. Wnt signaling gradients establish planar cell polarity by inducing Vangl2 phosphorylation through Ror2. *Developmental cell* **20**:163–176. doi:10.1016/j.devcel.2011.01.001

Gray RS, Roszko I, Solnica-Krezel L. 2011. Planar cell polarity: coordinating morphogenetic cell behaviors with embryonic polarity. *Developmental cell* **21**:120–133. doi:10.1016/j.devcel.2011.06.011

Gros J, Serralbo O, Marcelle C. 2009. WNT11 acts as a directional cue to organize the elongation of early muscle fibres. *Nature* **457**:589–593. doi:10.1038/nature07564

Guirao B, Meunier A, Mortaud S, Aguilar A, Corsi J-M, Strehl L, Hirota Y, Desoeuvre A, Boutin C, Han Y-G, Mirzadeh Z, Cremer H, Montcouquiol M, Sawamoto K, Spassky N. 2010. Coupling between hydrodynamic forces and planar cell polarity orients mammalian motile cilia. *Nature cell biology* **12**:341–350. doi:10.1038/ncb2040

Haeckel E. 1873. Die Gastraea-Theorie, die phylogenetische Classification des Thierreichs und die Homologie der Keimblätter. Jena.

Hale R, Strutt D. 2015. Conservation of Planar Polarity Pathway Function Across the Animal Kingdom. *Annu Rev Genet* **49**:529–551. doi:10.1146/annurev-genet-112414-055224

Heisenberg CP, Tada M, Rauch GJ, Saúde L, Concha ML, Geisler R, Stemple DL, Smith JC, Wilson SW. 2000. Silberblick/Wnt11 mediates convergent extension

movements during zebrafish gastrulation. *Nature* **405**:76–81. doi:10.1038/35011068

Holstein TW, Watanabe H, Özbek S. 2011. Signaling pathways and axis formation in the lower metazoa. *Current topics in developmental biology* **97**:137–177. doi:10.1016/b978-0-12-385975-4.00012-7

Hoops H. 1993. Flagellar, cellular and organismal polarity in Volvox carteri. *Journal of cell science*.

Houliston E, Leclère L, Munro C, Copley RR, Momose T. 2022. Past, present and future of Clytia hemisphaerica as a laboratory jellyfish. *Curr Top Dev Biol* **147**:121–151. doi:10.1016/bs.ctdb.2021.12.014

Hyman LH. 1951. The invertebrates□: Platyhelminthes and Rhynchocoela, the acoelomate Bilateria.. New York□: McGraw-Hill Book Company Inc.

Jager M, Dayraud C, Mialot A, Quéinnec E, Guyader H le, Manuel M. 2013. Evidence for Involvement of Wnt Signalling in Body Polarities, Cell Proliferation, and the Neuro-Sensory System in an Adult Ctenophore. *Plos One* **8**:e84363. doi:10.1371/journal.pone.0084363

Kilian B, Mansukoski H, Barbosa FC, Ulrich F, Tada M, Heisenberg C-P. 2003. The role of Ppt/Wnt5 in regulating cell shape and movement during zebrafish gastrulation. *Mechanisms of development* **120**:467–476. doi:10.1016/s0925-4773(03)00004-2

Korinek V, Barker N, Morin PJ, Wichen D van, Weger R de, Kinzler KW, Vogelstein B, Clevers H. 1997. Constitutive transcriptional activation by a beta-catenin-Tcf complex in APC-/- colon carcinoma. *Science* **275**:1784–1787. doi:10.1126/science.275.5307.1784

Kraus Y, Chevalier S, Houliston E. 2020. Cell shape changes during larval body plan development in Clytia hemisphaerica. *Developmental biology* **468**:59–79. doi:10.1016/j.ydbio.2020.09.013

Kumburegama S, Wijesena N, Xu R, Wikramanayake AH. 2011. Strabismus-mediated primary archenteron invagination is uncoupled from Wnt/β-catenin-dependent endoderm cell fate specification in Nematostella vectensis (Anthozoa, Cnidaria): Implications for the evolution of gastrulation. *EvoDevo* **2**:2. doi:10.1186/2041-9139-2-2

Kusserow A, Pang K, Sturm C, Hrouda M, Lentfer J, Schmidt HA, Technau U, Haeseler A von, Hobmayer B, Martindale MQ, Holstein TW. 2005. Unexpected complexity of the Wnt gene family in a sea anemone. *Nature* **433**:156–160. doi:10.1038/nature03158

Lapébie P, Borchiellini C, Houliston E. 2011. Dissecting the PCP pathway: one or more pathways?: Does a separate Wnt-Fz-Rho pathway drive morphogenesis? *Bioessays* **33**:759–768. doi:10.1002/bies.201100023

Lapébie P, Ruggiero A, Barreau C, Chevalier S, Chang P, Dru P, Houliston E, Momose T. 2014. Differential Responses to Wnt and PCP Disruption Predict Expression and Developmental Function of Conserved and Novel Genes in a Cnidarian. *PLoS genetics* **10**:e1004590. doi:10.1371/journal.pgen.1004590

Lechable M, Jan A, Duchene A, Uveira J, Weissbourd B, Gissat L, Collet S, Gilletta L, Chevalier S, Leclère L, Peron S, Barreau C, Lasbleiz R, Houliston E, Momose T. 2020. An improved whole life cycle culture protocol for the hydrozoan genetic model Clytia hemisphaerica. *Biology open* **9**:bio051268. doi:10.1242/bio.051268

Livshits A, Shani-Zerbib L, Maroudas-Sacks Y, Braun E, Keren K. 2017. Structural

Inheritance of the Actin Cytoskeletal Organization Determines the Body Axis in

Regenerating Hydra. *Cell Rep* **18**:1410–1421. doi:10.1016/j.celrep.2017.01.036

Loh KM, Amerongen R van, Nusse R. 2016. Generating Cellular Diversity and

Spatial Form: Wnt Signaling and the Evolution of Multicellular Animals.

Developmental cell **38**:643–655. doi:10.1016/j.devcel.2016.08.011

Maldonado M. 2004. Choanoflagellates, choanocytes, and animal multicellularity.

Invertebrate Biology **123**:1–22. doi:10.1111/j.1744-7410.2004.tb00138.x

May-Simera H, Kelley MW. 2012. Planar cell polarity in the inner ear. *Current topics

in developmental biology* **101**:111–140. doi:10.1016/b978-0-12-394592-1.00006-

Minegishi K, Hashimoto M, Ajima R, Takaoka K, Shinohara K, Ikawa Y, Nishimura H,

McMahon AP, Willert K, Okada Y, Sasaki H, Shi D, Fujimori T, Ohtsuka T,

Igarashi Y, Yamaguchi TP, Shimono A, Shiratori H, Hamada H. 2017. A Wnt5

Activity Asymmetry and Intercellular Signaling via PCP Proteins Polarize Node

Cells for Left-Right Symmetry Breaking. *Developmental cell* **40**:439-452.e4.

doi:10.1016/j.devcel.2017.02.010

Mitchell B, Jacobs R, Li J, Chien S, Kintner C. 2007. A positive feedback mechanism

governs the polarity and motion of motile cilia. *Nature* **447**:97–101.

Momose T, Derelle R, Houliston E. 2008. A maternally localised Wnt ligand required

for axial patterning in the cnidarian Clytia hemisphaerica. *Development

(Cambridge, England)* **135**:2105–2113. doi:10.1242/dev.021543

Momose T, Houliston E. 2007. Two oppositely localised frizzled RNAs as axis

determinants in a cnidarian embryo. *PLoS biology* **5**:e70.

doi:10.1371/journal.pbio.0050070

Momose T, Kraus Y, Houliston E. 2012. A conserved function for Strabismus in

establishing planar cell polarity in the ciliated ectoderm during cnidarian larval

development. *Development (Cambridge, England)* **139**:4374–4382.

doi:10.1242/dev.084251

Momose T, Schmid V. 2006. Animal pole determinants define oral-aboral axis

polarity and endodermal cell-fate in hydrozoan jellyfish Podocoryne carnea.

Developmental biology **292**:371–380. doi:10.1016/j.ydbio.2006.01.012

Negishi T, Miyazaki N, Murata K, Yasuo H, Ueno N. 2016. Physical association

between a novel plasma-membrane structure and centrosome orients cell

division. *eLife* **5**:e16550. doi:10.7554/elife.16550

Nichols SA, Dirks W, Pearse JS, King N. 2006. Early evolution of animal cell

signaling and adhesion genes. *Proc National Acad Sci* **103**:12451–12456.

doi:10.1073/pnas.0604065103

Nielsen C. 2008. Six major steps in animal evolution: are we derived sponge larvae?

Evolution & development **10**:241–257. doi:10.1111/j.1525-142x.2008.00231.x

Olguin P, Glavic A, Mlodzik M. 2011. Intertissue mechanical stress affects Frizzled-

mediated planar cell polarity in the Drosophila notum epidermis. *Current

biology : CB* **21**:236–242. doi:10.1016/j.cub.2011.01.001

Petersen CP, Reddien PW. 2009. Wnt signaling and the polarity of the primary body

axis. *Cell* **139**:1056–1068. doi:10.1016/j.cell.2009.11.035

Reid PJW, Matveev E, McClymont A, Posfai D, Hill AL, Leys SP. 2018. Wnt

signaling and polarity in freshwater sponges. *BMC Evol Biol* **18**:12.

doi:10.1186/s12862-018-1118-0

Sande M van der, Kraus Y, Houliston E, Kaandorp J. 2020. A cell-based boundary

model of gastrulation by unipolar ingression in the hydrozoan cnidarian Clytia

hemisphaerica. *Developmental biology* **460**:176–186.

doi:10.1016/j.ydbio.2019.12.012

Schenkelaars Q, Fierro-Constain L, Renard E, Borchiellini C. 2016. Retracing the

path of planar cell polarity. *BMC evolutionary biology* **16**:69–13.

doi:10.1186/s12862-016-0641-0

Sebé-Pedrós A, Degnan BM, Ruiz-Trillo I. 2017. The origin of Metazoa: a unicellular

perspective. *Nat Rev Genet* **18**:498–512. doi:10.1038/nrg.2017.21

Seifert JRK, Mlodzik M. 2007. Frizzled/PCP signalling: a conserved mechanism

regulating cell polarity and directed motility. *Nat Rev Genet* **8**:126–138.

doi:10.1038/nrg2042

Sinigaglia C, Peron S, Eichelbrenner J, Chevalier S, Steger J, Barreau C, Houliston

E, Leclère L. 2020. Pattern regulation in a regenerating jellyfish. *Elife* **9**:e54868.

doi:10.7554/elife.54868

Stokkermans A, Chakrabarti A, Subramanian K, Wang L, Yin S, Moghe P,

Steenbergen P, Mönke G, Hiiragi T, Prevedel R, Mahadevan L, Ikmi A. 2022.

Muscular hydraulics drive larva-polyp morphogenesis. *Curr Biol* **32**:4707-4718.e8.

doi:10.1016/j.cub.2022.08.065

Strutt H, Strutt D. 2021. How do the Fat–Dachsous and core planar polarity

pathways act together and independently to coordinate polarized cell

behaviours? *Open Biol* **11**:200356. doi:10.1098/rsob.200356

Tada M, Smith JC. 2000. Xwnt11 is a target of Xenopus Brachyury: regulation of

gastrulation movements via Dishevelled, but not through the canonical Wnt

pathway. *Development (Cambridge, England)* **127**:2227–2238.

Tao Q, Yokota C, Puck H, Kofron M, Birsoy B, Yan D, Asashima M, Wylie CC, Lin X,

Heasman J. 2005. Maternal Wnt11 Activates the Canonical Wnt Signaling

Pathway Required for Axis Formation in Xenopus Embryos. *Cell* **120**:857–871. doi:10.1016/j.cell.2005.01.013

Vladar EK, Antic D, Axelrod JD. 2009. Planar Cell Polarity Signaling: The Developing Cell's Compass. *Cold Spring Harbor Perspectives in Biology* **1**:a002964–a002964. doi:10.1101/cshperspect.a002964

Wallingford JB. 2012. Planar cell polarity and the developmental control of cell behavior in vertebrate embryos. **28**:627–653. doi:10.1146/annurev-cellbio-092910-154208

Wallingford JB, Harland RM. 2001. Xenopus Dishevelled signaling regulates both neural and mesodermal convergent extension: parallel forces elongating the body axis. *Development (Cambridge, England)* **128**:2581–2592.

Weissbourd B, Momose T, Nair A, Kennedy A, Hunt B, Anderson DJ. 2021. A genetically tractable jellyfish model for systems and evolutionary neuroscience. *Cell* **184**:5854-5868.e20. doi:10.1016/j.cell.2021.10.021

Witzel S, Zimyanin V, Carreira-Barbosa F, Tada M, Heisenberg CP. 2006. Wnt11 controls cell contact persistence by local accumulation of Frizzled 7 at the plasma membrane. *The Journal of cell biology* **175**:791–802. doi:10.1083/jcb.200606017

Wu J, Roman A-C, Carvajal-Gonzalez JM, Mlodzik M. 2013. Wg and Wnt4 provide long-range directional input to planar cell polarity orientation in Drosophila. *Nature cell biology* **15**:1045–1055. doi:10.1038/ncb2806

Yang C, Axelrod J, Simon M. 2002. Regulation of Frizzled by fat-like cadherins during planar polarity signaling in the Drosophila compound eye. *Cell* **108**:675–688.

Yu JJS, Maugarny-Calès A, Pelletier S, Alexandre C, Bellaiche Y, Vincent J-P,

McGough IJ. 2020. Frizzled-Dependent Planar Cell Polarity without Secreted

Wnt Ligands. *Dev Cell* **54**:583-592.e5. doi:10.1016/j.devcel.2020.08.004

Zallen JA. 2007. Planar polarity and tissue morphogenesis. *Cell* **129**:1051–1063.

doi:10.1016/j.cell.2007.05.050

Figure 1

A multicellulari-sation → ? → innovation of body axis

B
0	7	8~9	11~12	18-20 (hpf)
egg	mid blastula	late blastula	early gastrula (EG)	late gastrula (LG)

Animal pole — Vegetal pole

Wnt3 mRNA

ingression

Oral — Aboral

swimming

C Aboral — Oral; Apical — Basal; Stbm; cilium; microvilli (actin); γ-Tubulin cloud; centrosome; Aboral — Oral

D γ-Tubulin/actin; basal body; centroid of apical surface

E Early Gastrula; Control-MO; Wnt3-MO; actin + γ-tubulin; 10 μm; cell orientation

F Non-injected; Control-MO; Wnt3-MO; -180 -90 0 90 180

G polarization index; Non-injected; Control-MO; Wnt3-MO; Stbm-MO; **** **** ns

H Wnt3 MO → Wnt3 RNA + Dextran-Alexa633 → EG

I

J "Control (Mock)" Wnt3-MO (egg) + Dextran-Alexa633 (4-cell); Wnt3 (−); Mock; Wnt3 (−)

"Wnt3 rescue" Wnt3-MO (egg)+ Wnt3-RNA / Dextran-Alexa633(4-cell); Wnt3 (+); Wnt3 (−)

Mock inj. side; circ. s.d. = 120.0°; Wnt3(+) side; circ. s.d. = 39.6°

K Non-injected; Wnt3-MO; Wnt3 rescue

Figure 2

A 8 hpf (mid-blastula) **B** 9 hpf (late-blastula)

Wnt3 Rescue

Wnt3 (+)

Wnt3 (−)

50 μm

Wnt3(+) side

circ. s.d. = 105.0°

Wnt3 (−)

Wnt3 (+)

(C)

(D)

Wnt3(+) side

circ. s.d. = 78.6°

C

90   0.1

180   −90

circ. s.d. = 85.7°

D

90   0.15
      0.1
      0.05

180   −90

circ. s.d. = 51.8°

F

*

0.1

0   θ: Angle between cleavage and OA axes   90°

O

θ

A

G

distance from oral border (μm)

0   30   60   90
time after last cleavage (min)

H subset of cells in G, with
       initial distance < 4 μm

distance from oral border (μm)

0   30   60   90
time after last cleavage (min)

aboral

oral

E

10 μm

T₀ =8.0 hpf   +19min   +43min   +51min   (9hpf) +59min   +66min   +73min   +83min

Figure 3

A Wnt3-rescue + dnTCF(egg)

Wnt3 (−)   Wnt3 (+)

Wnt3(+) side

circ. s.d. = 50.4°

50 µm

B Wnt3-MO (egg)
constitutively active β-catenin (4-cell)

Wnt3(+) side

CA-β-catenin

Wnt3 (−)

circ. s.d. = 117.0°

C Early gastrula (green: actin, magenta: nuclei)

Wnt3 mRNA-injected

Wnt3-MO	−	+	+	+
dnTCF	−	−	−	+
Wnt3-mRNA (4-cell)	−	−	+	+

D Elongation (Late gastrula)

L/W

4
3
2
1

****
****
****
ns
ns
****
ns

egg   4-cell

	−	Wnt3 mRNA	Wnt3MO			dnTCF	
Wnt3 mRNA	−	+	−	−	−	−	+
CA-β-catenin mRNA	−	−	L	H	−	−	−

E Wnt3-rescue + StbmMO (egg)

Wnt3 (+)
Stbm(−)

Wnt3 (−)
Stbm(−)

proximal

distal

circ. s.d. = 59.5°

circ. s.d. = 93.5°

F Wnt3-rescue + FmiMO (egg)

Wnt3 (+)
Fmi(−)

Wnt3 (−)
Fmi(−)

proxim

dist

circ. s.d. = 59.0°

circ. s.d. = 114.6°

Figure 4

Host: Wnt3-MO (egg) + Wnt3 mRNA (4-cell) Alexa 647

Donor: Wnt3-MO + Stbm-MO + Dextran Cascade Blue   Donor: Wnt3-MO + Fmi-MO + Dextran Cascade Blue

A: **Stbm-MO** patch **proximal** to Wnt3(+)

B: **Stbm-MO** patch **distal** from Wnt3(+)

C: **Fmi-MO** patch **proximal** to Wnt3(+)

D: **Fmi-MO** patch **distal** from Wnt3(+)

Figure 5

Host: Wnt3-MO (egg) + Wnt3 mRNA (4-cell)   Donner: Wnt3-MO (egg)

A Wnt3 mRNA (−)   Wnt3 mRNA (+)

B   10 μm

C

D

E egg   4-cell   32-cell   ≥64-cell

Wnt3 MO + Dextran–Alexa633

donnor

Wnt3 MO   Wnt3 RNA + Dextran-FITC

host

Wnt3 (−)
Wnt3 (+)

F Late Gastrula
Control   RhoABC DN   RhoABC CA2

Early Gastrula RhoABC CA2

G I

H I   Oral
Aboral

I   10 μm

Figure 6