Genetic diversity and demographic history of the largest remaining migratory population

of blue wildebeest (*Connochaetes taurinus taurinus*) in southern Africa

Stephanie J. Szarmach[1,2], Katherine C. Teeter[1], Jassiel M'soka[3], Egil Dröge[4,5], Hellen Ndakala[6],

Clive Chifunte[4,7], Matthew S. Becker[4,8], Alec R. Lindsay[1]

[1]Department of Biology, Northern Michigan University, Marquette, MI 49855, USA

[2]Department of Biology, Pennsylvania State University, State College, PA 16802, USA

[3]USAID Zambia, Lusaka, Zambia

[4]Zambian Carnivore Programme, Mfuwe, Eastern Province, Zambia

[5]Wildlife Conservation Research Unit, Department of Biology, The Recanati-Kaplan Centre,

Oxford University, Oxford, United Kingdom

[6]Zambia Department of National Parks and Wildlife, Liuwa Plain

[7]Zambia Department of National Parks and Wildlife, Mumbwa, Zambia

[8]Department of Ecology, Montana State University, Bozeman, MT 59735, USA

Corresponding author: Stephanie J. Szarmach, stephszarmach@gmail.com

ORCID: 0000-0003-1538-8127

ABSTRACT

The blue wildebeest (*Connochaetes taurinus taurinus*) is a keystone species in the savannahs of

southern Africa, where it maintains shortgrass plains and serves as an important prey source for

large carnivores. Despite being the second largest migratory wildebeest population, the

wildebeest of the Greater Liuwa Ecosystem (GLE) of western Zambia have remained largely

unstudied, until recently. While studies have increased understanding of recent demography,

migration, and population limiting factors, the level of genetic diversity, patterns of gene flow,

and long-term demographic history of blue wildebeest in the GLE remains unknown. Most

genetic studies of wildebeest have focused on small, heavily-managed populations, rather than

large, migratory populations of high conservation significance. We used restriction-site

associated DNA sequencing (RAD-seq) to assess genetic diversity, population structure, and

demographic history of blue wildebeest in the GLE. Using SNPs from 1,730 loci genotyped

across 75 individuals, we found moderate levels of genetic diversity in GLE blue wildebeest (H_e

$= 0.210$), no evidence of inbreeding ($F_{IS} = 0.033$), and an effective population size of about one

tenth the estimated population size. No genetic population structure was evident within the GLE.

Analyses of the site frequency spectrum found signatures of expansion during the Middle

Pleistocene followed by population decline in the Late Pleistocene and early Holocene, a pattern

previously observed in other African ungulates. These results will supplement field studies in

developing effective conservation plans for wildebeest as they face continued and increasing

threats of habitat loss, poaching, and other human impacts across their remaining range.

Keywords: Blue wildebeest, *Connochaetes taurinus*, population genetics, demographic history,

RAD-seq, Zambia

INTRODUCTION

The blue wildebeest (*Connochaetes taurinus*) is a large antelope that acts as a keystone species in the savannahs of southern and eastern Africa. Many wildebeest populations migrate seasonally in response to precipitation, availability of high-quality forage, and predation risk [1–4]. Grazing by wildebeest maintains shortgrass plains and facilitates nutrient cycling [5–7], which in turn supports communities of other grazing mammals, insects, and birds that depend on small herbaceous plants [8]. Carnivores, such as lions (*Panthera leo*), hyenas (*Crocuta crocuta*), wild dogs (*Lycaon pictus*), and cheetahs (*Acinonyx jubatus*), often depend on wildebeest as an important prey source [9–12]. Because of the significant ecological roles of wildebeest, changes in migration dynamics or population size could have dramatic impacts on other species and communities in the African grasslands [13].

In many parts of the wildebeest's range, habitat destruction and fragmentation, obstruction of migration routes, and poaching have impacted wildebeest population size and movement [14]. The conversion of grassland to agricultural fields has reduced available foraging area, contributing to population declines [15,16] and the construction of fences and roads has blocked migration routes and caused previously migratory populations to become sedentary [17,18]. Poaching for bushmeat, driven by deteriorating economic conditions in many communities in and around protected area networks, also threatens wildebeest populations [19–21]. To better understand and address the impact of these threats, there is a need to detect and evaluate their effects on populations.

In addition to experiencing recent declines, wildebeest populations may have undergone fluctuations in the more distant past due to changes in climate or human activity. Genetic analysis of other African ungulates, including African buffalo (*Syncerus caffer*) [22–24], African

elephant (*Loxodonta africana*) [25], and hippopotamus (*Hippopotamus amphibius*) [26,27],

revealed population declines during the late Pleistocene and early Holocene. These declines

coincide with increases in human population size and periods of aridity on the African continent,

during which longer dry seasons led to drought and changes in water levels and vegetation types

[28–30]. Because many wildebeest populations migrate in response to rainfall and vegetation

growth and populations have declined during recent droughts, wildebeest likely experienced

similar declines during these two epochs [4,19].

Large population declines during the Late Pleistocene and Holocene may have affected

the level of genetic diversity found in the species today, and an understanding of these changes

could help in evaluating future demographic effects of climate change. Today, the dryland

ecosystems comprising most of the remaining wildebeest range are characterized by extreme

seasonality in resource abundance and distribution, making mobility and access to these

resources key, particularly in the face of climate change [31]. Understanding how continued

habitat fragmentation, loss of migratory routes, and population declines affect genetic diversity

and the ability of wildebeest to adapt to the rapidly changing environments of the Anthropocene

is therefore of key importance.

The Greater Liuwa Ecosystem (GLE), located in western Zambia and encompassing

Liuwa Plain National Park (3,660 km^2) and the Upper West Zambezi Game Management Area

(22,156 km^2; Figure 1), supports a large, migratory blue wildebeest population. This flat, open

grassland experiences extensive seasonal flooding, which drives the wildebeest to migrate

between the drier, southern portion of the GLE, where they spend the wet season (November –

April), and the previously flooded northern areas for the dry season (May – October) [32,33].

Though this is thought to be the second largest wildebeest migration in Africa, little was known

about this population until recently, after it experienced a significant decline. Poaching, driven

by regional politics and civil war in neighboring Angola, led to severe declines of many species

in the GLE, including wildebeest, and notably led to the extirpation of lions [32,34].

Additionally, while the wildebeest are thought to have historically migrated between Zambia and

Angola, since the Angolan civil war (1975-2002) no transboundary migration has been observed

and radio-tracking has shown wildebeest movements to be largely confined to the GLE [33].

Restoration efforts in Liuwa began in 2003 with the signing of a 20-year management

agreement between the government of Zambia and African Parks Network. Ecosystem recovery

efforts have resulted in growth and stabilization of the wildebeest population, from an estimated

15,000 individuals in 2001 to 46,000 in 2013 [33–35]. Studies of movement, demography,

environmental influences, and predator-prey dynamics [11,32,33,36] have improved

understanding of wildebeest population dynamics in the GLE, but the amount of genetic

diversity present in the population and its long-term demographic history remain unknown. Past

changes in population size have important implications for conservation in the present day,

because historical bottlenecks can reduce genetic diversity and limit future adaptive potential,

and the comparison of present day to past effective population sizes can improve understanding

of current conservation risks for the population [37,38].

Despite the wildebeest being an iconic species of the African savannahs with high

conservation significance, until very recently [39], few genetic studies of wildebeest had been

published. Studies utilizing molecular markers can inform wildlife management by delineating

conservation units, revealing inbreeding, and divulging historical patterns of movement and

demographic change [40,41]. Most previous genetic studies of wildebeest focused on heavily

managed populations in South Africa, and few evaluated the potential impacts of habitat loss,

fragmentation, and population declines occurring across remaining wildebeest range. Early

studies assessing wildebeest genetic diversity used allozymes, microsatellites or mitochondrial

DNA primarily to inform management of small populations of wildebeest on game ranches in

South Africa [42–44]. Microsatellite and mtDNA studies that included samples from a greater

part of the blue wildebeest's range identified high levels of genetic divergence between

wildebeest in southern versus eastern Africa, as well as lower levels of significant differentiation

between pairs of populations within these regions [45,46].

More recently, whole-genome resequencing of blue wildebeest from much of the species'

range found significant genetic differentiation between all subspecies of blue wildebeest, as well

as between local populations of the western white bearded (*C. t. mearnsi*) subspecies in eastern

Africa and the brindled (*C. t. taurinus*) subspecies in southern Africa [39]. Importantly, Liu et al.

[39] found that wildebeest populations where migration had recently been disrupted showed

lower genetic diversity and higher levels of inbreeding compared to fully migratory populations.

No previous genetic studies have included wildebeest from the GLE, despite this being one of

the largest migratory wildebeest populations. The effects of past changes in population size, both

historical and recent, on genetic diversity in this recovering migratory population remain

unknown.

We assessed genetic diversity, population structure, and demographic history of the GLE

blue wildebeest population using a genome-wide SNP dataset generated through restriction-site

associated DNA sequencing (RAD-seq). We first evaluated the current level of genetic diversity

present in the GLE wildebeest population, estimated contemporary effective population size, and

assessed whether genetic population structure is present. We used both model-based

(STRUCTURE) and model-free (PCA) approaches to determine whether multiple populations

exhibiting significant differences in SNP allele frequencies are present within the GLE, or

whether this region is occupied by a single, panmictic population. We then inferred the past

demographic history of the GLE blue wildebeest population and tested the hypothesis that

genetic signatures of population contraction are present due to past climate change and

anthropogenic pressures. To do this, we tested demographic models using the coalescent

simulator fastsimcoal2 [47] and implemented the model-independent stairway plot method

[48,49] to estimate effective population size over time based on the site frequency spectrum.

MATERIALS AND METHODS

Sample collection and DNA extraction

As part of long-term field studies conducted with the Zambia Department of National Parks and

Wildlife (DNPW) and African Parks, we collected 97 tissue samples in the GLE between

2011−2014 from wildebeest that were anesthetized for radio collaring as part of long-term

studies, or from opportunistically sampled wildebeest carcasses detected during the course of

fieldwork. These tissue samples were transferred to ethanol for storage. All animal handling

procedures were performed following animal welfare standards and protocols required by the

Zambia Department of Veterinary and Livestock Services and the DNPW and were approved by

the Montana State University IACUC. Samples were exported with Zambia Wildlife Authority

Permit #7830 and imported with USDA Veterinary Permit #109174. We extracted genomic

DNA from the tissues using the DNeasy Blood & Tissue Kit (Qiagen, Valencia, CA, USA) and

quantified the concentration and purity of the extracted DNA using a Nanodrop 2000c. DNA

extractions with a concentration <10 ng/µL were combined with DNA from a second extraction

from the same tissue sample, if available, and concentrated using vacuum centrifugation.

RAD library preparation and sequencing

We prepared double-digest restriction site associated DNA (ddRAD) libraries following the

protocol of DaCosta and Sorenson [50]. First, ~1 µg of genomic DNA from each wildebeest was

digested using the restriction enzymes *EcoR*I-HF and *Sbf*I-HF (New England Biolabs, Ipswich,

MA, USA). Unique combinations of barcoded adapters were ligated to the digested DNA from

each individual, so that samples could later be pooled for sequencing. Restriction fragments

within a size range of 200−400 bp were selected through gel excision, and DNA was purified

using a Qiagen MinElute Gel Extraction Kit. The size-selected DNA fragments were PCR

amplified using Phusion High-Fidelity PCR Master Mix (Thermo Scientific, Waltham, MA,

USA). PCR products were purified with AMPure XP SPRI beads (Beckman Coulter, Brea, CA,

USA). The concentrations of the purified PCR products were quantified using real-time PCR,

and samples were pooled in equimolar amounts, with the exception of some low-concentration

samples. Ultimately, 83 samples were included in the multiplexed library, which was sequenced

on one lane of an Illumina HiSeq 2500 Sequencing System (Tufts University, Medford, MA,

USA), generating single-end reads of 100 bp in length.

Alignment, genotyping, and quality filtering

After sequencing, we used FastQC version 0.11.7 to assess per-base sequence quality for the

reads [51]. We used a computational pipeline developed by DaCosta and Sorenson [50] for

analyzing ddRAD-seq data to process the raw sequence reads (Python scripts available at

https://github.com/BU-RAD-seq/ddRAD-seq-Pipeline). After correcting sequencing errors in

restriction sites and barcodes, reads were assigned to individual wildebeest based on the

combination of barcodes associated with the read. The demultiplexed sequence data are available

from the NCBI Sequence Read Archive (SRA) under the following accession numbers:

BioProject ID PRJNA588629; BioSample IDs SAMN13254976−SAMN13255074. After

demultiplexing, identical sequences were collapsed and reads with low average Phred quality

scores (Q < 20) were filtered out. The condensed and filtered reads were then clustered into

putative loci across individuals based on a similarity threshold of 0.85 using the UCLUST

algorithm in the program USEARCH version 5 [52]. A representative sequence from each cluster

was aligned to a reference genome using the blastn program in BLAST version 2.8.0 [53] and the

following settings: evalue = 0.0001, word_size = 11, gapopen = 5, gapextend = 2, penalty = -3,

reward = 1, and dust = yes). Sequences were aligned to two reference genomes: version 4.0 of

the domestic sheep (*Ovis aries*) genome (2015; Assembly GCA_000298735.2), downloaded

from NCBI, and the blue wildebeest genome sequenced and assembled by Chen et al. [54],

downloaded from http://animal.nwsuaf.edu.cn/code/index.php/Ruminantia. Separate clusters that

generated BLAST hits to the same genomic location were combined into a single cluster, and

those that did not generate hits to the reference genome were retained but marked as anonymous

loci. Reads were aligned within each cluster using MUSCLE version 3.8.31 [55].

We genotyped individuals at the identified loci using DaCosta and Sorenson's

RADGenotypes.py script [50], which calls individuals as homozygous for a SNP if >93% of

reads exhibit the same allele, and as heterozygous if >29% of reads show a second allele. When a

lower proportion of reads exhibited a minor allele, the genotype was flagged as a "provisional

heterozygote" (20-29% of reads) or "ambiguous" (7-20%) and was required to pass additional

examination and quality filtering before being included in the final set of SNPs. A custom R

script utilizing the tidyverse collection of packages [56,57] was used to remove loci with

substantial amounts of missing data and putative "paralogous loci" formed by erroneous

clustering of sequences originating from repetitive elements in the genome (R script available at

https://github.com/sszarmach/NMU-wildebeest). Samples with an average sequencing depth of

<10 reads per locus and loci with <5 reads in >10% of samples were removed, since low

sequencing depth can lead to genotyping errors [58]. We filtered out loci that significantly

departed from Hardy-Weinberg equilibrium based on a chi-square goodness-of-fit test ($\alpha = 0.05$),

because such deviation can result from null alleles (heterozygote deficiency) or incorrectly

clustered paralogous loci (heterozygote excess). To eliminate putative paralogous loci, we also

removed clusters that exhibited more than two alleles, that had abnormally high read depths

(mean depth >700), that had >4 SNPs, or that generated multiple high-quality BLAST hits to the

reference genome [59,60]. For all downstream analyses except inference of demographic history

using the site frequency spectrum, a minor allele frequency filter (MAF >0.01) was applied to

remove loci containing singleton SNPs, which could be an artifact of sequencing error. These

quality filtering steps are outlined in Supplementary Table 1.

Genetic diversity

A final data set of 1,730 polymorphic loci genotyped across 75 individuals was used to calculate

summary statistics describing genetic diversity in GLE blue wildebeest. We randomly drew one

SNP from each RAD locus to avoid the inclusion of multiple SNPs that were linked due to

chromosomal proximity. We used PGDSpider version 2.1.1.5 to convert the SNP data file into

the correct format for each analysis program [61]. Allele frequencies and expected and observed

heterozygosities were calculated using the R package adegenet [62,63]. We tested SNPs for

deviation from Hardy-Weinberg equilibrium using the hw.test function in the R package pegas

(Fisher's exact test with 1,000 MCMC replicates) [64]. *P*-values from the Hardy-Weinberg tests

were corrected for multiple comparisons using the Benjamini and Hochberg method in the R

package sgof [65–67]. The inbreeding coefficient, F_{IS}, was estimated using the R package

hierfstat [68].

Estimation of effective population size

We estimated contemporary effective population size (N_e) for GLE blue wildebeest using the

bias-corrected linkage disequilibrium method [69,70] implemented in NeEstimator version 2.1

[71] assuming a random-mating model. 95% confidence intervals for N_e were constructed using

the parametric method of Waples [69] and the jackknifing method of Jones et al. [72], which

accounts for pseudoreplication resulting from physically linked loci and is recommended for data

sets including large numbers of SNPs.

Population structure

We explored whether genetic structure is present within the GLE wildebeest population using the

model-based clustering approach implemented in STRUCTURE version 2.3.4 [73–76]. We

assumed the admixture model and correlated allele frequencies [74,77]. Ten independent runs

were performed for each number of genetic clusters tested ($K = 1$ to $K = 4$), and each run

contained 10,000 burn-in steps and 100,000 MCMC steps. We used STRUCTURE

HARVESTER web version 0.6.94 [78,79] to summarize STRUCTURE results, and the best K

value was selected based on the log probability of the data given the number of clusters (ln

Pr(X|K)) and the Evanno ΔK method [80]. Visualizations of the STRUCTURE output were

created using the R package and Shiny app pophelper [81]. We also examined genetic structure

using a non-model-based approach by performing a principal components analysis (PCA) in the

R package adegenet [62].

Demographic history

Using the coalescent simulator fastsimcoal version 2.6 [47] we evaluated five models of

historical demographic change: constant population size, expansion, decline, bottleneck, and

expansion followed by decline (hereafter "expansion-decline"; Figure 2). Fastsimcoal2 simulates

the expected site frequency spectrum (SFS) for a demographic model provided by the user and

computes the composite likelihood of the SFS observed from the data given this model. We

computed the folded site frequency spectrum from the minor allele frequencies of all sites in

1,831 loci (2,946 SNPs and 172,603 monomorphic sites) that passed quality filtering without

applying the minor allele frequency filter, as excluding singletons can bias inferences drawn

from the site frequency spectrum [82].

For each model, we ran 50 replicate simulations, each with randomly selected initial

parameter values, to increase the likelihood that the global maximum solution was discovered.

Each run included 100,000 coalescent simulations and 40 cycles of Brent's ECM algorithm,

which is used to select parameters for the model that optimize the composite likelihood [47]. We

provided broad parameter search ranges of 10–100,000 gene copies (haploid N) for all N_e

estimates, except $N_{ancestral}$ in the decline and bottleneck models and N_{mid} in the expansion-decline

model, which were set to 10–1,000,000 gene copies. The search range for all event time

parameters was 10-100,000 generations. The mutation rate was set to 1.7×10^{-8} substitutions per

base pair per generation, based on the average mammalian mutation rate (2.22×10^{-9} substitutions

per base pair per year) [83] and the generation time of wildebeest (7.5 years) [84]. For each

demographic scenario, the replicate with the highest likelihood was retained as the best model.

Fastsimcoal2 handles effective population size as the number of gene copies, so N_e was

converted to the number of diploid individuals by dividing all estimated N_e parameters by two.

Time parameters were converted from number of generations to years before present by

multiplying T estimates by a generation time of 7.5 years. We calculated Akaike Information

Criterion (AIC) values [85] for the models with the highest maximum composite likelihood for

each demographic scenario and compared the demographic models using Akaike weights

(AICw) calculated using the akaike.weights function in the qpcR R package [86].

In addition to the model-based approach implemented in fastsimcoal2, we also inferred

historical changes in population size using the model-independent stairway plot method.

Stairway Plot 2 uses a flexible multi-epoch model to estimate population size over different time

intervals based on the site frequency spectrum [48,49]. The same folded site frequency spectrum,

mutation rate, and generation time used in the fastsimcoal2 analysis were used to generate the

stairway plot. Confidence intervals were created based on 200 bootstrapped site frequency

spectra drawn from the observed SFS.

RESULTS

Sequencing produced ~220 million reads that passed initial Illumina quality filtering, and

demultiplexing resulted in an average of 2.2 million reads assigned to each wildebeest. After

these reads were filtered for quality and clustered by similarity, 22,483 RAD loci were identified.

A similar number of loci aligned to each reference genome: 19,216 (85.5%) of sequences

generated at least one BLAST hit to the sheep genome, and 19,267 (85.7%) of loci aligned to the

blue wildebeest genome. Eight wildebeest samples were removed due to low sequencing depth

(<10 reads per locus), leaving 75 wildebeest that were genotyped. There were 3,417 polymorphic

loci that were genotyped in >90% of samples. Many putatively paralogous loci were removed

from the dataset, likely the result of the large proportion of repetitive sequence in bovid genomes

(~50%) [54,87]. After all quality filtering (Supplementary Table 1), the final dataset included

1,730 high-quality polymorphic RAD loci (mean length = 96 bp, mean depth = 355X) that

passed a minor allele frequency (MAF) threshold of 0.01, containing 2,921 SNPs. The number of

loci that aligned to each chromosome of the sheep genome was correlated with the length of the

chromosome (Spearman's $\rho = 0.683$, $P < 0.001$; Supplementary Figure 1), indicating that the

dataset contains loci evenly spaced throughout the genome.

We estimated genetic diversity and effective population size for GLE blue wildebeest (n

= 75) using 1,730 biallelic SNPs, with one SNP randomly drawn from each RAD locus. After

correcting for multiple comparisons, no SNPs in the final dataset significantly deviated from

Hardy-Weinberg equilibrium ($\alpha = 0.05$). The minor allele frequency per SNP ranged from 0.01

to 0.50, with a mean of 0.14 (SD = 0.13). The observed heterozygosity of the SNPs ranged from

0 to 0.597, and the average observed heterozygosity of GLE blue wildebeest was 0.204 (SD =

0.150; Figure 3). Expected heterozygosity per SNP ranged from 0.013 to 0.504, and the average

expected heterozygosity was 0.209 (SD = 0.151). The inbreeding coefficient (F_{IS}) for the

population was 0.031 (95% CI: 0.026 – 0.038). The estimated effective population size (N_e) was

5,075 (parametric 95% CI = 3,360–10,322; jackknifing 95% CI = 800–∞).

We did not detect any genetic population structure within the GLE blue wildebeest. The

most likely number of clusters identified using STRUCTURE was $K = 1$ (Supplementary Table

2), and for each K value >1 that was tested, almost all individuals were assigned membership to

multiple clusters (Figure 4a; Supplementary Figure 2). Principal component analysis also did not

show any genetic clustering of subpopulations within the GLE (Figure 4b). Rather, the GLE blue

wildebeest seem to compose a single, panmictic population.

The results of both the model-based fastsimcoal2 analysis and model-independent

stairway plot analysis suggest that the GLE blue wildebeest experienced population expansion

during the Pleistocene followed by large declines throughout the Late Pleistocene and Holocene.

Of the five demographic models simulated in fastsimcoal2 (Figure 2), the expansion-decline

model had the highest composite likelihood score and Akaike weight (w_i; Table 1). The

parameters of the highest-likelihood expansion-decline model suggest a population increase from

N_e=26,800 to N_e=92,700 occurring 635,000 years ago, followed by a population decline to

N_e=13,400 occurring 15,480 years ago (Supplementary Table 3). The stairway plot exhibited

similar changes in population size as the fastsimcoal2 expansion-decline model (Figure 5), with

an increase in population size from N_e=30,000 to N_e=90,000 from 1,000,000-500,000 years ago,

followed by a sharp decline to N_e=11,000 from 25,000-3,000 years ago.

DISCUSSION

Our results show that a single, panmictic blue wildebeest population occupies the GLE,

possessing moderate levels of genetic diversity, no evidence of inbreeding, and no genetic

population structure. Though relatively few population genomic studies of wild ungulates have

measured SNP-based heterozygosity, the level of heterozygosity seen in GLE blue wildebeest

(average per-SNP H_o = 0.204) was slightly higher than average per-SNP heterozygosity in

another African antelope, the springbok (*Antidorcas marsupialis*; H_o = 0.15) [88], and similar to

that seen in pronghorn (*Antilocapra americana*; H_o = 0.21) [89]. Heterozygosity in GLE

wildebeest was slightly lower than that found in African forest elephants (*Loxodonta cyclotis*; H_o

= 0.27) [90], although the elephant value was averaged over multiple sampling locations. One

other study has reported per-SNP heterozygosity for reduced-representation genomic data from

blue wildebeest (~20,000 SNPs generated using DArT-seq) [91]. From their dataset generated

for wildebeest from a South African private game ranch, van Deventer et al. [91] found a mean

minor allele frequency (0.19) and average observed heterozygosity (0.19) similar to those of the

GLE blue wildebeest. However, in the South African population, observed heterozygosity was

significantly lower than expected heterozygosity (0.27), potentially a sign of inbreeding that was

not seen in the GLE blue wildebeest.

While direct comparison to other previously published studies of genetic variation in blue

wildebeest is more challenging, especially those that utilized allozymes, microsatellites, and

mitochondrial DNA, the GLE blue wildebeest seem to have relatively greater genetic diversity

than some previously studied populations. Researchers found low allozyme polymorphism and

low heterozygosity in wildebeest from Tanzania and Kenya [92], and low mtDNA nucleotide

diversity [42] and allozyme heterozygosity [43] in South African wildebeest. Many of these early

studies focused on smaller, more heavily managed wildebeest populations that likely were

experiencing founder effects resulting from translocations of individuals from one reserve to

another. In contrast, wildebeest in the GLE are free-ranging and do not have a history of

translocations or intensive management. In a recent study of wildebeest populations throughout

their range using whole-genome data, Liu et al. [39] found lower genome-wide heterozygosity

estimates and more runs of homozygosity (a sign of past inbreeding) in nonmigratory wildebeest

populations compared to migratory ones. Though the GLE wildebeest numbers declined

drastically in the early 2000's and the migration no longer extends into Angola, it does not seem

that these changes have resulted in low genetic diversity in Liuwa compared to other

populations.

The GLE blue wildebeest exhibited no signs of inbreeding. Though F_{IS} was significantly

greater than zero (0.033; 95% CI: 0.026 – 0.039), indicating a slight excess of homozygosity, this

low value does not indicate extensive nonrandom mating in the population. Similar values of the

inbreeding coefficient have been seen in springbok (F_{IS} = 0.018 – 0.032) from several

populations in South Africa [88]. While the GLE blue wildebeest population experienced

significant declines in the early 2000's [34], it seems these declines were not great enough and

did not last long enough (occurring over a few generations) to lead to inbreeding and reduction

of genetic diversity in the population. We also found no evidence of population structure using

either STRUCTURE or PCA, indicating that a single, panmictic wildebeest population inhabits

the GLE. Visual observation and GPS tracking of collared wildebeest have found that the GLE

wildebeest spend the wet season in a concentrated area in the southern portion of the park [33],

and our genetic results corroborate that there is likely no spatial segregation of individuals during

breeding.

Our estimate of the current effective population size (N_e), the size of an idealized

population that would experience the same amount of genetic drift as the population being

studied [93], was 5,075 (95% CI = 3,360–10,322). Estimating a population's effective size is

important for conservation, because a population with low N_e could lose genetic diversity faster

than expected based on its census population size (N). The effective population size estimated for

wildebeest in the GLE is 11% of the total estimated population size determined using aerial

surveys in 2013 (N = 46,000) [35]—an effective to census population size ratio (N_e/N) similar to

estimates for many comparable mammals. Reviews have found the average N_e/N ratio in wildlife

to be 0.10-0.14 [94,95]. In African buffalo, another large, highly mobile ungulate with a

promiscuous mating system, researchers have estimated N_e/N ratios ranging from 0.10 to 0.20

[96]. When also considering the moderate level of heterozygosity and low inbreeding coefficient

for the Liuwa population, it seems unlikely that this estimate of current effective population size

is indicative of a loss in genetic diversity after the population decline in the early 2000's. Very

recent population declines can actually increase the N_e/N ratio, because census population size

decreases faster than the timescale required for N_e to be reduced by a loss of genetic diversity

[37,38,95]. By comparison, LaCava et al. [89] estimated a very low N_e/N ratio of 0.01-0.02 for

pronghorn (*Antilocaptra americana*), a North American ungulate that, in addition to having a

polygynous mating system, experienced precipitous declines over a century ago.

Using both model-based and model-independent methods, we detected signatures of

population expansion followed by decline in the site frequency spectrum of GLE blue

wildebeest. Both the highest-likelihood expansion-decline fastsimcoal2 model and the stairway

plot inferred a threefold increase in population size during the Middle Pleistocene, followed by

an ~87% reduction in population size during the Late Pleistocene and early Holocene. Evidence

of Pleistocene population expansions followed by Holocene declines has been found in a number

of other African mammal species using genetic data, including African buffalo [23,24],

hippopotamus [26,27], African elephant [25,97], and savannah baboon [98]. In a comparative

study of ruminant whole genomes, including 26 African species, Chen et al. [54] found using the

pairwise sequentially Markovian coalescent (PSMC) method that most of the African species

shared a similar demographic history of population expansion followed by decline in the late

Pleistocene. Similarly, Liu et al. [39] used PSMC to infer effective population size over time for

the five blue wildebeest subspecies, and the demographic history inferred for the brindled

wildebeest (which did not include any individuals from the GLE) is very similar to the results of

our stairway plot analysis in both the timing of the expansion and decline and the effective

populations sizes inferred.

Two primary explanations have been proposed to explain the population declines in

African ungulates during the late Pleistocene and early Holocene: climatic change and

anthropogenic impacts [99]. Ice core records and other paleontological data suggest that Africa

experienced large fluctuations in precipitation during the late Pleistocene and early Holocene,

with extensive periods of drought [28–30]. Such climatic changes could have a profound impact

on wildebeest, as modern-day wildebeest populations have fluctuated in response to drought [19]

and wildebeest migration is at least partially driven by changes in rainfall and forage availability

[4]. However, others have argued that climatic changes in the late Pleistocene were no more

extreme than those occurring earlier in the epoch, and thus do not alone explain population

declines and extinctions [99]. Human population growth and the advent of pastoralism during the

Neolithic revolution may have increased pressures on ungulate populations already stressed by

drought, leading to the observed declines [23,54]. Past population declines in the face of climatic

changes and intensified human activity reflect the modern-day pressures faced by blue

wildebeest, as habitat destruction, overhunting, and climate change all threaten the species today.

Our results show that—despite recent declines in population size and the disruption of

migration to Angola—the GLE blue wildebeest population does not seem to have experienced

recent reductions in genetic diversity or increased inbreeding. The ecosystem restoration efforts

implemented in the GLE likely allowed the wildebeest population to increase before the

bottleneck became severe enough to result in detectable reductions in genetic diversity.

However, wildebeest continue to face threats of habitat destruction, obstruction of migration

routes from fencing, and climate change, and continued genetic monitoring could detect whether

these impacts are influencing genetic diversity in the GLE in the future. This is especially

important given the evidence that blue wildebeest population size drastically decreased during

past periods of climate change and anthropogenic pressure in the late Pleistocene and early

Holocene.

Future sampling and genomic analysis of wildebeest from Kafue National Park in Zambia

could help to elucidate patterns of connectivity or differentiation between wildebeest in the GLE

and those in a neighboring population that is also of conservation significance. The Zambezi

River, which separates these two protected areas and currently acts as a barrier, may have

historically prevented gene flow, leading to genetic differentiation between these populations.

Analysis of four wildebeest genomes from Kafue by Liu et al. [39] suggested that the Kafue

population was genetically distinct from more southern brindled wildebeest populations, but the

relationship between the Kafue and GLE populations remains unknown. Understanding patterns

of genetic differentiation between GLE blue wildebeest and those in Kafue National Park would

have important implications for future management decisions such as translocations between

these populations. Our study provides a baseline understanding of genetic diversity and long-

term demographic history for Africa's second largest migratory blue wildebeest population,

which will assist the development of effective conservation plans for the species as wildebeest

face threats of habitat destruction and overhunting.

Acknowledgements

Our thanks to the Zambia Department of National Parks and Wildlife, African Parks Network, and the Barotse Royal Establishment for permission and collaboration in this research. We gratefully acknowledge the late Jeffrey Horn of the NMU Computer Science Department, who passed away in February 2024. Jeff provided the computing infrastructure and assistance that made the data analysis for this work possible. We appreciate and miss his enthusiasm for collaborating across disciplines. We also thank Grant Combs for computing assistance. We are very grateful to Michael Sorenson and Jeffrey DaCosta for guidance on the RAD-seq protocol and analysis pipeline. We also thank Kurt Galbreath for input and advice on the project and Catherine Sun for feedback on the manuscript.

Funding: The fieldwork and wildebeest tissue sample collection for this study was supported by the World Wildlife Fund–Netherlands and Zambia, National Science Foundation (IOS1145749), DOB Ecology, and the Lee Schink Memorial Scholarship from the Wildlife Conservation Network awarded to J. M'soka. The molecular laboratory work and genetic analyses were funded by a Northern Michigan University Northern PRIME grant awarded to A. Lindsay, K. Teeter, and J. Horn.

Ethics Statement: All animal handling procedures were performed following animal welfare standards and protocols required by the Zambia Department of Veterinary and Livestock Services and the Zambia Department of National Parks and Wildlife and were approved by the Montana State University IACUC. Samples were exported with Zambia Wildlife Authority Permit #7830 and imported with USDA Veterinary Permit #109174.

Competing Interests: The authors have no relevant financial or non-financial interests to disclose.

Author Contributions: KCT, ARL, MSB, and SJS contributed to the study conception and

design. JM and ED collected the blue wildebeest tissues. HN and CC performed project support,

oversight, and administration. RAD-seq library preparation and data analysis were performed by

SJS. The first draft of the manuscript was written by SJS and all authors commented on the

manuscript, contributed to subsequent drafts, and approved the final draft.

Data Availability: The sequence data generated for this study are available from the NCBI

Sequence Read Archive (SRA) under the following accession numbers: BioProject ID

PRJNA588629; BioSample IDs SAMN13254976−SAMN13255074

(https://www.ncbi.nlm.nih.gov/bioproject/PRJNA588629).

REFERENCES

1. Fryxell JM, Sinclair ARE. Causes and consequences of migration by large herbivores. Trends in Ecology & Evolution. 1988;3: 237–241. doi:10.1016/0169-5347(88)90166-8

2. Wilmshurst JF, Fryxell JM, Farm BP, Sinclair ARE, Henschel CP. Spatial distribution of Serengeti wildebeest in relation to resources. 1999;77: 10.

3. Boone RB, Thirgood SJ, Hopcraft JGC. Serengeti wildebeest migratory patterns modeled from rainfall and new vegetation growth. Ecology. 2006;87: 1987–1994. doi:10.2307/20069184

4. Holdo RM, Holt RD, Fryxell JM. Opposing rainfall and plant nutritional gradients best explain the wildebeest migration in the Serengeti. The American Naturalist. 2009;173: 431–445. doi:10.1086/597229

5. McNaughton SJ. Serengeti migratory wildebeest: facilitation of energy flow by grazing. Science. 1976;191: 92–94. doi:10.1126/science.191.4222.92

6. Holdo RM, Holt RD, Coughenour MB, Ritchie ME. Plant productivity and soil nitrogen as a function of grazing, migration and fire in an African savanna. Journal of Ecology. 2007;95: 115–128. doi:10.1111/j.1365-2745.2006.01192.x

7. Estes RD, Raghunathan TE, Vleck DV. The impact of horning by wildebeest on woody vegetation of the Serengeti ecosystem. The Journal of Wildlife Management. 2008;72: 1572–1578. doi:10.2193/2008-105

8. Sinclair ARE. Mammal population regulation, keystone processes and ecosystem dynamics. Philos Trans R Soc Lond B Biol Sci. 2003;358: 1729–1740. doi:10.1098/rstb.2003.1359

9. Hayward MW, Kerley GIH. Prey preferences of the lion (Panthera leo). Journal of Zoology. 2005;267: 309–322. doi:10.1017/S0952836905007508

10. Hayward MW. Prey preferences of the spotted hyaena (Crocuta crocuta) and degree of

dietary overlap with the lion (Panthera leo). Journal of Zoology. 2006;270: 606–614.

doi:10.1111/j.1469-7998.2006.00183.x

11. Creel S, Dröge E, M'soka J, Smit D, Becker M, Christianson D, et al. The relationship

between direct predation and antipredator responses: a test with multiple predators and

multiple prey. Ecology. 2017;98: 2081–2092. doi:10.1002/ecy.1885

12. Dröge E, Creel S, Becker MS, M'soka J. Spatial and temporal avoidance of risk within a

large carnivore guild. Ecology and Evolution. 2017;7: 189–199. doi:10.1002/ece3.2616

13. Lee DE, Kissui BM, Kiwango YA, Bond ML. Migratory herds of wildebeests and zebras

indirectly affect calf survival of giraffes. Ecology and Evolution. 2016;6: 8402–8411.

doi:10.1002/ece3.2561

14. Estes R. The gnu's world: Serengeti wildebeest ecology and life history. Berkeley:

University of California Press; 2014.

15. Serneels S, Lambin EF. Impact of land-use changes on the wildebeest migration in the

northern part of the Serengeti-Mara Ecosystem. Journal of Biogeography. 2001;28: 391–

407. doi:10.2307/2656203

16. Homewood K, Lambin EF, Coast E, Kariuki A, Kikula I, Kivelia J, et al. Long-term changes

in Serengeti-Mara wildebeest and land cover: pastoralism, population, or policies? PNAS.

2001;98: 12544–12549. doi:10.1073/pnas.221053998

17. Whyte CSJ, Joubert IJ. Blue wildebeest population trends in the Kruger National Park and

the effects of fencing. South African Journal of Wildlife Research - 24-month delayed open

access. 1988;18: 78–87.

18. Holdo RM, Fryxell JM, Sinclair ARE, Dobson A, Holt RD. Predicted impact of barriers to

migration on the Serengeti Wildebeest population. PLoS ONE. 2011;6: e16370.

doi:10.1371/journal.pone.0016370

19. Mduma SAR, Sinclair ARE, Hilborn R. Food regulates the Serengeti wildebeest: a 40-year

record. Journal of Animal Ecology. 1999;68: 1101–1122. doi:10.1046/j.1365-

2656.1999.00352.x

20. Thirgood S, Mosser A, Tham S, Hopcraft G, Mwangomo E, Mlengeya T, et al. Can parks

protect migratory ungulates? The case of the Serengeti wildebeest. Animal Conservation.

2004;7: 113–120. doi:10.1017/S1367943004001404

21. Harris G, Thirgood S, Hopcraft JGC, Cromsight JPGM, Berger J. Global decline in

aggregated migrations of large terrestrial mammals. Endangered Species Research. 2009;7:

55–76. doi:10.3354/esr00173

22. Heller R, Lorenzen ED, Okello JBA, Masembe C, Siegismund HR. Mid-Holocene decline in

African buffalos inferred from Bayesian coalescent-based analyses of microsatellites and

mitochondrial DNA. Molecular Ecology. 2008;17: 4845–4858. doi:10.1111/j.1365-

294X.2008.03961.x

23. Heller R, Brüniche-Olsen A, Siegismund HR. Cape buffalo mitogenomics reveals a

Holocene shift in the African human–megafauna dynamics. Molecular Ecology. 2012;21:

3947–3959. doi:10.1111/j.1365-294X.2012.05671.x

24. de Jager D, Glanzmann B, Möller M, Hoal E, van Helden P, Harper C, et al. High diversity,

inbreeding and a dynamic Pleistocene demographic history revealed by African buffalo

genomes. Sci Rep. 2021;11: 4540. doi:10.1038/s41598-021-83823-8

25. Okello JBA, Wittemyer G, Rasmussen HB, Arctander P, Nyakaana S, Douglas-Hamilton I,

et al. Effective population size dynamics reveal impacts of historic climatic events and

recent anthropogenic pressure in African elephants. Molecular Ecology. 2008;17: 3788–

3799. doi:10.1111/j.1365-294X.2008.03871.x

26. Okello JBA, Nyakaana S, Masembe C, Siegismund HR, Arctander P. Mitochondrial DNA

variation of the common hippopotamus: evidence for a recent population expansion.

Heredity. 2005;95: 206–215. doi:10.1038/sj.hdy.6800711

27. Stoffel C, Dufresnes C, Okello JBA, Noirard C, Joly P, Nyakaana S, et al. Genetic

consequences of population expansions and contractions in the common hippopotamus

(Hippopotamus amphibius) since the Late Pleistocene. Molecular Ecology. 2015;24: 2507–

2520. doi:10.1111/mec.13179

28. Thompson LG, Mosley-Thompson E, Davis ME, Henderson KA, Brecher HH, Zagorodnov

VS, et al. Kilimanjaro ice core records: evidence of Holocene climate change in tropical

Africa. Science. 2002;298: 589–593. doi:10.1126/science.1073198

29. Marchant R, Hooghiemstra H. Rapid environmental change in African and South American

tropics around 4000 years before present: a review. Earth-Science Reviews. 2004;66: 217–

260. doi:10.1016/j.earscirev.2004.01.003

30. Kiage LM, Liu K. Late Quaternary paleoenvironmental changes in East Africa: a review of

multiproxy evidence from palynology, lake sediments, and associated records. Progress in

Physical Geography. 2006;30: 633–658. doi:10.1177/0309133306071146

31. Durant SM, Becker MS, Creel S, Bashir S, Dickman AJ, Beudels-Jamar RC, et al.

Developing fencing policies for dryland ecosystems. Journal of Applied Ecology. 2015;52:

544–551. doi:10.1111/1365-2664.12415

32. M'soka J, Creel S, Becker MS, Murdoch JD. Ecological and anthropogenic effects on the

density of migratory and resident ungulates in a human-inhabited protected area. Afr J Ecol.

2017;55: 618–631. doi:10.1111/aje.12398

33. Watson F, Becker MS, Smit D, Droge E, Mukula T, Martens S, et al. Predation strongly

limits demography of a keystone migratory herbivore in a recovering transfrontier

ecosystem. Ecology and Evolution. 2022;12: e9414. doi:10.1002/ece3.9414

34. Nyirenda VR, Nkhata BA. Collaborative governance and benefit sharing in Liuwa Plain

National Park, western Zambia. PARKS. 2013;19: 103–114.

doi:10.2305/IUCN.CH.2013.PARKS-19-1.VRN.en

35. African Parks Network. Aerial survey confirms substantial increase in important indicator

species in Liuwa. 2013 [cited 18 Jun 2024]. Available: https://www.african-

parks.org/newsroom/press-releases/aerial-survey-confirms-substantial-increase-Important-

indicator-species-liuwa

36. Dröge E, Creel S, Becker M, Christianson D, M'Soka J, Watson F. Response of wildebeest

(Connochaetes taurinus) movements to spatial variation in long term risks from a complete

predator guild. Biological Conservation. 2019;233: 139–151.

doi:10.1016/j.biocon.2019.02.031

37. Dedato MN, Robert C, Taillon J, Shafer ABA, Côté SD. Demographic history and

conservation genomics of caribou (Rangifer tarandus) in Québec. Evolutionary

Applications. 2022;15: 2043–2053. doi:10.1111/eva.13495

38. Wootton E, Robert C, Taillon J, Côté SD, Shafer ABA. Genomic health is dependent on

long-term population demographic history. Molecular Ecology. 2023;32: 1943–1954.

doi:10.1111/mec.16863

39. Liu X, Lin L, Sinding M-HS, Bertola LD, Hanghøj K, Quinn L, et al. Introgression and

disruption of migration routes have shaped the genetic integrity of wildebeest populations.

Nat Commun. 2024;15: 2921. doi:10.1038/s41467-024-47015-y

40. Allendorf FW, Hohenlohe PA, Luikart G. Genomics and the future of conservation genetics.

2010.

41. Funk WC, McKay JK, Hohenlohe PA, Allendorf FW. Harnessing genomics for delineating

conservation units. Trends in Ecology & Evolution. 2012;27: 489–496.

doi:10.1016/j.tree.2012.05.012

42. Corbet SW, Robinson TJ. Genetic divergence in South African wildebeest: comparative

cytogenetics and analysis of mitochondrial DNA. J Hered. 1991;82: 447–452.

doi:10.1093/oxfordjournals.jhered.a111126

43. Grobler JP, Van der Bank FH. Genetic variability in South African blue wildebeest

(*Connochaetes taurinus*). Comparative Biochemistry and Physiology Part B: Comparative

Biochemistry. 1993;106: 755–759. doi:10.1016/0305-0491(93)90159-3

44. Corbet SW, Grant WS, Robinson TJ. Genetic divergence in South African wildebeest:

analysis of allozyme variability. J Hered. 1994;85: 479–483.

doi:10.1093/oxfordjournals.jhered.a111504

45. Arctander P, Johansen C, Coutellec-Vreto MA. Phylogeography of three closely related

African bovids (tribe Alcelaphini). Mol Biol Evol. 1999;16: 1724–1739.

doi:10.1093/oxfordjournals.molbev.a026085

46. Grobler JP, Hartl GB, Grobler N, Kotze A, Botha K, Tiedemann R. The genetic status of an

isolated black wildebeest (*Connochaetes gnou*) population from the Abe Bailey Nature

Reserve, South Africa: Microsatellite data on a putative past hybridization with blue

625  wildebeest (*C. taurinus*). Mammalian Biology - Zeitschrift für Säugetierkunde. 2005;70:

626  35–45. doi:10.1078/1616-5047-00174

627 47. Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC, Foll M. Robust demographic

628  inference from genomic and SNP data. PLOS Genetics. 2013;9: e1003905.

629  doi:10.1371/journal.pgen.1003905

630 48. Liu X, Fu Y-X. Exploring population size changes using SNP frequency spectra. Nat Genet.

631  2015;47: 555–559. doi:10.1038/ng.3254

632 49. Liu X, Fu Y-X. Stairway Plot 2: demographic history inference with folded SNP frequency

633  spectra. Genome Biology. 2020;21: 280. doi:10.1186/s13059-020-02196-9

634 50. DaCosta JM, Sorenson MD. Amplification Biases and Consistent Recovery of Loci in a

635  Double-Digest RAD-seq Protocol. PLOS ONE. 2014;9: e106713.

636  doi:10.1371/journal.pone.0106713

637 51. Andrews S. FastQC: a quality control tool for high throughput sequence data. In: Babraham

638  Bioinformatics [Internet]. 2010. Available:

639  https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

640 52. Edgar RC. Search and clustering orders of magnitude faster than BLAST. Bioinformatics.

641  2010;26: 2460–2461. doi:10.1093/bioinformatics/btq461

642 53. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J

643  Mol Biol. 1990;215: 403–410. doi:10.1016/S0022-2836(05)80360-2

644 54. Chen L, Qiu Q, Jiang Y, Wang K, Lin Z, Li Z, et al. Large-scale ruminant genome

645  sequencing provides insights into their evolution and distinct traits. Science. 2019;364:

646  eaav6202. doi:10.1126/science.aav6202

55. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput.

Nucleic Acids Res. 2004;32: 1792–1797. doi:10.1093/nar/gkh340

56. Wickham H. tidyverse: Easily Install and Load the "Tidyverse". In: R package   version

1.2.1. [Internet]. 2017. Available: https://CRAN.R-project.org/package=tidyverse

57. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R

Foundation for Statistical Computing; 2024. Available: https://www.R-project.org/

58. O'Leary SJ, Puritz JB, Willis SC, Hollenbeck CM, Portnoy DS. These aren't the loci you're

looking for: Principles of effective SNP filtering for molecular ecologists. Molecular

Ecology. 2018;27: 3193–3206. doi:10.1111/mec.14792

59. Emerson KJ, Merz CR, Catchen JM, Hohenlohe PA, Cresko WA, Bradshaw WE, et al.

Resolving postglacial phylogeography using high-throughput sequencing. PNAS.

2010;107: 16196–16200. doi:10.1073/pnas.1006538107

60. Willis SC, Hollenbeck CM, Puritz JB, Gold JR, Portnoy DS. Haplotyping RAD loci: an

efficient method to filter paralogs and account for physical linkage. Molecular Ecology

Resources. 2017;17: 955–965. doi:10.1111/1755-0998.12647

61. Lischer HEL, Excoffier L. PGDSpider: an automated data conversion tool for connecting

population genetics and genomics programs. Bioinformatics. 2012;28: 298–299.

doi:10.1093/bioinformatics/btr642

62. Jombart T. adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics. 2008;24: 1403–1405. doi:10.1093/bioinformatics/btn129

63. Jombart T, Ahmed I. adegenet 1.3-1: new tools for the analysis of genome-wide SNP data.

Bioinformatics. 2011;27: 3070–3071. doi:10.1093/bioinformatics/btr521

64. Paradis E. pegas: an R package for population genetics with an integrated–modular

approach. Bioinformatics. 2010;26: 419–420. doi:10.1093/bioinformatics/btp696

65. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful

approach to multiple testing. Journal of the Royal Statistical Society Series B

(Methodological). 1995;57: 289–300.

66. Conde IC, Alvarez J de U. sgof: multiple hypothesis testing. In: R package version 2.3

[Internet]. 2016. Available: https://CRAN.R-project.org/package=sgof

67. White T, van der Ende J, Nichols TE. Beyond Bonferroni revisited: concerns over inflated

false positive research findings in the fields of conservation genetics, biology, and

medicine. Conserv Genet. 2019;20: 927–937. doi:10.1007/s10592-019-01178-0

68. Goudet J. hierfstat, a package for r to compute and test hierarchical F-statistics. Molecular

Ecology Notes. 2005;5: 184–186. doi:10.1111/j.1471-8286.2004.00828.x

69. Waples RS. A bias correction for estimates of effective population size based on linkage

disequilibrium at unlinked gene loci. Conserv Genet. 2006;7: 167. doi:10.1007/s10592-005-

9100-y

70. Waples RS, Do C. ldne: a program for estimating effective population size from data on

linkage disequilibrium. Mol Ecol Resour. 2008;8: 753–756. doi:10.1111/j.1755-

0998.2007.02061.x

71. Do C, Waples RS, Peel D, Macbeth GM, Tillett BJ, Ovenden JR. NeEstimator v2: re-

implementation of software for the estimation of contemporary effective population size

(Ne) from genetic data. Mol Ecol Resour. 2014;14: 209–214. doi:10.1111/1755-0998.12157

72. Jones AT, Ovenden JR, Wang Y-G. Improved confidence intervals for the linkage

disequilibrium method for estimating effective population size. Heredity (Edinb). 2016;117:

217–223. doi:10.1038/hdy.2016.19

73. Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus

genotype data. Genetics. 2000;155: 945–959.

74. Falush D, Stephens M, Pritchard JK. Inference of population structure using multilocus

genotype data: linked loci and correlated allele frequencies. Genetics. 2003;164: 1567–

1587.

75. Falush D, Stephens M, Pritchard JK. Inference of population structure using multilocus

genotype data: dominant markers and null alleles. Molecular Ecology Notes. 2007;7: 574–

578. doi:10.1111/j.1471-8286.2007.01758.x

76. Hubisz MJ, Falush D, Stephens M, Pritchard JK. Inferring weak population structure with

the assistance of sample group information. Molecular Ecology Resources. 2009;9: 1322–

1332. doi:10.1111/j.1755-0998.2009.02591.x

77. Porras-Hurtado L, Ruiz Y, Santos C, Phillips C, Carracedo Á, Lareu MV. An overview of

STRUCTURE: applications, parameter settings, and supporting software. Front Genet.

2013;4. doi:10.3389/fgene.2013.00098

78. Earl DA, vonHoldt BM. STRUCTURE HARVESTER: a website and program for

visualizing STRUCTURE output and implementing the Evanno method. Conservation

Genet Resour. 2012;4: 359–361. doi:10.1007/s12686-011-9548-7

79. Earl D. Structure Harvester Web v0.6.94. 2014. Available:

http://taylor0.biology.ucla.edu/structureHarvester/

80. Evanno G, Regnaut S, Goudet J. Detecting the number of clusters of individuals using the

software STRUCTURE: a simulation study. Molecular Ecology. 2005;14: 2611–2620.

doi:10.1111/j.1365-294X.2005.02553.x

81. Francis RM. pophelper: an R package and web app to analyse and visualize population

structure. Molecular Ecology Resources. 2017;17: 27–32. doi:10.1111/1755-0998.12509

82. Brumfield RT, Beerli P, Nickerson DA, Edwards SV. The utility of single nucleotide

polymorphisms in inferences of population history. Trends in Ecology & Evolution.

2003;18: 249–256. doi:10.1016/S0169-5347(03)00018-1

83. Kumar S, Subramanian S. Mutation rates in mammalian genomes. Proc Natl Acad Sci U S

A. 2002;99: 803–808. doi:10.1073/pnas.022629899

84. IUCN. Connochaetes taurinus: IUCN SSC antelope specialist group: The IUCN red list of

threatened species 2008. International Union for Conservation of Nature; 2008.

doi:10.2305/IUCN.UK.2008.RLTS.T5229A11120316.en

85. Akaike H. A new look at the statistical model identification. IEEE Trans Automat Contr.

1974;19: 716–723. doi:10.1109/TAC.1974.1100705

86. Ritz C, Spiess A-N. qpcR: an R package for sigmoidal model selection in quantitative real-

time polymerase chain reaction analysis. Bioinformatics. 2008;24: 1549–1551.

doi:10.1093/bioinformatics/btn227

87. Adelson DL, Raison JM, Edgar RC. Characterization and distribution of retrotransposons

and simple sequence repeats in the bovine genome. PNAS. 2009;106: 12855–12860.

doi:10.1073/pnas.0901282106

88. Coetzer WG, Grobler JP. Genetic variation among different springbok (Antidorcas

marsupialis) colour variants. Mammalian Biology. 2019;99: 42–53.

doi:10.1016/j.mambio.2019.10.006

89. LaCava MEF, Gagne RB, Stowell SML, Gustafson KD, Buerkle CA, Knox L, et al.

Pronghorn population genomics show connectivity in the core of their range. J Mammal.

2020;101: 1061–1071. doi:10.1093/jmammal/gyaa054

90. Bourgeois S, Senn H, Kaden J, Taggart JB, Ogden R, Jeffery KJ, et al. Single-nucleotide

polymorphism discovery and panel characterization in the African forest elephant. Ecology

and Evolution. 2018;8: 2207–2217. doi:10.1002/ece3.3854

91. van Deventer R, Rhode C, Marx M, Roodt-Wilding R. The development of genome-wide

single nucleotide polymorphisms in blue wildebeest using the DArTseq platform.

Genomics. 2020;112: 3455–3464. doi:10.1016/j.ygeno.2020.04.032

92. Georgiadis NJ, Kat PW, Oketch H, Patton J. Allozyme divergence within the Bovidae.

Evolution. 1990;44: 2135–2149. doi:10.1111/j.1558-5646.1990.tb04317.x

93. Wright S. Evolution in Mendelian populations. Genetics. 1931;16: 97–159.

94. Frankham R. Relationship of genetic variation to population size in wildlife. Conservation

Biology. 1996;10: 1500–1508. doi:10.1046/j.1523-1739.1996.10061500.x

95. Palstra FP, Ruzzante DE. Genetic estimates of contemporary effective population size: what

can they tell us about the importance of genetic stochasticity for wild population

persistence? Molecular Ecology. 2008;17: 3428–3447. doi:10.1111/j.1365-

294X.2008.03842.x

96. O'Ryan C, Harley EH, Bruford MW, Beaumont M, Wayne RK, Cherry MI. Microsatellite

analysis of genetic diversity in fragmented South African buffalo populations. Animal

Conservation. 1998;1: 85–94. doi:10.1111/j.1469-1795.1998.tb00015.x

97. Palkopoulou E, Lipson M, Mallick S, Nielsen S, Rohland N, Baleka S, et al. A

comprehensive genomic history of extinct and living elephants. Proceedings of the National

Academy of Sciences. 2018;115: E2566–E2574. doi:10.1073/pnas.1720554115

98. Storz JF, Beaumont MA, Alberts SC. Genetic evidence for long-term population decline in a

savannah-dwelling primate: inferences from a hierarchical bayesian model. Molecular

Biology and Evolution. 2002;19: 1981–1990. doi:10.1093/oxfordjournals.molbev.a004022

99. Barnosky AD, Koch PL, Feranec RS, Wing SL, Shabel AB. Assessing the causes of late

Pleistocene extinctions on the continents. Science. 2004;306: 70–75.

doi:10.1126/science.1101476

Fig. 1. Location of study area. (a) Satellite map of Zambia and surrounding area with Liuwa

Plain National Park, where blue wildebeest tissue samples were collected for this study, outlined

in white and marked with a star. Basemap: Google Satellite (obtained through

QuickMapServices QGIS plugin), Map data © 2024 Google. (b) Image of a blue wildebeest

(Photograph by S. Szarmach). (c) Location of Zambia (green) within Africa.

Fig. 2. Schematic of the five demographic scenarios modeled in the coalescent simulator

fastsimcoal2. For each model, parameters describing effective population size (N) and select

times before present (T) were estimated.

Fig. 3. Expected and observed heterozygosity for 1,730 SNPs genotyped in Greater Liuwa

Ecosystem blue wildebeest (n = 75). Boxplots show median and interquartile range.

Fig. 4 Blue wildebeest show no signs of population subdivision within the Greater Liuwa

Ecosystem, based on (a) model-based clustering in STRUCTURE (shown are membership

coefficients for *K*=2 clusters for 75 individuals), and (b) principal component analysis (PCA)

based on 1,730 SNPs.

Fig. 5. Stairway plot representing inferred changes in effective population size (N_e) over time for

the Greater Liuwa Ecosystem blue wildebeest population. The red line shows the median

estimated effective population size, the dark grey lines encompass the 75% CI, and the light gray

lines contain the 95% CI, determined from 200 bootstrapped site frequency spectrums.

Table 1. Comparison of likelihood scores and Akaike information criterion (AIC) values for the

highest likelihood model produced for each demographic scenario tested using the coalescent

simulator fastsimcoal2. The best supported model is shown in bold.

Model	Log likelihood	k	AIC	ΔAIC	Akaike weight (w_i)
Constant size	-11,513.55	1	23,029.10	120.70	0.00
Expansion	-11,514.12	3	23,034.24	125.84	0.00
Decline	-11,463.62	3	22,933.25	24.85	0.00
Bottleneck	-11,462.90	4	22,933.79	25.39	0.00
Expansion-Decline	**-11,449.20**	**5**	**22,908.40**	**0.00**	**1.00**